

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12403335
Kind Code	B2
Date of Patent	September 02, 2025
Inventor(s)	Ciaramella; Giuseppe et al.

Betacoronavirus MRNA vaccines

Abstract

The disclosure relates to respiratory virus ribonucleic acid (RNA) vaccines and combination vaccines, as well as methods of using the vaccines and compositions comprising the vaccines.

Inventors: Ciaramella; Giuseppe (Sudbury, MA), Himansu; Sunny (Winchester, MA)

Applicant: ModernaTX, Inc. (Cambridge, MA)

Family ID: 58558175

Assignee: ModernaTX, Inc. (Cambridge, MA)

Appl. No.: 18/528323

Filed: December 04, 2023

Prior Publication Data

Document Identifier	Publication Date
US 20240335526 A1	Oct. 10, 2024

Related U.S. Application Data

continuation parent-doc US 16897734 20200610 US 11872278 child-doc US 18528323
continuation parent-doc US 16368270 20190328 US 10702599 child-doc US 16897734
continuation parent-doc US 16040981 20180720 US 10272150 child-doc US 16368270
continuation parent-doc US 15674599 20170811 US 10064934 child-doc US 16040981
continuation parent-doc WO PCT/US2016/058327 20161021 PENDING child-doc US 15674599
us-provisional-application US 62247394 20151028
us-provisional-application US 62247362 20151028
us-provisional-application US 62247483 20151028
us-provisional-application US 62247297 20151028
us-provisional-application US 62244946 20151022
us-provisional-application US 62244802 20151022

us-provisional-application US 62245031 20151022
us-provisional-application US 62244813 20151022
us-provisional-application US 62244837 20151022

Publication Classification

Int. Cl.: **A61P31/14** (20060101); **A61K39/12** (20060101); **A61K39/155** (20060101);
A61K39/215 (20060101); **A61P11/00** (20060101); **C07K16/10** (20060101); A61K39/00
(20060101)

U.S. Cl.:

CPC **A61K39/12** (20130101); **A61K39/155** (20130101); **A61K39/215** (20130101);
A61P11/00 (20180101); **A61P31/14** (20180101); **C07K16/10** (20130101); **C07K16/1002**
(20230801); **C07K16/1027** (20130101); A61K2039/53 (20130101); A61K2039/55511
(20130101); A61K2039/55555 (20130101); A61K2039/6018 (20130101); A61K2039/70
(20130101); C07K2317/76 (20130101); C12N2760/18034 (20130101);
C12N2760/18334 (20130101); C12N2760/18434 (20130101); C12N2760/18534
(20130101); C12N2760/18634 (20130101); C12N2770/20034 (20130101); Y02A50/30
(20180101)

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
3906092	12/1974	Hilleman et al.	N/A	N/A
4790987	12/1987	Compans et al.	N/A	N/A
5169628	12/1991	Wathen	N/A	N/A
5427782	12/1994	Compans et al.	N/A	N/A
6225091	12/2000	Klein et al.	N/A	N/A
6500419	12/2001	Hone et al.	N/A	N/A
6514948	12/2002	Raz et al.	N/A	N/A
7001890	12/2005	Wagner et al.	N/A	N/A
7208161	12/2006	Murphy et al.	N/A	N/A
7449324	12/2007	Fouchier et al.	N/A	N/A
7531342	12/2008	Fouchier et al.	N/A	N/A
7671186	12/2009	Klein et al.	N/A	N/A
7704720	12/2009	Tang et al.	N/A	N/A
8217016	12/2011	Hoerr et al.	N/A	N/A
8252289	12/2011	Eleouët et al.	N/A	N/A
8691966	12/2013	Kariko et al.	N/A	N/A
8710200	12/2013	Schrum et al.	N/A	N/A
8722341	12/2013	Fouchier et al.	N/A	N/A
8734853	12/2013	Sood et al.	N/A	N/A
8754062	12/2013	De Fougerolles et al.	N/A	N/A

8822663	12/2013	Schrum et al.	N/A	N/A
8841433	12/2013	Fouchier et al.	N/A	N/A
8889146	12/2013	Blais et al.	N/A	N/A
8927206	12/2014	De Jong et al.	N/A	N/A
8980864	12/2014	Hoge et al.	N/A	N/A
8999380	12/2014	Bancel et al.	N/A	N/A
9192651	12/2014	Chakraborty et al.	N/A	N/A
9192661	12/2014	Jain et al.	N/A	N/A
9221891	12/2014	Bancel et al.	N/A	N/A
9271996	12/2015	De Fougerolles et al.	N/A	N/A
9283287	12/2015	Bancel et al.	N/A	N/A
9295689	12/2015	De Fougerolles et al.	N/A	N/A
9303079	12/2015	Bancel et al.	N/A	N/A
9376726	12/2015	Fouchier et al.	N/A	N/A
9464124	12/2015	Bancel et al.	N/A	N/A
9512456	12/2015	Wang et al.	N/A	N/A
9533047	12/2016	de Fougerolles et al.	N/A	N/A
9567653	12/2016	Fouchier et al.	N/A	N/A
9572896	12/2016	Bancel et al.	N/A	N/A
9597380	12/2016	Chakraborty et al.	N/A	N/A
9623095	12/2016	Kallen et al.	N/A	N/A
9669089	12/2016	Thess et al.	N/A	N/A
9675668	12/2016	Bancel et al.	N/A	N/A
9790531	12/2016	Wang et al.	N/A	N/A
9803199	12/2016	Koizumi et al.	N/A	N/A
9868691	12/2017	Benenato et al.	N/A	N/A
9872900	12/2017	Ciaramella et al.	N/A	N/A
9937196	12/2017	Jain et al.	N/A	N/A
10023626	12/2017	Bolen et al.	N/A	N/A
10064934	12/2017	Ciaramella et al.	N/A	N/A
10064935	12/2017	Ciaramella et al.	N/A	N/A
10124055	12/2017	Ciaramella et al.	N/A	N/A
10207010	12/2018	Besin et al.	N/A	N/A
10232055	12/2018	Kariko et al.	N/A	N/A
10272150	12/2018	Ciaramella et al.	N/A	N/A
10273269	12/2018	Ciaramella	N/A	N/A
10286086	12/2018	Roy et al.	N/A	N/A
10323076	12/2018	Ellsworth et al.	N/A	N/A
10385088	12/2018	Fraley et al.	N/A	N/A
10449244	12/2018	Ciaramella et al.	N/A	N/A
10465190	12/2018	Chen et al.	N/A	N/A
10493143	12/2018	Ciaramella et al.	N/A	N/A
10526629	12/2019	Rabideau et al.	N/A	N/A
10543269	12/2019	Ciaramella et al.	N/A	N/A
10653712	12/2019	Hoge	N/A	N/A
10653767	12/2019	Ciaramella et al.	N/A	N/A
10695419	12/2019	Ciaramella et al.	N/A	N/A
10702599	12/2019	Ciaramella et al.	N/A	N/A
10702600	12/2019	Ciaramella et al.	N/A	N/A
10857105	12/2019	Benenato et al.	N/A	N/A

10906944	12/2020	He et al.	N/A	N/A
10912826	12/2020	Thess et al.	N/A	N/A
10925958	12/2020	Ciaramella	N/A	N/A
10960070	12/2020	Graham et al.	N/A	N/A
11027025	12/2020	Hoge et al.	N/A	N/A
11045540	12/2020	Ciaramella	N/A	N/A
11103578	12/2020	Ciaramella et al.	N/A	N/A
11266735	12/2021	Kallen et al.	N/A	N/A
11351242	12/2021	Lori et al.	N/A	N/A
11384352	12/2021	Miracco	N/A	N/A
11406703	12/2021	Kramarczyk et al.	N/A	N/A
11464848	12/2021	Ciaramella et al.	N/A	N/A
11485960	12/2021	Dousis et al.	N/A	N/A
11497807	12/2021	Ciaramella et al.	N/A	N/A
11564893	12/2022	Smith	N/A	N/A
11576961	12/2022	Ciaramella et al.	N/A	N/A
11638693	12/2022	Geall	N/A	N/A
11638694	12/2022	Geall	N/A	N/A
11643441	12/2022	Ciaramella et al.	N/A	N/A
11666534	12/2022	Geall	N/A	N/A
11696946	12/2022	Ciaramella	N/A	N/A
11752206	12/2022	Ciaramella et al.	N/A	N/A
11766401	12/2022	Geall	N/A	N/A
11786467	12/2022	Geall	N/A	N/A
11786607	12/2022	Hoge et al.	N/A	N/A
11851694	12/2022	Mauger et al.	N/A	N/A
11866696	12/2023	Issa et al.	N/A	N/A
11872278	12/2023	Ciaramella et al.	N/A	N/A
11905525	12/2023	Brito et al.	N/A	N/A
11911453	12/2023	Ciaramella et al.	N/A	N/A
11912982	12/2023	Issa et al.	N/A	N/A
2003/0092653	12/2002	Kisich et al.	N/A	N/A
2003/0232061	12/2002	Fouchier et al.	N/A	N/A
2004/0005545	12/2003	Fouchier et al.	N/A	N/A
2004/0096451	12/2003	Young et al.	N/A	N/A
2005/0032730	12/2004	Von Der Mulbe et al.	N/A	N/A
2005/0059624	12/2004	Hoerr et al.	N/A	N/A
2005/0250723	12/2004	Hoerr et al.	N/A	N/A
2006/0002958	12/2005	Naylor et al.	N/A	N/A
2006/0172003	12/2005	Meers et al.	N/A	N/A
2006/0228367	12/2005	Ulbrandt et al.	N/A	N/A
2007/0280929	12/2006	Hoerr et al.	N/A	N/A
2008/0025944	12/2007	Hoerr et al.	N/A	N/A
2008/0171079	12/2007	Hanon et al.	N/A	N/A
2008/0171711	12/2007	Hoerr et al.	N/A	N/A
2009/0123529	12/2008	Xiaomao	N/A	N/A
2009/0162395	12/2008	Crowe et al.	N/A	N/A
2010/0203076	12/2009	Fotin-Mleczeek et al.	N/A	N/A
2010/0239608	12/2009	Von Der Mulbe et al.	N/A	N/A
2010/0272747	12/2009	Chow et al.	N/A	N/A

2010/0291156	12/2009	Barner et al.	N/A	N/A
2010/0305196	12/2009	Probst et al.	N/A	N/A
2011/0135645	12/2010	Williamson et al.	N/A	N/A
2011/0250225	12/2010	Fotin-Mleczek et al.	N/A	N/A
2011/0269950	12/2010	Von Der Mulbe et al.	N/A	N/A
2012/0009221	12/2011	Hoerr et al.	N/A	N/A
2012/0045471	12/2011	Haller et al.	N/A	N/A
2012/0101148	12/2011	Aking et al.	N/A	N/A
2012/0177701	12/2011	Ilyinskii et al.	N/A	N/A
2012/0213812	12/2011	Lipford et al.	N/A	N/A
2012/0219573	12/2011	Baumhof et al.	N/A	N/A
2012/0276209	12/2011	Cullis et al.	N/A	N/A
2013/0022538	12/2012	Rossi	N/A	N/A
2013/0078281	12/2012	He et al.	N/A	N/A
2013/0102034	12/2012	Schrum et al.	N/A	N/A
2013/0121988	12/2012	Hoerr et al.	N/A	N/A
2013/0142818	12/2012	Baumhof et al.	N/A	N/A
2013/0183355	12/2012	Jain et al.	N/A	N/A
2013/0195867	12/2012	Hoerr et al.	N/A	N/A
2013/0195967	12/2012	Guild et al.	N/A	N/A
2013/0195969	12/2012	Geall et al.	N/A	N/A
2013/0202684	12/2012	Geall et al.	N/A	N/A
2013/0236974	12/2012	De Fougerolles	N/A	N/A
2013/0243848	12/2012	Lobovkina et al.	N/A	N/A
2013/0245103	12/2012	de Fougerolles et al.	N/A	N/A
2013/0259923	12/2012	Bancel et al.	N/A	N/A
2013/0266640	12/2012	De Fougerolles et al.	N/A	N/A
2013/0295043	12/2012	Kallen et al.	N/A	N/A
2013/0336998	12/2012	Kallen et al.	N/A	N/A
2014/0024076	12/2013	Tang et al.	N/A	N/A
2014/0037660	12/2013	Folin-Mleczek et al.	N/A	N/A
2014/0147432	12/2013	Bancel et al.	N/A	N/A
2014/0148502	12/2013	Bancel et al.	N/A	N/A
2014/0193482	12/2013	Bancel et al.	N/A	N/A
2014/0206752	12/2013	Afeyan et al.	N/A	N/A
2014/0255472	12/2013	Geall et al.	N/A	N/A
2014/0271829	12/2013	Lilja et al.	N/A	N/A
2014/0308304	12/2013	Manoharan et al.	N/A	N/A
2014/0370497	12/2013	Fouchier et al.	N/A	N/A
2014/0378538	12/2013	Bancel	N/A	N/A
2015/0051268	12/2014	Bancel et al.	N/A	N/A
2015/0056253	12/2014	Bancel et al.	N/A	N/A
2015/0093413	12/2014	Thess et al.	N/A	N/A
2015/0126589	12/2014	Geiger et al.	N/A	N/A
2015/0141499	12/2014	Bancel et al.	N/A	N/A
2015/0307542	12/2014	Roy et al.	N/A	N/A
2015/0315541	12/2014	Bancel et al.	N/A	N/A
2015/0335728	12/2014	Wong et al.	N/A	N/A
2016/0022580	12/2015	Ramsay et al.	N/A	N/A
2016/0024140	12/2015	Issa et al.	N/A	N/A

2016/0024141	12/2015	Issa et al.	N/A	N/A
2016/0032273	12/2015	Shahrokh et al.	N/A	N/A
2016/0038612	12/2015	Hoge et al.	N/A	N/A
2016/0039884	12/2015	Li et al.	N/A	N/A
2016/0046675	12/2015	Kwong et al.	N/A	N/A
2016/0151474	12/2015	Kallen et al.	N/A	N/A
2016/0194368	12/2015	Hoge et al.	N/A	N/A
2016/0194625	12/2015	Hoge et al.	N/A	N/A
2016/0243221	12/2015	Hoge et al.	N/A	N/A
2016/0271272	12/2015	Bancel et al.	N/A	N/A
2016/0317647	12/2015	Ciaramella et al.	N/A	N/A
2016/0331828	12/2015	Ciaramella et al.	N/A	N/A
2016/0361411	12/2015	Gindy et al.	N/A	N/A
2017/0043037	12/2016	Kariko et al.	N/A	N/A
2017/0065675	12/2016	Bancel et al.	N/A	N/A
2017/0130255	12/2016	Wang et al.	N/A	N/A
2017/0202979	12/2016	Chakraborty et al.	N/A	N/A
2017/0204152	12/2016	Nelson et al.	N/A	N/A
2017/0340724	12/2016	Ciaramella et al.	N/A	N/A
2018/0000953	12/2017	Almarsson et al.	N/A	N/A
2018/0002393	12/2017	Bancel et al.	N/A	N/A
2018/0008694	12/2017	Ciaramella et al.	N/A	N/A
2018/0028645	12/2017	Ciaramella et al.	N/A	N/A
2018/0028664	12/2017	Besin et al.	N/A	N/A
2018/0237849	12/2017	Thompson	N/A	N/A
2018/0243225	12/2017	Ciaramella	N/A	N/A
2018/0243230	12/2017	Smith	N/A	N/A
2018/0256628	12/2017	Hoge et al.	N/A	N/A
2018/0271795	12/2017	Martini et al.	N/A	N/A
2018/0271970	12/2017	Ciaramella et al.	N/A	N/A
2018/0273977	12/2017	Mousavi et al.	N/A	N/A
2018/0274009	12/2017	Marquardt et al.	N/A	N/A
2018/0280496	12/2017	Ciaramella et al.	N/A	N/A
2018/0289792	12/2017	Ciaramella et al.	N/A	N/A
2018/0303929	12/2017	Ciaramella et al.	N/A	N/A
2018/0311336	12/2017	Ciaramella et al.	N/A	N/A
2018/0311343	12/2017	Huang et al.	N/A	N/A
2018/0318409	12/2017	Valiante et al.	N/A	N/A
2018/0326039	12/2017	Haruta	N/A	N/A
2018/0363019	12/2017	Hoge	N/A	N/A
2018/0369374	12/2017	Frederick et al.	N/A	N/A
2018/0371047	12/2017	Ticho et al.	N/A	N/A
2019/0002890	12/2018	Martini et al.	N/A	N/A
2019/0008938	12/2018	Ciaramella et al.	N/A	N/A
2019/0085368	12/2018	Bancel et al.	N/A	N/A
2019/0099481	12/2018	Ciaramella et al.	N/A	N/A
2019/0125839	12/2018	Frederick et al.	N/A	N/A
2019/0175517	12/2018	Martini et al.	N/A	N/A
2019/0175727	12/2018	Huang et al.	N/A	N/A
2019/0192646	12/2018	Cohen et al.	N/A	N/A

2019/0192653	12/2018	Hoge et al.	N/A	N/A
2019/0240317	12/2018	Ciaramella et al.	N/A	N/A
2019/0241633	12/2018	Fotin-Mleczek et al.	N/A	N/A
2019/0275170	12/2018	Benenato et al.	N/A	N/A
2019/0298657	12/2018	Martini et al.	N/A	N/A
2019/0298658	12/2018	Benenato	N/A	N/A
2019/0300906	12/2018	Martini et al.	N/A	N/A
2019/0314292	12/2018	Benenato et al.	N/A	N/A
2019/0314493	12/2018	Ciaramella et al.	N/A	N/A
2019/0336452	12/2018	Brader	N/A	N/A
2019/0336595	12/2018	Ciaramella	N/A	N/A
2019/0351040	12/2018	Valiante et al.	N/A	N/A
2019/0351047	12/2018	Jasny et al.	N/A	N/A
2019/0382774	12/2018	Hoge et al.	N/A	N/A
2019/0390181	12/2018	Benenato et al.	N/A	N/A
2020/0030432	12/2019	Ciaramella et al.	N/A	N/A
2020/0032274	12/2019	Mauger et al.	N/A	N/A
2020/0038499	12/2019	Narayanan et al.	N/A	N/A
2020/0054737	12/2019	Ciaramella et al.	N/A	N/A
2020/0061185	12/2019	Graham et al.	N/A	N/A
2020/0069599	12/2019	Smith et al.	N/A	N/A
2020/0069793	12/2019	Ciaramella	N/A	N/A
2020/0069794	12/2019	Ciaramella et al.	N/A	N/A
2020/0071689	12/2019	Miracco	N/A	N/A
2020/0085916	12/2019	Martini et al.	N/A	N/A
2020/0109420	12/2019	Brito et al.	N/A	N/A
2020/0129445	12/2019	Patel et al.	N/A	N/A
2020/0129608	12/2019	Ciaramella et al.	N/A	N/A
2020/0129615	12/2019	Ciaramella et al.	N/A	N/A
2020/0163878	12/2019	Baumhof et al.	N/A	N/A
2020/0239869	12/2019	Issa et al.	N/A	N/A
2020/0254086	12/2019	Hoge et al.	N/A	N/A
2020/0282046	12/2019	Ciaramella et al.	N/A	N/A
2020/0282047	12/2019	Ciaramella et al.	N/A	N/A
2020/0297634	12/2019	Karmali et al.	N/A	N/A
2020/0306191	12/2019	Schariter et al.	N/A	N/A
2020/0338004	12/2019	Hansson et al.	N/A	N/A
2020/0368162	12/2019	Martini	N/A	N/A
2020/0399322	12/2019	Baumhof et al.	N/A	N/A
2020/0407402	12/2019	He et al.	N/A	N/A
2021/0023199	12/2020	Kallen et al.	N/A	N/A
2021/0030866	12/2020	Kallen et al.	N/A	N/A
2021/0046173	12/2020	Ciaramella et al.	N/A	N/A
2021/0060175	12/2020	Baumhof et al.	N/A	N/A
2021/0087135	12/2020	Benenato et al.	N/A	N/A
2021/0128716	12/2020	Probst et al.	N/A	N/A
2021/0139543	12/2020	He et al.	N/A	N/A
2021/0145982	12/2020	Hoge et al.	N/A	N/A
2021/0162037	12/2020	Jasny et al.	N/A	N/A
2021/0163919	12/2020	Issa et al.	N/A	N/A

2021/0187097	12/2020	Ciaramella et al.	N/A	N/A
2021/0206818	12/2020	Huang et al.	N/A	N/A
2021/0217484	12/2020	Giessel et al.	N/A	N/A
2021/0228707	12/2020	Mektar et al.	N/A	N/A
2021/0228708	12/2020	Smith et al.	N/A	N/A
2021/0268086	12/2020	Zhong et al.	N/A	N/A
2021/0309976	12/2020	Dousis et al.	N/A	N/A
2021/0378980	12/2020	Horhota et al.	N/A	N/A
2022/0031631	12/2021	Almarsson et al.	N/A	N/A
2022/0047518	12/2021	Hennessy et al.	N/A	N/A
2022/0054653	12/2021	Martini et al.	N/A	N/A
2022/0062175	12/2021	Smith et al.	N/A	N/A
2022/0125899	12/2021	Ashburn et al.	N/A	N/A
2022/0145381	12/2021	Elich et al.	N/A	N/A
2022/0236253	12/2021	Hopson	N/A	N/A
2022/0241399	12/2021	Lusso et al.	N/A	N/A
2022/0347292	12/2021	Panther et al.	N/A	N/A
2022/0348900	12/2021	Shamashkin et al.	N/A	N/A
2022/0349006	12/2021	Amato et al.	N/A	N/A
2023/0000970	12/2022	Nachbagauer et al.	N/A	N/A
2023/0142529	12/2022	White et al.	N/A	N/A
2023/0173060	12/2022	Benmohamed	N/A	N/A
2023/0181481	12/2022	White et al.	N/A	N/A
2023/0190761	12/2022	Brader et al.	N/A	N/A
2023/0212645	12/2022	Marquardt et al.	N/A	N/A
2023/0287437	12/2022	Smith et al.	N/A	N/A
2023/0338506	12/2022	Shaw et al.	N/A	N/A
2023/0338512	12/2022	Muik et al.	N/A	N/A
2023/0346914	12/2022	Stewart-Jones et al.	N/A	N/A
2023/0355743	12/2022	Stewart-Jones et al.	N/A	N/A
2024/0100145	12/2023	Bollman et al.	N/A	N/A
2024/0100151	12/2023	Carfi et al.	N/A	N/A
2024/0139309	12/2023	Carfi et al.	N/A	N/A
2024/0173400	12/2023	Ciaramella et al.	N/A	N/A
2024/0181030	12/2023	Himansu et al.	N/A	N/A
2025/0121220	12/2024	Ciaramella et al.	N/A	N/A
2025/0121221	12/2024	Ciaramella et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
2015210364	12/2016	AU	N/A
2473135	12/2002	CA	N/A
3194325	12/2021	CA	N/A
110951756	12/2019	CN	N/A
110974950	12/2019	CN	N/A
110974954	12/2019	CN	N/A
111088283	12/2019	CN	N/A
111218458	12/2019	CN	N/A
111218459	12/2019	CN	N/A
111303254	12/2019	CN	N/A

111518175	12/2019	CN	N/A
111592602	12/2019	CN	N/A
111848753	12/2019	CN	N/A
111939250	12/2019	CN	N/A
111978377	12/2019	CN	N/A
112023035	12/2019	CN	N/A
112028976	12/2019	CN	N/A
112043825	12/2019	CN	N/A
112048005	12/2019	CN	N/A
112094327	12/2019	CN	N/A
112220920	12/2020	CN	N/A
112226445	12/2020	CN	N/A
112266411	12/2020	CN	N/A
112321688	12/2020	CN	N/A
112358533	12/2020	CN	N/A
112480217	12/2020	CN	N/A
112546211	12/2020	CN	N/A
112626089	12/2020	CN	N/A
112794884	12/2020	CN	N/A
1026253	12/1999	EP	N/A
1083232	12/2004	EP	N/A
1905844	12/2007	EP	N/A
2548960	12/2012	EP	N/A
3608308	12/2019	EP	N/A
2015-518472	12/2014	JP	N/A
2016-515815	12/2015	JP	N/A
1987/005326	12/1986	WO	N/A
1990/011092	12/1989	WO	N/A
1993/014778	12/1992	WO	N/A
1995/024485	12/1994	WO	N/A
1995/026204	12/1994	WO	N/A
1995/033835	12/1994	WO	N/A
WO 1996/040945	12/1995	WO	N/A
WO 1998/058956	12/1997	WO	N/A
1999/33982	12/1998	WO	N/A
WO 1999/033982	12/1998	WO	N/A
WO 2003/072720	12/2002	WO	N/A
WO 2004/076645	12/2003	WO	N/A
2005/009346	12/2004	WO	N/A
WO 2005/027963	12/2004	WO	N/A
WO 2005/118813	12/2004	WO	N/A
WO 2006/056027	12/2005	WO	N/A
WO 2006/071903	12/2005	WO	N/A
WO 2006/095259	12/2005	WO	N/A
WO 2007/024708	12/2006	WO	N/A
WO 2007/038862	12/2006	WO	N/A
WO 2007/095976	12/2006	WO	N/A
WO 2007/094854	12/2006	WO	N/A
WO 2008/052770	12/2007	WO	N/A
WO 2009/030254	12/2008	WO	N/A

WO 2009/030481	12/2008	WO	N/A
WO 2009/095226	12/2008	WO	N/A
WO 2009/127230	12/2008	WO	N/A
WO 2010/037408	12/2009	WO	N/A
WO 2010/037539	12/2009	WO	N/A
WO 2010/042877	12/2009	WO	N/A
WO 2010/054401	12/2009	WO	N/A
WO 2010/054406	12/2009	WO	N/A
WO 2010/088927	12/2009	WO	N/A
WO 2010/129709	12/2009	WO	N/A
WO 2010/144740	12/2009	WO	N/A
WO 2010/149743	12/2009	WO	N/A
WO 2011/005799	12/2010	WO	N/A
WO 2011/026641	12/2010	WO	N/A
WO 2011/068810	12/2010	WO	N/A
WO 2011/069529	12/2010	WO	N/A
WO 2011/069586	12/2010	WO	N/A
WO 2011/140627	12/2010	WO	N/A
WO 2011/144358	12/2010	WO	N/A
WO 2012/006369	12/2011	WO	N/A
WO 2012/006372	12/2011	WO	N/A
WO 2012/006377	12/2011	WO	N/A
WO-2012/006378	12/2011	WO	N/A
WO 2012/019630	12/2011	WO	N/A
WO 2012/019780	12/2011	WO	N/A
WO 2012/030901	12/2011	WO	N/A
WO 2012/031043	12/2011	WO	N/A
WO 2012/031046	12/2011	WO	N/A
WO 2012/045075	12/2011	WO	N/A
WO 2012/045082	12/2011	WO	N/A
WO-2012/068295	12/2011	WO	N/A
WO 2012/116714	12/2011	WO	N/A
WO 2012/116715	12/2011	WO	N/A
WO 2012/116810	12/2011	WO	N/A
WO 2012/116811	12/2011	WO	N/A
WO 2013/052523	12/2012	WO	N/A
WO 2013/055905	12/2012	WO	N/A
WO 2013/090186	12/2012	WO	N/A
WO 2013/090648	12/2012	WO	N/A
WO 2013/102203	12/2012	WO	N/A
WO 2013/120628	12/2012	WO	N/A
WO 2013/120629	12/2012	WO	N/A
WO 2013/185069	12/2012	WO	N/A
WO 2014/089239	12/2013	WO	N/A
WO 2014/089486	12/2013	WO	N/A
WO 2014/144196	12/2013	WO	N/A
WO 2014/144711	12/2013	WO	N/A
WO 2014/152027	12/2013	WO	N/A
WO 2014/152030	12/2013	WO	N/A
WO 2014/152200	12/2013	WO	N/A

WO 2014/152774	12/2013	WO	N/A
WO 2014/152940	12/2013	WO	N/A
WO 2014/159813	12/2013	WO	N/A
WO 2014/160243	12/2013	WO	N/A
WO 2015/005253	12/2014	WO	N/A
WO 2015/024668	12/2014	WO	N/A
WO-2015/081155	12/2014	WO	N/A
WO 2015/085318	12/2014	WO	N/A
WO 2015/095340	12/2014	WO	N/A
WO 2015/095346	12/2014	WO	N/A
WO 2015/101414	12/2014	WO	N/A
WO 2015/101415	12/2014	WO	N/A
WO 2015/130584	12/2014	WO	N/A
WO 2015/143335	12/2014	WO	N/A
WO 2016/103238	12/2015	WO	N/A
WO 2016/123864	12/2015	WO	N/A
WO 2016/138160	12/2015	WO	N/A
WO 2016/164762	12/2015	WO	N/A
WO 2016/201377	12/2015	WO	N/A
WO 2017/011773	12/2016	WO	N/A
WO 2017/015457	12/2016	WO	N/A
WO 2017/015463	12/2016	WO	N/A
WO 2017/019935	12/2016	WO	N/A
WO 2017/020026	12/2016	WO	N/A
WO 2017/062513	12/2016	WO	N/A
WO 2017/066789	12/2016	WO	N/A
WO 2017/070601	12/2016	WO	N/A
WO 2017/070616	12/2016	WO	N/A
WO 2017/070618	12/2016	WO	N/A
WO 2017/070620	12/2016	WO	N/A
WO 2017/070622	12/2016	WO	N/A
WO 2017/070623	12/2016	WO	N/A
WO 2017/070626	12/2016	WO	N/A
WO 2017/075038	12/2016	WO	N/A
WO 2017/075531	12/2016	WO	N/A
WO 2017/081082	12/2016	WO	N/A
WO 2017/112865	12/2016	WO	N/A
WO 2017/127750	12/2016	WO	N/A
WO 2017/174564	12/2016	WO	N/A
WO 2017/201333	12/2016	WO	N/A
WO 2017/201340	12/2016	WO	N/A
WO 2017/201342	12/2016	WO	N/A
WO 2017/201347	12/2016	WO	N/A
WO 2017/201349	12/2016	WO	N/A
WO 2018/006052	12/2017	WO	N/A
WO 2018/053209	12/2017	WO	N/A
WO 2018/075980	12/2017	WO	N/A
WO 2018/081459	12/2017	WO	N/A
WO 2018/081462	12/2017	WO	N/A
WO 2018/089851	12/2017	WO	N/A

WO 2018/107088	12/2017	WO	N/A
WO 2018/111967	12/2017	WO	N/A
WO 2018/144082	12/2017	WO	N/A
WO 2018/144778	12/2017	WO	N/A
WO 2018/151816	12/2017	WO	N/A
WO 2018/157009	12/2017	WO	N/A
WO 2018/170245	12/2017	WO	N/A
WO 2018/170256	12/2017	WO	N/A
WO 2018/170260	12/2017	WO	N/A
WO 2018/170270	12/2017	WO	N/A
WO 2018/170347	12/2017	WO	N/A
WO 2018/175783	12/2017	WO	N/A
WO 2018/187590	12/2017	WO	N/A
WO 2018/200737	12/2017	WO	N/A
WO 2018/232355	12/2017	WO	N/A
WO 2018/232357	12/2017	WO	N/A
WO 2019/036670	12/2018	WO	N/A
WO 2019/036682	12/2018	WO	N/A
WO 2019/036683	12/2018	WO	N/A
WO 2019/036685	12/2018	WO	N/A
WO 2019/103993	12/2018	WO	N/A
WO 2019/148101	12/2018	WO	N/A
WO 2019/202035	12/2018	WO	N/A
WO 2020/006242	12/2019	WO	N/A
WO 2020/056370	12/2019	WO	N/A
WO 2020/061284	12/2019	WO	N/A
WO 2020/061295	12/2019	WO	N/A
WO 2020/061367	12/2019	WO	N/A
WO 2020/097291	12/2019	WO	N/A
WO 2020/172239	12/2019	WO	N/A
WO 2020/185811	12/2019	WO	N/A
WO 2020/190750	12/2019	WO	N/A
WO 2020/243561	12/2019	WO	N/A
WO 2021/030533	12/2020	WO	N/A
WO 2021/050864	12/2020	WO	N/A
WO 2021/055811	12/2020	WO	N/A
WO 2021/155243	12/2020	WO	N/A
WO 2021/155274	12/2020	WO	N/A
WO 2021/159040	12/2020	WO	N/A
WO 2021/159130	12/2020	WO	N/A
WO 2021/160346	12/2020	WO	N/A
WO 2021/163365	12/2020	WO	N/A
WO 2021/204175	12/2020	WO	N/A
WO 2021/211343	12/2020	WO	N/A
WO 2021/213945	12/2020	WO	N/A
WO 2021/222304	12/2020	WO	N/A
WO 2021/226436	12/2020	WO	N/A
WO 2021/231901	12/2020	WO	N/A
WO 2021/231929	12/2020	WO	N/A
WO 2021/231963	12/2020	WO	N/A

WO 2021/237084	12/2020	WO	N/A
WO 2021/247817	12/2020	WO	N/A
WO 2022/032154	12/2021	WO	N/A
WO 2022/043551	12/2021	WO	N/A
WO 2022/067010	12/2021	WO	N/A
WO 2022/076562	12/2021	WO	N/A
WO 2022/099003	12/2021	WO	N/A
WO 2022/101469	12/2021	WO	N/A
WO 2022/104265	12/2021	WO	N/A
WO 2022/137133	12/2021	WO	N/A
WO 2022/150717	12/2021	WO	N/A
WO 2022/155524	12/2021	WO	N/A
WO 2022/155530	12/2021	WO	N/A
WO 2022/187698	12/2021	WO	N/A
WO 2022/197624	12/2021	WO	N/A
WO 2022/204491	12/2021	WO	N/A
WO 2022/212191	12/2021	WO	N/A
WO 2022/212442	12/2021	WO	N/A
WO 2022/212711	12/2021	WO	N/A
WO 2022/221335	12/2021	WO	N/A
WO 2022/221336	12/2021	WO	N/A
WO 2022/221359	12/2021	WO	N/A
WO 2022/221440	12/2021	WO	N/A
WO 2022/226277	12/2021	WO	N/A
WO 2022/226318	12/2021	WO	N/A
WO 2022/232585	12/2021	WO	N/A
WO 2022/241103	12/2021	WO	N/A
WO 2022/245888	12/2021	WO	N/A
WO 2022/266010	12/2021	WO	N/A
WO 2022/266012	12/2021	WO	N/A
WO 2022/266389	12/2021	WO	N/A
WO 2023/283642	12/2022	WO	N/A
WO 2023/283645	12/2022	WO	N/A
WO 2023/283651	12/2022	WO	N/A
WO 2023/014649	12/2022	WO	N/A
WO 2023/018773	12/2022	WO	N/A
WO 2023/018923	12/2022	WO	N/A
WO 2023/019181	12/2022	WO	N/A
WO 2023/056401	12/2022	WO	N/A
WO 2023/069625	12/2022	WO	N/A
WO 2023/069895	12/2022	WO	N/A
WO 2023/069900	12/2022	WO	N/A
WO 2023/076658	12/2022	WO	N/A
WO 2023/081311	12/2022	WO	N/A
WO 2023/092069	12/2022	WO	N/A
WO 2023/094713	12/2022	WO	N/A
WO 2023/107999	12/2022	WO	N/A
WO 2023/114307	12/2022	WO	N/A
WO 2023/132885	12/2022	WO	N/A
WO 2023/135333	12/2022	WO	N/A

WO 2023/137149	12/2022	WO	N/A
WO 2023/147091	12/2022	WO	N/A
WO 2023/150256	12/2022	WO	N/A
WO 2023/154818	12/2022	WO	N/A
WO 2023/196914	12/2022	WO	N/A
WO 2023/201204	12/2022	WO	N/A
WO 2023/201294	12/2022	WO	N/A
WO 2023/201296	12/2022	WO	N/A
WO 2023/212696	12/2022	WO	N/A
WO 2023/225524	12/2022	WO	N/A
WO 2023/250119	12/2022	WO	N/A
WO 2024/010993	12/2023	WO	N/A
WO 2024/015890	12/2023	WO	N/A
WO 2024/026005	12/2023	WO	N/A
WO 2024/030369	12/2023	WO	N/A
WO 2024/050483	12/2023	WO	N/A
WO 2024/097874	12/2023	WO	N/A

OTHER PUBLICATIONS

Kurimoto S, et al. *Molecules*. Apr. 3, 2019;24(7):1303. (Year: 2019). cited by examiner

Ernsting MJ, et. al. *J Control Release*. Dec. 28, 2013;172(3):782-94. Epub Sep. 25, 2013 (Year: 2013). cited by examiner

Szebeni J, et al. *Adv Drug Deliv Rev*. Sep. 16, 2011;63(12):1020-30. Epub Jul. 14, 2011 (Year: 2011). cited by examiner

Szebeni J, et. al. *Biochem Biophys Res Commun*. Dec. 18, 2015;468(3):490-7. Epub Jul. 14, 2015. Review (Year: 2015). cited by examiner

Szebeni J. *Mol Immunol*. Oct. 2014;61(2):163-73. Epub Aug. 12, 2014 (Year: 2014). cited by examiner

[No Author Listed] fusion glycoprotein F0 [Human respirovirus 3]. GenBank Accession No. AHX22069. First seen on NCBI on May 14, 2014. 2 pages. cited by applicant

[No Author Listed] Lineage list. Last accessed Oct. 17, 2023 at https://cov-lineages.org/lineage_list.html. 77 pages. cited by applicant

[No. Author Listed], "Messenger RNA" definition. *Collins Dictionary of Science*. 2005:514. cited by applicant

[No Author Listed], "Messenger RNA" definition. *The Penguin Dictionary of Science*. 2009: 3 pages. cited by applicant

[No Author Listed], "Messenger RNA" definition. *Webster's New World Medical Dictionary*. 2008;3:271. cited by applicant

[No Author Listed], 1st International mRNA Health Conference Programme. Universitat Tubingen. Oct. 23-24, 2013: 32 pages. cited by applicant

[No Author Listed], 2022 Breakthrough Prize in Life Sciences Awarded to Penn Medicine mRNA Pioneers Drew Weissman and Katalin Kariko. Press Release. Sep. 9, 2021: 3 pages. cited by applicant

[No Author Listed], 3rd International mRNA Health Conference Program. mRNA Conference. Nov. 11, 2015: 28 pages. cited by applicant

[No Author Listed], A Phase 3, Randomized, Stratified, Observer-Blind, Placebo-Controlled Study to Evaluate the Efficacy, Safety, and Immunogenicity of mRNA-1273 SARS-CoV-2 Vaccine in Adults Aged 18 Years and Older. ModernaTX, Inc. Protocol mRNA-1273-P301, Amendment 3. Aug. 20, 2020. 135 pages. cited by applicant

[No Author Listed], An Audience with Moncef Slaoui. *Nat Rev Drug Discov* Jul. 1, 2015;14:452-

453. <https://doi.org/10.1038/nrd4669>. cited by applicant
[No Author Listed], Background document on the mRNA-1273 vaccine (Moderna) against COVID-19. World Health Organization. Feb. 3, 2021. Accessed from <[https://www.who.int/publications/i/item/background-document-on-the-mrna-1273-vaccine-\(moderna\)-against-covid-19](https://www.who.int/publications/i/item/background-document-on-the-mrna-1273-vaccine-(moderna)-against-covid-19)>. 41 pages. cited by applicant
[No Author Listed], BioNTech Expands Clinical Oncology Portfolio with First Patient Dosed in Phase 2 Trial of mRNA-based Individualized Immunotherapy BNT122 in Colorectal Cancer Patients. Press Release. BioNTech. Oct. 1, 2021: 2 pages. cited by applicant
[No Author Listed], BioNTech Fourth Quarter and Full-Year 2019 Results. BioNTech. Mar. 31, 2020: 15 pages. cited by applicant
[No Author Listed], BioNTech SE Annual Report (Form 20-F) 179, F-12. United States Securities and Exchange Commission. Mar. 30, 2021: 739 pages. cited by applicant
[No Author Listed], BioNTech SE Annual Report (Form 20-F) 81. United States Securities and Exchange Commission. Mar. 30, 2022: 700 pages. cited by applicant
[No Author Listed], BioNTech Signs Collaboration Agreement with Pfizer to Develop mRNA-based Vaccines for Prevention of Influenza. Press Release. Aug. 16, 2018: 3 pages. cited by applicant
[No Author Listed], BNT162 COVID-19 Vaccine Program Update. BioNTech. Apr. 23, 2020: 15 pages. cited by applicant
[No Author Listed], Canada Exercises Increased Option for 20 Million Doses of mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Sep. 22, 2020. 2 pages. cited by applicant
[No Author Listed], CDC. SARS-COV-2 Variant Classifications and Definitions. <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-classifications.html>. Updated Sep. 1, 2023. 5 pages. cited by applicant
[No Author Listed], Center for Drug Evaluation and Research Approval Package for New Drug Application No. 50-718/S-50 for Doxil®. Approved Dec. 28, 2015. 97 pages. cited by applicant
[No Author Listed], Clinical trial NCT04528719: A Dose Escalation Study to Evaluate Safety, Reactogenicity, and Immunogenicity of mRNA-1345 in Healthy Adults and in Children Who Are Respiratory Syncytial Virus Seropositive (ModernaTX, Inc.) First Posted Aug. 27, 2020. Retrieved online on Mar. 15, 2021 at <https://www.clinicaltrials.gov/ct2/show/NCT04528719?term=NCT04528719&draw=2&rank=1>. cited by applicant
[No Author Listed], Comirnaty (mRNA-1273) EMA SmPC (summary of product characteristics);. Published Dec. 1, 2021. 276 pages. cited by applicant
[No Author Listed], Comirnaty Dosing Schedule for People 12 Years of Age & Older (2023-2024 Formula). Oct. 2023: 2 pages. cited by applicant
[No Author Listed], Comirnaty Prescribing Information. Jul. 2022: 27 pages. cited by applicant
[No Author Listed], Comirnaty Prescribing Information. Comirnaty. Oct. 2023: 36 pages. cited by applicant
[No Author Listed], Coronavirus (COVID-19) Update: FDA Authorizes Changes to Simplify Use of Bivalent mRNA COVID-19 Vaccines. FDA News Release. Apr. 18, 2023: 3 pages. cited by applicant
[No Author Listed], Coronavirus (COVID-19) Update: FDA Issues Policies to Guide Medical Product Developers Addressing Virus Variants. FDA News Release. Feb. 22, 2021. 5 pages. cited by applicant
[No Author Listed], Could COVID-19 vaccines be tweaked to cover new coronavirus variants? I Gavi, the Vaccine Alliance. Published Jan. 14, 2021.
URL:<https://www.gavi.org/vaccineswork/could-covid-19-vaccines-be-tweaked-cover-new-coronavirus-variants>: Last accessed Apr. 26, 2022. 5 Pages. cited by applicant
[No Author Listed], COVID-19 Vaccine Development Program. Pfizer. Jul. 1, 2020: 18 pages. cited

by applicant

[No Author Listed], DARPA Awards Moderna Therapeutics a Grant for up to \$25 Million to Develop Messenger RNA Therapeutics. News Release. Moderna. Oct. 2, 2013: 1 page. cited by applicant

[No Author Listed], Delivery means to avoid degradation of RNA preparations: Lipid nanoparticles (LNPs) for RNA delivery. CosmoBio. English-Language Machine Translation. Retrieved from the Internet: https://www.cosmobio.co.jp/product/detail/lipid-nanoparticles-for-rna-delivery-ecl.asp?entry_id=43081. Last Accessed: Oct. 23, 2023. 10 pages. cited by applicant

[No Author Listed], Dosing Schedule for Ages 12 Years of Age and Older with Pfizer-BioNTech COVID-19 Vaccine, Bivalent (Original and Omicron BA.4/BA.5). Apr. 2023:7-9. cited by applicant

[No Author Listed], European Commission Authorizes COVID 19 Vaccine Moderna in Europe. Jan. 6, 2021. Accessed from <https://www.drugs.com/clinical_trials/european-commission-authorizes-covid-19-vaccine-moderna-europe-19152.html>. 4 pages. cited by applicant

[No Author Listed], Experimental COVID-19 Vaccine Protects Upper and Lower Airways in Nonhuman Primates. NIH News Release. Jul. 28, 2020. 4 pages. cited by applicant

[No Author Listed], Experimental COVID-19 Vaccine Safe, Generates Immune Response. NIH News Release. Jul. 14, 2020. 4 pages. cited by applicant

[No Author Listed], Fact Sheet for Healthcare Providers Administering Vaccine: Emergency Use Authorization of Pfizer-BioNTech COVID-19 Vaccine (2023-2024 Formula) for 6 Months through 11 Years of Age. Pfizer-BioNTech. Sep. 2023: 59 pages. cited by applicant

[No Author Listed], Goldman Sachs Virtual 41st Annual Global Healthcare Conference. Pfizer Inc. Jun. 9, 2020: 13 pages. cited by applicant

[No Author Listed], History of Changes for Study: NCT04283461, Safety and Immunogenicity Study of 2019-nCoV Vaccine (mRNA-1273) for Prophylaxis of SARS-COV-2 Infection (COVID-19). Version submitted Dec. 8, 2022. 8 pages. cited by applicant

[No Author Listed], History of Changes for Study: NCT04283461, Safety and Immunogenicity Study of 2019-nCoV Vaccine (mRNA-1273) for Prophylaxis of SARS-COV-2 Infection (COVID-19). Version submitted Mar. 11, 2021. 10 pages. cited by applicant

[No Author Listed], History of Changes for Study: NCT04405076, Dose-Confirmation Study to Evaluate the Safety, Reactogenicity, and Immunogenicity of mRNA-1273 COVID-19 Vaccine in Adults Aged 18 Years and Older. Version submitted Dec. 6, 2022. 6 pages. cited by applicant

[No Author Listed], Increase in Manufacturing Capacity for COVID-19 Vaccines from Janssen, Moderna, and BioNTech/Pfizer. European Medicines Agency. Dec. 16, 2021: 4 pages. cited by applicant

[No Author Listed], Influenza A virus A/Jiangxi/IPB13/2013(H10N8) hemagglutinin protein. UniProtKB Accession: A0A059T4A1. Sep. 3, 2014. 1 page. cited by applicant

[No Author Listed], Katalin Kariko and Drew Weissman Awarded Horwitz Prize for Pioneering Research on COVID-19 Vaccines. Columbia University. Aug. 16, 2021: 12 pages. cited by applicant

[No Author Listed], Katalin Kariko Laureate de la Grande Medaille 2021 de L'Academie des Sciences. Press Release. De L'Academie des Sciences. Sep. 28, 2021: 2 pages. cited by applicant

[No Author Listed], Messenger RNA Pioneer Katalin Kariko Receives European Inventor Award 2022 for Lifetime Achievement. Press Release. European Patent Office. Jun. 21, 2022: 4 pages. cited by applicant

[No Author Listed], Moderna Advances Late-Stage Development of its Vaccine (mRNA-1273) Against COVID-19. Moderna Press Release. Jun. 11, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna and Lonza Announce Worldwide Strategic Collaboration to Manufacture Moderna's Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. May 1, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces Award from U.S. Government Agency BARDA for up to \$483 Million to Accelerate Development of mRNA Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. Apr. 16, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces Expansion of BARDA Agreement to Support Larger Phase 3 Program for Vaccine (mRNA-1273) Against COVID-19. Moderna Press Release. Jul. 26, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces FDA Authorization of Moderna COVID 19 Vaccine in U.S. Press Release. Dec. 19, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces First Participant Dosed in NIH-led Phase 1 Study of mRNA Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. Mar. 16, 2020. 2 pages. cited by applicant

[No Author Listed], Moderna Announces First Participant Dosed in Phase 2 Study of Omicron-Specific Booster Candidate and Publication of Data on Booster Durability Against Omicron Variant. News Release. Moderna. Jan. 26, 2022: 4 pages. cited by applicant

[No Author Listed], Moderna Announces First Participant Dosed in Phase 2 Study of Omicron-Specific Booster Candidate. News Release. Mar. 10, 2022: 3 pages. cited by applicant

[No Author Listed], Moderna Announces First Participants in Each Age Cohort Dosed in Phase 2 Study of mRNA Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. May 29, 2020. 2 pages. cited by applicant

[No Author Listed], Moderna Announces Funding Award from CEPI to Accelerate Development of Messenger RNA (mRNA) Vaccine Against Novel Coronavirus. News Release, Moderna. Jan. 23, 2020: 2 pages. cited by applicant

[No Author Listed], Moderna Announces IND Submitted to U.S. FDA for Phase 2 Study of mRNA Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. Apr. 27, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces it has Shipped Variant-Specific Vaccine Candidate, mRNA-1273.351, to NIH for Clinical Study. Moderna Press Release. Feb. 24, 2021. 3 pages. cited by applicant

[No Author Listed], Moderna Announces Phase 3 COVE Study of mRNA Vaccine Against COVID-19 (mRNA-1273) Begins. Moderna Press Release. Jul. 27, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces Positive Interim Phase 1 Data for its mRNA Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. May 18, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces Progress Across Broad Portfolio and all Three Clinical Stage Therapeutic Areas at 2020 R&D Day. Moderna Press Release. Sep. 17, 2020. 4 pages. cited by applicant

[No Author Listed], Moderna Announces Publication in The New England Journal of Medicine of Interim Results From Phase 1 Study of Its mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Jul. 14, 2020. 5 pages. cited by applicant

[No Author Listed], Moderna Announces Publication in The New England Journal of Medicine of Non-Human Primate Preclinical Viral Challenge Study of its mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Jul. 28, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Announces Strategy to Address Omicron (B.1.1.529) Sars-CoV-2 Variant. Press Release. Moderna. Nov. 26, 2021: 2 pages. cited by applicant

[No Author Listed], Moderna Announces Supply Agreement with U.S. Government for Initial 100 Million Doses of mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Aug. 11, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Completes Enrollment of Phase 2 Study of its mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Jul. 8, 2020. 2 pages. cited by

applicant

[No Author Listed], Moderna Confirms Advanced Discussions with European Commission to Supply Europe with 80 Million Doses of mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Aug. 24, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna Confirms Discussions with the Ministry of Health, Labour and Welfare to Supply Japan with 40 Million Doses of mRNA Vaccine Against COVID-19 (mRNA-1273). Moderna Press Release. Aug. 28, 2020. 2 pages. cited by applicant

[No Author Listed], Moderna has No. plans to share its COVID-19 vaccine recipe. Associated Press. Oct. 11, 2021:8 pages. cited by applicant

[No Author Listed], Moderna Inc to Discuss Updates on Respiratory Syncytial Virus (RSV) Vaccine Program Call. Edited Transcript. Refinitiv. Oct. 8, 2020: 14 pages. cited by applicant

[No Author Listed], Moderna Patents. <https://www.modernatx.com/en-US/patents> [last accessed Jul. 1, 2024]. cited by applicant

[No Author Listed], Moderna Provides Business Update and Announces Three New Development Programs in Infectious Disease Vaccines. Press Release. Jan. 11, 2021. 5 pages. cited by applicant

[No Author Listed], Moderna Receives FDA Authorization for Emergency Use of Omicron-Targeting Bivalent COVID-19 Booster Vaccine for Adults 18 Years and Older. News Release. Moderna. Aug. 31, 2022: 5 pages. cited by applicant

[No Author Listed], Moderna Receives FDA Fast Track Designation for mRNA Vaccine (mRNA-1273) Against Novel Coronavirus. Moderna Press Release. May 12, 2020. 2 pages. cited by applicant

[No Author Listed], Moderna Reports Fourth Quarter and Fiscal Year 2020 Financial Results and Provides Business Updates. Press Release. Feb. 25, 2021. 9 pages. cited by applicant

[No Author Listed], Moderna Ships mRNA Vaccine Against Novel Coronavirus (mRNA-1273) for Phase 1 Study. Moderna Press Release. Feb. 24, 2020. 2 pages. cited by applicant

[No Author Listed], Moderna's COVID-19 Vaccine Candidate Meets its Primary Efficacy Endpoint in the First Interim Analysis of the Phase 3 COVE Study. Press Release. Nov. 16, 2020. 3 pages. cited by applicant

[No Author Listed], Moderna's Update Patent Pledge. Press Release. Moderna. Mar. 7, 2022: 2 pages. cited by applicant

[No Author Listed], Modified mRNA Vaccines. 2021 Lasker-DeBakey Clinical Medical Research Award. 17 pages. cited by applicant

[No Author Listed], NIH Clinical Trial of Investigational Vaccine for COVID-19 begins. NIH News Release. Mar. 16, 2020. 3 pages. cited by applicant

[No Author Listed], NIH-Moderna Investigational COVID-19 Vaccine Shows Promise in Mouse Studies. NIH News Release. Aug. 5, 2020. 4 pages. cited by applicant

[No Author Listed], Penn mRNA Scientists Drew Weissman and Katalin Kariko Receive 2021 Lasker Award, America's Top Biomedical Research Prize. Penn News Release. Sep. 24, 2021: 6 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Announce Early Positive Data from an Ongoing Phase 1/2 Study of mRNA-based Vaccine Candidate against SARS-CoV-2. Press Release. Jul. 1, 2020: 8 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Announce Early Positive Update from German Phase 1/2 Study of COVID-19 Vaccine Study, Including First T Cell Response Data. Press Release. Jul. 20, 2020: 6 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Announce Phase 3 Trial Data Showing High Efficacy of a Booster Dose of Their COVID-19 Vaccine. Press Release. Pfizer. Oct. 21, 2021: 10 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Choose Lead mRNA Vaccine Candidate Against COVID-19 and Commence Pivotal Phase 2/3 Global Study. Press Release. Jul. 27, 2020: 7 pages. cited by

applicant

[No Author Listed], Pfizer and BioNTech Granted FDA Emergency Use Authorization of Omicron BA.4/BA.5-Adapted Bivalent COVID-19 Vaccine Booster for Ages 12 Years and Older. Pfizer.

Aug. 31, 2022: 12 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Receive Expanded U.S. Emergency Use Authorization for an Additional COVID-19 Vaccine Booster in Individuals Aged 50 Years and Older. Press Release. Pfizer. Mar. 29, 2022: 12 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Submit Applications to U.S. FDA for Omicron XBB.1.5-Adapted Monovalent COVID-19 Vaccine. Pfizer. Jun. 23, 2023: 8 pages. cited by applicant

[No Author Listed], Pfizer and BioNTech Submit Supplemental Biologics License Application for U.S. FDA Approval of Omicron BA.4/BA.5-Adapted Bivalent COVID-19 Vaccine for Ages 12 Years and Older as Primary Series or Booster. Pfizer. Feb. 24, 2023: 12 pages. cited by applicant

[No Author Listed], Pfizer Inc to discuss data from an ongoing phase 1/2 study of mRNA-based vaccine candidate against SARS_CoV-2 Call. Edited Transcript. Thomson Reuters. Jul. 1, 2020: 15 pages. cited by applicant

[No Author Listed], Pfizer reports fourth-quarter and full-year 2021 results. Pfizer. Feb. 8, 2022: 41 pages. cited by applicant

[No Author Listed], Pfizer-BioNTech COVID-19 Vaccine Information Sheet. 3 pages. cited by applicant

[No Author Listed], Phase 3 Clinical Trial of Investigational Vaccine for COVID-19 Begins. NIH News Release. Jul. 27, 2020. 3 pages. cited by applicant

[No Author Listed], Pioneering Scientists Awarded the Albany Prize, America's Most Distinguished Prize in Medicine, for Research Leading to Covid-19 Vaccines. The Albany Prize. Aug. 26, 2021: 4 pages. cited by applicant

[No Author Listed], Press release, Moderna, "Moderna Announces Publication in The New England Journal of Medicine of Interim Results From Older Adult Age Cohorts in Phase 1 Study of its mRNA Vaccine Against COVID-19 (mRNA-1273)", Sep. 29, 2020. 7 pages. cited by applicant

[No Author Listed], Programme of the 1st International mRNA Health Conference held in Germany, Oct. 2013, 32 pages. cited by applicant

[No Author Listed], Q2 2020 Earnings Call. BioNTech SE. Aug. 11, 2020:26 pages. cited by applicant

[No Author Listed], RBC capital Markets Global Healthcare Conference. Pfizer Inc. May 19, 2020: 9 pages. cited by applicant

[No Author Listed], Remarks by President Trump and Members of the Coronavirus Task Force in Meeting with Pharmaceutical Companies. The White House. Mar. 2, 2022: 31 pages. cited by applicant

[No Author Listed], SARS-CoV-2 variants of concern as of May 4, 2023. European Centre for Disease Prevention and Control. Last accessed May 16, 2023 at

<https://www.ecdc.europa.eu/en/covid-19/variants-concern>. 7 pages. cited by applicant

[No Author Listed], Severe acute respiratory syndrome coronavirus 2 isolate Wuhan-Hu-1, complete genome. GenBank Accession No. 045512, published on Jul. 18, 2020. 15 pages. cited by applicant

[No Author Listed], Severe acute respiratory syndrome coronavirus 2 isolate Wuhan-Hu-1, complete genome. GenBank Accession No. MN908947.3. Jan. 13, 2020. 15 pages. cited by applicant

[No Author Listed], spike glycoprotein [Middle East respiratory syndrome coronavirus]. GenBank Accession No. AGN52936.1. Published Jun. 10, 2013. 2 pages. cited by applicant

[No Author Listed], Spike_CVHSA. UniProt KB Accession No. P59594. Published Oct. 14, 2015. 1 page. cited by applicant

[No Author Listed], Spikevax (mRNA-1273) EMA SmPC (summary of product characteristics).

Published Jan. 20, 2021. 137 pages. cited by applicant

[No Author Listed], Statement by Moderna on Intellectual Property Matters during the COVID-19 Pandemic. Press Release. Moderna. Oct. 8, 2020: 1 page. cited by applicant

[No Author Listed], Statement from NIH and BARDA on the FDA Emergency Use Authorization of the Moderna COVID-19 Vaccines. COVID.gov. Dec. 18, 2020: 3 pages. cited by applicant

[No Author Listed], Two Pioneering RNA Scientists Win the 2021 Dr. Paul Janssen Award for Biomedical Research. Press Release. Johnson & Johnson. Sep. 28, 2021: 2 pages. cited by applicant

[No Author Listed], Vaccine information fact sheet for recipients and caregivers about Comirnaty (COVID-19 Vaccine, mRNA) and the Pfizer-BioNTech COVID-19 vaccine to prevent coronavirus disease 2019 (COVID-19) for use in individuals 12 years of age and older. Jul. 8, 2022: 9 pages. cited by applicant

[No Author Listed], WHO Drug Information, vol. 35, No. 2, 2021, 341 pages. cited by applicant

[No Author Listed], Who Drug Information. 2021;35(2):578-9. cited by applicant

[No Author Listed], World Health Organization. 2021. Classification of Omicron(B.1.1.529): SARS-CoV-2 Variant of Concern. Nov. 26, 2021. 3 pages. cited by applicant

[No Author Listed]. Reflection paper on the regulatory requirements for vaccines intended to provide protection against variant strain(s) of SARS-CoV-2. European Medicines Agency. EMA/117973/2021. Feb. 23, 2021. 7 pages. cited by applicant

Abbasi, COVID-19 and mRNA Vaccines-First Large Test for a New Approach. JAMA. Sep. 22, 2020;324(12):1125-1127. doi: 10.1001/jama.2020.16866. cited by applicant

Adney et al., Efficacy of an Adjuvanted Middle East Respiratory Syndrome Coronavirus Spike Protein Vaccine in Dromedary Camels and Alpacas. Viruses. Mar. 2, 2019;11(3). pii: E212. doi: 10.3390/v11030212. cited by applicant

Ahmad et al., New multivalent cationic lipids reveal bell curve for transfection efficiency versus membrane charge density: lipid-DNA complexes for gene delivery. J Gene Med. Jun. 2005;7(6):739-48. doi: 10.1002/jgm.717. cited by applicant

Ahmed et al., Biochemistry, Lipids, Statpearls Publishing (2023) (<https://www.ncbi.nlm.nih.gov/books/NBK525952/> [last accessed Aug. 24, 2023]: 5 pages. cited by applicant

Akinc et al., A combinatorial library of lipid-like materials for delivery of RNAi therapeutics. Nat Biotechnol. May 2008;26(5):561-9. doi: 10.1038/nbt1402. Epub Apr. 27, 2008. cited by applicant

Al Kahlout et al., Comparative Serological Study for the Prevalence of Anti-MERS Coronavirus Antibodies in High- and Low-Risk Groups in Qatar. J Immunol Res. Feb. 18, 2019;2019:1386740. doi: 10.1155/2019/1386740. eCollection 2019. cited by applicant

Altieri et al., The influence of 4-thiouridine labeling on pre-mRNA splicing outcomes. PLoS One. Dec. 13, 2021;16(12):e0257503. doi: 10.1371/journal.pone.0257503. eCollection 2021. cited by applicant

Anderson et al., Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults. N Engl J Med. Sep. 29, 2020;NEJMoa2028436: 1-12. doi: 10.1056/NEJMoa2028436. cited by applicant

Andries et al., N(1)-methylpseudouridine-incorporated mRNA outperforms pseudouridine-incorporated mRNA by providing enhanced protein expression and reduced immunogenicity in mammalian cell lines and mice. J Control Release. Nov. 10, 2015;217:337-44. doi: 10.1016/j.jconrel.2015.08.051. Epub Sep. 3, 2015. cited by applicant

Arthur, Pfizer predicts \$54bn in 2022 revenue from Comirnaty and Paxlovid. BioPharmaReporter. Feb. 8, 2022: 2 pages. cited by applicant

Atsmon et al., Safety and immunogenicity of multimeric-001—a novel universal influenza vaccine. J Clin Immunol. Jun. 2012;32(3):595-603. doi: 10.1007/s10875-011-9632-5. Epub Feb. 9, 2012. cited by applicant

Baden et al., Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med*. Feb. 4, 2021;384(5):403-416. doi: 10.1056/NEJMoa2035389. Epub Dec. 30, 2020. cited by applicant

Bancel, Our Global Commitment to Vaccine Access. Moderna. Oct. 8, 2021: 1 page. cited by applicant

Bangham et al., Diffusion of univalent ions across the lamellae of swollen phospholipids. *J Mol Biol*. Aug. 1965;13(1):238-52. doi: 10.1016/s0022-2836(65)80093-6. cited by applicant

Bear, Gene expression and regulation. In: Lewin's Cells. 2011. Cassimeris et al., Eds. Chapter 4:105-68. cited by applicant

Bennett et al., Cholesterol enhances cationic liposome-mediated DNA transfection of human respiratory epithelial cells. *Biosci Rep*. Feb. 1995;15(1):47-53. doi: 10.1007/BF01200214. cited by applicant

Bermudez, We decided not to enforce our patent during the pandemic: Noubar Afeyan, co-founder of Moderna. *BBC News World*. May 6, 2021: 11 pages. cited by applicant

Bharali et al., Nanoparticles and cancer therapy: a concise review with emphasis on dendrimers. *Int J Nanomedicine*. 2009;4:1-7. Epub Apr. 1, 2009. cited by applicant

Bhavana et al., COVID-19: Pathophysiology, treatment options, nanotechnology approaches, and research agenda to combating the SARS-CoV2 pandemic. *Life Sci*. Nov. 15, 2020;261:118336. doi: 10.1016/j.lfs.2020.118336. Epub Aug. 23, 2020. cited by applicant

Bourla, An Open Letter from Pfizer Chairman and CEO to Colleagues. Pfizer. May 7, 2021: 3 pages. cited by applicant

Bruxvoort et al., Real-world effectiveness of the mRNA-1273 vaccine against COVID-19: Interim results from a prospective observational cohort study. *Lancet Reg Health Am*. Feb. 2022;6:100134. doi: 10.1016/j.lana.2021.100134. Epub Nov. 25, 2021. cited by applicant

Buschmann et al., Nanomaterial Delivery Systems for mRNA Vaccines. *Vaccines (Basel)*. Jan. 19, 2021;9(1):65. doi: 10.3390/vaccines9010065. cited by applicant

Callaway, Omicron likely to weaken COVID vaccine protection. *Nature*. Dec. 2021;600(7889):367-368. doi: 10.1038/d41586-021-03672-3. cited by applicant

Carvalho et al., The first 12 months of COVID-19: a timeline of immunological insights. *Nat Rev Immunol*. Apr. 2021;21(4):245-256. doi:10.1038/s41577-021-00522-1. Epub Mar. 15, 2021. cited by applicant

Cele et al., SARS-COV-2 Omicron has extensive but incomplete escape of Pfizer BNT162b2 elicited neutralization and requires ACE2 for infection. *medRxiv*. 2021 doi: 10.1101/2021.12.08.21267417. 2021.12.08.21267417. 15 pages. cited by applicant

Cerutti et al., Potent SARS-CoV-2 neutralizing antibodies directed against spike N-terminal domain target a single supersite. *Cell Host Microbe*. May 12, 2021;29(5):819-833.e7. doi: 10.1016/j.chom.2021.03.005. Epub Mar. 12, 2021. cited by applicant

Chan et al., Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect*. Jan. 28, 2020;9(1):221-236. doi: 10.1080/22221751.2020.1719902. eCollection 2020. cited by applicant

Chen et al., A novel neutralizing monoclonal antibody targeting the N-terminal domain of the MERS-CoV spike protein. *Emerg Microbes Infect*. Jun. 28, 2017;6(6):e60. doi: 10.1038/emi.2017.50. Erratum for: *Emerg Microbes Infect*. May 24, 2017;6(5):e37. cited by applicant

Chen et al., Emerging coronaviruses: Genome structure, replication, and pathogenesis. *J Med Virol*. Apr. 2020;92(4):418-423. doi: 10.1002/jmv.25681. Epub Feb. 7, 2020. Erratum in: *J Med Virol*. Oct. 2020;92(10):2249. cited by applicant

Chen et al., Vaccine design of hemagglutinin glycoprotein against influenza. *Trends Biotechnol*. Sep. 2011;29(9):426-34. doi: 10.1016/j.tibtech.2011.04.007. cited by applicant

Cheng et al., Selective ORgan Targeting (SORT) nanoparticles for tissue specific mRNA delivery and CRISPR/Cas gene editing. *Nat Nanotechnol*. Apr. 2020; 15(4): 313-320. cited by applicant

Chi et al., A neutralizing human antibody binds to the N-terminal domain of the Spike protein of SARS-CoV-2. *Science*. Aug. 7, 2020;369(6504):650-655. doi: 10.1126/science.abc6952. Epub Jun. 22, 2020. cited by applicant

Choi et al., 2'-O-methylation in mRNA disrupts tRNA decoding during translation elongation. *Nat Struct Mol Biol*. Mar. 2018;25(3):208-216. doi: 10.1038/s41594-018-0030-z. Epub Feb. 19, 2018. cited by applicant

Choi et al., Safety and immunogenicity of SARS-CoV-2 variant mRNA vaccine boosters in healthy adults: an interim analysis. *Nat Med*. Nov. 2021;27(11):2025-2031. doi: 10.1038/s41591-021-01527-y. Epub Sep. 15, 2021. cited by applicant

Choi et al., Serum Neutralizing Activity of mRNA-1273 against SARS-CoV-2 Variants, *J Virol*. Nov. 9, 2021;95(23):e0131321. doi: 10.1128/JVI.01313-21. Epub Sep. 22, 2021. cited by applicant

Chu et al., Immune Memory Response After a Booster Injection of mRNA-1273 for Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2). 2021 MedRxiv doi: <https://doi.org/10.1101/2021.09.29.21264089>. 52 pages. cited by applicant

Cohen, New crop of mRNA vaccines aim for accessibility. *Science*. Apr. 8, 2022;376(6589):120-121. doi: 10.1126/science.abq3935. Epub Apr. 7, 2022. cited by applicant

Cohen, Scientists are moving at record speed to create new coronavirus vaccines-but they may come too late. *Science Insider*. Jan. 27, 2020. 6 pages. cited by applicant

Cohen, Scientists question Moderna invention claim in COVID-19 vaccine dispute. *Science Insider*. Aug. 29, 2022: 3 pages. cited by applicant

Cooper et al., Chapters 1, 7, and 8 in *The Cell: a Molecular Approach*, 5th Ed. (2009), 163 pages. cited by applicant

Corbett et al., Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in Nonhuman Primates. *N Engl J Med* 2020; 383:1544-55. doi: 10.1056/NEJMoa2024671. cited by applicant

Corbett et al., SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen preparedness. *Nature*. Oct. 2020;586(7830):567-571. doi: 10.1038/s41586-020-2622-0. Epub Aug. 5, 2020. cited by applicant

Cox et al., FluBlok, a recombinant hemagglutinin influenza vaccine. *Influenza Other Respir Viruses*. Nov. 2008; 2(6): 211-219. cited by applicant

Dai et al., Viral targets for vaccines against COVID-19. *Nat Rev Immunol*. Feb. 2021;21(2):73-82. doi: 10.1038/s41577-020-00480-0. Epub Dec. 18, 2020. cited by applicant

Defrancesco, The 'anti-hype' vaccine. *Nat Biotechnol*. Mar. 2017;35(3):193-197. doi: 10.1038/nbt.3812. Epub Feb. 27, 2017. cited by applicant

Degroot et al., Part II—The Positive Sense Single Stranded RNA Viruses, Family Coronaviridae, in *Virus Taxonomy: Ninth Report of the International Committee on Taxonomy of Viruses* 806 (2012) 25 pages. cited by applicant

Dejnirattisai et al., Reduced neutralisation of SARS-CoV-2 omicron B.1.1.529 variant by post-immunisation serum. *Lancet*. Jan. 15, 2022;399(10321):234-236. doi: 10.1016/S0140-6736(21)02844-0. Epub Dec. 20, 2021. cited by applicant

Dimitriadis et al., Translation of rabbit globin mRNA introduced by liposomes into mouse lymphocytes. *Nature*. Aug. 31, 1978;274(5674):923-4. doi: 10.1038/274923a0. cited by applicant

Dolgin, Business: The billion-dollar biotech. *Nature*. Jun. 4, 2015;522(7554):26-8. doi: 10.1038/522026a. cited by applicant

Dormitzer, Synthetic Vaccinology at Novartis, Presentation at The National Academy of Sciences Forum on Synthetic Biology. Oct. 21, 2013, The Keck Center, Washington, DC. 12 pages. cited by applicant

Du et al., The spike protein of SARS-CoV—a target for vaccine and therapeutic development. *Nat Rev Microbiol*. Mar. 2009;7(3):226-36. doi: 10.1038/nrmicro2090. Epub Feb. 9, 2009. cited by applicant

Dufourc, Sterols and membrane dynamics. *J Chem Biol*. Nov. 2008;1(1-4):63-77. doi:

10.1007/s12154-008-0010-6. Epub Sep. 23, 2008. cited by applicant

El Sahly et al., Efficacy of the mRNA-1273 SARS-CoV-2 Vaccine at Completion of Blinded Phase. *N Engl J Med*. Nov. 4, 2021; 385:1774-1785. cited by applicant

Ellebedy et al., Influenza Vaccines. *Vaccine*. Nov. 5, 2009; 27(Suppl 4): D65-D68. cited by applicant

Engler et al., Half- vs full-dose trivalent inactivated influenza vaccine (2004-2005): age, dose, and sex effects on immune responses. *Arch Intern Med*. Dec. 8, 2008;168(22):2405-14. doi: 10.1001/archinternmed.2008.513. cited by applicant

Erasmus et al., A Nanostructured Lipid Carrier for Delivery of a Replicating Viral RNA Provides Single, Low-Dose Protection against Zika. *Mol Ther*. Oct. 3, 2018;26(10):2507-2522. doi: 10.1016/j.ymthe.2018.07.010. Epub Aug. 2, 2018. cited by applicant

Escriou et al., Protection from SARS coronavirus conferred by live measles vaccine expressing the spike glycoprotein. *Virology*. Mar. 2014;452-453:32-41. doi: 10.1016/j.virol.2014.01.002. Epub Jan. 28, 2014. cited by applicant

Fauquet, Taxonomy, Classification and Nomenclature of Viruses. *Encyclopedia of Virology*. 2008:9-23. doi: 10.1016/B978-012374410-4.00509-4. Epub Jul. 30, 2008. cited by applicant

Fujimori, How this Once Little-Known Molecule is Disrupting Medicine. *Pfizer*. Jun. 2022: 7 pages. cited by applicant

Furuichi et al., Viral and cellular mRNA capping: past and prospects. *Adv Virus Res*. 2000;55:135-84. doi: 10.1016/s0065-3527(00)55003-9. cited by applicant

Gallie, D.R., The cap and poly(A) tail function synergistically to regulate mRNA translational efficiency. *Genes Dev*. Nov. 1991;5(11):2108-16. cited by applicant

Garde et al., The story of mRNA: How a once-dismissed idea became a leading technology in the Covid vaccine race. *STAT News*. Nov. 10, 2020: 23 pages. cited by applicant

Geall et al., Using self-amplifying mRNA vaccines to facilitate a rapid response to pandemic influenza. *Eur Pharm Rev*. Jul. 3, 2014; 3. 5 pages. cited by applicant

Gilboa, E. et al., Cancer immunotherapy with mRNA-transfected dendritic cells. *Immunol Rev*. Jun. 2004;199:251-63. cited by applicant

Guevara et al., Advances in Lipid Nanoparticles for mRNA-Based Cancer Immunotherapy. *Front Chem*. 2020; 8: 589959. cited by applicant

Guo et al., An Engineered Receptor-Binding Domain Improves the Immunogenicity of Multivalent SARS-CoV-2 Vaccines. *mBio*. May-Jun. 2021; 12(3): e00930-21. cited by applicant

Guo et al., Nanoparticles escaping RES and endosome: Challenges for siRNA delivery for cancer therapy. *J Nanomat*. Jan. 2011;742895:1-12. cited by applicant

Hashiba et al., The use of design of experiments with multiple responses to determine optimal formulations for in vivo hepatic mRNA delivery. *J Control Release*. Nov. 10, 2020;327:467-476. doi: 10.1016/j.jconrel.2020.08.031. Epub Aug. 25, 2020. cited by applicant

Hatziantoniou et al., Anaphylactic reactions to mRNA COVID-19 vaccines: A call for further study. *Vaccine*. May 6, 2021;39(19):2605-2607. doi: 10.1016/j.vaccine.2021.03.073. Epub Apr. 6, 2021. PMID: 33846043; PMCID: PMC8023205. cited by applicant

He et al., Receptor-binding domain of SARS-CoV spike protein induces highly potent neutralizing antibodies: implication for developing subunit vaccine. *Biochem Biophys Res Commun*. Nov. 12, 2004;324(2):773-81. doi: 10.1016/j.bbrc.2004.09.106. cited by applicant

He et al., Single-component, self-assembling, protein nanoparticles presenting the receptor binding domain and stabilized spike as SARS-CoV-2 vaccine candidates. *Sci Adv*. Mar. 19, 2021;7(12):eabf1591. doi: 10.1126/sciadv.abf1591. Print Mar. 2021. cited by applicant

Heinz et al., Distinguishing features of current COVID-19 vaccines: knowns and unknowns of antigen presentation and modes of action. *NPJ Vaccines*. Aug. 16, 2021;6(1):104. doi: 10.1038/s41541-021-00369-6. cited by applicant

Hemmer et al., Minimal peptide length requirements for CD4(+) T cell clones—implications for

molecular mimicry and T cell survival. *Int Immunol.* Mar. 2000;12(3):375-83. doi: 10.1093/intimm/12.3.375. cited by applicant

Hoernes et al., Nucleotide modifications within bacterial messenger RNAs regulate their translation and are able to rewire the genetic code. *Nucleic Acids Res.* Jan. 29, 2016; 44(2): 852- 862. cited by applicant

Hope et al., Reduction of Liposome Size and Preparation of Unilamellar Vesicles by Extrusion Techniques. In: *Liposome Technology*. 1993. Gregoriadis et al., Ed. vol. 1:123. cited by applicant

Houser et al., Influenza Vaccines: Challenges and Solutions. *Cell Host Microbe.* Mar. 11, 2015; 17(3): 295-300. doi: 10.1016/j.chom.2015.02.012. cited by applicant

Hsieh et al., Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science.* Sep. 18, 2020;369(6510):1501-1505. doi: 10.1126/science.abd0826. Epub Jul. 23, 2020. cited by applicant

Huang et al., Asymmetric 1-alkyl-2-acyl phosphatidylcholine: a helper lipid for enhanced non-viral gene delivery. *Int J Pharm.* May 1, 2012;427(1):64-70. doi: 10.1016/j.ijpharm.2011.06.022. Epub Jun. 21, 2011. cited by applicant

Huang et al., Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* Jan. 24, 2020;395(10223)497-506. cited by applicant

Huang et al., Construction and biological characterisation of recombinant porcine circovirus type 2 expressing the V5 epitope tag. *Virus Res.* Nov. 2011; 161(2):115-23. doi: 10.1016/j.virusres.2011.05.015. Epub May 27, 2011. cited by applicant

Jackson et al., An mRNA Vaccine against SARS-CoV-2—Preliminary Report. *N Engl J Med.* Jul. 14, 2020; NEJMoa2022483. doi: 10.1056/NEJMoa2022483. Online ahead of print. cited by applicant

Javier et al., Phylogenetic Analysis and Structural Modeling of SARS-CoV-2 Spike Protein Reveals an Evolutionary Distinct and Proteolytically Sensitive Activation Loop. *J Mol Biol.* May 1, 2020;432(10):3309-3325. doi: 10.1016/j.jmb.2020.04.009. Epub Apr. 19, 2020. cited by applicant

Jayaraman et al., Maximizing the potency of siRNA lipid nanoparticles for hepatic gene silencing in vivo. *Angew Chem Int Ed Engl.* Aug. 20, 2012;51(34):8529-33. doi: 10.1002/anie.201203263. cited by applicant

Jiaming et al., The recombinant N-terminal domain of spike proteins is a potential vaccine against Middle East respiratory syndrome coronavirus (MERS-CoV) infection. *Vaccine.* Jan. 3, 2017;35(1):10-18. doi: 10.1016/j.vaccine.2016.11.064. Epub Nov. 26, 2016. cited by applicant

Jiang et al., The Variation of SARS-CoV-2 and Advanced Research on Current Vaccines. *Front Med (Lausanne).* Jan. 18, 2022;8:806641. doi: 10.3389/fmed.2021.806641. eCollection 2021. cited by applicant

Johanning et al., A Sindbis virus mRNA polynucleotide vector achieves prolonged and high level heterologous gene expression in vivo. *Nucleic Acids Res.* May 11, 1995; 23(9): 1495-1501. cited by applicant

Kanasty et al., Delivery materials for siRNA therapeutics. *Nat Mater.* Nov. 2013;12(11):967-77. doi: 10.1038/nmat3765. cited by applicant

Kariko et al., Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Res.* Nov. 2011;39(21):e142. doi: 10.1093/nar/gkr695. Epub Sep. 2, 2011. cited by applicant

Kariko et al., Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA. *Immunity.* Aug. 2005;23(2):165-75. doi: 10.1016/j.immuni.2005.06.008. cited by applicant

Kauffman et al., Materials for non-viral intracellular delivery of messenger RNA therapeutics. *J Control Release.* Oct. 28, 2016;240:227-234. doi: 10.1016/j.jconrel.2015.12.032. Epub Dec. 21, 2015. cited by applicant

Keegan, The Lasker Awards: A History of “America's Nobels”. *Cell Mentor.* Sep. 11, 2018: 7 pages. cited by applicant

Khoury et al., Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat Med.* Jul. 2021;27(7):1205-1211. doi: 10.1038/s41591-021-01377-8. Epub May 17, 2021. cited by applicant

King et al., *Virus Taxonomy: Classification and Nomenclature of Viruses*. Ninth Report of the International Committee on Taxonomy of Viruses. Int Union of Microbiol Soc. 2012: 50 pages. cited by applicant

Kirchdoerfer et al., Stabilized coronavirus spikes are resistant to conformational changes induced by receptor recognition or proteolysis. *Sci Rep.* Oct. 24, 2018;8(1):15701. doi: 10.1038/s41598-018-34171-7. cited by applicant

Kisich et al., Antimycobacterial agent based on mRNA encoding human beta-defensin 2 enables primary macrophages to restrict growth of *Mycobacterium tuberculosis*. *Infect Immun.* Apr. 2001;69(4):2692-9. cited by applicant

Kolata, Kati Kariko Helped Shield the World from the Coronavirus. *New York Times*. 2021 Sep. 24: 5 pages. cited by applicant

Koyama et al., Analysis on GENIE reveals novel recurrent variants that affect molecular diagnosis of sizable number of cancer patients. *BMC Cancer.* Feb. 1, 2019;19(1):114. doi: 10.1186/s12885-019-5313-1. cited by applicant

Koyama et al., Emergence of Drift Variants That May Affect COVID-19 Vaccine Development and Antibody Treatment. *Pathogens.* Apr. 26, 2020;9(5):324. doi: 10.3390/pathogens9050324. cited by applicant

Krammer et al., Chimeric hemagglutinin influenza virus vaccine constructs elicit broadly protective stalk-specific antibodies. *J Virol.* Jun. 2013;87(12):6542-50. doi: 10.1128/JVI.00641-13. Epub Apr. 10, 2013. cited by applicant

Kresge, The Messenger RNA Pioneers Everyone Ignored. *Bloomberg*. Nov. 23, 2021: 4 pages. cited by applicant

Kurimoto et al., PEG-OligoRNA Hybridization of mRNA for Developing Sterically Stable Lipid Nanoparticles toward In Vivo Administration. *Molecules.* Apr. 3, 2019;24(7): 1303. cited by applicant

Leung et al., Lipid nanoparticles for short interfering RNA delivery. *Adv Genet.* 2014;88:71-110. doi: 10.1016/B978-0-12-800148-6.00004-3. cited by applicant

Leuschner et al., Therapeutic siRNA silencing in inflammatory monocytes in mice. *Nat Biotechnol.* Oct. 9, 2011;29(11):1005-10. doi: 10.1038/nbt.1989. cited by applicant

Li et al., Application of ultra-high performance liquid chromatography for chemical characterization of liposome-based therapeutic small-interfering RNA, *13 American Pharmaceutical Review* 102 (2010). 13 pages. cited by applicant

Li et al., Cell Attachment Domains of the Porcine Epidemic Diarrhea Virus Spike Protein Are Key Targets of Neutralizing Antibodies. *J Virol.* May 26, 2017;91(12):e00273-17. doi: 10.1128/JVI.00273-17. Print Jun. 15, 2017. cited by applicant

Li, Receptor recognition mechanisms of coronaviruses: a decade of structural studies. *J Virol.* Feb. 2015;89(4):1954-64. doi: 10.1128/JVI.02615-14. Epub Nov. 26, 2014. cited by applicant

Li, Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol.* Sep. 29, 2016;3(1):237-261. doi: 10.1146/annurev-virology-110615-042301. Epub Aug. 25, 2016. cited by applicant

Lin et al., Glycan Masking of Epitopes in the NTD and RBD of the Spike Protein Elicits Broadly Neutralizing Antibodies Against SARS-CoV-2 Variants. *Front Immunol.* Dec. 2, 2021;12:795741. doi: 10.3389/fimmu.2021.795741. eCollection 2021. cited by applicant

Lin et al., Three-dimensional imaging of lipid gene-carriers: membrane charge density controls universal transfection behavior in lamellar cationic liposome-DNA complexes. *Biophys J.* May 2003;84(5):3307-16. doi: 10.1016/S0006-3495(03)70055-1. cited by applicant

Liu et al., Chimeric severe acute respiratory syndrome coronavirus (Sars-CoV) S glycoprotein and

influenza matrix 1 efficiently form virus-like particles (VLPs) that protect mice against challenge with SARS-CoV. *Vaccine*. Sep. 2, 2011;29(38):6606-13. doi: 10.1016/j.vaccine.2011.06.111. Epub Jul. 14, 2011. cited by applicant

Liu et al., Influence of polyethylene glycol density and surface lipid on pharmacokinetics and biodistribution of lipid-calcium-phosphate nanoparticles. *Biomaterials*. Mar. 2014;35(9):3027-34. doi: 10.1016/j.biomaterials.2013.12.022. Epub Jan. 2, 2014. cited by applicant

Liu et al., Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike. *Nature*. Aug. 2020;584(7821):450-456. doi: 10.1038/s41586-020-2571-7. Epub Jul. 22, 2020. cited by applicant

Loftus, Moderna Signals It May Enforce Covid-19 Vaccine Patents in Wealthy Nations. *Wall Street Journal*. Mar. 7, 2022. <https://www.wsj.com/articles/moderna-signals-it-may-enforce-covid-19-vaccine-patents-in-wealthy-nations-11646699609> [last accessed Jul. 1, 2024]. cited by applicant

Lu et al., Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet*. Feb. 22, 2020;395(10224):565-574. doi: 10.1016/S0140-6736(20)30251-8. Epub Jan. 30, 2020. cited by applicant

Lutz et al., Unmodified mRNA in LNPs constitutes a competitive technology for prophylactic vaccines. *NPJ Vaccines*. Oct. 19, 2017;2:29. doi: 10.1038/s41541-017-0032-6. eCollection 2017. cited by applicant

Ma et al., Intranasal vaccination with recombinant receptor-binding domain of MERS-CoV spike protein induces much stronger local mucosal immune responses than subcutaneous immunization: Implication for designing novel mucosal MERS vaccines. *Vaccine*. Apr. 11, 2014;32(18):2100-8. doi: 10.1016/j.vaccine.2014.02.004. Epub Feb. 18, 2014. cited by applicant

Mandal et al., Reprogramming human fibroblasts to pluripotency using modified mRNA. *Nat Protoc*. Mar. 2013;8(3):568-82. doi: 10.1038/nprot.2013.019. Epub Feb. 21, 2013. cited by applicant

Martinez-Flores et al., SARS-CoV-2 Vaccines Based on the Spike Glycoprotein and Implications of New Viral Variants. *Front Immunol*. Jul. 12, 2021;12:701501. doi: 10.3389/fimmu.2021.701501. eCollection 2021. cited by applicant

McCallum et al., N-terminal domain antigenic mapping reveals a site of vulnerability for SARS-CoV-2. *Cell*. Apr. 29, 2021; 184(9): 2332-2347.e16. cited by applicant

McKay et al., Self-amplifying Rna SARS-CoV-2 lipid nanoparticle vaccine candidate induces high neutralizing antibody titers in mice. *Nat Commun*. Jul. 9, 2020;11(1):3523. doi: 10.1038/s41467-020-17409-9. cited by applicant

McLellan et al., Structure and function of respiratory syncytial virus surface glycoproteins. *Curr Top Microbiol Immunol*. 2013;372:83-104. doi: 10.1007/978-3-642-38919-1_4. cited by applicant

McLellan et al., Structure of respiratory syncytial virus fusion glycoprotein in the postfusion conformation reveals preservation of neutralizing epitopes. *J Virol*. Aug. 2011;85(15):7788-96. doi: 10.1128/JVI.00555-11. Epub May 25, 2011. cited by applicant

Mendonca et al., Design of lipid-based nanoparticles for delivery of therapeutic nucleic acids. *Drug Discov Today*. Mar. 2023;28(3):103505. doi: 10.1016/j.drudis.2023.103505. Epub Jan. 25, 2023. cited by applicant

Miao et al., Delivery of mRNA vaccines with heterocyclic lipids increases anti-tumor efficacy by STING-mediated immune cell activation. *Nat Biotechnol* 37, 1174-1185 (2019). <https://doi.org/10.1038/s41587-019-0247-3>. cited by applicant

Modernatx, Inc et al.v. Pfizer Inc et al. (D. Mass. 22-11378-RGS) D.I. 105—District Court Memorandum and Order on Claim Construction, Aug. 1, 2023. 15 pages. cited by applicant

Modernatx, Inc et al.v. Pfizer Inc et al. (D. Mass. 22-11378-RGS) D.I. 1—Complaint for Patent Infringement, Aug. 26, 2022. 39 pages. cited by applicant

Modernatx, Inc et al.v. Pfizer Inc et al. (D. Mass. 22-11378-RGS) D.I. 45—Answer to the Complaint, Dec. 5, 2022. 81 pages. cited by applicant

Modjarrad., Mers-CoV vaccine candidates in development: The current landscape. *Vaccine*. Jun. 3, 2016;34(26):2982-2987. doi: 10.1016/j.vaccine.2016.03.104. Epub Apr. 12, 2016. cited by applicant

Mullard, COVID-19 vaccine success enables a bolder vision for mRNA cancer vaccines, says BioNTech CEO. *Nat Rev Drug Discov*. Jul. 2021;20(7):500-501. doi: 10.1038/d41573-021-00110-x. cited by applicant

Mulligan et al., Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. *Nature*. Oct. 2020;586(7830):589-593. doi: 10.1038/s41586-020-2639-4. Epub Aug. 12, 2020. cited by applicant

Murphy, Excerpts from Janeway's Immunobiology, 8th Ed. (2012). 250 pages. cited by applicant

Muthumani et al., A synthetic consensus anti-spike protein DNA vaccine induces protective immunity against Middle East respiratory syndrome coronavirus in nonhuman primates. *Sci Transl Med*. Aug. 19, 2015;7(301):301ra132. doi: 10.1126/scitranslmed.aac7462. cited by applicant

Nance et al., Modifications in an Emergency: The Role of N1-Methylpseudouridine in COVID-19 Vaccines. *ACS Cent Sci*. May 26, 2021;7(5):748-756. doi: 10.1021/acscentsci.1c00197. Epub Apr. 6, 2021. cited by applicant

Oostvogels et al., Phase 1 Assessment of the Safety and Immunogenicity of an mRNA-Lipid Nanoparticle Vaccine Candidate Against SARS-CoV-2 in Human Volunteers. *medRxiv* Nov. 9, 2020. 20228551;1-38. doi: <https://doi.org/10.1101/2020.11.09.20228551>. cited by applicant

Opposition against European Patent No. 3324979 B1, filed by BioNTech SE (Opponent) against ModernaTX, Inc. (Patentee) on Jul. 10, 2023. 86 pages. cited by applicant

Ostro et al., Evidence for translation of rabbit globin mRNA after liposome-mediated insertion into a human cell line. *Nature*. Aug. 31, 1978;274(5674):921-3. doi: 10.1038/274921a0. cited by applicant

Pallesen et al., Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc Natl Acad Sci U S A*. Aug. 29, 2017;114(35):E7348-E7357. doi: 10.1073/pnas.1707304114. Epub Aug. 14, 2017. cited by applicant

Pardi et al., Nucleoside-modified mRNA vaccines induce potent T follicular helper and germinal center B cell responses. *J Exp Med*. Jun. 4, 2018;215(6):1571-1588. doi: 10.1084/jem.20171450. Epub May 8, 2018. cited by applicant

Pardi et al., Zika virus protection by a single low-dose nucleoside-modified mRNA vaccination. *Nature*. Mar. 9, 2017;543(7644):248-251. doi: 10.1038/nature21428. Epub Feb. 2, 2017. cited by applicant

Park et al., The Miracle Workers. Heroes of the Year. *TIME Magazine*. 2021: 17 pages. cited by applicant

Pascolo., Messenger RNA-based vaccines. *Expert Opin Biol Ther*. Aug. 2004;4(8):1285-94. doi: 10.1517/14712598.4.8.1285. cited by applicant

Patel et al., Naturally-occurring cholesterol analogues in lipid nanoparticles induce polymorphic shape and enhance intracellular delivery of mRNA. *Nat Commun*. Feb. 20, 2020;11(1):983. doi: 10.1038/s41467-020-14527-2. cited by applicant

Pathak et al., Intra-host variability in global SARS-CoV-2 genomes as signatures of RNA editing: implications in viral and host response outcomes. *bioRxiv* Dec. 9, 2020; 2020.12.09.417519; doi: <https://doi.org/10.1101/2020.12.09.417519>. 32 pages. cited by applicant

Peabody et al., Termination-reinitiation occurs in the translation of mammalian cell mRNAs. *Mol Cell Biol*. Jul. 1986;6(7):2695-703. doi: 10.1128/mcb.6.7.2695-2703.1986. cited by applicant

Peek et al., Nanotechnology in vaccine delivery. *Adv Drug Deliv Rev*. May 22, 2008;60(8):915-28. doi: 10.1016/j.addr.2007.05.017. Epub Feb. 7, 2008. cited by applicant

Pegu et al., Durability of mRNA-1273 vaccine-induced antibodies against SARS-CoV-2 variants. *Science*. Sep. 17, 2021;373(6561):1372-1377. doi: 10.1126/science.abj4176. Epub Aug. 13, 2021. cited by applicant

Probst et al., Characterization of the ribonuclease activity on the skin surface. *Genet Vaccines Ther*.

May 29, 2006;4:4. doi:10.1186/1479-0556-4-4. cited by applicant

Pulliam et al., Increased risk of SARS-CoV-2 reinfection associated with emergence of the Omicron variant in South Africa. medRxiv 2021. doi:10.1101/2021.11.11.21266068:2021.11.11.21266068. 42 pages. cited by applicant

Raj et al., Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. *Nature*. Mar. 14, 2013;495(7440):251-4. doi: 10.1038/nature12005. cited by applicant

Regalado, Don't panic about the latest coronavirus mutations say drug companies. MIT Technology Review. Dec. 23, 2020. 5 pages. cited by applicant

Regalado, Moderna believes it could update its coronavirus vaccine without a big new trial. MIT Technology Review. Jan. 13, 2021. 4 pages. cited by applicant

Robbins et al., Moderna Backs down in its Vaccine Patent Fight with the NIH. *New York Times*. Dec. 17, 2021: 1 page. cited by applicant

Rodriguez-Gascon et al., Development of nucleic acid vaccines: use of self-amplifying RNA in lipid nanoparticles. *Int J Nanomedicine*. 2014; 9: 1833-1843. cited by applicant

Rossi et al., IRES-based Venezuelan equine encephalitis vaccine candidate elicits protective immunity in mice. *Virology*. Mar. 15, 2013;437(2):81-8. doi: 10.1016/j.virol.2012.11.013. Epub Jan. 22, 2013. cited by applicant

Sahin et al., Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer. *Nature*. Jul. 13, 2017;547(7662):222-226. doi: 10.1038/nature23003. Epub Jul. 5, 2017. cited by applicant

Salomon et al., A liposomal RNA vaccine inducing neoantigen-specific CD4+ T cells augments the antitumor activity of local radiotherapy in mice. *Oncoimmunology*. Jun. 22, 2020;9(1):1771925. doi: 10.1080/2162402X.2020.1771925. cited by applicant

Santoro et al., An efficient system for the evolution of aminoacyl-tRNA synthetase specificity. *Nat Biotechnol*. Oct. 2002;20(10):1044-8. doi: 10.1038/nbt742. Epub Sep. 16, 2002. cited by applicant

Sarkar et al., S glycoprotein diversity of the Omicron variant. medRxiv 2021. doi:10.1101/2021.12.04.21267284:2021.12.04.21267284. 30 pages. cited by applicant

Schoenmaker et al., mRNA-lipid nanoparticle COVID-19 vaccines: Structure and stability. *Int J Pharm*. May 15, 2021;601:120586. doi: 10.1016/j.ijpharm.2021.120586. Epub Apr. 9, 2021. cited by applicant

Schumann et al., Multiple links between 5-methylcytosine content of mRNA and translation. *BMC Biol*. Apr. 15, 2020;18(1):40. doi: 10.1186/s12915-020-00769-5. cited by applicant

Schuurhuis et al., Ins and outs of dendritic cells. *Int Arch Allergy Immunol*. 2006;140(1):53-72. doi: 10.1159/000092002. Epub Mar. 13, 2006. cited by applicant

Shidhadye et al., Solid lipid nanoparticles and nanostructured lipid carriers—innovative generations of solid lipid carriers. *Curr Drug Deliv*. Oct. 2008;5(4):324-31. doi: 10.2174/156720108785915087. cited by applicant

Shim et al. Intranasal immunization with plasmid DNA encoding spike protein of SARS-coronavirus/polyethylenimine nanoparticles elicits antigen-specific humoral and cellular immune responses. *BMC Immunol*. Dec. 31, 2010;11:65. doi: 10.1186/1471-2172-11-65. cited by applicant

Skidmore et al., Genomic Sequencing of SARS-CoV-2 E484K Variant B.1.243.1, Arizona, USA. *Emerg Infect Dis*. Oct. 2021;27(10):2718-2720. doi: 10.3201/eid2710.211189. cited by applicant

Stephenson et al., Cross-reactivity to highly pathogenic avian influenza H5N1 viruses after vaccination with nonadjuvanted and MF59-adjuvanted influenza A/Duck/Singapore/97 (H5N3) vaccine: a potential priming strategy. *J Infect Dis*. Apr. 15, 2005;191(8):1210-5. doi: 10.1086/428948. Epub Mar. 14, 2005. cited by applicant

Tegally et al., Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. medRxiv. Dec. 22, 2020; Preprint: 19 pages. doi: 10.1101/2020.12.21.20248640. Epub Dec. 22, 2020. cited by applicant

Tews et al., Chapter 2: Self-Replicating RNA. RNA Vaccines. 2017; 1499: 15-35. Published online Jun. 11, 2016. doi: 10.1007/978-1-4939-6481-9_2. cited by applicant

Thery et al., The cell biology of antigen presentation in dendritic cells. *Curr Opin Immunol*. Feb. 2001;13(1):45-51. doi: 10.1016/s0952-7915(00)00180-1. cited by applicant

Van Boheemen et al., Genomic characterization of a newly discovered coronavirus associated with acute respiratory distress syndrome in humans. *mBio*. Nov. 20, 2012;3(6):e00473-12. doi: 10.1128/mBio.00473-12. cited by applicant

Vardi et al., Moderna's Mysterious Medicines. *Forbes*. Dec. 14, 2016: 5 pages. cited by applicant

Vardi, Moderna's Mysterious Coronavirus Vaccine Delivery System. *Forbes*. Jul. 29, 2020: 9 pages. cited by applicant

Verma et al., Double-headed nucleosides: Synthesis and applications. *Beilstein J Org Chem*. Jun. 8, 2021;17:1392-1439. doi: 10.3762/bjoc.17.98. eCollection 2021. cited by applicant

Vogel et al., BNT162b vaccines protect rhesus macaques from SARS-CoV-2. *Nature*. Apr. 2021;592(7853):283-289. doi: 10.1038/s41586-021-03275-y. Epub Feb. 1, 2021. cited by applicant

Von Niessen et al., Improving mRNA-Based Therapeutic Gene Delivery by Expression-Augmenting 3' UTRs Identified by Cellular Library Screening. *Mol Ther*. Apr. 10, 2019;27(4): 824-836. doi: 10.1016/j.ymthe.2018.12.011. Epub Dec. 18, 2018. cited by applicant

Walls et al., Crucial steps in the structure determination of a coronavirus spike glycoprotein using cryo-electron microscopy. *Protein Sci*. Jan. 2017;26(1):113-121. doi: 10.1002/pro.3048. Epub Oct. 18, 2016. cited by applicant

Walls et al., Tectonic conformational changes of a coronavirus spike glycoprotein promote membrane fusion. *Proc Natl Acad Sci U S A*. Oct. 17, 2017;114(42):11157-11162. doi: 10.1073/pnas.1708727114. Epub Oct. 3, 2017. cited by applicant

Walsh et al., Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *N Engl J Med*. Oct. 14, 2020;NEJMoa2027906: 1-13. doi: 10.1056/NEJMoa2027906. Online ahead of print. cited by applicant

Wang et al., An Evidence Based Perspective on mRNA-SARS-COV-2 Vaccine Development. *Med Sci Monit*. May 5, 2020;26: e924700. doi: 10.12659/MSM.924700. cited by applicant

Wang et al., Antibody Resistance of SARS-COV-2 Variants B.1.351 and B.1.1.7. *bioRxiv* [Preprint]. Feb. 12, 2021:2021.01.25.428137. doi: 10.1101/2021.01.25.428137. 24 pages. Update in: *Nature*. May 2021;593(7857):130-135. cited by applicant

Wang et al., Chapter 3: Lipid Nanoparticles for the Delivery of Nucleic Acids. Book: *Nanoparticulate Drug Delivery Systems: Strategies, Technologies, and Applications*. 2013. 29 pages. cited by applicant

Wang et al., Increased Resistance of SARS-COV-2 Variants B.1.351 and B.1.1.7 to Antibody Neutralization. *bioRxiv*. Jan. 26, 2021; 23 pages. doi: 10.1101 /2021.01.25.428137. Preprint. cited by applicant

Wang et al., mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *Nature*. Apr. 2021;592(7855):616-622. doi: 10.1038/s41586-021-03324-6. Epub Feb. 1, 2021. PMID: 33567448; PMCID: PMC8503938. cited by applicant

Wang et al., Structural Definition of a Neutralization-sensitive Epitope on the Mers-CoV S1-NTD. *Cell Rep*. Sep. 24, 2019;28(13):3395-3405.e6. doi: 10.1016/j.celrep.2019.08.052. cited by applicant

Weissman, mRNA transcript therapy. *Expert Rev Vaccines*. Feb. 2015;14(2):265-81. doi: 10.1586/14760584.2015.973859. Epub Oct. 31, 2014. cited by applicant

Whitehead et al., Knocking down barriers: advances in siRNA delivery. *Nat Rev Drug Discov*. Feb. 2009;8(2):129-38. doi: 10.1038/nrd2742. cited by applicant

Whitt, Generation of VSV pseudotypes using recombinant ΔG-VSV for studies on virus entry, identification of entry inhibitors, and immune responses to vaccines. *J Virol Methods*. Nov. 2010;169(2):365-74. doi: 10.1016/j.jviromet.2010.08.006. Epub Aug. 13, 2010. cited by applicant

Wilhelm et al., Reduced Neutralization of SARS-CoV-2 Omicron Variant by Vaccine Sera and Monoclonal Antibodies. medRxiv. 2021. doi:10.1101/2021.12.07.21267432:2021.12.07.21267432. 9 pages. cited by applicant

Wrapp et al., Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science. Feb. 19, 2020. pii: eabb2507. doi: 10.1126/science.abb2507. cited by applicant

Wu et al., A new coronavirus associated with human respiratory disease in China. Nature. Mar. 2020;579(7798):265-269. doi: 10.1038/s41586-020-2008-3. Epub Feb. 3, 2020. cited by applicant

Wu et al., mRNA-1273 vaccine induces neutralizing antibodies against spike mutants from global SARS-CoV-2 variants. bioRxiv preprint. Jan. 25, 2021. doi: 10.1101/2021.01.25.427948. 20 pages. cited by applicant

Wu et al., Serum Neutralizing Activity Elicited by mRNA-1273 Vaccine. N Engl J Med. Apr. 15, 2021;384(15):1468-1470. doi: 10.1056/NEJMc2102179. Epub Mar. 17, 2021. cited by applicant

Wu et al., Variant SARS-CoV-2 mRNA vaccines confer broad neutralization as primary or booster series in mice. Vaccine. Dec. 17, 2021;39(51):7394-7400. cited by applicant

Xia, Detailed Dissection and Critical Evaluation of the Pfizer/BioNTech and Moderna mRNA Vaccines. Vaccines (Basel). Jul. 3, 2021;9(7):734. doi: 10.3390/vaccines9070734. cited by applicant

Yang et al., A DNA vaccine induces SARS coronavirus neutralization and protective immunity in mice. Nature. Apr. 1, 2004;428(6982):561-4. doi: 10.1038/nature02463. cited by applicant

Yang et al., The structure and functions of coronavirus genomic 3' and 5' ends. Virus Res. Aug. 3, 2015;206:120-33. doi: 10.1016/j.virusres.2015.02.025. Epub Feb. 28, 2015. cited by applicant

Ye et al., Rational development of a combined mRNA vaccine against COVID-19 and influenza. NPJ Vaccines. Jul. 26, 2022;7(1):84. doi: 10.1038/s41541-022-00478-w. cited by applicant

Yin et al., Immunogenicity and protective efficacy of recombinant fusion proteins containing spike protein of infectious bronchitis virus and hemagglutinin of H3N2 influenza virus in chickens. Virus Res. Sep. 2, 2016;223:206-12. doi: 10.1016/j.virusres.2016.07.010. Epub Aug. 3, 2016. cited by applicant

Yin et al., Non-viral vectors for gene-based therapy. Nat Rev Genet. Aug. 2014;15(8):541-55. doi: 10.1038/nrg3763. Epub Jul. 15, 2014. cited by applicant

Zangi et al., Modified mRNA directs the fate of heart progenitor cells and induces vascular regeneration after myocardial infarction. Nat Biotechnol. Oct. 2013;31(10):898-907. doi: 10.1038/nbt.2682. Epub Sep. 8, 2013. cited by applicant

Zeng et al., Formulation and Delivery Technologies for mRNA Vaccines. Curr Top Microbiol Immunol. Jun. 2, 2020;10.1007/82_2020_217. doi: 10.1007/82_2020_217. cited by applicant

Zhang et al., A Thermostable mRNA Vaccine against COVID-19. Cell. Sep. 3, 2020;182(5):1271-1283.e16. doi: 10.1016/j.cell.2020.07.024. Epub Jul. 23, 2020. cited by applicant

Zhang et al., Delivery of mRNA vaccine with a lipid-like material potentiates antitumor efficacy through Toll-like receptor 4 signaling. Proc Natl Acad Sci U S A. Feb. 9, 2021;118(6):e2005191118. doi: 10.1073/pnas.2005191118. cited by applicant

Zhang et al., Prefusion spike protein stabilization through computational mutagenesis. Proteins. Apr. 2021;89(4):399-408. doi: 10.1002/prot.26025. Epub Dec. 4, 2020. cited by applicant

Zhao et al., Key Aspects of Coronavirus Avian Infectious Bronchitis Virus. Pathogens. May 11, 2023;12(5):698. doi: 10.3390/pathogens12050698. cited by applicant

Zhou et al., Evidence of escape of SARS-CoV-2 variant B.1.351 from natural and vaccine-induced sera. Cell. Apr. 29, 2021; 184(9): 2348-2361.e6. cited by applicant

Zhukovsky et al., Heterogeneity of early intermediates in cell-liposome fusion mediated by influenza hemagglutinin. Biophys J. Nov. 1, 2006;91(9):3349-58. doi: 10.1529/biophysj.106.088005. Epub Aug. 11, 2006. cited by applicant

Ziganshina et al., Antibody-Dependent Enhancement with a Focus on SARS-CoV-2 and Anti-Glycan Antibodies. Viruses. Jul. 20, 2023;15(7):1584. doi: 10.3390/v15071584. cited by applicant

International Search Report and Written Opinion for Application No. PCT/US2016/058327 mailed Jun. 29, 2017. cited by applicant

[No Author Listed], “Messenger RNA”, Internet: Wikipedia. Jun. 19, 2013, XP002699196, Retrieved from the Internet: URL: [http://en.wikipedia.org/wiki/Messenger RNA](http://en.wikipedia.org/wiki/Messenger_RNA). cited by applicant

[No Author Listed], Geneseq Accession No. BC153556. First entry: Jan. 14, 2016. 3 pages. cited by applicant

[No Author Listed], Geneseq Accession No. BGP67180. First entry: Sep. 19, 2019. 3 pages. cited by applicant

[No Author Listed], Katalin Karikó and Drew Weissman awarded Horwitz Prize for pioneering research on COVID-19 vaccines, Columbia University Irving Medical Center, Aug. 16, 2021, <https://www.cuimc.columbia.edu/news/horwitz-prize-2021> (last accessed Aug. 24, 2023). 12 pages. cited by applicant

[No Author Listed], List of past and pending litigations. Prepared Oct. 24, 2023. 1 page. cited by applicant

Agrawal et al., Immunization with inactivated Middle East Respiratory Syndrome coronavirus vaccine leads to lung immunopathology on challenge with live virus. *Hum Vaccin Immunother.* Sep. 2016;12(9):2351-6. doi:10.1080/21645515.2016.1177688. Epub Jun. 7, 2016. cited by applicant

Aliprantis et al., A phase 1, randomized, placebo-controlled study to evaluate the safety and immunogenicity of an mRNA-based RSV prefusion F protein vaccine in healthy younger and older adults. *Hum Vaccin Immunother.* May 4, 2021;17(5):1248-1261. doi: 10.1080/21645515.2020.1829899. Epub Oct. 29, 2020. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc. (Fed. Circ. 2023-2357) D.I. 8—Docketing Statement, Sep. 21, 2023. 2 pages. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc et al. (D. Del. 23-cv-580-CFC) D.I. 1—Complaint for Patent Infringement, May 26, 2023. 28 pages. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc et al. (D. Del. 23-cv-580-CFC) D.I. 12—Answer to the Complaint, Jul. 17, 2023. 37 pages. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc et al. (D. Del. 22-cv-335-CFC) D.I. 1—Complaint for Patent Infringement, Mar. 17, 2022. 14 pages. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc et al. (D. Del. 22-cv-335-CFC) D.I. 128—Final Judgement, Aug. 30, 2023. 2 pages. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc et al. (D. Del. 22-cv-335-CFC) D.I. 130—Notice of Appeal to the U.S. Court of Appeals for the Federal Circuit, Aug. 31, 2023. 3 pages. cited by applicant

Alnylam Pharmaceuticals, Inc. v. Moderna, Inc et al. (D. Del. 22-cv-335-CFC) D.I. 87—Answer to the Complaint, May 10, 2023. 44 pages. cited by applicant

Arbutus BioPharma Corp et al. v. Moderna, Inc et al. (D. Del. 22-cv-252-MSG) D.I. 1—Complaint for Patent Infringement, Feb. 28, 2022. 51 pages. cited by applicant

Arbutus BioPharma Corp et al. v. Moderna, Inc et al. (D. Del. 22-cv-252-MSG) D.I. 31—District Court Memorandum, Nov. 2, 2022. 16 pages. cited by applicant

Arbutus BioPharma Corp et al. v. Moderna, Inc et al. (D. Del. 22-cv-252-MSG) D.I. 35—Answer to the Complaint, Nov. 30, 2022. 83 pages. cited by applicant

Arbutus BioPharma Corp et al. v. Moderna, Inc et al. (D. Del. 22-cv-252-MSG) D.I. 64—District Court Memorandum, Mar. 10, 2023. 4 pages. cited by applicant

Archer, S.J., Induction of a T-cell specific antigen on bone marrow lymphocytes with thymus RNA. *Immunology.* Jan. 1978;34(1):123-9. cited by applicant

Ashley, D.M. et al., Bone marrow-generated dendritic cells pulsed with tumor extracts or tumor RNA induce antitumor immunity against central nervous system tumors. *J Exp Med.* Oct. 6, 1997; 186(7): 1177-82. cited by applicant

Austin et al., Split-Dose Administration Enhances Immune Responses Elicited by a mRNA/Lipid Nanoparticle Vaccine Expressing Respiratory Syncytial Virus F Protein. *Mol Pharm.* Jan. 2, 2023;20(1):279-289. doi: 10.1021/acs.molpharmaceut.2c00635. Epub Oct. 17, 2022. cited by applicant

Barbier et al., The clinical progress of mRNA vaccines and immunotherapies. *Nat Biotechnol.* Jun. 2022;40(6):840-854. doi: 10.1038/s41587-022-01294-2. Epub May 9, 2022. cited by applicant

Bettinger, T. et al., Peptide-mediated RNA delivery: a novel approach for enhanced transfection of primary and post-mitotic cells. *Nucleic Acids Res.* Sep. 15, 2001;29(18):3882-91. cited by applicant

Bogers et al., Potent immune responses in rhesus macaques induced by nonviral delivery of a self-amplifying RNA vaccine expressing HIV type 1 envelope with a cationic nanoemulsion. *J Infect Dis.* Mar. 15, 2015;211(6):947-55. doi: 10.1093/infdis/jiu522. Epub Sep. 18, 2014. cited by applicant

Bonehill, A., et al., Single-step antigen loading and activation of dendritic cells by mRNA electroporation for the purpose of therapeutic vaccination in melanoma patients. *Clin Cancer Res.* May 2009; 15(10): 3366-3375. cited by applicant

Bose, S. et al., Role of nucleolin in human parainfluenza virus type 3 infection of human lung epithelial cells. *J Viral.* Aug. 2004;78(15):8146-58. cited by applicant

Byoung-Shik et al., Intranasal immunization with plasmid DNA encoding spike protein of SARS-coronavirus/polyethylenimine nanoparticles elicits antigen-specific humoral and cellular immune responses. *BMC Immunol.* Dec. 31, 2010;11:65. doi: 10.1186/1471-2172-11-65. cited by applicant

Chaudhary et al., mRNA vaccines for infectious diseases: principles, delivery and clinical translation. *Nature Reviews Drug Discovery* vol. 20, pp. 817-838 (2021). cited by applicant

Clancy et al., Translation: DNA to mRNA to Protein. *Nature Education* 2008;1(1):101. cited by applicant

Cobb et al., Who discovered messenger RNA? *Curr Biol.* Jun. 29, 2015;25(13):R526-32. doi:10.1016/j.cub.2015.05.032. cited by applicant

Conry, R.M. et al., Characterization of a messenger RNA polynucleotide vaccine vector. *Cancer Res.* Apr. 1, 1995 ;55 (7):1397-1400. cited by applicant

Cross, Without these lipid shells, there would be no mRNA vaccines for COVID-19. *Chem Eng News.* Mar. 6, 2021; 99(8). 8 pages. cited by applicant

Cullis et al., Lipid Nanoparticle Systems for Enabling Gene Therapies. *Mol Ther.* Jul. 5, 2017;25(7):1467-1475. doi: 10.1016/j.ymthe.2017.03.013. Epub Apr. 13, 2017. cited by applicant

Dahlman, James E. et al., In vivo endothelial siRNA delivery using polymeric nanoparticles with low molecular weight, *Nature Nanotechnology*, 2014, No. vol.#, pp. 1-8. cited by applicant

Deering et al., Nucleic acid vaccines: prospects for non-viral delivery of mRNA vaccines. *Expert Opin Drug Deliv.* Jun. 2014;11(6):885-99. doi: 10.1517/17425247.2014.901308. Epub Mar. 26, 2014. cited by applicant

Degroot et al., Part II-13 The Positive Sense Single Stranded RNA Viruses, Family Coronaviridae, in *Virus Taxonomy: Ninth Report of the International Committee on Taxonomy of Viruses* 806 (2012). 25 pages. cited by applicant

Del Pozo-Rodriguez et al., Lipid nanoparticles as vehicles for macromolecules: nucleic acids and peptides. *Recent Pat Drug Deliv Formul.* Sep. 2011;5(3):214-26. doi: 10.2174/18722111797200515. cited by applicant

Diken et al., Current Developments in Actively Personalized Cancer Vaccination with a Focus on RNA as the Drug Format. *Prog Tumor Res.* 2015;42:44-54. doi: 10.1159/000437184. Epub Sep. 4, 2015. Review. cited by applicant

Du et al., Recombinant adeno-associated virus expressing the receptor-binding domain of severe acute respiratory syndrome coronavirus S protein elicits neutralizing antibodies: Implication for developing SARS vaccines. *Virology.* Sep. 15, 2006;353(1):6-16. doi: 10.1016/j.virol.2006.03.049.

Epub Jun. 21, 2006. cited by applicant

Fleeton et al., Self-replicative RNA vaccines elicit protection against influenza A virus, respiratory syncytial virus, and a tickborne encephalitis virus. *J Infect Dis.* May 1, 2001;183(9):1395-8. Epub Mar. 30, 2001. cited by applicant

Furuichi, Caps on Eukaryotic mRNAs. eLS. John Wiley & Sons. Jul. 2014. 1-12. cited by applicant

Geall et al., Nonviral delivery of self-amplifying RNA vaccines. *Proc Natl Acad Sci U S A.* Sep. 4, 2012;109(36):14604-9. doi: 10.1073/pnas.1209367109. Epub Aug. 20, 2012. cited by applicant

Geall et al., RNA: the new revolution in nucleic acid vaccines. *Semin Immunol.* Apr. 2013;25(2):152-9. doi: 10.1016/j.smim.2013.05.001. Epub Jun. 2, 2013. cited by applicant

GenBank Accession No. AHX22069. First seen on NCBI on May 14, 2014. cited by applicant

Glenn et al., Safety and immunogenicity of a Sf9 insect cell-derived respiratory syncytial virus fusion protein nanoparticle vaccine. *Vaccine.* Jan. 7, 2013;31(3):524-32. doi: 10.1016/j.vaccine.2012.11.009. cited by applicant

Graham et al., Novel antigens for RSV vaccines. *Curr Opin Immunol.* Aug. 2015;35:30-8. doi: 10.1016/j.coi.2015.04.005. Epub Jun. 10, 2015. cited by applicant

Graham et al., Novel Vaccine Technologies: Essential Components of an Adequate Response to Emerging Viral Diseases. *JAMA.* Apr. 10, 2018;319(14):1431-1432. doi: 10.1001/jama.2018.0345. cited by applicant

Greer et al., Long-term protection in hamsters against human parainfluenza virus type 3 following mucosal or combinations of mucosal and systemic immunizations with chimeric alphavirus-based replicon particles. *Scand J Immunol.* Dec. 2007;66(6):645-53. Epub Oct. 17, 2007. cited by applicant

Hartmann et al., Handbook of RNA Biochemistry, Second Edition, "Part I RNA Synthesis and Detection." 2014. p 1-27. cited by applicant

Hecker, J.G. et al., Non-Viral DNA and mRNA Gene Delivery to the CNS Pre-Operatively for Neuroprotection and Following Neurotrauma. *Molecular Therapy.* 2004; 9, S258-S258. cited by applicant

Heiser, A. et al., Induction of polyclonal prostate cancer-specific CTL using dendritic cells transfected with amplified tumor RNA. *J Immunol.* Mar. 1, 2001; 166(5):2953-60. cited by applicant

Heyes et al., Cationic lipid saturation influences intracellular delivery of encapsulated nucleic acids. *J Control Release.* Oct. 3, 2005;107(2):276-87. cited by applicant

Hoerr, I. et al., In vivo application of RNA leads to induction of specific cytotoxic T lymphocytes and antibodies. *Eur J Immunol.* Jan. 2000;30(1):1-7. cited by applicant

Hoerr, I. et al., Stabilized Messenger RNA (RNAActive™) as a Tool for Innovative Gene Delivery. *Tissue Engineering.* Apr. 2007; 13(4): 865-925. cited by applicant

Hoerr, More than a messenger: A new class of drugs-mRNA-based therapeutics. *Genetic Engineering & Biotechnology News.* Jun. 18, 2013. <http://www.genengnews.com/gen-articles/more-than-a-messenger-a-new-class-of-drugs-mrna-based-therapeutics/4916/> [last accessed Mar. 25, 2016]. cited by applicant

Holtkamp, S. et al., Modification of antigen-encoding RNA increases stability, translational efficacy, and T-cell stimulatory capacity of dendritic cells. *Blood.* Dec. 15, 2006;108(13):4009-17. cited by applicant

Hörr, RNA vaccine for the induction of specific cytotoxic T lymphocytes (CTL) and antibodies. Eberhard Karls University of Tübingen. Dissertation. English-Language Translation. 1999. 138 Pages. cited by applicant

Hou et al., Lipid nanoparticles for mRNA delivery. *Nature Reviews Materials* vol. 6, pp. 1078-1094 (2021). cited by applicant

Human Metapneumovirus (HM PV) Clinical Features, CDC Website, <https://www.cdc.gov/surveillance/nrevss/hmpv/clinical.html>, Sep. 5, 2019. (Year: 2019). cited by

applicant

Jirikowski, G.F., et al., Reversal of diabetes insipidus in Brattleboro Rats: Intrahypothalamic injection of vasopressin mRNA. *Science*. Feb. 1992; 255(5047): 996-998. cited by applicant
Jorquera et al., Nanoparticle vaccines encompassing the respiratory syncytial virus (RSV) G protein CX3C chemokine motif induce robust immunity protecting from challenge and disease. *PLoS One*. Sep. 10, 2013;8(9):e74905. doi: 10.1371/journal.pone.0074905. eCollection 2013. cited by applicant

Kallen et al., A development that may evolve into a revolution in medicine: mRNA as the basis for novel, nucleotide-based vaccines and drugs. *Ther Adv Vaccines*. Jan. 2014;2(1):10-31. doi: 10.1177/2051013613508729. cited by applicant

Kallen et al., A novel, disruptive vaccination technology: self-adjuvanted RActive(®) vaccines. *Hum Vaccin Immunother*. Oct. 2013;9(10):2263-76. doi: 10.4161/hv.25181. Epub Jun. 4, 2013. Review. cited by applicant

Kalra et al., Virosomes: As a Drug Delivery Carrier. *American Journal of Advanced Drug Delivery*. 2013;1:29-35. cited by applicant

Kanapathipillai, et al., Nanoparticle targeting of anti-cancer drugs that alter intracellular signaling or influence the tumor microenvironment, *Adv. Drug Deliv. Rev.* (2014), , pp. 1-12. cited by applicant

Kariko et al., Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability. *Mol Ther*. Nov. 2008; 16(11):1833-40. Epub Sep. 16, 2008. cited by applicant

Kariko, K., et al., Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA, *Nucleic Acids Research*, Oxford University Press, GB, vol. 39, No. 21, Sep. 2, 2011 (Sep. 2, 2011), e142. doi: 10.1093/nar/gkr695. Epub Sep. 2, 2011. cited by applicant

Kauffman et al., Optimization of Lipid Nanoparticle Formulations for mRNA Delivery in Vivo with Fractional Factorial and Definitive Screening Designs. *Nano Lett*. Nov. 11, 2015;15(11):7300-6. doi: 10.1021/acs.nanolett.5b02497. Epub Oct. 20, 2015. cited by applicant

Keshwara et al., Rabies-based vaccine induces potent immune responses against Nipah virus. *NPJ Vaccines*. Apr. 15, 2019;4:15. Erratum in: *NPJ Vaccines*. May 13, 2019;4:18. cited by applicant

Kozielski et al., Bio-reducible cationic polymer-based nanoparticles for efficient and environmentally triggered cytoplasmic siRNA delivery to primary human brain cancer cells. *ACS Nano*. Apr. 22, 2014;8(4):3232-41. doi: 10.1021/nn500704t. Epub Apr. 3, 2014. cited by applicant

Kramps et al., Messenger RNA-based vaccines: progress, challenges, applications. *Wiley Interdiscip Rev RNA*. Nov.-Dec. 2013;4(6):737-49. doi: 10.1002/wrna.1189. Epub Jul. 25, 2013. cited by applicant

Kreiter, S., et al., Intranodal vaccination with naked antigen-encoding RNA elicits potent prophylactic and therapeutic antitumoral immunity. *Cancer Res*. 2010; 70: 9031-9040. cited by applicant

Kreiter, S., et al., Tumor vaccination using messenger RNA: prospects of a future therapy. *Curr Opinion in Immun*. Jun. 2011; 23(3): 399-406. cited by applicant

Kresge, The messenger RNA pioneers everyone ignored. *Bloomberg*, Nov. 23, 2021, <https://www.bloomberg.com/news/newsletters/2021-11-23/themessenger-rna-pioneers-everyone-ignored> (last accessed Aug. 24, 2023). 8 pages. cited by applicant

Kuhn, A.N., et al., mRNA as a versatile tool for exogenous protein expression. *Current Gene Therapy*. Oct. 2012; 12 (5): 347-361. cited by applicant

Leitner, W.W. et al., DNA and RNA-based vaccines: principles, progress and prospects. *Vaccine*. Dec. 10, 1999;18 (9-10):765-77. cited by applicant

Li et al., Effects of local structural transformation of lipid-like compounds on delivery of messenger RNA. *Sci Rep*. Feb. 26, 2016;6:22137. doi: 10.1038/srep22137. cited by applicant

Li et al., Lipid-based nanoparticles for nucleic acid delivery. *Pharm Res.* Mar. 2007;24(3):438-49. doi: 10.1007/s11095-006-9180-5. cited by applicant

Li, L. et al., Overcoming obstacles to develop effective and safe siRNA therapeutics. *Expert Opin Biol Ther.* May 2009; 9(5): 609-19. cited by applicant

Liang et al., Enhanced Neutralizing Antibody Response Induced by Respiratory Syncytial Virus Prefusion F Protein Expressed by a Vaccine Candidate. *J Virol.* Sep. 2015;89(18):9499-510. doi: 10.1128/JVI.01373-15. Epub Jul. 8, 2015. cited by applicant

Lin et al., Progress and challenges of mRNA vaccines. *Interdisc Med.* Dec. 22, 2022;1(1):e20220008. doi: 10.1002/INMD.20220008. cited by applicant

Lorenzi, J.C., et al., Intranasal vaccination with messenger RNA as a new approach in gene therapy: Use against tuberculosis. *BMC Biotechnol.* Oct. 2010; 10(77): 1-11. cited by applicant

Lu et al., Bat-to-human: spike features determining 'host jump' of coronaviruses SARS-CoV, MERS-CoV, and beyond. *Trends Microbiol.* Aug. 2015;23(8):468-78. doi: 10.1016/j.tim.2015.06.003. Epub Jul. 21, 2015. cited by applicant

MacLachlan, Lipid Nanoparticle-mediated delivery of messenger RNA. Presentation. 1st International mRNA Health Conference. Tübingen, Germany. Oct. 24, 2013. http://files.shareholder.com/downloads/ABEA-50QJTB/2628241206x0x699789/47543d12-db34-4e6e-88a9-f3ae5d97b1d2/MacLachlan_mRNA_Conf_2013.pdf. Last accessed Dec. 22, 2016. cited by applicant

Magini et al., Self-Amplifying mRNA Vaccines Expressing Multiple Conserved Influenza Antigens Confer Protection against Homologous and Heterosubtypic Viral Challenge. *PLoS One.* Aug. 15, 2016;11(8):e0161193. doi: 10.1371/journal.pone.0161193. eCollection 2016. cited by applicant

Martinon, F. et al., Induction of virus-specific cytotoxic T lymphocytes in vivo by liposome-entrapped mRNA. *Eur J Immunol.* Jul. 1993;23(7):1719-22. cited by applicant

Mas et al., Engineering, Structure and Immunogenicity of the Human Metapneumovirus F Protein in the Postfusion Conformation. *PLoS Pathog.* Sep. 9, 2016;12(9):e1005859. doi:10.1371/journal.ppat.1005859. eCollection Sep. 2016. cited by applicant

McDonnell et al. DNA vaccines. *N Engl J Med.* Jan. 4, 1996;334(1):42-5. doi: 10.1056/NEJM199601043340110. cited by applicant

McLellan et al., Structure-based design of a fusion glycoprotein vaccine for respiratory syncytial virus. *Science.* Nov. 1, 2013;342(6158):592-8. doi: 10.1126/science.1243283. cited by applicant

Midoux et al., Lipid-based mRNA vaccine delivery systems. *Expert Rev Vaccines.* Feb. 2015;14(2):221-34. doi: 10.1586/14760584.2015.986104. Epub Dec. 26, 2014. Review. cited by applicant

Mitchell, DA et al., RNA transfected dendritic cells as cancer vaccines. *Curr Opin Mol Ther.* Apr. 2000;2(2):176-81. cited by applicant

Mitchell, DA et al., RNA-transfected dendritic cells in cancer immunotherapy. *J Clin Invest.* Nov. 2000;106 (9):1065-9. cited by applicant

Mockey et al., mRNA-based cancer vaccine: prevention of B16 melanoma progression and metastasis by systemic injection of MART1 mRNA histidylated lipopolyplexes, *Cancer Gene Therapy*, 2007, 14, pp. 802-814. cited by applicant

Mok et al., An Alphavirus Replicon-Based Human Metapneumovirus Vaccine Is Immunogenic and Protective in Mice and Cotton Rats. *J Virol.* Nov. 2008; 82(22): 11410-11418. cited by applicant

Muller, M.R. et al., Transfection of dendritic cells with RNA induces CD4- and COB-mediated T cell immunity against breast carcinomas and reveals the immunodominance of presented T cell epitopes. *J Immunol.* Jun. 15, 2003;170(12):5892-6. cited by applicant

Narayanan et al., Interplay between viruses and host mRNA degradation. *Biochim Biophys Acta.* Jun.-Jul. 2013;1829(6-7):732-41. doi: 10.1016/j.bbagr.2012.12.003. Epub Dec. 26, 2012. cited by applicant

Pardi et al., mRNA vaccines—a new era in vaccinology. *Nat Rev Drug Discov.* Apr.

2018;17(4):261-279. doi: 10.1038/nrd.2017.243. Epub Jan. 12, 2018. cited by applicant

Pascolo, S. Vaccination with messenger RNA (mRNA). *Handb Exp Pharmacol*. 2008; 183:221-235. cited by applicant

Petition For Inter Partes Review of U.S. Pat. No. 10,702,600. *BioNTech SE and Pfizer Inc.* (Petitioners) v. *ModernaTX, Inc.* (Patent Owner). Filed Aug. 28, 2023. 87 pages. cited by applicant

Petition For Inter Partes Review of U.S. Pat. No. 10,933,127. *BioNTech SE and Pfizer Inc.* (Petitioners) v. *ModernaTX, Inc.* (Patent Owner). Filed Aug. 28, 2023. 89 pages. cited by applicant

Petition For Inter Partes Review of U.S. Pat. No. 8,058,069 (IPR2019-00554). *Moderna Therapeutics, Inc.* (Petitioner) v. *Arbutus Biopharma Corporation* (Patent Owner). Filed Nov. 15, 2011. 81 pages. cited by applicant

Petsch et al., Protective efficacy of in vitro synthesized, specific mRNA vaccines against influenza A virus infection. *Nat Biotechnol*. Dec. 2012;30(12):1210-6. doi: 10.1038/nbt.2436. Epub Nov. 25, 2012. cited by applicant

Phua et al., Messenger RNA (mRNA) nanoparticle tumour vaccination. *Nanoscale*. Jul. 21, 2014;6(14):7715-29. doi: 10.1039/c4nr01346h. Review. cited by applicant

Promosome, LLC v. Moderna, Inc. (S.D. Cal. 23-cv-1047-JES-DDL) D.I. 1—Complaint for Patent Infringement, Jun. 6, 2023. 51 pages. cited by applicant

Promosome, LLC v. Moderna, Inc. (S.D. Cal. 23-cv-1047-JES-DDL) D.I. 30—Notice of Dismissal, Sep. 19, 2023. 4 pages. cited by applicant

Pulford, B., et al., Liposome-siRNA-peptide complexes cross the blood-brain barrier and significantly decrease PrP^C on neuronal cells and PrP^{RES} in infected cell cultures. *PloS ONE*. 2010; 5(6): e11085. cited by applicant

Rabinovich, P.M., et al., Synthetic messenger RNA as a tool for gene therapy. *Hum. Gene Ther*. Oct. 2006; 17: 1027-1035. cited by applicant

Ramamoorth et al., Nonviral vectors in gene therapy—an overview. *J Clin Diagn Res*. Jan. 2015; 9(1): GE01-GE06. cited by applicant

Reichmuth et al., mRNA Vaccine Delivery Using Lipid Nanoparticles. *Ther Deliv*. 2016;7(5):319-34. doi: 10.4155/tde-2016-0006. cited by applicant

Ren et al., Recent vaccine development for human metapneumovirus. *J Gen Virol*. Jul. 2015;96(Pt 7):1515-20. doi: 10.1099/vir.0.000083. Epub Feb. 9, 2015. cited by applicant

Rittig et al., Intradermal vaccinations with RNA coding for TAA generate CD8⁺ and CD4⁺ immune responses and induce clinical benefit in vaccinated patients. *Mol Ther*. May 2011;19(5):990-9. doi: 10.1038/mt.2010.289. Epub Dec. 28, 2010. cited by applicant

Sahin et al., mRNA-based therapeutics—developing a new class of drugs. *Nat Rev Drug Discov*. Oct. 2014; 13(10):759-80. doi: 10.1038/nrd4278. Epub Sep. 19, 2014. cited by applicant

Schlake et al., Developing mRNA-vaccine technologies. *RNA Biol*. Nov. 2012;9(11):1319-30. doi: 10.4161/rna.22269. Epub Oct. 12, 2012. cited by applicant

Schmitt, W.E. et al., In vitro induction of a bladder cancer-specific T-cell response by mRNA-transfected dendritic cells. *J Cancer Res Clin Oncol*. 2001 ;127(3):203-6. cited by applicant

Schott, J.W., et al., Viral and non-viral approaches for transient delivery of mRNA and proteins. *Current Gene Ther*. 2011; 11 (5): 382-398. cited by applicant

Segura, J., et al., Monitoring gene therapy by external imaging of mRNA: Pilot study on murine erythropoietin. *Ther Drug Monit*. Oct. 2007; 29(5): 612-8. cited by applicant

Shin et al., Recent Advances in RNA Therapeutics and RNA Delivery Systems Based on Nanoparticles. *Adv. Therap.*, Nov. 2018; 1(7):1800065. Review. cited by applicant

Smits, E., et al., RNA-based gene transfer for adult stem cells and T cells. *Leukemia*. 2004; 18: 1898-1902. cited by applicant

Sohn, R.L., et al., In-vivo particle mediated delivery of mRNA to mammalian tissues: ballistic and biological effects. *Wound Rep and Regen*. Jul.-Aug. 2001; 287-296. cited by applicant

Strong, V.T. et al., Incorporation of beta-globin untranslated regions into a Sindbis virus vector for

augmentation of heterologous mRNA expression. *Gene Ther.* Jun. 1997;4(6):624-7. cited by applicant

Sullenger, Ba et al., Emerging clinical applications of RNA. *Nature.* Jul. 11, 2002;418(6894):252-8. cited by applicant

Tam et al., Advances in Lipid Nanoparticles for siRNA Delivery. *Pharmaceutics.* Sep. 18, 2013;5(3):498-507. doi: 10.3390/pharmaceutics5030498. cited by applicant

Tavernier, G., et al., mRNA as gene therapeutic: How to control protein expression. *J. of Controlled Release.* Mar. 2011; 150(3): 238-247. cited by applicant

Tenchov et al., Lipid Nanoparticles—From Liposomes to mRNA Vaccine Delivery, a Landscape of Research Diversity and Advancement. *ACS Nano.* Nov. 23, 2021;15(11):16982-17015. doi: 10.1021/acsnano.1c04996. Epub Jun. 28, 2021. cited by applicant

Terada et al., Characterization of Lipid Nanoparticles Containing Ionizable Cationic Lipids Using Design-of-Experiments Approach. *Langmuir.* Jan. 26, 2021;37(3):1120-1128. doi: 10.1021/acs.langmuir.0c03039. Epub Jan. 13, 2021. cited by applicant

Teufel, R. et al., Human peripheral blood mononuclear cells transfected with messenger RNA stimulate antigen-specific cytotoxic T-lymphocytes in vitro. *Cell Mol Life Sci.* Aug. 2005;62(15):1755-62. cited by applicant

Thess et al., Sequence-engineered mRNA Without Chemical Nucleoside Modifications Enables an Effective Protein Therapy in Large Animals. *Mol Ther.* Sep. 2015;23(9):1456-64. doi: 10.1038/mt.2015.103. Epub Jun. 8, 2015. cited by applicant

To et al., An overview of rational design of mRNA-based therapeutics and vaccines. *Expert Opin Drug Discov.* Nov. 2021;16(11):1307-1317. doi: 10.1080/17460441.2021.1935859. Epub Jul. 19, 2021. cited by applicant

Ulmer et al., RNA-based vaccines. *Vaccine.* Jun. 22, 2012;30(30):4414-8. doi: 10.1016/j.vaccine.2012.04.060. Epub Apr. 28, 2012. cited by applicant

Wang et al., Delivery of oligonucleotides with lipid nanoparticles. *Adv Drug Deliv Rev.* Jun. 29, 2015;87:68-80. doi: 10.1016/j.addr.2015.02.007. Epub Feb. 27, 2015. cited by applicant

Wang et al., Systemic delivery of modified mRNA encoding herpes simplex virus 1 thymidine kinase for targeted cancer gene therapy. *Mol Ther.* Feb. 2013;21(2):358-67. doi: 10.1038/mt.2012.250. Epub Dec. 11, 2012. cited by applicant

Wen et al., New Approaches for Immunization and Therapy against Human Metapneumovirus. *Clin Vaccine Immunol.* Aug. 2015;22(8):858-66. doi: 10.1128/CI.00230-15. Epub Jun. 10, 2015. cited by applicant

Williams et al., Human metapneumovirus isolate TN/92-4 fusion protein gene, complete eds. GenBank: EF051124.1. Nov. 29, 2007. cited by applicant

Wong et al., An mRNA vaccine for influenza. *Nat Biotechnol.* Dec. 2012;30(12):1202-4. doi: 10.1038/nbt.2439. cited by applicant

Yamamoto et al., Current prospects for mRNA gene delivery, *European Journal of Pharmaceutics and Biopharmaceutics* 71 (2009) 484-489. cited by applicant

Youn et al., Modified mRNA as an alternative to plasmid DNA (pDNA) for transcript replacement and vaccination therapy. *Expert Opin Biol Ther.* 2015;15(9):1337-48. doi: 10.1517/14712598.2015.1057563. Epub Jun. 30, 2015. cited by applicant

Yuan et al., Cryo-EM structures of MERS-CoV and SARS-CoV spike glycoproteins reveal the dynamic receptor binding domains. *Nat Commun.* Apr. 10, 2017;8:15092. doi: 10.1038/ncomms15092. cited by applicant

Zhao et al., Chapter Two: Lipid Nanoparticles for Gene Delivery. Book: *Advances in Genetics.* Elsevier, 2014. 24 pages. cited by applicant

Zhao et al., Nanoparticle vaccines. *Vaccine.* Jan. 9, 2014;32(3):327-37. doi: 10.1016/j.vaccine.2013.11.069. Epub Dec. 2, 2013. cited by applicant

Zhou, W.Z. et al., RNA melanoma vaccine: induction of antitumor immunity by human

glycoprotein 100 mRNA immunization. Hum Gene Ther. Nov. 1, 1999;10(16):2719-24. cited by applicant

Zumla et al., Middle East respiratory syndrome. Lancet. Sep. 5, 2015;386(9997):995-1007. doi: 10.1016/S0140-6736(15)60454-8. Epub Jun. 3, 2015. cited by applicant

[IPR-2023-01358] Institution Decision in IPR2023-01358 dated Mar. 19, 2024. cited by applicant

[IPR-2023-01358] Patent Owner's Preliminary Response in IPR2023-01358 dated Dec. 8, 2023. cited by applicant

[IPR-2023-01358] Patent Owner's Preliminary Statement Regarding Petitioner's Inconsistent Positions in IPR2023-01358 dated Jan. 16, 2024. cited by applicant

[IPR-2023-01358] Patent Owner's Response in IPR2023-01358 dated May 31, 2024. cited by applicant

[IPR-2023-01358] Patent Owner's Sur-Reply in IPR2023-01358 dated Oct. 16, 2024. cited by applicant

[IPR-2023-01358] Patent Owner's Sur-Reply in Support of Preliminary Response in IPR2023-01358 dated Jan. 25, 2024. cited by applicant

[IPR-2023-01358] Petitioner's Preliminary Statement Regarding Patent Owner's Inconsistent Positions in IPR-2023-01358 dated Jan. 23, 2024. cited by applicant

[IPR-2023-01358] Petitioner's Reply to Patent Owner's Preliminary Response in IPR2023-01358 dated Jan. 12, 2024. cited by applicant

[IPR-2023-01358] Petitioner's Reply to Patent Owner's Response in IPR2023-01358 dated Aug. 30, 2024. cited by applicant

[IPR2023-01359] Institution Decision dated Mar. 19, 2024 in IPR2023-01359. cited by applicant

[IPR2023-01359] Patent Owner's Preliminary Response in IPR2023-01359 dated Dec. 7, 2023. cited by applicant

[IPR2023-01359] Patent Owner's Preliminary Statement Regarding Alleged Inconsistent Positions in IPR2023-01359 dated Jan. 16, 2024. cited by applicant

[IPR2023-01359] Patent Owner's Response in IPR2023-01359 dated May 31, 2024. cited by applicant

[IPR2023-01359] Patent Owner's Sur-Reply in IPR2023-01359 dated Oct. 16, 2024. cited by applicant

[IPR2023-01359] Patent Owner's Sur-Reply in Support of Preliminary Response in IPR2023-01359 dated Jan. 25, 2024. cited by applicant

[IPR2023-01359] Petitioner's Reply to Patent Owner's Preliminary Response in IPR2024-01359 dated Jan. 12, 2024. cited by applicant

[IPR2023-01359] Petitioner's Reply to Patent Owner's Response in IPR2023-01359 dated Aug. 30, 2024. cited by applicant

Agnihothram, S., Review Committee Chair, Summary Basis for Regulatory Action, 2022, STN 125752/0 (30 Pages). cited by applicant

Ahmed et al., Biochemistry, Lipids, Statpearls Publishing (2023) (available at <https://www.ncbi.nlm.nih.gov/books/NBK525952/>). cited by applicant

Akinc et al., "Targeted Delivery of RNAi Therapeutics With Endogenous and Exogenous Ligand-Based Mechanisms," 2010, vol. 18, Issue 7, Molecular Therapy (8 pages). cited by applicant

Allen et al., "Liposomal drug delivery systems: From concept to clinical applications," 2013, Adv. Drug Deliv. Rev., pp. 36-48. cited by applicant

Atkins et al., "The molecular pathogenesis of Semliki Forest virus: a model virus made useful?," Journal of General Virology 2287, 1999 (11 Pages). cited by applicant

Bahl et al., "Preclinical and Clinical Demonstration of Immunogenicity by mRNA Vaccines against H10N8 and H7N9 Influenza Viruses," 2017, vol. 25, Issue 6, Mol Ther. cited by applicant

Bisht et al., "Severe acute respiratory syndrome coronavirus spike protein expressed by attenuated vaccinia virus protectively immunizes mice," 2004, Biological Sciences (6 Pages). cited by

applicant

Brito et al., "A Cationic Nanoemulsion for the Delivery of Next-generation RNA Vaccines," 2014, The American Society of Gene & Cell Therapy (12 Pages). cited by applicant

Buchholz et al., "Contributions of the structural proteins of severe acute respiratory syndrome coronavirus to protective immunity," 2004, vol. 101, Issue 26, Biological Sciences. cited by applicant

Buschmann et al., "Chitosans for Delivery of Nucleic Acids," 2013, Advanced Drug Delivery Reviews, pp. 1234-1270. cited by applicant

Cassimeris et al., "Lewin's Cells," 2011, Jones and Bartlett Publishers, vol. 2 (109 Pages). cited by applicant

Center for Drug Evaluation and Research Approval Package for New Drug Application No. 50-718/S-50 for Doxil (97 Pages). cited by applicant

Chalkias et al., "Safety, immunogenicity and antibody persistence of a bivalent Beta-containing booster vaccine against COVID-19: a phase 2/3 trial," 2022, Nature Medicine, pp. 2388-2397. cited by applicant

Chang et al., Recent Insights into the Developments of Therapeutics Against Coronavirus Disease by Targeting N Protein, 2016, Drug Discovery Today, No. 4, pp. 562-572. cited by applicant

Cherry et al., "SARS: The First Pandemic of the 21st Century," 2004, Pediatric Research, vol. 56, Issue 1, pp. 1-5. cited by applicant

Coelho et al., "Safety and Efficacy of RNAi Therapy for Transthyretin Amyloidosis," 2013, vol. 369, Issue 9, New England J. Med. (11 Pages). cited by applicant

Curriculum Vitae of Daniel O. Griffin, M.D., Ph.D. cited by applicant

Curriculum Vitae of James J. Moon, Ph.D. cited by applicant

Curriculum Vitae of Deborah Fuller (Deposition Exhibit). cited by applicant

Declaration of Amy K. Wigmore in IPR2023-01358 dated Apr. 16, 2024. cited by applicant

Declaration of Amy K. Wigmore in IPR2023-01359 dated Apr. 16, 2024. cited by applicant

Declaration of Andrew J. Danford in IPR2023-01358 dated Jul. 1, 2024. cited by applicant

Declaration of Daniel O. Griffin, M.D., Ph.D. in IPR2023-01358 dated Aug. 27, 2023. cited by applicant

Declaration of Daniel O. Griffin, M.D., Ph.D. in IPR2023-01359 dated Aug. 27, 2023. cited by applicant

Declaration of Deborah H. Fuller, Ph.D., in IPR2023-01358 dated May 30, 2024 (public redacted version of Ex. 2199). cited by applicant

Declaration of Deborah H. Fuller, Ph.D., in IPR2023-01359 dated May 30, 2024. (public redacted version of Ex. 2199). cited by applicant

Declaration of Douglas L. Sawyer in IPR2023-01358 dated Jul. 17, 2024. cited by applicant

Declaration of Douglas L. Sawyer in IPR2023-01359 dated Jul. 17, 2024. cited by applicant

Declaration of Gabrielle Landes in IPR2023-01358 dated Apr. 25, 2024. cited by applicant

Declaration of Gabrielle Landes in IPR2023-01359 dated Apr. 25, 2024. cited by applicant

Declaration of James E. Malackowski in IPR2023-01358 dated May 29, 2024. cited by applicant

Declaration of James E. Malackowski in IPR2023-01359 dated May 29, 2024. cited by applicant

Declaration of James J. Moon, Ph.D. in IPR2023-01358 dated Apr. 25, 2023. cited by applicant

Declaration of James J. Moon, Ph.D. in IPR2023-01359 dated Apr. 25, 2023. cited by applicant

Declaration of Joshua L. Stern in IPR2023-01358 dated Apr. 17, 2024. cited by applicant

Declaration of Joshua L. Stern in IPR2023-01359 dated Apr. 17, 2024. cited by applicant

Declaration of Kathryn S. Kayali in IPR2023-01358 dated Apr. 16, 2024. cited by applicant

Declaration of Kathryn S. Kayali in IPR2023-01359 dated Apr. 26, 2024. cited by applicant

Declaration of Kevin S. Prussia in IPR2023-01358 dated Apr. 17, 2024. cited by applicant

Declaration of Kevin S. Prussia in IPR2023-01359 dated Apr. 17, 2024. cited by applicant

Declaration of L. Ross Pierce in IPR2023-01358 dated Aug. 30, 2024 (Redacted public version).

cited by applicant
Declaration of L. Ross Pierce in IPR2023-01359 dated Aug. 30, 2024. (Redacted public version).
cited by applicant
Declaration of M. Ryan Meuth in IPR2023-01358 dated Apr. 25, 2024. cited by applicant
Declaration of M. Ryan Meuth in IPR2023-01359 dated Apr. 25, 2024. cited by applicant
Declaration of Philp Krause, M.D., in IPR2023-01359 dated May 28, 2024. (public redacted version of Ex. 2200). cited by applicant
Declaration of Philp Krause, M.D., in IPR2023-01358 dated May 28, 2024. (public redacted version of Ex. 2200). cited by applicant
Declaration of Thomas H. L. Selby in IPR2023-01358 dated Apr. 26, 2024. cited by applicant
Declaration of Thomas H. L. Selby in IPR2023-01359 dated Apr. 26, 2024. cited by applicant
Declaration of W. Christopher Bakewell in IPR2023-01358 dated Aug. 30, 2024. cited by applicant
Declaration of W. Christopher Bakewell in IPR2023-01359 dated Aug. 30, 2024. cited by applicant
Declaration of Warren C.W. Chan, Ph.D. in IPR2023-01358 dated May 29, 2024. cited by applicant
Declaration of Warren C.W. Chan, Ph.D. in IPR2023-01359 dated May 29, 2024. cited by applicant
Defendants' Supplemental Patent-Related Disclosures, Exhibit C—Invalidity Contentions of U.S. Pat. No. 10,702,600 (Jun. 27, 2023)(Excerpts). cited by applicant
Defendants' Patent-Related Disclosures, Exhibit D—Invalidity Contentions of U.S. Pat. No. 10,933,127, 2023, pp. 1-5 (Excerpts). cited by applicant
Defendants' Patent-Related Disclosures, Exhibit C—Invalidity Contentions of U.S. Pat. No. 10,702,600 (Mar. 31, 2023) (Excerpts). cited by applicant
Defendants' Supplemental Patent-Related Disclosures, Exhibit D—Invalidity Contentions of U.S. Pat. No. 10,933,127, 2023, pp. 1-110 (Excerpts). cited by applicant
Deming et al., “Vaccine Efficacy in Senescent Mice Challenged with Recombinant SARS-CoV Bearing Epidemic and Zoonotic Spike Variants,” 2006, vol. 3, Issue 12, PLOS Medicine. cited by applicant
Deposition of Daniel O. Griffin, M.D., Ph.D. (May 22, 2024), in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01358 and -01359. cited by applicant
Deposition of Daniel O. Griffin, M.D., Ph.D. (Sep. 20, 2024) in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01358 and -01359. cited by applicant
Deposition of Deborah Fuller, Ph.D., dated Aug. 2, 2024. cited by applicant
Deposition of James E. Malackowski dated Jul. 22, 2024 and Errata Sheet dated Aug. 27, 2024. cited by applicant
Deposition of James J. Moon, Ph. D. (May 10, 2024), in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01358 and -01359. cited by applicant
Deposition of James J. Moon, Ph.D. (Sep. 13, 2024) in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01358 and -01359. cited by applicant
Deposition of L. Ross Pierce, M.D., (Oct. 4, 2024) in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01358. (redacted version of Ex. 2256). cited by applicant
Deposition of L. Ross Pierce, M.D., (Oct. 4, 2024) in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01359. (redacted version of Ex. 2256). cited by applicant
Deposition of Philip Krause, M.D., dated Aug. 14, 2024 (redacted). cited by applicant
Deposition of W. Christopher Bakewell (Oct. 3, 2024) in *BioNTech SE and Pfizer Inc. v. Moderna TX, Inc.*, in IPR2023-01358 and -01359. cited by applicant
Deposition of Warren C.W. Chan Ph.D., dated Jul. 30, 2024. cited by applicant
Division of Microbiology and Infectious Diseases (DMID) Protocol, Phase 1, Open-Label, Dose-Ranging Study of the Safety and Immunogenicity of 2019-nCoV Vaccine (mRNA-1273) in Healthy Adults, 2020, pp. 1-10 (Excerpts). cited by applicant
Division of Microbiology and Infectious Diseases (DMID) Protocol, “Phase I, Open-Label, Dose-Ranging Study of the Safety and Immunogenicity of 2019-nCoV Vaccine (mRNA-1273) in Healthy

Adults,” 2020, pp. 1-12 (Excerpts). cited by applicant

Division of Microbiology and Infectious Diseases (DMID) Protocol (Feb. 14, 2020); MOD_000477491-MOD_000477561 at MOD_000477491-MOD_000477498; MOD_000477504-MOD_000477509 (Excerpts). cited by applicant

Division of Microbiology and Infectious Diseases (DMID) Protocol (Mar. 30, 2020); MOD_000477134-MOD_000477212 at MOD_000477134-MOD_000477139, MOD_000477148-MOD_000477154 (Excerpts). cited by applicant

Division of Microbiology and Infectious Diseases (DMID) Protocol (May 20, 2020); MOD_000471975-MOD_000472059 at MOD_000471975-MOD_000471983; MOD_000471990-MOD_000471992; MOD_000471994-MOD_000471997 (Excerpts). cited by applicant

Division of Microbiology and Infectious Diseases (DMID) Protocol, “Phase I, Open-Label, Dose-Ranging Study of the Safety and Immunogenicity of 2019-nCoV Vaccine (mRNA-1273) in Healthy Adults,” 2020, pp. 1-8 (Excerpts). cited by applicant

Drummond et al., “The Evolutionary Consequences of Erroneous Protein Synthesis,” 2009, Nature Reviews Genetics (10 Pages). cited by applicant

European Medicines Agency Assessment Report for COVID-19 Vaccine Moderna, 2021, pp. 1-169. cited by applicant

European Medicines Agency, Assessment Report for Spikevax (Sep. 14, 2023). cited by applicant

European Medicines Agency, Nuvaxovid (available at <https://www.ema.europa.eu/en/medicines/human/EPAR/nuvaxovid>). cited by applicant

FDA, E6(R2) Good Clinical Practice: Integrated Addendum to ICH E6(R1) (Mar. 2018). cited by applicant

FDA, Guidance for Industry E6 Good Clinical Practice: Consolidated Guidance (Apr. 1996). cited by applicant

Felgner et al., “Lipofection: A highly efficient, lipid-mediated DNA-transfection procedure,” 1987, 84 Proc. Natl. Acad. Sci. USA (5 Pages). cited by applicant

Fuller, Deborah, The Conversation, “How mRNA and DNA vaccines could soon treat cancers, HIV, autoimmune disorders and genetic diseases,” 2022, pp. 1-5. cited by applicant

Gandhi et al., “Immunization of HIV-1-Infected Persons with Autologous Dendritic Cells Transfected with mRNA Encoding HIV-1 Gag and Nef: Results of a Randomized, Placebo-Controlled Clinical Trial, 71 J,” 2016, Acquired Immune Deficiency Syndrome, Basic Translational Science. cited by applicant

Gao et al., “Nonviral gene delivery: What we know and what is next,” 2007, The AAPS Journal (13 Pages). cited by applicant

Gilleron et al., “Image-Based Analysis of Lipid Nanoparticle-Mediated siRNA Delivery, Intracellular Trafficking and Endosomal Escape,” 2013, Nature Biotech. 638. cited by applicant

Ginn et al., “Gene Therapy Clinical Trials Worldwide to 2017: An Update,” 2018, 20 J, Gene Med, Article e3015. cited by applicant

Gorbalenya, A., E., “The Species Severe Acute Respiratory Syndrome-related Coronavirus: Classifying 2019-nCoV and Naming it SARS-CoV-2, 5,” 2020, Nature Microbiology, pp. 536-544. cited by applicant

Griffin et al., “Co-infection with Hepatitis C Virus Increases Mortality in HIV-1 Infected Patients Through Increased Liver-Related Deaths Rather Than By Increasing Malignancy Related Deaths,” 2016, vol. 7, Issue 4, J. Aids and Clin. Res. (3 Pages). cited by applicant

Griffin et al., “Cytokine storm of a different flavour: the different cytokine signature of SARS-CoV-2 the cause of COVID-19 from the original SARS outbreak,” Journal of Global Antimicrobial Resistance, pp. 1-3. cited by applicant

Griffin et al., “Human B1 cells in umbilical cord and adult peripheral blood express the novel phenotype CD20 CD27 CD43 CD70-,” 2011, vol. 208, Issue 1, J. Exp. Med. (14 Pages). cited by applicant

Griffin et al., A small CD11b human B1 cell subpopulation stimulates T cells and is expanded in lupus, 2011, vol. 208, Issue 13, J. Exp. Med (8 Pages). cited by applicant

Hattinger, E., et al. "Prophylactic mRNA Vaccination against Allergy Confers Long-Term Memory Responses and Persistent Protection in Mice," 2015, Journal of Immunology Research 2015, Article 797421. cited by applicant

Hekele et al., "Rapidly produced SAM? vaccine against H7N9 influenza is immunogenic in mice," 2013, vol. 2, Issue 8, Emerging Microbes and Infections (7 Pages). cited by applicant

Heyes et al., "Cationic lipid saturation influences intracellular delivery of encapsulated nucleic acids," 2005, 107 J .. Controlled Release, pp. 276-287. cited by applicant

Hoerr et al., "In vivo application of RNA leads to induction of specific cytotoxic T lymphocytes and antibodies," 2000, 30 Eur. J. Immunol. cited by applicant

Holtkamp et al., "Modification of antigen-encoding RNA increases stability, translational efficacy, and T-cell stimulatory capacity of dendritic cells," 2006, vol. 108, Issue 13, Gene Therapy. cited by applicant

Hu et al., "Cytosolic delivery mediated via electrostatic surface binding of protein, virus, or siRNA cargos to pH-responsive core-shell gel particles," 2009, vol. 10, Issue 4, Biomacromolecules (10 Pages). cited by applicant

Huang, "In Vivo Delivery of RNAi with Lipid-Based Nanoparticles," 2011, 13 Ann. Rev. Biomed. Eng'g. cited by applicant

International Publication No. WO 2012/006378 A1 (Asserted in *GlaxoSmithKline Biologicals SA et al.v. Pfizer Inc et al.*, C.A. No. 24-512 (GBW) (D. Del.)). cited by applicant

Japanese Package Insert for Spikevax Intramuscular Injection (Monovalent: Omicron XBB. 1.5). cited by applicant

Jaume et al., "Anti-Severe Acute Respiratory Syndrome Coronavirus Spike Antibodies Trigger Infection of Human Immune Cells via a pH- and Cysteine Protease-Independent FcγR Pathway," 2011, Journal of Virology, pp. 10582-10597. cited by applicant

Jeffs et al., "A Scalable, Extrusion-Free Method for Efficient Liposomal Encapsulation of Plasmid DNA," 2005, 22 Pharmaceutical Research (11 Pages). cited by applicant

Jesse H. Erasmus et al., "An Alphavirus-derived replicon RNA vaccine induces SARS-CoV-2 neutralizing antibody and T cell responses in mice and nonhuman primates," 2020, Sci. Transl. Med. (11 Pages). cited by applicant

Johansson et al., "Intradermal Electroporation of Naked Replicon RNA Elicits Strong Immune Responses," 2012, PLoS ONE, Article e29732. cited by applicant

Kam et al., "Antibodies against trimeric S glycoprotein protect hamsters against SARS-CoV challenge despite their capacity to mediate FcγRII-dependent entry into B cells in vitro," 2007, Vaccine (13 Pages). cited by applicant

Kariko et al., "Modified Uridines Are the Key to a Successful Message," 2021, Nature Reviews Immunology (2 Pages). cited by applicant

Kariko et al., "Naturally occurring nucleoside modifications suppress the immunostimulatory activity of RNA: Implication for therapeutic RNA development," Curr. Opin. Drug Discov. Devel., pp. 523-532. cited by applicant

Kariko et al., "Suppression of RNA recognition by Toll-like receptors: The impact of nucleoside modification and the evolutionary origin of RNA," 2005, Immunity (8 Pages). cited by applicant

Kariko et al., Increased erythropoiesis in mice injected with submicrogram quantities of pseudouridine-containing mRNA encoding erythropoietin, 20 Molecular Therapy 948 (2012) (Ex. 1028). cited by applicant

Kim et al., "Generation and Characterization of DNA Vaccines Targeting the Nucleocapsid Protein of Severe Acute Respiratory Syndrome Coronavirus," 2004, vol. 78 J, Issue 9, Virology, pp. 4638-4645. cited by applicant

King et al., "Virus Taxonomy: Classification and Nomenclature of Viruses," 2012, Elsevier

Academic Press (50 Pages). cited by applicant

Kranz et al., "Systemic RNA Delivery to Dendritic Cells Exploits Antiviral Defense for Cancer Immunotherapy," 2016, *Nature*, vol. 534, pp. 396-411. cited by applicant

Kreiter et al., "Intranodal Vaccination with Naked Antigen-Encoding RNA Elicits Potent Prophylactic and Therapeutic Antitumoral Immunity," 2010, *Cancer Research*, pp. 9301-9040. cited by applicant

Kutzler et al., "DNA Vaccines: Ready for Prime Time?," 2008, *Nature Reviews Genetics*, vol. 9, pp. 776-788. cited by applicant

Langer, R., "New methods of drug delivery," 1990, *Science*, 1527-1533. cited by applicant

Leal et al., "Phase I clinical trial of an intranodally administered mRNA-based therapeutic vaccine against HIV-1 infection," 2018, *AIDS*, vol. 32, Issue 17, Pages. cited by applicant

Leung et al., "Chapter Four—Lipid nanoparticles for short interfering RNA delivery," 2014, in *Advances in Genetic*, vol. 88 (40 Pages). cited by applicant

Liang et al., "Efficient Targeting and Activation of Antigen-Presenting Cells In Vivo after Modified mRNA Vaccine Administration in Rhesus Macaques," 2017, *25 Mol Therapy*, pp. 2635-2647. cited by applicant

Lingshu Wang et al., "Evaluation of candidate vaccine approaches for MERS-CoV," 2015, *Nature Communications* (11 Pages). cited by applicant

Liu et al., "The Membrane Protein of Severe Acute Respiratory Syndrome Coronavirus Acts as a Dominant Immunogen Revealed by a Clustering Region of Novel Functionally and Structurally Defined Cytotoxic T-Lymphocyte Epitopes," 2010, vol. 202 *J, Infectious Diseases*, pp. 1171-1180. cited by applicant

Lu et al., "Immunogenicity of DNA Vaccines Expressing Human Immunodeficiency Virus Type 1 Envelope Glycoprotein With and Without Deletions in the V1/2 and V3 Regions," *AIDS Research & Human Retroviruses*, pp. 151-155. cited by applicant

Marshall et al., "Ch. 79: Regulation and Testing of Vaccines," 2018, *Plotkin's Vaccines*, 7th edition, pp. 1547-1565. cited by applicant

Martin et al., "A SARS DNA Vaccine Induces Neutralizing Antibody and Cellular Immune Responses in Healthy Adults in a Phase I Clinical Trial," 2008, *Vaccine* (6 Pages). cited by applicant

Matthew G. Stanton & Kerry E. Murphy-Benenato, "Messenger RNA as a Novel Therapeutic Approach," 2017, *Topics in Medicinal Chemistry RNA Therapeutics*, pp. 1-2 (Excerpts). cited by applicant

Moderna Form 10-K for the fiscal year ended Dec. 31, 2018. cited by applicant

Moderna Form 10-K for the fiscal year ended Dec. 31, 2022. cited by applicant

Moderna Form 10-Q for the quarterly period ended Jun. 30, 2024. cited by applicant

Moderna Form 8-K, Mar. 16, 2020. cited by applicant

Moderna, Inc. 2020 Form 10-K. cited by applicant

Moderna, Inc. 2023 Form 10-K. cited by applicant

Moderna, Inc., Patents, <https://www.modernatx.com/patents> (3 Pages). cited by applicant

Moderna, Science & Technology Day dated May 17, 2022. cited by applicant

Moderna, Vaccines and Related Biological Products Advisory Committee Presentation: Moderna COVID-19 Variant Vaccines (Jun. 15, 2023); MOD_000767652-MOD_000767686. cited by applicant

Nakayama et al., "Comparison of Current Regulatory Status for Gene-Based Vaccines in the U.S., Europe and Japan," 2015, *Vaccines*, pp. 186-202. cited by applicant

National Human Genome Research Institute, "Double Helix," 2023, NIH, <https://www.genome.gov/geneticsglossary/Double-Helix> (2 Pages). cited by applicant

National Human Genome Research Institute, Central Dogma, 2023, NIH, <https://www.genome.gov/geneticsglossary/Central-Dogma> (4 Pages). cited by applicant

ONPATRO Prescribing Label (available at

https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/210922s000lbl.pdf). cited by applicant
Openshaw et al., “Immune Responses and Disease Enhancement During Respiratory Syncytial Virus Infection,” 2005, *Clinical Microbiology Reviews*, vol. 18, Issue 13, pp. 541-555. cited by applicant

Package Insert for Spikevax (COVID-19 Vaccine, mRNA) – 2023-2024 Formula (45 Pages). cited by applicant

Package Insert for Spikevax (COVID-19 Vaccine, mRNA) – 2023- 2024 Formula (Apr. 2024). cited by applicant

Paoletti, E., “Applications of Pox Virus Vectors to Vaccination: An Update,” 1996, *Proceedings of the Natl. Acad. of Sci.*, pp. 11349-11353. cited by applicant

Paramasivam et al., “Endosomal Escape of Delivered mRNA from Endosomal Recycling Tubules Visualized at the Nanoscale,” 221 *J. Cell Biology* e202110137. cited by applicant

Park et al., “Non-Viral COVID-19 Vaccine Delivery Systems,” *Advanced Drug Delivery Reviews*, pp. 137-151. cited by applicant

Pascolo, S., “Vaccination with Messenger RNA (mRNA),” 2008, *Handbook of Experimental Pharmacology*, pp. 221-235. cited by applicant

Patent Sublicense Agreement Between Cellscript, LLC and ModernaTx, Inc. cited by applicant

Plotkin, S., A., “Vaccines: Past, Present and Future, 11 *Nature Medicine Supplement* 55,” 2005, *Historical Perspective, Nature Medicine*, pp. S5-S11. cited by applicant

Regla-Nava et al., “Severe Acute Respiratory Syndrome Coronaviruses with Mutations in the E Protein Are Attenuated and Promising Vaccine Candidates,” 2015, vol. 89, *Journal of Virology* (18 Pages). cited by applicant

Reply Declaration of Daniel Griffin in IPR2023-01358 dated Aug. 30, 2024. (Redacted public version). cited by applicant

Reply Declaration of Daniel Griffin in IPR2023-01359 dated Aug. 30, 2024. (Redacted public version). cited by applicant

Reply Declaration of James Moon in IPR2023-01358 dated Aug. 30, 2024. (Redacted public version). cited by applicant

Reply Declaration of James Moon in IPR2023-01359 dated Aug. 30, 2024. (Redacted public version). cited by applicant

Sahdev et al., *Biomaterials for Nanoparticle Vaccine Delivery Systems, Pharmaceutical Research*, 2014, pp. 2563-2582. cited by applicant

Sahin et al., “mRNA-based therapeutics Developing a new class of drugs,” 2014, *Nature Reviews Drug Discovery*. cited by applicant

Semple et al., “Efficient encapsulation of antisense oligonucleotides in lipid vesicles using ionizable aminolipids: Formation of novel small multilamellar vesicle structures,” 2001, *Biochimica Et Biophysica Acta*, pp. 152-166. cited by applicant

Semple et al., “Rational design of cationic lipids for siRNA delivery,” 2010, *Nature Biotechnology*, pp. 172-178. cited by applicant

Shaw et al., “The Path to an RSV Vaccine,” 2013, *Current Opinion in Virology*, pp. 332-342. cited by applicant

Slimko et al., “Codon optimization of *Caenorhabditis elegans* GluCI ion channel genes for mammalian cells dramatically improves expression levels,” 2003, *Journal of Neuroscience Methods*. cited by applicant

Song et al., “Middle East Respiratory Syndrome Coronavirus Spike Protein Delivered by Modified Vaccinia Virus Ankara Efficiently Induces Virus-Neutralizing Antibodies,” 2013, 87 *J. of Virology*, pp. 1-5. cited by applicant

Steitz et al., “Effective Induction of Anti-Melanoma Immunity Following Genetic Vaccination with Synthetic mRNA Coding for the Fusion Protein EGFP. TRP2,” 2006, *Cancer Immunology*, pp. 246-

253. cited by applicant

Stenler et al., "Safety and efficacy of DNA vaccines: Plasmids vs. minicircles," 2014, vol. 10, Issue 5, Human Vaccines & Immunotherapeutics. cited by applicant

Tenchov et al., "Lipid nanoparticles 2014 From liposomes to mRNA vaccine delivery, a landscape of research diversity and advancement," 2021, ACS NANO (34 Pages). cited by applicant

Updated Curriculum Vitae of Daniel O. Griffin, Ph.D. cited by applicant

Vennema et al., "Early Death After Feline Infectious Peritonitis Virus Challenge due to Recombinant Vaccinia Virus Immunization," 1990, Journal of Virology, pp. 1407-1409. cited by applicant

Vergati et al., "Strategies for Cancer Vaccine Development," 2010, Journal of Biomedicine and Biotechnology, (13 Pages). cited by applicant

Wan et al., "Comparison of Immunogenicity Between Codon Optimized HIV-1 Thailand Subtype B gp140 and gp145 Vaccines," 2007, Vaccine, vol. 25, pp. 4949-4959. cited by applicant

Weide et al., "Direct Injection of Protamine-protected mRNA: Results of a Phase 1/2 Vaccination Trial in Metastatic Melanoma Patients," 2009, Immunotherapy (11 Pages). cited by applicant

Weide et al., "Results of the First Phase I/II Clinical Vaccination Trial with Direct Injection of mRNA," 2008, J. Immunotherapy, pp. 180-188. cited by applicant

Withoff et al., "Replication-defective Recombinant Semliki Forest Virus Encoding GM-CSF as a Vector System for Rapid and Facile Generation of Autologous Human Tumor Cell Vaccines," 2001, Gene Therapy, pp. 1515-1523. cited by applicant

Wolff et al., Direct gene transfer into mouse muscle in vivo, 247 Science 1465 (1990) (Ex. 1013). cited by applicant

Yang et al., "Evasion of Antibody Neutralization in Emerging Severe Acute Respiratory Syndrome Coronaviruses," 2005, PNAS, vol. 102, Issue 3, pp. 797-801. cited by applicant

Yoo et al., Synthesis and Processing of the Haemagglutinin-Esterase Glycoprotein of Bovine Coronavirus Encoded in the E3 Region of Adenovirus, vol. 73, J. Gen. Virology, pp. 2591-2600. cited by applicant

Zhou et al., "Current RNA-Based Therapeutics in Clinical Trials," 2019, Current Gene Therapy, pp. 172-196. cited by applicant

"Final Written Decision" IPR2023-01358 dated Mar. 5, 2025. cited by applicant

"Final Written Decision" IPR2023-01359 dated Mar. 5, 2025. cited by applicant

Alberer et al., "Safety and Immunogenicity of a mRNA Rabies Vaccine in Healthy Adults: an Open-label, Nonrandomised, Prospective, First-in-human Phase 1 Clinical Trial," 2017, The Lancet, pp. 1-10. cited by applicant

Berkeley Lovelace Jr. & Noah Higgins-Dunn, Dr. Anthony Fauci says chance of coronavirus vaccine being highly effective is 'not great', CNBC.com (last updated Aug. 8, 2020), <https://www.cnbc.com/2020/08/07/coronavirus-vaccine-dr-fauci-says-chances-of-it-being-highly-effective-is-not-great.html>. cited by applicant

Fraiser Kansteiner, It's the end of the line for J&J's COVID shot in the US, CDC says, Fiercepharma (May 16, 2023). cited by applicant

Karin Bok, Accelerated COVID-19 vaccine development: milestones, lessons, and prospects, 54 Immunity 1636 (2021). cited by applicant

Liu, Agnus, AstraZeneca withdraws US COVID vaccine application, shifts focus to antibody treatments, Fierce Pharma (Nov. 10, 2022). cited by applicant

Monika Kumari et al., A critical overview of current progress for COVID-19: development of vaccines, antiviral drugs, and therapeutic antibodies, J. Biomed Sci 29(68) (2022). cited by applicant

Prescribing Information for Comirnaty (Pfizer). cited by applicant

Prescribing Information for Spikevax (Moderna). cited by applicant

Press Release, Novavax, U.S. FDA Grants Emergency Use Authorization for Novavax COVID-19

Vaccine, Adjuvanted for Individuals Aged 18 and Over (Jul. 13, 2022). cited by applicant
Walter et al., “Evaluation of the BNT162b2 COVID-19 Vaccine in Children 5 to 11 Years of Age,”
New England J. Med., 2022; 386: 35-46 (published online Nov. 9, 2021). cited by applicant

Primary Examiner: White; Nicole Kinsey

Attorney, Agent or Firm: Foley & Lardner LLP

Background/Summary

RELATED APPLICATIONS (1) This application is a continuation of U.S. application Ser. No. 16/897,734, now U.S. Pat. No. 11,872,278, filed Jun. 10, 2020, which is a continuation of U.S. application Ser. No. 16/368,270, now U.S. Pat. No. 10,702,599, filed Mar. 28, 2019, which is a continuation of U.S. application Ser. No. 16/040,981, now U.S. Pat. No. 10,272,150, filed Jul. 20, 2018, which is a continuation of U.S. application Ser. No. 15/674,599, now U.S. Pat. No. 10,064,934, filed Aug. 11, 2017, which is a continuation of international application number PCT/US2016/058327, which claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional application No. 62/244,802, filed Oct. 22, 2015, U.S. provisional application No. 62/247,297, filed Oct. 28, 2015, U.S. provisional application No. 62/244,946, filed Oct. 22, 2015, U.S. provisional application No. 62/247,362, filed Oct. 28, 2015, U.S. provisional application No. 62/244,813, filed Oct. 22, 2015, U.S. provisional application No. 62/247,394, filed Oct. 28, 2015, U.S. provisional application No. 62/244,837, filed Oct. 22, 2015, U.S. provisional application No. 62/247,483, filed Oct. 28, 2015, and U.S. provisional application No. 62/245,031, filed Oct. 22, 2015, each of which is incorporated by reference herein in its entirety.

REFERENCE TO AN ELECTRONIC SEQUENCE LISTING

(1) The contents of the electronic sequence listing (M137870053US11-SEQ-JXV.xml; Size: 387,115 bytes; and Date of Creation: Nov. 29, 2023) is herein incorporated by reference in its entirety.

BACKGROUND

(2) Respiratory disease is a medical term that encompasses pathological conditions affecting the organs and tissues that make gas exchange possible in higher organisms, and includes conditions of the upper respiratory tract, trachea, bronchi, bronchioles, alveoli, pleura and pleural cavity, and the nerves and muscles of breathing. Respiratory diseases range from mild and self-limiting, such as the common cold, to life-threatening entities like bacterial pneumonia, pulmonary embolism, acute asthma and lung cancer. Respiratory disease is a common and significant cause of illness and death around the world. In the US, approximately 1 billion “common colds” occur each year. Respiratory conditions are among the most frequent reasons for hospital stays among children.

(3) The human metapneumovirus (hMPV) is a negative-sense, single-stranded RNA virus of the genus Pneumovirinae and of the family Paramyxoviridae and is closely related to the avian metapneumovirus (AMPV) subgroup C. It was isolated for the first time in 2001 in the Netherlands by using the RAP-PCR (RNA arbitrarily primed PCR) technique for identification of unknown viruses growing in cultured cells. hPMV is second only to RSV as an important cause of viral lower respiratory tract illness (LRI) in young children. The seasonal epidemiology of hMPV appears to be similar to that of RSV, but the incidence of infection and illness appears to be substantially lower.

(4) Parainfluenza virus type 3 (PIV3), like hMPV, is also a negative-sense, single-stranded sense RNA virus of the genus Pneumovirinae and of the family Paramyxoviridae and is a major cause of ubiquitous acute respiratory infections of infancy and early childhood. Its incidence peaks around

4-12 months of age, and the virus is responsible for 3-10% of hospitalizations, mainly for bronchiolitis and pneumonia. PIV3 can be fatal, and in some instances is associated with neurologic diseases, such as febrile seizures. It can also result in airway remodeling, a significant cause of morbidity. In developing regions of the world, infants and young children are at the highest risk of mortality, either from primary PIV3 viral infection or a secondary consequences, such as bacterial infections. Human parainfluenza viruses (hPIV) types 1, 2 and 3 (hPIV1, hPIV2 and hPIV3, respectively), also like hMPV, are second only to RSV as important causes of viral LRI in young children.

(5) RSV, too, is a negative-sense, single-stranded RNA virus of the genus Pneumovirinae and of the family Paramyxoviridae. Symptoms in adults typically resemble a sinus infection or the common cold, although the infection may be asymptomatic. In older adults (e.g., >60 years), RSV infection may progress to bronchiolitis or pneumonia. Symptoms in children are often more severe, including bronchiolitis and pneumonia. It is estimated that in the United States, most children are infected with RSV by the age of three. The RSV virion consists of an internal nucleocapsid comprised of the viral RNA bound to nucleoprotein (N), phosphoprotein (P), and large polymerase protein (L). The nucleocapsid is surrounded by matrix protein (M) and is encapsulated by a lipid bilayer into which the viral fusion (F) and attachment (G) proteins as well as the small hydrophobic protein (SH) are incorporated. The viral genome also encodes two nonstructural proteins (NS1 and NS2), which inhibit type I interferon activity as well as the M-2 protein.

(6) The continuing health problems associated with hMPV, PIV3 and RSV are of concern internationally, reinforcing the importance of developing effective and safe vaccine candidates against these virus.

(7) Despite decades of research, no vaccines currently exist (Sato and Wright, *Pediatr. Infect. Dis. J.* 2008; 27 (10 Suppl): S123-5). Recombinant technology, however, has been used to target the formation of vaccines for hPIV-1, 2 and 3 serotypes, for example, and has taken the form of several live-attenuated intranasal vaccines. Two vaccines in particular were found to be immunogenic and well tolerated against hPIV-3 in phase I trials. hPIV1 and hPIV2 vaccine candidates remain less advanced (Durbin and Karron, *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 2003; 37(12):1668-77).

(8) Measles virus (MeV), like hMPV, PIV3 and RSV, is a negative-sense, single-stranded RNA virus that is the cause of measles, an infection of the respiratory system. MeV is of the genus Morbillivirus within the family Paramyxoviridae. Humans are the natural hosts of the virus; no animal reservoirs are known to exist. Symptoms of measles include fever, cough, runny nose, red eyes and a generalized, maculopapular, erythematous rash. The virus is highly contagious and is spread by coughing

(9) In addition to hMPV, PIV, RSV and MeV, betacoronaviruses are known to cause respiratory illnesses. Betacoronaviruses (BetaCoVs) are one of four genera of coronaviruses of the subfamily Coronavirinae in the family Coronaviridae, of the order Nidovirales. They are enveloped, positive-sense, single-stranded RNA viruses of zoonotic origin. The coronavirus genera are each composed of varying viral lineages, with the betacoronavirus genus containing four such lineages. The BetaCoVs of the greatest clinical importance concerning humans are OC43 and HKU1 of the A lineage, SARS-CoV of the B lineage, and MERS-CoV of the C lineage. MERS-CoV is the first betacoronavirus belonging to lineage C that is known to infect humans.

(10) The Middle East respiratory syndrome coronavirus (MERS-CoV), or EMC/2012 (HCoV-EMC/2012), initially referred to as novel coronavirus 2012 or simply novel coronavirus, was first reported in 2012 after genome sequencing of a virus isolated from sputum samples from a person who fell ill during a 2012 outbreak of a new flu. As of July 2015, MERS-CoV cases have been reported in over 21 countries. The outbreaks of MERS-CoV have raised serious concerns worldwide, reinforcing the importance of developing effective and safe vaccine candidates against MERS-CoV.

(11) Severe acute respiratory syndrome (SARS) emerged in China in 2002 and spread to other countries before brought under control. Because of a concern for reemergence or a deliberate release of the SARS coronavirus, vaccine development was initiated.

(12) Deoxyribonucleic acid (DNA) vaccination is one technique used to stimulate humoral and cellular immune responses to foreign antigens, such as hMPV antigens and/or PIV antigens and/or RSV antigens. The direct injection of genetically engineered DNA (e.g., naked plasmid DNA) into a living host results in a small number of its cells directly producing an antigen, resulting in a protective immunological response. With this technique, however, comes potential problems, including the possibility of insertional mutagenesis, which could lead to the activation of oncogenes or the inhibition of tumor suppressor genes.

SUMMARY

(13) Provided herein are ribonucleic acid (RNA) vaccines that build on the knowledge that RNA (e.g., messenger RNA (mRNA)) can safely direct the body's cellular machinery to produce nearly any protein of interest, from native proteins to antibodies and other entirely novel protein constructs that can have therapeutic activity inside and outside of cells. The RNA (e.g., mRNA) vaccines of the present disclosure may be used to induce a balanced immune response against hMPV, PIV, RSV, MeV, and/or BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1), or any combination of two or more of the foregoing viruses, comprising both cellular and humoral immunity, without risking the possibility of insertional mutagenesis, for example. hMPV, PIV, RSV, MeV, BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1) and combinations thereof are referred to herein as “respiratory viruses.” Thus, the term “respiratory virus RNA vaccines” encompasses hMPV RNA vaccines, PIV RNA vaccines, RSV RNA vaccines, MeV RNA vaccines, BetaCoV RNA vaccines, and any combination of two or more of hMPV RNA vaccines, PIV RNA vaccines, RSV RNA vaccines, MeV RNA vaccines, and BetaCoV RNA vaccines.

(14) The RNA (e.g., mRNA) vaccines may be utilized in various settings depending on the prevalence of the infection or the degree or level of unmet medical need. The RNA (e.g., mRNA) vaccines may be utilized to treat and/or prevent a hMPV, PIV, RSV, MeV, a BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV—HKU1), or any combination of two or more of the foregoing viruses, of various genotypes, strains, and isolates. The RNA (e.g., mRNA) vaccines have superior properties in that they produce much larger antibody titers and produce responses earlier than commercially available anti-viral therapeutic treatments. While not wishing to be bound by theory, it is believed that the RNA (e.g., mRNA) vaccines, as mRNA polynucleotides, are better designed to produce the appropriate protein conformation upon translation as the RNA (e.g., mRNA) vaccines co-opt natural cellular machinery. Unlike traditional vaccines, which are manufactured ex vivo and may trigger unwanted cellular responses, RNA (e.g., mRNA) vaccines are presented to the cellular system in a more native fashion.

(15) In some aspects the invention is a respiratory virus vaccine, comprising at least one RNA polynucleotide having an open reading frame encoding at least one respiratory virus antigenic polypeptide, formulated in a cationic lipid nanoparticle.

(16) Surprisingly, in some aspects, it has also been shown that efficacy of mRNA vaccines can be significantly enhanced when combined with a flagellin adjuvant, in particular, when one or more antigen-encoding mRNAs is combined with an mRNA encoding flagellin.

(17) RNA (e.g., mRNA) vaccines combined with the flagellin adjuvant (e.g., mRNA-encoded flagellin adjuvant) have superior properties in that they may produce much larger antibody titers and produce responses earlier than commercially available vaccine formulations. While not wishing to be bound by theory, it is believed that the RNA (e.g., mRNA) vaccines, for example, as mRNA polynucleotides, are better designed to produce the appropriate protein conformation upon

translation, for both the antigen and the adjuvant, as the RNA (e.g., mRNA) vaccines co-opt natural cellular machinery. Unlike traditional vaccines, which are manufactured ex vivo and may trigger unwanted cellular responses, RNA (e.g., mRNA) vaccines are presented to the cellular system in a more native fashion.

(18) Some embodiments of the present disclosure provide RNA (e.g., mRNA) vaccines that include at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one antigenic polypeptide or an immunogenic fragment thereof (e.g., an immunogenic fragment capable of inducing an immune response to the antigenic polypeptide) and at least one RNA (e.g., mRNA polynucleotide) having an open reading frame encoding a flagellin adjuvant.

(19) In some embodiments, at least one flagellin polypeptide (e.g., encoded flagellin polypeptide) is a flagellin protein. In some embodiments, at least one flagellin polypeptide (e.g., encoded flagellin polypeptide) is an immunogenic flagellin fragment. In some embodiments, at least one flagellin polypeptide and at least one antigenic polypeptide are encoded by a single RNA (e.g., mRNA) polynucleotide. In other embodiments, at least one flagellin polypeptide and at least one antigenic polypeptide are each encoded by a different RNA polynucleotide.

(20) In some embodiments at least one flagellin polypeptide has at least 80%, at least 85%, at least 90%, or at least 95% identity to a flagellin polypeptide having a sequence identified by any one of SEQ ID NO: 54-56.

(21) Provided herein, in some embodiments, is a ribonucleic acid (RNA) (e.g., mRNA) vaccine, comprising at least one (e.g., at least 2, 3, 4 or 5) RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one (e.g., at least 2, 3, 4 or 5) hMPV, PIV, RSV, MeV, or a BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV—HKU1) antigenic polypeptide, or any combination of two or more of the foregoing antigenic polypeptides. Herein, use of the term “antigenic polypeptide” encompasses immunogenic fragments of the antigenic polypeptide (an immunogenic fragment that induces (or is capable of inducing) an immune response to hMPV, PIV, RSV, MeV, or a BetaCoV), unless otherwise stated.

(22) Also provided herein, in some embodiments, is a RNA (e.g., mRNA) vaccine comprising at least one (e.g., at least 2, 3, 4 or 5) RNA polynucleotide having an open reading frame encoding at least one (e.g., at least 2, 3, 4 or 5) hMPV, PIV, RSV, MeV, and/or a BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV—HKU1) antigenic polypeptide or an immunogenic fragment thereof, linked to a signal peptide.

(23) Further provided herein, in some embodiments, is a nucleic acid (e.g., DNA) encoding at least one (e.g., at least 2, 3, 4 or 5) hMPV, PIV, RSV, MeV, and/or a BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV—HKU1) RNA (e.g., mRNA) polynucleotide.

(24) Further still, provided herein, in some embodiments, is a method of inducing an immune response in a subject, the method comprising administering to the subject a vaccine comprising at least one (e.g., at least 2, 3, 4 or 5) RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one (e.g., at least 2, 3, 4 or 5) hMPV, PIV, RSV, MeV, and/or a BetaCoV (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV—HKU1) antigenic polypeptide, or any combination of two or more of the foregoing antigenic polypeptides.

(25) hMPV/PIV3/RSV

(26) In some embodiments, a RNA (e.g., mRNA) vaccine comprises at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one hMPV, PIV3 or RSV antigenic polypeptide. In some embodiments, at least one antigenic polypeptide is a hMPV, PIV3 or RSV polyprotein. In some embodiments, at least one antigenic polypeptide is major surface glycoprotein G or an immunogenic fragment thereof. In some embodiments, at least one antigenic polypeptide is Fusion (F) glycoprotein (e.g., Fusion glycoprotein F0, F1 or F2) or an immunogenic

fragment thereof. In some embodiments, at least one antigenic polypeptide is major surface glycoprotein G or an immunogenic fragment thereof and F glycoprotein or an immunogenic fragment thereof. In some embodiments, the antigenic polypeptide is nucleoprotein (N) or an immunogenic fragment thereof, phosphoprotein (P) or an immunogenic fragment thereof, large polymerase protein (L) or an immunogenic fragment thereof, matrix protein (M) or an immunogenic fragment thereof, small hydrophobic protein (SH) or an immunogenic fragment thereof nonstructural protein 1 (NS1) or an immunogenic fragment thereof, or nonstructural protein 2 (NS2) and an immunogenic fragment thereof.

(27) In some embodiments, at least one hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 (Table 3; see also amino acid sequences of Table 4). In some embodiments, the amino acid sequence of the hMPV antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino acid sequence identified by any one of SEQ ID NO: 5-8 (Table 3; see also amino acid sequences of Table 4).

(28) In some embodiments, at least one hMPV antigenic polypeptide is encoded by a nucleic acid sequence identified by any one of SEQ ID NO: 1-4 (Table 2).

(29) In some embodiments, at least one hMPV RNA (e.g., mRNA) polynucleotide is encoded by a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 1-4 (Table 2). In some embodiments, at least one hMPV RNA (e.g., mRNA) polynucleotide comprises a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 57-60 (Table 2).

(30) In some embodiments, at least one antigenic polypeptide is obtained from hMPV strain CAN98-75 (CAN75) or the hMPV strain CAN97-83 (CAN83).

(31) In some embodiments, at least one PIV3 antigenic polypeptide comprises hemagglutinin-neuraminidase, Fusion (F) glycoprotein, matrix protein (M), nucleocapsid protein (N), viral replicase (L), non-structural V protein, or an immunogenic fragment thereof.

(32) In some embodiments, at least one PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 (Table 6; see also amino acid sequences of Table 7). In some embodiments, the amino acid sequence of the PIV3 antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino acid sequence identified by any one of SEQ ID NO: 12-13 (Table 6; see also amino acid sequences of Table 7).

(33) In some embodiments, at least one PIV3 antigenic polypeptide is encoded by a nucleic acid sequence identified by any one of SEQ ID NO: 9-12 (Table 5; see also nucleic acid sequences of Table 7).

(34) In some embodiments, at least one PIV3 RNA (e.g., mRNA) polynucleotide is encoded by a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 9-12 (Table 5; see also nucleic acid sequences of Table 7). In some embodiments, at least one PIV3 RNA (e.g., mRNA) polynucleotide comprises a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 61-64 (Table 5).

(35) In some embodiments, at least one antigenic polypeptide is obtained from PIV3 strain HPIV3/*Homo sapiens*/PER/FLA4815/2008.

(36) In some embodiments, at least one RSV antigenic polypeptide comprises at least one antigenic polypeptide that comprises glycoprotein G, glycoprotein F, or an immunogenic fragment thereof. In some embodiments, at least one RSV antigenic polypeptide comprises at least one antigenic polypeptide that comprises glycoprotein F and at least one or at least two antigenic polypeptide selected from G, M, N, P, L, SH, M2, NS1 and NS2.

(37) MeV

(38) In some embodiments, a RNA (e.g., mRNA) vaccine comprises at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one MeV antigenic

polypeptide. In some embodiments, at least one antigenic polypeptide is a hemagglutinin (HA) protein or an immunogenic fragment thereof. The HA protein may be from MeV strain D3 or B8, for example. In some embodiments, at least one antigenic polypeptide is a Fusion (F) protein or an immunogenic fragment thereof. The F protein may be from MeV strain D3 or B8, for example. In some embodiments, a MeV RNA (e.g., mRNA) vaccine comprises at least one RNA polynucleotide encoding a HA protein and a F protein. The HA and F proteins may be from MeV strain D3 or B8, for example.

(39) In some embodiments, at least one MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 (Table 14). In some embodiments, the amino acid sequence of the MeV antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino acid sequence identified by any one of SEQ ID NO: 47-50 (Table 14).

(40) In some embodiments, at least one MeV antigenic polypeptide is encoded by a nucleic acid sequence of SEQ ID NO: 35-46 (Table 13).

(41) In some embodiments, at least one MeV RNA (e.g., mRNA) polynucleotide is encoded by a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 35-46 (Table 13). In some embodiments, at least one MeV RNA (e.g., mRNA) polynucleotide comprises a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 69-80 (Table 13).

(42) In some embodiments, at least one antigenic polypeptide is obtained from MeV strain B3/B3.1, C2, D4, D6, D7, D8, G3, H1, Moraten, Rubeovax, MVi/New Jersey.USA/45.05, MVi/Texas.USA/4.07, AIK-C, MVi/New York.USA/26.09/3, MVi/California.USA/16.03, MVi/Virginia.USA/15.09, MVi/California.USA/8.04, or MVi/Pennsylvania.USA/20.09.

(43) BetaCoV

(44) In some embodiments, a RNA (e.g., mRNA) vaccine comprises at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one BetaCoV antigenic polypeptide. In some embodiments, the BetaCoV is MERS-CoV. In some embodiments, the BetaCoV is SARS-CoV. In some embodiments, the BetaCoV is HCoV—OC43. In some embodiments, the BetaCoV is HCoV-229E. In some embodiments, the BetaCoV is HCoV-NL63. In some embodiments, the BetaCoV is HCoV—HKU1. In some embodiments, at least one antigenic polypeptide is a betacoronavirus structural protein. For example, a betacoronavirus structural protein may be spike protein (S), envelope protein (E), nucleocapsid protein (N), membrane protein (M) or an immunogenic fragment thereof. In some embodiments, a betacoronavirus structural protein is a spike protein (S). In some embodiments, a betacoronavirus structural protein is a S1 subunit or a S2 subunit of spike protein (S) or an immunogenic fragment thereof.

(45) BetaCoV RNA (e.g., mRNA) polynucleotides of the vaccines provided herein may encode viral protein components of betacoronaviruses, for example, accessory proteins, replicase proteins and the like are encompassed by the present disclosure. RNA (e.g., mRNA) vaccines may include RNA polynucleotides encoding at least one accessory protein (e.g., protein 3, protein 4a, protein 4b, protein 5), at least one replicase protein (e.g., protein 1a, protein 1b), or a combination of at least one accessory protein and at least one replicase protein. The present disclosure also encompasses RNA (e.g., mRNA) vaccines comprising RNA (e.g., mRNA) polynucleotides encoding an accessory protein and/or a replicase protein in combination with at least one structural protein. Due to their surface expression properties, vaccines featuring RNA polynucleotides encoding structural proteins are believed to have preferred immunogenic activity and, hence, may be most suitable for use in the vaccines of the present disclosure.

(46) Some embodiments of the present disclosure provide betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV-HKU1 or a combination thereof) vaccines that include at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV

—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH, HCoV—HKU1) antigenic polypeptide. Also provided herein are pan-betacoronavirus vaccines. Thus, a betacoronavirus vaccine comprising a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding any one, two, three or four of MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, and HCoV—HKU1, for example, may be effective against any one of, any combination of, or all of, MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1. Other betacoronaviruses are encompassed by the present disclosure.

(47) In some embodiments, at least one antigenic polypeptide is a MERS-CoV structural protein. For example, a MERS-CoV structural protein may be spike protein (S), envelope protein (E), nucleocapsid protein (N), membrane protein (M) or an immunogenic fragment thereof. In some embodiments, the MERS-CoV structural protein is a spike protein (S) (see, e.g., Coleman C M et al. *Vaccine* 2014; 32:3169-74, incorporated herein by reference). In some embodiments, the MERS-CoV structural protein is a S1 subunit or a S2 subunit of spike protein (S) or an immunogenic fragment thereof (Li J et al. *Viral Immunol* 2013; 26(2):126-32; He Y et al. *Biochem Biophys Res Commun* 2004; 324(2):773-81, each of which is incorporated herein by reference).

(48) In some embodiments, at least one MERS-CoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-28 or 33 (Table 11). In some embodiments, the amino acid sequence of the MERS-CoV antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino acid sequence identified by any one of SEQ ID NO: 24-28 or 33 (Table 11).

(49) In some embodiments, at least one MERS-CoV antigenic polypeptide is encoded by a nucleic acid sequence identified by any one of SEQ ID NO: 20-23 (Table 10).

(50) In some embodiments, at least one MERS-CoV RNA (e.g., mRNA) polynucleotide is encoded by a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 20-23 (Table 10). In some embodiments, at least one MERS-CoV RNA (e.g., mRNA) polynucleotide comprises a nucleic acid sequence, or a fragment of a nucleotide sequence, identified by any one of SEQ ID NO: 65-68 (Table 10).

(51) In some embodiments, at least one antigenic polypeptide is obtained from MERS-CoV strain Riyadh_14_2013, 2cEMC/2012, or Hasa_1_2013.

(52) In some embodiments, at least one antigenic polypeptide is a SARS-CoV structural protein. For example, a SARS-CoV structural protein may be spike protein (S), envelope protein (E), nucleocapsid protein (N), membrane protein (M) or an immunogenic fragment thereof. In some embodiments, the SARS-CoV structural protein is a spike protein (S). In some embodiments, the SARS-CoV structural protein is a S1 subunit or a S2 subunit of spike protein (S) or an immunogenic fragment thereof.

(53) In some embodiments, at least one SARS-CoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 29, 32 or 34 (Table 11). In some embodiments, the amino acid sequence of the SARS-CoV antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino acid sequence identified by any one of SEQ ID NO: 29, 32 or 34 (Table 11).

(54) In some embodiments, at least one antigenic polypeptide is a HCoV—OC43 structural protein. For example, a HCoV—OC43 structural protein may be spike protein (S), envelope protein (E), nucleocapsid protein (N), membrane protein (M) or an immunogenic fragment thereof. In some embodiments, the HCoV—OC43 structural protein is a spike protein (S). In some embodiments, the HCoV—OC43 structural protein is a S1 subunit or a S2 subunit of spike protein (S) or an immunogenic fragment thereof.

(55) In some embodiments, at least one HCoV—OC43 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 30 (Table 11). In some embodiments, the amino acid sequence of the HCoV—OC43 antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino

acid sequence identified by any one of SEQ ID NO: 30 (Table 11).

(56) In some embodiments, an antigenic polypeptide is a HCoV—HKU1 structural protein. For example, a HCoV—HKU1 structural protein may be spike protein (S), envelope protein (E), nucleocapsid protein (N), membrane protein (M) or an immunogenic fragment thereof. In some embodiments, the HCoV—HKU1 structural protein is a spike protein (S). In some embodiments, the HCoV—HKU1 structural protein is a S1 subunit or a S2 subunit of spike protein (S) or an immunogenic fragment thereof.

(57) In some embodiments, at least one HCoV—HKU1 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 31 (Table 11). In some embodiments, the amino acid sequence of the HCoV—HKU1 antigenic polypeptide is, or is a fragment of, or is a homolog or variant having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to, the amino acid sequence identified by any one of SEQ ID NO: 31 (Table 11).

(58) In some embodiments, an open reading frame of a RNA (e.g., mRNA) vaccine is codon-optimized. In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15) and is codon optimized mRNA.

(59) In some embodiments, a RNA (e.g., mRNA) vaccine further comprising an adjuvant.

(60) Tables 4, 7, 12 and 15 provide National Center for Biotechnology Information (NCBI) accession numbers of interest. It should be understood that the phrase “an amino acid sequence of Tables 4, 7, 12 and 15” refers to an amino acid sequence identified by one or more NCBI accession numbers listed in Tables 4, 7, 12 and 15. Each of the amino acid sequences, and variants having greater than 95% identity or greater than 98% identity to each of the amino acid sequences encompassed by the accession numbers of Tables 4, 7, 12 and 15 are included within the constructs (polynucleotides/polypeptides) of the present disclosure.

(61) In some embodiments, at least one mRNA polynucleotide is encoded by a nucleic acid having a sequence identified by any one of SEQ ID NO: 1-4, 9-12, 20-23, or 35-46 (Tables 2, 5, 10 and 13; see also nucleic acid sequences of Table 7) and having less than 80% identity to wild-type mRNA sequence. In some embodiments, at least one mRNA polynucleotide is encoded by a nucleic acid having a sequence identified by any one of SEQ ID NO: 1-4, 9-12, 20-23, or 35-46 (Tables 2, 5, 10 and 13; see also nucleic acid sequences of Table 7) and having less than 75%, 85% or 95% identity to a wild-type mRNA sequence. In some embodiments, at least one mRNA polynucleotide is encoded by a nucleic acid having a sequence identified by any one of SEQ ID NO: 1-4, 9-12, 20-23, or 35-46 (Tables 2, 5, 10 and 13; see also nucleic acid sequences of Table 7) and having less than 50-80%, 60-80%, 40-80%, 30-80%, 70-80%, 75-80% or 78-80% identity to wild-type mRNA sequence. In some embodiments, at least one mRNA polynucleotide is encoded by a nucleic acid having a sequence identified by any one of SEQ ID NO: 1-4, 9-12, 20-23, or 35-46 (Tables 2, 5, 10 and 13; see also nucleic acid sequences of Table 7) and having less than 40-85%, 50-85%, 60-85%, 30-85%, 70-85%, 75-85% or 80-85% identity to wild-type mRNA sequence. In some embodiments, at least one mRNA polynucleotide is encoded by a nucleic acid having a sequence identified by any one of SEQ ID NO: 1-4, 9-12, 20-23, or 35-46 (Tables 2, 5, 10 and 13; see also nucleic acid sequences of Table 7) and having less than 40-90%, 50-90%, 60-90%, 30-90%, 70-90%, 75-90%, 80-90%, or 85-90% identity to wild-type mRNA sequence.

(62) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15) and having at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) identity to wild-type mRNA sequence, but does not include wild-type mRNA sequence.

(63) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-

34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15) and has less than 95%, 90%, 85%, 80% or 75% identity to wild-type mRNA sequence. In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15) and has 30-80%, 40-80%, 50-80%, 60-80%, 70-80%, 75-80% or 78-80%, 30-85%, 40-85%, 50-805%, 60-85%, 70-85%, 75-85% or 78-85%, 30-90%, 40-90%, 50-90%, 60-90%, 70-90%, 75-90%, 80-90% or 85-90% identity to wild-type mRNA sequence.

(64) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15). In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having 95%-99% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15).

(65) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15) and having membrane fusion activity. In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide having 95%-99% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, or 47-50 (Tables 3, 6, 11 and 14; see also amino acid sequences of Tables 4, 7, 12 and 15) and having membrane fusion activity.

(66) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides) that attaches to cell receptors.

(67) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides) that causes fusion of viral and cellular membranes.

(68) In some embodiments, at least one RNA polynucleotide encodes at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides) that is responsible for binding of the virus to a cell being infected.

(69) Some embodiments of the present disclosure provide a vaccine that includes at least one ribonucleic acid (RNA) (e.g., mRNA) polynucleotide having an open reading frame encoding at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides), at least one 5' terminal

cap and at least one chemical modification, formulated within a lipid nanoparticle.

(70) In some embodiments, a 5' terminal cap is 7mG(5')ppp(5')NlmpNp.

(71) In some embodiments, at least one chemical modification is selected from pseudouridine, N1-methylpseudouridine, N1-ethylpseudouridine, 2-thiouridine, 4'-thiouridine, 5-methylcytosine, 5-methyluridine, 2-thio-1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-pseudouridine, 2-thio-5-aza-uridine, 2-thio-dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-pseudouridine, 4-methoxy-2-thio-pseudouridine, 4-methoxy-pseudouridine, 4-thio-1-methyl-pseudouridine, 4-thio-pseudouridine, 5-aza-uridine, dihydropseudouridine, 5-methoxyuridine and 2'-O-methyl uridine. In some embodiments, the chemical modification is in the 5-position of the uracil. In some embodiments, the chemical modification is a N1-methylpseudouridine. In some embodiments, the chemical modification is a N1-ethylpseudouridine.

(72) In some embodiments, a lipid nanoparticle comprises a cationic lipid, a PEG-modified lipid, a sterol and a non-cationic lipid. In some embodiments, a cationic lipid is an ionizable cationic lipid and the non-cationic lipid is a neutral lipid, and the sterol is a cholesterol. In some embodiments, a cationic lipid is selected from the group consisting of 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), (12Z,15Z)—N,N-dimethyl-2-nonylhenicosa-12,15-dien-1-amine (L608), and N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]heptadecan-8-amine (L530).

(73) In some embodiments, the lipid is

(74) ##STR00001##

(75) In some embodiments, the lipid is

(76) ##STR00002##

(77) In some embodiments, a lipid nanoparticle comprises compounds of Formula (I) and/or Formula (II), discussed below.

(78) In some embodiments, a respiratory virus RNA (e.g., mRNA) vaccine is formulated in a lipid nanoparticle that comprises a compound selected from Compounds 3, 18, 20, 25, 26, 29, 30, 60, 108-112 and 122, described below.

(79) Some embodiments of the present disclosure provide a vaccine that includes at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides), wherein at least 80% (e.g., 85%, 90%, 95%, 98%, 99%) of the uracil in the open reading frame have a chemical modification, optionally wherein the vaccine is formulated in a lipid nanoparticle (e.g., a lipid nanoparticle comprises a cationic lipid, a PEG-modified lipid, a sterol and a non-cationic lipid).

(80) In some embodiments, 100% of the uracil in the open reading frame have a chemical modification. In some embodiments, a chemical modification is in the 5-position of the uracil. In some embodiments, a chemical modification is a N1-methyl pseudouridine. In some embodiments, 100% of the uracil in the open reading frame have a N1-methyl pseudouridine in the 5-position of the uracil.

(81) In some embodiments, an open reading frame of a RNA (e.g., mRNA) polynucleotide encodes at least two antigenic polypeptides (e.g., at least two hMPV antigenic polypeptides, at least two PIV3 antigenic polypeptides, at least two RSV antigenic polypeptides, at least two MeV antigenic polypeptides, or at least two BetaCoV antigenic polypeptides, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides). In some embodiments, the open reading frame encodes at least five or at least ten antigenic polypeptides. In

some embodiments, the open reading frame encodes at least 100 antigenic polypeptides. In some embodiments, the open reading frame encodes 2-100 antigenic polypeptides.

(82) In some embodiments, a vaccine comprises at least two RNA (e.g., mRNA) polynucleotides, each having an open reading frame encoding at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides). In some embodiments, the vaccine comprises at least five or at least ten RNA (e.g., mRNA) polynucleotides, each having an open reading frame encoding at least one antigenic polypeptide or an immunogenic fragment thereof. In some embodiments, the vaccine comprises at least 100 RNA (e.g., mRNA) polynucleotides, each having an open reading frame encoding at least one antigenic polypeptide. In some embodiments, the vaccine comprises 2-100 RNA (e.g., mRNA) polynucleotides, each having an open reading frame encoding at least one antigenic polypeptide.

(83) In some embodiments, at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides) is fused to a signal peptide. In some embodiments, the signal peptide is selected from: a HuIgGk signal peptide (METPAQLLFLLLLWLPDTTG; SEQ ID NO: 15); IgE heavy chain epsilon-1 signal peptide (MDWTWILFLVAAATRVHS; SEQ ID NO: 16); Japanese encephalitis PRM signal sequence (MLGSNSGQRVVFTILLLLVAPAYS; SEQ ID NO: 17), VSVg protein signal sequence (MKCLLYLAFLFIGVNCA; SEQ ID NO: 18) and Japanese encephalitis JEV signal sequence (MWLVSLAIVTACAGA; SEQ ID NO: 19).

(84) In some embodiments, the signal peptide is fused to the N-terminus of at least one antigenic polypeptide. In some embodiments, a signal peptide is fused to the C-terminus of at least one antigenic polypeptide.

(85) In some embodiments, at least one antigenic polypeptide (e.g., at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, or at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides) comprises a mutated N-linked glycosylation site.

(86) Also provided herein is a RNA (e.g., mRNA) vaccine of any one of the foregoing paragraphs (e.g., a hMPV vaccine, a PIV3 vaccine, a RSV vaccine, a MeV vaccine, or a BetaCoV vaccine, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing vaccines), formulated in a nanoparticle (e.g., a lipid nanoparticle).

(87) In some embodiments, the nanoparticle has a mean diameter of 50-200 nm. In some embodiments, the nanoparticle is a lipid nanoparticle. In some embodiments, the lipid nanoparticle comprises a cationic lipid, a PEG-modified lipid, a sterol and a non-cationic lipid. In some embodiments, the lipid nanoparticle comprises a molar ratio of about 20-60% cationic lipid, 0.5-15% PEG-modified lipid, 25-55% sterol, and 25% non-cationic lipid. In some embodiments, the cationic lipid is an ionizable cationic lipid and the non-cationic lipid is a neutral lipid, and the sterol is a cholesterol. In some embodiments, the cationic lipid is selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319).

- (88) In some embodiments, a lipid nanoparticle comprises compounds of Formula (I) and/or Formula (II), as discussed below.
- (89) In some embodiments, a lipid nanoparticle comprises Compounds 3, 18, 20, 25, 26, 29, 30, 60, 108-112, or 122, as discussed below.
- (90) In some embodiments, the nanoparticle has a polydispersity value of less than 0.4 (e.g., less than 0.3, 0.2 or 0.1).
- (91) In some embodiments, the nanoparticle has a net neutral charge at a neutral pH value.
- (92) In some embodiments, the respiratory virus vaccine is multivalent.
- (93) Some embodiments of the present disclosure provide methods of inducing an antigen specific immune response in a subject, comprising administering to the subject any of the RNA (e.g., mRNA) vaccine as provided herein in an amount effective to produce an antigen-specific immune response. In some embodiments, the RNA (e.g., mRNA) vaccine is a hMPV vaccine, a PIV3 vaccine, a RSV vaccine, a MeV vaccine, or a BetaCoV vaccine, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1 vaccines. In some embodiments, the RNA (e.g., mRNA) vaccine is a combination vaccine comprising a combination of any two or more of the foregoing vaccines.
- (94) In some embodiments, an antigen-specific immune response comprises a T cell response or a B cell response.
- (95) In some embodiments, a method of producing an antigen-specific immune response comprises administering to a subject a single dose (no booster dose) of a RNA (e.g., mRNA) vaccine of the present disclosure. In some embodiments, the RNA (e.g., mRNA) vaccine is a hMPV vaccine, a PIV3 vaccine, a RSV vaccine, a MeV vaccine, or a BetaCoV vaccine, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1 vaccines. In some embodiments, the RNA (e.g., mRNA) vaccine is a combination vaccine comprising a combination of any two or more of the foregoing vaccines.
- (96) In some embodiments, a method further comprises administering to the subject a second (booster) dose of a RNA (e.g., mRNA) vaccine. Additional doses of a RNA (e.g., mRNA) vaccine may be administered.
- (97) In some embodiments, the subjects exhibit a seroconversion rate of at least 80% (e.g., at least 85%, at least 90%, or at least 95%) following the first dose or the second (booster) dose of the vaccine. Seroconversion is the time period during which a specific antibody develops and becomes detectable in the blood. After seroconversion has occurred, a virus can be detected in blood tests for the antibody. During an infection or immunization, antigens enter the blood, and the immune system begins to produce antibodies in response. Before seroconversion, the antigen itself may or may not be detectable, but antibodies are considered absent. During seroconversion, antibodies are present but not yet detectable. Any time after seroconversion, the antibodies can be detected in the blood, indicating a prior or current infection.
- (98) In some embodiments, a RNA (e.g., mRNA) vaccine is administered to a subject by intradermal or intramuscular injection.
- (99) Some embodiments, of the present disclosure provide methods of inducing an antigen specific immune response in a subject, including administering to a subject a RNA (e.g., mRNA) vaccine in an effective amount to produce an antigen specific immune response in a subject. Antigen-specific immune responses in a subject may be determined, in some embodiments, by assaying for antibody titer (for titer of an antibody that binds to a hMPV, PIV3, RSV, MeV and/or BetaCoV antigenic polypeptide) following administration to the subject of any of the RNA (e.g., mRNA) vaccines of the present disclosure. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased by at least 1 log relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased by 1-3 log relative to a control.
- (100) In some embodiments, the anti-antigenic polypeptide antibody titer produced in a subject is

increased at least 2 times relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased at least 5 times relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased at least 10 times relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased 2-10 times relative to a control.

(101) In some embodiments, the control is an anti-antigenic polypeptide antibody titer produced in a subject who has not been administered a RNA (e.g., mRNA) vaccine of the present disclosure. In some embodiments, the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine (see, e.g., Ren J. et al. *J of Gen. Virol.* 2015; 96: 1515-1520), or wherein the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. In some embodiments, the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a hMPV, PIV3, RSV, MeV and/or BetaCoV virus-like particle (VLP) vaccine (see, e.g., Cox R G et al., *J Virol.* 2014 June; 88(11): 6368-6379).

(102) A RNA (e.g., mRNA) vaccine of the present disclosure is administered to a subject in an effective amount (an amount effective to induce an immune response). In some embodiments, the effective amount is a dose equivalent to an at least 2-fold, at least 4-fold, at least 10-fold, at least 100-fold, at least 1000-fold reduction in the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, wherein the anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, a purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, a live attenuated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine, an inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine, or a hMPV, PIV3, RSV, MeV and/or BetaCoV VLP vaccine. In some embodiments, the effective amount is a dose equivalent to 2-1000-fold reduction in the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, wherein the anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, a purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, a live attenuated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine, an inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine, or a hMPV, PIV3, RSV, MeV and/or BetaCoV VLP vaccine.

(103) In some embodiments, the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a virus-like particle (VLP) vaccine comprising structural proteins of hMPV, PIV3, RSV, MeV and/or BetaCoV.

(104) In some embodiments, the RNA (e.g., mRNA) vaccine is formulated in an effective amount to produce an antigen specific immune response in a subject.

(105) In some embodiments, the effective amount is a total dose of 25 µg to 1000 µg, or 50 µg to 1000 µg. In some embodiments, the effective amount is a total dose of 100 µg. In some embodiments, the effective amount is a dose of 25 µg administered to the subject a total of two times. In some embodiments, the effective amount is a dose of 100 µg administered to the subject a total of two times. In some embodiments, the effective amount is a dose of 400 µg administered to the subject a total of two times. In some embodiments, the effective amount is a dose of 500 µg administered to the subject a total of two times.

(106) In some embodiments, the efficacy (or effectiveness) of a RNA (e.g., mRNA) vaccine is greater than 60%. In some embodiments, the RNA (e.g., mRNA) polynucleotide of the vaccine at least one hMPV antigenic polypeptide, at least one PIV3 antigenic polypeptide, at least one RSV antigenic polypeptide, at least one MeV antigenic polypeptide, at least one BetaCoV antigenic polypeptide, e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-

NL63, HCoV-NL, HCoV—NH and HCoV—HKU1, or any combination of two or more of the foregoing antigenic polypeptides.

(107) Vaccine efficacy may be assessed using standard analyses (see, e.g., Weinberg et al., *J Infect Dis.* 2010 Jun. 1; 201(11):1607-10). For example, vaccine efficacy may be measured by double-blind, randomized, clinical controlled trials. Vaccine efficacy may be expressed as a proportionate reduction in disease attack rate (AR) between the unvaccinated (ARU) and vaccinated (ARV) study cohorts and can be calculated from the relative risk (RR) of disease among the vaccinated group with use of the following formulas:

(108) $\text{Efficacy} = (\text{ARU} - \text{ARV}) / \text{ARU} \times 100$; and $\text{Efficacy} = (1 - \text{RR}) \times 100$.

(109) Likewise, vaccine effectiveness may be assessed using standard analyses (see, e.g., Weinberg et al., *J Infect Dis.* 2010 Jun. 1; 201(11):1607-10). Vaccine effectiveness is an assessment of how a vaccine (which may have already proven to have high vaccine efficacy) reduces disease in a population. This measure can assess the net balance of benefits and adverse effects of a vaccination program, not just the vaccine itself, under natural field conditions rather than in a controlled clinical trial. Vaccine effectiveness is proportional to vaccine efficacy (potency) but is also affected by how well target groups in the population are immunized, as well as by other non-vaccine-related factors that influence the ‘real-world’ outcomes of hospitalizations, ambulatory visits, or costs. For example, a retrospective case control analysis may be used, in which the rates of vaccination among a set of infected cases and appropriate controls are compared. Vaccine effectiveness may be expressed as a rate difference, with use of the odds ratio (OR) for developing infection despite vaccination:

(110) $\text{Effectiveness} = (1 - \text{OR}) \times 100$.

(111) In some embodiments, the efficacy (or effectiveness) of a RNA (e.g., mRNA) vaccine is at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, or at least 90%.

(112) In some embodiments, the vaccine immunizes the subject against hMPV, PIV3, RSV, MeV, BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1), or any combination of two or more of the foregoing viruses for up to 2 years. In some embodiments, the vaccine immunizes the subject against hMPV, PIV3, RSV, MeV, BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1), or any combination of two or more of the foregoing viruses for more than 2 years, more than 3 years, more than 4 years, or for 5-10 years.

(113) In some embodiments, the subject is about 5 years old or younger. For example, the subject may be between the ages of about 1 year and about 5 years (e.g., about 1, 2, 3, 5 or 5 years), or between the ages of about 6 months and about 1 year (e.g., about 6, 7, 8, 9, 10, 11 or 12 months). In some embodiments, the subject is about 12 months or younger (e.g., 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 months or 1 month). In some embodiments, the subject is about 6 months or younger.

(114) In some embodiments, the subject was born full term (e.g., about 37-42 weeks). In some embodiments, the subject was born prematurely, for example, at about 36 weeks of gestation or earlier (e.g., about 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26 or 25 weeks). For example, the subject may have been born at about 32 weeks of gestation or earlier. In some embodiments, the subject was born prematurely between about 32 weeks and about 36 weeks of gestation. In such subjects, a RNA (e.g., mRNA) vaccine may be administered later in life, for example, at the age of about 6 months to about 5 years, or older.

(115) In some embodiments, the subject is pregnant (e.g., in the first, second or third trimester) when administered an RNA (e.g., mRNA) vaccine. Viruses such as hMPV, PIV3 and RSV causes infections of the lower respiratory tract, mainly in infants and young children. One-third of RSV related deaths, for example, occur in the first year of life, with 99 percent of these deaths occurring in low-resource countries. It's so widespread in the United States that nearly all children become infected with the virus before their second birthdays. Thus, the present disclosure provides RNA

(e.g., mRNA) vaccines for maternal immunization to improve mother-to-child transmission of protection against the virus.

(116) In some embodiments, the subject is a young adult between the ages of about 20 years and about 50 years (e.g., about 20, 25, 30, 35, 40, 45 or 50 years old).

(117) In some embodiments, the subject is an elderly subject about 60 years old, about 70 years old, or older (e.g., about 60, 65, 70, 75, 80, 85 or 90 years old).

(118) In some embodiments, the subject has a chronic pulmonary disease (e.g., chronic obstructive pulmonary disease (COPD) or asthma). Two forms of COPD include chronic bronchitis, which involves a long-term cough with mucus, and emphysema, which involves damage to the lungs over time. Thus, a subject administered a RNA (e.g., mRNA) vaccine may have chronic bronchitis or emphysema.

(119) In some embodiments, the subject has been exposed to hMPV, PIV3, RSV, MeV, BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1), or any combination of two or more of the foregoing viruses; the subject is infected with hMPV, PIV3, RSV, MeV, BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1), or any combination of two or more of the foregoing viruses; or subject is at risk of infection by hMPV, PIV3, RSV, MeV, BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1), or any combination of two or more of the foregoing viruses.

(120) In some embodiments, the subject is immunocompromised (has an impaired immune system, e.g., has an immune disorder or autoimmune disorder).

(121) In some embodiments the nucleic acid vaccines described herein are chemically modified. In other embodiments the nucleic acid vaccines are unmodified.

(122) Yet other aspects provide compositions for and methods of vaccinating a subject comprising administering to the subject a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a first respiratory virus antigenic polypeptide, wherein the RNA polynucleotide does not include a stabilization element, and wherein an adjuvant is not coformulated or co-administered with the vaccine.

(123) In other aspects the invention is a composition for or method of vaccinating a subject comprising administering to the subject a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide wherein a dosage of between 10 µg/kg and 400 µg/kg of the nucleic acid vaccine is administered to the subject. In some embodiments the dosage of the RNA polynucleotide is 1-5 µg, 5-10 µg, 10-15 µg, 15-20 µg, 10-25 µg, 20-25 µg, 20-50 µg, 30-50 µg, 40-50 µg, 40-60 µg, 60-80 µg, 60-100 µg, 50-100 µg, 80-120 µg, 40-120 µg, 40-150 µg, 50-150 µg, 50-200 µg, 80-200 µg, 100-200 µg, 120-250 µg, 150-250 µg, 180-280 µg, 200-300 µg, 50-300 µg, 80-300 µg, 100-300 µg, 40-300 µg, 50-350 µg, 100-350 µg, 200-350 µg, 300-350 µg, 320-400 µg, 40-380 µg, 40-100 µg, 100-400 µg, 200-400 µg, or 300-400 µg per dose. In some embodiments, the nucleic acid vaccine is administered to the subject by intradermal or intramuscular injection. In some embodiments, the nucleic acid vaccine is administered to the subject on day zero. In some embodiments, a second dose of the nucleic acid vaccine is administered to the subject on day twenty one.

(124) In some embodiments, a dosage of 25 micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, a dosage of 100 micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, a dosage of 50 micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, a dosage of 75 micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, a dosage of 150 micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, a dosage of 400

micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, a dosage of 200 micrograms of the RNA polynucleotide is included in the nucleic acid vaccine administered to the subject. In some embodiments, the RNA polynucleotide accumulates at a 100 fold higher level in the local lymph node in comparison with the distal lymph node. In other embodiments the nucleic acid vaccine is chemically modified and in other embodiments the nucleic acid vaccine is not chemically modified.

(125) Aspects of the invention provide a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide, wherein the RNA polynucleotide does not include a stabilization element, and a pharmaceutically acceptable carrier or excipient, wherein an adjuvant is not included in the vaccine. In some embodiments, the stabilization element is a histone stem-loop. In some embodiments, the stabilization element is a nucleic acid sequence having increased GC content relative to wild type sequence.

(126) Aspects of the invention provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide, wherein the RNA polynucleotide is present in the formulation for in vivo administration to a host, which confers an antibody titer superior to the criterion for seroprotection for the first antigen for an acceptable percentage of human subjects. In some embodiments, the antibody titer produced by the mRNA vaccines of the invention is a neutralizing antibody titer. In some embodiments the neutralizing antibody titer is greater than a protein vaccine. In other embodiments the neutralizing antibody titer produced by the mRNA vaccines of the invention is greater than an adjuvanted protein vaccine. In yet other embodiments the neutralizing antibody titer produced by the mRNA vaccines of the invention is 1,000-10,000, 1,200-10,000, 1,400-10,000, 1,500-10,000, 1,000-5,000, 1,000-4,000, 1,800-10,000, 2000-10,000, 2,000-5,000, 2,000-3,000, 2,000-4,000, 3,000-5,000, 3,000-4,000, or 2,000-2,500. A neutralization titer is typically expressed as the highest serum dilution required to achieve a 50% reduction in the number of plaques.

(127) Also provided are nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide, wherein the RNA polynucleotide is present in a formulation for in vivo administration to a host for eliciting a longer lasting high antibody titer than an antibody titer elicited by an mRNA vaccine having a stabilizing element or formulated with an adjuvant and encoding the first antigenic polypeptide. In some embodiments, the RNA polynucleotide is formulated to produce a neutralizing antibodies within one week of a single administration. In some embodiments, the adjuvant is selected from a cationic peptide and an immunostimulatory nucleic acid. In some embodiments, the cationic peptide is protamine.

(128) Aspects provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification or optionally no nucleotide modification, the open reading frame encoding a first antigenic polypeptide, wherein the RNA polynucleotide is present in the formulation for in vivo administration to a host such that the level of antigen expression in the host significantly exceeds a level of antigen expression produced by an mRNA vaccine having a stabilizing element or formulated with an adjuvant and encoding the first antigenic polypeptide.

(129) Other aspects provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification or optionally no nucleotide modification, the open reading frame encoding a first antigenic polypeptide, wherein the vaccine has at least 10 fold less RNA polynucleotide than is required for an unmodified mRNA vaccine to produce an equivalent antibody titer. In some embodiments, the RNA polynucleotide is present in a dosage of 25-100 micrograms.

(130) Aspects of the invention also provide a unit of use vaccine, comprising between 10 ug and 400 ug of one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification or optionally no nucleotide modification, the open reading frame encoding a

first antigenic polypeptide, and a pharmaceutically acceptable carrier or excipient, formulated for delivery to a human subject. In some embodiments, the vaccine further comprises a cationic lipid nanoparticle.

(131) Aspects of the invention provide methods of creating, maintaining or restoring antigenic memory to a respiratory virus strain in an individual or population of individuals comprising administering to said individual or population an antigenic memory booster nucleic acid vaccine comprising (a) at least one RNA polynucleotide, said polynucleotide comprising at least one chemical modification or optionally no nucleotide modification and two or more codon-optimized open reading frames, said open reading frames encoding a set of reference antigenic polypeptides, and (b) optionally a pharmaceutically acceptable carrier or excipient. In some embodiments, the vaccine is administered to the individual via a route selected from the group consisting of intramuscular administration, intradermal administration and subcutaneous administration. In some embodiments, the administering step comprises contacting a muscle tissue of the subject with a device suitable for injection of the composition. In some embodiments, the administering step comprises contacting a muscle tissue of the subject with a device suitable for injection of the composition in combination with electroporation.

(132) Aspects of the invention provide methods of vaccinating a subject comprising administering to the subject a single dosage of between 25 ug/kg and 400 ug/kg of a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a first antigenic polypeptide in an effective amount to vaccinate the subject.

(133) Other aspects provide nucleic acid vaccines comprising one or more RNA polynucleotides having an open reading frame comprising at least one chemical modification, the open reading frame encoding a first antigenic polypeptide, wherein the vaccine has at least 10 fold less RNA polynucleotide than is required for an unmodified mRNA vaccine to produce an equivalent antibody titer. In some embodiments, the RNA polynucleotide is present in a dosage of 25-100 micrograms.

(134) Other aspects provide nucleic acid vaccines comprising an LNP formulated RNA polynucleotide having an open reading frame comprising no nucleotide modifications (unmodified), the open reading frame encoding a first antigenic polypeptide, wherein the vaccine has at least 10 fold less RNA polynucleotide than is required for an unmodified mRNA vaccine not formulated in a LNP to produce an equivalent antibody titer. In some embodiments, the RNA polynucleotide is present in a dosage of 25-100 micrograms.

(135) The data presented in the Examples demonstrate significant enhanced immune responses using the formulations of the invention. Both chemically modified and unmodified RNA vaccines are useful according to the invention. Surprisingly, in contrast to prior art reports that it was preferable to use chemically unmodified mRNA formulated in a carrier for the production of vaccines, it is described herein that chemically modified mRNA-LNP vaccines required a much lower effective mRNA dose than unmodified mRNA, i.e., tenfold less than unmodified mRNA when formulated in carriers other than LNP. Both the chemically modified and unmodified RNA vaccines of the invention produce better immune responses than mRNA vaccines formulated in a different lipid carrier.

(136) In other aspects the invention encompasses a method of treating an elderly subject age 60 years or older comprising administering to the subject a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a respiratory virus antigenic polypeptide in an effective amount to vaccinate the subject.

(137) In other aspects the invention encompasses a method of treating a young subject age 17 years or younger comprising administering to the subject a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a respiratory virus antigenic polypeptide in an effective amount to vaccinate the subject.

(138) In other aspects the invention encompasses a method of treating an adult subject comprising

administering to the subject a nucleic acid vaccine comprising one or more RNA polynucleotides having an open reading frame encoding a respiratory virus antigenic polypeptide in an effective amount to vaccinate the subject.

(139) In some aspects the invention is a method of vaccinating a subject with a combination vaccine including at least two nucleic acid sequences encoding respiratory antigens wherein the dosage for the vaccine is a combined therapeutic dosage wherein the dosage of each individual nucleic acid encoding an antigen is a sub therapeutic dosage. In some embodiments, the combined dosage is 25 micrograms of the RNA polynucleotide in the nucleic acid vaccine administered to the subject. In some embodiments, the combined dosage is 100 micrograms of the RNA polynucleotide in the nucleic acid vaccine administered to the subject. In some embodiments the combined dosage is 50 micrograms of the RNA polynucleotide in the nucleic acid vaccine administered to the subject. In some embodiments, the combined dosage is 75 micrograms of the RNA polynucleotide in the nucleic acid vaccine administered to the subject. In some embodiments, the combined dosage is 150 micrograms of the RNA polynucleotide in the nucleic acid vaccine administered to the subject. In some embodiments, the combined dosage is 400 micrograms of the RNA polynucleotide in the nucleic acid vaccine administered to the subject. In some embodiments, the sub therapeutic dosage of each individual nucleic acid encoding an antigen is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 micrograms. In other embodiments the nucleic acid vaccine is chemically modified and in other embodiments the nucleic acid vaccine is not chemically modified.

(140) The RNA polynucleotide is one of SEQ ID NO: 1-4, 9-12, 20-23, 35-46, 57-61, and 64-80 and includes at least one chemical modification. In other embodiments the RNA polynucleotide is one of SEQ ID NO: 1-4, 9-12, 20-23, 35-46, 57-61, and 64-80 and does not include any nucleotide modifications, or is unmodified. In yet other embodiments the at least one RNA polynucleotide encodes an antigenic protein of any of SEQ ID NO: 5-8, 12-13, 24-34, and 47-50 and includes at least one chemical modification. In other embodiments the RNA polynucleotide encodes an antigenic protein of any of SEQ ID NO: 5-8, 12-13, 24-34, and 47-50 and does not include any nucleotide modifications, or is unmodified.

(141) In preferred aspects, vaccines of the invention (e.g., LNP-encapsulated mRNA vaccines) produce prophylactically- and/or therapeutically-efficacious levels, concentrations and/or titers of antigen-specific antibodies in the blood or serum of a vaccinated subject. As defined herein, the term antibody titer refers to the amount of antigen-specific antibody produces in s subject, e.g., a human subject. In exemplary embodiments, antibody titer is expressed as the inverse of the greatest dilution (in a serial dilution) that still gives a positive result. In exemplary embodiments, antibody titer is determined or measured by enzyme-linked immunosorbent assay (ELISA). In exemplary embodiments, antibody titer is determined or measured by neutralization assay, e.g., by microneutralization assay. In certain aspects, antibody titer measurement is expressed as a ratio, such as 1:40, 1:100, etc.

(142) In exemplary embodiments of the invention, an efficacious vaccine produces an antibody titer of greater than 1:40, greater that 1:100, greater than 1:400, greater than 1:1000, greater than 1:2000, greater than 1:3000, greater than 1:4000, greater than 1:500, greater than 1:6000, greater than 1:7500, greater than 1:10000. In exemplary embodiments, the antibody titer is produced or reached by 10 days following vaccination, by 20 days following vaccination, by 30 days following vaccination, by 40 days following vaccination, or by 50 or more days following vaccination. In exemplary embodiments, the titer is produced or reached following a single dose of vaccine administered to the subject. In other embodiments, the titer is produced or reached following multiple doses, e.g., following a first and a second dose (e.g., a booster dose.)

(143) In exemplary aspects of the invention, antigen-specific antibodies are measured in units of µg/ml or are measured in units of IU/L (International Units per liter) or mIU/ml (milli International Units per ml). In exemplary embodiments of the invention, an efficacious vaccine produces >0.5 µg/ml, >0.1 µg/ml, >0.2 µg/ml, >0.35 µg/ml, >0.5 µg/ml, >1 µg/ml, >2 µg/ml, >5 µg/ml or >10

µg/ml. In exemplary embodiments of the invention, an efficacious vaccine produces >10 mIU/ml, >20 mIU/ml, >50 mIU/ml, >100 mIU/ml, >200 mIU/ml, >500 mIU/ml or >1000 mIU/ml. In exemplary embodiments, the antibody level or concentration is produced or reached by 10 days following vaccination, by 20 days following vaccination, by 30 days following vaccination, by 40 days following vaccination, or by 50 or more days following vaccination. In exemplary embodiments, the level or concentration is produced or reached following a single dose of vaccine administered to the subject. In other embodiments, the level or concentration is produced or reached following multiple doses, e.g., following a first and a second dose (e.g., a booster dose.) In exemplary embodiments, antibody level or concentration is determined or measured by enzyme-linked immunosorbent assay (ELISA). In exemplary embodiments, antibody level or concentration is determined or measured by neutralization assay, e.g., by microneutralization assay.

(144) The details of various embodiments of the disclosure are set forth in the description below. Other features, objects, and advantages of the disclosure will be apparent from the description and from the claims.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

- (1) The foregoing and other objects, features and advantages will be apparent from the following description of particular embodiments of the disclosure, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating the principles of various embodiments of the disclosure.
- (2) FIG. 1 shows a schematic of one example of a RNA (e.g. mRNA) vaccine construct of the present disclosure. The construct depicts a human metapneumovirus and human respiratory syncytial virus full length fusion protein obtained from wild-type strains (*The Journal of General Virology*. 2008; 89 (Pt 12):3113-3118, incorporated herein by reference).
- (3) FIGS. 2A-2C are graphs showing the levels of anti-hMPV fusion protein-specific antibodies in the serum of mice immunized with hMPV mRNA vaccines on day 0 (FIG. 2A), day 14 (FIG. 2B) and day 35 (FIG. 2C) post immunization. The mice were immunized with a single dose (2 µg or 10 µg) on day 0 and were given a boost dose (2 µg or 10 µg) on day 21. hMPV fusion protein-specific antibodies were detected at up to 1:10000 dilution of serum on day 35 for both doses.
- (4) FIGS. 3A-3C are graphs showing the result of IgG isotyping in the serum of mice immunized with hMPV mRNA vaccines. The levels of hMPV fusion protein-specific IgG2a (FIG. 3A) and IgG1 (FIG. 3B) antibodies in the serum are measured by ELISA. FIG. 3C shows that hMPV fusion protein mRNA vaccine induced a mixed Th1/Th2 cytokine response with a Th1 bias.
- (5) FIG. 4 is a graph showing in vitro neutralization of a hMPV B2 strain (TN/91-316) using the sera of mice immunized with a mRNA vaccine encoding hMPV fusion protein. Mouse serum obtained from mice receiving a 10 µg or a 2 µg dose contained hMPV-neutralizing antibodies.
- (6) FIGS. 5A-5C are graphs showing a Th1 cytokine response induced by a hMPV fusion peptide pool (15-mers-50 (overlap)) in splenocytes isolated from mice immunized with the hMPV mRNA vaccines. Virus-free media was used as a negative control and Concanavalin A (ConA, a positive control for splenocyte stimulation) was included. The cytokines tested included IFN-γ (FIG. 5A), IL-2 (FIG. 5B) and IL12 (FIG. 5C).
- (7) FIGS. 6A-6E are graphs showing the Th2 cytokine response induced by a hMPV fusion peptide pool (15-mers-50) in splenocytes isolated from mice immunized with the hMPV mRNA vaccines. Virus-free media was used as a negative control and Concanavalin A was also included. The cytokines tested included IL-10 (FIG. 6A), TNF-α (FIG. 6B), IL4 (FIG. 6C), IL-5 (FIG. 6D) and IL-6 (FIG. 6E).

(8) FIGS. 7A-7C are graphs showing the Th1 response induced by inactivated hMPV virus in splenocytes isolated from mice immunized with hMPV mRNA vaccines. Virus-free media was used as a negative control and Concanavalin A was included. The cytokines tested included IFN-7 (FIG. 7A), IL-2 (FIG. 7B) and IL12 (FIG. 7C).

(9) FIGS. 8A-8E are graphs showing the Th2 response induced by inactivated hMPV virus in splenocytes isolated from mice immunized with the hMPV mRNA vaccines. Virus-free media was used as a negative control and Concanavalin A was included. The cytokines tested include IL-10 (FIG. 8A), TNF- α (FIG. 8B), IL4 (FIG. 8C), IL-5 (FIG. 8D) and IL-6 (FIG. 8E).

(10) FIGS. 9A-9B are graphs showing the results of cotton rat challenge experiments. Two different doses of the hMPV mRNA vaccines were used (2 μ g or 10 μ g doses) to immunize the cotton rats before challenge. The hMPV mRNA vaccines reduced the viral titer in the lung and nose of the cotton rat, with the 10 μ g dose being more effective in reducing viral titer. Use of a 10 μ g dose resulted in 100% protection in the lung and a ~2 log reduction in nose viral titer. Use of a 2 μ g dose resulted in a 1 log reduction in lung viral titer and no reduction in nose viral titer. The vaccine was administered on Day 0, and a boost was administered on Day 21.

(11) FIG. 10 is a graph showing the lung histopathology of cotton rats that received hMPV mRNA vaccines. Pathology associated with vaccine-enhanced disease was not observed in immunized groups.

(12) FIG. 11 is a graph showing hMPV neutralization antibody titers in cotton rats that received hMPV mRNA vaccines (2 μ g or 10 μ g doses) on days 35 and 42 post immunization.

(13) FIG. 12 is a graph showing the lung and nose viral load in cotton rats challenged with a hMPV/A2 strain after immunization with the indicated mRNA vaccines (hMPV mRNA vaccine or hMPV/PIV mRNA combination vaccine). Vaccinated cotton rats showed reduced lung and nose viral loads after challenge, compared to control.

(14) FIG. 13 is a graph showing the lung and nose viral load in cotton rats challenged with PIV3 strain after immunization with indicated mRNA vaccines (PIV mRNA vaccine or hMPV/PIV combination vaccine). Vaccinated cotton rats showed reduced lung and nose viral loads after challenge, compared to control.

(15) FIG. 14 is a graph showing hMPV neutralizing antibody titers in cotton rats that received different dosages of hMPV mRNA vaccines or hMPV/PIV combination mRNA vaccines on day 42 post immunization. The dosages of the vaccine are indicated in Table 9.

(16) FIG. 15 is a graph showing PIV3 neutralizing antibody titers in cotton rats that received different dosages of PIV mRNA vaccines or hMPV/PIV combination mRNA vaccines on day 42 post immunization. The dosages of the vaccine are indicated in Table 9.

(17) FIG. 16 is a graph showing the lung histopathology score of cotton rats immunized with hMPV mRNA vaccines, PIV mRNA vaccines or hMPV/PIV combination mRNA vaccines as indicated in Table 9. Low occurrence of alveolitis and interstitial pneumonia was observed, indicating no antibody-dependent enhancement (ADE) of hMPV associated diseases.

(18) FIG. 17 is a graph showing the reciprocal MERS-CoV neutralizing antibody titers in mice immunized with betacoronavirus mRNA vaccine encoding the MERS-CoV full-length Spike protein, on days 0, 21, 42, and 56 post immunization.

(19) FIG. 18 is a graph showing the reciprocal MERS-CoV neutralizing antibody titers in mice immunized with betacoronavirus mRNA vaccine encoding either the MERS-CoV full-length Spike protein, or the S2 subunit of the Spike protein. The full length spike protein induced a stronger immune response compared to the S2 subunit alone.

(20) FIGS. 19A-19C are graphs showing the viral load in the nose and throat, the bronchoalveolar lavage (BAL), or the lungs of New Zealand white rabbits 4 days post challenge with MERS-CoV. The New Zealand white rabbits were immunized with one 20 μ g-dose (on day 0) or two 20 μ g-doses (on day 0 and 21) of MERS-CoV mRNA vaccine encoding the full-length Spike protein before challenge. FIG. 19A shows that two doses of MERS-CoV mRNA vaccine resulted in a 3 log

reduction of viral load in the nose and led to complete protection in the throat of the New Zealand white rabbits. FIG. 19B shows that two doses of MERS-CoV mRNA vaccine resulted in a 4 log reduction of viral load in the BAL of the New Zealand white rabbits. FIG. 19C show one dose of MERS-CoV mRNA vaccine resulted in a 2 log reduction of viral load, while two doses of MERS-CoV mRNA vaccine resulted in an over 4 log reduction of viral load in the lungs of the New Zealand white rabbits.

(21) FIGS. 20A-20B are images and graphs showing viral load or replicating virus detected by PCR in the lungs of New Zealand white rabbits 4 days post challenge with MERS-CoV. The New Zealand white rabbits were immunized with a single 20 µg dose (on day 0, Group 1a) of MERS-CoV mRNA vaccine encoding the full-length Spike protein, two 20 µg doses (on day 0 and 21, Group 1b) of MERS-CoV mRNA vaccine encoding the full-length Spike protein, or placebo (Group 2) before challenge. FIG. 20A shows that two doses of 20 µg a MERS-CoV mRNA vaccine reduced over 99% (2 log) of viruses in the lungs of New Zealand white rabbits. FIG. 20B shows that the group of New Zealand white rabbits that received 2 doses of 20 µg MERS-CoV mRNA vaccine did not have any detectable replicating MERS-CoV virus in their lungs.

(22) FIG. 21 is a graph showing the MERS-CoV neutralizing antibody titers in New Zealand white rabbits immunized with MERS-CoV mRNA vaccine encoding the full-length Spike protein. The results show that two doses of 20 µg MERS-CoV mRNA vaccine induced a significant amount of neutralizing antibodies against MERS-CoV (EC₅₀ between 500-1000). The MERS-CoV mRNA vaccine induced antibody titer is 3-5 fold better than any other vaccines tested in the same model.

DETAILED DESCRIPTION

(23) The present disclosure provides, in some embodiments, vaccines that comprise RNA (e.g., mRNA) polynucleotides encoding a human metapneumovirus (hMPV) antigenic polypeptide, a parainfluenza virus type 3 (PIV3) antigenic polypeptide, a respiratory syncytial virus (RSV) antigenic polypeptide, a measles virus (MeV) antigenic polypeptide, or a betacoronavirus antigenic polypeptide (e.g., Middle East respiratory syndrome coronavirus (MERS-CoV), SARS-CoV, human coronavirus (HCoV)—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH (New Haven) and HCoV—HKU1) (see, e.g., Esper F. et al. *Emerging Infectious Diseases*, 12(5), 2006; and Pyre K. et al. *Journal of Virology*, 81(7):3051-57, 2007, the contents of each of which is here incorporated by reference in their entirety). The present disclosure also provides, in some embodiments, combination vaccines that comprise at least one RNA (e.g., mRNA) polynucleotide encoding at least two antigenic polypeptides selected from hMPV antigenic polypeptides, PIV3 antigenic polypeptides, RSV antigenic polypeptides, MeV antigenic polypeptides and BetaCoV antigenic polypeptides. Also provided herein are methods of administering the RNA (e.g., mRNA) vaccines, methods of producing the RNA (e.g., mRNA) vaccines, compositions (e.g., pharmaceutical compositions) comprising the RNA (e.g., mRNA) vaccines, and nucleic acids (e.g., DNA) encoding the RNA (e.g., mRNA) vaccines. In some embodiments, a RNA (e.g., mRNA) vaccine comprises an adjuvant, such as a flagellin adjuvant, as provided herein.

(24) The RNA (e.g., mRNA) vaccines (e.g., hMPV, PIV3, RSV, MeV, BetaCoV RNA vaccines and combinations thereof), in some embodiments, may be used to induce a balanced immune response, comprising both cellular and humoral immunity, without many of the risks associated with DNA vaccination.

(25) The entire contents of International Application No. PCT/US2015/02740 is incorporated herein by reference.

(26) Human Metapneumovirus (hMPV)

(27) hMPV shares substantial homology with respiratory syncytial virus (RSV) in its surface glycoproteins. hMPV fusion protein (F) is related to other paramyxovirus fusion proteins and appears to have homologous regions that may have similar functions. The hMPV fusion protein amino acid sequence contains features characteristic of other paramyxovirus F proteins, including a

putative cleavage site and potential N-linked glycosylation sites. Paramyxovirus fusion proteins are synthesized as inactive precursors (F0) that are cleaved by host cell proteases into the biologically fusion-active F1 and F2 domains (see, e.g., Cseke G. et al. *Journal of Virology* 2007; 81(2):698-707, incorporated herein by reference). hMPV has one putative cleavage site, in contrast to the two sites established for RSV F, and only shares 34% amino acid sequence identity with RSV F. F2 is extracellular and disulfide linked to F1. Fusion proteins are type I glycoproteins existing as trimers, with two 4-3 heptad repeat domains at the N- and C-terminal regions of the protein (HR1 and HR2), which form coiled-coil alpha-helices. These coiled coils become apposed in an antiparallel fashion when the protein undergoes a conformational change into the fusogenic state. There is a hydrophobic fusion peptide N proximal to the N-terminal heptad repeat, which is thought to insert into the target cell membrane, while the association of the heptad repeats brings the transmembrane domain into close proximity, inducing membrane fusion (see, e.g., Baker, K A et al. *Mol. Cell* 1999; 3:309-319). This mechanism has been proposed for a number of different viruses, including RSV, influenza virus, and human immunodeficiency virus. Fusion proteins are major antigenic determinants for all known paramyxoviruses and for other viruses that possess similar fusion proteins such as human immunodeficiency virus, influenza virus, and Ebola virus.

(28) In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding hMPV fusion protein (F). In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding a F1 or F2 subunit of a hMPV F protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding hMPV glycoprotein (G). In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding hMPV matrix protein (M). In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding hMPV phosphoprotein (P). In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding hMPV nucleoprotein (N). In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding hMPV SH protein (SH).

(29) In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein, M protein, P protein, N protein and SH protein.

(30) In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and G protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and M protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and P protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and N protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and SH protein.

(31) In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and M protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and P protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and N protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and SH protein.

(32) In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and M protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and P protein. In some embodiments, a hMPV vaccine of the present disclosure

comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and N protein. In some embodiments, a hMPV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and SH protein.

(33) A hMPV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one hMPV antigenic polypeptide identified by any one of SEQ ID NO: 5-8 (Table 3; see also amino acid sequences of Table 4).

(34) A hMPV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide encoded by a nucleic acid (e.g., DNA) identified by any one of SEQ ID NO: 1-4 (Table 2).

(35) The present disclosure is not limited by a particular strain of hMPV. The strain of hMPV used in a vaccine may be any strain of hMPV. Non-limiting examples of strains of hMPV for use as provide herein include the CAN98-75 (CAN75) and the CAN97-83 (CAN83) hMPV strains (Skiadopoulos M H et al. *J Virol.* 20014; 78 (13) 6927-37, incorporated herein by reference), a hMPV A1, A2, B1 or B2 strain (see, e.g., de Graaf M et al. *The Journal of General Virology* 2008; 89:975-83; Peret T C T et al. *The Journal of Infectious Disease* 2002; 185:1660-63, incorporated herein by reference), a hMPV isolate TN/92-4 (e.g., SEQ ID NO: 1 and 5), a hMPV isolate NL/1/99 (e.g., SEQ ID NO: 2 and 6), or a hMPV isolate PER/CFI0497/2010/B (e.g., SEQ ID NO: 3 and 7).

(36) In some embodiments, at least one hMPV antigenic polypeptide is obtained from a hMPV A1, A2, B1 or B2 strain (see, e.g., de Graaf M et al. *The Journal of General Virology* 2008; 89:975-83; Peret T C T et al. *The Journal of Infectious Disease* 2002; 185:1660-63, incorporated herein by reference). In some embodiments, at least one antigenic polypeptide is obtained from the CAN98-75 (CAN75) hMPV strain. In some embodiments, at least one antigenic polypeptide is obtained from the CAN97-83 (CAN83) hMPV strain. In some embodiments, at least one antigenic polypeptide is obtained from hMPV isolate TN/92-4 (e.g., SEQ ID NO: 1 and 5). In some embodiments, at least one antigenic polypeptide is obtained from hMPV isolate NL/1/99 (e.g., SEQ ID NO: 2 and 6). In some embodiments, at least one antigenic polypeptide is obtained from hMPV isolate PER/CFI0497/2010/B (e.g., SEQ ID NO: 3 and 7).

(37) In some embodiments, hMPV vaccines comprise RNA (e.g., mRNA) polynucleotides encoding a hMPV antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with hMPV F protein and having F protein activity.

(38) A protein is considered to have F protein activity if, for example, the protein acts to fuse the viral envelope and host cell plasma membrane, mediates viral entry into a host cell via an interaction with arginine-glycine-aspartate RGD-binding integrins, or a combination thereof (see, e.g., Cox R G et al. *J Virol.* 2012; 88(22):12148-60, incorporated herein by reference).

(39) In some embodiments, hMPV vaccines comprise RNA (e.g., mRNA) polynucleotides encoding hMPV antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with hMPV G protein and having G protein activity.

(40) A protein is considered to have G protein activity if, for example, the protein acts to modulate (e.g., inhibit) hMPV-induced cellular (immune) responses (see, e.g., Bao X et al. *PLoS Pathog.* 2008; 4 (5):e1000077, incorporated herein by reference).

(41) Human Parainfluenza Virus Type 3 (PIV3)

(42) Parainfluenza viruses belong to the family Paramyxoviridae. These are enveloped viruses with a negative-sense single-stranded RNA genome. Parainfluenza viruses belong to the subfamily Paramyxoviridae, which is subdivided into three genera: Respirovirus (PIV-1, PIV-3, and Sendai virus (SeV)), Rubulavirus (PIV-2, PIV-4 and mumps virus) and Morbillivirus (measles virus, rinderpest virus and canine distemper virus (CDV)). Their genome, a ~15 500 nucleotide-long negative-sense RNA molecule, encodes two envelope glycoproteins, the hemagglutinin-neuraminidase (HN), the fusion protein (F or F0), which is cleaved into F1 and F2 subunits, a matrix protein (M), a nucleocapsid protein (N) and several nonstructural proteins including the viral replicase (L). All parainfluenza viruses, except for PIV-1, express a non-structural V protein that blocks IFN signaling in the infected cell and acts therefore as a virulence factor (see, e.g.,

Nishio M et al. *J Virol.* 2008; 82(13):6130-38).

(43) PIV3 hemagglutinin-neuraminidase (HN), a structural protein, is found on the viral envelope, where it is necessary for attachment and cell entry. It recognizes and binds to sialic acid-containing receptors on the host cell's surface. As a neuroaminidase, HN removes sialic acid from virus particles, preventing self-aggregation of the virus, and promoting the efficient spread of the virus. Furthermore, HN promotes the activity of the fusion (F or F0) protein, contributing to the penetration of the host cell's surface.

(44) PIV3 fusion protein (PIV3 F) is located on the viral envelope, where it facilitates the viral fusion and cell entry. The F protein is initially inactive, but proteolytic cleavage leads to its active forms, F1 and F2, which are linked by disulfide bonds. This occurs when the HN protein binds its receptor on the host cell's surface. During early phases of infection, the F glycoprotein mediates penetration of the host cell by fusion of the viral envelope to the plasma membrane. In later stages of the infection, the F protein facilitates the fusion of the infected cells with neighboring uninfected cells, which leads to the formation of a syncytium and spread of the infection.

(45) PIV3 matrix protein (M) is found within the viral envelope and assists with viral assembly. It interacts with the nucleocapsid and envelope glycoproteins, where it facilitates the budding of progeny viruses through its interactions with specific sites on the cytoplasmic tail of the viral glycoproteins and nucleocapsid. It also plays a role in transporting viral components to the budding site.

(46) PIV3 phosphoprotein (P) and PIV3 large polymerase protein (L) are found in the nucleocapsid where they form part of the RNA polymerase complex. The L protein, a viral RNA-dependent RNA polymerase, facilitates genomic transcription, while the host cell's ribosomes translate the viral mRNA into viral proteins.

(47) PIV3 V is a non-structural protein that blocks IFN signaling in the infected cell, therefore acting as a virulence factor.

(48) PIV3 nucleoprotein (N) encapsidates the genome in a ratio of 1 N per 6 ribonucleotides, protecting it from nucleases. The nucleocapsid (NC) has a helical structure. The encapsidated genomic RNA is termed the NC and serves as template for transcription and replication. During replication, encapsidation by PIV3 N is coupled to RNA synthesis and all replicative products are resistant to nucleases. PIV3 N homo-multimerizes to form the nucleocapsid and binds to viral genomic RNA. PIV3 N binds the P protein and thereby positions the polymerase on the template.

(49) In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding PIV3 fusion protein (F). In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding a F1 or F2 subunit of a PIV3 F protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding PIV3 hemagglutinin-neuraminidase (HN) (see, e.g., van Wyke Coelingh K L et al. *J Virol.* 1987; 61(5):1473-77, incorporated herein by reference). In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding PIV3 matrix protein (M). In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding PIV3 phosphoprotein (P). In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding PIV3 nucleoprotein (N).

(50) In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, HN protein, M protein, P protein, and N protein.

(51) In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and HN protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and M protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and P protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and N

protein.

(52) In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HN protein and M protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HN protein and P protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HN protein and N protein.

(53) In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, HN protein and M protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, HN protein and P protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, HN protein and N protein.

(54) A PIV3 vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one PIV3 antigenic polypeptide identified by any one of SEQ ID NO: 12-13 (Table 6; see also amino acid sequences of Table 7).

(55) A PIV3 vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide encoded by a nucleic acid (e.g., DNA) identified by any one of SEQ ID NO: 9-12 (Table 5; see also nucleic acid sequences of Table 7).

(56) The present disclosure is not limited by a particular strain of PIV3. The strain of PIV3 used in a vaccine may be any strain of PIV3. A non-limiting example of a strain of PIV3 for use as provide herein includes HPIV3/*Homo sapiens*/PER/FLA4815/2008.

(57) In some embodiments, PIV3 vaccines comprise RNA (e.g., mRNA) polynucleotides encoding a PIV3 antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with PIV3 F protein and having F protein activity.

(58) In some embodiments, PIV3 vaccines comprise RNA (e.g., mRNA) polynucleotides encoding PIV3 antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with PIV3 hemagglutinin-neuraminidase (HN) and having hemagglutinin-neuraminidase activity.

(59) A protein is considered to have hemagglutinin-neuraminidase activity if, for example, it is capable of both receptor binding and receptor cleaving. Such proteins are major surface glycoproteins that have functional sites for cell attachment and for neuraminidase activity. They are able to cause red blood cells to agglutinate and to cleave the glycosidic linkages of neuraminic acids, so they have the potential to both bind a potential host cell and then release the cell if necessary, for example, to prevent self-aggregation of the virus.

(60) In some embodiments, PIV3 vaccines comprise RNA (e.g., mRNA) polynucleotides encoding PIV3 antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with PIV3 HN, F (e.g., F, F1 or F2), M, N, L or V and having HN, F (e.g., F, F1 or F2), M, N, L or V activity, respectively.

(61) Respiratory Syncytial Virus (RSV)

(62) RSV is a negative-sense, single-stranded RNA virus of the genus Pneumovirinae. The virus is present in at least two antigenic subgroups, known as Group A and Group B, primarily resulting from differences in the surface G glycoproteins. Two RSV surface glycoproteins—G and F—mediate attachment with and attachment to cells of the respiratory epithelium. F surface glycoproteins mediate coalescence of neighboring cells. This results in the formation of syncytial cells. RSV is the most common cause of bronchiolitis. Most infected adults develop mild cold-like symptoms such as congestion, low-grade fever, and wheezing. Infants and small children may suffer more severe symptoms such as bronchiolitis and pneumonia. The disease may be transmitted among humans via contact with respiratory secretions.

(63) The genome of RSV encodes at least three surface glycoproteins, including F, G, and SH, four nucleocapsid proteins, including L, P, N, and M2, and one matrix protein, M. Glycoprotein F directs viral penetration by fusion between the virion and the host membrane. Glycoprotein G is a

type II transmembrane glycoprotein and is the major attachment protein. SH is a short integral membrane protein. Matrix protein M is found in the inner layer of the lipid bilayer and assists virion formation. Nucleocapsid proteins L, P, N, and M2 modulate replication and transcription of the RSV genome. It is thought that glycoprotein G tethers and stabilizes the virus particle at the surface of bronchial epithelial cells, while glycoprotein F interacts with cellular glycosaminoglycans to mediate fusion and delivery of the RSV virion contents into the host cell (Krzyzaniak M A et al. *PLoS Pathog* 2013; 9 (4)).

(64) In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding L protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding P protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding N protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding M2 protein. In some embodiments, a PIV3 vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding M protein.

(65) In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein, L protein, P protein, N protein, M2 protein and M protein.

(66) In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and G protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and L protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and P protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and N protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and M2 protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and M protein.

(67) In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and L protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and P protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and N protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and M2 protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding G protein and M protein.

(68) In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and L protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and P protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and N protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and M2 protein. In some embodiments, a RSV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein, G protein and M protein.

(69) The present disclosure is not limited by a particular strain of RSV. The strain of RSV used in a vaccine may be any strain of RSV.

(70) In some embodiments, RSV vaccines comprise RNA (e.g., mRNA) polynucleotides encoding

a RSV antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with RSV F protein and having F protein activity.

(71) In some embodiments, RSV vaccines comprise RNA (e.g., mRNA) polynucleotides encoding RSV antigenic polypeptides having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with RSV G protein and having G protein activity.

(72) A protein is considered to have G protein activity if, for example, the protein acts to modulate (e.g., inhibit) hMPV-induced cellular (immune) responses (see, e.g., Bao X et al. *PLoS Pathog.* 2008; 4 (5):e1000077, incorporated herein by reference).

(73) Measles Virus (MeV)

(74) Molecular epidemiologic investigations and virologic surveillance contribute notably to the control and prevention of measles. Nearly half of measles-related deaths worldwide occur in India, yet virologic surveillance data are incomplete for many regions of the country. Previous studies have documented the presence of measles virus genotypes D4, D7, and D8 in India, and genotypes D5, D9, D11, H1, and G3 have been detected in neighboring countries. Recently, MeV genotype B3 was detected in India (Kuttiatt V S et al. *Emerg Infect Dis.* 2014; 20(10): 1764-66).

(75) The glycoprotein complex of paramyxoviruses mediates receptor binding and membrane fusion. In particular, the MeV fusion (F) protein executes membrane fusion, after receptor binding by the hemagglutinin (HA) protein (Muhlebach M D et al. *Journal of Virology* 2008; 82(22):11437-45). The MeV P gene codes for three proteins: P, an essential polymerase cofactor, and V and C, which have multiple functions but are not strictly required for viral propagation in cultured cells. V shares the amino-terminal domain with P but has a zinc-binding carboxyl-terminal domain, whereas C is translated from an overlapping reading frame. The MeV C protein is an infectivity factor. During replication, the P protein binds incoming monomeric nucleocapsid (N) proteins with its amino-terminal domain and positions them for assembly into the nascent ribonucleocapsid. The P protein amino-terminal domain is natively unfolded (Deveaux P et al. *Journal of Virology* 2004; 78(21):11632-40).

(76) In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding P protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding V protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding C protein.

(77) In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein, F protein, P protein, V protein and C protein.

(78) In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein and F protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein and P protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein and V protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein and C protein.

(79) In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and P protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and V protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding F protein and C protein.

(80) In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein, F protein and P protein. In some embodiments, a

MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein, F protein and V protein. In some embodiments, a MeV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding HA protein, F protein and C protein.

(81) In some embodiments, MeV vaccines comprise RNA (e.g., mRNA) encoding a MeV antigenic polypeptide having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with MeV HA protein and having MeV HA protein activity.

(82) In some embodiments, MeV vaccines comprise RNA (e.g., mRNA) encoding a MeV antigenic polypeptide having at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity with MeV F protein and having MeV F protein activity.

(83) A protein is considered to have HA protein activity if the protein mediates receptor binding and/or membrane fusion. MeV F protein executes membrane fusion, after receptor binding by the MeV HA protein.

(84) A MeV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one MeV antigenic polypeptide identified by any one of SEQ ID NO: 47-50 (Table 14; see also amino acid sequences of Table 15).

(85) A MeV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide identified by any one of SEQ ID NO: 37, 40, 43, 46 (Table 13).

(86) A MeV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide encoded by a nucleic acid (e.g., DNA) identified by any one of SEQ ID NO: 35, 36, 38, 39, 41, 42, 44 and 45 (Table 13).

(87) The present disclosure is not limited by a particular strain of MeV. The strain of MeV used in a vaccine may be any strain of MeV. Non-limiting examples of strains of MeV for use as provide herein include B3/B3.1, C2, D4, D6, D7, D8, G3, H1, Moraten, Rubeovax, MVi/New Jersey.USA/45.05, MVi/Texas.USA/4.07, AIK-C, MVi/New York.USA/26.09/3, MVi/California.USA/16.03, MVi/Virginia.USA/15.09, MVi/California.USA/8.04, and MVi/Pennsylvania.USA/20.09.

(88) MeV proteins may be from MeV genotype D4, D5, D7, D8, D9, D11, H1, G3 or B3. In some embodiments, a MeV HA protein or a MeV F protein is from MeV genotype D8. In some embodiments, a MeV HA protein or a MeV F protein is from MeV genotype B3.

(89) Betacoronaviruses (BetaCoV)

(90) MERS-CoV. MERS-CoV is a positive-sense, single-stranded RNA virus of the genus Betacoronavirus. The genomes are phylogenetically classified into two clades, clade A and clade B. It has a strong tropism for non-ciliated bronchial epithelial cells, evades the innate immune response and antagonizes interferon (IFN) production in infected cells. Dipeptyl peptidase 4 (DDP4, also known as CD26) has been identified as a functional cellular receptor for MERS-CoV. Its enzymatic activity is not required for infection, although its amino acid sequence is highly conserved across species and is expressed in the human bronchial epithelium and kidneys. Most infected individuals develop severe acute respiratory illnesses, including fever, cough, and shortness of breath, and the virus can be fatal. The disease may be transmitted among humans, generally among those in close contact.

(91) The genome of MERS-CoV encodes at least four unique accessory proteins, such as 3, 4a, 4b and 5, two replicase proteins (open reading frame 1a and 1b), and four major structural proteins, including spike (S), envelope (E), nucleocapsid (N), and membrane (M) proteins (Almazan F et al. *MBio* 2013; 4 (5):e00650-13). The accessory proteins play nonessential roles in MERS-CoV replication, but they are likely structural proteins or interferon antagonists, modulating in vivo replication efficiency and/or pathogenesis, as in the case of SARS-CoV (Almazan F et al. *MBio* 2013; 4 (5):e00650-13; Totura A L et al. *Curr Opin Virol* 2012; 2(3):264-75; Scobey T et al. *Proc Natl Acad Sci USA* 2013; 110(40):16157-62). The other proteins of MERS-CoV maintain different functions in virus replication. The E protein, for example, involves in virulence, and deleting the E-

coding gene results in replication-competent and propagation-defective viruses or attenuated viruses (Almazan F et al. *MBio* 2013; 4 (5):e00650-13). The S protein is particularly essential in mediating virus binding to cells expressing receptor dipeptidyl peptidase-4 (DPP4) through receptor-binding domain (RBD) in the S1 subunit, whereas the S2 subunit subsequently mediates virus entry via fusion of the virus and target cell membranes (Li F. *J Virol* 2015; 89(4):1954-64; Raj V S et al. *Nature* 2013; 495(7440):251-4).

(92) In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding the S1 subunit of the S protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding the S2 subunit of the S protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding E protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding N protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding M protein.

(93) In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), E protein, N protein and M protein.

(94) In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2) and E protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2) and N protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2) and M protein.

(95) In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), E protein and M protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), E protein and N protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), M protein and N protein. In some embodiments, a MERS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding E protein, M protein and N protein.

(96) A MERS-CoV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one MERS-CoV antigenic polypeptide identified by any one of SEQ ID NO: 24-38 or 33 (Table 11; see also amino acid sequences of Table 12).

(97) A MERS-CoV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide encoded by a nucleic acid (e.g., DNA) identified by any one of SEQ ID NO: 20-23 (Table 10).

(98) The present disclosure is not limited by a particular strain of MERS-CoV. The strain of MERS-CoV used in a vaccine may be any strain of MERS-CoV. Non-limiting examples of strains of MERS-CoV for use as provide herein include Riyadh_14_2013, and 2cEMC/2012, Hasa_1_2013.

(99) SARS-CoV. The genome of SARS-CoV includes of a single, positive-strand RNA that is approximately 29,700 nucleotides long. The overall genome organization of SARS-CoV is similar to that of other coronaviruses. The reference genome includes 13 genes, which encode at least 14 proteins. Two large overlapping reading frames (ORFs) encompass 71% of the genome. The remainder has 12 potential ORFs, including genes for structural proteins S (spike), E (small envelope), M (membrane), and N (nucleocapsid). Other potential ORFs code for unique putative SARS-CoV-specific polypeptides that lack obvious sequence similarity to known proteins. A

detailed analysis of the SARS-CoV genome has been published in *J Mol Biol* 2003; 331: 991-1004.

(100) In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), E protein, N protein and M protein.

(101) In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2) and E protein. In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2) and N protein. In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2) and M protein.

(102) In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), E protein and M protein. In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), E protein and N protein. In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding S protein (S, S1 and/or S2), M protein and N protein. In some embodiments, a SARS-CoV vaccine of the present disclosure comprises a RNA (e.g., mRNA) polynucleotide encoding E protein, M protein and N protein.

(103) A SARS-CoV vaccine may comprise, for example, at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one SARS-CoV antigenic polypeptide identified by any one of SEQ ID NO: 29, 32 or 34 (Table 11; see also amino acid sequences of Table 12).

(104) The present disclosure is not limited by a particular strain of SARS-CoV. The strain of SARS-CoV used in a vaccine may be any strain of SARS-CoV.

(105) HCoV—OC43. Human coronavirus OC43 is an enveloped, positive-sense, single-stranded RNA virus in the species Betacoronavirus-1 (genus Betacoronavirus, subfamily Coronavirinae, family Coronaviridae, order Nidovirales). Four HCoV—OC43 genotypes (A to D), have been identified with genotype D most likely arising from recombination. The complete genome sequencing of two genotype C and D strains and bootscan analysis shows recombination events between genotypes B and C in the generation of genotype D. Of 29 strains identified, none belong to the more ancient genotype A. Along with HCoV-229E, a species in the Alphacoronavirus genus, HCoV—OC43 are among the known viruses that cause the common cold. Both viruses can cause severe lower respiratory tract infections, including pneumonia in infants, the elderly, and immunocompromised individuals such as those undergoing chemotherapy and those with HIV-AIDS.

(106) HCoV—HKU1. Human coronavirus HKU1 (HCoV—HKU1) is a positive-sense, single-stranded RNA virus with the HE gene, which distinguishes it as a group 2, or betacoronavirus. It was discovered in January 2005 in two patients in Hong Kong. The genome of HCoV—HKU1 is a 29,926-nucleotide, polyadenylated RNA. The GC content is 32%, the lowest among all known coronaviruses. The genome organization is the same as that of other group II coronaviruses, with the characteristic gene order 1a, 1b, HE, S, E, M, and N. Furthermore, accessory protein genes are present between the S and E genes (ORF4) and at the position of the N gene (ORF8). The TRS is presumably located within the AAUCUAAAC sequence, which precedes each ORF except E. As in sialodacryoadenitis virus and mouse hepatitis virus (MHV), translation of the E protein possibly occurs via an internal ribosomal entry site. The 3' untranslated region contains a predicted stem-loop structure immediately downstream of the N ORF (nucleotide position 29647 to 29711). Further downstream, a pseudoknot structure is present at nucleotide position 29708 to 29760. Both RNA structures are conserved in group II coronaviruses and are critical for virus replication.

(107) HCoV-NL63. The RNA genome of human coronavirus NL63 (HCoV-NL63) is 27,553 nucleotides, with a poly(A) tail (FIG. 1). With a GC content of 34%, HCoV-NL63 has one of the

lowest GC contents of the coronaviruses, for which GC content ranges from 32 to 42%. Untranslated regions of 286 and 287 nucleotides are present at the 5' and 3' termini, respectively. Genes predicted to encode the S, E, M, and N proteins are found in the 3' part of the HCoV-NL63 genome. The HE gene, which is present in some group II coronaviruses, is absent, and there is only a single, monocistronic accessory protein ORF (ORF3) located between the S and E genes. Subgenomic mRNAs are generated for all ORFs (S, ORF3, E, M, and N), and the core sequence of the TRS of HCoV-NL63 is defined as AACUAAA. This sequence is situated upstream of every ORF except for the E ORF, which contains the suboptimal core sequence AACUAUA. Interestingly, a 13-nucleotide sequence with perfect homology to the leader sequence is situated upstream of the suboptimal E TRS. Annealing of this 13-nucleotide sequence to the leader sequence may act as a compensatory mechanism for the disturbed leader-TRS/body-TRS interaction.

(108) HCoV-229E. Human coronavirus 229E (HCoV-229E) is a single-stranded, positive-sense, RNA virus species in the Alphacoronavirus genus of the subfamily Coronavirinae, in the family Coronaviridae, of the order Nidovirales. Along with Human coronavirus OC43, it is responsible for the common cold. HCoV-NL63 and HCoV-229E are two of the four human coronaviruses that circulate worldwide. These two viruses are unique in their relationship towards each other. Phylogenetically, the viruses are more closely related to each other than to any other human coronavirus, yet they only share 65% sequence identity. Moreover, the viruses use different receptors to enter their target cell. HCoV-NL63 is associated with croup in children, whereas all signs suggest that the virus probably causes the common cold in healthy adults. HCoV-229E is a proven common cold virus in healthy adults, so it is probable that both viruses induce comparable symptoms in adults, even though their mode of infection differs (HCoV-NL63 and HCoV-229E are two of the four human coronaviruses that circulate worldwide. These two viruses are unique in their relationship towards each other. Phylogenetically, the viruses are more closely related to each other than to any other human coronavirus, yet they only share 65% sequence identity. Moreover, the viruses use different receptors to enter their target cell. HCoV-NL63 is associated with croup in children, whereas all signs suggest that the virus probably causes the common cold in healthy adults. HCoV-229E is a proven common cold virus in healthy adults, so it is probable that both viruses induce comparable symptoms in adults, even though their mode of infection differs (Dijkman R. et al. *J Formos Med Assoc.* 2009 April; 108(4):270-9, the contents of which is incorporated herein by reference in their entirety).

(109) Combination Vaccines

(110) Embodiments of the present disclosure also provide combination RNA (e.g., mRNA) vaccines. A “combination RNA (e.g., mRNA) vaccine” of the present disclosure refers to a vaccine comprising at least one (e.g., at least 2, 3, 4, or 5) RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a combination of any two or more (or all of) antigenic polypeptides selected from hMPV antigenic polypeptides, PIV3 antigenic polypeptides, RSV antigenic polypeptides, MeV antigenic polypeptides, and BetaCoV antigenic polypeptides (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(111) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide, a RSV antigenic polypeptide, a MeV antigenic polypeptide, and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(112) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide and a PIV3 antigenic polypeptide.

(113) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide and a RSV antigenic polypeptide.

- (114) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide and a MeV antigenic polypeptide.
- (115) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide and a BetaCoV antigenic polypeptide.
- (116) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a PIV3 antigenic polypeptide and a RSV antigenic polypeptide.
- (117) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a PIV3 antigenic polypeptide and a MeV antigenic polypeptide.
- (118) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a PIV3 antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (119) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a RSV antigenic polypeptide and a MeV antigenic polypeptide.
- (120) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a RSV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (121) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a MeV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (122) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide, a RSV antigenic polypeptide and a MeV antigenic polypeptide.
- (123) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide, a RSV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (124) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide, a MeV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (125) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a RSV antigenic polypeptide, a MeV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (126) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a PIV3 antigenic polypeptide, a RSV antigenic polypeptide, a MeV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).
- (127) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide and a RSV antigenic polypeptide.
- (128) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g.,

mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide and a MeV antigenic polypeptide.

(129) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a PIV3 antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(130) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a RSV antigenic polypeptide and a MeV antigenic polypeptide.

(131) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a RSV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(132) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a hMPV antigenic polypeptide, a MeV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(133) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a PIV3 antigenic polypeptide, a RSV antigenic polypeptide and a MeV antigenic polypeptide.

(134) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a PIV3 antigenic polypeptide, a RSV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(135) In some embodiments, a combination RNA (e.g., mRNA) vaccine comprises a RNA (e.g., mRNA) polynucleotide encoding a RSV antigenic polypeptide, a MeV antigenic polypeptide and a BetaCoV antigenic polypeptide (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1).

(136) Other combination respiratory virus RNA (e.g., mRNA) vaccines are encompassed by the present disclosure.

(137) It has been discovered that the mRNA vaccines described herein are superior to current vaccines in several ways. First, the lipid nanoparticle (LNP) delivery is superior to other formulations including a protamine base approach described in the literature and no additional adjuvants are to be necessary. The use of LNPs enables the effective delivery of chemically modified or unmodified mRNA vaccines. Additionally it has been demonstrated herein that both modified and unmodified LNP formulated mRNA vaccines were superior to conventional vaccines by a significant degree. In some embodiments the mRNA vaccines of the invention are superior to conventional vaccines by a factor of at least 10 fold, 20 fold, 40 fold, 50 fold, 100 fold, 500 fold or 1,000 fold.

(138) Although attempts have been made to produce functional RNA vaccines, including mRNA vaccines and self-replicating RNA vaccines, the therapeutic efficacy of these RNA vaccines have not yet been fully established. Quite surprisingly, the inventors have discovered, according to aspects of the invention a class of formulations for delivering mRNA vaccines in vivo that results in significantly enhanced, and in many respects synergistic, immune responses including enhanced antigen generation and functional antibody production with neutralization capability. These results can be achieved even when significantly lower doses of the mRNA are administered in comparison with mRNA doses used in other classes of lipid based formulations. The formulations of the invention have demonstrated significant unexpected in vivo immune responses sufficient to establish the efficacy of functional mRNA vaccines as prophylactic and therapeutic agents. Additionally, self-replicating RNA vaccines rely on viral replication pathways to deliver enough

RNA to a cell to produce an immunogenic response. The formulations of the invention do not require viral replication to produce enough protein to result in a strong immune response. Thus, the mRNA of the invention are not self-replicating RNA and do not include components necessary for viral replication.

(139) The invention involves, in some aspects, the surprising finding that lipid nanoparticle (LNP) formulations significantly enhance the effectiveness of mRNA vaccines, including chemically modified and unmodified mRNA vaccines. The efficacy of mRNA vaccines formulated in LNP was examined in vivo using several distinct antigens. The results presented herein demonstrate the unexpected superior efficacy of the mRNA vaccines formulated in LNP over other commercially available vaccines.

(140) In addition to providing an enhanced immune response, the formulations of the invention generate a more rapid immune response with fewer doses of antigen than other vaccines tested. The mRNA-LNP formulations of the invention also produce quantitatively and qualitatively better immune responses than vaccines formulated in a different carriers.

(141) The data described herein demonstrate that the formulations of the invention produced significant unexpected improvements over existing antigen vaccines. Additionally, the mRNA-LNP formulations of the invention are superior to other vaccines even when the dose of mRNA is lower than other vaccines. Mice immunized with either 10 µg or 2 µg doses of an hMPV fusion protein mRNA LNP vaccine or a PIV3 mRNA LNP vaccine produced neutralizing antibodies which for instance, successfully neutralized the hMPV B2 virus. A 10 µg dose of mRNA vaccine protected 100% of mice from lethal challenge and drastically reduced the viral titer after challenge (~2 log reduction).

(142) Two 20 µg doses of MERS-CoV mRNA LNP vaccine significantly reduced viral load and induced significant amount of neutralizing antibodies against MERS-CoV (EC.sub.50 between 500-1000). The MERS-CoV mRNA vaccine induced antibody titer was 3-5 fold better than any other vaccines tested in the same model.

(143) The LNP used in the studies described herein has been used previously to deliver siRNA in various animal models as well as in humans. In view of the observations made in association with the siRNA delivery of LNP formulations, the fact that LNP is useful in vaccines is quite surprising. It has been observed that therapeutic delivery of siRNA formulated in LNP causes an undesirable inflammatory response associated with a transient IgM response, typically leading to a reduction in antigen production and a compromised immune response. In contrast to the findings observed with siRNA, the LNP-mRNA formulations of the invention are demonstrated herein to generate enhanced IgG levels, sufficient for prophylactic and therapeutic methods rather than transient IgM responses.

(144) Nucleic Acids/Polynucleotides

(145) Respiratory virus vaccines, as provided herein, comprise at least one (one or more) ribonucleic acid (RNA) (e.g., mRNA) polynucleotide having an open reading frame encoding at least one antigenic polypeptide selected from hMPV, PIV3, RSV, MeV and BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1) antigenic polypeptides. The term “nucleic acid” includes any compound and/or substance that comprises a polymer of nucleotides (nucleotide monomer). These polymers are referred to as polynucleotides. Thus, the terms “nucleic acid” and “polynucleotide” are used interchangeably.

(146) Nucleic acids may be or may include, for example, ribonucleic acids (RNAs), deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked nucleic acids (LNAs, including LNA having a β-D-ribo configuration, α-LNA having an α-L-ribo configuration (a diastereomer of LNA), 2'-amino-LNA having a 2'-amino functionalization, and 2'-amino-α-LNA having a 2'-amino functionalization), ethylene nucleic acids (ENA), cyclohexenyl nucleic acids (CeNA) or chimeras or combinations

thereof.

(147) In some embodiments, polynucleotides of the present disclosure function as messenger RNA (mRNA). “Messenger RNA” (mRNA) refers to any polynucleotide that encodes a (at least one) polypeptide (a naturally-occurring, non-naturally-occurring, or modified polymer of amino acids) and can be translated to produce the encoded polypeptide in vitro, in vivo, in situ or ex vivo. The skilled artisan will appreciate that, except where otherwise noted, polynucleotide sequences set forth in the instant application will recite “T”s in a representative DNA sequence but where the sequence represents RNA (e.g., mRNA), the “T”s would be substituted for “U”s. Thus, any of the RNA polynucleotides encoded by a DNA identified by a particular sequence identification number may also comprise the corresponding RNA (e.g., mRNA) sequence encoded by the DNA, where each “T” of the DNA sequence is substituted with “U.”

(148) The basic components of an mRNA molecule typically include at least one coding region, a 5′ untranslated region (UTR), a 3′ UTR, a 5′ cap and a poly-A tail. Polynucleotides of the present disclosure may function as mRNA but can be distinguished from wild-type mRNA in their functional and/or structural design features, which serve to overcome existing problems of effective polypeptide expression using nucleic-acid based therapeutics.

(149) In some embodiments, a RNA polynucleotide of an RNA (e.g., mRNA) vaccine encodes 2-10, 2-9, 2-8, 2-7, 2-6, 2-5, 2-4, 2-3, 3-10, 3-9, 3-8, 3-7, 3-6, 3-5, 3-4, 4-10, 4-9, 4-8, 4-7, 4-6, 4-5, 5-10, 5-9, 5-8, 5-7, 5-6, 6-10, 6-9, 6-8, 6-7, 7-10, 7-9, 7-8, 8-10, 8-9 or 9-10 antigenic polypeptides. In some embodiments, a RNA (e.g., mRNA) polynucleotide of a respiratory virus vaccine encodes at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 antigenic polypeptides. In some embodiments, a RNA (e.g., mRNA) polynucleotide of a respiratory virus vaccine encodes at least 100 or at least 200 antigenic polypeptides. In some embodiments, a RNA polynucleotide of an respiratory virus vaccine encodes 1-10, 5-15, 10-20, 15-25, 20-30, 25-35, 30-40, 35-45, 40-50, 1-50, 1-100, 2-50 or 2-100 antigenic polypeptides.

(150) Polynucleotides of the present disclosure, in some embodiments, are codon optimized. Codon optimization methods are known in the art and may be used as provided herein. Codon optimization, in some embodiments, may be used to match codon frequencies in target and host organisms to ensure proper folding; bias GC content to increase mRNA stability or reduce secondary structures; minimize tandem repeat codons or base runs that may impair gene construction or expression; customize transcriptional and translational control regions; insert or remove protein trafficking sequences; remove/add post translation modification sites in encoded protein (e.g. glycosylation sites); add, remove or shuffle protein domains; insert or delete restriction sites; modify ribosome binding sites and mRNA degradation sites; adjust translational rates to allow the various domains of the protein to fold properly; or to reduce or eliminate problem secondary structures within the polynucleotide. Codon optimization tools, algorithms and services are known in the art—non-limiting examples include services from GeneArt (Life Technologies), DNA2.0 (Menlo Park CA) and/or proprietary methods. In some embodiments, the open reading frame (ORF) sequence is optimized using optimization algorithms.

(151) In some embodiments, a codon optimized sequence shares less than 95% sequence identity, less than 90% sequence identity, less than 85% sequence identity, less than 80% sequence identity, or less than 75% sequence identity to a naturally-occurring or wild-type sequence (e.g., a naturally-occurring or wild-type mRNA sequence encoding a polypeptide or protein of interest (e.g., an antigenic protein or antigenic polypeptide)).

(152) In some embodiments, a codon-optimized sequence shares between 65% and 85% (e.g., between about 67% and about 85%, or between about 67% and about 80%) sequence identity to a naturally-occurring sequence or a wild-type sequence (e.g., a naturally-occurring or wild-type mRNA sequence encoding a polypeptide or protein of interest (e.g., an antigenic protein or polypeptide)). In some embodiments, a codon-optimized sequence shares between 65% and 75%, or about 80% sequence identity to a naturally-occurring sequence or wild-type sequence (e.g., a

naturally-occurring or wild-type mRNA sequence encoding a polypeptide or protein of interest (e.g., an antigenic protein or polypeptide)).

(153) In some embodiments a codon-optimized RNA (e.g., mRNA) may, for instance, be one in which the levels of G/C are enhanced. The G/C-content of nucleic acid molecules may influence the stability of the RNA. RNA having an increased amount of guanine (G) and/or cytosine (C) residues may be functionally more stable than nucleic acids containing a large amount of adenine (A) and thymine (T) or uracil (U) nucleotides. WO02/098443 discloses a pharmaceutical composition containing an mRNA stabilized by sequence modifications in the translated region. Due to the degeneracy of the genetic code, the modifications work by substituting existing codons for those that promote greater RNA stability without changing the resulting amino acid. The approach is limited to coding regions of the RNA.

(154) Antigens/Antigenic Polypeptides

(155) In some embodiments, an antigenic polypeptide (e.g., a hMPV, PIV3, RSV, MeV or BetaCoV antigenic polypeptide) is longer than 25 amino acids and shorter than 50 amino acids. Polypeptides include gene products, naturally occurring polypeptides, synthetic polypeptides, homologs, orthologs, paralogs, fragments and other equivalents, variants, and analogs of the foregoing. A polypeptide may be a single molecule or may be a multi-molecular complex such as a dimer, trimer or tetramer. Polypeptides may also comprise single chain polypeptides or multichain polypeptides, such as antibodies or insulin, and may be associated or linked to each other. Most commonly, disulfide linkages are found in multichain polypeptides. The term “polypeptide” may also apply to amino acid polymers in which at least one amino acid residue is an artificial chemical analogue of a corresponding naturally-occurring amino acid.

(156) A “polypeptide variant” is a molecule that differs in its amino acid sequence relative to a native sequence or a reference sequence. Amino acid sequence variants may possess substitutions, deletions, insertions, or a combination of any two or three of the foregoing, at certain positions within the amino acid sequence, as compared to a native sequence or a reference sequence. Ordinarily, variants possess at least 50% identity to a native sequence or a reference sequence. In some embodiments, variants share at least 80% identity or at least 90% identity with a native sequence or a reference sequence.

(157) In some embodiments “variant mimics” are provided. A “variant mimic” contains at least one amino acid that would mimic an activated sequence. For example, glutamate may serve as a mimic for phosphoro-threonine and/or phosphoro-serine. Alternatively, variant mimics may result in deactivation or in an inactivated product containing the mimic. For example, phenylalanine may act as an inactivating substitution for tyrosine, or alanine may act as an inactivating substitution for serine.

(158) “Orthologs” refers to genes in different species that evolved from a common ancestral gene by speciation. Normally, orthologs retain the same function in the course of evolution. Identification of orthologs is important for reliable prediction of gene function in newly sequenced genomes.

(159) “Analog” is meant to include polypeptide variants that differ by one or more amino acid alterations, for example, substitutions, additions or deletions of amino acid residues that still maintain one or more of the properties of the parent or starting polypeptide.

(160) The present disclosure provides several types of compositions that are polynucleotide or polypeptide based, including variants and derivatives. These include, for example, substitutional, insertional, deletion and covalent variants and derivatives. The term “derivative” is synonymous with the term “variant” and generally refers to a molecule that has been modified and/or changed in any way relative to a reference molecule or a starting molecule.

(161) As such, polynucleotides encoding peptides or polypeptides containing substitutions, insertions and/or additions, deletions and covalent modifications with respect to reference sequences, in particular the polypeptide sequences disclosed herein, are included within the scope

of this disclosure. For example, sequence tags or amino acids, such as one or more lysines, can be added to peptide sequences (e.g., at the N-terminal or C-terminal ends). Sequence tags can be used for peptide detection, purification or localization. Lysines can be used to increase peptide solubility or to allow for biotinylation. Alternatively, amino acid residues located at the carboxy and amino terminal regions of the amino acid sequence of a peptide or protein may optionally be deleted providing for truncated sequences. Certain amino acids (e.g., C-terminal residues or N-terminal residues) alternatively may be deleted depending on the use of the sequence, as for example, expression of the sequence as part of a larger sequence that is soluble, or linked to a solid support.

(162) “Substitutional variants” when referring to polypeptides are those that have at least one amino acid residue in a native or starting sequence removed and a different amino acid inserted in its place at the same position. Substitutions may be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more (e.g., 3, 4 or 5) amino acids have been substituted in the same molecule.

(163) As used herein the term “conservative amino acid substitution” refers to the substitution of an amino acid that is normally present in the sequence with a different amino acid of similar size, charge, or polarity. Examples of conservative substitutions include the substitution of a non-polar (hydrophobic) residue such as isoleucine, valine and leucine for another non-polar residue.

Likewise, examples of conservative substitutions include the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, and between glycine and serine. Additionally, the substitution of a basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue such as aspartic acid or glutamic acid for another acidic residue are additional examples of conservative substitutions. Examples of non-conservative substitutions include the substitution of a non-polar (hydrophobic) amino acid residue such as isoleucine, valine, leucine, alanine, methionine for a polar (hydrophilic) residue such as cysteine, glutamine, glutamic acid or lysine and/or a polar residue for a non-polar residue.

(164) “Features” when referring to polypeptide or polynucleotide are defined as distinct amino acid sequence-based or nucleotide-based components of a molecule respectively. Features of the polypeptides encoded by the polynucleotides include surface manifestations, local conformational shape, folds, loops, half-loops, domains, half-domains, sites, termini and any combination(s) thereof.

(165) As used herein when referring to polypeptides the term “domain” refers to a motif of a polypeptide having one or more identifiable structural or functional characteristics or properties (e.g., binding capacity, serving as a site for protein-protein interactions).

(166) As used herein when referring to polypeptides the terms “site” as it pertains to amino acid based embodiments is used synonymously with “amino acid residue” and “amino acid side chain.” As used herein when referring to polynucleotides the terms “site” as it pertains to nucleotide based embodiments is used synonymously with “nucleotide.” A site represents a position within a peptide or polypeptide or polynucleotide that may be modified, manipulated, altered, derivatized or varied within the polypeptide-based or polynucleotide-based molecules.

(167) As used herein the terms “termini” or “terminus” when referring to polypeptides or polynucleotides refers to an extremity of a polypeptide or polynucleotide respectively. Such extremity is not limited only to the first or final site of the polypeptide or polynucleotide but may include additional amino acids or nucleotides in the terminal regions. Polypeptide-based molecules may be characterized as having both an N-terminus (terminated by an amino acid with a free amino group (NH₂)) and a C-terminus (terminated by an amino acid with a free carboxyl group (COOH)). Proteins are in some cases made up of multiple polypeptide chains brought together by disulfide bonds or by non-covalent forces (multimers, oligomers). These proteins have multiple N- and C-termini. Alternatively, the termini of the polypeptides may be modified such that they begin or end, as the case may be, with a non-polypeptide based moiety such as an organic conjugate.

(168) As recognized by those skilled in the art, protein fragments, functional protein domains, and

homologous proteins are also considered to be within the scope of polypeptides of interest. For example, provided herein is any protein fragment (meaning a polypeptide sequence at least one amino acid residue shorter than a reference polypeptide sequence but otherwise identical) of a reference protein having a length of 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or longer than 100 amino acids. In another example, any protein that includes a stretch of 20, 30, 40, 50, or 100 (contiguous) amino acids that are 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100% identical to any of the sequences described herein can be utilized in accordance with the disclosure. In some embodiments, a polypeptide includes 2, 3, 4, 5, 6, 7, 8, 9, 10, or more mutations as shown in any of the sequences provided herein or referenced herein. In another example, any protein that includes a stretch of 20, 30, 40, 50, or 100 amino acids that are greater than 80%, 90%, 95%, or 100% identical to any of the sequences described herein, wherein the protein has a stretch of 5, 10, 15, 20, 25, or 30 amino acids that are less than 80%, 75%, 70%, 65% to 60% identical to any of the sequences described herein can be utilized in accordance with the disclosure.

(169) Polypeptide or polynucleotide molecules of the present disclosure may share a certain degree of sequence similarity or identity with the reference molecules (e.g., reference polypeptides or reference polynucleotides), for example, with art-described molecules (e.g., engineered or designed molecules or wild-type molecules). The term “identity,” as known in the art, refers to a relationship between the sequences of two or more polypeptides or polynucleotides, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between two sequences as determined by the number of matches between strings of two or more amino acid residues or nucleic acid residues. Identity measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (e.g., “algorithms”). Identity of related peptides can be readily calculated by known methods. “% identity” as it applies to polypeptide or polynucleotide sequences is defined as the percentage of residues (amino acid residues or nucleic acid residues) in the candidate amino acid or nucleic acid sequence that are identical with the residues in the amino acid sequence or nucleic acid sequence of a second sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent identity.

Methods and computer programs for the alignment are well known in the art. Identity depends on a calculation of percent identity but may differ in value due to gaps and penalties introduced in the calculation. Generally, variants of a particular polynucleotide or polypeptide have at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% but less than 100% sequence identity to that particular reference polynucleotide or polypeptide as determined by sequence alignment programs and parameters described herein and known to those skilled in the art. Such tools for alignment include those of the BLAST suite (Stephen F. Altschul, et al. (1997).” Gapped BLAST and PSI-BLAST: a new generation of protein database search programs,” *Nucleic Acids Res.* 25:3389-3402). Another popular local alignment technique is based on the Smith-Waterman algorithm (Smith, T. F. & Waterman, M. S. (1981) “Identification of common molecular subsequences.” *J. Mol. Biol.* 147:195-197). A general global alignment technique based on dynamic programming is the Needleman-Wunsch algorithm (Needleman, S. B. & Wunsch, C. D. (1970) “A general method applicable to the search for similarities in the amino acid sequences of two proteins.” *J. Mol. Biol.* 48:443-453). More recently, a Fast Optimal Global Sequence Alignment Algorithm (FOGSAA) was developed that purportedly produces global alignment of nucleotide and protein sequences faster than other optimal global alignment methods, including the Needleman-Wunsch algorithm. Other tools are described herein, specifically in the definition of “identity” below.

(170) As used herein, the term “homology” refers to the overall relatedness between polymeric molecules, e.g. between nucleic acid molecules (e.g. DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Polymeric molecules (e.g. nucleic acid molecules (e.g. DNA molecules and/or RNA molecules) and/or polypeptide molecules) that share a threshold level

of similarity or identity determined by alignment of matching residues are termed homologous. Homology is a qualitative term that describes a relationship between molecules and can be based upon the quantitative similarity or identity. Similarity or identity is a quantitative term that defines the degree of sequence match between two compared sequences. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identical or similar. The term “homologous” necessarily refers to a comparison between at least two sequences (polynucleotide or polypeptide sequences). Two polynucleotide sequences are considered homologous if the polypeptides they encode are at least 50%, 60%, 70%, 80%, 90%, 95%, or even 99% for at least one stretch of at least 20 amino acids. In some embodiments, homologous polynucleotide sequences are characterized by the ability to encode a stretch of at least 4-5 uniquely specified amino acids. For polynucleotide sequences less than 60 nucleotides in length, homology is determined by the ability to encode a stretch of at least 4-5 uniquely specified amino acids. Two protein sequences are considered homologous if the proteins are at least 50%, 60%, 70%, 80%, or 90% identical for at least one stretch of at least 20 amino acids.

(171) Homology implies that the compared sequences diverged in evolution from a common origin. The term “homolog” refers to a first amino acid sequence or nucleic acid sequence (e.g., gene (DNA or RNA) or protein sequence) that is related to a second amino acid sequence or nucleic acid sequence by descent from a common ancestral sequence. The term “homolog” may apply to the relationship between genes and/or proteins separated by the event of speciation or to the relationship between genes and/or proteins separated by the event of genetic duplication.

“Orthologs” are genes (or proteins) in different species that evolved from a common ancestral gene (or protein) by speciation. Typically, orthologs retain the same function in the course of evolution. “Paralogs” are genes (or proteins) related by duplication within a genome. Orthologs retain the same function in the course of evolution, whereas paralogs evolve new functions, even if these are related to the original one.

(172) The term “identity” refers to the overall relatedness between polymeric molecules, for example, between polynucleotide molecules (e.g. DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of the percent identity of two polynucleic acid sequences, for example, can be performed by aligning the two sequences for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second nucleic acid sequences for optimal alignment and non-identical sequences can be disregarded for comparison purposes). In certain embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% of the length of the reference sequence. The nucleotides at corresponding nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. For example, the percent identity between two nucleic acid sequences can be determined using methods such as those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; each of which is incorporated herein by reference. For example, the percent identity between two nucleic acid sequences can be determined using the algorithm of Meyers and Miller (CABIOS,

1989, 4:11-17), which has been incorporated into the ALIGN program (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. The percent identity between two nucleic acid sequences can, alternatively, be determined using the GAP program in the GCG software package using an NWSgapdna.CMP matrix. Methods commonly employed to determine percent identity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., *SIAM J Applied Math.*, 48:1073 (1988); incorporated herein by reference. Techniques for determining identity are codified in publicly available computer programs. Exemplary computer software to determine homology between two sequences include, but are not limited to, GCG program package, Devereux, J., et al., *Nucleic Acids Research*, 12(1), 387 (1984)), BLASTP, BLASTN, and FASTA Altschul, S. F. et al., *J. Molec. Biol.*, 215, 403 (1990)).

(173) Multiprotein and Multicomponent Vaccines

(174) The present disclosure encompasses respiratory virus vaccines comprising multiple RNA (e.g., mRNA) polynucleotides, each encoding a single antigenic polypeptide, as well as respiratory virus vaccines comprising a single RNA polynucleotide encoding more than one antigenic polypeptide (e.g., as a fusion polypeptide). Thus, a vaccine composition comprising a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a first antigenic polypeptide and a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a second antigenic polypeptide encompasses (a) vaccines that comprise a first RNA polynucleotide encoding a first antigenic polypeptide and a second RNA polynucleotide encoding a second antigenic polypeptide, and (b) vaccines that comprise a single RNA polynucleotide encoding a first and second antigenic polypeptide (e.g., as a fusion polypeptide). RNA (e.g., mRNA) vaccines of the present disclosure, in some embodiments, comprise 2-10 (e.g., 2, 3, 4, 5, 6, 7, 8, 9 or 10), or more, RNA polynucleotides having an open reading frame, each of which encodes a different antigenic polypeptide (or a single RNA polynucleotide encoding 2-10, or more, different antigenic polypeptides). The antigenic polypeptides may be selected from hMPV, PIV3, RSV, MEV and BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1) antigenic polypeptides.

(175) In some embodiments, a respiratory virus vaccine comprises a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a viral capsid protein, a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a viral premembrane/membrane protein, and a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a viral envelope protein. In some embodiments, a respiratory virus vaccine comprises a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a viral fusion (F) protein and a RNA polynucleotide having an open reading frame encoding a viral major surface glycoprotein (G protein). In some embodiments, a vaccine comprises a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a viral F protein. In some embodiments, a vaccine comprises a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a viral G protein. In some embodiments, a vaccine comprises a RNA (e.g., mRNA) polynucleotide having an open reading frame encoding a HN protein.

(176) In some embodiments, a multicomponent vaccine comprises at least one RNA (e.g., mRNA) polynucleotide encoding at least one antigenic polypeptide fused to a signal peptide (e.g., any one of SEQ ID NO: 15-19). The signal peptide may be fused at the N-terminus or the C-terminus of an antigenic polypeptide. An antigenic polypeptide fused to a signal peptide may be selected from hMPV, PIV3, RSV, MEV and BetaCoV (e.g., selected from MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and HCoV—HKU1) antigenic polypeptides.

(177) Signal Peptides

(178) In some embodiments, antigenic polypeptides encoded by respiratory virus RNA (e.g., mRNA) polynucleotides comprise a signal peptide. Signal peptides, comprising the N-terminal 15-

60 amino acids of proteins, are typically needed for the translocation across the membrane on the secretory pathway and, thus, universally control the entry of most proteins both in eukaryotes and prokaryotes to the secretory pathway. Signal peptides generally include three regions: an N-terminal region of differing length, which usually comprises positively charged amino acids; a hydrophobic region; and a short carboxy-terminal peptide region. In eukaryotes, the signal peptide of a nascent precursor protein (pre-protein) directs the ribosome to the rough endoplasmic reticulum (ER) membrane and initiates the transport of the growing peptide chain across it for processing. ER processing produces mature proteins, wherein the signal peptide is cleaved from precursor proteins, typically by a ER-resident signal peptidase of the host cell, or they remain uncleaved and function as a membrane anchor. A signal peptide may also facilitate the targeting of the protein to the cell membrane. The signal peptide, however, is not responsible for the final destination of the mature protein. Secretory proteins devoid of additional address tags in their sequence are by default secreted to the external environment. During recent years, a more advanced view of signal peptides has evolved, showing that the functions and immunodominance of certain signal peptides are much more versatile than previously anticipated.

(179) Respiratory virus vaccines of the present disclosure may comprise, for example, RNA (e.g., mRNA) polynucleotides encoding an artificial signal peptide, wherein the signal peptide coding sequence is operably linked to and is in frame with the coding sequence of the antigenic polypeptide. Thus, respiratory virus vaccines of the present disclosure, in some embodiments, produce an antigenic polypeptide comprising an antigenic polypeptide (e.g., hMPV, PIV3, RSV, MeV or BetaCoV) fused to a signal peptide. In some embodiments, a signal peptide is fused to the N-terminus of the antigenic polypeptide. In some embodiments, a signal peptide is fused to the C-terminus of the antigenic polypeptide.

(180) In some embodiments, the signal peptide fused to the antigenic polypeptide is an artificial signal peptide. In some embodiments, an artificial signal peptide fused to the antigenic polypeptide encoded by the RNA (e.g., mRNA) vaccine is obtained from an immunoglobulin protein, e.g., an IgE signal peptide or an IgG signal peptide. In some embodiments, a signal peptide fused to the antigenic polypeptide encoded by a RNA (e.g., mRNA) vaccine is an Ig heavy chain epsilon-1 signal peptide (IgE HC SP) having the sequence of: MDWTWILFLVAAATRVHS (SEQ ID NO: 16). In some embodiments, a signal peptide fused to the antigenic polypeptide encoded by the (e.g., mRNA) RNA (e.g., mRNA) vaccine is an IgGk chain V-III region HAH signal peptide (IgGk SP) having the sequence of METPAQLLFLLLWLPDTTG (SEQ ID NO: 15). In some embodiments, the signal peptide is selected from: Japanese encephalitis PRM signal sequence (MLGSNSGQRVVFTILLLLVAPAYS; SEQ ID NO: 17), VSVg protein signal sequence (MKCLLYLAFLFIGVNCA; SEQ ID NO: 18) and Japanese encephalitis JEV signal sequence (MWLVSLAIVTACAGA; SEQ ID NO: 19).

(181) In some embodiments, the antigenic polypeptide encoded by a RNA (e.g., mRNA) vaccine comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8, 12-13, 24-34, 47-50 or 54-56 (Tables 3, 6, 11, 14 or 17; see also amino acid sequences of Tables 4, 7, 12 or 15) fused to a signal peptide identified by any one of SEQ ID NO: 15-19 (Table 8). The examples disclosed herein are not meant to be limiting and any signal peptide that is known in the art to facilitate targeting of a protein to ER for processing and/or targeting of a protein to the cell membrane may be used in accordance with the present disclosure.

(182) A signal peptide may have a length of 15-60 amino acids. For example, a signal peptide may have a length of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, or 60 amino acids. In some embodiments, a signal peptide has a length of 20-60, 25-60, 30-60, 35-60, 40-60, 45-60, 50-60, 55-60, 15-55, 20-55, 25-55, 30-55, 35-55, 40-55, 45-55, 50-55, 15-50, 20-50, 25-50, 30-50, 35-50, 40-50, 45-50, 15-45, 20-45, 25-45, 30-45, 35-45, 40-45, 15-40, 20-40, 25-40, 30-40, 35-40, 15-35, 20-35, 25-35, 30-35, 15-30, 20-30, 25-30, 15-25, 20-25, or 15-20 amino acids.

(183) A signal peptide is typically cleaved from the nascent polypeptide at the cleavage junction during ER processing. The mature antigenic polypeptide produced by a respiratory virus RNA (e.g., mRNA) vaccine of the present disclosure typically does not comprise a signal peptide.

(184) Chemical Modifications

(185) Respiratory virus vaccines of the present disclosure, in some embodiments, comprise at least RNA (e.g. mRNA) polynucleotide having an open reading frame encoding at least one antigenic polypeptide that comprises at least one chemical modification.

(186) The terms “chemical modification” and “chemically modified” refer to modification with respect to adenosine (A), guanosine (G), uridine (U), thymidine (T) or cytidine (C) ribonucleosides or deoxyribonucleosides in at least one of their position, pattern, percent or population. Generally, these terms do not refer to the ribonucleotide modifications in naturally occurring 5'-terminal mRNA cap moieties. With respect to a polypeptide, the term “modification” refers to a modification relative to the canonical set 20 amino acids. Polypeptides, as provided herein, are also considered “modified” if they contain amino acid substitutions, insertions or a combination of substitutions and insertions.

(187) Polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides), in some embodiments, comprise various (more than one) different modifications. In some embodiments, a particular region of a polynucleotide contains one, two or more (optionally different) nucleoside or nucleotide modifications. In some embodiments, a modified RNA polynucleotide (e.g., a modified mRNA polynucleotide), introduced to a cell or organism, exhibits reduced degradation in the cell or organism, respectively, relative to an unmodified polynucleotide. In some embodiments, a modified RNA polynucleotide (e.g., a modified mRNA polynucleotide), introduced into a cell or organism, may exhibit reduced immunogenicity in the cell or organism, respectively (e.g., a reduced innate response).

(188) Modifications of polynucleotides include, without limitation, those described herein.

Polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) may comprise modifications that are naturally-occurring, non-naturally-occurring or the polynucleotide may comprise a combination of naturally-occurring and non-naturally-occurring modifications. Polynucleotides may include any useful modification, for example, of a sugar, a nucleobase, or an internucleoside linkage (e.g., to a linking phosphate, to a phosphodiester linkage or to the phosphodiester backbone).

(189) Polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides), in some embodiments, comprise non-natural modified nucleotides that are introduced during synthesis or post-synthesis of the polynucleotides to achieve desired functions or properties. The modifications may be present on an internucleotide linkages, purine or pyrimidine bases, or sugars. The modification may be introduced with chemical synthesis or with a polymerase enzyme at the terminal of a chain or anywhere else in the chain. Any of the regions of a polynucleotide may be chemically modified.

(190) The present disclosure provides for modified nucleosides and nucleotides of a polynucleotide (e.g., RNA polynucleotides, such as mRNA polynucleotides). A “nucleoside” refers to a compound containing a sugar molecule (e.g., a pentose or ribose) or a derivative thereof in combination with an organic base (e.g., a purine or pyrimidine) or a derivative thereof (also referred to herein as “nucleobase”). A nucleotide” refers to a nucleoside, including a phosphate group. Modified nucleotides may be synthesized by any useful method, such as, for example, chemically, enzymatically, or recombinantly, to include one or more modified or non-natural nucleosides. Polynucleotides may comprise a region or regions of linked nucleosides. Such regions may have variable backbone linkages. The linkages may be standard phosphodiester linkages, in which case the polynucleotides would comprise regions of nucleotides.

(191) Modified nucleotide base pairing encompasses not only the standard adenosine-thymine, adenosine-uracil, or guanosine-cytosine base pairs, but also base pairs formed between nucleotides

and/or modified nucleotides comprising non-standard or modified bases, wherein the arrangement of hydrogen bond donors and hydrogen bond acceptors permits hydrogen bonding between a non-standard base and a standard base or between two complementary non-standard base structures. One example of such non-standard base pairing is the base pairing between the modified nucleotide inosine and adenine, cytosine or uracil. Any combination of base/sugar or linker may be incorporated into polynucleotides of the present disclosure.

(192) Modifications of polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) that are useful in the vaccines of the present disclosure include, but are not limited to the following: 2-methylthio-N6-(cis-hydroxyisopentenyl)adenosine; 2-methylthio-N6-methyladenosine; 2-methylthio-N6-threonyl carbamoyladenosine; N6-glyciny carbamoyladenosine; N6-isopentenyladenosine; N6-methyladenosine; N6-threonyl carbamoyladenosine; 1,2'-O-dimethyladenosine; 1-methyladenosine; 2'-O-methyladenosine; 2'-O-ribosyladenosine (phosphate); 2-methyladenosine; 2-methylthio-N6 isopentenyladenosine; 2-methylthio-N6-hydroxynorvalyl carbamoyladenosine; 2'-O-methyladenosine; 2'-O-ribosyladenosine (phosphate); Isopentenyladenosine; N6-(cis-hydroxyisopentenyl)adenosine; N6,2'-O-dimethyladenosine; N6,2'-O-dimethyladenosine; N6,N6,2'-O-trimethyladenosine; N6,N6-dimethyladenosine; N6-acetyladenosine; N6-hydroxynorvalyl carbamoyladenosine; N6-methyl-N6-threonyl carbamoyladenosine; 2-methyladenosine; 2-methylthio-N6-isopentenyladenosine; 7-deaza-adenosine; N1-methyladenosine; N6, N6 (dimethyl)adenine; N6-cis-hydroxy-isopentenyl-adenosine; α -thio-adenosine; 2 (amino)adenine; 2 (aminopropyl)adenine; 2 (methylthio) N6 (isopentenyl)adenine; 2-(alkyl)adenine; 2-(aminoalkyl)adenine; 2-(aminopropyl)adenine; 2-(halo)adenine; 2-(halo)adenine; 2-(propyl)adenine; 2'-Amino-2'-deoxy-ATP; 2'-Azido-2'-deoxy-ATP; 2'-Deoxy-2'-a-aminoadenosine TP; 2'-Deoxy-2'-a-azidoadenosine TP; 6 (alkyl)adenine; 6 (methyl)adenine; 6-(alkyl)adenine; 6-(methyl)adenine; 7 (deaza)adenine; 8 (alkenyl)adenine; 8 (alkynyl)adenine; 8 (amino)adenine; 8 (thioalkyl)adenine; 8-(alkenyl)adenine; 8-(alkyl)adenine; 8-(alkynyl)adenine; 8-(amino)adenine; 8-(halo)adenine; 8-(hydroxyl)adenine; 8-(thioalkyl)adenine; 8-(thiol)adenine; 8-azido-adenosine; aza adenine; deaza adenine; N6 (methyl)adenine; N6-(isopentyl)adenine; 7-deaza-8-aza-adenosine; 7-methyladenine; 1-Deazaadenosine TP; 2'Fluoro-N6-Bz-deoxyadenosine TP; 2'-OMe-2-Amino-ATP; 2'O-methyl-N6-Bz-deoxyadenosine TP; 2'-a-Ethynyladenosine TP; 2-aminoadenine; 2-Aminoadenosine TP; 2-Amino-ATP; 2'-a-Trifluoromethyladenosine TP; 2-Azidoadenosine TP; 2'-b-Ethynyladenosine TP; 2-Bromoadenosine TP; 2'-b-Trifluoromethyladenosine TP; 2-Chloroadenosine TP; 2'-Deoxy-2',2'-difluoroadenosine TP; 2'-Deoxy-2'-a-mercaptoadenosine TP; 2'-Deoxy-2'-a-thiomethoxyadenosine TP; 2'-Deoxy-2'-b-aminoadenosine TP; 2'-Deoxy-2'-b-azidoadenosine TP; 2'-Deoxy-2'-b-bromoadenosine TP; 2'-Deoxy-2'-b-chloroadenosine TP; 2'-Deoxy-2'-b-fluoroadenosine TP; 2'-Deoxy-2'-b-iodoadenosine TP; 2'-Deoxy-2'-b-mercaptoadenosine TP; 2'-Deoxy-2'-b-thiomethoxyadenosine TP; 2-Fluoroadenosine TP; 2-Iodoadenosine TP; 2-Mercaptoadenosine TP; 2-methoxy-adenine; 2-methylthio-adenine; 2-Trifluoromethyladenosine TP; 3-Deaza-3-bromoadenosine TP; 3-Deaza-3-chloroadenosine TP; 3-Deaza-3-fluoroadenosine TP; 3-Deaza-3-iodoadenosine TP; 3-Deazaadenosine TP; 4'-Azidoadenosine TP; 4'-Carbocyclic adenosine TP; 4'-Ethynyladenosine TP; 5'-Homo-adenosine TP; 8-Aza-ATP; 8-bromo-adenosine TP; 8-Trifluoromethyladenosine TP; 9-Deazaadenosine TP; 2-aminopurine; 7-deaza-2,6-diaminopurine; 7-deaza-8-aza-2,6-diaminopurine; 7-deaza-8-aza-2-aminopurine; 2,6-diaminopurine; 7-deaza-8-aza-adenine, 7-deaza-2-aminopurine; 2-thiocytidine; 3-methylcytidine; 5-formylcytidine; 5-hydroxymethylcytidine; 5-methylcytidine; N4-acetylcytidine; 2'-O-methylcytidine; 2'-O-methylcytidine; 5,2'-O-dimethylcytidine; 5-formyl-2'-O-methylcytidine; Lysidine; N4,2'-O-dimethylcytidine; N4-acetyl-2'-O-methylcytidine; N4-methylcytidine; N4,N4-Dimethyl-2'-OMe-Cytidine TP; 4-methylcytidine; 5-aza-cytidine; Pseudo-iso-cytidine; pyrrolo-cytidine; α -thio-cytidine; 2-(thio)cytosine; 2'-Amino-2'-deoxy-CTP; 2'-Azido-2'-deoxy-CTP; 2'-Deoxy-2'-a-aminocytidine TP; 2'-Deoxy-2'-a-azidocytidine TP; 3 (deaza) 5

(aza)cytosine; 3 (methyl)cytosine; 3-(alkyl)cytosine; 3-(deaza) 5 (aza)cytosine; 3-(methyl)cytidine; 4,2'-O-dimethylcytidine; 5 (halo)cytosine; 5 (methyl)cytosine; 5 (propynyl)cytosine; 5 (trifluoromethyl)cytosine; 5-(alkyl)cytosine; 5-(alkynyl)cytosine; 5-(halo)cytosine; 5-(propynyl)cytosine; 5-(trifluoromethyl)cytosine; 5-bromo-cytidine; 5-iodo-cytidine; 5-propynyl cytosine; 6-(azo)cytosine; 6-aza-cytidine; aza cytosine; deaza cytosine; N4 (acetyl)cytosine; 1-methyl-1-deaza-pseudoisocytidine; 1-methyl-pseudoisocytidine; 2-methoxy-5-methyl-cytidine; 2-methoxy-cytidine; 2-thio-5-methyl-cytidine; 4-methoxy-1-methyl-pseudoisocytidine; 4-methoxy-pseudoisocytidine; 4-thio-1-methyl-1-deaza-pseudoisocytidine; 4-thio-1-methyl-pseudoisocytidine; 4-thio-pseudoisocytidine; 5-aza-zebularine; 5-methyl-zebularine; pyrrolo-pseudoisocytidine; Zebularine; (E)-5-(2-Bromo-vinyl)cytidine TP; 2,2'-anhydro-cytidine TP hydrochloride; 2'Fluoro-N4-Bz-cytidine TP; 2'Fluoro-N4-Acetyl-cytidine TP; 2'-O-Methyl-N4-Acetyl-cytidine TP; 2'O-methyl-N4-Bz-cytidine TP; 2'-a-Ethynylcytidine TP; 2'-a-Trifluoromethylcytidine TP; 2'-b-Ethynylcytidine TP; 2'-b-Trifluoromethylcytidine TP; 2'-Deoxy-2',2'-difluorocytidine TP; 2'-Deoxy-2'-a-mercaptopcytidine TP; 2'-Deoxy-2'-a-thiomethoxycytidine TP; 2'-Deoxy-2'-b-aminocytidine TP; 2'-Deoxy-2'-b-azidocytidine TP; 2'-Deoxy-2'-b-bromocytidine TP; 2'-Deoxy-2'-b-chlorocytidine TP; 2'-Deoxy-2'-b-fluorocytidine TP; 2'-Deoxy-2'-b-iodocytidine TP; 2'-Deoxy-2'-b-mercaptopcytidine TP; 2'-Deoxy-2'-b-thiomethoxycytidine TP; 2'-O-Methyl-5-(1-propynyl)cytidine TP; 3'-Ethynylcytidine TP; 4'-Azidocytidine TP; 4'-Carbocyclic cytidine TP; 4'-Ethynylcytidine TP; 5-(1-Propynyl)ara-cytidine TP; 5-(2-Chloro-phenyl)-2-thiocytidine TP; 5-(4-Amino-phenyl)-2-thiocytidine TP; 5-Aminoallyl-CTP; 5-Cyanocytidine TP; 5-Ethynylara-cytidine TP; 5-Ethynylcytidine TP; 5'-Homo-cytidine TP; 5-Methoxycytidine TP; 5-Trifluoromethyl-Cytidine TP; N4-Amino-cytidine TP; N4-Benzoyl-cytidine TP; Pseudoisocytidine; 7-methylguanosine; N2,2'-O-dimethylguanosine; N2-methylguanosine; Wyosine; 1,2'-O-dimethylguanosine; 1-methylguanosine; 2'-O-methylguanosine; 2'-O-ribosylguanosine (phosphate); 2'-O-methylguanosine; 2'-O-ribosylguanosine (phosphate); 7-aminomethyl-7-deazaguanosine; 7-cyano-7-deazaguanosine; Archaeosine; Methylwyosine; N2,7-dimethylguanosine; N2,N2,2'-O-trimethylguanosine; N2,N2,7-trimethylguanosine; N2,N2-dimethylguanosine; N2,7,2'-O-trimethylguanosine; 6-thio-guanosine; 7-deaza-guanosine; 8-oxo-guanosine; N1-methyl-guanosine; α -thio-guanosine; 2 (propyl)guanine; 2-(alkyl)guanine; 2'-Amino-2'-deoxy-GTP; 2'-Azido-2'-deoxy-GTP; 2'-Deoxy-2'-a-aminoguanosine TP; 2'-Deoxy-2'-a-azidoguanosine TP; 6 (methyl)guanine; 6-(alkyl)guanine; 6-(methyl)guanine; 6-methyl-guanosine; 7 (alkyl)guanine; 7 (deaza)guanine; 7 (methyl)guanine; 7-(alkyl)guanine; 7-(deaza)guanine; 7-(methyl)guanine; 8 (alkyl)guanine; 8 (alkynyl)guanine; 8 (halo)guanine; 8 (thioalkyl)guanine; 8-(alkenyl)guanine; 8-(alkyl)guanine; 8-(alkynyl)guanine; 8-(amino)guanine; 8-(halo)guanine; 8-(hydroxyl)guanine; 8-(thioalkyl)guanine; 8-(thiol)guanine; aza guanine; deaza guanine; N (methyl)guanine; N-(methyl)guanine; 1-methyl-6-thio-guanosine; 6-methoxy-guanosine; 6-thio-7-deaza-8-aza-guanosine; 6-thio-7-deaza-guanosine; 6-thio-7-methyl-guanosine; 7-deaza-8-aza-guanosine; 7-methyl-8-oxo-guanosine; N2,N2-dimethyl-6-thio-guanosine; N2-methyl-6-thio-guanosine; 1-Me-GTP; 2'Fluoro-N2-isobutyl-guanosine TP; 2'O-methyl-N2-isobutyl-guanosine TP; 2'-a-Ethynylguanosine TP; 2'-a-Trifluoromethylguanosine TP; 2'-b-Ethynylguanosine TP; 2'-b-Trifluoromethylguanosine TP; 2'-Deoxy-2',2'-difluoroguanosine TP; 2'-Deoxy-2'-a-mercaptopguanosine TP; 2'-Deoxy-2'-a-thiomethoxyguanosine TP; 2'-Deoxy-2'-b-aminoguanosine TP; 2'-Deoxy-2'-b-azidoguanosine TP; 2'-Deoxy-2'-b-bromoguanosine TP; 2'-Deoxy-2'-b-chloroguanosine TP; 2'-Deoxy-2'-b-fluoroguanosine TP; 2'-Deoxy-2'-b-iodoguanosine TP; 2'-Deoxy-2'-b-mercaptopguanosine TP; 2'-Deoxy-2'-b-thiomethoxyguanosine TP; 4'-Azidoguanosine TP; 4'-Carbocyclic guanosine TP; 4'-Ethynylguanosine TP; 5'-Homo-guanosine TP; 8-bromo-guanosine TP; 9-Deazaguanosine TP; N2-isobutyl-guanosine TP; 1-methylinosine; Inosine; 1,2'-O-dimethylinosine; 2'-O-methylinosine; 7-methylinosine; 2'-O-methylinosine; Epoxyqueuosine; galactosyl-queuosine; Mannosylqueuosine; Queuosine; allylamino-thymidine; aza thymidine; deaza thymidine; deoxy-thymidine; 2'-O-methyluridine; 2-thiouridine; 3-methyluridine; 5-

carboxymethyluridine; 5-hydroxyuridine; 5-methyluridine; 5-taurinomethyl-2-thiouridine; 5-taurinomethyluridine; Dihydrouridine; Pseudouridine; (3-(3-amino-3-carboxypropyl)uridine; 1-methyl-3-(3-amino-5-carboxypropyl)pseudouridine; 1-methylpseudouridine; 1-methylpseudouridine; 2'-O-methyluridine; 2'-O-methylpseudouridine; 2'-O-methyluridine; 2-thio-2'-O-methyluridine; 3-(3-amino-3-carboxypropyl)uridine; 3,2'-O-dimethyluridine; 3-Methyl-pseudo-Uridine TP; 4-thiouridine; 5-(carboxyhydroxymethyl)uridine; 5-(carboxyhydroxymethyl)uridine methyl ester; 5,2'-O-dimethyluridine; 5,6-dihydro-uridine; 5-aminomethyl-2-thiouridine; 5-carbamoylmethyl-2'-O-methyluridine; 5-carbamoylmethyluridine; 5-carboxyhydroxymethyluridine; 5-carboxyhydroxymethyluridine methyl ester; 5-carboxymethylaminomethyl-2'-O-methyluridine; 5-carboxymethylaminomethyl-2-thiouridine; 5-carboxymethylaminomethyl-2-thiouridine; 5-carboxymethylaminomethyluridine; 5-carboxymethylaminomethyluridine; 5-Carbamoylmethyluridine TP; 5-methoxycarbonylmethyl-2'-O-methyluridine; 5-methoxycarbonylmethyl-2-thiouridine; 5-methoxycarbonylmethyluridine; 5-methoxyuridine; 5-methyl-2-thiouridine; 5-methylaminomethyl-2-selenouridine; 5-methylaminomethyl-2-thiouridine; 5-methylaminomethyluridine; 5-Methyldihydrouridine; 5-Oxyacetic acid-Uridine TP; 5-Oxyacetic acid-methyl ester-Uridine TP; N1-methyl-pseudo-uridine; uridine 5-oxyacetic acid; uridine 5-oxyacetic acid methyl ester; 3-(3-Amino-3-carboxypropyl)-Uridine TP; 5-(iso-Pentenylaminomethyl)-2-thiouridine TP; 5-(iso-Pentenylaminomethyl)-2'-O-methyluridine TP; 5-(iso-Pentenylaminomethyl)uridine TP; 5-propynyl uracil; α -thio-uridine; 1 (aminoalkylamino-carbonylethylenyl)-2(thio)-pseudouracil; 1 (aminoalkylaminocarbonylethylenyl)-2,4-(dithio)pseudouracil; 1 (aminoalkylaminocarbonylethylenyl)-4 (thio)pseudouracil; 1 (aminoalkylaminocarbonylethylenyl)-pseudouracil; 1 (aminocarbonylethylenyl)-2(thio)-pseudouracil; 1 (aminocarbonylethylenyl)-2,4-(dithio)pseudouracil; 1 (aminocarbonylethylenyl)-4 (thio)pseudouracil; 1 (aminocarbonylethylenyl)-pseudouracil; 1 substituted 2(thio)-pseudouracil; 1 substituted 2,4-(dithio)pseudouracil; 1 substituted 4 (thio)pseudouracil; 1 substituted pseudouracil; 1-(aminoalkylamino-carbonylethylenyl)-2-(thio)-pseudouracil; 1-Methyl-3-(3-amino-3-carboxypropyl) pseudouridine TP; 1-Methyl-3-(3-amino-3-carboxypropyl)pseudo-UTP; 1-Methyl-pseudo-UTP; 2 (thio)pseudouracil; 2' deoxy uridine; 2' fluorouridine; 2-(thio)uracil; 2,4-(dithio)pseudouracil; 2' methyl, 2' amino, 2' azido, 2' fluoro-guanosine; 2'-Amino-2'-deoxy-UTP; 2'-Azido-2'-deoxy-UTP; 2'-Azido-deoxyuridine TP; 2'-O-methylpseudouridine; 2' deoxy uridine; 2' fluorouridine; 2'-Deoxy-2'-a-aminouridine TP; 2'-Deoxy-2'-a-azidouridine TP; 2-methylpseudouridine; 3 (3 amino-3 carboxypropyl)uracil; 4 (thio)pseudouracil; 4-(thio)pseudouracil; 4-(thio)uracil; 4-thiouracil; 5 (1,3-diazole-1-alkyl)uracil; 5 (2-aminopropyl)uracil; 5 (aminoalkyl)uracil; 5 (dimethylaminoalkyl)uracil; 5 (guanidiniumalkyl)uracil; 5 (methoxycarbonylmethyl)-2-(thio)uracil; 5 (methoxycarbonylmethyl)uracil; 5 (methyl) 2 (thio)uracil; 5 (methyl) 2,4 (dithio)uracil; 5 (methyl) 4 (thio)uracil; 5 (methylaminomethyl)-2 (thio)uracil; 5 (methylaminomethyl)-2,4 (dithio)uracil; 5 (methylaminomethyl)-4 (thio)uracil; 5 (propynyl)uracil; 5 (trifluoromethyl)uracil; 5-(2-aminopropyl)uracil; 5-(alkyl)-2-(thio)pseudouracil; 5-(alkyl)-2,4 (dithio)pseudouracil; 5-(alkyl)-4 (thio)pseudouracil; 5-(alkyl)pseudouracil; 5-(alkyl)uracil; 5-(alkynyl)uracil; 5-(allylamino)uracil; 5-(cyanoalkyl)uracil; 5-(dialkylaminoalkyl)uracil; 5-(dimethylaminoalkyl)uracil; 5-(guanidiniumalkyl)uracil; 5-(halo)uracil; 5-(1,3-diazole-1-alkyl)uracil; 5-(methoxy)uracil; 5-(methoxycarbonylmethyl)-2-(thio)uracil; 5-(methoxycarbonylmethyl)uracil; 5-(methyl) 2(thio)uracil; 5-(methyl) 2,4 (dithio)uracil; 5-(methyl) 4 (thio)uracil; 5-(methyl)-2-(thio)pseudouracil; 5-(methyl)-2,4 (dithio)pseudouracil; 5-(methyl)-4 (thio)pseudouracil; 5-(methyl)pseudouracil; 5-(methylaminomethyl)-2 (thio)uracil; 5-(methylaminomethyl)-2,4(dithio)uracil; 5-(methylaminomethyl)-4-(thio)uracil; 5-(propynyl)uracil; 5-(trifluoromethyl)uracil; 5-aminoallyl-uridine; 5-bromo-uridine; 5-iodo-uridine; 5-uracil; 6 (azo)uracil; 6-(azo)uracil; 6-aza-uridine; allylamino-uracil; aza uracil; deaza uracil; N3

(methyl)uracil; Pseudo-UTP-1-2-ethanoic acid; Pseudouracil; 4-Thio-pseudo-UTP; 1-carboxymethyl-pseudouridine; 1-methyl-1-deaza-pseudouridine; 1-propynyl-uridine; 1-taurinomethyl-1-methyl-uridine; 1-taurinomethyl-4-thio-uridine; 1-taurinomethyl-pseudouridine; 2-methoxy-4-thio-pseudouridine; 2-thio-1-methyl-1-deaza-pseudouridine; 2-thio-1-methyl-pseudouridine; 2-thio-5-aza-uridine; 2-thio-dihydropseudouridine; 2-thio-dihydrouridine; 2-thio-pseudouridine; 4-methoxy-2-thio-pseudouridine; 4-methoxy-pseudouridine; 4-thio-1-methyl-pseudouridine; 4-thio-pseudouridine; 5-aza-uridine; Dihydropseudouridine; ($\pm$)1-(2-Hydroxypropyl)pseudouridine TP; (2R)-1-(2-Hydroxypropyl)pseudouridine TP; (2S)-1-(2-Hydroxypropyl)pseudouridine TP; (E)-5-(2-Bromo-vinyl)ara-uridine TP; (E)-5-(2-Bromo-vinyl)uridine TP; (Z)-5-(2-Bromo-vinyl)ara-uridine TP; (Z)-5-(2-Bromo-vinyl)uridine TP; 1-(2,2,2-Trifluoroethyl)-pseudo-UTP; 1-(2,2,3,3,3-Pentafluoropropyl)pseudouridine TP; 1-(2,2-Diethoxyethyl)pseudouridine TP; 1-(2,4,6-Trimethylbenzyl)pseudouridine TP; 1-(2,4,6-Trimethylbenzyl)pseudo-UTP; 1-(2,4,6-Trimethyl-phenyl)pseudo-UTP; 1-(2-Amino-2-carboxyethyl)pseudo-UTP; 1-(2-Amino-ethyl)pseudo-UTP; 1-(2-Hydroxyethyl)pseudouridine TP; 1-(2-Methoxyethyl)pseudouridine TP; 1-(3,4-Bis-trifluoromethoxybenzyl)pseudouridine TP; 1-(3,4-Dimethoxybenzyl)pseudouridine TP; 1-(3-Amino-3-carboxypropyl)pseudo-UTP; 1-(3-Aminopropyl)pseudo-UTP; 1-(3-Cyclopropyl-prop-2-ynyl)pseudouridine TP; 1-(4-Amino-4-carboxybutyl)pseudo-UTP; 1-(4-Amino-benzyl)pseudo-UTP; 1-(4-Amino-butyl)pseudo-UTP; 1-(4-Amino-phenyl)pseudo-UTP; 1-(4-Azidobenzyl)pseudouridine TP; 1-(4-Bromobenzyl)pseudouridine TP; 1-(4-Chlorobenzyl)pseudouridine TP; 1-(4-Fluorobenzyl)pseudouridine TP; 1-(4-Iodobenzyl)pseudouridine TP; 1-(4-Methanesulfonylbenzyl)pseudouridine TP; 1-(4-Methoxybenzyl)pseudouridine TP; 1-(4-Methoxybenzyl)pseudo-UTP; 1-(4-Methoxy-phenyl)pseudo-UTP; 1-(4-Methylbenzyl)pseudouridine TP; 1-(4-Methyl-benzyl)pseudo-UTP; 1-(4-Nitrobenzyl)pseudouridine TP; 1-(4-Nitro-benzyl)pseudo-UTP; 1-(4-Nitro-phenyl)pseudo-UTP; 1-(4-Thiomethoxybenzyl)pseudouridine TP; 1-(4-Trifluoromethoxybenzyl)pseudouridine TP; 1-(4-Trifluoromethylbenzyl)pseudouridine TP; 1-(5-Amino-pentyl)pseudo-UTP; 1-(6-Amino-hexyl)pseudo-UTP; 1,6-Dimethyl-pseudo-UTP; 1-[3-(2-{2-[2-(2-Aminoethoxy)-ethoxy]-ethoxy}-ethoxy)-propionyl]pseudouridine TP; 1-{3-[2-(2-Aminoethoxy)-ethoxy]-propionyl} pseudouridine TP; 1-Acetylpsudouridine TP; 1-Alkyl-6-(1-propynyl)-pseudo-UTP; 1-Alkyl-6-(2-propynyl)-pseudo-UTP; 1-Alkyl-6-allyl-pseudo-UTP; 1-Alkyl-6-ethynyl-pseudo-UTP; 1-Alkyl-6-homoallyl-pseudo-UTP; 1-Alkyl-6-vinyl-pseudo-UTP; 1-Allylpsudouridine TP; 1-Aminomethyl-pseudo-UTP; 1-Benzoylpsudouridine TP; 1-Benzyloxymethylpsudouridine TP; 1-Benzyl-pseudo-UTP; 1-Biotinyl-PEG2-pseudouridine TP; 1-Biotinypseudouridine TP; 1-Butyl-pseudo-UTP; 1-Cyanomethylpsudouridine TP; 1-Cyclobutylmethyl-pseudo-UTP; 1-Cyclobutyl-pseudo-UTP; 1-Cycloheptylmethyl-pseudo-UTP; 1-Cycloheptyl-pseudo-UTP; 1-Cyclohexylmethyl-pseudo-UTP; 1-Cyclohexyl-pseudo-UTP; 1-Cyclooctylmethyl-pseudo-UTP; 1-Cyclooctyl-pseudo-UTP; 1-Cyclopentylmethyl-pseudo-UTP; 1-Cyclopentyl-pseudo-UTP; 1-Cyclopropylmethyl-pseudo-UTP; 1-Cyclopropyl-pseudo-UTP; 1-Ethyl-pseudo-UTP; 1-Hexyl-pseudo-UTP; 1-Homoallylpsudouridine TP; 1-Hydroxymethylpsudouridine TP; 1-iso-propyl-pseudo-UTP; 1-Me-2-thio-pseudo-UTP; 1-Me-4-thio-pseudo-UTP; 1-Me-alpha-thio-pseudo-UTP; 1-Methanesulfonylmethylpsudouridine TP; 1-Methoxymethylpsudouridine TP; 1-Methyl-6-(2,2,2-Trifluoroethyl)pseudo-UTP; 1-Methyl-6-(4-morpholino)-pseudo-UTP; 1-Methyl-6-(4-thiomorpholino)-pseudo-UTP; 1-Methyl-6-(substituted phenyl)pseudo-UTP; 1-Methyl-6-amino-pseudo-UTP; 1-Methyl-6-azido-pseudo-UTP; 1-Methyl-6-bromo-pseudo-UTP; 1-Methyl-6-butyl-pseudo-UTP; 1-Methyl-6-chloro-pseudo-UTP; 1-Methyl-6-cyano-pseudo-UTP; 1-Methyl-6-dimethylamino-pseudo-UTP; 1-Methyl-6-ethoxy-pseudo-UTP; 1-Methyl-6-ethylcarboxylate-pseudo-UTP; 1-Methyl-6-ethyl-pseudo-UTP; 1-Methyl-6-fluoro-pseudo-UTP; 1-Methyl-6-formyl-pseudo-UTP; 1-Methyl-6-hydroxyamino-pseudo-UTP; 1-Methyl-6-hydroxy-pseudo-UTP; 1-Methyl-6-iodo-pseudo-UTP; 1-Methyl-6-iso-propyl-pseudo-UTP; 1-Methyl-6-methoxy-pseudo-UTP; 1-Methyl-6-methylamino-pseudo-UTP; 1-Methyl-6-phenyl-

pseudo-UTP; 1-Methyl-6-propyl-pseudo-UTP; 1-Methyl-6-tert-butyl-pseudo-UTP; 1-Methyl-6-trifluoromethoxy-pseudo-UTP; 1-Methyl-6-trifluoromethyl-pseudo-UTP; 1-Morpholinomethylpseudouridine TP; 1-Pentyl-pseudo-UTP; 1-Phenyl-pseudo-UTP; 1-Pivaloylpseudouridine TP; 1-Propargylpseudouridine TP; 1-Propyl-pseudo-UTP; 1-propynyl-pseudouridine; 1-p-tolyl-pseudo-UTP; 1-tert-Butyl-pseudo-UTP; 1-Thiomethoxymethylpseudouridine TP; 1-Thiomorpholinomethylpseudouridine TP; 1-Trifluoroacetyl-pseudouridine TP; 1-Trifluoromethyl-pseudo-UTP; 1-Vinylpseudouridine TP; 2,2'-anhydro-uridine TP; 2'-bromo-deoxyuridine TP; 2'-F-5-Methyl-2'-deoxy-UTP; 2'-OMe-5-Me-UTP; 2'-OMe-pseudo-UTP; 2'-a-Ethynyluridine TP; 2'-a-Trifluoromethyluridine TP; 2'-b-Ethynyluridine TP; 2'-b-Trifluoromethyluridine TP; 2'-Deoxy-2',2'-difluorouridine TP; 2'-Deoxy-2'-a-mercaptopuridine TP; 2'-Deoxy-2'-a-thiomethoxyuridine TP; 2'-Deoxy-2'-b-aminouridine TP; 2'-Deoxy-2'-b-azidouridine TP; 2'-Deoxy-2'-b-bromouridine TP; 2'-Deoxy-2'-b-chlorouridine TP; 2'-Deoxy-2'-b-fluorouridine TP; 2'-Deoxy-2'-b-iodouridine TP; 2'-Deoxy-2'-b-mercaptopuridine TP; 2'-Deoxy-2'-b-thiomethoxyuridine TP; 2-methoxy-4-thio-uridine; 2-methoxyuridine; 2'-O-Methyl-5-(1-propynyl)uridine TP; 3-Alkyl-pseudo-UTP; 4'-Azidouridine TP; 4'-Carbocyclic uridine TP; 4'-Ethynyluridine TP; 5-(1-Propynyl)ara-uridine TP; 5-(2-Furanyl)uridine TP; 5-Cyanouridine TP; 5-Dimethylaminouridine TP; 5'-Homo-uridine TP; 5-iodo-2'-fluoro-deoxyuridine TP; 5-Phenylethynyluridine TP; 5-Trideuteromethyl-6-deuterouridine TP; 5-Trifluoromethyl-Uridine TP; 5-Vinylarauridine TP; 6-(2,2,2-Trifluoroethyl)-pseudo-UTP; 6-(4-Morpholino)-pseudo-UTP; 6-(4-Thiomorpholino)-pseudo-UTP; 6-(Substituted-Phenyl)-pseudo-UTP; 6-Amino-pseudo-UTP; 6-Azido-pseudo-UTP; 6-Bromo-pseudo-UTP; 6-Butyl-pseudo-UTP; 6-Chloro-pseudo-UTP; 6-Cyano-pseudo-UTP; 6-Dimethylamino-pseudo-UTP; 6-Ethoxy-pseudo-UTP; 6-Ethylcarboxylate-pseudo-UTP; 6-Ethyl-pseudo-UTP; 6-Fluoro-pseudo-UTP; 6-Formyl-pseudo-UTP; 6-Hydroxyamino-pseudo-UTP; 6-Hydroxy-pseudo-UTP; 6-Iodo-pseudo-UTP; 6-iso-Propyl-pseudo-UTP; 6-Methoxy-pseudo-UTP; 6-Methylamino-pseudo-UTP; 6-Methyl-pseudo-UTP; 6-Phenyl-pseudo-UTP; 6-Phenyl-pseudo-UTP; 6-Propyl-pseudo-UTP; 6-tert-Butyl-pseudo-UTP; 6-Trifluoromethoxy-pseudo-UTP; 6-Trifluoromethyl-pseudo-UTP; Alpha-thio-pseudo-UTP; Pseudouridine 1-(4-methylbenzenesulfonic acid) TP; Pseudouridine 1-(4-methylbenzoic acid) TP; Pseudouridine TP 1-[3-(2-ethoxy)]propionic acid; Pseudouridine TP 1-[3-{2-(2-[2-(2-ethoxy)-ethoxy]-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-[3-{2-(2-[2-{2(2-ethoxy)-ethoxy}-ethoxy]-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-[3-{2-(2-[2-ethoxy]-ethoxy)-ethoxy}]propionic acid; Pseudouridine TP 1-[3-{2-(2-ethoxy)-ethoxy}] propionic acid; Pseudouridine TP 1-methylphosphonic acid; Pseudouridine TP 1-methylphosphonic acid diethyl ester; Pseudo-UTP-N1-3-propionic acid; Pseudo-UTP-N1-4-butanoic acid; Pseudo-UTP-N1-5-pentanoic acid; Pseudo-UTP-N1-6-hexanoic acid; Pseudo-UTP-N1-7-heptanoic acid; Pseudo-UTP-N1-methyl-p-benzoic acid; Pseudo-UTP-N1-p-benzoic acid; Wybutosine; Hydroxywybutosine; Isowyosine; Peroxywybutosine; undermodified hydroxywybutosine; 4-demethylwyosine; 2,6-(diamino)purine; 1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 1,3-(diaz)-2-(oxo)-phenthiazin-1-yl; 1,3-(diaz)-2-(oxo)-phenoxazin-1-yl; 1,3,5-(triaz)-2,6-(diox)-naphthalene; 2 (amino)purine; 2,4,5-(trimethyl)phenyl; 2' methyl, 2' amino, 2'azido, 2'fluro-cytidine; 2' methyl, 2' amino, 2'azido, 2'fluro-adenine; 2'methyl, 2' amino, 2'azido, 2'fluro-uridine; 2'-amino-2'-deoxyribose; 2-amino-6-Chloro-purine; 2-aza-inosinyl; 2'-azido-2'-deoxyribose; 2'fluoro-2'-deoxyribose; 2'-fluoro-modified bases; 2'-O-methyl-ribose; 2-oxo-7-aminopyridopyrimidin-3-yl; 2-oxo-pyridopyrimidine-3-yl; 2-pyridinone; 3 nitropyrrole; 3-(methyl)-7-(propynyl)isocarbostyrylyl; 3-(methyl)isocarbostyrylyl; 4-(fluoro)-6-(methyl)benzimidazole; 4-(methyl)benzimidazole; 4-(methyl)indolyl; 4,6-(dimethyl)indolyl; 5 nitroindole; 5 substituted pyrimidines; 5-(methyl)isocarbostyrylyl; 5-nitroindole; 6-(aza)pyrimidine; 6-(azo)thymine; 6-(methyl)-7-(aza)indolyl; 6-chloro-purine; 6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; 7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenthiazin-1-yl; 7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-(aminoalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenoxazin-1-yl; 7-(aminoalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenthiazin-1-yl; 7-

(aminoalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenoxazin-1-yl; 7-(aza)indolyl; 7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenthiazin-1-yl; 7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-(guanidiniumalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenoxazin-1-yl; 7-(guanidiniumalkylhydroxy)-1,3-(diaz)-2-(oxo)-phenthiazin-1-yl; 7-(propynyl)isocarbostyryl; 7-(propynyl)isocarbostyryl, propynyl-7-(aza)indolyl; 7-deaza-inosinyl; 7-substituted 1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl; 7-substituted 1,3-(diaz)-2-(oxo)-phenoxazin-1-yl; 9-(methyl)-imidizopyridinyl; Aminoindolyl; Anthracenyl; bis-ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; bis-ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; Difluorotolyl; Hypoxanthine; Imidizopyridinyl; Inosinyl; Isocarbostyryl; Isoguanisine; N2-substituted purines; N6-methyl-2-amino-purine; N6-substituted purines; N-alkylated derivative; Napthalenyl; Nitrobenzimidazolyl; Nitroimidazolyl; Nitroindazolyl; Nitropyrazolyl; Nubularine; O6-substituted purines; O-alkylated derivative; ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; Oxoformycin TP; para-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; para-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl; Pentacenyl; Phenanthracenyl; Phenyl; propynyl-7-(aza)indolyl; Pyrenyl; pyridopyrimidin-3-yl; pyridopyrimidin-3-yl, 2-oxo-7-amino-pyridopyrimidin-3-yl; pyrrolo-pyrimidin-2-on-3-yl; Pyrrolopyrimidinyl; Pyrrolopyrizinyl; Stilbenzyl; substituted 1,2,4-triazoles; Tetracenyl; Tubercidine; Xanthine; Xanthosine-5'-TP; 2-thio-zebularine; 5-aza-2-thio-zebularine; 7-deaza-2-amino-purine; pyridin-4-one ribonucleoside; 2-Amino-riboside-TP; Formycin A TP; Formycin B TP; Pyrrolosine TP; 2'-OH-ara-adenosine TP; 2'-OH-ara-cytidine TP; 2'-OH-ara-uridine TP; 2'-OH-ara-guanosine TP; 5-(2-carbomethoxyvinyl)uridine TP; and N6-(19-Amino-pentaoxanonadecyl)adenosine TP.

(193) In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) include a combination of at least two (e.g., 2, 3, 4 or more) of the aforementioned modified nucleobases.

(194) In some embodiments, modified nucleobases in polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) are selected from the group consisting of pseudouridine (ψ), N1-methylpseudouridine (m.sup.1 ψ), N1-ethylpseudouridine, 2-thiouridine, 4'-thiouridine, 5-methylcytosine, 2-thio-1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-pseudouridine, 2-thio-5-aza-uridine, 2-thio-dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-pseudouridine, 4-methoxy-2-thio-pseudouridine, 4-methoxy-pseudouridine, 4-thio-1-methyl-pseudouridine, 4-thio-pseudouridine, 5-aza-uridine, dihydropseudouridine, 5-methoxyuridine and 2'-O-methyl uridine. In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) include a combination of at least two (e.g., 2, 3, 4 or more) of the aforementioned modified nucleobases.

(195) In some embodiments, modified nucleobases in polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) are selected from the group consisting of 1-methyl-pseudouridine (m.sup.1 ψ), 5-methoxy-uridine (mo.sup.5U), 5-methyl-cytidine (m.sup.5C), pseudouridine (ψ), α -thio-guanosine and α -thio-adenosine. In some embodiments, polynucleotides includes a combination of at least two (e.g., 2, 3, 4 or more) of the aforementioned modified nucleobases.

(196) In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise pseudouridine (ψ) and 5-methyl-cytidine (m.sup.5C). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 1-methyl-pseudouridine (m.sup.1 ψ). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 1-methyl-pseudouridine (m.sup.1 ψ) and 5-methyl-cytidine (m.sup.5C). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 2-thiouridine (s.sup.2U). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 2-

thiouridine and 5-methyl-cytidine (m.sup.5C). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise methoxy-uridine (mo.sup.5U). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 5-methoxy-uridine (mo.sup.5U) and 5-methyl-cytidine (m.sup.5C). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 2'-O-methyl uridine. In some embodiments polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise 2'-O-methyl uridine and 5-methyl-cytidine (m.sup.5C). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise N6-methyl-adenosine (m.sup.6A). In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) comprise N6-methyl-adenosine (m.sup.6A) and 5-methyl-cytidine (m.sup.5C).

(197) In some embodiments, polynucleotides (e.g., RNA polynucleotides, such as mRNA polynucleotides) are uniformly modified (e.g., fully modified, modified throughout the entire sequence) for a particular modification. For example, a polynucleotide can be uniformly modified with 5-methyl-cytidine (m.sup.5C), meaning that all cytosine residues in the mRNA sequence are replaced with 5-methyl-cytidine (m.sup.5C). Similarly, a polynucleotide can be uniformly modified for any type of nucleoside residue present in the sequence by replacement with a modified residue such as those set forth above.

(198) Exemplary nucleobases and nucleosides having a modified cytosine include N4-acetyl-cytidine (ac4C), 5-methyl-cytidine (m5C), 5-halo-cytidine (e.g., 5-iodo-cytidine), 5-hydroxymethyl-cytidine (hm5C), 1-methyl-pseudoisocytidine, 2-thio-cytidine (s2C), and 2-thio-5-methyl-cytidine.

(199) In some embodiments, a modified nucleobase is a modified uridine. Exemplary nucleobases and In some embodiments, a modified nucleobase is a modified cytosine. nucleosides having a modified uridine include 5-cyano uridine, and 4'-thio uridine.

(200) In some embodiments, a modified nucleobase is a modified adenine. Exemplary nucleobases and nucleosides having a modified adenine include 7-deaza-adenine, 1-methyl-adenosine (m1A), 2-methyl-adenine (m2A), and N6-methyl-adenosine (m6A).

(201) In some embodiments, a modified nucleobase is a modified guanine. Exemplary nucleobases and nucleosides having a modified guanine include inosine (I), 1-methyl-inosine (m1I), wyosine (imG), methylwyosine (mimG), 7-deaza-guanosine, 7-cyano-7-deaza-guanosine (preQ0), 7-aminomethyl-7-deaza-guanosine (preQ1), 7-methyl-guanosine (m7G), 1-methyl-guanosine (m1G), 8-oxo-guanosine, 7-methyl-8-oxo-guanosine.

(202) The polynucleotides of the present disclosure may be partially or fully modified along the entire length of the molecule. For example, one or more or all or a given type of nucleotide (e.g., purine or pyrimidine, or any one or more or all of A, G, U, C) may be uniformly modified in a polynucleotide of the disclosure, or in a given predetermined sequence region thereof (e.g., in the mRNA including or excluding the polyA tail). In some embodiments, all nucleotides X in a polynucleotide of the present disclosure (or in a given sequence region thereof) are modified nucleotides, wherein X may any one of nucleotides A, G, U, C, or any one of the combinations A+G, A+U, A+C, G+U, G+C, U+C, A+G+U, A+G+C, G+U+C or A+G+C.

(203) The polynucleotide may contain from about 1% to about 100% modified nucleotides (either in relation to overall nucleotide content, or in relation to one or more types of nucleotide, i.e., any one or more of A, G, U or C) or any intervening percentage (e.g., from 1% to 20%, from 1% to 25%, from 1% to 50%, from 1% to 60%, from 1% to 70%, from 1% to 80%, from 1% to 90%, from 1% to 95%, from 10% to 20%, from 10% to 25%, from 10% to 50%, from 10% to 60%, from 10% to 70%, from 10% to 80%, from 10% to 90%, from 10% to 95%, from 10% to 100%, from 20% to 25%, from 20% to 50%, from 20% to 60%, from 20% to 70%, from 20% to 80%, from 20% to 90%, from 20% to 95%, from 20% to 100%, from 50% to 60%, from 50% to 70%, from 50% to 80%, from 50% to 90%, from 50% to 95%, from 50% to 100%, from 70% to 80%, from

70% to 90%, from 70% to 95%, from 70% to 100%, from 80% to 90%, from 80% to 95%, from 80% to 100%, from 90% to 95%, from 90% to 100%, and from 95% to 100%). Any remaining percentage is accounted for by the presence of unmodified A, G, U, or C.

(204) The polynucleotides may contain at a minimum 1% and at maximum 100% modified nucleotides, or any intervening percentage, such as at least 5% modified nucleotides, at least 10% modified nucleotides, at least 25% modified nucleotides, at least 50% modified nucleotides, at least 80% modified nucleotides, or at least 90% modified nucleotides. For example, the polynucleotides may contain a modified pyrimidine such as a modified uracil or cytosine. In some embodiments, at least 5%, at least 10%, at least 25%, at least 50%, at least 80%, at least 90% or 100% of the uracil in the polynucleotide is replaced with a modified uracil (e.g., a 5-substituted uracil). The modified uracil can be replaced by a compound having a single unique structure, or can be replaced by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures). In some embodiments, at least 5%, at least 10%, at least 25%, at least 50%, at least 80%, at least 90% or 100% of the cytosine in the polynucleotide is replaced with a modified cytosine (e.g., a 5-substituted cytosine). The modified cytosine can be replaced by a compound having a single unique structure, or can be replaced by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures).

(205) Thus, in some embodiments, the RNA (e.g., mRNA) vaccines comprise a 5'UTR element, an optionally codon optimized open reading frame, and a 3'UTR element, a poly(A) sequence and/or a polyadenylation signal wherein the RNA is not chemically modified.

(206) In some embodiments, the modified nucleobase is a modified uracil. Exemplary nucleobases and nucleosides having a modified uracil include pseudouridine (ψ), pyridin-4-one ribonucleoside, 5-aza-uridine, 6-aza-uridine, 2-thio-5-aza-uridine, 2-thio-uridine (s.sup.2U), 4-thio-uridine (s.sup.4U), 4-thio-pseudouridine, 2-thio-pseudouridine, 5-hydroxy-uridine (hosU), 5-aminoallyl-uridine, 5-halo-uridine (e.g., 5-iodo-uridine or 5-bromo-uridine), 3-methyl-uridine (m.sup.3U), 5-methoxy-uridine (mo.sup.5U), uridine 5-oxyacetic acid (cmo.sup.5U), uridine 5-oxyacetic acid methyl ester (mcmo.sup.5U), 5-carboxymethyl-uridine (cm.sup.5U), 1-carboxymethyl-pseudouridine, 5-carboxyhydroxymethyl-uridine (chm.sup.5U), 5-carboxyhydroxymethyl-uridine methyl ester (mchm.sup.5U), 5-methoxycarbonylmethyl-uridine (mcm.sup.5U), 5-methoxycarbonylmethyl-2-thio-uridine (mcm.sup.5s.sup.2U), 5-aminomethyl-2-thio-uridine (nm.sup.5s.sup.2U), 5-methylaminomethyl-uridine (mnm.sup.5U), 5-methylaminomethyl-2-thio-uridine (mnm.sup.5s.sup.2U), 5-methylaminomethyl-2-seleno-uridine (mnm.sup.5se.sup.2U), 5-carbamoylmethyl-uridine (ncm.sup.5U), 5-carboxymethylaminomethyl-uridine (cmnm.sup.5U), 5-carboxymethylaminomethyl-2-thio-uridine (cmnm.sup.5s.sup.2U), 5-propynyl-uridine, 1-propynyl-pseudouridine, 5-aurinomethyl-uridine (τ m.sup.5U), 1-aurinomethyl-pseudouridine, 5-aurinomethyl-2-thio-uridine (τ m.sup.5s.sup.2U), 1-aurinomethyl-4-thio-pseudouridine, 5-methyl-uridine (m.sup.5U, i.e., having the nucleobase deoxythymine), 1-methyl-pseudouridine (m.sup.1 ψ), 5-methyl-2-thio-uridine (m.sup.5s.sup.2U), 1-methyl-4-thio-pseudouridine (m.sup.1s.sup.4 ψ), 4-thio-1-methyl-pseudouridine, 3-methyl-pseudouridine (m.sup.3 ψ), 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydrouridine (D), dihydropseudouridine, 5,6-dihydrouridine, 5-methyldihydrouridine (m.sup.5D), 2-thio-dihydrouridine, 2-thio-dihydropseudouridine, 2-methoxy-uridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, 4-methoxy-2-thio-pseudouridine, N1-methyl-pseudouridine, 3-(3-amino-3-carboxypropyl)uridine (acp.sup.3U), 1-methyl-3-(3-amino-3-carboxypropyl)pseudouridine (acp.sup.3 ψ), 5-(isopentenylaminomethyl)uridine (inm.sup.5U), 5-(isopentenylaminomethyl)-2-thio-uridine (inm.sup.5s.sup.2U), α -thio-uridine, 2'-O-methyl-uridine (Um), 5,2'-O-dimethyl-uridine (m.sup.5Um), 2'-O-methyl-pseudouridine (ψ m), 2-thio-2'-O-methyl-uridine (s.sup.2Um), 5-methoxycarbonylmethyl-2'-O-methyl-uridine (mcm.sup.5Um), 5-carbamoylmethyl-2'-O-methyl-uridine (ncm.sup.5Um), 5-carboxymethylaminomethyl-2'-O-methyl-uridine (cmnm.sup.5Um), 3,2'-O-dimethyl-uridine (m.sup.3Um), and 5-(isopentenylaminomethyl)-2'-O-methyl-uridine

(imm.sup.5Um), 1-thio-uridine, deoxythymidine, 2'-F-ara-uridine, 2'-OH-ara-uridine, 5-(2-carbomethoxyvinyl) uridine, and 5-[3-(1-E-propenylamino)]uridine.

(207) In some embodiments, the modified nucleobase is a modified cytosine. Exemplary nucleobases and nucleosides having a modified cytosine include 5-aza-cytidine, 6-aza-cytidine, pseudoisocytidine, 3-methyl-cytidine (m.sup.3C), N4-acetyl-cytidine (ac.sup.4C), 5-formylcytidine (f.sup.5C), N4-methyl-cytidine (m.sup.4C), 5-methyl-cytidine (m.sup.5C), 5-halo-cytidine (e.g., 5-iodo-cytidine), 5-hydroxymethyl-cytidine (hm.sup.5C), 1-methyl-pseudoisocytidine, pyrrolo-cytidine, pyrrolo-pseudoisocytidine, 2-thio-cytidine (s.sup.2C), 2-thio-5-methyl-cytidine, 4-thio-pseudoisocytidine, 4-thio-1-methyl-pseudoisocytidine, 4-thio-1-methyl-1-deaza-pseudoisocytidine, 1-methyl-1-deaza-pseudoisocytidine, zebularine, 5-aza-zebularine, 5-methyl-zebularine, 5-aza-2-thio-zebularine, 2-thio-zebularine, 2-methoxy-cytidine, 2-methoxy-5-methyl-cytidine, 4-methoxy-pseudoisocytidine, 4-methoxy-1-methyl-pseudoisocytidine, lysidine (k.sub.2C), α -thio-cytidine, 2'-O-methyl-cytidine (Cm), 5,2'-O-dimethylcytidine (m.sup.5Cm), N4-acetyl-2'-O-methyl-cytidine (ac.sup.4Cm), N4,2'-O-dimethylcytidine (m.sup.4Cm), 5-formyl-2'-O-methyl-cytidine (f.sup.5Cm), N4,N4,2'-O-trimethyl-cytidine (m.sup.4.sub.2Cm), 1-thio-cytidine, 2'-F-ara-cytidine, 2'-F-cytidine, and 2'-OH-ara-cytidine.

(208) In some embodiments, the modified nucleobase is a modified adenine. Exemplary nucleobases and nucleosides having a modified adenine include 2-amino-purine, 2, 6-diaminopurine, 2-amino-6-halo-purine (e.g., 2-amino-6-chloro-purine), 6-halo-purine (e.g., 6-chloro-purine), 2-amino-6-methyl-purine, 8-azido-adenosine, 7-deaza-adenine, 7-deaza-8-aza-adenine, 7-deaza-2-amino-purine, 7-deaza-8-aza-2-amino-purine, 7-deaza-2,6-diaminopurine, 7-deaza-8-aza-2,6-diaminopurine, 1-methyl-adenosine (m.sup.1A), 2-methyl-adenine (m.sup.2A), N6-methyl-adenosine (m.sup.6A), 2-methylthio-N6-methyl-adenosine (ms.sup.2m.sup.6A), N6-isopentenyl-adenosine (i.sup.6A), 2-methylthio-N6-isopentenyl-adenosine (ms.sup.2i.sup.6A), N6-(cis-hydroxyisopentenyl)adenosine (io.sup.6A), 2-methylthio-N6-(cis-hydroxyisopentenyl)adenosine (ms.sup.2io.sup.6A), N6-glycinylicarbamoyl-adenosine (g.sup.6A), N6-threonylicarbamoyl-adenosine (t.sup.6A), N6-methyl-N6-threonylicarbamoyl-adenosine (m.sup.6t.sup.6A), 2-methylthio-N6-threonylicarbamoyl-adenosine (ms.sup.2g.sup.6A), N6,N6-dimethyl-adenosine (m.sup.6.sub.2A), N6-hydroxynorvalylcarbamoyl-adenosine (hn.sup.6A), 2-methylthio-N6-hydroxynorvalylcarbamoyl-adenosine (ms.sup.2hn.sup.6A), N6-acetyl-adenosine (ac.sup.6A), 7-methyl-adenine, 2-methylthio-adenine, 2-methoxy-adenine, α -thio-adenosine, 2'-O-methyl-adenosine (Am), N6,2'-O-dimethyl-adenosine (m.sup.6Am), N6,N6,2'-O-trimethyl-adenosine (m.sup.6.sub.2Am), 1,2'-O-dimethyl-adenosine (m.sup.1Am), 2'-O-ribosyladenosine (phosphate) (Ar(p)), 2-amino-N6-methyl-purine, 1-thio-adenosine, 8-azido-adenosine, 2'-F-ara-adenosine, 2'-F-adenosine, 2'-OH-ara-adenosine, and N6-(19-amino-pentaoxonadecyl)-adenosine.

(209) In some embodiments, the modified nucleobase is a modified guanine. Exemplary nucleobases and nucleosides having a modified guanine include inosine (I), 1-methyl-inosine (m.sup.1I), wyosine (imG), methylwyosine (mimG), 4-demethyl-wyosine (imG-14), isowyosine (imG2), wybutosine (yW), peroxywybutosine (o.sub.2yW), hydroxywybutosine (OhyW), undermodified hydroxywybutosine (OhyW*), 7-deaza-guanosine, queuosine (Q), epoxyqueuosine (oQ), galactosyl-queuosine (galQ), mannosyl-queuosine (manQ), 7-cyano-7-deaza-guanosine (preQ.sub.0), 7-aminomethyl-7-deaza-guanosine (preQ.sub.1), archaeosine (G.sup.+), 7-deaza-8-aza-guanosine, 6-thio-guanosine, 6-thio-7-deaza-guanosine, 6-thio-7-deaza-8-aza-guanosine, 7-methyl-guanosine (m.sup.7G), 6-thio-7-methyl-guanosine, 7-methyl-inosine, 6-methoxy-guanosine, 1-methyl-guanosine (m.sup.1G), N2-methyl-guanosine (m.sup.2G), N2,N2-dimethyl-guanosine (m.sup.2.sub.2G), N2,7-dimethyl-guanosine (m.sup.2,7G), N2, N2,7-dimethyl-guanosine (m.sup.2,2,7G), 8-oxo-guanosine, 7-methyl-8-oxo-guanosine, 1-methyl-6-thio-guanosine, N2-methyl-6-thio-guanosine, N2,N2-dimethyl-6-thio-guanosine, α -thio-guanosine, 2'-O-methyl-guanosine (Gm), N2-methyl-2'-O-methyl-guanosine (m.sup.2Gm), N2,N2-dimethyl-2'-

O-methyl-guanosine (m.sup.2.sub.2Gm), 1-methyl-2'-O-methyl-guanosine (mGm), N2,7-dimethyl-2'-O-methyl-guanosine (m.sup.2,7Gm), 2'-O-methyl-inosine (Im), 1,2'-O-dimethyl-inosine (m.sup.1Im), 2'-O-ribosylguanosine (phosphate) (Gr(p)), 1-thio-guanosine, O6-methyl-guanosine, 2'-F-ara-guanosine, and 2'-F-guanosine.

(210) N-Linked Glycosylation Site Mutants

(211) N-linked glycans of viral proteins play important roles in modulating the immune response. Glycans can be important for maintaining the appropriate antigenic conformations, shielding potential neutralization epitopes, and may alter the proteolytic susceptibility of proteins. Some viruses have putative N-linked glycosylation sites. Deletion or modification of an N-linked glycosylation site may enhance the immune response. Thus, the present disclosure provides, in some embodiments, RNA (e.g., mRNA) vaccines comprising nucleic acids (e.g., mRNA) encoding antigenic polypeptides that comprise a deletion or modification at one or more N-linked glycosylation sites.

(212) In Vitro Transcription of RNA (e.g., mRNA)

(213) Respiratory virus vaccines of the present disclosure comprise at least one RNA polynucleotide, such as a mRNA (e.g., modified mRNA). mRNA, for example, is transcribed in vitro from template DNA, referred to as an "in vitro transcription template." In some embodiments, an in vitro transcription template encodes a 5' untranslated (UTR) region, contains an open reading frame, and encodes a 3' UTR and a polyA tail. The particular nucleic acid sequence composition and length of an in vitro transcription template will depend on the mRNA encoded by the template.

(214) A "5' untranslated region" (5'UTR) refers to a region of an mRNA that is directly upstream (i.e., 5') from the start codon (i.e., the first codon of an mRNA transcript translated by a ribosome) that does not encode a polypeptide.

(215) A "3' untranslated region" (3'UTR) refers to a region of an mRNA that is directly downstream (i.e., 3') from the stop codon (i.e., the codon of an mRNA transcript that signals a termination of translation) that does not encode a polypeptide.

(216) An "open reading frame" is a continuous stretch of DNA beginning with a start codon (e.g., methionine (ATG)), and ending with a stop codon (e.g., TAA, TAG or TGA) and encodes a polypeptide.

(217) A "polyA tail" is a region of mRNA that is downstream, e.g., directly downstream (i.e., 3'), from the 3' UTR that contains multiple, consecutive adenosine monophosphates. A polyA tail may contain 10 to 300 adenosine monophosphates. For example, a polyA tail may contain 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290 or 300 adenosine monophosphates. In some embodiments, a polyA tail contains 50 to 250 adenosine monophosphates. In a relevant biological setting (e.g., in cells, in vivo) the poly(A) tail functions to protect mRNA from enzymatic degradation, e.g., in the cytoplasm, and aids in transcription termination, export of the mRNA from the nucleus and translation.

(218) In some embodiments, a polynucleotide includes 200 to 3,000 nucleotides. For example, a polynucleotide may include 200 to 500, 200 to 1000, 200 to 1500, 200 to 3000, 500 to 1000, 500 to 1500, 500 to 2000, 500 to 3000, 1000 to 1500, 1000 to 2000, 1000 to 3000, 1500 to 3000, or 2000 to 3000 nucleotides.

(219) Flagellin Adjuvants

(220) Flagellin is an approximately 500 amino acid monomeric protein that polymerizes to form the flagella associated with bacterial motion. Flagellin is expressed by a variety of flagellated bacteria (*Salmonella typhimurium* for example) as well as non-flagellated bacteria (such as *Escherichia coli*). Sensing of flagellin by cells of the innate immune system (dendritic cells, macrophages, etc.) is mediated by the Toll-like receptor 5 (TLR5) as well as by Nod-like receptors (NLRs) Ipaf and Naip5. TLRs and NLRs have been identified as playing a role in the activation of innate immune response and adaptive immune response. As such, flagellin provides an adjuvant

effect in a vaccine.

(221) The nucleotide and amino acid sequences encoding known flagellin polypeptides are publicly available in the NCBI GenBank database. The flagellin sequences from *S. typhimurium*, *H. pylori*, *V. cholera*, *S. marcesens*, *S. flexneri*, *T. pallidum*, *L. pneumophila*, *B. burgdorferi*, *C. difficile*, *R. meliloti*, *A. tumefaciens*, *R. lupini*, *B. clarridgeiae*, *P. mirabilis*, *B. subtilis*, *L. monocytogenes*, *P. aeruginosa*, and *E. coli*, among others are known.

(222) A flagellin polypeptide, as used herein, refers to a full length flagellin protein, immunogenic fragments thereof, and peptides having at least 50% sequence identify to a flagellin protein or immunogenic fragments thereof. Exemplary flagellin proteins include flagellin from *Salmonella typhi* (UniPro Entry number: Q56086), *Salmonella typhimurium* (A0A0C9DG09), *Salmonella enteritidis* (A0A0C9BAB7), and *Salmonella choleraesuis* (Q6V2X8), and SEQ ID NO: 54-56 (Table 17). In some embodiments, the flagellin polypeptide has at least 60%, 70%, 75%, 80%, 90%, 95%, 97%, 98%, or 99% sequence identify to a flagellin protein or immunogenic fragments thereof.

(223) In some embodiments, the flagellin polypeptide is an immunogenic fragment. An immunogenic fragment is a portion of a flagellin protein that provokes an immune response. In some embodiments, the immune response is a TLR5 immune response. An example of an immunogenic fragment is a flagellin protein in which all or a portion of a hinge region has been deleted or replaced with other amino acids. For example, an antigenic polypeptide may be inserted in the hinge region. Hinge regions are the hypervariable regions of a flagellin. Hinge regions of a flagellin are also referred to as “D3 domain or region,” “propeller domain or region,” “hypervariable domain or region” and “variable domain or region.” “At least a portion of a hinge region,” as used herein, refers to any part of the hinge region of the flagellin, or the entirety of the hinge region. In other embodiments an immunogenic fragment of flagellin is a 20, 25, 30, 35, or 40 amino acid C-terminal fragment of flagellin.

(224) The flagellin monomer is formed by domains DO through D3. DO and D1, which form the stem, are composed of tandem long alpha helices and are highly conserved among different bacteria. The D1 domain includes several stretches of amino acids that are useful for TLR5 activation. The entire D1 domain or one or more of the active regions within the domain are immunogenic fragments of flagellin. Examples of immunogenic regions within the D1 domain include residues 88-114 and residues 411-431 (in *Salmonella typhimurium* FliC flagellin. Within the 13 amino acids in the 88-100 region, at least 6 substitutions are permitted between *Salmonella* flagellin and other flagellins that still preserve TLR5 activation. Thus, immunogenic fragments of flagellin include flagellin like sequences that activate TLR5 and contain a 13 amino acid motif that is 53% or more identical to the *Salmonella* sequence in 88-100 of FliC (LQRVRELAVQSAN; SEQ ID NO: 84).

(225) In some embodiments, the RNA (e.g., mRNA) vaccine includes an RNA that encodes a fusion protein of flagellin and one or more antigenic polypeptides. A “fusion protein” as used herein, refers to a linking of two components of the construct. In some embodiments, a carboxy-terminus of the antigenic polypeptide is fused or linked to an amino terminus of the flagellin polypeptide. In other embodiments, an amino-terminus of the antigenic polypeptide is fused or linked to a carboxy-terminus of the flagellin polypeptide. The fusion protein may include, for example, one, two, three, four, five, six or more flagellin polypeptides linked to one, two, three, four, five, six or more antigenic polypeptides. When two or more flagellin polypeptides and/or two or more antigenic polypeptides are linked such a construct may be referred to as a “multimer.”

(226) Each of the components of a fusion protein may be directly linked to one another or they may be connected through a linker. For instance, the linker may be an amino acid linker. The amino acid linker encoded for by the RNA (e.g., mRNA) vaccine to link the components of the fusion protein may include, for instance, at least one member selected from the group consisting of a lysine residue, a glutamic acid residue, a serine residue and an arginine residue. In some embodiments the

linker is 1-30, 1-25, 1-25, 5-10, 5, 15, or 5-20 amino acids in length.

(227) In other embodiments the RNA (e.g., mRNA) vaccine includes at least two separate RNA polynucleotides, one encoding one or more antigenic polypeptides and the other encoding the flagellin polypeptide. The at least two RNA polynucleotides may be co-formulated in a carrier such as a lipid nanoparticle.

(228) Broad Spectrum RNA (e.g., mRNA) Vaccines

(229) There may be situations where persons are at risk for infection with more than one strain of hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1). RNA (e.g., mRNA) therapeutic vaccines are particularly amenable to combination vaccination approaches due to a number of factors including, but not limited to, speed of manufacture, ability to rapidly tailor vaccines to accommodate perceived geographical threat, and the like. Moreover, because the vaccines utilize the human body to produce the antigenic protein, the vaccines are amenable to the production of larger, more complex antigenic proteins, allowing for proper folding, surface expression, antigen presentation, etc. in the human subject. To protect against more than one strain of hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1), a combination vaccine can be administered that includes RNA (e.g., mRNA) encoding at least one antigenic polypeptide protein (or antigenic portion thereof) of a first respiratory virus and further includes RNA encoding at least one antigenic polypeptide protein (or antigenic portion thereof) of a second respiratory virus. RNA (e.g., mRNA) can be co-formulated, for example, in a single lipid nanoparticle (LNP) or can be formulated in separate LNPs for co-administration.

(230) Methods of Treatment

(231) Provided herein are compositions (e.g., pharmaceutical compositions), methods, kits and reagents for prevention and/or treatment of respiratory diseases/infections in humans and other mammals. Respiratory virus RNA (e.g. mRNA) vaccines can be used as therapeutic or prophylactic agents, alone or in combination with other vaccine(s). They may be used in medicine to prevent and/or treat respiratory disease/infection. In exemplary aspects, the RNA (e.g., mRNA) vaccines of the present disclosure are used to provide prophylactic protection from hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1). Prophylactic protection from hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) can be achieved following administration of a RNA (e.g., mRNA) vaccine of the present disclosure. Respiratory virus RNA (e.g., mRNA) vaccines of the present disclosure may be used to treat or prevent viral “co-infections” containing two or more respiratory infections. Vaccines can be administered once, twice, three times, four times or more, but it is likely sufficient to administer the vaccine once (optionally followed by a single booster). It is possible, although less desirable, to administer the vaccine to an infected individual to achieve a therapeutic response. Dosing may need to be adjusted accordingly.

(232) A method of eliciting an immune response in a subject against hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) is provided in aspects of the present disclosure. The method involves administering to the subject a respiratory virus RNA (e.g., mRNA) vaccine comprising at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) antigenic polypeptide thereof, thereby inducing in the subject an immune response specific to hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) antigenic polypeptide or an immunogenic fragment thereof, wherein anti-antigenic polypeptide antibody titer in the subject is

increased following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1). An “anti-antigenic polypeptide antibody” is a serum antibody that binds specifically to the antigenic polypeptide.

(233) In some embodiments, a RNA (e.g., mRNA) vaccine (e.g., a hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1 RNA vaccine) capable of eliciting an immune response is administered intramuscularly via a composition including a compound according to Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIe) (e.g., Compound 3, 18, 20, 25, 26, 29, 30, 60, 108-112, or 122).

(234) A prophylactically effective dose is a therapeutically effective dose that prevents infection with the virus at a clinically acceptable level. In some embodiments the therapeutically effective dose is a dose listed in a package insert for the vaccine. A traditional vaccine, as used herein, refers to a vaccine other than the RNA (e.g., mRNA) vaccines of the present disclosure. For instance, a traditional vaccine includes but is not limited to live/attenuated microorganism vaccines, killed/inactivated microorganism vaccines, subunit vaccines, protein antigen vaccines, DNA vaccines, VLP vaccines, etc. In exemplary embodiments, a traditional vaccine is a vaccine that has achieved regulatory approval and/or is registered by a national drug regulatory body, for example the Food and Drug Administration (FDA) in the United States or the European Medicines Agency (EMA).

(235) In some embodiments the anti-antigenic polypeptide antibody titer in the subject is increased 1 log to 10 log following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1).

(236) In some embodiments the anti-antigenic polypeptide antibody titer in the subject is increased 1 log, 2 log, 3 log, 5 log or 10 log following vaccination relative to anti-antigenic polypeptide antibody titer in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1).

(237) A method of eliciting an immune response in a subject against hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) is provided in other aspects of the disclosure. The method involves administering to the subject a respiratory virus RNA (e.g., mRNA) vaccine comprising at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) antigenic polypeptide or an immunogenic fragment thereof, thereby inducing in the subject an immune response specific to hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) antigenic polypeptide or an immunogenic fragment thereof, wherein the immune response in the subject is equivalent to an immune response in a subject vaccinated with a traditional vaccine against the hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) at 2 times to 100 times the dosage level relative to the RNA (e.g., mRNA) vaccine.

(238) In some embodiments, the immune response in the subject is equivalent to an immune response in a subject vaccinated with a traditional vaccine at 2, 3, 4, 5, 10, 50, 100 times the dosage level relative to the hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) RNA

(e.g., mRNA) vaccine.

(239) In some embodiments the immune response in the subject is equivalent to an immune response in a subject vaccinated with a traditional vaccine at 10-100 times, or 100-1000 times, the dosage level relative to the hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) RNA (e.g., mRNA) vaccine.

(240) In some embodiments the immune response is assessed by determining [protein] antibody titer in the subject.

(241) Some aspects of the present disclosure provide a method of eliciting an immune response in a subject against a In some embodiments the immune response in the subject is equivalent to an immune response in a subject vaccinated with a traditional vaccine at 2, 3, 4, 5, 10, 50, 100 times the dosage level relative to the hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) RNA (e.g., mRNA) vaccine by administering to the subject a respiratory virus RNA (e.g., mRNA) vaccine comprising at least one RNA (e.g., mRNA) polynucleotide having an open reading frame encoding at least one hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) antigenic polypeptide, thereby inducing in the subject an immune response specific to the antigenic polypeptide or an immunogenic fragment thereof, wherein the immune response in the subject is induced 2 days to 10 weeks earlier relative to an immune response induced in a subject vaccinated with a prophylactically effective dose of a traditional vaccine against the hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1). In some embodiments, the immune response in the subject is induced in a subject vaccinated with a prophylactically effective dose of a traditional vaccine at 2 times to 100 times the dosage level relative to the RNA (e.g., mRNA) vaccine.

(242) In some embodiments, the immune response in the subject is induced 2 days earlier, or 3 days earlier, relative to an immune response induced in a subject vaccinated with a prophylactically effective dose of a traditional vaccine.

(243) In some embodiments the immune response in the subject is induced 1 week, 2 weeks, 3 weeks, 5 weeks, or 10 weeks earlier relative to an immune response induced in a subject vaccinated with a prophylactically effective dose of a traditional vaccine.

(244) Also provided herein is a method of eliciting an immune response in a subject against hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) by administering to the subject a respiratory virus RNA (e.g., mRNA) vaccine having an open reading frame encoding a first antigenic polypeptide, wherein the RNA polynucleotide does not include a stabilization element, and wherein an adjuvant is not co-formulated or co-administered with the vaccine.

(245) Therapeutic and Prophylactic Compositions

(246) Provided herein are compositions (e.g., pharmaceutical compositions), methods, kits and reagents for prevention, treatment or diagnosis of hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1) in humans and other mammals, for example. Respiratory virus RNA (e.g. mRNA) vaccines can be used as therapeutic or prophylactic agents. They may be used in medicine to prevent and/or treat infectious disease. In some embodiments, the respiratory RNA (e.g., mRNA) vaccines of the present disclosure are used for the priming of immune effector cells, for example, to activate peripheral blood mononuclear cells (PBMCs) ex vivo, which are then infused (re-infused) into a subject.

(247) In some embodiments, respiratory virus vaccine containing RNA (e.g., mRNA) polynucleotides as described herein can be administered to a subject (e.g., a mammalian subject,

such as a human subject), and the RNA (e.g., mRNA) polynucleotides are translated in vivo to produce an antigenic polypeptide.

(248) The respiratory virus RNA (e.g., mRNA) vaccines may be induced for translation of a polypeptide (e.g., antigen or immunogen) in a cell, tissue or organism. In some embodiments, such translation occurs in vivo, although such translation may occur ex vivo, in culture or in vitro. In some embodiments, the cell, tissue or organism is contacted with an effective amount of a composition containing a respiratory virus RNA (e.g., mRNA) vaccine that contains a polynucleotide that has at least one a translatable region encoding an antigenic polypeptide.

(249) An “effective amount” of an respiratory virus RNA (e.g. mRNA) vaccine is provided based, at least in part, on the target tissue, target cell type, means of administration, physical characteristics of the polynucleotide (e.g., size, and extent of modified nucleosides) and other components of the vaccine, and other determinants. In general, an effective amount of the respiratory virus RNA (e.g., mRNA) vaccine composition provides an induced or boosted immune response as a function of antigen production in the cell, preferably more efficient than a composition containing a corresponding unmodified polynucleotide encoding the same antigen or a peptide antigen. Increased antigen production may be demonstrated by increased cell transfection (the percentage of cells transfected with the RNA, e.g., mRNA, vaccine), increased protein translation from the polynucleotide, decreased nucleic acid degradation (as demonstrated, for example, by increased duration of protein translation from a modified polynucleotide), or altered antigen specific immune response of the host cell.

(250) In some embodiments, RNA (e.g. mRNA) vaccines (including polynucleotides their encoded polypeptides) in accordance with the present disclosure may be used for treatment of hMPV, PIV3, RSV, MeV and/or BetaCoV (including MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH and/or HCoV—HKU1).

(251) Respiratory RNA (e.g. mRNA) vaccines may be administered prophylactically or therapeutically as part of an active immunization scheme to healthy individuals or early in infection during the incubation phase or during active infection after onset of symptoms. In some embodiments, the amount of RNA (e.g., mRNA) vaccine of the present disclosure provided to a cell, a tissue or a subject may be an amount effective for immune prophylaxis.

(252) Respiratory virus RNA (e.g. mRNA) vaccines may be administered with other prophylactic or therapeutic compounds. As a non-limiting example, a prophylactic or therapeutic compound may be an adjuvant or a booster. As used herein, when referring to a prophylactic composition, such as a vaccine, the term “booster” refers to an extra administration of the prophylactic (vaccine) composition. A booster (or booster vaccine) may be given after an earlier administration of the prophylactic composition. The time of administration between the initial administration of the prophylactic composition and the booster may be, but is not limited to, 1 minute, 2 minutes, 3 minutes, 4 minutes, 5 minutes, 6 minutes, 7 minutes, 8 minutes, 9 minutes, 10 minutes, 15 minutes, 20 minutes, 35 minutes, 40 minutes, 45 minutes, 50 minutes, 55 minutes, 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 1 day, 36 hours, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 10 days, 2 weeks, 3 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 1 year, 18 months, 2 years, 3 years, 4 years, 5 years, 6 years, 7 years, 8 years, 9 years, 10 years, 11 years, 12 years, 13 years, 14 years, 15 years, 16 years, 17 years, 18 years, 19 years, 20 years, 25 years, 30 years, 35 years, 40 years, 45 years, 50 years, 55 years, 60 years, 65 years, 70 years, 75 years, 80 years, 85 years, 90 years, 95 years or more than 99 years. In some embodiments, the time of administration between the initial administration of the prophylactic composition and the booster may be, but is not limited to, 1 week, 2 weeks, 3 weeks, 1 month, 2 months, 3 months, 6 months or 1 year.

(253) In some embodiments, respiratory virus RNA (e.g. mRNA) vaccines may be administered

intramuscularly or intradermally, similarly to the administration of inactivated vaccines known in the art.

(254) Respiratory virus RNA (e.g. mRNA) vaccines may be utilized in various settings depending on the prevalence of the infection or the degree or level of unmet medical need. As a non-limiting example, the RNA (e.g., mRNA) vaccines may be utilized to treat and/or prevent a variety of respiratory infections. RNA (e.g., mRNA) vaccines have superior properties in that they produce much larger antibody titers and produce responses early than commercially available anti-viral agents/compositions.

(255) Provided herein are pharmaceutical compositions including respiratory virus RNA (e.g. mRNA) vaccines and RNA (e.g. mRNA) vaccine compositions and/or complexes optionally in combination with one or more pharmaceutically acceptable excipients.

(256) Respiratory virus RNA (e.g. mRNA) vaccines may be formulated or administered alone or in conjunction with one or more other components. For instance, hMPV/PIV3/RSV RNA (e.g., mRNA) vaccines (vaccine compositions) may comprise other components including, but not limited to, adjuvants.

(257) In some embodiments, respiratory virus (e.g. mRNA) vaccines do not include an adjuvant (they are adjuvant free).

(258) Respiratory virus RNA (e.g. mRNA) vaccines may be formulated or administered in combination with one or more pharmaceutically-acceptable excipients. In some embodiments, vaccine compositions comprise at least one additional active substances, such as, for example, a therapeutically-active substance, a prophylactically-active substance, or a combination of both. Vaccine compositions may be sterile, pyrogen-free or both sterile and pyrogen-free. General considerations in the formulation and/or manufacture of pharmaceutical agents, such as vaccine compositions, may be found, for example, in Remington: The Science and Practice of Pharmacy 21st ed., Lippincott Williams & Wilkins, 2005 (incorporated herein by reference in its entirety).

(259) In some embodiments, respiratory virus RNA (e.g. mRNA) vaccines are administered to humans, human patients or subjects. For the purposes of the present disclosure, the phrase “active ingredient” generally refers to the RNA (e.g., mRNA) vaccines or the polynucleotides contained therein, for example, RNA polynucleotides (e.g., mRNA polynucleotides) encoding antigenic polypeptides.

(260) Formulations of the respiratory virus vaccine compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of bringing the active ingredient (e.g., mRNA polynucleotide) into association with an excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, dividing, shaping and/or packaging the product into a desired single- or multi-dose unit.

(261) Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the disclosure will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, the composition may comprise between 0.1% and 100%, e.g., between 0.5 and 50%, between 1-30%, between 5-80%, at least 80% (w/w) active ingredient.

(262) Respiratory virus RNA (e.g. mRNA) vaccines can be formulated using one or more excipients to: (1) increase stability; (2) increase cell transfection; (3) permit the sustained or delayed release (e.g., from a depot formulation); (4) alter the biodistribution (e.g., target to specific tissues or cell types); (5) increase the translation of encoded protein in vivo; and/or (6) alter the release profile of encoded protein (antigen) in vivo. In addition to traditional excipients such as any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, excipients can include, without limitation, lipidoids, liposomes, lipid nanoparticles, polymers, lipoplexes,

core-shell nanoparticles, peptides, proteins, cells transfected with respiratory virus RNA (e.g. mRNA) vaccines (e.g., for transplantation into a subject), hyaluronidase, nanoparticle mimics and combinations thereof.

(263) Stabilizing Elements

(264) Naturally-occurring eukaryotic mRNA molecules have been found to contain stabilizing elements, including, but not limited to untranslated regions (UTR) at their 5'-end (5'UTR) and/or at their 3'-end (3'UTR), in addition to other structural features, such as a 5'-cap structure or a 3'-poly(A) tail. Both the 5'UTR and the 3'UTR are typically transcribed from the genomic DNA and are elements of the premature mRNA. Characteristic structural features of mature mRNA, such as the 5'-cap and the 3'-poly(A) tail are usually added to the transcribed (premature) mRNA during mRNA processing. The 3'-poly(A) tail is typically a stretch of adenine nucleotides added to the 3'-end of the transcribed mRNA. It can comprise up to about 400 adenine nucleotides. In some embodiments the length of the 3'-poly(A) tail may be an essential element with respect to the stability of the individual mRNA.

(265) In some embodiments the RNA (e.g., mRNA) vaccine may include one or more stabilizing elements. Stabilizing elements may include for instance a histone stem-loop. A stem-loop binding protein (SLBP), a 32 kDa protein has been identified. It is associated with the histone stem-loop at the 3'-end of the histone messages in both the nucleus and the cytoplasm. Its expression level is regulated by the cell cycle; it peaks during the S-phase, when histone mRNA levels are also elevated. The protein has been shown to be essential for efficient 3'-end processing of histone pre-mRNA by the U7 snRNP. SLBP continues to be associated with the stem-loop after processing, and then stimulates the translation of mature histone mRNAs into histone proteins in the cytoplasm. The RNA binding domain of SLBP is conserved through metazoa and protozoa; its binding to the histone stem-loop depends on the structure of the loop. The minimum binding site includes at least three nucleotides 5' and two nucleotides 3' relative to the stem-loop.

(266) In some embodiments, the RNA (e.g., mRNA) vaccines include a coding region, at least one histone stem-loop, and optionally, a poly(A) sequence or polyadenylation signal. The poly(A) sequence or polyadenylation signal generally should enhance the expression level of the encoded protein. The encoded protein, in some embodiments, is not a histone protein, a reporter protein (e.g. Luciferase, GFP, EGFP, β -Galactosidase, EGFP), or a marker or selection protein (e.g. alpha-Globin, Galactokinase and Xanthine:guanine phosphoribosyl transferase (GPT)).

(267) In some embodiments, the combination of a poly(A) sequence or polyadenylation signal and at least one histone stem-loop, even though both represent alternative mechanisms in nature, acts synergistically to increase the protein expression beyond the level observed with either of the individual elements. It has been found that the synergistic effect of the combination of poly(A) and at least one histone stem-loop does not depend on the order of the elements or the length of the poly(A) sequence.

(268) In some embodiments, the RNA (e.g., mRNA) vaccine does not comprise a histone downstream element (HDE). "Histone downstream element" (HDE) includes a purine-rich polynucleotide stretch of approximately 15 to 20 nucleotides 3' of naturally occurring stem-loops, representing the binding site for the U7 snRNA, which is involved in processing of histone pre-mRNA into mature histone mRNA. Ideally, the inventive nucleic acid does not include an intron.

(269) In some embodiments, the RNA (e.g., mRNA) vaccine may or may not contain an enhancer and/or promoter sequence, which may be modified or unmodified or which may be activated or inactivated. In some embodiments, the histone stem-loop is generally derived from histone genes, and includes an intramolecular base pairing of two neighbored partially or entirely reverse complementary sequences separated by a spacer, including (e.g., consisting of) a short sequence, which forms the loop of the structure. The unpaired loop region is typically unable to base pair with either of the stem loop elements. It occurs more often in RNA, as is a key component of many RNA secondary structures, but may be present in single-stranded DNA as well. Stability of the

stem-loop structure generally depends on the length, number of mismatches or bulges, and base composition of the paired region. In some embodiments, wobble base pairing (non-Watson-Crick base pairing) may result. In some embodiments, the at least one histone stem-loop sequence comprises a length of 15 to 45 nucleotides.

(270) In other embodiments the RNA (e.g., mRNA) vaccine may have one or more AU-rich sequences removed. These sequences, sometimes referred to as AURES are destabilizing sequences found in the 3'UTR. The AURES may be removed from the RNA (e.g., mRNA) vaccines. Alternatively the AURES may remain in the RNA (e.g., mRNA) vaccine.

(271) Nanoparticle Formulations

(272) In some embodiments, respiratory virus RNA (e.g. mRNA) vaccines are formulated in a nanoparticle. In some embodiments, respiratory virus RNA (e.g. mRNA) vaccines are formulated in a lipid nanoparticle. In some embodiments, respiratory virus RNA (e.g. mRNA) vaccines are formulated in a lipid-polycation complex, referred to as a cationic lipid nanoparticle. As a non-limiting example, the polycation may include a cationic peptide or a polypeptide such as, but not limited to, polylysine, polyornithine and/or polyarginine. In some embodiments, respiratory virus RNA (e.g., mRNA) vaccines are formulated in a lipid nanoparticle that includes a non-cationic lipid such as, but not limited to, cholesterol or dioleoyl phosphatidylethanolamine (DOPE).

(273) A lipid nanoparticle formulation may be influenced by, but not limited to, the selection of the cationic lipid component, the degree of cationic lipid saturation, the nature of the PEGylation, ratio of all components and biophysical parameters such as size. In one example by Semple et al. (*Nature Biotech.* 2010 28:172-176), the lipid nanoparticle formulation is composed of 57.1% cationic lipid, 7.1% dipalmitoylphosphatidylcholine, 34.3% cholesterol, and 1.4% PEG-c-DMA. As another example, changing the composition of the cationic lipid can more effectively deliver siRNA to various antigen presenting cells (Basha et al. *Mol Ther.* 2011 19:2186-2200).

(274) In some embodiments, lipid nanoparticle formulations may comprise 35 to 45% cationic lipid, 40% to 50% cationic lipid, 50% to 60% cationic lipid and/or 55% to 65% cationic lipid. In some embodiments, the ratio of lipid to RNA (e.g., mRNA) in lipid nanoparticles may be 5:1 to 20:1, 10:1 to 25:1, 15:1 to 30:1 and/or at least 30:1.

(275) In some embodiments, the ratio of PEG in the lipid nanoparticle formulations may be increased or decreased and/or the carbon chain length of the PEG lipid may be modified from C14 to C18 to alter the pharmacokinetics and/or biodistribution of the lipid nanoparticle formulations. As a non-limiting example, lipid nanoparticle formulations may contain 0.5% to 3.0%, 1.0% to 3.5%, 1.5% to 4.0%, 2.0% to 4.5%, 2.5% to 5.0% and/or 3.0% to 6.0% of the lipid molar ratio of PEG-c-DOMG (R-3-[(ω-methoxy-poly(ethyleneglycol)2000)carbamoyl]-1,2-dimyristyloxypropyl-3-amine) (also referred to herein as PEG-DOMG) as compared to the cationic lipid, DSPC and cholesterol. In some embodiments, the PEG-c-DOMG may be replaced with a PEG lipid such as, but not limited to, PEG-DSG (1,2-Distearoyl-sn-glycerol, methoxypolyethylene glycol), PEG-DMG (1,2-Dimyristoyl-sn-glycerol) and/or PEG-DPG (1,2-Dipalmitoyl-sn-glycerol, methoxypolyethylene glycol). The cationic lipid may be selected from any lipid known in the art such as, but not limited to, DLin-MC3-DMA, DLin-DMA, C12-200 and DLin-KC2-DMA.

(276) In some embodiments, an respiratory virus RNA (e.g. mRNA) vaccine formulation is a nanoparticle that comprises at least one lipid. The lipid may be selected from, but is not limited to, DLin-DMA, DLin-K-DMA, 98N12-5, C12-200, DLin-MC3-DMA, DLin-KC2-DMA, DODMA, PLGA, PEG, PEG-DMG, PEGylated lipids and amino alcohol lipids. In some embodiments, the lipid may be a cationic lipid such as, but not limited to, DLin-DMA, DLin-D-DMA, DLin-MC3-DMA, DLin-KC2-DMA, DODMA and amino alcohol lipids. The amino alcohol cationic lipid may be the lipids described in and/or made by the methods described in U.S. Patent Publication No. US20130150625, herein incorporated by reference in its entirety. As a non-limiting example, the cationic lipid may be 2-amino-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[(9Z,2Z)-octadeca-9,12-dien-1-yloxy]methyl}propan-1-ol (Compound 1 in US20130150625); 2-amino-3-[(9Z)-

octadec-9-en-1-yloxy]-2-[[[(9Z)-octadeca-9-en-1-yloxy]methyl]propan-1-ol (Compound 2 in US20130150625); 2-amino-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[(octyloxy)methyl]propan-1-ol (Compound 3 in US20130150625); and 2-(dimethylamino)-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[[[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]methyl]propan-1-ol (Compound 4 in US20130150625); or any pharmaceutically acceptable salt or stereoisomer thereof.

(277) Lipid nanoparticle formulations typically comprise a lipid, in particular, an ionizable cationic lipid, for example, 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), or di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), and further comprise a neutral lipid, a sterol and a molecule capable of reducing particle aggregation, for example a PEG or PEG-modified lipid.

(278) In some embodiments, a lipid nanoparticle formulation consists essentially of (i) at least one lipid selected from the group consisting of 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319); (ii) a neutral lipid selected from DSPC, DPPC, POPC, DOPE and SM; (iii) a sterol, e.g., cholesterol; and (iv) a PEG-lipid, e.g., PEG-DMG or PEG-cDMA, in a molar ratio of 20-60% cationic lipid: 5-25% neutral lipid: 25-55% sterol; 0.5-15% PEG-lipid.

(279) In some embodiments, a lipid nanoparticle formulation includes 25% to 75% on a molar basis of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), e.g., 35 to 65%, 45 to 65%, 60%, 57.5%, 50% or 40% on a molar basis.

(280) In some embodiments, a lipid nanoparticle formulation includes 0.5% to 15% on a molar basis of the neutral lipid, e.g., 3 to 12%, 5 to 10% or 15%, 10%, or 7.5% on a molar basis. Examples of neutral lipids include, without limitation, DSPC, POPC, DPPC, DOPE and SM. In some embodiments, the formulation includes 5% to 50% on a molar basis of the sterol (e.g., 15 to 45%, 20 to 40%, 40%, 38.5%, 35%, or 31% on a molar basis. A non-limiting example of a sterol is cholesterol. In some embodiments, a lipid nanoparticle formulation includes 0.5% to 20% on a molar basis of the PEG or PEG-modified lipid (e.g., 0.5 to 10%, 0.5 to 5%, 1.5%, 0.5%, 1.5%, 3.5%, or 5% on a molar basis. In some embodiments, a PEG or PEG modified lipid comprises a PEG molecule of an average molecular weight of 2,000 Da. In some embodiments, a PEG or PEG modified lipid comprises a PEG molecule of an average molecular weight of less than 2,000, for example around 1,500 Da, around 1,000 Da, or around 500 Da. Non-limiting examples of PEG-modified lipids include PEG-distearoyl glycerol (PEG-DMG) (also referred herein as PEG-C14 or C14-PEG), PEG-cDMA (further discussed in Reyes et al. J. Controlled Release, 107, 276-287 (2005) the contents of which are herein incorporated by reference in their entirety).

(281) In some embodiments, lipid nanoparticle formulations include 25-75% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 0.5-15% of the neutral lipid, 5-50% of the sterol, and 0.5-20% of the PEG or PEG-modified lipid on a molar basis.

(282) In some embodiments, lipid nanoparticle formulations include 35-65% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 3-12% of the neutral lipid, 15-45% of the sterol, and 0.5-10% of the PEG or PEG-modified lipid on a molar basis.

(283) In some embodiments, lipid nanoparticle formulations include 45-65% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-

(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 5-10% of the neutral lipid, 25-40% of the sterol, and 0.5-10% of the PEG or PEG-modified lipid on a molar basis.

(284) In some embodiments, lipid nanoparticle formulations include 60% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 7.5% of the neutral lipid, 31% of the sterol, and 1.5% of the PEG or PEG-modified lipid on a molar basis.

(285) In some embodiments, lipid nanoparticle formulations include 50% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 10% of the neutral lipid, 38.5% of the sterol, and 1.5% of the PEG or PEG-modified lipid on a molar basis.

(286) In some embodiments, lipid nanoparticle formulations include 50% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 10% of the neutral lipid, 35% of the sterol, 4.5% or 5% of the PEG or PEG-modified lipid, and 0.5% of the targeting lipid on a molar basis.

(287) In some embodiments, lipid nanoparticle formulations include 40% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 15% of the neutral lipid, 40% of the sterol, and 5% of the PEG or PEG-modified lipid on a molar basis.

(288) In some embodiments, lipid nanoparticle formulations include 57.2% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 7.1% of the neutral lipid, 34.3% of the sterol, and 1.4% of the PEG or PEG-modified lipid on a molar basis.

(289) In some embodiments, lipid nanoparticle formulations include 57.5% of a cationic lipid selected from the PEG lipid is PEG-cDMA (PEG-cDMA is further discussed in Reyes et al. (J. Controlled Release, 107, 276-287 (2005), the contents of which are herein incorporated by reference in their entirety), 7.5% of the neutral lipid, 31.5% of the sterol, and 3.5% of the PEG or PEG-modified lipid on a molar basis.

(290) In some embodiments, lipid nanoparticle formulations consists essentially of a lipid mixture in molar ratios of 20-70% cationic lipid: 5-45% neutral lipid: 20-55% cholesterol: 0.5-15% PEG-modified lipid. In some embodiments, lipid nanoparticle formulations consists essentially of a lipid mixture in a molar ratio of 20-60% cationic lipid: 5-25% neutral lipid: 25-55% cholesterol: 0.5-15% PEG-modified lipid.

(291) In some embodiments, the molar lipid ratio is 50/10/38.5/1.5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG, PEG-DSG or PEG-DPG), 57.2/7.1134.3/1.4 (mol % cationic lipid/neutral lipid, e.g., DPPC/Chol/PEG-modified lipid, e.g., PEG-cDMA), 40/15/40/5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG), 50/10/35/4.5/0.5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DSG), 50/10/35/5 (cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG), 40/10/40/10 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG or PEG-cDMA), 35/15/40/10 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG or PEG-cDMA) or 52/13/30/5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG or PEG-cDMA).

(292) Non-limiting examples of lipid nanoparticle compositions and methods of making them are

described, for example, in Semple et al. (2010) *Nat Biotechnol.* 28:172-176; Jayarama et al. (2012), *Angew. Chem. Int. Ed.*, 51: 8529-8533; and Maier et al. (2013) *Molecular Therapy* 21, 1570-1578 (the contents of each of which are incorporated herein by reference in their entirety).

(293) In some embodiments, lipid nanoparticle formulations may comprise a cationic lipid, a PEG lipid and a structural lipid and optionally comprise a non-cationic lipid. As a non-limiting example, a lipid nanoparticle may comprise 40-60% of cationic lipid, 5-15% of a non-cationic lipid, 1-2% of a PEG lipid and 30-50% of a structural lipid. As another non-limiting example, the lipid nanoparticle may comprise 50% cationic lipid, 10% non-cationic lipid, 1.5% PEG lipid and 38.5% structural lipid. As yet another non-limiting example, a lipid nanoparticle may comprise 55% cationic lipid, 10% non-cationic lipid, 2.5% PEG lipid and 32.5% structural lipid. In some embodiments, the cationic lipid may be any cationic lipid described herein such as, but not limited to, DLin-KC2-DMA, DLin-MC3-DMA and L319.

(294) In some embodiments, the lipid nanoparticle formulations described herein may be 4 component lipid nanoparticles. The lipid nanoparticle may comprise a cationic lipid, a non-cationic lipid, a PEG lipid and a structural lipid. As a non-limiting example, the lipid nanoparticle may comprise 40-60% of cationic lipid, 5-15% of a non-cationic lipid, 1-2% of a PEG lipid and 30-50% of a structural lipid. As another non-limiting example, the lipid nanoparticle may comprise 50% cationic lipid, 10% non-cationic lipid, 1.5% PEG lipid and 38.5% structural lipid. As yet another non-limiting example, the lipid nanoparticle may comprise 55% cationic lipid, 10% non-cationic lipid, 2.5% PEG lipid and 32.5% structural lipid. In some embodiments, the cationic lipid may be any cationic lipid described herein such as, but not limited to, DLin-KC2-DMA, DLin-MC3-DMA and L319.

(295) In some embodiments, the lipid nanoparticle formulations described herein may comprise a cationic lipid, a non-cationic lipid, a PEG lipid and a structural lipid. As a non-limiting example, the lipid nanoparticle comprise 50% of the cationic lipid DLin-KC2-DMA, 10% of the non-cationic lipid DSPC, 1.5% of the PEG lipid PEG-DOMG and 38.5% of the structural lipid cholesterol. As a non-limiting example, the lipid nanoparticle comprise 50% of the cationic lipid DLin-MC3-DMA, 10% of the non-cationic lipid DSPC, 1.5% of the PEG lipid PEG-DOMG and 38.5% of the structural lipid cholesterol. As a non-limiting example, the lipid nanoparticle comprise 50% of the cationic lipid DLin-MC3-DMA, 10% of the non-cationic lipid DSPC, 1.5% of the PEG lipid PEG-DMG and 38.5% of the structural lipid cholesterol. As yet another non-limiting example, the lipid nanoparticle comprise 55% of the cationic lipid L319, 10% of the non-cationic lipid DSPC, 2.5% of the PEG lipid PEG-DMG and 32.5% of the structural lipid cholesterol.

(296) Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a vaccine composition may vary, depending upon the identity, size, and/or condition of the subject being treated and further depending upon the route by which the composition is to be administered. For example, the composition may comprise between 0.1% and 99% (w/w) of the active ingredient. By way of example, the composition may comprise between 0.1% and 100%, e.g., between 0.5 and 50%, between 1-30%, between 5-80%, at least 80% (w/w) active ingredient.

(297) In some embodiments, the respiratory virus RNA (e.g. mRNA) vaccine composition may comprise the polynucleotide described herein, formulated in a lipid nanoparticle comprising MC3, Cholesterol, DSPC and PEG2000-DMG, the buffer trisodium citrate, sucrose and water for injection. As a non-limiting example, the composition comprises: 2.0 mg/mL of drug substance (e.g., polynucleotides encoding H10N8 hMPV), 21.8 mg/mL of MC3, 10.1 mg/mL of cholesterol, 5.4 mg/mL of DSPC, 2.7 mg/mL of PEG2000-DMG, 5.16 mg/mL of trisodium citrate, 71 mg/mL of sucrose and 1.0 mL of water for injection.

(298) In some embodiments, a nanoparticle (e.g., a lipid nanoparticle) has a mean diameter of 10-500 nm, 20-400 nm, 30-300 nm, 40-200 nm. In some embodiments, a nanoparticle (e.g., a lipid nanoparticle) has a mean diameter of 50-150 nm, 50-200 nm, 80-100 nm or 80-200 nm.

(299) Liposomes, Lipoplexes, and Lipid Nanoparticles

(300) The RNA (e.g., mRNA) vaccines of the disclosure can be formulated using one or more liposomes, lipoplexes, or lipid nanoparticles. In some embodiments, pharmaceutical compositions of RNA (e.g., mRNA) vaccines include liposomes. Liposomes are artificially-prepared vesicles which may primarily be composed of a lipid bilayer and may be used as a delivery vehicle for the administration of nutrients and pharmaceutical formulations. Liposomes can be of different sizes such as, but not limited to, a multilamellar vesicle (MLV) which may be hundreds of nanometers in diameter and may contain a series of concentric bilayers separated by narrow aqueous compartments, a small unicellular vesicle (SUV) which may be smaller than 50 nm in diameter, and a large unilamellar vesicle (LUV) which may be between 50 and 500 nm in diameter. Liposome design may include, but is not limited to, opsonins or ligands in order to improve the attachment of liposomes to unhealthy tissue or to activate events such as, but not limited to, endocytosis. Liposomes may contain a low or a high pH in order to improve the delivery of the pharmaceutical formulations.

(301) The formation of liposomes may depend on the physicochemical characteristics such as, but not limited to, the pharmaceutical formulation entrapped and the liposomal ingredients, the nature of the medium in which the lipid vesicles are dispersed, the effective concentration of the entrapped substance and its potential toxicity, any additional processes involved during the application and/or delivery of the vesicles, the optimization size, polydispersity and the shelf-life of the vesicles for the intended application, and the batch-to-batch reproducibility and possibility of large-scale production of safe and efficient liposomal products.

(302) In some embodiments, pharmaceutical compositions described herein may include, without limitation, liposomes such as those formed from 1,2-dioleoyloxy-N,N-dimethylaminopropane (DODMA) liposomes, DiLa2 liposomes from Marina Biotech (Bothell, WA), 1,2-dilinoleoyloxy-3-dimethylaminopropane (DLin-DMA), 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), and MC3 (US20100324120; herein incorporated by reference in its entirety) and liposomes which may deliver small molecule drugs such as, but not limited to, DOXIL® from Janssen Biotech, Inc. (Horsham, PA).

(303) In some embodiments, pharmaceutical compositions described herein may include, without limitation, liposomes such as those formed from the synthesis of stabilized plasmid-lipid particles (SPLP) or stabilized nucleic acid lipid particle (SNALP) that have been previously described and shown to be suitable for oligonucleotide delivery in vitro and in vivo (see Wheeler et al. Gene Therapy. 1999 6:271-281; Zhang et al. Gene Therapy. 1999 6:1438-1447; Jeffs et al. Pharm Res. 2005 22:362-372; Morrissey et al., Nat Biotechnol. 2005 2:1002-1007; Zimmermann et al., Nature. 2006 441:111-114; Heyes et al. J Contr Rel. 2005 107:276-287; Semple et al. Nature Biotech. 2010 28:172-176; Judge et al. J Clin Invest. 2009 119:661-673; deFougerolles Hum Gene Ther. 2008 19:125-132; U.S. Patent Publication No US20130122104; all of which are incorporated herein in their entireties). The original manufacture method by Wheeler et al. was a detergent dialysis method, which was later improved by Jeffs et al. and is referred to as the spontaneous vesicle formation method. The liposome formulations are composed of 3 to 4 lipid components in addition to the polynucleotide. As an example a liposome can contain, but is not limited to, 55% cholesterol, 20% distearylphosphatidyl choline (DSPC), 10% PEG-S-DSG, and 15% 1,2-dioleoyloxy-N,N-dimethylaminopropane (DODMA), as described by Jeffs et al. As another example, certain liposome formulations may contain, but are not limited to, 48% cholesterol, 20% DSPC, 2% PEG-c-DMA, and 30% cationic lipid, where the cationic lipid can be 1,2-distearloxy-N,N-dimethylaminopropane (DSDMA), DODMA, DLin-DMA, or 1,2-dilinolenyloxy-3-dimethylaminopropane (DLenDMA), as described by Heyes et al.

(304) In some embodiments, liposome formulations may comprise from about 25.0% cholesterol to about 40.0% cholesterol, from about 30.0% cholesterol to about 45.0% cholesterol, from about 35.0% cholesterol to about 50.0% cholesterol and/or from about 48.5% cholesterol to about 60%

cholesterol. In some embodiments, formulations may comprise a percentage of cholesterol selected from the group consisting of 28.5%, 31.5%, 33.5%, 36.5%, 37.0%, 38.5%, 39.0% and 43.5%. In some embodiments, formulations may comprise from about 5.0% to about 10.0% DSPC and/or from about 7.0% to about 15.0% DSPC.

(305) In some embodiments, the RNA (e.g., mRNA) vaccine pharmaceutical compositions may be formulated in liposomes such as, but not limited to, DiLa2 liposomes (Marina Biotech, Bothell, WA), SMARTICLES® (Marina Biotech, Bothell, WA), neutral DOPC (1,2-dioleoyl-sn-glycero-3-phosphocholine) based liposomes (e.g., siRNA delivery for ovarian cancer (Landen et al. Cancer Biology & Therapy 2006 5 (12) 1708-1713); herein incorporated by reference in its entirety) and hyaluronan-coated liposomes (Quiet Therapeutics, Israel).

(306) In some embodiments, the cationic lipid may be a low molecular weight cationic lipid such as those described in U.S. Patent Application No. 20130090372, the contents of which are herein incorporated by reference in their entirety.

(307) In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in a lipid vesicle, which may have crosslinks between functionalized lipid bilayers.

(308) In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in a lipid-polycation complex. The formation of the lipid-polycation complex may be accomplished by methods known in the art and/or as described in U.S. Pub. No. 20120178702, herein incorporated by reference in its entirety. As a non-limiting example, the polycation may include a cationic peptide or a polypeptide such as, but not limited to, polylysine, polyornithine and/or polyarginine. In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in a lipid-polycation complex, which may further include a non-cationic lipid such as, but not limited to, cholesterol or dioleoyl phosphatidylethanolamine (DOPE).

(309) In some embodiments, the ratio of PEG in the lipid nanoparticle (LNP) formulations may be increased or decreased and/or the carbon chain length of the PEG lipid may be modified from C14 to C18 to alter the pharmacokinetics and/or biodistribution of the LNP formulations. As a non-limiting example, LNP formulations may contain from about 0.5% to about 3.0%, from about 1.0% to about 3.5%, from about 1.5% to about 4.0%, from about 2.0% to about 4.5%, from about 2.5% to about 5.0% and/or from about 3.0% to about 6.0% of the lipid molar ratio of PEG-c-DOMG (R-3-[(ω-methoxy-poly(ethyleneglycol)2000)carbamoyl]-1,2-dimyristyloxypropyl-3-amine) (also referred to herein as PEG-DOMG) as compared to the cationic lipid, DSPC and cholesterol. In some embodiments, the PEG-c-DOMG may be replaced with a PEG lipid such as, but not limited to, PEG-DSG (1,2-Distearoyl-sn-glycerol, methoxypolyethylene glycol), PEG-DMG (1,2-Dimyristoyl-sn-glycerol) and/or PEG-DPG (1,2-Dipalmitoyl-sn-glycerol, methoxypolyethylene glycol). The cationic lipid may be selected from any lipid known in the art such as, but not limited to, DLin-MC3-DMA, DLin-DMA, C12-200 and DLin-KC2-DMA.

(310) In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in a lipid nanoparticle.

(311) In some embodiments, the RNA (e.g., mRNA) vaccine formulation comprising the polynucleotide is a nanoparticle which may comprise at least one lipid. The lipid may be selected from, but is not limited to, DLin-DMA, DLin-K-DMA, 98N12-5, C12-200, DLin-MC3-DMA, DLin-KC2-DMA, DODMA, PLGA, PEG, PEG-DMG, PEGylated lipids and amino alcohol lipids. In another aspect, the lipid may be a cationic lipid such as, but not limited to, DLin-DMA, DLin-D-DMA, DLin-MC3-DMA, DLin-KC2-DMA, DODMA and amino alcohol lipids. The amino alcohol cationic lipid may be the lipids described in and/or made by the methods described in U.S. Patent Publication No. US20130150625, herein incorporated by reference in its entirety. As a non-limiting example, the cationic lipid may be 2-amino-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[[[(9Z,2Z)-octadeca-9,12-dien-1-yloxy]methyl]propan-1-ol (Compound 1 in US20130150625); 2-amino-3-[(9Z)-octadec-9-en-1-yloxy]-2-[[[(9Z)-octadec-9-en-1-yloxy]methyl]propan-1-ol (Compound 2 in US20130150625); 2-amino-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[(octyloxy)methyl]propan-

1-ol (Compound 3 in US20130150625); and 2-(dimethylamino)-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-2-[[[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]methyl]propan-1-ol (Compound 4 in US20130150625); or any pharmaceutically acceptable salt or stereoisomer thereof.

(312) Lipid nanoparticle formulations typically comprise a lipid, in particular, an ionizable cationic lipid, for example, 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), or di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), and further comprise a neutral lipid, a sterol and a molecule capable of reducing particle aggregation, for example a PEG or PEG-modified lipid.

(313) In some embodiments, the lipid nanoparticle formulation consists essentially of (i) at least one lipid selected from the group consisting of 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319); (ii) a neutral lipid selected from DSPC, DPPC, POPC, DOPE and SM; (iii) a sterol, e.g., cholesterol; and (iv) a PEG-lipid, e.g., PEG-DMG or PEG-cDMA, in a molar ratio of about 20-60% cationic lipid: 5-25% neutral lipid: 25-55% sterol; 0.5-15% PEG-lipid.

(314) In some embodiments, the formulation includes from about 25% to about 75% on a molar basis of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), e.g., from about 35 to about 65%, from about 45 to about 65%, about 60%, about 57.5%, about 50% or about 40% on a molar basis.

(315) In some embodiments, the formulation includes from about 0.5% to about 15% on a molar basis of the neutral lipid e.g., from about 3 to about 12%, from about 5 to about 10% or about 15%, about 10%, or about 7.5% on a molar basis. Examples of neutral lipids include, but are not limited to, DSPC, POPC, DPPC, DOPE and SM. In some embodiments, the formulation includes from about 5% to about 50% on a molar basis of the sterol (e.g., about 15 to about 45%, about 20 to about 40%, about 40%, about 38.5%, about 35%, or about 31% on a molar basis. An exemplary sterol is cholesterol. In some embodiments, the formulation includes from about 0.5% to about 20% on a molar basis of the PEG or PEG-modified lipid (e.g., about 0.5 to about 10%, about 0.5 to about 5%, about 1.5%, about 0.5%, about 1.5%, about 3.5%, or about 5% on a molar basis. In some embodiments, the PEG or PEG modified lipid comprises a PEG molecule of an average molecular weight of 2,000 Da. In other embodiments, the PEG or PEG modified lipid comprises a PEG molecule of an average molecular weight of less than 2,000, for example around 1,500 Da, around 1,000 Da, or around 500 Da. Examples of PEG-modified lipids include, but are not limited to, PEG-distearoyl glycerol (PEG-DMG) (also referred herein as PEG-C14 or C14-PEG), PEG-cDMA (further discussed in Reyes et al. *J. Controlled Release*, 107, 276-287 (2005) the contents of which are herein incorporated by reference in their entirety)

(316) In some embodiments, the formulations of the present disclosure include 25-75% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 0.5-15% of the neutral lipid, 5-50% of the sterol, and 0.5-20% of the PEG or PEG-modified lipid on a molar basis.

(317) In some embodiments, the formulations of the present disclosure include 35-65% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 3-12% of the neutral lipid, 15-45% of the sterol, and 0.5-10% of the PEG or PEG-modified lipid on a molar basis.

(318) In some embodiments, the formulations of the present disclosure include 45-65% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-

DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), 5-10% of the neutral lipid, 25-40% of the sterol, and 0.5-10% of the PEG or PEG-modified lipid on a molar basis.

(319) In some embodiments, the formulations of the present disclosure include about 60% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), about 7.5% of the neutral lipid, about 31% of the sterol, and about 1.5% of the PEG or PEG-modified lipid on a molar basis.

(320) In some embodiments, the formulations of the present disclosure include about 50% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), about 10% of the neutral lipid, about 38.5% of the sterol, and about 1.5% of the PEG or PEG-modified lipid on a molar basis.

(321) In some embodiments, the formulations of the present disclosure include about 50% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), about 10% of the neutral lipid, about 35% of the sterol, about 4.5% or about 5% of the PEG or PEG-modified lipid, and about 0.5% of the targeting lipid on a molar basis.

(322) In some embodiments, the formulations of the present disclosure include about 40% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), about 15% of the neutral lipid, about 40% of the sterol, and about 5% of the PEG or PEG-modified lipid on a molar basis.

(323) In some embodiments, the formulations of the present disclosure include about 57.2% of a cationic lipid selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319), about 7.1% of the neutral lipid, about 34.3% of the sterol, and about 1.4% of the PEG or PEG-modified lipid on a molar basis.

(324) In some embodiments, the formulations of the present disclosure include about 57.5% of a cationic lipid selected from the PEG lipid is PEG-cDMA (PEG-cDMA is further discussed in Reyes et al. (J. Controlled Release, 107, 276-287 (2005), the contents of which are herein incorporated by reference in their entirety), about 7.5% of the neutral lipid, about 31.5% of the sterol, and about 3.5% of the PEG or PEG-modified lipid on a molar basis.

(325) In some embodiments, lipid nanoparticle formulation consists essentially of a lipid mixture in molar ratios of about 20-70% cationic lipid: 5-45% neutral lipid: 20-55% cholesterol: 0.5-15% PEG-modified lipid; more preferably in a molar ratio of about 20-60% cationic lipid: 5-25% neutral lipid: 25-55% cholesterol: 0.5-15% PEG-modified lipid.

(326) In some embodiments, the molar lipid ratio is approximately 50/10/38.5/1.5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG, PEG-DSG or PEG-DPG), 57.2/7.1/34.3/1.4 (mol % cationic lipid/neutral lipid, e.g., DPPC/Chol/PEG-modified lipid, e.g., PEG-cDMA), 40/15/40/5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG), 50/10/35/4.5/0.5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DSG), 50/10/35/5 (cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG), 40/10/40/10 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG or PEG-cDMA), 35/15/40/10 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG or PEG-cDMA) or 52/13/30/5 (mol % cationic lipid/neutral lipid, e.g., DSPC/Chol/PEG-modified lipid, e.g., PEG-DMG or PEG-cDMA).

(327) Examples of lipid nanoparticle compositions and methods of making same are described, for

example, in Semple et al. (2010) *Nat. Biotechnol.* 28:172-176; Jayarama et al. (2012), *Angew. Chem. Int. Ed.*, 51: 8529-8533; and Maier et al. (2013) *Molecular Therapy* 21, 1570-1578 (the contents of each of which are incorporated herein by reference in their entirety).

(328) In some embodiments, the lipid nanoparticle formulations described herein may comprise a cationic lipid, a PEG lipid and a structural lipid and optionally comprise a non-cationic lipid. As a non-limiting example, the lipid nanoparticle may comprise about 40-60% of cationic lipid, about 5-15% of a non-cationic lipid, about 1-2% of a PEG lipid and about 30-50% of a structural lipid. As another non-limiting example, the lipid nanoparticle may comprise about 50% cationic lipid, about 10% non-cationic lipid, about 1.5% PEG lipid and about 38.5% structural lipid. As yet another non-limiting example, the lipid nanoparticle may comprise about 55% cationic lipid, about 10% non-cationic lipid, about 2.5% PEG lipid and about 32.5% structural lipid. In some embodiments, the cationic lipid may be any cationic lipid described herein such as, but not limited to, DLin-KC2-DMA, DLin-MC3-DMA and L319.

(329) In some embodiments, the lipid nanoparticle formulations described herein may be 4 component lipid nanoparticles. The lipid nanoparticle may comprise a cationic lipid, a non-cationic lipid, a PEG lipid and a structural lipid. As a non-limiting example, the lipid nanoparticle may comprise about 40-60% of cationic lipid, about 5-15% of a non-cationic lipid, about 1-2% of a PEG lipid and about 30-50% of a structural lipid. As another non-limiting example, the lipid nanoparticle may comprise about 50% cationic lipid, about 10% non-cationic lipid, about 1.5% PEG lipid and about 38.5% structural lipid. As yet another non-limiting example, the lipid nanoparticle may comprise about 55% cationic lipid, about 10% non-cationic lipid, about 2.5% PEG lipid and about 32.5% structural lipid. In some embodiments, the cationic lipid may be any cationic lipid described herein such as, but not limited to, DLin-KC2-DMA, DLin-MC3-DMA and L319.

(330) In some embodiments, the lipid nanoparticle formulations described herein may comprise a cationic lipid, a non-cationic lipid, a PEG lipid and a structural lipid. As a non-limiting example, the lipid nanoparticle comprise about 50% of the cationic lipid DLin-KC2-DMA, about 10% of the non-cationic lipid DSPC, about 1.5% of the PEG lipid PEG-DOMG and about 38.5% of the structural lipid cholesterol. As a non-limiting example, the lipid nanoparticle comprise about 50% of the cationic lipid DLin-MC3-DMA, about 10% of the non-cationic lipid DSPC, about 1.5% of the PEG lipid PEG-DOMG and about 38.5% of the structural lipid cholesterol. As a non-limiting example, the lipid nanoparticle comprise about 50% of the cationic lipid DLin-MC3-DMA, about 10% of the non-cationic lipid DSPC, about 1.5% of the PEG lipid PEG-DMG and about 38.5% of the structural lipid cholesterol. As yet another non-limiting example, the lipid nanoparticle comprise about 55% of the cationic lipid L319, about 10% of the non-cationic lipid DSPC, about 2.5% of the PEG lipid PEG-DMG and about 32.5% of the structural lipid cholesterol.

(331) As a non-limiting example, the cationic lipid may be selected from (20Z,23Z)—N,N-dimethylnonacos-20,23-dien-10-amine, (17Z,20Z)—N,N-dimethylhexacos-17,20-dien-9-amine, (1Z,19Z)—N,N-dimethylpentacos-1,16,19-dien-8-amine, (13Z,16Z)—N,N-dimethyldocos-13,16-dien-5-amine, (12Z,15Z)—N,N-dimethylheneicos-12,15-dien-4-amine, (14Z,17Z)—N,N-dimethyltricos-14,17-dien-6-amine, (15Z,18Z)—N,N-dimethyltetracos-15,18-dien-7-amine, (18Z,21Z)—N,N-dimethylheptacos-18,21-dien-10-amine, (15Z,18Z)—N,N-dimethyltetracos-15,18-dien-5-amine, (14Z,17Z)—N,N-dimethyltricos-14,17-dien-4-amine, (19Z,22Z)—N,N-dimeihyloctacos-19,22-dien-9-amine, (18Z,21 Z)—N,N-dimethylheptacos-18,21-dien-8-amine, (17Z,20Z)—N,N-dimethylhexacos-17,20-dien-7-amine, (16Z,19Z)—N,N-dimethylpentacos-16,19-dien-6-amine, (22Z,25Z)—N,N-dimethylhentriaconta-22,25-dien-10-amine, (21 Z,24Z)—N,N-dimethyltriaconta-21,24-dien-9-amine, (18Z)—N,N-dimetylheptacos-18-en-10-amine, (17Z)—N,N-dimethylhexacos-17-en-9-amine, (19Z,22Z)—N,N-dimethyloctacos-19,22-dien-7-amine, N,N-dimethylheptacos-10-amine, (20Z,23Z)—N-ethyl-N-methylnonacos-20,23-dien-10-amine, 1-[(11Z,14Z)-1-nonylicos-11,14-dien-1-yl] pyrrolidine, (20Z)—N,N-dimethylheptacos-20-en-1

O-amine, (15Z)—N,N-dimethyl eptacos-15-en-10-amine, (14Z)—N,N-dimethylnonacos-14-en-10-amine, (17Z)—N,N-dimethylnonacos-17-en-10-amine, (24Z)—N,N-dimethyltritriacont-24-en-10-amine, (20Z)—N,N-dimethylnonacos-20-en-1 O-amine, (22Z)—N,N-dimethylhentriacont-22-en-10-amine, (16Z)—N,N-dimethylpentacos-16-en-8-amine, (12Z,15Z)—N,N-dimethyl-2-nonylhenicosa-12,15-dien-1-amine, (13Z,16Z)—N,N-dimethyl-3-nonyldocosa-13,16-dien-1-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl] eptadecan-8-amine, 1-[(1S,2R)-2-hexylcyclopropyl]-N,N-dimethylnonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]nonadecan-10-amine, N,N-dimethyl-21-[(1S,2R)-2-octylcyclopropyl]henicosan-10-amine, N,N-dimethyl-1-[(1S,2S)-2-[(1R,2R)-2-pentylcyclopropyl]methyl}cyclopropyl]nonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]hexadecan-8-amine, N,N-dimethyl-[(1R,2S)-2-undecylcyclopropyl]tetradecan-5-amine, N,N-dimethyl-3-{7-[(1S,2R)-2-octylcyclopropyl]heptyl} dodecan-1-amine, 1-[(1R,2S)-2-heptylcyclopropyl]-N,N-dimethyloctadecan-9-amine, 1-[(1S,2R)-2-decylcyclopropyl]-N,N-dimethylpentadecan-6-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]pentadecan-8-amine, R—N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, S—N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}pyrrolidine, (2S)—N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-[(5Z)-oct-5-en-1-yloxy]propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}azetidine, (2S)-1-(hexyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2S)-1-(heptyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(nonyloxy)-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-[(9Z)-octadec-9-en-1-yloxy]-3-(octyloxy)propan-2-amine; (2S)—N,N-dimethyl-1-[(6Z,9Z,12Z)-octadeca-6,9,12-trien-1-yloxy]-3-(octyloxy)propan-2-amine, (2S)-1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(pentyloxy)propan-2-amine, (2S)-1-(hexyloxy)-3-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethylpropan-2-amine, 1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2S)-1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, (2S)-1-[(13Z)-docos-13-en-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, 1-[(13Z)-docos-13-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(9Z)-hexadec-9-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2R)—N,N-dimethyl-H(1-metoyloctyl)oxy]-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2R)-1-[(3,7-dimethyloctyl)oxy]-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(octyloxy)-3-({8-[(1S,2S)-2-[(1R,2R)-2-pentylcyclopropyl]methyl}cyclopropyl)octyl}oxy)propan-2-amine, N,N-dimethyl-1-{{8-(2-octylcyclopropyl)octyl}oxy}-3-(octyloxy)propan-2-amine and (1E,20Z,23Z)—N,N-dimethylnonacos-11,20,2-trien-10-amine or a pharmaceutically acceptable salt or stereoisomer thereof.

(332) In some embodiments, the LNP formulations of the RNA (e.g., mRNA) vaccines may contain PEG-c-DOMG at 3% lipid molar ratio. In some embodiments, the LNP formulations of the RNA (e.g., mRNA) vaccines may contain PEG-c-DOMG at 1.5% lipid molar ratio.

(333) In some embodiments, the pharmaceutical compositions of the RNA (e.g., mRNA) vaccines may include at least one of the PEGylated lipids described in International Publication No. WO2012099755, the contents of which are herein incorporated by reference in their entirety.

(334) In some embodiments, the LNP formulation may contain PEG-DMG 2000 (1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000]). In some embodiments, the LNP formulation may contain PEG-DMG 2000, a cationic lipid known in the art and at least one other component. In some embodiments, the LNP formulation may contain PEG-DMG 2000, a cationic lipid known in the art, DSPC and cholesterol. As a non-limiting example, the LNP formulation may contain PEG-DMG 2000, DLin-DMA, DSPC and cholesterol. As another non-

limiting example the LNP formulation may contain PEG-DMG 2000, DLin-DMA, DSPC and cholesterol in a molar ratio of 2:40:10:48 (see e.g., Geall et al., Nonviral delivery of self-amplifying RNA (e.g., mRNA) vaccines, PNAS 2012; PMID: 22908294, the contents of each of which are herein incorporated by reference in their entirety).

(335) The lipid nanoparticles described herein may be made in a sterile environment.

(336) In some embodiments, the LNP formulation may be formulated in a nanoparticle such as a nucleic acid-lipid particle. As a non-limiting example, the lipid particle may comprise one or more active agents or therapeutic agents; one or more cationic lipids comprising from about 50 mol % to about 85 mol % of the total lipid present in the particle; one or more non-cationic lipids comprising from about 13 mol % to about 49.5 mol % of the total lipid present in the particle; and one or more conjugated lipids that inhibit aggregation of particles comprising from about 0.5 mol % to about 2 mol % of the total lipid present in the particle.

(337) The nanoparticle formulations may comprise a phosphate conjugate. The phosphate conjugate may increase in vivo circulation times and/or increase the targeted delivery of the nanoparticle. As a non-limiting example, the phosphate conjugates may include a compound of any one of the formulas described in International Application No. WO2013033438, the contents of which are herein incorporated by reference in its entirety.

(338) The nanoparticle formulation may comprise a polymer conjugate. The polymer conjugate may be a water soluble conjugate. The polymer conjugate may have a structure as described in U.S. Patent Application No. 20130059360, the contents of which are herein incorporated by reference in its entirety. In some embodiments, polymer conjugates with the polynucleotides of the present disclosure may be made using the methods and/or segmented polymeric reagents described in U.S. Patent Application No. 20130072709, the contents of which are herein incorporated by reference in its entirety. In some embodiments, the polymer conjugate may have pendant side groups comprising ring moieties such as, but not limited to, the polymer conjugates described in U.S. Patent Publication No. US20130196948, the contents which are herein incorporated by reference in its entirety.

(339) The nanoparticle formulations may comprise a conjugate to enhance the delivery of nanoparticles of the present disclosure in a subject. Further, the conjugate may inhibit phagocytic clearance of the nanoparticles in a subject. In one aspect, the conjugate may be a “self” peptide designed from the human membrane protein CD47 (e.g., the “self” particles described by Rodriguez et al. (*Science* 2013 339, 971-975), herein incorporated by reference in its entirety). As shown by Rodriguez et al., the self peptides delayed macrophage-mediated clearance of nanoparticles which enhanced delivery of the nanoparticles. In another aspect, the conjugate may be the membrane protein CD47 (e.g., see Rodriguez et al. *Science* 2013 339, 971-975, herein incorporated by reference in its entirety). Rodriguez et al. showed that, similarly to “self” peptides, CD47 can increase the circulating particle ratio in a subject as compared to scrambled peptides and PEG coated nanoparticles.

(340) In some embodiments, the RNA (e.g., mRNA) vaccines of the present disclosure are formulated in nanoparticles which comprise a conjugate to enhance the delivery of the nanoparticles of the present disclosure in a subject. The conjugate may be the CD47 membrane or the conjugate may be derived from the CD47 membrane protein, such as the “self” peptide described previously. In some embodiments, the nanoparticle may comprise PEG and a conjugate of CD47 or a derivative thereof. In some embodiments, the nanoparticle may comprise both the “self” peptide described above and the membrane protein CD47.

(341) In some embodiments, a “self” peptide and/or CD47 protein may be conjugated to a virus-like particle or pseudovirion, as described herein for delivery of the RNA (e.g., mRNA) vaccines of the present disclosure.

(342) In some embodiments, RNA (e.g., mRNA) vaccine pharmaceutical compositions comprising the polynucleotides of the present disclosure and a conjugate that may have a degradable linkage.

Non-limiting examples of conjugates include an aromatic moiety comprising an ionizable hydrogen atom, a spacer moiety, and a water-soluble polymer. As a non-limiting example, pharmaceutical compositions comprising a conjugate with a degradable linkage and methods for delivering such pharmaceutical compositions are described in U.S. Patent Publication No. US20130184443, the contents of which are herein incorporated by reference in their entirety.

(343) The nanoparticle formulations may be a carbohydrate nanoparticle comprising a carbohydrate carrier and a RNA (e.g., mRNA) vaccine. As a non-limiting example, the carbohydrate carrier may include, but is not limited to, an anhydride-modified phytoglycogen or glycogen-type material, phytoglycogen octenyl succinate, phytoglycogen beta-dextrin, anhydride-modified phytoglycogen beta-dextrin. (See e.g., International Publication No. WO2012109121; the contents of which are herein incorporated by reference in their entirety).

(344) Nanoparticle formulations of the present disclosure may be coated with a surfactant or polymer in order to improve the delivery of the particle. In some embodiments, the nanoparticle may be coated with a hydrophilic coating such as, but not limited to, PEG coatings and/or coatings that have a neutral surface charge. The hydrophilic coatings may help to deliver nanoparticles with larger payloads such as, but not limited to, RNA (e.g., mRNA) vaccines within the central nervous system. As a non-limiting example nanoparticles comprising a hydrophilic coating and methods of making such nanoparticles are described in U.S. Patent Publication No. US20130183244, the contents of which are herein incorporated by reference in their entirety.

(345) In some embodiments, the lipid nanoparticles of the present disclosure may be hydrophilic polymer particles. Non-limiting examples of hydrophilic polymer particles and methods of making hydrophilic polymer particles are described in U.S. Patent Publication No. US20130210991, the contents of which are herein incorporated by reference in their entirety.

(346) In some embodiments, the lipid nanoparticles of the present disclosure may be hydrophobic polymer particles.

(347) Lipid nanoparticle formulations may be improved by replacing the cationic lipid with a biodegradable cationic lipid which is known as a rapidly eliminated lipid nanoparticle (reLNP). Ionizable cationic lipids, such as, but not limited to, DlinDMA, Dlin-KC2-DMA, and Dlin-MC3-DMA, have been shown to accumulate in plasma and tissues over time and may be a potential source of toxicity. The rapid metabolism of the rapidly eliminated lipids can improve the tolerability and therapeutic index of the lipid nanoparticles by an order of magnitude from a 1 mg/kg dose to a 10 mg/kg dose in rat. Inclusion of an enzymatically degraded ester linkage can improve the degradation and metabolism profile of the cationic component, while still maintaining the activity of the reLNP formulation. The ester linkage can be internally located within the lipid chain or it may be terminally located at the terminal end of the lipid chain. The internal ester linkage may replace any carbon in the lipid chain.

(348) In some embodiments, the internal ester linkage may be located on either side of the saturated carbon.

(349) In some embodiments, an immune response may be elicited by delivering a lipid nanoparticle which may include a nanospecies, a polymer and an immunogen. (U.S. Publication No. 20120189700 and International Publication No. WO2012099805; each of which is herein incorporated by reference in their entirety). The polymer may encapsulate the nanospecies or partially encapsulate the nanospecies. The immunogen may be a recombinant protein, a modified RNA and/or a polynucleotide described herein. In some embodiments, the lipid nanoparticle may be formulated for use in a vaccine such as, but not limited to, against a pathogen.

(350) Lipid nanoparticles may be engineered to alter the surface properties of particles so the lipid nanoparticles may penetrate the mucosal barrier. Mucus is located on mucosal tissue such as, but not limited to, oral (e.g., the buccal and esophageal membranes and tonsil tissue), ophthalmic, gastrointestinal (e.g., stomach, small intestine, large intestine, colon, rectum), nasal, respiratory (e.g., nasal, pharyngeal, tracheal and bronchial membranes), genital (e.g., vaginal, cervical and

urethral membranes). Nanoparticles larger than 10-200 nm which are preferred for higher drug encapsulation efficiency and the ability to provide the sustained delivery of a wide array of drugs have been thought to be too large to rapidly diffuse through mucosal barriers. Mucus is continuously secreted, shed, discarded or digested and recycled so most of the trapped particles may be removed from the mucosa tissue within seconds or within a few hours. Large polymeric nanoparticles (200 nm-500 nm in diameter) which have been coated densely with a low molecular weight polyethylene glycol (PEG) diffused through mucus only 4 to 6-fold lower than the same particles diffusing in water (Lai et al. PNAS 2007 104(5):1482-487; Lai et al. *Adv Drug Deliv Rev.* 2009 61(2): 158-171; each of which is herein incorporated by reference in their entirety). The transport of nanoparticles may be determined using rates of permeation and/or fluorescent microscopy techniques including, but not limited to, fluorescence recovery after photobleaching (FRAP) and high resolution multiple particle tracking (MPT). As a non-limiting example, compositions which can penetrate a mucosal barrier may be made as described in U.S. Pat. No. 8,241,670 or International Patent Publication No. WO2013110028, the contents of each of which are herein incorporated by reference in its entirety.

(351) The lipid nanoparticle engineered to penetrate mucus may comprise a polymeric material (i.e. a polymeric core) and/or a polymer-vitamin conjugate and/or a tri-block co-polymer. The polymeric material may include, but is not limited to, polyamines, polyethers, polyamides, polyesters, polycarbamates, polyureas, polycarbonates, poly(styrenes), polyimides, polysulfones, polyurethanes, polyacetylenes, polyethylenes, polyethyleneimines, polyisocyanates, polyacrylates, polymethacrylates, polyacrylonitriles, and polyarylates. The polymeric material may be biodegradable and/or biocompatible. Non-limiting examples of biocompatible polymers are described in International Patent Publication No. WO2013116804, the contents of which are herein incorporated by reference in their entirety. The polymeric material may additionally be irradiated. As a non-limiting example, the polymeric material may be gamma irradiated (see e.g., International App. No. WO201282165, herein incorporated by reference in its entirety). Non-limiting examples of specific polymers include poly(caprolactone) (PCL), ethylene vinyl acetate polymer (EVA), poly(lactic acid) (PLA), poly(L-lactic acid) (PLLA), poly(glycolic acid) (PGA), poly(lactic acid-co-glycolic acid) (PLGA), poly(L-lactic acid-co-glycolic acid) (PLLGA), poly(D,L-lactide) (PDLA), poly(L-lactide) (PLLA), poly(D,L-lactide-co-caprolactone), poly(D,L-lactide-co-caprolactone-co-glycolide), poly(D,L-lactide-co-PEO-co-D,L-lactide), poly(D,L-lactide-co-PPO-co-D,L-lactide), polyalkyl cyanoacrylate, polyurethane, poly-L-lysine (PLL), hydroxypropyl methacrylate (HPMA), polyethyleneglycol, poly-L-glutamic acid, poly(hydroxy acids), polyanhydrides, polyorthoesters, poly(ester amides), polyamides, poly(ester ethers), polycarbonates, polyalkylenes such as polyethylene and polypropylene, polyalkylene glycols such as poly(ethylene glycol) (PEG), polyalkylene oxides (PEO), polyalkylene terephthalates such as poly(ethylene terephthalate), polyvinyl alcohols (PVA), polyvinyl ethers, polyvinyl esters such as poly(vinyl acetate), polyvinyl halides such as poly(vinyl chloride) (PVC), polyvinylpyrrolidone, polysiloxanes, polystyrene (PS), polyurethanes, derivatized celluloses such as alkyl celluloses, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, hydroxypropylcellulose, carboxymethylcellulose, polymers of acrylic acids, such as poly(methyl(meth)acrylate) (PMMA), poly(ethyl(meth)acrylate), poly(butyl(meth)acrylate), poly(isobutyl(meth)acrylate), poly(hexyl(meth)acrylate), poly(isodecyl(meth)acrylate), poly(lauryl(meth)acrylate), poly(phenyl(meth)acrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate) and copolymers and mixtures thereof, polydioxanone and its copolymers, polyhydroxyalkanoates, polypropylene fumarate, polyoxymethylene, poloxamers, poly(ortho)esters, poly(butyric acid), poly(valeric acid), poly(lactide-co-caprolactone), PEG-PLGA-PEG and trimethylene carbonate, polyvinylpyrrolidone. The lipid nanoparticle may be coated or associated with a co-polymer such as, but not limited to, a block co-polymer (such as a branched polyether-polyamide block copolymer described in International Publication No.

WO2013012476, herein incorporated by reference in its entirety), and (poly(ethylene glycol))- (poly(propylene oxide))-(poly(ethylene glycol)) triblock copolymer (see e.g., U.S. Publication 20120121718 and U.S. Publication 20100003337 and U.S. Pat. No. 8,263,665, the contents of each of which is herein incorporated by reference in their entirety). The co-polymer may be a polymer that is generally regarded as safe (GRAS) and the formation of the lipid nanoparticle may be in such a way that no new chemical entities are created. For example, the lipid nanoparticle may comprise poloxamers coating PLGA nanoparticles without forming new chemical entities which are still able to rapidly penetrate human mucus (Yang et al. *Angew. Chem. Int. Ed.* 2011 50:2597-2600; the contents of which are herein incorporated by reference in their entirety). A non-limiting scalable method to produce nanoparticles which can penetrate human mucus is described by Xu et al. (see, e.g., *J Control Release* 2013, 170(2):279-86; the contents of which are herein incorporated by reference in their entirety).

(352) The vitamin of the polymer-vitamin conjugate may be vitamin E. The vitamin portion of the conjugate may be substituted with other suitable components such as, but not limited to, vitamin A, vitamin E, other vitamins, cholesterol, a hydrophobic moiety, or a hydrophobic component of other surfactants (e.g., sterol chains, fatty acids, hydrocarbon chains and alkylene oxide chains).

(353) The lipid nanoparticle engineered to penetrate mucus may include surface altering agents such as, but not limited to, polynucleotides, anionic proteins (e.g., bovine serum albumin), surfactants (e.g., cationic surfactants such as for example dimethyldioctadecylammonium bromide), sugars or sugar derivatives (e.g., cyclodextrin), nucleic acids, polymers (e.g., heparin, polyethylene glycol and poloxamer), mucolytic agents (e.g., N-acetylcysteine, mugwort, bromelain, papain, clerodendrum, acetylcysteine, bromhexine, carbocysteine, eprazinone, mesna, ambroxol, sobrerol, domiodol, letosteine, stepronin, tiopronin, gelsolin, thymosin 34 dornase alfa, neltenexine, erdosteine) and various DNases including rhDNase. The surface altering agent may be embedded or enmeshed in the particle's surface or disposed (e.g., by coating, adsorption, covalent linkage, or other process) on the surface of the lipid nanoparticle. (see e.g., U.S. Publication 20100215580 and U.S. Publication 20080166414 and US20130164343; the contents of each of which are herein incorporated by reference in their entirety).

(354) In some embodiments, the mucus penetrating lipid nanoparticles may comprise at least one polynucleotide described herein. The polynucleotide may be encapsulated in the lipid nanoparticle and/or disposed on the surface of the particle. The polynucleotide may be covalently coupled to the lipid nanoparticle. Formulations of mucus penetrating lipid nanoparticles may comprise a plurality of nanoparticles. Further, the formulations may contain particles which may interact with the mucus and alter the structural and/or adhesive properties of the surrounding mucus to decrease mucoadhesion, which may increase the delivery of the mucus penetrating lipid nanoparticles to the mucosal tissue.

(355) In some embodiments, the mucus penetrating lipid nanoparticles may be a hypotonic formulation comprising a mucosal penetration enhancing coating. The formulation may be hypotonic for the epithelium to which it is being delivered. Non-limiting examples of hypotonic formulations may be found in International Patent Publication No. WO2013110028, the contents of which are herein incorporated by reference in their entirety.

(356) In some embodiments, in order to enhance the delivery through the mucosal barrier the RNA (e.g., mRNA) vaccine formulation may comprise or be a hypotonic solution. Hypotonic solutions were found to increase the rate at which mucoinert particles such as, but not limited to, mucus-penetrating particles, were able to reach the vaginal epithelial surface (see e.g., Ensign et al. *Biomaterials* 2013 34(28):6922-9, the contents of which are herein incorporated by reference in their entirety).

(357) In some embodiments, the RNA (e.g., mRNA) vaccine is formulated as a lipoplex, such as, without limitation, the ATUPLEX™ system, the DACC system, the DBTC system and other siRNA-lipoplex technology from Silence Therapeutics (London, United Kingdom), STEMFECT™

from STEMAGENT® (Cambridge, MA), and polyethylenimine (PEI) or protamine-based targeted and non-targeted delivery of nucleic acids (Aleku et al. *Cancer Res.* 2008 68:9788-9798; Strumberg et al. *Int J Clin Pharmacol Ther* 2012 50:76-78; Santel et al., *Gene Ther* 2006 13:1222-1234; Santel et al., *Gene Ther* 2006 13:1360-1370; Gutbier et al., *Pulm Pharmacol. Ther.* 2010 23:334-344; Kaufmann et al. *Microvasc Res* 2010 80:286-293 Weide et al. *J Immunother.* 2009 32:498-507; Weide et al. *J Immunother.* 2008 31:180-188; Pascolo *Expert Opin. Biol. Ther.* 4:1285-1294; Fotin-Mleczek et al., 2011 *J. Immunother.* 34:1-15; Song et al., *Nature Biotechnol.* 2005, 23:709-717; Peer et al., *Proc Natl Acad Sci USA.* 2007 6; 104:4095-4100; deFougerolles *Hum Gene Ther.* 2008 19:125-132, the contents of each of which are incorporated herein by reference in their entirety).

(358) In some embodiments, such formulations may also be constructed or compositions altered such that they passively or actively are directed to different cell types in vivo, including but not limited to hepatocytes, immune cells, tumor cells, endothelial cells, antigen presenting cells, and leukocytes (Akinc et al. *Mol Ther.* 2010 18:1357-1364; Song et al., *Nat Biotechnol.* 2005 23:709-717; Judge et al., *J Clin Invest.* 2009 119:661-673; Kaufmann et al., *Microvasc Res* 2010 80:286-293; Santel et al., *Gene Ther* 2006 13:1222-1234; Santel et al., *Gene Ther* 2006 13:1360-1370; Gutbier et al., *Pulm Pharmacol. Ther.* 2010 23:334-344; Basha et al., *Mol. Ther.* 2011 19:2186-2200; Fenske and Cullis, *Expert Opin Drug Deliv.* 2008 5:25-44; Peer et al., *Science.* 2008 319:627-630; Peer and Lieberman, *Gene Ther.* 2011 18:1127-1133, the contents of each of which are incorporated herein by reference in their entirety). One example of passive targeting of formulations to liver cells includes the DLin-DMA, DLin-KC2-DMA and DLin-MC3-DMA-based lipid nanoparticle formulations, which have been shown to bind to apolipoprotein E and promote binding and uptake of these formulations into hepatocytes in vivo (Akinc et al. *Mol Ther.* 2010 18:1357-1364, the contents of which are incorporated herein by reference in their entirety). Formulations can also be selectively targeted through expression of different ligands on their surface as exemplified by, but not limited by, folate, transferrin, N-acetylgalactosamine (GalNAc), and antibody targeted approaches (Kolhatkar et al., *Curr Drug Discov Technol.* 2011 8:197-206; Musacchio and Torchilin, *Front Biosci.* 2011 16:1388-1412; Yu et al., *Mol Membr Biol.* 2010 27:286-298; Patil et al., *Crit Rev Ther Drug Carrier Syst.* 2008 25:1-61; Benoit et al., *Biomacromolecules.* 2011 12:2708-2714; Zhao et al., *Expert Opin Drug Deliv.* 2008 5:309-319; Akinc et al., *Mol Ther.* 2010 18:1357-1364; Srinivasan et al., *Methods Mol Biol.* 2012 820:105-116; Ben-Arie et al., *Methods Mol Biol.* 2012 757:497-507; Peer 2010 *J Control Release.* 20:63-68; Peer et al., *Proc Natl Acad Sci USA.* 2007 104:4095-4100; Kim et al., *Methods Mol Biol.* 2011 721:339-353; Subramanya et al., *Mol Ther.* 2010 18:2028-2037; Song et al., *Nat Biotechnol.* 2005 23:709-717; Peer et al., *Science.* 2008 319:627-630; Peer and Lieberman, *Gene Ther.* 2011 18:1127-1133, the contents of each of which are incorporated herein by reference in their entirety).

(359) In some embodiments, the RNA (e.g., mRNA) vaccine is formulated as a solid lipid nanoparticle. A solid lipid nanoparticle (SLN) may be spherical with an average diameter between 10 to 1000 nm. SLN possess a solid lipid core matrix that can solubilize lipophilic molecules and may be stabilized with surfactants and/or emulsifiers. In some embodiments, the lipid nanoparticle may be a self-assembly lipid-polymer nanoparticle (see Zhang et al., *ACS Nano*, 2008, 2 (8), pp 1696-1702; the contents of which are herein incorporated by reference in their entirety). As a non-limiting example, the SLN may be the SLN described in International Patent Publication No. WO2013105101, the contents of which are herein incorporated by reference in their entirety. As another non-limiting example, the SLN may be made by the methods or processes described in International Patent Publication No. WO2013105101, the contents of which are herein incorporated by reference in their entirety.

(360) Liposomes, lipoplexes, or lipid nanoparticles may be used to improve the efficacy of polynucleotides directed protein production as these formulations may be able to increase cell transfection by the RNA (e.g., mRNA) vaccine; and/or increase the translation of encoded protein.

One such example involves the use of lipid encapsulation to enable the effective systemic delivery of polyplex plasmid DNA (Heyes et al., Mol Ther. 2007 15:713-720; the contents of which are incorporated herein by reference in their entirety). The liposomes, lipoplexes, or lipid nanoparticles may also be used to increase the stability of the polynucleotide.

(361) In some embodiments, the RNA (e.g., mRNA) vaccines of the present disclosure can be formulated for controlled release and/or targeted delivery. As used herein, “controlled release” refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome. In some embodiments, the RNA (e.g., mRNA) vaccines may be encapsulated into a delivery agent described herein and/or known in the art for controlled release and/or targeted delivery. As used herein, the term “encapsulate” means to enclose, surround or encase. As it relates to the formulation of the compounds of the disclosure, encapsulation may be substantial, complete or partial. The term “substantially encapsulated” means that at least greater than 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.9 or greater than 99.999% of the pharmaceutical composition or compound of the disclosure may be enclosed, surrounded or encased within the delivery agent. “Partially encapsulation” means that less than 10, 10, 20, 30, 40 50 or less of the pharmaceutical composition or compound of the disclosure may be enclosed, surrounded or encased within the delivery agent. Advantageously, encapsulation may be determined by measuring the escape or the activity of the pharmaceutical composition or compound of the disclosure using fluorescence and/or electron micrograph. For example, at least 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the pharmaceutical composition or compound of the disclosure are encapsulated in the delivery agent.

(362) In some embodiments, the controlled release formulation may include, but is not limited to, tri-block co-polymers. As a non-limiting example, the formulation may include two different types of tri-block co-polymers (International Pub. No. WO2012131104 and WO2012131106, the contents of each of which are incorporated herein by reference in their entirety).

(363) In some embodiments, the RNA (e.g., mRNA) vaccines may be encapsulated into a lipid nanoparticle or a rapidly eliminated lipid nanoparticle and the lipid nanoparticles or a rapidly eliminated lipid nanoparticle may then be encapsulated into a polymer, hydrogel and/or surgical sealant described herein and/or known in the art. As a non-limiting example, the polymer, hydrogel or surgical sealant may be PLGA, ethylene vinyl acetate (EVAc), poloxamer, GELSTTE® (Nanotherapeutics, Inc. Alachua, FL), HYLENEX® (Halozyne Therapeutics, San Diego CA), surgical sealants such as fibrinogen polymers (Ethicon Inc. Cornelia, GA), TISSELL® (Baxter International, Inc Deerfield, IL), PEG-based sealants, and COSEAL® (Baxter International, Inc Deerfield, IL).

(364) In some embodiments, the lipid nanoparticle may be encapsulated into any polymer known in the art which may form a gel when injected into a subject. As another non-limiting example, the lipid nanoparticle may be encapsulated into a polymer matrix which may be biodegradable.

(365) In some embodiments, the RNA (e.g., mRNA) vaccine formulation for controlled release and/or targeted delivery may also include at least one controlled release coating. Controlled release coatings include, but are not limited to, OPADRY®, polyvinylpyrrolidone/vinyl acetate copolymer, polyvinylpyrrolidone, hydroxypropyl methylcellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, EUDRAGIT RL®, EUDRAGIT RS® and cellulose derivatives such as ethylcellulose aqueous dispersions (AQUACOAT® and SURELEASE®).

(366) In some embodiments, the RNA (e.g., mRNA) vaccine controlled release and/or targeted delivery formulation may comprise at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In some embodiments, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

(367) In some embodiments, the RNA (e.g., mRNA) vaccine controlled release and/or targeted delivery formulation comprising at least one polynucleotide may comprise at least one PEG and/or PEG related polymer derivatives as described in U.S. Pat. No. 8,404,222, the contents of which are incorporated herein by reference in their entirety.

(368) In some embodiments, the RNA (e.g., mRNA) vaccine controlled release delivery formulation comprising at least one polynucleotide may be the controlled release polymer system described in US20130130348, the contents of which are incorporated herein by reference in their entirety.

(369) In some embodiments, the RNA (e.g., mRNA) vaccines of the present disclosure may be encapsulated in a therapeutic nanoparticle, referred to herein as “therapeutic nanoparticle RNA (e.g., mRNA) vaccines.” Therapeutic nanoparticles may be formulated by methods described herein and known in the art such as, but not limited to, International Pub Nos. WO2010005740, WO2010030763, WO2010005721, WO2010005723, WO2012054923, U.S. Publication Nos. US20110262491, US20100104645, US20100087337, US20100068285, US20110274759, US20100068286, US20120288541, US20130123351 and US20130230567 and U.S. Pat. Nos. 8,206,747, 8,293,276, 8,318,208 and 8,318,211; the contents of each of which are herein incorporated by reference in their entirety. In some embodiments, therapeutic polymer nanoparticles may be identified by the methods described in US Pub No. US20120140790, the contents of which are herein incorporated by reference in their entirety.

(370) In some embodiments, the therapeutic nanoparticle RNA (e.g., mRNA) vaccine may be formulated for sustained release. As used herein, “sustained release” refers to a pharmaceutical composition or compound that conforms to a release rate over a specific period of time. The period of time may include, but is not limited to, hours, days, weeks, months and years. As a non-limiting example, the sustained release nanoparticle may comprise a polymer and a therapeutic agent such as, but not limited to, the polynucleotides of the present disclosure (see International Pub No. 2010075072 and US Pub No. US20100216804, US20110217377 and US20120201859, the contents of each of which are incorporated herein by reference in their entirety). In another non-limiting example, the sustained release formulation may comprise agents which permit persistent bioavailability such as, but not limited to, crystals, macromolecular gels and/or particulate suspensions (see U.S. Patent Publication No US20130150295, the contents of each of which are incorporated herein by reference in their entirety).

(371) In some embodiments, the therapeutic nanoparticle RNA (e.g., mRNA) vaccines may be formulated to be target specific. As a non-limiting example, the therapeutic nanoparticles may include a corticosteroid (see International Pub. No. WO2011084518, the contents of which are incorporated herein by reference in their entirety). As a non-limiting example, the therapeutic nanoparticles may be formulated in nanoparticles described in International Pub No. WO2008121949, WO2010005726, WO2010005725, WO2011084521 and US Pub No. US20100069426, US20120004293 and US20100104655, the contents of each of which are incorporated herein by reference in their entirety.

(372) In some embodiments, the nanoparticles of the present disclosure may comprise a polymeric matrix. As a non-limiting example, the nanoparticle may comprise two or more polymers such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester) or combinations thereof.

(373) In some embodiments, the therapeutic nanoparticle comprises a diblock copolymer. In some embodiments, the diblock copolymer may include PEG in combination with a polymer such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids,

polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester) or combinations thereof. In yet another embodiment, the diblock copolymer may be a high-X diblock copolymer such as those described in International Patent Publication No. WO2013120052, the contents of which are incorporated herein by reference in their entirety.

(374) As a non-limiting example the therapeutic nanoparticle comprises a PLGA-PEG block copolymer (see U.S. Publication No. US20120004293 and U.S. Pat. No. 8,236,330, each of which is herein incorporated by reference in their entirety). In another non-limiting example, the therapeutic nanoparticle is a stealth nanoparticle comprising a diblock copolymer of PEG and PLA or PEG and PLGA (see U.S. Pat. No. 8,246,968 and International Publication No. WO2012166923, the contents of each of which are herein incorporated by reference in their entirety). In yet another non-limiting example, the therapeutic nanoparticle is a stealth nanoparticle or a target-specific stealth nanoparticle as described in U.S. Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in their entirety.

(375) In some embodiments, the therapeutic nanoparticle may comprise a multiblock copolymer (see e.g., U.S. Pat. Nos. 8,263,665 and 8,287,910 and U.S. Patent Pub. No. US20130195987, the contents of each of which are herein incorporated by reference in their entirety).

(376) In yet another non-limiting example, the lipid nanoparticle comprises the block copolymer PEG-PLGA-PEG (see e.g., the thermosensitive hydrogel (PEG-PLGA-PEG) was used as a TGF-beta1 gene delivery vehicle in Lee et al. Thermosensitive Hydrogel as a Tgf- β 1 Gene Delivery Vehicle Enhances Diabetic Wound Healing. *Pharmaceutical Research*, 2003 20(12): 1995-2000; as a controlled gene delivery system in Li et al. Controlled Gene Delivery System Based on Thermosensitive Biodegradable Hydrogel. *Pharmaceutical Research* 2003 20(6):884-888; and Chang et al., Non-ionic amphiphilic biodegradable PEG-PLGA-PEG copolymer enhances gene delivery efficiency in rat skeletal muscle. *J Controlled Release*. 2007 118:245-253, the contents of each of which are herein incorporated by reference in their entirety). The RNA (e.g., mRNA) vaccines of the present disclosure may be formulated in lipid nanoparticles comprising the PEG-PLGA-PEG block copolymer.

(377) In some embodiments, the therapeutic nanoparticle may comprise a multiblock copolymer (see e.g., U.S. Pat. Nos. 8,263,665 and 8,287,910 and U.S. Patent Pub. No. US20130195987, the contents of each of which are herein incorporated by reference in their entirety).

(378) In some embodiments, the block copolymers described herein may be included in a polyion complex comprising a non-polymeric micelle and the block copolymer. (see e.g., U.S. Publication No. 20120076836, the contents of which are herein incorporated by reference in their entirety).

(379) In some embodiments, the therapeutic nanoparticle may comprise at least one acrylic polymer. Acrylic polymers include but are not limited to, acrylic acid, methacrylic acid, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, amino alkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), polycyanoacrylates and combinations thereof.

(380) In some embodiments, the therapeutic nanoparticles may comprise at least one poly(vinyl ester) polymer. The poly(vinyl ester) polymer may be a copolymer such as a random copolymer. As a non-limiting example, the random copolymer may have a structure such as those described in International Application No. WO2013032829 or U.S. Patent Publication No US20130121954, the contents of each of which are herein incorporated by reference in their entirety. In some embodiments, the poly(vinyl ester) polymers may be conjugated to the polynucleotides described herein.

(381) In some embodiments, the therapeutic nanoparticle may comprise at least one diblock copolymer. The diblock copolymer may be, but it not limited to, a poly(lactic) acid-

poly(ethylene)glycol copolymer (see, e.g., International Patent Publication No. WO2013044219, the contents of which are herein incorporated by reference in their entirety). As a non-limiting example, the therapeutic nanoparticle may be used to treat cancer (see International publication No. WO2013044219, the contents of which are herein incorporated by reference in their entirety).

(382) In some embodiments, the therapeutic nanoparticles may comprise at least one cationic polymer described herein and/or known in the art.

(383) In some embodiments, the therapeutic nanoparticles may comprise at least one amine-containing polymer such as, but not limited to polylysine, polyethylene imine, poly(amidoamine) dendrimers, poly(beta-amino esters) (see, e.g., U.S. Pat. No. 8,287,849, the contents of which are herein incorporated by reference in their entirety) and combinations thereof.

(384) In some embodiments, the nanoparticles described herein may comprise an amine cationic lipid such as those described in International Patent Application No. WO2013059496, the contents of which are herein incorporated by reference in their entirety.

(385) In some embodiments, the cationic lipids may have an amino-amine or an amino-amide moiety. In some embodiments, the therapeutic nanoparticles may comprise at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In some embodiments, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

(386) In some embodiments, the synthetic nanocarriers may contain an immunostimulatory agent to enhance the immune response from delivery of the synthetic nanocarrier. As a non-limiting example, the synthetic nanocarrier may comprise a Th1 immunostimulatory agent, which may enhance a Th1-based response of the immune system (see International Pub No. WO2010123569 and U.S. Publication No. US20110223201, the contents of each of which are herein incorporated by reference in their entirety).

(387) In some embodiments, the synthetic nanocarriers may be formulated for targeted release. In some embodiments, the synthetic nanocarrier is formulated to release the polynucleotides at a specified pH and/or after a desired time interval. As a non-limiting example, the synthetic nanoparticle may be formulated to release the RNA (e.g., mRNA) vaccines after 24 hours and/or at a pH of 4.5 (see International Publication Nos. WO2010138193 and WO2010138194 and US Pub Nos. US20110020388 and US20110027217, each of which is herein incorporated by reference in their entireties).

(388) In some embodiments, the synthetic nanocarriers may be formulated for controlled and/or sustained release of the polynucleotides described herein. As a non-limiting example, the synthetic nanocarriers for sustained release may be formulated by methods known in the art, described herein and/or as described in International Pub No. WO2010138192 and US Pub No. 20100303850, each of which is herein incorporated by reference in their entirety.

(389) In some embodiments, the RNA (e.g., mRNA) vaccine may be formulated for controlled and/or sustained release wherein the formulation comprises at least one polymer that is a crystalline side chain (CYSC) polymer. CYSC polymers are described in U.S. Pat. No. 8,399,007, herein incorporated by reference in its entirety.

(390) In some embodiments, the synthetic nanocarrier may be formulated for use as a vaccine. In some embodiments, the synthetic nanocarrier may encapsulate at least one polynucleotide which encode at least one antigen. As a non-limiting example, the synthetic nanocarrier may include at least one antigen and an excipient for a vaccine dosage form (see International Publication No. WO2011150264 and U.S. Publication No. US20110293723, the contents of each of which are herein incorporated by reference in their entirety). As another non-limiting example, a vaccine dosage form may include at least two synthetic nanocarriers with the same or different antigens and an excipient (see International Publication No. WO2011150249 and U.S. Publication No. US20110293701, the contents of each of which are herein incorporated by reference in their

entirety). The vaccine dosage form may be selected by methods described herein, known in the art and/or described in International Publication No. WO2011150258 and U.S. Publication No. US20120027806, the contents of each of which are herein incorporated by reference in their entirety).

(391) In some embodiments, the synthetic nanocarrier may comprise at least one polynucleotide which encodes at least one adjuvant. As a non-limiting example, the adjuvant may comprise dimethyldioctadecylammonium-bromide, dimethyldioctadecylammonium-chloride, dimethyldioctadecylammonium-phosphate or dimethyldioctadecylammonium-acetate (DDA) and an apolar fraction or part of said apolar fraction of a total lipid extract of a *mycobacterium* (see, e.g., U.S. Pat. No. 8,241,610, the content of which is herein incorporated by reference in its entirety). In some embodiments, the synthetic nanocarrier may comprise at least one polynucleotide and an adjuvant. As a non-limiting example, the synthetic nanocarrier comprising and adjuvant may be formulated by the methods described in International Publication No. WO2011150240 and U.S. Publication No. US20110293700, the contents of each of which are herein incorporated by reference in their entirety.

(392) In some embodiments, the synthetic nanocarrier may encapsulate at least one polynucleotide that encodes a peptide, fragment or region from a virus. As a non-limiting example, the synthetic nanocarrier may include, but is not limited to, any of the nanocarriers described in International Publication No. WO2012024621, WO201202629, WO2012024632 and U.S. Publication No. US20120064110, US20120058153 and US20120058154, the contents of each of which are herein incorporated by reference in their entirety.

(393) In some embodiments, the synthetic nanocarrier may be coupled to a polynucleotide which may be able to trigger a humoral and/or cytotoxic T lymphocyte (CTL) response (see, e.g., International Publication No. WO2013019669, the contents of which are herein incorporated by reference in their entirety).

(394) In some embodiments, the RNA (e.g., mRNA) vaccine may be encapsulated in, linked to and/or associated with zwitterionic lipids. Non-limiting examples of zwitterionic lipids and methods of using zwitterionic lipids are described in U.S. Patent Publication No. US20130216607, the contents of which are herein incorporated by reference in their entirety. In some aspects, the zwitterionic lipids may be used in the liposomes and lipid nanoparticles described herein.

(395) In some embodiments, the RNA (e.g., mRNA) vaccine may be formulated in colloid nanocarriers as described in U.S. Patent Publication No. US20130197100, the contents of which are herein incorporated by reference in their entirety.

(396) In some embodiments, the nanoparticle may be optimized for oral administration. The nanoparticle may comprise at least one cationic biopolymer such as, but not limited to, chitosan or a derivative thereof. As a non-limiting example, the nanoparticle may be formulated by the methods described in U.S. Publication No. 20120282343, the contents of which are herein incorporated by reference in their entirety.

(397) In some embodiments, LNPs comprise the lipid KL52 (an amino-lipid disclosed in U.S. Application Publication No. 2012/0295832, the contents of which are herein incorporated by reference in their entirety. Activity and/or safety (as measured by examining one or more of ALT/AST, white blood cell count and cytokine induction, for example) of LNP administration may be improved by incorporation of such lipids. LNPs comprising KL52 may be administered intravenously and/or in one or more doses. In some embodiments, administration of LNPs comprising KL52 results in equal or improved mRNA and/or protein expression as compared to LNPs comprising MC3.

(398) In some embodiments, RNA (e.g., mRNA) vaccine may be delivered using smaller LNPs. Such particles may comprise a diameter from below 0.1 μm up to 100 nm such as, but not limited to, less than 0.1 μm , less than 1.0 μm , less than 5 μm , less than 10 μm , less than 15 μm , less than 20 μm , less than 25 μm , less than 30 μm , less than 35 μm , less than 40 μm , less than 50 μm , less than

55 um, less than 60 um, less than 65 um, less than 70 um, less than 75 um, less than 80 um, less than 85 um, less than 90 um, less than 95 um, less than 100 um, less than 125 um, less than 150 um, less than 175 um, less than 200 um, less than 225 um, less than 250 um, less than 275 um, less than 300 um, less than 325 um, less than 350 um, less than 375 um, less than 400 um, less than 425 um, less than 450 um, less than 475 um, less than 500 um, less than 525 um, less than 550 um, less than 575 um, less than 600 um, less than 625 um, less than 650 um, less than 675 um, less than 700 um, less than 725 um, less than 750 um, less than 775 um, less than 800 um, less than 825 um, less than 850 um, less than 875 um, less than 900 um, less than 925 um, less than 950 um, less than 975 um, or less than 1000 um.

(399) In some embodiments, RNA (e.g., mRNA) vaccines may be delivered using smaller LNPs, which may comprise a diameter from about 1 nm to about 100 nm, from about 1 nm to about 10 nm, about 1 nm to about 20 nm, from about 1 nm to about 30 nm, from about 1 nm to about 40 nm, from about 1 nm to about 50 nm, from about 1 nm to about 60 nm, from about 1 nm to about 70 nm, from about 1 nm to about 80 nm, from about 1 nm to about 90 nm, from about 5 nm to about 100 nm, from about 5 nm to about 10 nm, about 5 nm to about 20 nm, from about 5 nm to about 30 nm, from about 5 nm to about 40 nm, from about 5 nm to about 50 nm, from about 5 nm to about 60 nm, from about 5 nm to about 70 nm, from about 5 nm to about 80 nm, from about 5 nm to about 90 nm, about 10 to about 50 nm, from about 20 to about 50 nm, from about 30 to about 50 nm, from about 40 to about 50 nm, from about 20 to about 60 nm, from about 30 to about 60 nm, from about 40 to about 60 nm, from about 20 to about 70 nm, from about 30 to about 70 nm, from about 40 to about 70 nm, from about 50 to about 70 nm, from about 60 to about 70 nm, from about 20 to about 80 nm, from about 30 to about 80 nm, from about 40 to about 80 nm, from about 50 to about 80 nm, from about 60 to about 80 nm, from about 20 to about 90 nm, from about 30 to about 90 nm, from about 40 to about 90 nm, from about 50 to about 90 nm, from about 60 to about 90 nm and/or from about 70 to about 90 nm.

(400) In some embodiments, such LNPs are synthesized using methods comprising microfluidic mixers. Examples of microfluidic mixers may include, but are not limited to, a slit interdigital micromixer including, but not limited to those manufactured by Microinnova (Allerheiligen bei Wildon, Austria) and/or a staggered herringbone micromixer (SHM) (Zhigaltsev, I. V. et al., Bottom-up design and synthesis of limit size lipid nanoparticle systems with aqueous and triglyceride cores using millisecond microfluidic mixing have been published (Langmuir. 2012. 28:3633-40; Belliveau, N. M. et al., Microfluidic synthesis of highly potent limit-size lipid nanoparticles for in vivo delivery of siRNA. Molecular Therapy-Nucleic Acids. 2012. 1:e37; Chen, D. et al., Rapid discovery of potent siRNA-containing lipid nanoparticles enabled by controlled microfluidic formulation. J Am Chem Soc. 2012. 134(16):6948-51, the contents of each of which are herein incorporated by reference in their entirety). In some embodiments, methods of LNP generation comprising SHM, further comprise the mixing of at least two input streams wherein mixing occurs by microstructure-induced chaotic advection (MICA). According to this method, fluid streams flow through channels present in a herringbone pattern causing rotational flow and folding the fluids around each other. This method may also comprise a surface for fluid mixing wherein the surface changes orientations during fluid cycling. Methods of generating LNPs using SHM include those disclosed in U.S. Application Publication Nos. 2004/0262223 and 2012/0276209, the contents of each of which are herein incorporated by reference in their entirety.

(401) In some embodiments, the RNA (e.g., mRNA) vaccine of the present disclosure may be formulated in lipid nanoparticles created using a micromixer such as, but not limited to, a Slit Interdigital Microstructured Mixer (SIMM-V2) or a Standard Slit Interdigital Micro Mixer (SSIMM) or Caterpillar (CPMM) or Impinging-jet (IJMM) from the Institut fir Mikrotechnik Mainz GmbH, Mainz Germany).

(402) In some embodiments, the RNA (e.g., mRNA) vaccines of the present disclosure may be formulated in lipid nanoparticles created using microfluidic technology (see, e.g., Whitesides,

George M. The Origins and the Future of Microfluidics. *Nature*, 2006 442: 368-373; and Abraham et al. Chaotic Mixer for Microchannels. *Science*, 2002 295: 647-651; each of which is herein incorporated by reference in its entirety). As a non-limiting example, controlled microfluidic formulation includes a passive method for mixing streams of steady pressure-driven flows in micro channels at a low Reynolds number (see, e.g., Abraham et al. Chaotic Mixer for Microchannels. *Science*, 2002 295: 647-651, the contents of which are herein incorporated by reference in their entirety).

(403) In some embodiments, the RNA (e.g., mRNA) vaccines of the present disclosure may be formulated in lipid nanoparticles created using a micromixer chip such as, but not limited to, those from Harvard Apparatus (Holliston, MA) or Dolomite Microfluidics (Royston, UK). A micromixer chip can be used for rapid mixing of two or more fluid streams with a split and recombine mechanism.

(404) In some embodiments, the RNA (e.g., mRNA) vaccines of the disclosure may be formulated for delivery using the drug encapsulating microspheres described in International Patent Publication No. WO2013063468 or U.S. Pat. No. 8,440,614, the contents of each of which are herein incorporated by reference in their entirety. The microspheres may comprise a compound of the formula (I), (II), (III), (IV), (V) or (VI) as described in International Patent Publication No. WO2013063468, the contents of which are herein incorporated by reference in their entirety. In some embodiments, the amino acid, peptide, polypeptide, lipids (APPL) are useful in delivering the RNA (e.g., mRNA) vaccines of the disclosure to cells (see International Patent Publication No. WO2013063468, the contents of which are herein incorporated by reference in their entirety).

(405) In some embodiments, the RNA (e.g., mRNA) vaccines of the disclosure may be formulated in lipid nanoparticles having a diameter from about 10 to about 100 nm such as, but not limited to, about 10 to about 20 nm, about 10 to about 30 nm, about 10 to about 40 nm, about 10 to about 50 nm, about 10 to about 60 nm, about 10 to about 70 nm, about 10 to about 80 nm, about 10 to about 90 nm, about 20 to about 30 nm, about 20 to about 40 nm, about 20 to about 50 nm, about 20 to about 60 nm, about 20 to about 70 nm, about 20 to about 80 nm, about 20 to about 90 nm, about 20 to about 100 nm, about 30 to about 40 nm, about 30 to about 50 nm, about 30 to about 60 nm, about 30 to about 70 nm, about 30 to about 80 nm, about 30 to about 90 nm, about 30 to about 100 nm, about 40 to about 50 nm, about 40 to about 60 nm, about 40 to about 70 nm, about 40 to about 80 nm, about 40 to about 90 nm, about 40 to about 100 nm, about 50 to about 60 nm, about 50 to about 70 nm, about 50 to about 80 nm, about 50 to about 90 nm, about 50 to about 100 nm, about 60 to about 70 nm, about 60 to about 80 nm, about 60 to about 90 nm, about 60 to about 100 nm, about 70 to about 80 nm, about 70 to about 90 nm, about 70 to about 100 nm, about 80 to about 90 nm, about 80 to about 100 nm and/or about 90 to about 100 nm.

(406) In some embodiments, the lipid nanoparticles may have a diameter from about 10 to 500 nm.

(407) In some embodiments, the lipid nanoparticle may have a diameter greater than 100 nm, greater than 150 nm, greater than 200 nm, greater than 250 nm, greater than 300 nm, greater than 350 nm, greater than 400 nm, greater than 450 nm, greater than 500 nm, greater than 550 nm, greater than 600 nm, greater than 650 nm, greater than 700 nm, greater than 750 nm, greater than 800 nm, greater than 850 nm, greater than 900 nm, greater than 950 nm or greater than 1000 nm.

(408) In some embodiments, the lipid nanoparticle may be a limit size lipid nanoparticle described in International Patent Publication No. WO2013059922, the contents of which are herein incorporated by reference in their entirety. The limit size lipid nanoparticle may comprise a lipid bilayer surrounding an aqueous core or a hydrophobic core; where the lipid bilayer may comprise a phospholipid such as, but not limited to, diacylphosphatidylcholine, a diacylphosphatidylethanolamine, a ceramide, a sphingomyelin, a dihydrosphingomyelin, a cephalin, a cerebroside, a C8-C20 fatty acid diacylphosphatidylcholine, and 1-palmitoyl-2-oleoyl phosphatidylcholine (POPC). In some embodiments, the limit size lipid nanoparticle may comprise a polyethylene glycol-lipid such as, but not limited to, DLPE-PEG, DMPE-PEG, DPPC-PEG and

DSPE-PEG.

(409) In some embodiments, the RNA (e.g., mRNA) vaccines may be delivered, localized and/or concentrated in a specific location using the delivery methods described in International Patent Publication No. WO2013063530, the contents of which are herein incorporated by reference in their entirety. As a non-limiting example, a subject may be administered an empty polymeric particle prior to, simultaneously with or after delivering the RNA (e.g., mRNA) vaccines to the subject. The empty polymeric particle undergoes a change in volume once in contact with the subject and becomes lodged, embedded, immobilized or entrapped at a specific location in the subject.

(410) In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in an active substance release system (see, e.g., U.S. Patent Publication No. US20130102545, the contents of which are herein incorporated by reference in their entirety). The active substance release system may comprise 1) at least one nanoparticle bonded to an oligonucleotide inhibitor strand which is hybridized with a catalytically active nucleic acid and 2) a compound bonded to at least one substrate molecule bonded to a therapeutically active substance (e.g., polynucleotides described herein), where the therapeutically active substance is released by the cleavage of the substrate molecule by the catalytically active nucleic acid.

(411) In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in a nanoparticle comprising an inner core comprising a non-cellular material and an outer surface comprising a cellular membrane. The cellular membrane may be derived from a cell or a membrane derived from a virus. As a non-limiting example, the nanoparticle may be made by the methods described in International Patent Publication No. WO2013052167, the contents of which are herein incorporated by reference in their entirety. As another non-limiting example, the nanoparticle described in International Patent Publication No. WO2013052167, the contents of which are herein incorporated by reference in their entirety, may be used to deliver the RNA (e.g., mRNA) vaccines described herein.

(412) In some embodiments, the RNA (e.g., mRNA) vaccines may be formulated in porous nanoparticle-supported lipid bilayers (protocells). Protocells are described in International Patent Publication No. WO2013056132, the contents of which are herein incorporated by reference in their entirety.

(413) In some embodiments, the RNA (e.g., mRNA) vaccines described herein may be formulated in polymeric nanoparticles as described in or made by the methods described in U.S. Pat. Nos. 8,420,123 and 8,518,963 and European Patent No. EP2073848B1, the contents of each of which are herein incorporated by reference in their entirety. As a non-limiting example, the polymeric nanoparticle may have a high glass transition temperature such as the nanoparticles described in or nanoparticles made by the methods described in U.S. Pat. No. 8,518,963, the contents of which are herein incorporated by reference in their entirety. As another non-limiting example, the polymer nanoparticle for oral and parenteral formulations may be made by the methods described in European Patent No. EP2073848B1, the contents of which are herein incorporated by reference in their entirety.

(414) In some embodiments, the RNA (e.g., mRNA) vaccines described herein may be formulated in nanoparticles used in imaging. The nanoparticles may be liposome nanoparticles such as those described in U.S. Patent Publication No US20130129636, herein incorporated by reference in its entirety. As a non-limiting example, the liposome may comprise gadolinium(III)2-{4,7-bis-carboxymethyl-10-[(N,N-distearylamidomethyl-N'-amido-methyl]-1,4,7,10-tetra-azacyclododec-1-yl}-acetic acid and a neutral, fully saturated phospholipid component (see, e.g., U.S. Patent Publication No US20130129636, the contents of which are herein incorporated by reference in their entirety).

(415) In some embodiments, the nanoparticles which may be used in the present disclosure are formed by the methods described in U.S. Patent Application No. US20130130348, the contents of

which are herein incorporated by reference in their entirety.

(416) The nanoparticles of the present disclosure may further include nutrients such as, but not limited to, those which deficiencies can lead to health hazards from anemia to neural tube defects (see, e.g., the nanoparticles described in International Patent Publication No WO2013072929, the contents of which are herein incorporated by reference in their entirety). As a non-limiting example, the nutrient may be iron in the form of ferrous, ferric salts or elemental iron, iodine, folic acid, vitamins or micronutrients.

(417) In some embodiments, the RNA (e.g., mRNA) vaccines of the present disclosure may be formulated in a swellable nanoparticle. The swellable nanoparticle may be, but is not limited to, those described in U.S. Pat. No. 8,440,231, the contents of which are herein incorporated by reference in their entirety. As a non-limiting embodiment, the swellable nanoparticle may be used for delivery of the RNA (e.g., mRNA) vaccines of the present disclosure to the pulmonary system (see, e.g., U.S. Pat. No. 8,440,231, the contents of which are herein incorporated by reference in their entirety).

(418) The RNA (e.g., mRNA) vaccines of the present disclosure may be formulated in polyanhydride nanoparticles such as, but not limited to, those described in U.S. Pat. No. 8,449,916, the contents of which are herein incorporated by reference in their entirety.

(419) The nanoparticles and microparticles of the present disclosure may be geometrically engineered to modulate macrophage and/or the immune response. In some embodiments, the geometrically engineered particles may have varied shapes, sizes and/or surface charges in order to incorporate the polynucleotides of the present disclosure for targeted delivery such as, but not limited to, pulmonary delivery (see, e.g., International Publication No WO2013082111, the contents of which are herein incorporated by reference in their entirety). Other physical features the geometrically engineering particles may have include, but are not limited to, fenestrations, angled arms, asymmetry and surface roughness, charge which can alter the interactions with cells and tissues. As a non-limiting example, nanoparticles of the present disclosure may be made by the methods described in International Publication No WO2013082111, the contents of which are herein incorporated by reference in their entirety.

(420) In some embodiments, the nanoparticles of the present disclosure may be water soluble nanoparticles such as, but not limited to, those described in International Publication No. WO2013090601, the contents of which are herein incorporated by reference in their entirety. The nanoparticles may be inorganic nanoparticles which have a compact and zwitterionic ligand in order to exhibit good water solubility. The nanoparticles may also have small hydrodynamic diameters (HD), stability with respect to time, pH, and salinity and a low level of non-specific protein binding.

(421) In some embodiments the nanoparticles of the present disclosure may be developed by the methods described in U.S. Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in their entirety.

(422) In some embodiments, the nanoparticles of the present disclosure are stealth nanoparticles or target-specific stealth nanoparticles such as, but not limited to, those described in U.S. Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in their entirety. The nanoparticles of the present disclosure may be made by the methods described in U.S. Patent Publication No. US20130172406, the contents of which are herein incorporated by reference in their entirety.

(423) In some embodiments, the stealth or target-specific stealth nanoparticles may comprise a polymeric matrix. The polymeric matrix may comprise two or more polymers such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polyesters,

polyanhydrides, polyethers, polyurethanes, polymethacrylates, polyacrylates, polycyanoacrylates or combinations thereof.

(424) In some embodiments, the nanoparticle may be a nanoparticle-nucleic acid hybrid structure having a high density nucleic acid layer. As a non-limiting example, the nanoparticle-nucleic acid hybrid structure may be made by the methods described in U.S. Patent Publication No.

US20130171646, the contents of which are herein incorporated by reference in their entirety. The nanoparticle may comprise a nucleic acid such as, but not limited to, polynucleotides described herein and/or known in the art.

(425) At least one of the nanoparticles of the present disclosure may be embedded in the core a nanostructure or coated with a low density porous 3-D structure or coating which is capable of carrying or associating with at least one payload within or on the surface of the nanostructure. Non-limiting examples of the nanostructures comprising at least one nanoparticle are described in International Patent Publication No. WO2013123523, the contents of which are herein incorporated by reference in their entirety.

(426) In some embodiments the RNA (e.g., mRNA) vaccine may be associated with a cationic or polycationic compounds, including protamine, nucleoline, spermine or spermidine, or other cationic peptides or proteins, such as poly-L-lysine (PLL), polyarginine, basic polypeptides, cell penetrating peptides (CPPs), including HIV-binding peptides, HIV-1 Tat (HIV), Tat-derived peptides, Penetratin, VP.sup.22 derived or analog peptides, Pestivirus Erns, HSV, VP.sup.22 (Herpes simplex), MAP, KALA or protein transduction domains (PTDs), PpT620, prolin-rich peptides, arginine-rich peptides, lysine-rich peptides, MPG-peptide(s), Pep-1, L-oligomers, Calcitonin peptide(s), Antennapedia-derived peptides (particularly from *Drosophila* antennapedia), pAntp, pIsl, FGF, Lactoferrin, Transportan, Buforin-2, Bac715-24, SynB, SynB(1), pVEC, hCT-derived peptides, SAP, histones, cationic polysaccharides, for example chitosan, polybrene, cationic polymers, e.g. polyethyleneimine (PEI), cationic lipids, e.g. DOTMA: [1-(2,3-sioleyloxy)propyl]-N,N,N-trimethylammonium chloride, DMRIE, di-C14-amidine, DOTIM, SAINT, DC-Chol, BGTC, CTAP, DOPC, DODAP, DOPE: Dioleoyl phosphatidylethanol-amine, DOSPA, DODAB, DOIC, DMEPC, DOGS: Dioctadecylamidoglycylspermin, DIMRI: Dimyristooxypropyl dimethyl hydroxyethyl ammonium bromide, DOTAP: dioleoyloxy-3-(trimethylammonio)propane, DC-6-14: O,O-ditetradecanoyl-N—. alpha.-trimethylammonioacetyl)diethanolamine chloride, CLIP1: rac-[(2,3-dioctadecyloxypropyl)(2-hydroxyethyl)]-dimethylammonium chloride, CLIP6: rac-[2(2,3-dihexadecyloxypropyloxymethyloxy)ethyl]-trimethylammonium, CLIP9: rac-[2(2,3-dihexadecyloxypropyloxysuccinyloxy)ethyl]-trimethylammonium, oligofectamine, or cationic or polycationic polymers, e.g. modified polyaminoacids, such as beta-aminoacid-polymers or reversed polyamides, etc., modified polyethylenes, such as PVP (poly(N-ethyl-4-vinylpyridinium bromide)), etc., modified acrylates, such as pDMAEMA (poly(dimethylaminoethyl methylacrylate)), etc., modified amidoamines such as pAMAM (poly(amidoamine)), etc., modified polybetaminoester (PBAE), such as diamine end modified 1,4 butanediol diacrylate-co-5-amino-1-pentanol polymers, etc., dendrimers, such as polypropylamine dendrimers or pAMAM based dendrimers, etc., polyimine(s), such as PEI: poly(ethyleneimine), poly(propyleneimine), etc., polyallylamine, sugar backbone based polymers, such as cyclodextrin based polymers, dextran based polymers, chitosan, etc., silan backbone based polymers, such as PMOXA-PDMS copolymers, etc., blockpolymers consisting of a combination of one or more cationic blocks (e.g. selected from a cationic polymer as mentioned above) and of one or more hydrophilic or hydrophobic blocks (e.g. polyethyleneglycole), etc.

(427) In other embodiments the RNA (e.g., mRNA) vaccine is not associated with a cationic or polycationic compounds.

(428) In some embodiments, a nanoparticle comprises compounds of Formula (I):

(429) ##STR00003## or a salt or isomer thereof, wherein: R.sub.1 is selected from the group consisting of C.sub.5-30 alkyl, C.sub.5-20 alkenyl, —R*YR'', —YR'', and —R''M'R'; R.sub.2 and

R.sub.3 are independently selected from the group consisting of H, C.sub.1-14 alkyl, C.sub.2-14 alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R.sub.2 and R.sub.3, together with the atom to which they are attached, form a heterocycle or carbocycle; R.sub.4 is selected from the group consisting of a C.sub.3-6 carbocycle, $-(CH.sub.2).sub.nQ$, $-(CH.sub.2).sub.nCHQR$, $CHQR$, $-CQ(R).sub.2$, and unsubstituted C.sub.1-6 alkyl, where Q is selected from a carbocycle, heterocycle, $-OR$, $-O(CH.sub.2).sub.nN(R).sub.2$, $-C(O)OR$, $-OC(O)R$, $-CX.sub.3$, $-CX.sub.2H$, $-CXH.sub.2$, $-CN$, $-N(R).sub.2$, $-C(O)N(R).sub.2$, $-N(R)C(O)R$, $-N(R)S(O).sub.2R$, $-N(R)C(O)N(R).sub.2$, $-N(R)C(S)N(R).sub.2$, $-N(R)R.sub.8$, $-O(CH.sub.2).sub.nOR$, $-N(R)C(=NR.sub.9)N(R).sub.2$, $-N(R)C(=CHR.sub.9)N(R).sub.2$, $-OC(O)N(R).sub.2$, $-N(R)C(O)OR$, $-N(OR)C(O)R$, $-N(OR)S(O).sub.2R$, $-N(OR)C(O)OR$, $-N(OR)C(O)N(R).sub.2$, $-N(OR)C(S)N(R).sub.2$, $-N(OR)C(=NR.sub.9)N(R).sub.2$, $-N(OR)C(=CHR.sub.9)N(R).sub.2$, $-C(=NR.sub.9)N(R).sub.2$, $-C(=NR.sub.9)R$, $-C(O)N(R)OR$, and $-C(R)N(R).sub.2C(O)OR$, and each n is independently selected from 1, 2, 3, 4, and 5; each R.sub.5 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R.sub.6 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O).sub.2-$, $-S-S-$, an aryl group, and a heteroaryl group; R.sub.7 is selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; R.sub.8 is selected from the group consisting of C.sub.3-6 carbocycle and heterocycle; R.sub.9 is selected from the group consisting of H, CN, $NO.sub.2$, C.sub.1-6 alkyl, $-OR$, $-S(O).sub.2R$, $-S(O).sub.2N(R).sub.2$, C.sub.2-6 alkenyl, C.sub.3-6 carbocycle and heterocycle; each R is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R' is independently selected from the group consisting of C.sub.1-18 alkyl, C.sub.2-18 alkenyl, $-R^*YR''$, $-YR''$, and H; each R'' is independently selected from the group consisting of C.sub.3-14 alkyl and C.sub.3-14 alkenyl; each R* is independently selected from the group consisting of C.sub.1-12 alkyl and C.sub.2-12 alkenyl; each Y is independently a C.sub.3-6 carbocycle; each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13.

(430) In some embodiments, a subset of compounds of Formula (I) includes those in which when R.sub.4 is $-(CH.sub.2).sub.nQ$, $-(CH.sub.2).sub.nCHQR$, $-CHQR$, or $-CQ(R).sub.2$, then (i) Q is not-N(R).sub.2 when n is 1, 2, 3, 4 or 5, or (ii) Q is not 5, 6, or 7-membered heterocycloalkyl when n is 1 or 2.

(431) In some embodiments, another subset of compounds of Formula (I) includes those in which R.sub.1 is selected from the group consisting of C.sub.5-30 alkyl, C.sub.5-20 alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$; R.sub.2 and R.sub.3 are independently selected from the group consisting of H, C.sub.1-14 alkyl, C.sub.2-14 alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R.sub.2 and R.sub.3, together with the atom to which they are attached, form a heterocycle or carbocycle; R.sub.4 is selected from the group consisting of a C.sub.3-6 carbocycle, $-(CH.sub.2).sub.nQ$, $-(CH.sub.2).sub.nCHQR$, $-CHQR$, $-CQ(R).sub.2$, and unsubstituted C1.6 alkyl, where Q is selected from a C.sub.3-6 carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, $-OR$, $-O(CH.sub.2).sub.nN(R).sub.2$, $-C(O)OR$, $-OC(O)R$, $-CX.sub.3$, $-CX.sub.2H$, $-CXH.sub.2$, $-CN$, $-C(O)N(R).sub.2$, $-N(R)C(O)R$, $-N(R)S(O).sub.2R$, $-N(R)C(O)N(R).sub.2$, $-N(R)C(S)N(R).sub.2$, $-CRN(R).sub.2C(O)OR$, $-N(R)R.sub.8$, $-O(CH.sub.2).sub.nOR$, $-N(R)C(=NR.sub.9)N(R).sub.2$, $-N(R)C(=CHR.sub.9)N(R).sub.2$, $-OC(O)N(R).sub.2$, $-N(R)C(O)OR$, $-N(OR)C(O)R$, $-N(OR)S(O).sub.2R$, $-N(OR)C(O)OR$, $-N(OR)C(O)N(R).sub.2$, $-N(OR)C(S)N(R).sub.2$, $-N(OR)C(=NR.sub.9)N(R).sub.2$, $-N(OR)C(=CHR.sub.9)N(R).sub.2$, $-C(=NR.sub.9)N(R).sub.2$, $-C(=NR.sub.9)R$, $-C(O)N(R)OR$, and a 5- to 14-membered heterocycloalkyl having one or more heteroatoms selected from N, O, and S which is substituted

with one or more substituents selected from oxo ($=O$), OH, amino, mono- or di-alkylamino, and C.sub.1-3 alkyl, and each n is independently selected from 1, 2, 3, 4, and 5; each R.sub.5 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R.sub.6 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O).sub.2-$, $-S-S-$, an aryl group, and a heteroaryl group; R.sub.7 is selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; R.sub.8 is selected from the group consisting of C.sub.3-6 carbocycle and heterocycle; R.sub.9 is selected from the group consisting of H, CN, NO.sub.2, C.sub.1-6 alkyl, $-OR$, $-S(O).sub.2R$, $-S(O).sub.2N(R).sub.2$, C.sub.2-6 alkenyl, C.sub.3-6 carbocycle and heterocycle; each R is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R' is independently selected from the group consisting of C.sub.1-8 alkyl, C.sub.2-18 alkenyl, $-R^*YR''$, $-YR''$, and H; each R'' is independently selected from the group consisting of C.sub.3-14 alkyl and C.sub.3-14 alkenyl; each R* is independently selected from the group consisting of C.sub.1-12 alkyl and C.sub.2-12 alkenyl; each Y is independently a C.sub.3-6 carbocycle; each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13, or salts or isomers thereof.

(432) In some embodiments, another subset of compounds of Formula (I) includes those in which R.sub.1 is selected from the group consisting of C.sub.5-30 alkyl, C.sub.5-20 alkenyl, $-R^*YR''$, $-YR''$, and $-R''M'R'$; R.sub.2 and R.sub.3 are independently selected from the group consisting of H, C1.14 alkyl, C.sub.2-14 alkenyl, $-R^*YR''$, $-YR''$, and $-R^*OR''$, or R.sub.2 and R.sub.3, together with the atom to which they are attached, form a heterocycle or carbocycle; R.sub.4 is selected from the group consisting of a C.sub.3-6 carbocycle, $-(CH.sub.2).sub.nQ$, $-(CH.sub.2).sub.nCHQR$, $-CHQR$, $-CQ(R).sub.2$, and unsubstituted C.sub.1-6 alkyl, where Q is selected from a C.sub.3-6 carbocycle, a 5- to 14-membered heterocycle having one or more heteroatoms selected from N, O, and S, $-OR$, $-O(CH.sub.2).sub.nN(R).sub.2$, $-C(O)OR$, $-OC(O)R$, $-CX.sub.3$, $-CX.sub.2H$, $-CXH.sub.2$, $-CN$, $-C(O)N(R).sub.2$, $-N(R)C(O)R$, $-N(R)S(O).sub.2R$, $-N(R)C(O)N(R).sub.2$, $-N(R)C(S)N(R).sub.2$, $-CRN(R).sub.2C(O)OR$, $-N(R)R.sub.8$, $-O(CH.sub.2).sub.nOR$, $-N(R)C(=NR.sub.9)N(R).sub.2$, $-N(R)C(=CHR.sub.9)N(R).sub.2$, $-OC(O)N(R).sub.2$, $-N(R)C(O)OR$, $-N(OR)C(O)R$, $-N(OR)S(O).sub.2R$, $-N(OR)C(O)OR$, $-N(OR)C(O)N(R).sub.2$, $-N(OR)C(S)N(R).sub.2$, $-N(OR)C(=NR.sub.9)N(R).sub.2$, $-N(OR)C(=CHR.sub.9)N(R).sub.2$, $-C(=NR.sub.9)R$, $-C(O)N(R)OR$, and $-C(=NR.sub.9)N(R).sub.2$, and each n is independently selected from 1, 2, 3, 4, and 5; and when Q is a 5- to 14-membered heterocycle and (i) R.sub.4 is $-(CH.sub.2).sub.nQ$ in which n is 1 or 2, or (ii) R.sub.4 is $-(CH.sub.2).sub.nCHQR$ in which n is 1, or (iii) R.sub.4 is $-CHQR$, and $-CQ(R).sub.2$, then Q is either a 5- to 14-membered heteroaryl or 8- to 14-membered heterocycloalkyl; each R.sub.5 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R.sub.6 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; M and M' are independently selected from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O).sub.2-$, $-S-S-$, an aryl group, and a heteroaryl group; R.sub.7 is selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; R.sub.8 is selected from the group consisting of C.sub.3-6 carbocycle and heterocycle; R.sub.9 is selected from the group consisting of H, CN, NO.sub.2, C.sub.1-6 alkyl, $-OR$, $-S(O).sub.2R$, $-S(O).sub.2N(R).sub.2$, C.sub.2-6 alkenyl, C.sub.3-6 carbocycle and heterocycle; each R is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R' is independently selected from the group consisting of C.sub.1-8 alkyl, C.sub.2-18 alkenyl, $-R^*YR''$, $-YR''$, and H; each R'' is independently selected from the group consisting of C.sub.3-14 alkyl and C.sub.3-14 alkenyl; each R* is independently selected from the group consisting of

C.sub.1-12 alkyl and C.sub.2-12 alkenyl; each Y is independently a C.sub.3-6 carbocycle; each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13, or salts or isomers thereof.

(433) In some embodiments, another subset of compounds of Formula (I) includes those in which R.sub.1 is selected from the group consisting of C.sub.5-30 alkyl, C.sub.5-20 alkenyl, —R*YR'', —YR'', and —R''M'R'; R.sub.2 and R.sub.3 are independently selected from the group consisting of H, C.sub.1-14 alkyl, C.sub.2-14 alkenyl, —R*YR'', —YR'', and —R*OR'', or R.sub.2 and R.sub.3, together with the atom to which they are attached, form a heterocycle or carbocycle; R.sub.4 is selected from the group consisting of a C.sub.3-6 carbocycle, —(CH.sub.2).sub.nQ, —(CH.sub.2).sub.nCHQR, —CHQR, —CQ(R).sub.2, and unsubstituted C.sub.1-6 alkyl, where Q is selected from a C.sub.3-6 carbocycle, a 5- to 14-membered heteroaryl having one or more heteroatoms selected from N, O, and S, —OR, —O(CH.sub.2).sub.nN(R).sub.2, —C(O)OR, —OC(O)R, —CX.sub.3, —CX.sub.2H, —CXH.sub.2, —CN, —C(O)N(R).sub.2, —N(R)C(O)R, —N(R)S(O).sub.2R, —N(R)C(O)N(R).sub.2, —N(R)C(S)N(R).sub.2, —CRN(R).sub.2C(O)OR, —N(R)R.sub.8, —O(CH.sub.2).sub.nOR, —N(R)C(=NR.sub.9)N(R).sub.2, —N(R)C(=CHR.sub.9)N(R).sub.2, —OC(O)N(R).sub.2, —N(R)C(O)OR, —N(OR)C(O)R, —N(OR)S(O).sub.2R, —N(OR)C(O)OR, —N(OR)C(O)N(R).sub.2, —N(OR)C(S)N(R).sub.2, —N(OR)C(=NR.sub.9)N(R).sub.2, —N(OR)C(=CHR.sub.9)N(R).sub.2, —C(=NR.sub.9)R, —C(O)N(R)OR, and —C(=NR.sub.9)N(R).sub.2, and each n is independently selected from 1, 2, 3, 4, and 5; each R.sub.5 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R.sub.6 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; M and M' are independently selected from —C(O)O—, —OC(O)—, —C(O)N(R')—, —N(R')C(O)—, —C(O)—, —C(S)—, —C(S)S—, —SC(S)—, —CH(OH)—, —P(O)(OR')O—, —S(O).sub.2—, —S—S—, an aryl group, and a heteroaryl group; R.sub.7 is selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; R.sub.8 is selected from the group consisting of C.sub.3-6 carbocycle and heterocycle; R.sub.9 is selected from the group consisting of H, CN, NO.sub.2, C.sub.1-6 alkyl, —OR, —S(O).sub.2R, —S(O).sub.2N(R).sub.2, C.sub.2-6 alkenyl, C.sub.3-6 carbocycle and heterocycle; each R is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R' is independently selected from the group consisting of C.sub.1-8 alkyl, C.sub.2-18 alkenyl, —R*YR'', —YR'', and H; each R'' is independently selected from the group consisting of C.sub.3-14 alkyl and C.sub.3-14 alkenyl; each R* is independently selected from the group consisting of C.sub.1-12 alkyl and C.sub.2-12 alkenyl; each Y is independently a C.sub.3-6 carbocycle; each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13, or salts or isomers thereof.

(434) In some embodiments, another subset of compounds of Formula (I) includes those in which R.sub.1 is selected from the group consisting of C.sub.5-30 alkyl, C.sub.5-20 alkenyl, —R*YR'', —YR'', and —R''M'R'; R.sub.2 and R.sub.3 are independently selected from the group consisting of H, C.sub.2-14 alkyl, C.sub.2-14 alkenyl, —R*YR'', —YR'', and —R*OR'', or R.sub.2 and R.sub.3, together with the atom to which they are attached, form a heterocycle or carbocycle; R.sub.4 is —(CH.sub.2).sub.nQ or —(CH.sub.2).sub.nCHQR, where Q is —N(R).sub.2, and n is selected from 3, 4, and 5; each R.sub.5 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R.sub.6 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; M and M' are independently selected from —C(O)O—, —OC(O)—, —C(O)N(R')—, —N(R')C(O)—, —C(O)—, —C(S)—, —C(S)S—, —SC(S)—, —CH(OH)—, —P(O)(OR')O—, —S(O).sub.2—, —S—S—, an aryl group, and a heteroaryl group; R.sub.7 is selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R' is independently selected from the group consisting of C.sub.1-8 alkyl, C.sub.2-18 alkenyl, —R*YR'', —YR'', and H; each R'' is independently selected from the

group consisting of C.sub.3-14 alkyl and C.sub.3-14 alkenyl; each R* is independently selected from the group consisting of C.sub.1-12 alkyl and C.sub.1-12 alkenyl; each Y is independently a C.sub.3-6 carbocycle; each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13, or salts or isomers thereof.

(435) In some embodiments, another subset of compounds of Formula (I) includes those in which R.sub.1 is selected from the group consisting of C.sub.5-30 alkyl, C.sub.5-20 alkenyl, —R*YR'', —YR'', and —R''M'R'; R.sub.2 and R.sub.3 are independently selected from the group consisting of C.sub.1-14 alkyl, C.sub.2-14 alkenyl, —R*YR'', —YR'', and —R*OR'', or R.sub.2 and R.sub.3, together with the atom to which they are attached, form a heterocycle or carbocycle; R.sub.4 is selected from the group consisting of —(CH.sub.2).sub.nQ, —(CH.sub.2).sub.nCHQR, —CHQR, and —CQ(R).sub.2, where Q is —N(R).sub.2, and n is selected from 1, 2, 3, 4, and 5; each R.sub.5 is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R.sub.6 is independently selected from the group consisting of C1.3 alkyl, C.sub.2-3 alkenyl, and H; M and M' are independently selected from —C(O)O—, —OC(O)—, —C(O)N(R')—, —N(R')C(O)—, —C(O)—, —C(S)—, —C(S)S—, —SC(S)—, —CH(OH)—, —P(O)(OR')O—, —S(O).sub.2—, —S—S—, an aryl group, and a heteroaryl group; R.sub.7 is selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R is independently selected from the group consisting of C.sub.1-3 alkyl, C.sub.2-3 alkenyl, and H; each R' is independently selected from the group consisting of C18 alkyl, C.sub.2-18 alkenyl, —R*YR'', —YR'', and H; each R'' is independently selected from the group consisting of C.sub.3-14 alkyl and C.sub.3-14 alkenyl; each R* is independently selected from the group consisting of C.sub.1-12 alkyl and C.sub.1-12 alkenyl; each Y is independently a C.sub.3-6 carbocycle; each X is independently selected from the group consisting of F, Cl, Br, and I; and m is selected from 5, 6, 7, 8, 9, 10, 11, 12, and 13, or salts or isomers thereof.

(436) In some embodiments, a subset of compounds of Formula (I) includes those of Formula (IA):

(437) ##STR00004## or a salt or isomer thereof, wherein 1 is selected from 1, 2, 3, 4, and 5; m is selected from 5, 6, 7, 8, and 9; M1 is a bond or M'; R.sub.4 is unsubstituted C.sub.1-3 alkyl, or —(CH.sub.2).sub.nQ, in which Q is OH, —NHC(S)N(R).sub.2, —NHC(O)N(R).sub.2, —N(R)C(O)R, —N(R)S(O).sub.2R, —N(R)R.sub.8, —NHC(=NR.sub.9)N(R).sub.2, —NHC(=CHR.sub.9)N(R).sub.2, —OC(O)N(R).sub.2, —N(R)C(O)OR, heteroaryl or heterocycloalkyl; M and M' are independently selected from —C(O)O—, —OC(O)—, —C(O)N(R')—, —P(O)(OR')O—, —S—S—, an aryl group, and a heteroaryl group; and R.sub.2 and R.sub.3 are independently selected from the group consisting of H, C.sub.1-14 alkyl, and C.sub.2-14 alkenyl.

(438) In some embodiments, a subset of compounds of Formula (I) includes those of Formula (II):

(439) ##STR00005##

or a salt or isomer thereof, wherein 1 is selected from 1, 2, 3, 4, and 5; M.sub.1 is a bond or M'; R.sub.4 is unsubstituted C.sub.1-3 alkyl, or —(CH.sub.2).sub.nQ, in which n is 2, 3, or 4, and Q is OH, —NHC(S)N(R).sub.2, —NHC(O)N(R).sub.2, —N(R)C(O)R, —N(R)S(O).sub.2R, —N(R)R.sub.8, —NHC(=NR.sub.9)N(R).sub.2, —NHC(=CHR.sub.9)N(R).sub.2, —OC(O)N(R).sub.2, —N(R)C(O)OR, heteroaryl or heterocycloalkyl; M and M' are independently selected from —C(O)O—, —OC(O)—, —C(O)N(R')—, —P(O)(OR')O—, —S—S—, an aryl group, and a heteroaryl group; and R.sub.2 and R.sub.3 are independently selected from the group consisting of H, C.sub.1-14 alkyl, and C.sub.2-14 alkenyl.

(440) In some embodiments, a subset of compounds of Formula (I) includes those of Formula (IIa), (IIb), (IIc), or (IIe):

(441) ##STR00006## or a salt or isomer thereof, wherein R.sub.4 is as described herein.

(442) In some embodiments, a subset of compounds of Formula (I) includes those of Formula (IId):

(443) ##STR00007## or a salt or isomer thereof, wherein n is 2, 3, or 4; and m, R', R'', and R.sub.2 through R.sub.6 are as described herein. For example, each of R.sub.2 and R.sub.3 may be

independently selected from the group consisting of C.sub.5-14 alkyl and C.sub.5-14 alkenyl.

(444) In some embodiments, a subset of compounds of Formula (I) includes those of Formula (IIa), (IIb), (IIc), or (IIe):

(445) ##STR00008## or a salt or isomer thereof, wherein R.sub.4 is as described herein.

(446) In some embodiments, a subset of compounds of Formula (I) includes those of Formula (IIc):

(447) ##STR00009## or a salt or isomer thereof, wherein n is 2, 3, or 4; and m, R', R'', and R.sub.2 through R.sub.6 are as described herein. For example, each of R.sub.2 and R.sub.3 may be independently selected from the group consisting of C.sub.5-14 alkyl and C.sub.5-14 alkenyl.

(448) In some embodiments, the compound of Formula (I) is selected from the group consisting of:

(449) ##STR00010## ##STR00011## ##STR00012## ##STR00013## ##STR00014##
##STR00015## ##STR00016## ##STR00017## ##STR00018##

(450) In further embodiments, the compound of Formula (I) is selected from the group consisting of:

(451) ##STR00019##

(452) In some embodiments, the compound of Formula (I) is selected from the group consisting of:

(453) ##STR00020## ##STR00021## ##STR00022## ##STR00023## ##STR00024##
##STR00025## ##STR00026## ##STR00027## ##STR00028## ##STR00029## ##STR00030##
##STR00031## ##STR00032## ##STR00033## ##STR00034## ##STR00035## ##STR00036##
##STR00037## ##STR00038## ##STR00039## ##STR00040## ##STR00041## ##STR00042##
##STR00043## ##STR00044## ##STR00045## ##STR00046## ##STR00047##

and salts and isomers thereof.

(454) In some embodiments, a nanoparticle comprises the following compound:

(455) ##STR00048##

or salts and isomers thereof.

(456) In some embodiments, the disclosure features a nanoparticle composition including a lipid component comprising a compound as described herein (e.g., a compound according to Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IIe), (IIe) or (IIe)).

(457) In some embodiments, the disclosure features a pharmaceutical composition comprising a nanoparticle composition according to the preceding embodiments and a pharmaceutically acceptable carrier. For example, the pharmaceutical composition is refrigerated or frozen for storage and/or shipment (e.g., being stored at a temperature of 4° C. or lower, such as a temperature between about -150° C. and about 0° C. or between about -80° C. and about -20° C. (e.g., about -5° C., -10° C., -15° C., -20° C., -25° C., -30° C., -40° C., -50° C., -60° C., -70° C., -80° C., -90° C., -130° C. or -150° C.). For example, the pharmaceutical composition is a solution that is refrigerated for storage and/or shipment at, for example, about -20° C., -30° C., -40° C., -50° C., -60° C., -70° C., or -80° C.

(458) In some embodiments, the disclosure provides a method of delivering a therapeutic and/or prophylactic (e.g., RNA, such as mRNA) to a cell (e.g., a mammalian cell). This method includes the step of administering to a subject (e.g., a mammal, such as a human) a nanoparticle composition including (i) a lipid component including a phospholipid (such as a polyunsaturated lipid), a PEG lipid, a structural lipid, and a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IIe) or (IIe) and (ii) a therapeutic and/or prophylactic, in which administering involves contacting the cell with the nanoparticle composition, whereby the therapeutic and/or prophylactic is delivered to the cell.

(459) In some embodiments, the disclosure provides a method of producing a polypeptide of interest in a cell (e.g., a mammalian cell). The method includes the step of contacting the cell with a nanoparticle composition including (i) a lipid component including a phospholipid (such as a polyunsaturated lipid), a PEG lipid, a structural lipid, and a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IIe) or (IIe) and (ii) an mRNA encoding the polypeptide of interest, whereby the mRNA is capable of being translated in the cell to produce the polypeptide.

(460) In some embodiments, the disclosure provides a method of treating a disease or disorder in a

mammal (e.g., a human) in need thereof. The method includes the step of administering to the mammal a therapeutically effective amount of a nanoparticle composition including (i) a lipid component including a phospholipid (such as a polyunsaturated lipid), a PEG lipid, a structural lipid, and a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIE) and (ii) a therapeutic and/or prophylactic (e.g., an mRNA). In some embodiments, the disease or disorder is characterized by dysfunctional or aberrant protein or polypeptide activity. For example, the disease or disorder is selected from the group consisting of rare diseases, infectious diseases, cancer and proliferative diseases, genetic diseases (e.g., cystic fibrosis), autoimmune diseases, diabetes, neurodegenerative diseases, cardio- and reno-vascular diseases, and metabolic diseases.

(461) In some embodiments, the disclosure provides a method of delivering (e.g., specifically delivering) a therapeutic and/or prophylactic to a mammalian organ (e.g., a liver, spleen, lung, or femur). This method includes the step of administering to a subject (e.g., a mammal) a nanoparticle composition including (i) a lipid component including a phospholipid, a PEG lipid, a structural lipid, and a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIE) and (ii) a therapeutic and/or prophylactic (e.g., an mRNA), in which administering involves contacting the cell with the nanoparticle composition, whereby the therapeutic and/or prophylactic is delivered to the target organ (e.g., a liver, spleen, lung, or femur).

(462) In some embodiments, the disclosure features a method for the enhanced delivery of a therapeutic and/or prophylactic (e.g., an mRNA) to a target tissue (e.g., a liver, spleen, lung, or femur). This method includes administering to a subject (e.g., a mammal) a nanoparticle composition, the composition including (i) a lipid component including a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIE), a phospholipid, a structural lipid, and a PEG lipid; and (ii) a therapeutic and/or prophylactic, the administering including contacting the target tissue with the nanoparticle composition, whereby the therapeutic and/or prophylactic is delivered to the target tissue.

(463) In some embodiments, the disclosure features a method of lowering immunogenicity comprising introducing the nanoparticle composition of the disclosure into cells, wherein the nanoparticle composition reduces the induction of the cellular immune response of the cells to the nanoparticle composition, as compared to the induction of the cellular immune response in cells induced by a reference composition which comprises a reference lipid instead of a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIE). For example, the cellular immune response is an innate immune response, an adaptive immune response, or both.

(464) The disclosure also includes methods of synthesizing a compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIE) and methods of making a nanoparticle composition including a lipid component comprising the compound of Formula (I), (IA), (II), (IIa), (IIb), (IIc), (IId) or (IIE).

(465) Modes of Vaccine Administration

(466) Respiratory virus RNA (e.g. mRNA) vaccines may be administered by any route which results in a therapeutically effective outcome. These include, but are not limited, to intradermal, intramuscular, and/or subcutaneous administration. The present disclosure provides methods comprising administering RNA (e.g., mRNA) vaccines to a subject in need thereof. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease, the particular composition, its mode of administration, its mode of activity, and the like. Respiratory virus RNA (e.g., mRNA) vaccines compositions are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of RNA (e.g., mRNA) vaccine compositions may be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective, prophylactically effective, or appropriate imaging dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex

and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts. (467) In some embodiments, respiratory virus RNA (e.g., mRNA) vaccine compositions may be administered at dosage levels sufficient to deliver 0.0001 mg/kg to 100 mg/kg, 0.001 mg/kg to 0.05 mg/kg, 0.005 mg/kg to 0.05 mg/kg, 0.001 mg/kg to 0.005 mg/kg, 0.05 mg/kg to 0.5 mg/kg, 0.01 mg/kg to 50 mg/kg, 0.1 mg/kg to 40 mg/kg, 0.5 mg/kg to 30 mg/kg, 0.01 mg/kg to 10 mg/kg, 0.1 mg/kg to 10 mg/kg, or 1 mg/kg to 25 mg/kg, of subject body weight per day, one or more times a day, per week, per month, etc. to obtain the desired therapeutic, diagnostic, prophylactic, or imaging effect (see, e.g., the range of unit doses described in International Publication No WO2013078199, the contents of which are herein incorporated by reference in their entirety). The desired dosage may be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, every four weeks, every 2 months, every three months, every 6 months, etc. In some embodiments, the desired dosage may be delivered using multiple administrations (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations). When multiple administrations are employed, split dosing regimens such as those described herein may be used. In exemplary embodiments, respiratory virus RNA (e.g., mRNA) vaccine compositions may be administered at dosage levels sufficient to deliver 0.0005 mg/kg to 0.01 mg/kg, e.g., about 0.0005 mg/kg to about 0.0075 mg/kg, e.g., about 0.0005 mg/kg, about 0.001 mg/kg, about 0.002 mg/kg, about 0.003 mg/kg, about 0.004 mg/kg or about 0.005 mg/kg.

(468) In some embodiments, respiratory virus RNA (e.g., mRNA) vaccine compositions may be administered once or twice (or more) at dosage levels sufficient to deliver 0.025 mg/kg to 0.250 mg/kg, 0.025 mg/kg to 0.500 mg/kg, 0.025 mg/kg to 0.750 mg/kg, or 0.025 mg/kg to 1.0 mg/kg.

(469) In some embodiments, respiratory virus RNA (e.g., mRNA) vaccine compositions may be administered twice (e.g., Day 0 and Day 7, Day 0 and Day 14, Day 0 and Day 21, Day 0 and Day 28, Day 0 and Day 60, Day 0 and Day 90, Day 0 and Day 120, Day 0 and Day 150, Day 0 and Day 180, Day 0 and 3 months later, Day 0 and 6 months later, Day 0 and 9 months later, Day 0 and 12 months later, Day 0 and 18 months later, Day 0 and 2 years later, Day 0 and 5 years later, or Day 0 and 10 years later) at a total dose of or at dosage levels sufficient to deliver a total dose of 0.0100 mg, 0.025 mg, 0.050 mg, 0.075 mg, 0.100 mg, 0.125 mg, 0.150 mg, 0.175 mg, 0.200 mg, 0.225 mg, 0.250 mg, 0.275 mg, 0.300 mg, 0.325 mg, 0.350 mg, 0.375 mg, 0.400 mg, 0.425 mg, 0.450 mg, 0.475 mg, 0.500 mg, 0.525 mg, 0.550 mg, 0.575 mg, 0.600 mg, 0.625 mg, 0.650 mg, 0.675 mg, 0.700 mg, 0.725 mg, 0.750 mg, 0.775 mg, 0.800 mg, 0.825 mg, 0.850 mg, 0.875 mg, 0.900 mg, 0.925 mg, 0.950 mg, 0.975 mg, or 1.0 mg. Higher and lower dosages and frequency of administration are encompassed by the present disclosure. For example, a respiratory virus RNA (e.g., mRNA) vaccine composition may be administered three or four times.

(470) In some embodiments, respiratory virus RNA (e.g., mRNA) vaccine compositions may be administered twice (e.g., Day 0 and Day 7, Day 0 and Day 14, Day 0 and Day 21, Day 0 and Day 28, Day 0 and Day 60, Day 0 and Day 90, Day 0 and Day 120, Day 0 and Day 150, Day 0 and Day 180, Day 0 and 3 months later, Day 0 and 6 months later, Day 0 and 9 months later, Day 0 and 12 months later, Day 0 and 18 months later, Day 0 and 2 years later, Day 0 and 5 years later, or Day 0 and 10 years later) at a total dose of or at dosage levels sufficient to deliver a total dose of 0.010 mg, 0.025 mg, 0.100 mg or 0.400 mg.

(471) In some embodiments, the respiratory virus RNA (e.g., mRNA) vaccine for use in a method of vaccinating a subject is administered to the subject as a single dosage of between 10 µg/kg and 400 µg/kg of the nucleic acid vaccine (in an effective amount to vaccinate the subject). In some embodiments the RNA (e.g., mRNA) vaccine for use in a method of vaccinating a subject is administered to the subject as a single dosage of between 10 µg and 400 µg of the nucleic acid vaccine (in an effective amount to vaccinate the subject). In some embodiments, a respiratory virus

RNA (e.g., mRNA) vaccine for use in a method of vaccinating a subject is administered to the subject as a single dosage of 25-1000 μg (e.g., a single dosage of mRNA encoding hMPV, PIV3, RSV, MeV and/or BetaCoV antigen). In some embodiments, a respiratory virus RNA (e.g., mRNA) vaccine is administered to the subject as a single dosage of 25, 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950 or 1000 μg . For example, a respiratory virus RNA (e.g., mRNA) vaccine may be administered to a subject as a single dose of 25-100, 25-500, 50-100, 50-500, 50-1000, 100-500, 100-1000, 250-500, 250-1000, or 500-1000 μg . In some embodiments, a respiratory virus RNA (e.g., mRNA) vaccine for use in a method of vaccinating a subject is administered to the subject as two dosages, the combination of which equals 25-1000 μg of the respiratory virus RNA (e.g., mRNA) vaccine.

(472) A respiratory virus RNA (e.g. mRNA) vaccine pharmaceutical composition described herein can be formulated into a dosage form described herein, such as an intranasal, intratracheal, or injectable (e.g., intravenous, intraocular, intravitreal, intramuscular, intradermal, intracardiac, intraperitoneal, and subcutaneous).

(473) Respiratory Virus RNA (e.g., mRNA) Vaccine Formulations and Methods of Use

(474) Some aspects of the present disclosure provide formulations of the respiratory virus RNA (e.g., mRNA) vaccine, wherein the RNA (e.g., mRNA) vaccine is formulated in an effective amount to produce an antigen specific immune response in a subject (e.g., production of antibodies specific to an hMPV, PIV3, RSV, MeV and/or BetaCoV antigenic polypeptide). “An effective amount” is a dose of an RNA (e.g., mRNA) vaccine effective to produce an antigen-specific immune response. Also provided herein are methods of inducing an antigen-specific immune response in a subject.

(475) In some embodiments, the antigen-specific immune response is characterized by measuring an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide antibody titer produced in a subject administered a respiratory virus RNA (e.g., mRNA) vaccine as provided herein. An antibody titer is a measurement of the amount of antibodies within a subject, for example, antibodies that are specific to a particular antigen (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) or epitope of an antigen. Antibody titer is typically expressed as the inverse of the greatest dilution that provides a positive result. Enzyme-linked immunosorbent assay (ELISA) is a common assay for determining antibody titers, for example.

(476) In some embodiments, an antibody titer is used to assess whether a subject has had an infection or to determine whether immunizations are required. In some embodiments, an antibody titer is used to determine the strength of an autoimmune response, to determine whether a booster immunization is needed, to determine whether a previous vaccine was effective, and to identify any recent or prior infections. In accordance with the present disclosure, an antibody titer may be used to determine the strength of an immune response induced in a subject by the respiratory virus RNA (e.g., mRNA) vaccine.

(477) In some embodiments, an anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject is increased by at least 1 log relative to a control. For example, anti-antigenic polypeptide antibody titer produced in a subject may be increased by at least 1.5, at least 2, at least 2.5, or at least 3 log relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased by 1, 1.5, 2, 2.5 or 3 log relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased by 1-3 log relative to a control. For example, the anti-antigenic polypeptide antibody titer produced in a subject may be increased by 1-1.5, 1-2, 1-2.5, 1-3, 1.5-2, 1.5-2.5, 1.5-3, 2-2.5, 2-3, or 2.5-3 log relative to a control.

(478) In some embodiments, the anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject is

increased at least 2 times relative to a control. For example, the anti-antigenic polypeptide antibody titer produced in a subject may be increased at least 3 times, at least 4 times, at least 5 times, at least 6 times, at least 7 times, at least 8 times, at least 9 times, or at least 10 times relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is increased 2, 3, 4, 5, 6, 7, 8, 9, or 10 times relative to a control. In some embodiments, the anti-antigenic polypeptide antibody titer produced in a subject is increased 2-10 times relative to a control. For example, the anti-antigenic polypeptide antibody titer produced in a subject may be increased 2-10, 2-9, 2-8, 2-7, 2-6, 2-5, 2-4, 2-3, 3-10, 3-9, 3-8, 3-7, 3-6, 3-5, 3-4, 4-10, 4-9, 4-8, 4-7, 4-6, 4-5, 5-10, 5-9, 5-8, 5-7, 5-6, 6-10, 6-9, 6-8, 6-7, 7-10, 7-9, 7-8, 8-10, 8-9, or 9-10 times relative to a control.

(479) A control, in some embodiments, is the anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject who has not been administered a respiratory virus RNA (e.g., mRNA) vaccine of the present disclosure. In some embodiments, a control is an anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject who has been administered a live attenuated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine. An attenuated vaccine is a vaccine produced by reducing the virulence of a viable (live). An attenuated virus is altered in a manner that renders it harmless or less virulent relative to live, unmodified virus. In some embodiments, a control is an anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject administered inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine. In some embodiments, a control is an anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject administered a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. Recombinant protein vaccines typically include protein antigens that either have been produced in a heterologous expression system (e.g., bacteria or yeast) or purified from large amounts of the pathogenic organism. In some embodiments, a control is an anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject who has been administered an hMPV, PIV3, RSV, MeV and/or BetaCoV virus-like particle (VLP) vaccine. For example, an hMPV VLP vaccine used as a control may be a hMPV VLPs, comprising (or consisting of) viral matrix (M) and fusion (F) proteins, generated by expressing viral proteins in suspension-adapted human embryonic kidney epithelial (293-F) cells (see, e.g., Cox R G et al., *J Virol.* 2014 June; 88(11): 6368-6379, the contents of which are herein incorporated by reference).

(480) In some embodiments, an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose that is reduced compared to the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. A “standard of care,” as provided herein, refers to a medical or psychological treatment guideline and can be general or specific. “Standard of care” specifies appropriate treatment based on scientific evidence and collaboration between medical professionals involved in the treatment of a given condition. It is the diagnostic and treatment process that a physician/clinician should follow for a certain type of patient, illness or clinical circumstance. A “standard of care dose,” as provided herein, refers to the dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, or a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine, that a physician/clinician or other medical professional would administer to a subject to treat or prevent hMPV, PIV3, RSV, MeV and/or BetaCoV, or a hMPV-, PIV3-, RSV—, MeV—and/or BetaCoV-related condition, while following the standard of care guideline for treating or preventing hMPV, PIV3, RSV, MeV and/or BetaCoV, or a hMPV-, PIV3-, RSV—, MeV—and/or BetaCoV-related condition.

(481) In some embodiments, the anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a subject

administered an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is equivalent to an anti-antigenic polypeptide (e.g., an anti-hMPV, anti-PIV3, anti-RSV, anti-MeV and/or anti-BetaCoV antigenic polypeptide) antibody titer produced in a control subject administered a standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine or a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine.

(482) In some embodiments, an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose equivalent to an at least 2-fold reduction in a standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. For example, an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine may be a dose equivalent to an at least 3-fold, at least 4-fold, at least 5-fold, at least 6-fold, at least 7-fold, at least 8-fold, at least 9-fold, or at least 10-fold reduction in a standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. In some embodiments, an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose equivalent to an at least at least 100-fold, at least 500-fold, or at least 1000-fold reduction in a standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. In some embodiments, an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose equivalent to a 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 20-, 50-, 100-, 250-, 500-, or 1000-fold reduction in a standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. In some embodiments, the anti-antigenic polypeptide antibody titer produced in a subject administered an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant or protein hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine or a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine. In some embodiments, an effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose equivalent to a 2-fold to 1000-fold (e.g., 2-fold to 100-fold, 10-fold to 1000-fold) reduction in the standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine, wherein the anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine or a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine.

(483) In some embodiments, the effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose equivalent to a 2 to 1000-, 2 to 900-, 2 to 800-, 2 to 700-, 2 to 600-, 2 to 500-, 2 to 400-, 2 to 300-, 2 to 200-, 2 to 100-, 2 to 90-, 2 to 80-, 2 to 70-, 2 to 60-, 2 to 50-, 2 to 40-, 2 to 30-, 2 to 20-, 2 to 10-, 2 to 9-, 2 to 8-, 2 to 7-, 2 to 6-, 2 to 5-, 2 to 4-, 2 to 3-, 3 to 1000-, 3 to 900-, 3 to 800-, 3 to 700-, 3 to 600-, 3 to 500-, 3 to 400-, 3 to 3 to 00-, 3 to 200-, 3 to 100-, 3 to 90-, 3 to 80-, 3 to 70-, 3 to 60-, 3 to 50-, 3 to 40-, 3 to 30-, 3 to 20-, 3 to 10-, 3 to 9-, 3 to 8-, 3 to 7-, 3 to 6-, 3 to 5-, 3 to 4-, 4 to 1000-, 4 to 900-, 4 to 800-, 4 to 700-, 4 to 600-, 4 to 500-, 4 to 400-, 4 to 4 to 00-, 4 to 200-, 4 to 100-, 4 to 90-, 4 to 80-, 4 to 70-, 4 to 60-, 4 to 50-, 4 to 40-, 4 to 30-, 4 to 20-, 4 to 10-, 4 to 9-, 4 to 8-, 4 to 7-, 4 to 6-, 4 to 5-, 4 to 4-, 5 to 1000-, 5 to 900-, 5 to 800-, 5 to 700-, 5 to 600-, 5 to 500-, 5 to 400-, 5 to 300-, 5 to 200-, 5 to 100-, 5 to 90-, 5 to 80-, 5 to 70-, 5 to 60-, 5 to 50-, 5 to 40-, 5 to 30-, 5 to 20-, 5 to 10-, 5 to 9-, 5 to 8-, 5 to 7-, 5 to 6-, 6 to 1000-, 6 to 900-, 6 to 800-, 6 to 700-, 6 to 600-, 6 to 500-, 6 to 400-, 6 to 300-, 6 to 200-, 6 to 100-, 6 to 90-, 6 to 80-, 6 to 70-, 6 to 60-, 6 to 50-, 6 to 40-, 6 to 30-, 6 to 20-, 6 to 10-, 6 to 9-, 6 to 8-, 6 to 7-, 7 to 1000-, 7 to 900-, 7 to 800-, 7 to 700-, 7 to 600-, 7 to 500-, 7 to 400-, 7 to 300-, 7 to 200-, 7 to 100-, 7 to 90-, 7 to 80-, 7 to 70-, 7 to 60-, 7 to 50-, 7 to 40-, 7 to 30-, 7 to 20-, 7 to 10-, 7 to 9-, 7 to 8-, 8 to 1000-, 8 to 900-, 8 to 800-, 8 to 700-, 8 to 600-, 8 to 500-, 8 to 400-, 8 to 300-, 8 to 200-, 8 to 100-, 8 to 90-, 8 to 80-, 8 to 70-, 8 to 60-, 8 to 50-, 8 to 40-, 8 to 30-, 8 to 20-, 8 to 10-, 8 to 9-, 9 to 1000-, 9 to 900-, 9 to 800-, 9 to 700-, 9 to 600-, 9 to 500-, 9 to 400-, 9 to 300-, 9 to 200-, 9 to 100-, 9 to 90-, 9 to 80-, 9 to 70-, 9 to 60-, 9 to 50-, 9 to 40-, 9 to 30-, 9 to 20-, 9 to 10-, 10 to 1000-, 10 to 900-, 10 to 800-, 10 to

700-, 10 to 600-, 10 to 500-, 10 to 400-, 10 to 300-, 10 to 200-, 10 to 100-, 10 to 90-, 10 to 80-, 10 to 70-, 10 to 60-, 10 to 50-, 10 to 40-, 10 to 30-, 10 to 20-, 20 to 1000-, 20 to 900-, 20 to 800-, 20 to 700-, 20 to 600-, 20 to 500-, 20 to 400-, 20 to 300-, 20 to 200-, 20 to 100-, 20 to 90-, 20 to 80-, 20 to 70-, 20 to 60-, 20 to 50-, 20 to 40-, 20 to 30-, 30 to 1000-, 30 to 900-, 30 to 800-, 30 to 700-, 30 to 600-, 30 to 500-, 30 to 400-, 30 to 300-, 30 to 200-, 30 to 100-, 30 to 90-, 30 to 80-, 30 to 70-, 30 to 60-, 30 to 50-, 30 to 40-, 40 to 1000-, 40 to 900-, 40 to 800-, 40 to 700-, 40 to 600-, 40 to 500-, 40 to 400-, 40 to 300-, 40 to 200-, 40 to 100-, 40 to 90-, 40 to 80-, 40 to 70-, 40 to 60-, 40 to 50-, 50 to 1000-, 50 to 900-, 50 to 800-, 50 to 700-, 50 to 600-, 50 to 500-, 50 to 400-, 50 to 300-, 50 to 200-, 50 to 100-, 50 to 90-, 50 to 80-, 50 to 70-, 50 to 60-, 60 to 1000-, 60 to 900-, 60 to 800-, 60 to 700-, 60 to 600-, 60 to 500-, 60 to 400-, 60 to 300-, 60 to 200-, 60 to 100-, 60 to 90-, 60 to 80-, 60 to 70-, 70 to 1000-, 70 to 900-, 70 to 800-, 70 to 700-, 70 to 600-, 70 to 500-, 70 to 400-, 70 to 300-, 70 to 200-, 70 to 100-, 70 to 90-, 70 to 80-, 80 to 1000-, 80 to 900-, 80 to 800-, 80 to 700-, 80 to 600-, 80 to 500-, 80 to 400-, 80 to 300-, 80 to 200-, 80 to 100-, 80 to 90-, 90 to 1000-, 90 to 900-, 90 to 800-, 90 to 700-, 90 to 600-, 90 to 500-, 90 to 400-, 90 to 300-, 90 to 200-, 90 to 100-, 100 to 1000-, 100 to 900-, 100 to 800-, 100 to 700-, 100 to 600-, 100 to 500-, 100 to 400-, 100 to 300-, 100 to 200-, 200 to 1000-, 200 to 900-, 200 to 800-, 200 to 700-, 200 to 600-, 200 to 500-, 200 to 400-, 200 to 300-, 300 to 1000-, 300 to 900-, 300 to 800-, 300 to 700-, 300 to 600-, 300 to 500-, 300 to 400-, 400 to 1000-, 400 to 900-, 400 to 800-, 400 to 700-, 400 to 600-, 400 to 500-, 500 to 1000-, 500 to 900-, 500 to 800-, 500 to 700-, 500 to 600-, 600 to 1000-, 600 to 900-, 600 to 800-, 600 to 700-, 700 to 1000-, 700 to 900-, 700 to 800-, 800 to 1000-, 800 to 900-, or 900 to 1000-fold reduction in the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. In some embodiments, the anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine or a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine. In some embodiments, the effective amount is a dose equivalent to (or equivalent to an at least) 2-, 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 20-, 30-, 40-, 50-, 60-, 70-, 80-, 90-, 100-, 110-, 120-, 130-, 140-, 150-, 160-, 170-, 1280-, 190-, 200-, 210-, 220-, 230-, 240-, 250-, 260-, 270-, 280-, 290-, 300-, 310-, 320-, 330-, 340-, 350-, 360-, 370-, 380-, 390-, 400-, 410-, 420-, 430-, 440-, 450-, 4360-, 470-, 480-, 490-, 500-, 510-, 520-, 530-, 540-, 550-, 560-, 5760-, 580-, 590-, 600-, 610-, 620-, 630-, 640-, 650-, 660-, 670-, 680-, 690-, 700-, 710-, 720-, 730-, 740-, 750-, 760-, 770-, 780-, 790-, 800-, 810-, 820-, 830-, 840-, 850-, 860-, 870-, 880-, 890-, 900-, 910-, 920-, 930-, 940-, 950-, 960-, 970-, 980-, 990-, or 1000-fold reduction in the standard of care dose of a recombinant hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine. In some embodiments, an anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant or purified hMPV, PIV3, RSV, MeV and/or BetaCoV protein vaccine or a live attenuated or inactivated hMPV, PIV3, RSV, MeV and/or BetaCoV vaccine.

(484) In some embodiments, the effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a total dose of 50-1000 µg. In some embodiments, the effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a total dose of 50-1000, 50-900, 50-800, 50-700, 50-600, 50-500, 50-400, 50-300, 50-200, 50-100, 50-90, 50-80, 50-70, 50-60, 60-1000, 60-900, 60-800, 60-700, 60-600, 60-500, 60-400, 60-300, 60-200, 60-100, 60-90, 60-80, 60-70, 70-1000, 70-900, 70-800, 70-700, 70-600, 70-500, 70-400, 70-300, 70-200, 70-100, 70-90, 70-80, 80-1000, 80-900, 80-800, 80-700, 80-600, 80-500, 80-400, 80-300, 80-200, 80-100, 80-90, 90-1000, 90-900, 90-800, 90-700, 90-600, 90-500, 90-400, 90-300, 90-200, 90-100, 100-1000, 100-900, 100-800, 100-700, 100-600, 100-500, 100-400, 100-300, 100-200, 200-1000, 200-900, 200-800, 200-700, 200-600, 200-500, 200-400, 200-300, 300-1000, 300-900, 300-800, 300-700, 300-600, 300-500, 300-400, 400-1000, 400-900, 400-800, 400-700, 400-600, 400-500, 500-1000, 500-900, 500-800, 500-700, 500-600, 600-1000, 600-900, 600-800, 600-700, 700-1000, 700-900, 700-800, 800-1000, 800-900, or 900-

1000 µg. In some embodiments, the effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a total dose of 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950 or 1000 µg. In some embodiments, the effective amount is a dose of 25-500 g administered to the subject a total of two times. In some embodiments, the effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a dose of 25-500, 25-400, 25-300, 25-200, 25-100, 25-50, 50-500, 50-400, 50-300, 50-200, 50-100, 100-500, 100-400, 100-300, 100-200, 150-500, 150-400, 150-300, 150-200, 200-500, 200-400, 200-300, 250-500, 250-400, 250-300, 300-500, 300-400, 350-500, 350-400, 400-500 or 450-500 µg administered to the subject a total of two times. In some embodiments, the effective amount of a respiratory virus RNA (e.g., mRNA) vaccine is a total dose of 25, 50, 100, 150, 200, 250, 300, 350, 400, 450, or 500 µg administered to the subject a total of two times.

Examples of Additional Embodiments of the Disclosure

(485) Additional embodiments of the present disclosure are encompassed by the following numbered paragraphs: 1. A respiratory virus vaccine, comprising: at least one ribonucleic acid (RNA) polynucleotide having an open reading frame encoding at least one, at least two, at least three, at least four or at least five antigenic polypeptides selected from human metapneumovirus (hMPV) antigenic polypeptides or immunogenic fragments thereof, human parainfluenza virus type 3 (PIV3) antigenic polypeptides or immunogenic fragments thereof, respiratory syncytial virus (RSV) antigenic polypeptides or immunogenic fragments thereof, measles virus (MeV) antigenic polypeptides or immunogenic fragments thereof, and betacoronavirus (BetaCoV) antigenic polypeptides or immunogenic fragments thereof. 2. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and a PIV3 antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof. 3. The respiratory virus vaccine of paragraph 2, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, and/or wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13. 4. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and a RSV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof. 5. The respiratory virus vaccine of paragraph 4, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8. 6. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 7. The respiratory virus vaccine of paragraph 6, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, and/or wherein the MeV antigenic polypeptide

comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 8. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 9. The respiratory virus vaccine of paragraph 8, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 10. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof and a RSV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof. 11. The respiratory virus vaccine of paragraph 10, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13. 12. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof and a MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 13. The respiratory virus vaccine of paragraph 12, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 14. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 15. The respiratory virus vaccine of paragraph 14, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 16. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof and a MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof

and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 17. The respiratory virus vaccine of paragraph 16, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 18. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 19. The respiratory virus vaccine of paragraph 18, wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 20. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two RNA polynucleotides, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 21. The respiratory virus vaccine of paragraph 20, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 22. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, and a RSV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof. 23. The respiratory virus vaccine of paragraph 22, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, and/or wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13. 24. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, and a MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 25. The respiratory virus vaccine of paragraph 24, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID

NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 26. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 27. The respiratory virus vaccine of paragraph 26, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13 and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 23-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 23-34. 28. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, and a MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 29. The respiratory virus vaccine of paragraph 28, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, and/or wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 30. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 31. The respiratory virus vaccine of paragraph 30, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 23-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 23-34. 32. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic

polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 33. The respiratory virus vaccine of paragraph 32, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 23-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 23-34. 34. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, and a MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 35. The respiratory virus vaccine of paragraph 34, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 36. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 37. The respiratory virus vaccine of paragraph 36, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 23-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 23-34. 38. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 39. The respiratory virus vaccine of paragraph 38, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified

by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 23-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 23-34. 40. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two or three RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 41. The respiratory virus vaccine of paragraph 40, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 23-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 23-34. 42. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, and a MeV antigenic polypeptide or an immunogenic fragment thereof; or at least two, three or four RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof. 43. The respiratory virus vaccine of paragraph 42, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50. 44. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two, three or four RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 45. The respiratory virus vaccine of paragraph 44, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an

amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 46. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two, three or four RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 47. The respiratory virus vaccine of paragraph 46, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 48. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two, three or four RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 49. The respiratory virus vaccine of paragraph 48, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 50. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two, three or four

RNA polynucleotides, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 51. The respiratory virus vaccine of paragraph 50, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 52. The respiratory virus vaccine of paragraph 1, comprising: at least one RNA polynucleotide having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, a PIV3 antigenic polypeptide or an immunogenic fragment thereof, a RSV antigenic polypeptide or an immunogenic fragment thereof, a MeV antigenic polypeptide or an immunogenic fragment thereof, and a BetaCoV antigenic polypeptide or an immunogenic fragment thereof; or at least two, three, four or five RNA polynucleotides, one having an open reading frame encoding a hMPV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a PIV3 antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a RSV antigenic polypeptide or an immunogenic fragment thereof, one having an open reading frame encoding a MeV antigenic polypeptide or an immunogenic fragment thereof, and one having an open reading frame encoding a BetaCoV antigenic polypeptide or an immunogenic fragment thereof. 53. The respiratory virus vaccine of paragraph 52, wherein the hMPV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 5-8 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 5-8, wherein the PIV3 antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 12-13 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 12-13, wherein the MeV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 47-50 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 47-50, and/or wherein the BetaCoV antigenic polypeptide comprises an amino acid sequence identified by any one of SEQ ID NO: 24-34 or an amino acid sequence having at least 90% or 95% identity to an amino acid sequence identified by any one of SEQ ID NO: 24-34. 54. The vaccine of any one of paragraphs 1-53, wherein at least one RNA polynucleotide has less than 80% identity to wild-type mRNA sequence. 55. The vaccine of any one of paragraphs 1-53, wherein at least one RNA polynucleotide has at least 80% identity to wild-type mRNA sequence, but does not include wild-type mRNA sequence. 56. The vaccine of any one of paragraphs 1-55, wherein at least one antigenic polypeptide has membrane fusion activity, attaches to cell receptors, causes fusion of viral and cellular membranes, and/or is responsible for binding of the virus to a cell being infected. 57. The vaccine of any one of paragraphs 1-56, wherein at least one RNA polynucleotide comprises at least one chemical modification. 58. The vaccine of paragraph 57, wherein the chemical modification is selected from pseudouridine, N1-methylpseudouridine, N1-ethylpseudouridine, 2-thiouridine, 4'-thiouridine, 5-methylcytosine, 5-methyluridine, 2-thio-1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-pseudouridine, 2-thio-5-aza-uridine, 2-thio-dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-pseudouridine, 4-methoxy-2-thio-pseudouridine, 4-methoxy-pseudouridine, 4-thio-1-methyl-pseudouridine, 4-thio-pseudouridine, 5-aza-uridine, dihydropseudouridine, 5-methoxyuridine and

2'-O-methyl uridine. 59. The vaccine of paragraph 57 or 58, wherein the chemical modification is in the 5-position of the uracil. 60. The vaccine of any one of paragraphs 57-59, wherein the chemical modification is a N1-methylpseudouridine or N1-ethylpseudouridine. 61. The vaccine of any one of paragraphs 57-60, wherein at least 80%, at least 90% or 100% of the uracil in the open reading frame have a chemical modification. 62. The vaccine of any one of paragraphs 1-61, wherein at least one RNA polynucleotide further encodes at least one 5' terminal cap, optionally wherein the 5' terminal cap is 7mG(5')ppp(5')NlmpNp. 63. The vaccine of any one of paragraphs 1-62, wherein at least one antigenic polypeptide or immunogenic fragment thereof is fused to a signal peptide selected from: a HuIgGk signal peptide (METPAQLLFLLLLWLPDTTG; SEQ ID NO: 15); IgE heavy chain epsilon-1 signal peptide (MDWTWILFLVAAATRVHS; SEQ ID NO: 16); Japanese encephalitis PRM signal sequence (MLGSNSGQRVVFTILLVAPAYS; SEQ ID NO: 17), VSVg protein signal sequence (MKCLLYLAFLFIGVNCA; SEQ ID NO: 18) and Japanese encephalitis JEV signal sequence (MWLVSLAIVTACAGA; SEQ ID NO: 19). 64. The vaccine of paragraph 63, wherein the signal peptide is fused to the N-terminus or the C-terminus of at least one antigenic polypeptide. 65. The vaccine of any one of paragraphs 1-64, wherein the antigenic polypeptide or immunogenic fragment thereof comprises a mutated N-linked glycosylation site. 66. The vaccine of any one of paragraphs 1-65 formulated in a nanoparticle, optionally a lipid nanoparticle. 67. The vaccine of paragraph 66, wherein the lipid nanoparticle comprises a cationic lipid, a PEG-modified lipid, a sterol and a non-cationic lipid; optionally wherein the lipid nanoparticle carrier comprises a molar ratio of about 20-60% cationic lipid, 0.5-15% PEG-modified lipid, 25-55% sterol, and 25% non-cationic lipid; optionally wherein the cationic lipid is an ionizable cationic lipid and the non-cationic lipid is a neutral lipid, and the sterol is a cholesterol; and optionally wherein the cationic lipid is selected from 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA), dilinoleyl-methyl-4-dimethylaminobutyrate (DLin-MC3-DMA), and di((Z)-non-2-en-1-yl) 9-((4-(dimethylamino)butanoyl)oxy)heptadecanedioate (L319). Formula (II) 68. The vaccine of paragraph 66 or 67, wherein the nanoparticle (e.g., lipid nanoparticle) comprises a compound of Formula (I) and/or Formula (II), optionally Compound 3, 18, 20, 25, 26, 29, 30, 60, 108-112, or 122. 69. The vaccine of any one of paragraphs 1-68 further comprising an adjuvant, optionally a flagellin protein or peptide that optionally comprises an amino acid sequence identified by any one of SEQ ID NO: 54-56. 70. The vaccine of any one of paragraphs 1-69, wherein the open reading frame is codon-optimized. 71. The vaccine of any one of paragraphs 1-70 formulated in an effective amount to produce an antigen-specific immune response. 72. A method of inducing an immune response in a subject, the method comprising administering to the subject the vaccine of any one of paragraphs 1-71 in an amount effective to produce an antigen-specific immune response in the subject. 73. The method of paragraph 72, wherein the subject is administered a single dose of the vaccine, or wherein the subject is administered a first dose and then a booster dose of the vaccine. 74. The method of paragraph 72 or 73, wherein the vaccine is administered to the subject by intradermal injection or intramuscular injection. 75. The method of any one of paragraphs 72-74, wherein an anti-antigenic polypeptide antibody titer produced in the subject is increased by at least 1 log relative to a control, and/or wherein the anti-antigenic polypeptide antibody titer produced in the subject is increased at least 2 times relative to a control. 76. The method of any one of paragraphs 72-75, wherein the control is an anti-antigenic polypeptide antibody titer produced in a subject who has not been administered a vaccine against the virus, and/or wherein the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a live attenuated vaccine or an inactivated vaccine against the virus, and/or, wherein the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a recombinant protein vaccine or purified protein vaccine against the virus, and/or wherein the control is an anti-antigenic polypeptide antibody titer produced in a subject who has been administered a VLP vaccine against the virus. 77. The method of any one of paragraphs 72-76, wherein the effective amount is a dose equivalent to an at least 2-

fold reduction in the standard of care dose of a recombinant protein vaccine or a purified protein vaccine against the virus, and wherein an anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a recombinant protein vaccine or a purified protein vaccine against the virus, respectively; and/or wherein the effective amount is a dose equivalent to an at least 2-fold reduction in the standard of care dose of a live attenuated vaccine or an inactivated vaccine against the virus, and wherein an anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a live attenuated vaccine or an inactivated vaccine against the virus, respectively; and/or wherein the effective amount is a dose equivalent to an at least 2-fold reduction in the standard of care dose of a VLP vaccine against the virus, and wherein an anti-antigenic polypeptide antibody titer produced in the subject is equivalent to an anti-antigenic polypeptide antibody titer produced in a control subject administered the standard of care dose of a VLP vaccine against the virus. 78. The method of any one of paragraphs 72-77, wherein the effective amount is a total dose of 50 μ g-1000 μ g, optionally wherein the effective amount is a dose of 25 μ g, 100 μ g, 400 μ g, or 500 μ g administered to the subject a total of two times. 79. The method of any one of paragraphs 72-78, wherein the efficacy of the vaccine against the virus is greater than 65%; and/or wherein the vaccine immunizes the subject against the virus for up to 2 years or wherein the vaccine immunizes the subject against the virus for more than 2 years. 80. The method of any one of paragraphs 72-79, wherein the subject has an age of about 5 years old or younger or wherein the subject has an age of about 60 years old or older; and/or wherein the subject has a chronic pulmonary disease; and/or the subject has been exposed to the virus, wherein the subject is infected with the virus, or wherein the subject is at risk of infection by the virus; and/or wherein the subject is immunocompromised. 81. The respiratory virus vaccine of any one of paragraphs 1-71, comprising at least one (e.g., at least two, at least three, at least four, or at least five) RNA polynucleotide having an open reading frame encoding at least one (e.g., at least two, at least three, at least four, or at least five) antigenic polypeptide selected from hMPV antigenic polypeptides (SEQ ID NO: 5-8), PIV3 antigenic polypeptides (SEQ ID NO: 12-13), RSV antigenic polypeptides, MeV antigenic polypeptides (SEQ ID NO: 47-50) and BetaCoV antigenic polypeptides (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1; (SEQ ID NO: 24-34)), formulated in a cationic lipid nanoparticle (a) having a molar ratio of about 20-60% cationic lipid, about 5-25% non-cationic lipid, about 25-55% sterol, and about 0.5-15% PEG-modified lipid, and/or (b) comprising a compound of Formula (I) and/or Formula (II), wherein the at least one (e.g., at least two, at least three, at least four, or at least five) RNA polynucleotide comprises at least one chemical modification. 82. The respiratory virus vaccine of any one of paragraphs 1-71, comprising at least one (e.g., at least two, at least three, at least four, or at least five) RNA polynucleotide having an open reading frame encoding at least one (e.g., at least two, at least three, at least four, or at least five) antigenic polypeptide selected from hMPV antigenic polypeptides (SEQ ID NO: 5-8), PIV3 antigenic polypeptides (SEQ ID NO: 12-13), RSV antigenic polypeptides, MeV antigenic polypeptides (SEQ ID NO: 47-50) and BetaCoV antigenic polypeptides (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1; (SEQ ID NO: 24-34)), formulated in a cationic lipid nanoparticle (a) having a molar ratio of about 20-60% cationic lipid, about 5-25% non-cationic lipid, about 25-55% sterol, and about 0.5-15% PEG-modified lipid, and/or (b) comprising at least one (e.g., at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14) Compound selected from Compounds 3, 18, 20, 25, 26, 29, 30, 60, 108-112 and 122. 83. The respiratory virus vaccine of paragraphs 81 or 82, wherein the at least one antigenic polypeptide is selected from hMPV antigenic polypeptides (e.g., SEQ ID NO: 5-8). 84. The respiratory virus vaccine of any one of paragraphs 81-83, wherein the at least one antigenic polypeptide is selected from PIV3 antigenic polypeptides (e.g., SEQ ID NO: 12-13). 85. The

respiratory virus vaccine of any one of paragraphs 81-84, wherein the at least one antigenic polypeptide is selected from RSV antigenic polypeptides. 86. The respiratory virus vaccine of any one of paragraphs 81-85, wherein the at least one antigenic polypeptide is selected from MeV antigenic polypeptides (e.g., SEQ ID NO: 47-50). 87. The respiratory virus vaccine of any one of paragraphs 81-86, wherein the at least one antigenic polypeptide is selected from BetaCoV antigenic polypeptides (e.g., SEQ ID NO: 24-34). 88. The respiratory virus vaccine of paragraph 87, wherein the BetaCoV antigenic polypeptides are MERS antigenic polypeptides. 89. The respiratory virus vaccine of paragraph 87, wherein the BetaCoV antigenic polypeptides are SARS antigenic polypeptides. 90. The respiratory virus vaccine of any one of paragraphs 81-89, wherein the at least one (e.g., at least two, at least three, at least four, or at least five) RNA polynucleotide comprises at least one chemical modification (e.g., selected from pseudouridine, N1-methylpseudouridine, N1-ethylpseudouridine, 2-thiouridine, 4'-thiouridine, 5-methylcytosine, 5-methyluridine, 2-thio-1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-pseudouridine, 2-thio-5-aza-uridine, 2-thio-dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-pseudouridine, 4-methoxy-2-thio-pseudouridine, 4-methoxy-pseudouridine, 4-thio-1-methyl-pseudouridine, 4-thio-pseudouridine, 5-aza-uridine, dihydropseudouridine, 5-methoxyuridine and 2'-O-methyl uridine). 91. A respiratory virus vaccine, comprising: at least one messenger ribonucleic acid (mRNA) polynucleotide having a 5' terminal cap, an open reading frame encoding at least one respiratory virus antigenic polypeptide, and a 3' polyA tail. 92. The vaccine of paragraph 91, wherein the at least one mRNA polynucleotide comprises a sequence identified by any one of SEQ ID NO: 57-80. 93. The vaccine of paragraph 91 or 92, wherein the 5' terminal cap is or comprises 7mG(5')ppp(5')NlmpNp. 94. The vaccine of any one of paragraphs 91-93, wherein 100% of the uracil in the open reading frame is modified to include N1-methyl pseudouridine at the 5-position of the uracil. 95. The vaccine of any one of paragraphs 91-94, wherein the vaccine is formulated in a lipid nanoparticle comprising: DLin-MC3-DMA; cholesterol; 1,2-Distearoyl-sn-glycero-3-phosphocholine (DSPC); and polyethylene glycol (PEG)2000-DMG. 96. The vaccine of paragraph 95, wherein the lipid nanoparticle further comprises trisodium citrate buffer, sucrose and water. 97. A respiratory syncytial virus (RSV) vaccine, comprising: at least one messenger ribonucleic acid (mRNA) polynucleotide having a 5' terminal cap 7mG(5')ppp(5')NlmpNp, a sequence identified by any one of SEQ ID NO: 57-80 and a 3' polyA tail, formulated in a lipid nanoparticle comprising DLin-MC3-DMA, cholesterol, 1,2-Distearoyl-sn-glycero-3-phosphocholine (DSPC), and polyethylene glycol (PEG)2000-DMG, wherein the uracil nucleotides of the sequence identified by any one of SEQ ID NO: 57-80 are modified to include N1-methyl pseudouridine at the 5-position of the uracil nucleotide.

(486) This disclosure is not limited in its application to the details of construction and the arrangement of components set forth in the following description or illustrated in the drawings. The disclosure is capable of other embodiments and of being practiced or of being carried out in various ways. Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of “including,” “comprising,” or “having,” “containing,” “involving,” and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

EXAMPLES

Example 1: Manufacture of Polynucleotides

(487) According to the present disclosure, the manufacture of polynucleotides and/or parts or regions thereof may be accomplished utilizing the methods taught in International Publication WO2014/152027, entitled “Manufacturing Methods for Production of RNA Transcripts,” the contents of which is incorporated herein by reference in its entirety.

(488) Purification methods may include those taught in International Publication WO2014/152030 and International Publication WO2014/152031, each of which is incorporated herein by reference in its entirety.

(489) Detection and characterization methods of the polynucleotides may be performed as taught in International Publication WO2014/144039, which is incorporated herein by reference in its entirety.

(490) Characterization of the polynucleotides of the disclosure may be accomplished using polynucleotide mapping, reverse transcriptase sequencing, charge distribution analysis, detection of RNA impurities, or any combination of two or more of the foregoing. "Characterizing" comprises determining the RNA transcript sequence, determining the purity of the RNA transcript, or determining the charge heterogeneity of the RNA transcript, for example. Such methods are taught in, for example, International Publication WO2014/144711 and International Publication WO2014/144767, the content of each of which is incorporated herein by reference in its entirety.

Example 2: Chimeric Polynucleotide Synthesis

(491) According to the present disclosure, two regions or parts of a chimeric polynucleotide may be joined or ligated using triphosphate chemistry. A first region or part of 100 nucleotides or less is chemically synthesized with a 5' monophosphate and terminal 3'desOH or blocked OH, for example. If the region is longer than 80 nucleotides, it may be synthesized as two strands for ligation.

(492) If the first region or part is synthesized as a non-positionally modified region or part using in vitro transcription (IVT), conversion the 5'monophosphate with subsequent capping of the 3' terminus may follow.

(493) Monophosphate protecting groups may be selected from any of those known in the art.

(494) The second region or part of the chimeric polynucleotide may be synthesized using either chemical synthesis or IVT methods. IVT methods may include an RNA polymerase that can utilize a primer with a modified cap. Alternatively, a cap of up to 130 nucleotides may be chemically synthesized and coupled to the IVT region or part.

(495) For ligation methods, ligation with DNA T4 ligase, followed by treatment with DNase should readily avoid concatenation.

(496) The entire chimeric polynucleotide need not be manufactured with a phosphate-sugar backbone. If one of the regions or parts encodes a polypeptide, then such region or part may comprise a phosphate-sugar backbone.

(497) Ligation is then performed using any known click chemistry, orthoclick chemistry, solulink, or other bioconjugate chemistries known to those in the art.

(498) Synthetic Route

(499) The chimeric polynucleotide may be made using a series of starting segments. Such segments include: (a) a capped and protected 5' segment comprising a normal 3'OH (SEG. 1) (b) a 5' triphosphate segment, which may include the coding region of a polypeptide and a normal 3'OH (SEG. 2) (c) a 5' monophosphate segment for the 3' end of the chimeric polynucleotide (e.g., the tail) comprising cordycepin or no 3'OH (SEG. 3)

(500) After synthesis (chemical or IVT), segment 3 (SEG. 3) may be treated with cordycepin and then with pyrophosphatase to create the 5' monophosphate.

(501) Segment 2 (SEG. 2) may then be ligated to SEG. 3 using RNA ligase. The ligated polynucleotide is then purified and treated with pyrophosphatase to cleave the diphosphate. The treated SEG.2-SEG. 3 construct may then be purified and SEG. 1 is ligated to the 5' terminus. A further purification step of the chimeric polynucleotide may be performed.

(502) Where the chimeric polynucleotide encodes a polypeptide, the ligated or joined segments may be represented as: 5'UTR (SEG. 1), open reading frame or ORF (SEG. 2) and 3'UTR+PolyA (SEG. 3).

(503) The yields of each step may be as much as 90-95%.

Example 3: PCR for cDNA Production

(504) PCR procedures for the preparation of cDNA may be performed using 2×KAPA HIFI™ HotStart ReadyMix by Kapa Biosystems (Woburn, MA). This system includes 2×KAPA ReadyMix 12.5 µl; Forward Primer (10 µM) 0.75 µl; Reverse Primer (10 PM) 0.75 µl; Template cDNA 100

ng; and dH.sub.2O diluted to 25.0 µl. The reaction conditions may be at 95° C. for 5 min. The reaction may be performed for 25 cycles of 98° C. for 20 sec, then 58° C. for 15 sec, then 72° C. for 45 sec, then 72° C. for 5 min, then 4° C. to termination.

(505) The reaction may be cleaned up using Invitrogen's PURELINK™ PCR Micro Kit (Carlsbad, CA) per manufacturer's instructions (up to 5 µg). Larger reactions may require a cleanup using a product with a larger capacity. Following the cleanup, the cDNA may be quantified using the NANODROP™ and analyzed by agarose gel electrophoresis to confirm that the cDNA is the expected size. The cDNA may then be submitted for sequencing analysis before proceeding to the in vitro transcription reaction.

Example 4: In Vitro Transcription (IVT)

(506) The in vitro transcription reaction generates RNA polynucleotides. Such polynucleotides may comprise a region or part of the polynucleotides of the disclosure, including chemically modified RNA (e.g., mRNA) polynucleotides. The chemically modified RNA polynucleotides can be uniformly modified polynucleotides. The in vitro transcription reaction utilizes a custom mix of nucleotide triphosphates (NTPs). The NTPs may comprise chemically modified NTPs, or a mix of natural and chemically modified NTPs, or natural NTPs.

(507) A typical in vitro transcription reaction includes the following:

(508) TABLE-US-00001 1) Template cDNA 1.0 µg 2) 10× transcription buffer 2.0 µl (400 mM Tris-HCl pH 8.0, 190 mM MgCl.sub.2, 50 mM DTT, 10 mM Spermidine) 3) Custom NTPs (25 mM each) 0.2 µl 4) RNase Inhibitor 20 U 5) T7 RNA polymerase 3000 U 6) dH.sub.2O up to 20.0 µl. and 7) Incubation at 37° C. for 3 hr-5 hrs.

(509) The crude IVT mix may be stored at 4° C. overnight for cleanup the next day. 1 U of RNase-free DNase may then be used to digest the original template. After 15 minutes of incubation at 37° C., the mRNA may be purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. This kit can purify up to 500 µg of RNA. Following the cleanup, the RNA polynucleotide may be quantified using the NanoDrop and analyzed by agarose gel electrophoresis to confirm the RNA polynucleotide is the proper size and that no degradation of the RNA has occurred.

Example 5: Enzymatic Capping

(510) Capping of a RNA polynucleotide is performed as follows where the mixture includes: IVT RNA 60 µg-180 µg and dH.sub.2O up to 72 µl. The mixture is incubated at 65° C. for 5 minutes to denature RNA, and then is transferred immediately to ice.

(511) The protocol then involves the mixing of 10× Capping Buffer (0.5 M Tris-HCl (pH 8.0), 60 mM KCl, 12.5 mM MgCl.sub.2) (10.0 µl); 20 mM GTP (5.0 µl); 20 mM S-Adenosyl Methionine (2.5 µl); RNase Inhibitor (100 U); 2'-O-Methyltransferase (400U); Vaccinia capping enzyme (Guanylyl transferase) (40 U); dH.sub.2O (Up to 28 µl); and incubation at 37° C. for 30 minutes for 60 µg RNA or up to 2 hours for 180 µg of RNA.

(512) The RNA polynucleotide may then be purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. Following the cleanup, the RNA may be quantified using the NANODROP™ (ThermoFisher, Waltham, MA) and analyzed by agarose gel electrophoresis to confirm the RNA polynucleotide is the proper size and that no degradation of the RNA has occurred. The RNA polynucleotide product may also be sequenced by running a reverse-transcription-PCR to generate the cDNA for sequencing.

Example 6: PolyA Tailing Reaction

(513) Without a poly-T in the cDNA, a poly-A tailing reaction must be performed before cleaning the final product. This is done by mixing capped IVT RNA (100 µl); RNase Inhibitor (20 U); 10× Tailing Buffer (0.5 M Tris-HCl (pH 8.0), 2.5 M NaCl, 100 mM MgCl.sub.2) (12.0 µl); 20 mM ATP (6.0 µl); Poly-A Polymerase (20 U); dH.sub.2O up to 123.5 µl and incubation at 37° C. for 30 min. If the poly-A tail is already in the transcript, then the tailing reaction may be skipped and proceed directly to cleanup with Ambion's MEGACLEAR™ kit (Austin, TX) (up to 500 µg). Poly-A

Polymerase may be a recombinant enzyme expressed in yeast.

(514) It should be understood that the processivity or integrity of the polyA tailing reaction may not always result in an exact size polyA tail. Hence, polyA tails of approximately between 40-200 nucleotides, e.g., about 40, 50, 60, 70, 80, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 150-165, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164 or 165 are within the scope of the present disclosure.

Example 7: Natural 5' Caps and 5' Cap Analogues

(515) 5'-capping of polynucleotides may be completed concomitantly during the in vitro-transcription reaction using the following chemical RNA cap analogs to generate the 5'-guanosine cap structure according to manufacturer protocols: 3'-O-Me-m⁷G(5')ppp(5') G [the ARCA cap]; G(5')ppp(5')A; G(5')ppp(5')G; m⁷G(5')ppp(5')A; m⁷G(5')ppp(5')G (New England BioLabs, Ipswich, MA). 5'-capping of modified RNA may be completed post-transcriptionally using a Vaccinia Virus Capping Enzyme to generate the "Cap 0" structure: m⁷G(5')ppp(5')G (New England BioLabs, Ipswich, MA). Cap 1 structure may be generated using both Vaccinia Virus Capping Enzyme and a 2'-O methyl-transferase to generate: m⁷G(5')ppp(5')G-2'-O-methyl. Cap 2 structure may be generated from the Cap 1 structure followed by the 2'-O-methylation of the 5'-antepenultimate nucleotide using a 2'-O methyl-transferase. Cap 3 structure may be generated from the Cap 2 structure followed by the 2'-O-methylation of the 5'-preantepenultimate nucleotide using a 2'-O methyl-transferase. Enzymes are preferably derived from a recombinant source.

(516) When transfected into mammalian cells, the modified mRNAs have a stability of between 12-18 hours or more than 18 hours, e.g., 24, 36, 48, 60, 72 or greater than 72 hours.

Example 8: Capping Assays

(517) Protein Expression Assay

(518) Polynucleotides (e.g., mRNA) encoding a polypeptide, containing any of the caps taught herein, can be transfected into cells at equal concentrations. The amount of protein secreted into the culture medium can be assayed by ELISA at 6, 12, 24 and/or 36 hours post-transfection. Synthetic polynucleotides that secrete higher levels of protein into the medium correspond to a synthetic polynucleotide with a higher translationally-competent cap structure.

(519) Purity Analysis Synthesis

(520) RNA (e.g., mRNA) polynucleotides encoding a polypeptide, containing any of the caps taught herein can be compared for purity using denaturing Agarose-Urea gel electrophoresis or HPLC analysis. RNA polynucleotides with a single, consolidated band by electrophoresis correspond to the higher purity product compared to polynucleotides with multiple bands or streaking bands. Chemically modified RNA polynucleotides with a single HPLC peak also correspond to a higher purity product. The capping reaction with a higher efficiency provides a more pure polynucleotide population.

(521) Cytokine Analysis

(522) RNA (e.g., mRNA) polynucleotides encoding a polypeptide, containing any of the caps taught herein can be transfected into cells at multiple concentrations. The amount of pro-inflammatory cytokines, such as TNF-alpha and IFN-beta, secreted into the culture medium can be assayed by ELISA at 6, 12, 24 and/or 36 hours post-transfection. RNA polynucleotides resulting in the secretion of higher levels of pro-inflammatory cytokines into the medium correspond to a polynucleotides containing an immune-activating cap structure.

(523) Capping Reaction Efficiency

(524) RNA (e.g., mRNA) polynucleotides encoding a polypeptide, containing any of the caps taught herein can be analyzed for capping reaction efficiency by LC-MS after nuclease treatment. Nuclease treatment of capped polynucleotides yield a mixture of free nucleotides and the capped 5'-5'-triphosphate cap structure detectable by LC-MS. The amount of capped product on the LC-MS spectra can be expressed as a percent of total polynucleotide from the reaction and correspond to capping reaction efficiency. The cap structure with a higher capping reaction efficiency has a

higher amount of capped product by LC-MS.

Example 9: Agarose Gel Electrophoresis of Modified RNA or RT PCR Products

(525) Individual RNA polynucleotides (200-400 ng in a 20 µl volume) or reverse transcribed PCR products (200-400 ng) may be loaded into a well on a non-denaturing 1.2% Agarose E-Gel (Invitrogen, Carlsbad, CA) and run for 12-15 minutes, according to the manufacturer protocol.

Example 10: Nanodrop Modified RNA Quantification and UV Spectral Data

(526) Chemically modified RNA polynucleotides in TE buffer (1 µl) are used for Nanodrop UV absorbance readings to quantitate the yield of each polynucleotide from an chemical synthesis or in vitro transcription reaction.

Example 11: Formulation of Modified mRNA Using Lipidoids

(527) RNA (e.g., mRNA) polynucleotides may be formulated for in vitro experiments by mixing the polynucleotides with the lipidoid at a set ratio prior to addition to cells. In vivo formulation may require the addition of extra ingredients to facilitate circulation throughout the body. To test the ability of these lipidoids to form particles suitable for in vivo work, a standard formulation process used for siRNA-lipidoid formulations may be used as a starting point. After formation of the particle, polynucleotide is added and allowed to integrate with the complex. The encapsulation efficiency is determined using a standard dye exclusion assays.

Example 12: Immunogenicity Study

(528) The instant study is designed to test the immunogenicity in mice of candidate hMPV vaccines comprising a mRNA polynucleotide encoding Fusion (F) glycoprotein, major surface glycoprotein G, or a combination thereof, obtained from hMPV.

(529) Mice are immunized intravenously (IV), intramuscularly (IM), or intradermally (ID) with candidate vaccines. Candidate vaccines are chemically modified or unmodified. A total of four immunizations are given at 3-week intervals (i.e., at weeks 0, 3, 6, and 9), and sera are collected after each immunization until weeks 33-51. Serum antibody titers against Fusion (F) glycoprotein or major surface glycoprotein (G) protein are determined by ELISA. Sera collected from each mouse during weeks 10-16 are pooled, and total IgG purified. Purified antibodies are used for immunoelectron microscopy, antibody-affinity testing, and in vitro protection assays.

Example 13: hMPV Rodent Challenge

(530) The instant study is designed to test the efficacy in cotton rats of candidate hMPV vaccines against a lethal challenge using an hMPV vaccine comprising mRNA encoding Fusion (F) glycoprotein, major surface glycoprotein G, or a combination of both antigens obtained from hMPV. Cotton rats are challenged with a lethal dose of the hMPV.

(531) Animals are immunized intravenously (IV), intramuscularly (IM), or intradermally (ID) at week 0 and week 3 with candidate hMPV vaccines with and without adjuvant. Candidate vaccines are chemically modified or unmodified. The animals are then challenged with a lethal dose of hMPV on week 7 via IV, IM or ID. Endpoint is day 13 post infection, death or euthanasia. Animals displaying severe illness as determined by >30% weight loss, extreme lethargy or paralysis are euthanized. Body temperature and weight are assessed and recorded daily.

(532) In experiments where a lipid nanoparticle (LNP) formulation is used, the formulation may include a cationic lipid, non-cationic lipid, PEG lipid and structural lipid in the ratios 50:10:1.5:38.5. The cationic lipid is DLin-KC2-DMA (50 mol %) or DLin-MC3-DMA (50 mol %), the non-cationic lipid is DSPC (10 mol %), the PEG lipid is PEG-DOMG (1.5 mol %) and the structural lipid is cholesterol (38.5 mol %), for example.

Example 14: Immunogenicity of hMPV mRNA Vaccine in BALB/c Mice

(533) The instant study was designed to test the immunogenicity in BALB/c mice of hMPV vaccines comprising an mRNA polynucleotide encoding the hMPV Fusion (F) glycoprotein. The mRNA polynucleotide encodes the full-length fusion protein and comprises the wild-type nucleotide sequence obtained from the hMPV A2a strain. Mice were divided into 3 groups (n=8 for each group) and immunized intramuscularly (IM) with PBS, a 10 µg dose of mRNA vaccines

encoding hMPV fusion protein, or a 2 µg dose of mRNA vaccines encoding hMPV fusion protein. A total of two immunizations were given at 3-week intervals (i.e., at weeks 0, and 3 weeks), and sera were collected after each immunization according to the schedule described in Table 1. Serum antibody titers against hMPV fusion glycoprotein were determined by ELISA and antibodies were detected in the sera collected on day 14 onward. Both vaccine doses tested induced comparable levels of immune response in mice (FIGS. 2A-2C).

(534) Additionally, mice sera were used for IgG isotyping (FIGS. 3A-3C). Both hMPV fusion protein-specific IgG1 and IgG2a were detected in mice sera. hMPV fusion protein mRNA vaccine also induced Th1 and Th2 cytokine responses, with a Th1 bias.

(535) Sera from mice immunized with either 10 µg or 2 µg doses of the hMPV fusion protein mRNA vaccine contain neutralizing antibodies. The ability of these antibodies to neutralize hMPV B2 strain was also tested. The antibody-containing sera successfully neutralized the hMPV B2 virus (FIG. 4).

Example 15: T-Cell Stimulation

(536) The instant study was designed to test T-cell stimulation in the splenocytes of mice immunized with mRNA vaccines encoding hMPV fusion protein, as described herein.

Immunization of BALB/c mice was performed as described in Example 14. The splenocytes for each group were pooled and split into two parts. One part of splenocytes from each group of mice was stimulated with hMPV-free media, Concanavalin A or a hMPV fusion protein peptide pool comprising 15-mers (15 amino acids long); while the other part of splenocytes from each group of mice was stimulated with hMPV-free media, Concanavalin A or inactivated hMPV virus. Secreted mouse cytokines were measured using the Meso Scale Discovery (MSD) assay.

(537) Cytokines specific to Th1 or Th2 responses were measured. For Th1 response, IFN- γ , IL2 and IL12 were detected from splenocytes stimulated with the hMPV fusion protein peptide pool at a level comparable to that of Concanavalin A (FIGS. 5A-5C). For a Th2 response, the hMPV fusion protein peptide pool induced the secretion of detectable IL10, TNF- α , IL4 and IL6, but not IL5, while Concanavalin A stimulated the secretion of all the above-mentioned Th2 cytokines (FIGS. 6A-6E) at a much higher level.

(538) In contrast, inactivated hMPV virus only induced the secretion of IL2 in the Th1 response comparable to that of Concanavalin A (FIGS. 7A-7C). For the Th2 response, the inactivated hMPV virus induced the secretion of detectable IL10, TNF- α , IL4 and IL6, but not IL5, while Concanavalin A stimulated the secretion of all the above-mentioned Th2 cytokines (FIGS. 8A-8E) at a much higher level.

Example 16: hMPV Rodent Challenge in Cotton Rats Immunized with mRNA Vaccine Encoding hMPV Fusion Protein

(539) The instant study was designed to test the efficacy in cotton rats of hMPV vaccines against a lethal challenge. mRNA vaccines encoding hMPV fusion protein were used. The mRNA polynucleotide encodes a full-length fusion protein and comprises the wild-type nucleotide sequence obtained from the hMPV A2a strain.

(540) Cotton rats were immunized intramuscularly (IM) at week 0 and week 3 with the mRNA vaccines encoding hMPV fusion protein with either 2 µg or 10 µg doses for each immunization. The animals were then challenged with a lethal dose of hMPV in week 7 post initial immunization via IV, IM or ID. The endpoint was day 13 post infection, death or euthanasia. Viral titers in the noses and lungs of the cotton rats were measured. The results (FIGS. 9A and 9B) show that a 10 µg dose of mRNA vaccine protected the cotton mice 100% in the lung and drastically reduced the viral titer in the nose after challenge (~2 log reduction). Moreover, a 2 µg dose of mRNA vaccine showed a 1 log reduction in lung viral titer in the cotton mice challenged.

(541) Further, the histopathology of the lungs of the cotton mice immunized and challenged showed no pathology associated with vaccine-enhanced disease (FIG. 10).

Example 17: Immunogenicity Study

(542) The instant study is designed to test the immunogenicity in mice of candidate PIV3 vaccines comprising a mRNA polynucleotide encoding hemagglutinin-neuraminidase or fusion protein (F or F0) obtained from PIV3.

(543) Mice are immunized intravenously (IV), intramuscularly (IM), or intradermally (ID) with candidate vaccines. Candidate vaccines are chemically modified or unmodified. A total of four immunizations are given at 3-week intervals (i.e., at weeks 0, 3, 6, and 9), and sera are collected after each immunization until weeks 33-51. Serum antibody titers against hemagglutinin-neuraminidase or fusion protein (F or F0) are determined by ELISA. Sera collected from each mouse during weeks 10-16 are, optionally, pooled, and total IgGs are purified. Purified antibodies are used for immunoelectron microscopy, antibody-affinity testing, and in vitro protection assays.

Example 18: PIV3 Rodent Challenge

(544) The instant study is designed to test the efficacy in cotton rats of candidate PIV3 vaccines against a lethal challenge using a PIV3 vaccine comprising mRNA encoding hemagglutinin-neuraminidase or fusion protein (F or F0) obtained from PIV3. Cotton rats are challenged with a lethal dose of the PIV3.

(545) Animals are immunized intravenously (IV), intramuscularly (IM), or intradermally (ID) at week 0 and week 3 with candidate PIV3 vaccines with and without adjuvant. Candidate vaccines are chemically modified or unmodified. The animals are then challenged with a lethal dose of PIV3 on week 7 via IV, IM or ID. Endpoint is day 13 post infection, death or euthanasia. Animals displaying severe illness as determined by >30% weight loss, extreme lethargy or paralysis are euthanized. Body temperature and weight are assessed and recorded daily.

(546) In experiments where a lipid nanoparticle (LNP) formulation is used, the formulation may include a cationic lipid, non-cationic lipid, PEG lipid and structural lipid in the ratios 50:10:1.5:38.5. The cationic lipid is DLin-KC2-DMA (50 mol %) or DLin-MC3-DMA (50 mol %), the non-cationic lipid is DSPC (10 mol %), the PEG lipid is PEG-DOMG (1.5 mol %) and the structural lipid is cholesterol (38.5 mol %), for example.

Example 19: hMPV/PIV Cotton Rat Challenge

(547) The instant study was designed to test the efficacy in cotton rats of candidate hMPV mRNA vaccines, PIV3 mRNA vaccines, or hMPV/PIV combination mRNA vaccines against a lethal challenge using PIV3 strain or hMPV/A2 strain. The study design is shown in Table 9.

(548) Cotton rats of 10-12 weeks old were divided into 12 groups (n=5), and each group was vaccinated with mRNA vaccines indicated in Table 9. The PIV3 vaccine comprises mRNA encoding hemagglutinin-neuraminidase or fusion protein (F or F0) obtained from PIV3. The hMPV mRNA vaccine encodes the full-length hMPV fusion protein. The hMPV/PIV combination mRNA vaccine is a mixture of the PIV3 vaccine and hMPV vaccine at a 1:1 ratio.

(549) Cotton rats were immunized intramuscularly (IM) at week 0 and week 3 with candidate vaccines with the doses indicated in Table 9. Cotton rats immunized with hMPV mRNA vaccines or hMPV/PIV combination mRNA vaccines were challenged with a lethal dose of hMPV/A2 strain on week 7 via IM. Cotton rats immunized with PIV mRNA vaccines or hMPV/PIV combination mRNA vaccines were challenged with a lethal dose of PIV3 strain on week 7 via IM.

(550) The endpoint was day 13 post infection, death or euthanasia. Animals displaying severe illness as determined by >30% weight loss, extreme lethargy or paralysis were euthanized. Body temperature and weight were assessed and recorded daily.

(551) Lung and nose hMPV/A2 (FIG. 12) or PIV3 (FIG. 13) viral titers were assessed. Lung histopathology of the immunized and challenged cotton rat immunized and challenged were assessed to determine pathology associated with vaccine enhance disease. Neutralization antibody titers in the serum of immunized cotton rats on day 0 and 42 post immunization were assessed (FIG. 11).

(552) hMPV/A2 (FIG. 14) or PIV3 (FIG. 15) neutralizing antibody titers in the serum samples of the immunized cotton rat 42 days post immunization were measured. All mRNA vaccines tested

induced strong neutralizing antibodies cotton rats. Lung histopathology of the immunized cotton rats were also evaluated (FIG. 16). Low occurrence of alevolitis and interstitial pneumonia was observed, indicating no antibody-dependent enhancement (ADE) of hMPV or PIV associated diseases.

Example 20: Betacoronavirus Immunogenicity Study

(553) The instant study is designed to test the immunogenicity in rabbits of candidate betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1 or a combination thereof) vaccines comprising a mRNA polynucleotide encoding the spike (S) protein, the S1 subunit (S1) of the spike protein, or the S2 subunit (S2) of the spike protein obtained from a betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1).

(554) Rabbits are vaccinated on week 0 and 3 via intravenous (IV), intramuscular (IM), or intradermal (ID) routes. One group remains unvaccinated and one is administered inactivated betacoronavirus. Serum is collected from each rabbit on weeks 1, 3 (pre-dose) and 5. Individual bleeds are tested for anti-S, anti-S1 or anti-S2 activity via a virus neutralization assay from all three time points, and pooled samples from week 5 only are tested by Western blot using inactivated betacoronavirus (e.g., inactivated MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1).

(555) In experiments where a lipid nanoparticle (LNP) formulation is used, the formulation may include a cationic lipid, non-cationic lipid, PEG lipid and structural lipid in the ratios 50:10:1.5:38.5. The cationic lipid is DLin-KC2-DMA (50 mol %) or DLin-MC3-DMA (50 mol %), the non-cationic lipid is DSPC (10 mol %), the PEG lipid is PEG-DOMG (1.5 mol %) and the structural lipid is cholesterol (38.5 mol %), for example.

Example 21: Betacoronavirus Challenge

(556) The instant study is designed to test the efficacy in rabbits of candidate betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV—HKU1 or a combination thereof) vaccines against a lethal challenge using a betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV—HKU1 or a combination thereof) vaccine comprising mRNA encoding the spike (S) protein, the S1 subunit (S1) of the spike protein, or the S2 subunit (S2) of the spike protein obtained from betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1). Rabbits are challenged with a lethal dose ($10 \times \text{LD}_{90}$; ~ 100 plaque-forming units; PFU) of betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1).

(557) The animals used are 6-8 week old female rabbits in groups of 10. Rabbits are vaccinated on weeks 0 and 3 via an IM, ID or IV route of administration. Candidate vaccines are chemically modified or unmodified. Rabbit serum is tested for microneutralization (see Example 14). Rabbits are then challenged with $\sim 1 \text{ LD}_{90}$ of betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1) on week 7 via an IN, IM, ID or IV route of administration. Endpoint is day 13 post infection, death or euthanasia. Animals displaying severe illness as determined by $>30\%$ weight loss, extreme lethargy or paralysis are euthanized. Body temperature and weight are assessed and recorded daily.

Example 22: Microneutralization Assay

(558) Nine serial 2-fold dilutions (1:50-1:12,800) of rabbit serum are made in 50 μl virus growth medium (VGM) with trypsin in 96 well microtiter plates. Fifty microliters of virus containing ~ 50 pfu of betacoronavirus (e.g., MERS-CoV, SARS-CoV, HCoV—OC43, HCoV-229E, HCoV-NL63, HCoV-NL, HCoV—NH or HCoV—HKU1) is added to the serum dilutions and allowed to incubate for 60 minutes at room temperature (RT). Positive control wells of virus without sera and negative control wells without virus or sera are included in triplicate on each plate. While the serum-virus mixtures incubate, a single cell suspension of Madin-Darby Canine-Kidney cells are prepared by trypsinizing (Gibco 0.5% bovine pancrease trypsin in EDTA) a confluent monolayer

and suspended cells are transferred to a 50 ml centrifuge tube, topped with sterile PBS and gently mixed. The cells are then pelleted at 200 g for 5 minutes, supernatant aspirated and cells resuspended in PBS. This procedure is repeated once and the cells are resuspended at a concentration of 3×10^5 /ml in VGM with porcine trypsin. Then, 100 μ l of cells are added to the serum-virus mixtures and the plates incubated at 35° C. in CO.sub.2 for 5 days. The plates are fixed with 80% acetone in phosphate buffered saline (PBS) for 15 minutes at RT, air dried and then blocked for 30 minutes containing PBS with 0.5% gelatin and 2% FCS. An antibody to the S proteins, S1 protein or S2 protein is diluted in PBS with 0.5% gelatin/2% FCS/0.5% Tween 20 and incubated at RT for 2 hours. Wells are washed and horseradish peroxidase-conjugated goat anti-mouse IgG added, followed by another 2 hour incubation. After washing, O-phenylenediamine dihydrochloride is added and the neutralization titer is defined as the titer of serum that reduced color development by 50% compared to the positive control wells.

Example 23: MERS CoV Vaccine Immunogenicity Study in Mice

(559) The instant study was designed to test the immunogenicity in mice of candidate MERS-CoV vaccines comprising a mRNA polynucleotide encoding the full-length Spike (S) protein, or the S2 subunit (S2) of the Spike protein obtained from MERS-CoV.

(560) Mice were vaccinated with a 10 μ g dose of MERS-CoV mRNA vaccine encoding either the full-length MERS-CoV Spike (S) protein, or the S2 subunit (S2) of the Spike protein on days 0 and 21. Sera were collected from each mice on days 0, 21, 42, and 56. Individual bleeds were tested for anti-S, anti-S2 activity via a virus neutralization assay from all four time points.

(561) As shown in FIG. 17, the MERS-CoV vaccine encoding the full-length S protein induced strong immune response after the boost dose on day 21. Further, full-length S protein vaccine generated much higher neutralizing antibody titers as compared to S2 alone (FIG. 18).

Example 24: MERS CoV Vaccine Immunogenicity Study in New Zealand White Rabbits

(562) The instant study was designed to test the immunogenicity of candidate MERS-CoV mRNA vaccines encoding the full-length Spike (S) protein. The New Zealand white rabbits used in this study weighed about 4-5 kg. The rabbits were divided into three groups (Group 1a, Group 1b, and Group 2, n=8). Rabbits in Group 1a were immunized intramuscularly (IM) with one 20 μ g dose of the MERS-CoV mRNA vaccine encoding the full-length Spike protein on day 0. Rabbits in Group 1b were immunized intramuscularly (IM) with one 20 μ g dose of the MERS-CoV mRNA vaccine encoding the full-length Spike protein on day 0, and again on day 21 (booster dose). Group 2 received placebo (PBS). The immunized rabbits were then challenged and samples were collected 4 days after challenge. The viral loads in the lungs, bronchoalveolar lavage (Bal), nose, and throat of the rabbits were determined, e.g., via quantitative PCR. Replicating virus in the lung tissues of the rabbits were also detected. Lung histopathology were evaluated and the neutralizing antibody titers in serum samples of the rabbits were determined.

(563) Two 20 μ g doses of MERS-CoV mRNA vaccine resulted in a 3 log reduction of viral load in the nose and led to complete protection in the throat of the New Zealand white rabbits (FIG. 19A). Two 20 μ g doses of MERS-CoV mRNA vaccine also resulted in a 4 log reduction of viral load in the BAL of the New Zealand white rabbits (FIG. 19B). One 20 μ g dose of MERS-CoV mRNA vaccine resulted in a 2 log reduction of viral load, while two 20 μ g doses of MERS-CoV mRNA vaccine resulted in an over 4 log reduction of viral load in the lungs of the New Zealand white rabbits (FIG. 19C).

(564) Quantitative PCR results show that two 20 μ g doses of MERS-CoV mRNA vaccine reduced over 99% (2 log) of viruses in the lungs of New Zealand white rabbits (FIG. 20A). No replicating virus were detected in the lungs (FIG. 20B).

(565) Further, as shown in FIG. 21, two 20 μ g doses of MERS-CoV mRNA vaccine induced significant amount of neutralizing antibodies against MERS-CoV (EC.sub.50 between 500-1000). The MERS-CoV mRNA vaccine induced antibody titer is 3-5 fold better than any other vaccines tested in the same model.

Example 25: Immunogenicity Study

(566) The instant study is designed to test the immunogenicity in mice of candidate MeV vaccines comprising a mRNA polynucleotide encoding MeV hemagglutinin (HA) protein, MeV Fusion (F) protein or a combination of both.

(567) Mice are immunized intravenously (IV), intramuscularly (IM), or intradermally (ID) with candidate vaccines. Up to three immunizations are given at 3-week intervals (i.e., at weeks 0, 3, 6, and 9), and sera are collected after each immunization until weeks 33-51. Serum antibody titers against MeV HA protein or MeV F protein are determined by ELISA.

Example 26: MeV Rodent Challenge

(568) The instant study is designed to test the efficacy in transgenic mice of candidate MeV vaccines against a lethal challenge using a MeV vaccine comprising mRNA encoding MeV HA protein or MeV F protein. The transgenic mice express human receptor CD46 or signaling lymphocyte activation molecule (SLAM) (also referred to as CD150). Humans are the only natural host for MeV infection, thus transgenic lines are required for this study. CD46 is a complement regulatory protein that protects host tissue from complement deposition by binding to complement components C3b and C4b. Its expression on murine fibroblast and lymphoid cell lines renders these otherwise refractory cells permissive for MeV infection, and the expression of CD46 on primate cells parallels the clinical tropism of MeV infection in humans and nonhuman primates (Rall G F et al. *PNAS USA* 1997; 94(9):4659-63). SLAM is a type 1 membrane glycoprotein belonging to the immunoglobulin superfamily. It is expressed on the surface of activated lymphocytes, macrophages, and dendritic cells and is thought to play an important role in lymphocyte signaling. SLAM is a receptor for both wild-type and vaccine MeV strains (Sellin C I et al. *J Virol.* 2006; 80(13):6420-29).

(569) CD46 or SLAM/CD150 transgenic mice are challenged with a lethal dose of the MeV. Animals are immunized intravenously (IV), intramuscularly (IM), or intradermally (ID) at week 0 and week 3 with candidate MeV vaccines with and without adjuvant. The animals are then challenged with a lethal dose of MeV on week 7 via IV, IM or ID. Endpoint is day 13 post infection, death or euthanasia. Animals displaying severe illness as determined by >30% weight loss, extreme lethargy or paralysis are euthanized. Body temperature and weight are assessed and recorded daily.

(570) In experiments where a lipid nanoparticle (LNP) formulation is used, the formulation may include a cationic lipid, non-cationic lipid, PEG lipid and structural lipid in the ratios 50:10:1.5:38.5. The cationic lipid is DLin-KC2-DMA (50 mol %), the non-cationic lipid is DSPC (10 mol %), the PEG lipid is PEG-DOMG (1.5 mol %) and the structural lipid is cholesterol (38.5 mol %), for example.

(571) TABLE-US-00002 TABLE 1 hMPV Immunogenicity studies bleeding schedule Animal groups Day (n = 8) vaccine -2 0 7 14 21 28 35 56 Placebo Group PBS Pre- Prime Bleeds Bleeds Bleeds/ Bleeds Bleeds Bleeds Harvest 1 (n = 8) (IM) Bleed Boost Splens/ 10 µg Group 10 µg Terminal Dose 2 (n = 8) (IM) Bleeds 2 µg Group 2 µg Dose 3 (n = 8) (IM) Total n = 24

(572) Each of the sequences described herein encompasses a chemically modified sequence or an unmodified sequence which includes no nucleotide modifications.

(573) TABLE-US-00003 TABLE 2 SEQ ID Description Sequence NO: hMPV Nucleic Acid Sequences gi|122891979|gb|

ATGAGCTGGAAGGTGGTGATTATCTTCAGCCTGCTGATTA 1 EF051124.1| Human CACCTCAACACGGCCTGAAGGAGAGCTACCTGGAAGAGA metapneumovirus GCTGCTCCACCATCACCGAGGGCTACCTGAGCGTGCTGC isolate TN/92-4 GGACCGGCTGGTACACCAACGTGTTACCCCTGGAGGTGG fusion protein gene, GCGACGTGGAGAACCTGACCTGCAGCGACGGCCCTAGCC complete genome TGATCAAGACCGAGCTGGACCTGACCAAGAGCGCTCTGA GAGAGCTGAAGACCGTGTCCGCCGACCAGCTGGCCAGAG

AGGAACAGACCTCGGCAGAGCAGATTTCGTGC
TGGGCGCCATCGCTCTGGGAGTCGCCGCTGCCGCTGCAG
TGACAGCTGGAGTGGCCATTGCTAAGACCATCAGACTGG
AAAGCGAGGTGACAGCCATCAACAATGCCCTGAAGAAG
ACCAACGAGGCCGTGAGCACCTGGGCAATGGAGTGAGA
GTGCTGGCCACAGCCGTGCGGGAGCTGAAGGACTTCGTG
AGCAAGAACCTGACCAGAGCCATCAACAAGAACAAGTG
CGACATCGATGACCTGAAGATGGCCGTGAGCTTCTCCA
GTTCAACAGACGGTTCCTGAACGTGGTGAGACAGTTCTC
CGACAACGCTGGAATCACACCTGCCATTAGCCTGGACCT
GATGACCGACGCCGAGCTGGCTAGAGCCGTGCCCAACAT
GCCCACCAGCGCTGGCCAGATCAAGCTGATGCTGGAGAA
CAGAGCCATGGTGCGGAGAAAGGGCTTCGGCATCCTGAT
TGGGGTGTATGGAAGCTCCGTGATCTACATGGTGCAGCT
GCCCATCTTCGGCGTGATCGACACACCCTGCTGGATCGTG
AAGGCCGCTCCTAGCTGCTCCGAGAAGAAAGGAAACTAT
GCCTGTCTGCTGAGAGAGGACCAGGGCTGGTACTGCCAG
AACGCCGGAAGCACAGTGTACTATCCCAACGAGAAGGAC
TGCGAGACCAGAGGCGACCACGTGTTCTGCGACACCGCT
GCCGGAATCAACGTGGCCGAGCAGAGCAAGGAGTGCAA
CATCAACATCAGCACAACCAACTACCCCTGCAAGGTGAG
CACCGGACGGCACCCCATCAGCATGGTGGCTCTGAGCCC
TCTGGGCGCTCTGGTGGCCTGCTATAAGGGCGTGTCTGT
AGCATCGGCAGCAATCGGGTGGGCATCATCAAGCAGCTG
AACAAGGGATGCTCCTACATCACCAACCAGGACGCCGAC
ACCGTGACCATCGACAACACCGTGTACCAGCTGAGCAAG
GTGGAGGGCGAGCAGCACGTGATCAAGGGCAGACCCGT
GAGCTCCAGCTTCGACCCCATCAAGTTCCCTGAGGACCA
GTTCAACGTGGCCCTGGACCAGGTGTTTGAGAACATCGA
GAACAGCCAGGCCCTGGTGGACCAGAGCAACAGAATCCT
GTCCAGCGCTGAGAAGGGCAACACCGGCTTCATCATTGT
GATCATTCTGATCGCCGTGCTGGGCAGCTCCATGATCCTG
GTGAGCATCTTCATCATTATCAAGAAGACCAAGAAACC
ACCGGAGCCCCTCCTGAGCTGAGCGGCGTGACCAACAAT
GGCTTCATTCCCCACAACCTGA gb|AY525843.1|:

ATGTCTTGGAAAGTGATGATCATCATTTTCGTTACTCATAA 2 3065-4684 Human
CACCCCAGCACGGGCTAAAGGAGAGTTATTTGGAAGAAT metapneumovirus
CATGTAGTACTATAACTGAGGGATACCTCAGTGTTTTAAG isolate NL/1/99,
AACAGGCTGGTACACTAATGTCTTCACATTAGAAGTTGGT complete genome
GATGTTGAAAATCTTACATGTACTGATGGACCTAGCTTAA
TCAAAACAGAACTTGATCTAACAAAAAGTGCTTTAAGGG
AACTCAAAACAGTCTCTGCTGATCAGTTGGCGAGAGAGG
AGCAAATTGAAAATCCCAGACAATCAAGATTTGTCTTAG
GTGCGATAGCTCTCGGAGTTGCTACAGCAGCAGCAGTCA
CAGCAGGCATTGCAATAGCCAAAACCATAAGGCTTGAGA
GTGAGGTGAATGCAATTAAAGGTGCTCTCAAACAACTA
ATGAAGCAGTATCCACATTAGGGAATGGTGTGCGGGTCC
TAGCCACTGCAGTGAGAGAGCTAAAAGAATTTGTGAGCA
AAAACCTGACTAGTGCAATCAACAGGAACAAATGTGACA
TTGCTGATCTGAAGATGGCTGTCAGCTTCAGTCAATTCAA

CAGAGATTCTTCTGTTGTCGGCAGTTTTTCAGACAAT
GCAGGGATAACACCAGCAATATCATTGGACCTGATGACT
GATGCTGAGTTGGCCAGAGCTGTATCATACATGCCAACA
TCTGCAGGGCAGATAAAACTGATGTTGGAGAACCGCGCA
ATGGTAAGGAGAAAAGGATTTGGAATCCTGATAGGGGTC
TACGGAAGCTCTGTGATTTACATGGTTCAATTGCCGATCT
TTGGTGTCTAGATACACCTTGTTGGATCATCAAGGCAGC
TCCCTCTTGCTCAGAAAAAACGGGAATTATGCTTGCCTC
CTAAGAGAGGATCAAGGGTGGTATTGTAAAAATGCAGGA
TCTACTGTTTACTACCCAAATGAAAAAGACTGCGAAACA
AGAGGTGATCATGTTTTTTGTGACACAGCAGCAGGGATC
AATGTTGCTGAGCAATCAAGAGAATGCAACATCAACATA
TCTACTACCAACTACCCATGCAAAGTCAGCACAGGAAGA
CACCTATAAGCATGGTTGCACTATCACCTCTCGGTGCTT
TGGTGGCTTGCTATAAAGGGGTAAAGCTGCTCGATTGGCA GCAATTGGGT
TGGAATCATCAAACAATTACCCAAAGGCTGCTCATACAT
AACCAACCAGGATGCAGACACTGTAACAATTGACAATAC
CGTGTATCAACTAAGCAAAGTTGAAGGTGAACAGCATGT
AATAAAAGGGGAGACCAGTTTCAAGCAGTTTTTGATCCAAT
CAAGTTTCCTGAGGATCAGTTCAATGTTGCGCTTGATCAA
GTCTTCGAAAGCATTGAGAACAGTCAGGCACTAGTGGAC
CAGTCAAACAAAATTCTAAACAGTGCAGAAAAAGGAAA
CACTGGTTTTATTATCGTAGTAATTTTGGTTGCTGTTCTTG
GTCTAACCATGATTTCAGTGAGCATCATCATATAATCAA
GAAAACAAGGAAGCCACAGGAGCACCTCCAGAGCTGA
ATGGTGTACCAACGGCGGTTTTCATACCACATAGTTA gb|KJ627414.1|:
ATGTCTTGGAAGTGATGATTATCATTTTCGTTACTCATAA 3 3015-4634 Human
CACCTCAGCATGGACTAAAAGAAAGTTATTTAGAAGAAT metapneumovirus
CATGTAGTACTATAACTGAAGGATATCTCAGTGTTTTAAG strain hMPV/*Homo*
AACAGGTTGGTACACCAATGTCTTTACATTAGAAGTTGGT *sapiens*/PER/
GATGTTGAAAATCTTACATGTACTGATGGACCTAGCTTAA CFI0497/2010/B,
TCAAAACAGAACTTGACCTAACCAAAAGTGCTTTAAGAG complete genome
AACTCAAAACAGTTTCTGCTGATCAGTTAGCGAGAGAAG
AACAAATTGAAAATCCCAGACAATCAAGGTTTGTCTTAG
GTGCAATAGCTCTTGAGTTGCCACAGCAGCAGCAGTCA
CAGCAGGCATTGCAATAGCCAAAACCTATAAGGCTTGAGA
GTGAAGTGAATGCAATCAAAGGTGCTCTCAAAACAACCA
ATGAGGCAGTATCAACACTAGGAAATGGAGTGCGGGTCC
TAGCCACTGCAGTAAGAGAGCTGAAAGAATTTGTGAGCA
AAAACCTGACTAGTGCGATCAACAAGAACAAGTGTGACA
TTGCTGATTTGAAGATGGCTGTCAGCTTCAGTCAGTTCAA
CAGAAGATTCCTAAATGTTGTGCGGCAGTTTTTCAGACAAT
GCAGGGATAACACCAGCAATATCATTGGACCTGATGAAT
GATGCTGAGCTGGCCAGAGCTGTATCATACATGCCAACA
TCTGCAGGACAGATAAAACTAATGTTAGAGAACCGTGCA
ATGGTGAGGAGAAAAGGATTTGGAATCTTGATAGGGGTC
TACGGAAGCTCTGTGATTTACATGGTCCAGCTGCCGATCT
TTGGTGTCTATAAATACACCTTGTTGGATAATCAAGGCAGC
TCCCTCTTGTTTCAGAAAAAGATGGAAATTATGCTTGCCTC
CTAAGAGAGGATCAAGGGTGGTATTGTAAAAATGCAGGA

TCCACTGTTTACTGAAATAAGACTGCGAAACA
AGAGGTGATCATGTTTTTTGTGACACAGCAGCAGGGATC
AATGTTGCTGAGCAATCAAGAGAATGCAACATCAACATA
TCTACCACCAACTACCCATGCAAAGTCAGCACAGGAAGA
CACCTATCAGCATGGTTGCACTATCACCTCTCGGTGCTT
TGGTAGCTTGCTACAAAGGGGTTAGCTGCTCGACTGGCA
GTAATCAGGTTGGAATAATCAAACAACCTAAAGGCT
GCTCATACATAACTAACCAGGACGCAGACACTGTAACAA
TTGACAACACTGTGTATCAACTAAGCAAAGTTGAGGGTG
AACAGCATGTAATAAAAAGGGAGACCAGTTTCAAGCAGTT
TTGATCCAATCAGGTTTCCTGAGGATCAGTTCAATGTTGC
GCTTGATCAAGTCTTTGAAAGCATTGAAAACAGTCAAGC
ACTAGTGGACCAGTCAAACAAAATTCTGAACAGTGCAGA AAAAGGAAACACTGGT
TTCATTATTGTAATAATTTTGATTGCTGTTCTTGGGTAAAC
CATGATTTTCAGTGAGCATCATCATATAATCAAAAAAAC
AAGGAAGCCCACAGGGGCACCTCCGGAGCTGAATGGTGT
TACCAACGGCGGTTTCATACCGCATAGTTAG gb|KJ723483.1|:
ATGGAGTTGCCAATCCTCAAAACAAATGCAATTACCACA 4 5586-7310 Human
ATCCTTGCTGCAGTCACACTCTGTTTCGCTTCCAGTCAAA respiratory
ACATCACTGAAGAATTTTATCAATCAACATGCAGTGCAG syncytial virus
TTAGCAAAGGCTATCTTAGTGCTCTAAGAACTGGTTGGTA strain RSVA/Homo
TACTAGTGTTATAACTATAGAATTAAGTAATATCAAGGA *sapiens*/USA/84I-
AAATAAGTGTAATGGAACAGATGCTAAGGTAAAATTGAT 215A-01/1984,
AAAACAAGAATTAGATAAATATAAAAATGCTGTAACAGA complete genome
ATTGCAGTTGCTCATGCAAAGCACACCAGCAGCCAACAA
TCGAGCCAGAAGAGAACTACCAAGGTTTATGAATTATAC
ACTCAATAATACCAAAAATACCAATGTAACATTAAGCAA
GAAAAGGAAAAGAAGATTTCTTGGCTTTTTGTAGGTGTT
GGATCTGCAATCGCCAGTGGCATTGCTGTATCTAAGGTCC
TGCACCTAGAAGGGGAAGTGAACAAAATCAAAAGTGCTC
TACTATCCACAAACAAGGCTGTAGTCAGCTTATCAAATG
GAGTTAGTGTCTTAACCAGCAAAGTGTTAGACCTCAAAA
ACTATATAGATAAACAGTTGTTACCTATTGTGAACAAGC
AAAGCTGCAGCATATCAAACATTGAAACTGTGATAGAGT
TCCAACAAAAGAACAACAGACTACTAGAGATTACCAGGG
AATTTAGTGTTAATGCAGGTGTAACCTACACCTGTAAGCAC
TTATATGTTAACTAATAGTGAATTATTATCATTAATCAAT
GATATGCCTATAACAAATGATCAGAAAAAGTTAATGTCC
AACAATGTTCAAATAGTTAGACAGCAAAGTTACTCTATC
ATGTCCATAATAAAGGAGGAAGTCTTAGCATATGTAGTA
CAATTACCACTATATGGTGTAATAGATACACCCTGTTGGA
AACTGCACACATCCCCTCTATGTACAACCAACACAAAGG
AAGGGTCCAACATCTGCTTAACAAGAACCGACAGAGGAT
GGTATTGTGACAATGCAGGATCAGTATCTTTCTTCCCACA
AGCTGAAACATGTAAAGTTCAATCGAATCGGGTATTTTGT
GACACAATGAACAGTTTAAACATTACCAAGTGAAGTAAAT
CTCTGCAACATTGACATATTCAACCCCAAATATGATTGCA
AAATTATGACTTCAAAAACAGATGTAAGCAGCTCCGTTA
TCACATCTCTAGGAGCCATTGTGTCATGCTATGGCAAAAC
TAAATGTACAGCATCCAATAAAAATCGTGGGATCATAAA

GACATTCTTTCTGGTGGTATTATGTATCAAATAAGGG
GGTGGATACTGTGTCTGTAGGTAATACATTATATTATGTA
AATAAGCAAGAAGGCCAAAAGTCTCTATGTAAAAGGTGAA
CCAATAATAAATTTCTATGACCCATTAGTGTTCCCCTCTG
ATGAATTTGATGCATCAATATCTCAAGTCAATGAGAAGA
TTAACCAGAGCCTAGCATTATTTCGTAAATCCGATGAATT
ATTACATAATGTAAATGCTGGTAAATCCACCACAAATAT
CATGATAACTACTATAATTATAGTGATTATAGTAATATTG
TTATCATTAATTGCAGTTGGACTGCTCCTATACTGCAAGG
CCAGAAGCACACCAGTCACACTAAGTAAGGATCAACTGA
GTGGTATAAATAATATTGCATTTAGTAACTGA hMPV mRNA Sequences gi|122891979|gb|
AUGAGCUGGAAGGUGGUGAUUAUCUUCAGCCUGCUGAU 57 EF051124.1| Human
UACACCUCAACACGGCCUGAAGGAGAGCUACCUGGAAG metapneumovirus
AGAGCUGCUCCACCAUCACCGAGGGCUACCUGAGCGUG isolate TN/92-4
CUGCGGACCGGCUGGUACACCAACGUGUUCACCCUGGA fusion protein gene,
GGUGGGCGACGUGGAGAACCUGACCUGCAGCGACGGCC complete genome
CUAGCCUGAUCAAGACCGAGCUGGACCUGACCAAGAGC
GCUCUGAGAGAGCUGAAGACCGUGUCCGCCGACCAGCU
GGCCAGAGAGGAACAGAUUCGAGAACCCUCGGCAGAGCA
GAUUCGUGCUGGGCGCCAUCGCUCUGGGAGUCGCCGCU
GCCGCUGCAGUGACAGCUGGAGUGGCCAUUGCUAAGAC
CAUCAGACUGGAAAGCGAGGUGACAGCCAUCAACAAUG
CCCUGAAGAAGACCAACGAGGCCGUGAGCACCCUGGGC
AAUGGAGUGAGAGUGCUGGCCACAGCCGUGCGGGAGCU
GAAGGACUUCGUGAGCAAGAACCUGACCAGAGCCAUCA
ACAAGAACAAGUGCGACAUCGAUGACCUGAAGAUGGCC
GUGAGCUUCUCCCAGUUCAACAGACGGUUCCUGAACGU
GGUGAGACAGUUCUCCGACAACGCUGGAAUCACACCUG
CCAUUAGCCUGGACCUGAUGACCGACGCCGAGCUGGCU
AGAGCCGUGCCCAACAUGCCCACCAGCGCUGGCCAGAU
CAAGCUGAUGCUGGAGAACAGAGCCAUGGUGCGGAGAA
AGGGCUUCCGCAUCCUGAUUUGGGGUGUAUGGAAGCUCC
GUGAUCUACAUGGUGCAGCUGCCCAUCUUCGGCGUGAU
CGACACACCCUGCUGGAUCGUGAAGGCCGCUCCUAGCU
GCUCCGAGAAGAAAGGAAACUAUGCCUGUCUGCUGAGA
GAGGACCAGGGCUGGUACUGCCAGAACGCCGGAAGCAC
AGUGUACUAUCCCAACGAGAAGGACUGCGAGACCAGAG
GCGACCACGUGUUCUGCGACACCGCUGCCGGAUCAAAC
GUGGCCGAGCAGAGCAAGGAGUGCAACAUCAACAUCAG
CACAACCAACUACCCUGCAAGGUGAGCACCGGACGGC
ACCCCAUCAGCAUGGUGGCUCUGAGCCUCUGGGCGCU
CUGGUGGCCUGCUAUAAGGGCGUGUCCUGUAGCAUCGG
CAGCAAUCGGGUGGGCAUCAUAAGCAGCUGAACAAGG
GAUGCUCUACAUCACCAACCAGGACGCCGACACCGUG
ACCAUCGACAACACCGUGUACCAGCUGAGCAAGGUGGA
GGGCGAGCAGCAGUGAUAAGGGCAGACCCGUGAGCU
CCAGCUUCGACCCCAUCAAGUUCCUGAGGACCAGUUC
AACGUGGCCCUGGACCAGGUGUUGAGAACAUCGAGAA
CAGCCAGGCCUGGUGGACCAGAGCAACAGAAUCCUGU
CCAGCGCUGAGAAGGGCAACACCGGCUUCAUCAUUGUG

AAUCAUUCGAGCUGGCGUCCAUUGAUCCU
GGUGAGCAUCUUCAUCAUUAUCAAGAAGACCAAGAAAC
CCACCGGAGCCCCUCCUGAGCUGAGCGGCGUGACCAAC
AAUGGCUUCAUUCCCCACAACUGA gb|AY525843.1|:
AUGUCUUGGAAAGUGAUGAUCAUCAUUCGUUACUCAU 58 3065-4684 Human
AACACCCCAGCACGGGCUAAAGGAGAGUUAUUUGGAAG metapneumovirus
AAUCAUGUAGUACUAUAACUGAGGGAUACCUCAGUGUU isolate NL/1/99,
UUAAGAACAGGCUGGUACACUAAUGUCUUCACAUUAGA complete genome
AGUUGGUGAUGUUGAAAAUCUUACAUGUACUGAUGGA
CCUAGCUUAAAUCAAAACAGAACUUGAUCUAACAAAAAG
UGC UUUAAGGGAACUCAAAACAGUCUCUGCUGAUCAGU
UGGCGAGAGAGGAGCAAAUUGAAAAUCCCAGACAAUCA
AGAUUUGUCUUAAGGUGCGAUAGCUCUCGGAGUUGCUAC
AGCAGCAGCAGUCACAGCAGGCAUUGCAAUAGCCAAAA
CCAUAAGGCUUGAGAGUGAGGUGAAUGCAAUUAAGG
UGCUCUCAAAACAAACUAAUGAAGCAGUAUCCACAUUAG
GGAAUGGUGUGCGGGUCCUAGCCACUGCAGUGAGAGAG
CUAAAAGAAUUUGUGAGCAAAAACCUGACUAGUGCAAU
CAACAGGAACAAAUUGUGACAUUGCUGAUCUGAAGAUGG
CUGUCAGCUUCAGUCAAUUCAACAGAAGAUUUCUAAAU
GUUGUGCGGCAGUUUUCAGACAAUGCAGGGAUAAACACC
AGCAAUAUCAUUGGACCUGAUGACUGAUGCUGAGUUGG
CCAGAGCUGUAUCAUACAUGCCAACAUCUGCAGGGCAG
AUAAAACUGAUGUUGGAGAACCGCGCAAUGGUAAGGAG
AAAAGGAUUUGGAAUCCUGAUAGGGGUCUACGGAAGCU
CUGUGAUUUACAUGGUUCAAUUGCCGAUCUUUGGUGUC
AUAGAUACACCUUGUUGGAUCAUCAAGGCAGCUCUCCUC
UUGCUCAGAAAAAAACGGGAAUUUAUGCUUGCCUCCUAA
GAGAGGAUCAAGGGUGGUUAUUUGUAAAAAUGCAGGAUC
UACUGUUUACUACCCAAAUUGAAAAAGACUGCGAAACAA
GAGGUGAUCAUGUUUUUGUGACACAGCAGCAGGGAUUC
AAUGUUGCUGAGCAAUCAAGAGAAUGCAACAUCAACAU
AUCUACUACCAACUACCCAUGCAAAGUCAGCACAGGAA
GACACCCUAUAAGCAUGGUUGCACUAUCACCUCUCGGU
GCUUUGGUGGCUUGCUAUAAAGGGGUAAGCUGCUCGAU UGGCAGCAAUUGGGU
UGGAAUCAUCAAAACAAUUACCCAAAGGCUGCUCAUACA
UAACCAACCAGGAUGCAGACACUGUAACAAUUGACAAU
ACCGUGUAUCAACUAAGCAAAGUUGAAGGUGAACAGCA
UGUAAUAAAAGGGAGACCAGUUUCAAGCAGUUUUGAUC
CAAUCAAGUUUCCUGAGGAUCAGUUCAAUGUUGCGCUU
GAUCAAGUCUUCGAAAGCAUUGAGAACAGUCAGGCACU
AGUGGACCAGUCAAAACAAAAUUCUAAACAGUGCAGAAA
AAGGAAACACUGGUUUCAUUAUCGUAGUAAUUUUGGU
UGCUGUUCUUGGUCUAACCAUGAUUUUCAGUGAGCAUCA
UCAUCAUAAUCAAGAAAACAAGGAAGCCCACAGGAGCA
CCUCCAGAGCUGAAUGGUGUCACCAACGGCGGUUUCAU ACCACAUAAGUUAG
gb|KJ627414.1|: AUGUCUUGGAAAGUGAUGAUUAUCAUUCGUUACUCAU 59 3015-
4634 Human AACACCUCAGCAUGGACUAAAAGAAAGUUAUUUAGAAG
metapneumovirus AAUCAUGUAGUACUAUAACUGAAGGAUAUCUCAGUGUU strain
hMPV/*Homo* UUAAGAACAGGUUGGUACACCAAUGUCUUUACAUAUAGA *sapiens*/PER/

AGUUGUGUAGUAGUUAACUUGAUCUGAUGGA CF10497/2010/B,
CCUAGCUUAAUCAAACAGAACUUGACCUAACCAAAAG complete genome
UGC UUAAGAGAACUCAAACAGUUUCUGCUGAUCAGU
UAGCGAGAGAAGAACAAAUUGAAAAUCCCAGACAAUCA
AGGUUUGUCCUAGGUGCAAUAGCUCUUGGAGUUGCCAC
AGCAGCAGCAGUCACAGCAGGCAUUGCAAUAGCCAAAA
CUAUAAAGGCUUGAGAGUGAAGUGAAUGCAAUCAAAAGG
UGCUCUCAAAACAACCAAUGAGGCAGUAUCAACACUAG
GAAAUUGGAGUGCGGGUCCUAGCCACUGCAGUAAGAGAG
CUGAAAGAAUUUGUGAGCAAAAACCUGACUAGUGCGAU
CAACAAGAACAAGUGUGACA UUGCUGAUUUGAAGAUGG
CUGUCAGCUUCAGUCAGUUCAACAGAAGAUUCCUAAAU
GUUGUGCGGCAGUUUUCAGACAAUGCAGGGAUAACACC
AGCAAUAUCAUUGGACCUGAUGAAUGAUGCUGAGCUGG
CCAGAGCUGUAUCAUACAUGCCAACAUCUGCAGGACAG
AUAAAACUAAUGUAGAGAACCGUGCAAUGGUGAGGA
GAAAAGGAUUUGGAAUCUUGAUAGGGGUCUACGGAAG
CUCUGUGAUUUACAUGGUCCAGCUGCCGAUCUUUGGUG
UCAUAAAUACACCUUGUUGGAUAAUCAAGGCAGCUCCC
UCUUGUUCAGAAAAAGAUGGAAAUUAUGCUUGCCUCCU
AAGAGAGGAUCAAGGGUGGUAUUGUAAAAAUGCAGGA
UCCACUGUUUACUACCCAAAUGAAAAAGACUGCGAAAC
AAGAGGUGAUCAUGUUUUUUGUGACACAGCAGCAGGGA
UCAAUGUUGCUGAGCAAUCAAGAGAAUGCAACAUCAAC
AUAUUCUACCACCAACUACCCAUGCAAAGUCAGCACAGG
AAGACACCCUAUCAGCAUGGUUGCACUAUCACCUCUCG
GUGCUUUGGUAGCUUGCUACAAAGGGGUUAGCUGCUCG
ACUGGCAGUAAUCAGGUUGGAAUAAUCAAAACAACUACC
UAAAGGCUGCUCAUACAUAACUAACCAGGACGCAGACA
CUGUAACAAUUGACAACACUGUGUAUCAACUAAGCAAA
GUUGAGGGUGAACAGCAUGUAAUAAAAGGGAGACCAG
UUUCAAGCAGUUUUGAUCCAUCAGGUUUCUGAGGAU
CAGUUCA AUGUUGCGCUUGAUCAAGUCUUUGAAAGCAU
UGAAAACAGUCAAGCACUAGUGGACCAGUCAAAACAAA
UUCUGAACAGUGCAGAAAAAGGAAACACUGGU
UUCAUUAUUGUAAUAAUUUUGAUUUGCUGUUCUUGGGU
UAACCAUGAUUUCAGUGAGCAUCAUCAUCAAUCAA
AAAACAAGGAAGCCCACAGGGGCACCUCCGGAGCUGAA
UGGUGUUACCAACGGCGGUUCAUACCGCAUAGUUAG gb|KJ723483.1|:
AUGGAGUUGCCAAUCCUCAAACAAAUGCAAUUAACCAC 60 5586-7310 Human
AAUCCUUGCUGCAGUCACACUCUGUUUCGCUUCCAGUC respiratory
AAAACAUCACUGAAGAAUUUUAUCAAUCAACAUGCAGU syncytial virus
GCAGUUAGCAAAGGCUAUCUAGUGCUCUAAGAACUGG strain RSVA/Homo
UUGGUUAUACUAGUGUUUAUAAUUAAGUAAU sapiens/USA/84I-
AUCAAGGAAAAUAAGUGUAAUGGAACAGAUGC UAAGG 215A-01/1984,
UAAAAUUGAUAAAACAAGAAUUAAGAUAAAUAUAAAAA complete genome
UGCUGUAACAGAAUUGCAGUUGCUC AUGCAAAGCACAC
CAGCAGCCAACAUCGAGCCAGAAGAGAACUACCAAGG
UUUAUGAAUUAUACACUCAAUAAUACCAAAAAUACCAA
UGUAACAUAAGCAAGAAAAGGAAAAGAAGAUUUCU

GGCUUUUUUUGGUAUCGCGCCAGUGG
CAUUGCUGUAUCUAAGGUCCUGCACCUAGAAGGGGAAG
UGAACAAAAUCAAAAGUGCUCUACUAUCCACAAACAAG
GCUGUAGUCAGCUUAUCAA AUGGAGUUAGUGUCUUAAC
CAGCAAAGUGUUAGACCUCAAAAACUAUAUAGAUAAAC
AGUUGUUACCUAUUGUGAACAAGCAAAGCUGCAGCAUA
UCAAACAUUGAAACUGUGAUAGAGUUCCAACAAAAGAA
CAACAGACUACUAGAGAUUACCAGGGAAUUUAGUGUUA
AUGCAGGUGUAACUACACCUGUAAGCACUUAUAUGUUA
ACUAAUAGUGAAUUAUUAUCAUUAUCAAUGAUUAGCC
UAUAACAAAUGAUCAGAAAAAGUUAAUGUCCAACAAG
UUCAAAUAGUUAGACAGCAAAGUUACUCUAUCAUGUCC
AUAUAAAAGGAGGAAGUCUUAAGCAUAUGUAGUACAAU
UACCACUAUAUGGUGUAAUAGAUACACCCUGUUGGAAA
CUGCACACAUCCCCUCUAUGUACAACCAACACAAAGGA
AGGGUCCAACAUCUGCUUAACAAGAACCGACAGAGGAU
GGUAUUGUGACAAUGCAGGAUCAGUAUCUUUCUCCCCA
CAAGCUGAAACAUGUAAAGUUCAAUCGAAUCGGGUAAU
UUGUGACACAAUGAACAGUUUAACAUAUACCAAGUGAAG
UAAAUCUCUGCAACAUAUGACAUAUUAACCCCAAUAU
GAUUGCAAAAUUAUGACUUCAAAAACAGAUGUAAGCAG
CUCCGUUAUCACAUCUCUAGGAGCCAUAUGUGUCAUGCU
AUGGCAAAACUAAAUGUACAGCAUCCAUAUAAAAAUCGU
GGGAUCAUAAAGACAUAUUUCUAACGGGUGUGAUUAUG
UAUCAAAUAAGGGGGUGGAUACUGUGUCUGUAGGUAA
UACAUUAUAUUAUGUAAAUAAGCAAGAAGGCAAAAGU
CUCUAUGUAAAAGGUGAACCAUAUAUAAAUUUCUAUGA
CCCAUUAGUGUUCUUUCUGAUGAAUUUGAUGCAUCAA
UAUCUCAAGUCAAUUGAGAAGAUUAACCAGAGCCUAGCA
UUUAUUCGUAAAUCCGAUGAAUUAUUAACAUAUUGUAA
AUGCUGGUAAAUCCACCACAAAUAUCAUGAUAAACUACU
AUAAUUAUAGUGAUUAUAGUAAUAUUGUUAUCAUUA
UUGCAGUUGGACUGCUCCUAUACUGCAAGGCCAGAAGC
ACACCAGUCACACUAAGUAAGGAUCAACUGAGUGGUAAU
AAAUAAUAUUGCAUUUAGUAACUGA

(574) TABLE-US-00004 TABLE 3 hMPV Amino Acid Sequences SEQ ID Description
Sequence NO: gi|122891979|gb|

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGW 5 EF051124.1| Human
YTNVFTLEVGDVENLTCSDGPSLIKTELDTLSALRELKTVS metapneumovirus
ADQLAREEQIENPRQSRFVLGAIALGVAAAAAVTAGVAIAK isolate TN/92-4
TIRLESEVTAINNALKKTNEAVSTLGNGVRVLATAVRELKD fusion protein gene,
FVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS complete cds
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRA
MVRKRGFGILIGVYGSSVIYMVQLPIFGVIDTPCWIVKAAPS
CSEKKGN YACLLREDQGWYCQNAGSTVYYPNEKDCETRG
DHVFCDTAAGINVAEQSKECNINISTTNYPCKVSTGRHPISM
VALSPLGALVACYKGVSCSIGSNRVGIIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQF
NVALDQVFENIENSQALVDQSNRILSSAEKGNTGFIIIVILIAV
LGSSMILVSIFIIKKTKKPTGAPPELSGVTNNGFIPHN gb|AY525843.1|:

MSWKVMIISLLITPQHGLKESYLEESCSTITEGYLSVLRTGW 6 3065-4684 Human
YTNVFTLEVGDVENLTCTDGP SLIKTELDLT KSALRELKTVS metapneumovirus
ADQLAREEQIENPRQSRFVLGAIALGVATAAAV TAGIAIAKT isolate NL/1/99,
IRLESEVNAIKGALKQTNEAVSTLGN GVRVLATAVRELKEF complete cds
VSKNLTSAINRNKCDIADLKMAVSFSQFNRRFLNVVRQFSD
NAGITPAISLDLMTDAELARAVSYMPTSAGQIKLMLENRAM
VRRKGFGILIGVYGSSVIYMVQLPIFGVIDTPCWIIKAAPSCS
EKNGNYACLLREDQG WYCKNAGSTVYYPNEKDCETR GDH
VFCDTAAGINVAEQSREC NINISTTNYPCKVSTGRHPISMVA
LSPLGALVACYKGVSCSIGSNWVGIIKQLPKGCSYITNQDAD
TVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFNV
ALDQVFESIENSQALVDQSNKILNSAEKGNTGFIIVVILVAVL
GLTMISVSIIIIKKTRKPTGAPPELNGVTNGGFIPHS gb|KJ627414.1|:

MSWKVMIISLLITPQHGLKESYLEESCSTITEGYLSVLRTGW 7 3015-4634 Human
YTNVFTLEVGDVENLTCTDGP SLIKTELDLT KSALRELKTVS metapneumovirus
ADQLAREEQIENPRQSRFVLGAIALGVATAAAV TAGIAIAKT strain hMPV/*Homo*
IRLESEVNAIKGALKTTNEAVSTLGN GVRVLATAVRELKEF *sapiens*/PER/
VSKNLTSAINKNKCDIADLKMAVSFSQFNRRFLNVVRQFSD CFI0497/2010/B,
NAGITPAISLDL MND AELARAVSYMPTSAGQIKLMLENRAM complete cds
VRRKGFGILIGVYGSSVIYMVQLPIFGVINTPCWIIKAAPSCS
EKDGN YACLLREDQG WYCKNAGSTVYYPNEKDCETR GDH
VFCDTAAGINVAEQSREC NINISTTNYPCKVSTGRHPISMVA
LSPLGALVACYKGVSCSTG SNQVGIIKQLPKGCSYITNQDAD
TVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIRFPEDQFNV
ALDQVFESIENSQALVDQSNKILNSAEKGNTGFIIVIIILIAVLG
LTMISVSIIIIKKTRKPTGAPPELNGVTNGGFIPHS gb|KJ723483.1|:

MELPILKTNAITTLAAVTLCFASSQNITEEFYQSTCSAVSKG 8 5586-7310 Human
YLSALRTGWYTSVITIELSNIKENKCNGTDAKV KLIKQELDK respiratory
YKNAVTELQLLMQSTPAANNRARELPRFMNYTLNNTKNT syncytial virus
NVTLSKKRKRRFLGLLG VGSIAIASGIAVSKVLHLEGEVNKI strain RSA/*Homo*
KSALLSTNKAVVSLNGVSVLTSKVLDLKNYIDKQLLPVN *sapiens*/USA/84I-
KQSCSISNIETVIEFQQKNRLL EITREFSVNAGVTTTPVSTYM 215A-01/1984,
LTNSELLSLINDMPITNDQKKLMSNNVQIVRQQSYSIMSIIKE complete cds
EVLAYVVQLPLYGVIDTPCWKLHTSPLCTTNTKEGSNICLTR
TDRGWYCDNAGSVSFFPQAETCKVQSNRVFCDTMNSLTLP
SEVNL CNIDIFNPKYDCKIMTSKTDVSSSVITSLGAIVSCYGK
TKCTASNKNRGIKTF SNGCDYVSNKGVDTVSVGNTLYYVN
KQEGKS LYVKGEPIINFYDPLVFP SDEFDASISQVNEKINQSL
AFIRKSDELLHNVNAGKSTTNIMITTHIIVIVILLSLIAVGLLL
YCKARSTPVTLSKDQLSGINNI AFSN

(575) TABLE-US-00005 TABLE 4 hMPV NCBI Accession Numbers (Amino Acid Sequences)
GenBank Virus Accession F [Human metapneumovirus] [Human metapneumovirus] AEK26895.1
fusion glycoprotein [Human metapneumovirus] ACJ53565.1 fusion glycoprotein [Human
metapneumovirus] ACJ53566.1 fusion glycoprotein [Human metapneumovirus] ACJ53569.1
fusion protein [Human metapneumovirus] AEZ52347.1 fusion glycoprotein [Human
metapneumovirus] ACJ53574.1 fusion glycoprotein [Human metapneumovirus] AHV79473.1
fusion glycoprotein [Human metapneumovirus] ACJ53570.1 fusion glycoprotein [Human
metapneumovirus] ACJ53567.1 fusion protein [Human metapneumovirus] AAS22125.1 fusion
glycoprotein [Human metapneumovirus] AHV79795.1 fusion glycoprotein [Human
metapneumovirus] AHV79455.1 fusion glycoprotein [Human metapneumovirus] ACJ53568.1

fusion protein [Human metapneumovirus] AAS22109.1 fusion glycoprotein [Human metapneumovirus] AGU68417.1 fusion glycoprotein [Human metapneumovirus] AGJ74228.1 fusion glycoprotein [Human metapneumovirus] ACJ53575.1 fusion protein [Human metapneumovirus] AAU25820.1 fusion glycoprotein [Human metapneumovirus] AGU68377.1 fusion glycoprotein [Human metapneumovirus] AGU68371.1 fusion glycoprotein [Human metapneumovirus] AGJ74087.1 fusion glycoprotein [Human metapneumovirus] ACJ53560.1 fusion glycoprotein [Human metapneumovirus] AHV79858.1 fusion glycoprotein [Human metapneumovirus] ACJ53577.1 fusion protein [Human metapneumovirus] AAS22085.1 fusion protein [Human metapneumovirus] AEZ52348.1 fusion glycoprotein [Human metapneumovirus] AGJ74044.1 fusion glycoprotein [Human metapneumovirus] ACJ53563.1 fusion glycoprotein precursor [Human metapneumovirus] YP_012608.1 fusion glycoprotein [Human metapneumovirus] AGJ74053.1 fusion protein [Human metapneumovirus] BAM37562.1 fusion glycoprotein [Human metapneumovirus] ACJ53561.1 fusion glycoprotein [Human metapneumovirus] AGU68387.1 fusion [Human metapneumovirus] AGL74060.1 fusion glycoprotein precursor [Human metapneumovirus] AAV88364.1 fusion protein [Human metapneumovirus] AAN52910.1 fusion protein [Human metapneumovirus] AAN52915.1 fusion protein [Human metapneumovirus] BAM37564.1 fusion glycoprotein precursor [Human metapneumovirus] BAH59618.1 fusion protein [Human metapneumovirus] AAQ90144.1 fusion glycoprotein [Human metapneumovirus] AHV79446.1 fusion protein [Human metapneumovirus] AEL87260.1 fusion glycoprotein [Human metapneumovirus] AHV79867.1 fusion protein [Human metapneumovirus] ABQ66027.2 fusion glycoprotein [Human metapneumovirus] ACJ53621.1 fusion protein [Human metapneumovirus] AAN52911.1 fusion glycoprotein [Human metapneumovirus] AHV79536.1 fusion glycoprotein [Human metapneumovirus] AGU68411.1 fusion protein [Human metapneumovirus] AEZ52346.1 fusion protein [Human metapneumovirus] AAN52913.1 fusion protein [Human metapneumovirus] AAN52908.1 fusion glycoprotein [Human metapneumovirus] ACJ53553.1 fusion glycoprotein [Human metapneumovirus] AIY25727.1 fusion protein [Human metapneumovirus] ABM67072.1 fusion protein [Human metapneumovirus] AEZ52361.1 fusion protein [Human metapneumovirus] AAS22093.1 fusion glycoprotein [Human metapneumovirus] AGH27049.1 fusion protein [metapneumovirus] AAK62968.2 fusion glycoprotein [Human metapneumovirus] ACJ53556.1 fusion glycoprotein [Human metapneumovirus] ACJ53620.1 fusion protein [Human metapneumovirus] ABQ58820.1 F [Human metapneumovirus] [Human metapneumovirus] AEK26886.1 fusion glycoprotein [Human metapneumovirus] ACJ53619.1 fusion glycoprotein [Human metapneumovirus] ACJ53555.1 fusion [Human metapneumovirus] AGL74057.1 fusion protein [Human metapneumovirus] ABD27850.1 fusion protein [Human metapneumovirus] AEZ52349.1 fusion protein [Human metapneumovirus] ABD27848.1 fusion protein [Human metapneumovirus] ABD27846.1 fusion protein [Human metapneumovirus] ABQ66021.1 fusion protein [Human metapneumovirus] AFM57710.1 fusion protein [Human metapneumovirus] AFM57709.1 fusion protein [Human metapneumovirus] ABH05968.1 fusion protein [Human metapneumovirus] AEZ52350.1 fusion protein [Human metapneumovirus] AFM57712.1 fusion protein [Human metapneumovirus] AEZ52364.1 fusion protein [Human metapneumovirus] AAN52912.1 fusion protein [Human metapneumovirus] AEZ52363.1 fusion [Human metapneumovirus] AGL74059.1 fusion glycoprotein [Human metapneumovirus] ACJ53583.1 fusion protein [Human metapneumovirus] AEZ52356.1 fusion protein [Human metapneumovirus] AEZ52353.1 fusion glycoprotein [Human metapneumovirus] ACJ53581.1 fusion glycoprotein [Human metapneumovirus] ACJ53578.1 fusion protein [Human metapneumovirus] AAS22117.1 fusion protein [Human metapneumovirus] BAN75965.1 fusion protein [Human metapneumovirus] AGF92105.1 fusion protein [Human metapneumovirus] AAS22077.1 fusion protein [Human metapneumovirus] AAN52909.1 fusion glycoprotein [Human metapneumovirus] ACJ53586.1 fusion protein [Human metapneumovirus] AAQ90145.1 fusion glycoprotein [Human metapneumovirus] AGT75042.1 fusion [Human

metapneumovirus] AGL74058.1 fusion protein [Human metapneumovirus] AEL87263.1 fusion glycoprotein [Human metapneumovirus] AGH27057.1 fusion glycoprotein [Human metapneumovirus] AHV79491.1 F [Human metapneumovirus] [Human metapneumovirus] AEK26906.1 fusion glycoprotein [Human metapneumovirus] ACJ53580.1 fusion protein [Human metapneumovirus] AEZ52354.1 fusion protein [Human metapneumovirus] AAN52914.1 G [Human metapneumovirus] [Human metapneumovirus] AEK26901.1 glycoprotein [Human metapneumovirus] AFI56738.1 glycoprotein [Human metapneumovirus] AFI56739.1 glycoprotein [Human metapneumovirus] AFI56745.1 G protein [Human metapneumovirus] AAQ62718.1 G protein [Human metapneumovirus] AAQ62719.1 attachment glycoprotein G [Human metapneumovirus] AGH27104.1 G protein [Human metapneumovirus] AAQ62729.1 G protein [Human metapneumovirus] AAQ62728.1 glycoprotein [Human metapneumovirus] AFI56753.1 glycoprotein [Human metapneumovirus] AFI56746.1 glycoprotein [Human metapneumovirus] AFI56750.1 glycoprotein [Human metapneumovirus] AFI56747.1 G protein [Human metapneumovirus] AAQ62721.1 glycoprotein [Human metapneumovirus] AAT46573.1 glycoprotein [Human metapneumovirus] AFI56748.1 glycoprotein [Human metapneumovirus] AFI56736.1 glycoprotein [Human metapneumovirus] AFI56749.1 attachment glycoprotein G [Human metapneumovirus] AGH27131.1 attachment glycoprotein G [Human metapneumovirus] AHV79558.1 glycoprotein [Human metapneumovirus] AFI56740.1 glycoprotein [Human metapneumovirus] AFI56741.1 glycoprotein [Human metapneumovirus] AFI56744.1 attachment glycoprotein G [Human metapneumovirus] AHV79790.1 attachment glycoprotein G [Human metapneumovirus] AGH27122.1 attachment glycoprotein G [Human metapneumovirus] AHV79763.1 attachment glycoprotein G [Human metapneumovirus] AGZ48849.1 glycoprotein [Human metapneumovirus] AFI56743.1 attachment glycoprotein G [Human metapneumovirus] AHV79450.1 glycoprotein [Human metapneumovirus] AFI56751.1 attachment glycoprotein [Human metapneumovirus] AAS48482.1 attachment glycoprotein G [Human metapneumovirus] AHV79889.1 attachment surface glycoprotein [Human AGW43050.1 metapneumovirus] glycoprotein [Human metapneumovirus] AFI56754.1 attachment glycoprotein G [Human metapneumovirus] AHV79601.1 glycoprotein [Human metapneumovirus] AFI56752.1 attachment glycoprotein G [Human metapneumovirus] AHV79871.1 G protein [Human metapneumovirus] AEZ68099.1 attachment glycoprotein G [Human metapneumovirus] AHV79817.1 attachment glycoprotein G [Human metapneumovirus] AHV79943.1 attachment glycoprotein G [Human metapneumovirus] BAN75968.1 attachment surface glycoprotein [Human AGW43045.1 metapneumovirus] attachment glycoprotein G [Human metapneumovirus] AHV79628.1 attachment glycoprotein [Human metapneumovirus] AFK49783.1 G protein [Human metapneumovirus] AAQ62723.1 attachment glycoprotein [Human metapneumovirus] ABD27839.1 attachment surface glycoprotein [Human AGW43046.1 metapneumovirus] G protein [Human metapneumovirus] AAQ62717.1 glycoprotein [Human metapneumovirus] AFI56742.1 attachment protein [Human metapneumovirus] ABQ44522.1 glycoprotein [Human metapneumovirus] AFI56735.1 attachment surface glycoprotein [Human AGW43065.1 metapneumovirus] G protein [Human metapneumovirus] AAQ62724.1 attachment surface glycoprotein [Human AGW43075.1 metapneumovirus] attachment surface glycoprotein [Human AGW43062.1 metapneumovirus] glycoprotein [Human metapneumovirus] AAT46579.1 attachment surface glycoprotein [Human AGW43064.1 metapneumovirus] attachment surface glycoprotein [Human AGW43054.1 metapneumovirus] attachment surface glycoprotein [Human AGW43042.1 metapneumovirus] attachment surface glycoprotein [Human AGW43078.1 metapneumovirus] attachment surface glycoprotein [Human AGW43067.1 metapneumovirus] G protein [Human metapneumovirus] AAQ62722.1 attachment surface glycoprotein metapneumovirus] AGW43063.1 glycoprotein [Human metapneumovirus] AAT46571.1 glycoprotein [Human metapneumovirus] AAT46578.1 attachment glycoprotein G [Human metapneumovirus] AGJ74232.1 glycoprotein [Human metapneumovirus] AAT46580.1 glycoprotein [Human metapneumovirus] AAT46574.1 attachment

surface glycoprotein [Human AGW43061.1 metapneumovirus] attachment glycoprotein [Human metapneumovirus] AFK49791.1 attachment surface glycoprotein [Human AGW43047.1 metapneumovirus] glycoprotein [Human metapneumovirus] ABC26386.1 attachment glycoprotein [Human metapneumovirus] AAS48466.1 attachment surface glycoprotein [Human AGW43048.1 metapneumovirus] attachment glycoprotein G [Human metapneumovirus] AGH27140.1 attachment surface glycoprotein [Human AGW43049.1 metapneumovirus] attachment glycoprotein G [Human metapneumovirus] AGJ74082.1 attachment glycoprotein G [Human metapneumovirus] AHV79442.1 attachment glycoprotein G [Human metapneumovirus] AGJ74091.1 attachment glycoprotein G [Human metapneumovirus] AHV79477.1 attachment surface glycoprotein [Human AGW43056.1 metapneumovirus] attachment protein [Human metapneumovirus] ABQ44523.1 attachment glycoprotein G [Human metapneumovirus] BAH59622.1 attachment surface glycoprotein [Human AGW43070.1 metapneumovirus] glycoprotein [Human metapneumovirus] AAT46585.1 attachment glycoprotein G [Human metapneumovirus] AGU68409.1 attachment glycoprotein G [Human metapneumovirus] AGJ74223.1 attachment glycoprotein [Human metapneumovirus] AAS22129.1 attachment glycoprotein G [Human metapneumovirus] AGJ74048.1 G protein [Human metapneumovirus] AAQ62725.1 glycoprotein [Human metapneumovirus] ABC26384.1 attachment protein [Human metapneumovirus] ABQ44525.1 attachment glycoprotein G [Human metapneumovirus] YP_012612.1 attachment surface glycoprotein [Human AGW43071.1 metapneumovirus] attachment glycoprotein G [Human metapneumovirus] AGJ74162.1 attachment glycoprotein G [Human metapneumovirus] AGH27095.1 attachment glycoprotein G [Human metapneumovirus] AHV79531.1 G protein [Human metapneumovirus] AAQ62726.1 attachment glycoprotein [Human metapneumovirus] AAS48465.1 attachment surface glycoprotein [Human AGW43058.1 metapneumovirus] P [Human metapneumovirus] [Human metapneumovirus] AEK26894.1 phosphoprotein [Human metapneumovirus] AHV79631.1 phosphoprotein [Human metapneumovirus] AHV79901.1 phosphoprotein [Human metapneumovirus] AHV79570.1 phosphoprotein [Human metapneumovirus] AGJ74076.1 phosphoprotein [Human metapneumovirus] AAS22123.1 phosphoprotein [Human metapneumovirus] ABB16895.1 phosphoprotein [Human metapneumovirus] AHV79579.1 phosphoprotein [Human metapneumovirus] AGJ74244.1 phosphoprotein [Human metapneumovirus] AHV79856.1 phosphoprotein [Human metapneumovirus] ACJ70113.1 phosphoprotein [Human metapneumovirus] AGZ48843.1 phosphoprotein [Human metapneumovirus] AHV79498.1 phosphoprotein [Human metapneumovirus] AHV79480.1 phosphoprotein [Human metapneumovirus] ABQ43382.1 phosphoprotein [Human metapneumovirus] AAS22107.1 phosphoprotein [Human metapneumovirus] ABB16898.1 phosphoprotein [Human metapneumovirus] AGH27134.1 phosphoprotein [Human metapneumovirus] ABB16899.1 phosphoprotein [Human metapneumovirus] AGH27098.1 phosphoprotein [Human metapneumovirus] AAN52866.1 phosphoprotein [Human metapneumovirus] AAS22083.1 phosphoprotein [Human metapneumovirus] YP_012606.1 phosphoprotein [Human metapneumovirus] AHV79973.1 phosphoprotein [Human metapneumovirus] AHV79462.1 phosphoprotein [Human metapneumovirus] AGJ74042.1 phosphoprotein [Human metapneumovirus] AAV88362.1 P [Human metapneumovirus] [Human metapneumovirus] AIL23591.1 phosphoprotein [Human metapneumovirus] AHV79453.1 phosphoprotein [Human metapneumovirus] AGJ74261.1 phosphoprotein [Human metapneumovirus] AGH27116.1 phosphoprotein [Human metapneumovirus] ABB16444.1 phosphoprotein [Human metapneumovirus] ABB16445.1 phosphoprotein [Human metapneumovirus] AHV79507.1 phosphoprotein [Human metapneumovirus] BAH59616.1 phosphoprotein [Human metapneumovirus] ABB16443.1 phosphoprotein [Human metapneumovirus] ABQ43388.1 phosphoprotein [Human metapneumovirus] ABQ43389.1 phosphoprotein [Human metapneumovirus] ABQ43395.1 phosphoprotein [Human metapneumovirus] ABQ43385.1 phosphoprotein [Human

metapneumovirus] AAP84042.1 phosphoprotein [Human metapneumovirus] AAN52868.1 phosphoprotein [Human metapneumovirus] AAP84041.1 phosphoprotein [Human metapneumovirus] AGH27080.1 phosphoprotein [Human metapneumovirus] ABQ43387.1 phosphoprotein [Human metapneumovirus] AAS22099.1 phosphoprotein [Human metapneumovirus] ABB16896.1 phosphoprotein [Human metapneumovirus] AGJ74094.1 phosphoprotein [Human metapneumovirus] AEZ68089.1 phosphoprotein [Human metapneumovirus] ABK97002.1 phosphoprotein [Human metapneumovirus] AAP13486.1 phosphoprotein [Human metapneumovirus] AHV79444.1 phosphoprotein [Human metapneumovirus] AHV79865.1 phosphoprotein [Human metapneumovirus] AGJ74226.1 phosphoprotein [Human metapneumovirus] ABQ43383.1 phosphoprotein [Human metapneumovirus] AAN52863.1 phosphoprotein [Human metapneumovirus] AHV79775.1 phosphoprotein [Human metapneumovirus] AEZ68094.1 phosphoprotein [Human metapneumovirus] AHV79883.1 phosphoprotein [Human metapneumovirus] AEZ68092.1 phosphoprotein [Human metapneumovirus] ABQ43390.1 phosphoprotein [Human metapneumovirus] ABQ43386.1 phosphoprotein [Human metapneumovirus] ABQ43391.1 phosphoprotein [Human metapneumovirus] ACS16062.1 phosphoprotein [Human metapneumovirus] AEZ68090.1 phosphoprotein [Human metapneumovirus] AAK62967.1 phosphoprotein [Human metapneumovirus] AEZ68093.1 phosphoprotein [Human metapneumovirus] AEZ68088.1 phosphoprotein [Human metapneumovirus] ABQ43392.1 phosphoprotein [Human metapneumovirus] ABQ43393.1 phosphoprotein [Human metapneumovirus] ABQ43384.1 phosphoprotein [Human metapneumovirus] ABQ43394.1 phosphoprotein [Human metapneumovirus] ABK96999.1 phosphoprotein [Human metapneumovirus] AHV79489.1 phosphoprotein [Human metapneumovirus] AGJ74235.1 phosphoprotein [Human metapneumovirus] AAS22075.1 phosphoprotein [Human metapneumovirus] AAS22115.1 phosphoprotein [Human metapneumovirus] AII17601.1 phosphoprotein [Human metapneumovirus] ABK97000.1 phosphoprotein [Human metapneumovirus] AHV79561.1 phosphoprotein [Human metapneumovirus] AGT75040.1 phosphoprotein [Human metapneumovirus] AAN52864.1 phosphoprotein [Human metapneumovirus] ABK97001.1 phosphoprotein [Human metapneumovirus] AGT74979.1 phosphoprotein [Human metapneumovirus] AHV79955.1 phosphoprotein [Human metapneumovirus] AGH27055.1 phosphoprotein [Human metapneumovirus] AAV88361.1 phosphoprotein [Human metapneumovirus] ABQ43397.1 phosphoprotein [Human metapneumovirus] AGJ74173.1 P [Human metapneumovirus] [Human metapneumovirus] AEK26904.1 phosphoprotein [Human metapneumovirus] ACJ70104.1 phosphoprotein [Human metapneumovirus] ABK97003.1 phosphoprotein [Human metapneumovirus] AGT74955.1 phosphoprotein [Human metapneumovirus] AAN52856.1 phosphoprotein [Human metapneumovirus] AAN52862.1 phosphoprotein [Human metapneumovirus] AGJ74138.1 phosphoprotein [Human metapneumovirus] AHV79613.1 phosphoprotein [Human metapneumovirus] AGJ74060.1 phosphoprotein [Human metapneumovirus] AAQ67684.1 phosphoprotein [Human metapneumovirus] AEA02278.1 N [Human metapneumovirus] [Human metapneumovirus] AEK26899.1 nucleoprotein [Human metapneumovirus] ACS16061.1 nucleoprotein [Human metapneumovirus] AAS88425.1 nucleoprotein [Human metapneumovirus] YP_012605.1 nucleoprotein [Human metapneumovirus] AHV79882.1 nucleoprotein [Human metapneumovirus] AHV79774.1 nucleocapsid protein [Human metapneumovirus] AAN52886.1 nucleoprotein [Human metapneumovirus] AAS22082.1 nucleoprotein [Human metapneumovirus] AHV79864.1 nucleoprotein [Human metapneumovirus] AHV79828.1 nucleoprotein [Human metapneumovirus] AGJ74084.1 nucleocapsid protein [Human metapneumovirus] AAN52888.1 N [Human metapneumovirus] [Human metapneumovirus] AIL23590.1 nucleoprotein [Human metapneumovirus] AAK62966.1 nucleoprotein [Human metapneumovirus] AHV79972.1 nucleoprotein [Human metapneumovirus] AHV79470.1 nucleoprotein [Human metapneumovirus]

AHV79452.1 nucleoprotein [Human metapneumovirus] AGJ74243.1 nucleoprotein [Human metapneumovirus] AHV79533.1 nucleoprotein [Human metapneumovirus] AGJ74181.1 nucleoprotein [Human metapneumovirus] AHV79497.1 nucleoprotein [Human metapneumovirus] AHV79702.1 nucleoprotein [Human metapneumovirus] AHV79648.1 nucleoprotein [Human metapneumovirus] AHV79435.1 putative nucleoprotein [Human metapneumovirus] AGJ74260.1 nucleocapsid protein [Human metapneumovirus] AAN52887.1 nucleoprotein [Human metapneumovirus] AGU68386.1 nucleocapsid protein [Human metapneumovirus] AAN52899.1 nucleoprotein [Human metapneumovirus] AAR17673.1 nucleocapsid protein [Human metapneumovirus] AAN52898.1 nucleoprotein [Human metapneumovirus] AEA02277.1 nucleoprotein [Human metapneumovirus] AHV79612.1 nucleoprotein [Human metapneumovirus] AGU68416.1 nucleoprotein [Human metapneumovirus] AGU68408.1 nucleoprotein [Human metapneumovirus] AGU68370.1 nucleoprotein [Human metapneumovirus] AAQ67683.1 nucleoprotein [Human metapneumovirus] AGJ74137.1 nucleoprotein [Human metapneumovirus] AGU68344.1 nucleocapsid protein [Human metapneumovirus] ABK96997.1 nucleoprotein [Human metapneumovirus] AGU68413.1 nucleocapsid protein [Human metapneumovirus] AAN52891.1 nucleoprotein [Human metapneumovirus] AGU68360.1 nucleoprotein [Human metapneumovirus] AGU68353.1 nucleocapsid protein [Human metapneumovirus] ABK96996.1 nucleoprotein [Human metapneumovirus] AAR17666.1 N [Human metapneumovirus] [Human metapneumovirus] AEK26903.1 nucleoprotein [Human metapneumovirus] AGT75039.1 nucleoprotein [Human metapneumovirus] AGU68410.1 nucleoprotein [Human metapneumovirus] AAS22074.1 nucleoprotein [Human metapneumovirus] AHV79560.1 nucleoprotein [Human metapneumovirus] AGT74978.1 nucleoprotein [Human metapneumovirus] AGJ74128.1 nucleoprotein [Human metapneumovirus] AAR17663.1 nucleoprotein [Human metapneumovirus] AAR17662.1 nucleoprotein [Human metapneumovirus] AAR17664.1 nucleoprotein [Human metapneumovirus] AAR17657.1 nucleoprotein [Human metapneumovirus] AAR17659.1 nucleoprotein [Human metapneumovirus] AAR17661.1 nucleoprotein [Human metapneumovirus] AGU68352.1 nucleoprotein [Human metapneumovirus] AGU68373.1 nucleoprotein [Human metapneumovirus] AGU68376.1 nucleoprotein [Human metapneumovirus] AGU68342.1 nucleoprotein [Human metapneumovirus] AGU68365.1 nucleoprotein [Human metapneumovirus] AGU68363.1 nucleoprotein [Human metapneumovirus] AGU68398.1 nucleoprotein [Human metapneumovirus] AGU68348.1 nucleoprotein [Human metapneumovirus] AGU68354.1 nucleoprotein [Human metapneumovirus] AGU68391.1 nucleoprotein [Human metapneumovirus] AGU68389.1 nucleoprotein [Human metapneumovirus] AGU68399.1 nucleoprotein [Human metapneumovirus] AGU68337.1 nucleoprotein [Human metapneumovirus] AAR17660.1 nucleoprotein [Human metapneumovirus] AAR17667.1 nucleoprotein [Human metapneumovirus] AGU68402.1 nucleoprotein [Avian metapneumovirus type C] CDN30025.1 nucleoprotein [Avian metapneumovirus] AGZ87947.1 Nucleoprotein [Avian metapneumovirus type C] CAL25113.1 nucleocapsid protein [Avian metapneumovirus] ABO42286.1 nucleocapsid protein [Avian metapneumovirus] AAK38430.1 nucleocapsid protein [Avian metapneumovirus] AAK54155.1 nucleocapsid protein [Avian metapneumovirus] AAK38426.1 nucleocapsid protein [Avian metapneumovirus] AAK38425.1 nucleocapsid protein [Avian metapneumovirus] AAK38424.1 nucleocapsid protein [Avian metapneumovirus] AAF05909.1 nucleocapsid protein [Avian metapneumovirus] AAK38435.1 nucleocapsid protein [Avian metapneumovirus] AAK38428.1 nucleoprotein [Human metapneumovirus] AAR17669.1 nucleocapsid protein [Avian metapneumovirus] AAK38429.1 nucleocapsid protein [Avian metapneumovirus] AAK38427.1 nucleocapsid protein [Avian metapneumovirus] AAK38423.1 nucleocapsid protein [Avian metapneumovirus] AAK38434.1 nucleoprotein [Human metapneumovirus] AGU68338.1 nucleoprotein [Avian metapneumovirus] YP_443837.1 nucleoprotein [Human metapneumovirus] AGU68384.1 nucleocapsid protein [Avian metapneumovirus] AAK38431.1 nucleoprotein [Human metapneumovirus] AGU68405.1 nucleoprotein [Human metapneumovirus] AGU68382.1

nucleoprotein [Human metapneumovirus] AGU68395.1 nucleocapsid [Human metapneumovirus]
AAL35389.3 nucleoprotein [Human metapneumovirus] AEZ68064.1
(576) TABLE-US-00006 TABLE 5 SEQ ID Description Sequence NO: PIV3 Nucleic
Acid Sequences >gb|KJ672601.1|: ATGCCAATTTCAATACTGTTAATTATTACAACCATGATC
9 4990-6609 Human ATGGCATCACACTGCCAAATAGACATCACAAAACACTACA
parainfluenza virus GCATGTAGGTGTATTGGTCAACAGTCCCAAAGGGATGA 3 strain
AGATATCACAAAACCTTCGAAACAAGATATCTAATCCTGA HPIV3/Homo
GTCTCATACCAAAAATAGAAGATTCTAACTCTTGTGGTG sapiens/PER/
ACCAACAGATCAAGCAATACAAGAGGTTATTGGATAGA FLA4815/2008 [fusion
CTGATCATTCTTTTATATGATGGACTAAGATTACAGAAG glycoprotein F0]
GATGTGATAGTGACTAATCAAGAATCCAATGAAAACAC
TGATCCCAGAACAGAACGATTCTTTGGAGGGGTAATTGG
AACTATTGCTCTAGGAGTAGCAACCTCAGCACAAATTAC
AGCAGCAGTTGCTCTGGTTGAAGCCAAGCAGGCAAGAT
CAGACATTGAAAACTCAAGGAAGCAATCAGGGACACA
AATAAAGCAGTGCAGTCAGTTCAGAGCTCTGTAGGAAA
TTTGATAGTAGCAATTAAATCAGTCCAGGATTATGTCAA
CAAAGAAATCGTGCCATCGATTGCGAGACTAGGTTGTG
AAGCAGCAGGACTTCAGTTAGGGATTGCATTAACACAG
CATTACTCAGAATTAACAAATATATTTGGTGATAACATA
GGATCGTTACAAGAAAAAGGAATAAAATTACAAGGTAT
AGCATCATTATACCGTACAAATATCACAGAAATATTCAC
AACATCAACAGTTGACAAATATGATATTTATGATCTATT
ATTTACAGAATCAATAAAGGTGAGAGTTATAGATGTTGA
TTTGAATGATTACTCAATAACCCTCCAAGTCAGACTCCC
TTTATTGACCAGACTGCTGAACACTCAAATCTACAAAGT
AGATTCCATATCATACAATATCCAAAATAGAGAATGGTA
TATCCCTCTTCCCAGCCATATCATGACGAAAGGGGCATT
TCTAGGTGGAGCAGATGTCAAAGAATGCATAGAAGCAT
TCAGCAGTTATATATGCCCTTCTGATCCAGGATTTGTACT
AAACCATGAAATGGAGAGCTGTCTATCAGGAAACATAT
CCCAATGTCCAAGAACCACAGTCACATCAGACATAGTTC
CTAGGTATGCATTTGTCAATGGAGGAGTGGTTGCGAATT
GTATAACAACACTACATGTACATGCAATGGTATCGGTAATA
GAATCAACCAACCACCTGATCAAGGAGTCAAAATTATA
ACACATAAAGAATGTAATACAATAGGTATCAACGGAAT
GCTATTCAACACAAACAAAGAAGGAACTCTTGCATTCTA
CACACCAGACGACATAACATTAAACAATTCTGTTGCACT
TGATCCGATTGACATATCAATCGAGCTCAACAAGGCCAA
ATCAGATCTTGAGGAATCAAAGAATGGATAAGAAGGT
CAAATCAAAGCTAGATTCTATTGGAAGTTGGCATCAAT
CTAGCACTACAATCATAGTTATTTTGATAATGATGATTA
TATTGTTTATAATTAATATAACAATAATTACAATTGCAA
TTAAGTATTACAGAATTCAAAAGAGAAATCGAGTGGAT
CAAAATGATAAGCCGTATGTATTAACAAACAAG gi|612507167|gb|
ATGGAATACTGGAAGCACACCAACCACGGAAAGGATGC 10 AHX22430.1|
TGGTAATGAGCTGGAGACATCCACAGCCACTCATGGCA hemagglutinin-
ACAAGCTCACCAACAAGATAACATATATATTGTGGACG neuraminidase
ATAACCCTGGTGTATTATCAATAGTCTTCATCATAGTG [Human
CTAACTAATTCCATCAAAGTGAAAAGGCCCGCGAATC parainfluenza virus

ATTGCTACATGACATATAATGAGTTTATGGAAGTTAC 3]
AGAAAAGATCCAAGTGGCATCGGATAATACTAATGATC
TAATACAGTCAGGAGTGAATACAAGGCTTCTTACAATTC
AGAGTCATGTCCAGAATTATATACCAATATCATTGACAC
AACAAATATCGGATCTTAGGAAATTCATTAGTGAAATTA
CAATTAGAAATGATAATCAAGAAGTGCCACCACAAAGA
ATAACACATGATGTGGGTATAAAACCTTTAAATCCAGAT
GATTTCTGGAGATGCACGTCTGGTCTTCCATCTTTGATG
AAAACCTCCAAAAATAAGATTAATGCCGGGACCAGGATT
ATTAGCTATGCCAACGACTGTTGATGGCTGTGTCAGAAC
CCCGTCCTTAGTGATAAATGATCTGATTTATGCTTACAC
CTCAAATCTAATTACTCGAGGTTGCCAGGATATAGGGAA
ATCATATCAAGTATTACAGATAGGGATAATAACTGTAAA
CTCAGACTTGGTACCTGACTTAAATCCTAGGATCTCTCA
TACCTTCAACATAAATGACAATAGAAAGTCATGTTCTCT
AGCACTCCTAAATACAGATGTATATCAACTGTGTTCAAC
CCCAAAAGTTGATGAAAGATCAGATTATGCATCATCAG
GCATAGAAGATATTGTACTTGATATTGTCAATTATGATG
GCTCAATCTCGACAACAAGATTTAAGAATAATAATATAA
GTTTTGATCAACCATATGCGGCATTATACCCATCTGTTG
GACCAGGGATATACTACAAAGGCAAAATAATTTCTC
GGGTATGGAGGTCTTGAACATCCAATAAATGAGAATGC
AATCTGCAACACAACCTGGGTGTCCTGGGAAAACACAGA
GAGACTGTAATCAAGCATCTCATAGTCCATGGTTTTTCAG
ATAGAAGGATGGTCAACTCTATAATTGTTGTTGACAAGG
GCTTGAACCTCAGTTCCAAAATTGAAGGTATGGACGATAT
CTATGAGACAAAATTACTGGGGGTCAGAAGGAAGATTA
CTTCTACTAGGTAACAAGATCTACATATACACAAGATCT
ACAAGTTGGCACAGCAAGTTACAATTAGGAATAATTGA
CATTACTGACTACAGTGATATAAGGATAAAATGGACAT
GGCATAATGTGCTATCAAGACCAGGAAACAATGAATGT
CCATGGGGACATTCATGTCCGGATGGATGTATAACGGG
AGTATATACCGATGCATATCCACTCAATCCCACAGGAAG
CATTGTATCATCTGTTCATATTGGACTCACAAAAATCGAG
AGTCAACCCAGTCATAACTTACTCAACAGCAACCGAAA
GGGTAAACGAGCTGGCTATCCGAAACAAAACACTCTCA
GCTGGGTACACAACAACAAGCTGCATTACACACTATAA
CAAAGGGTATTGTTTTTCATATAGTAGAAATAAATCATAA
AAGCTTAAACACATTTCAACCCATGTTGTTCAAAACAGA GATTCCAAAAAGCTGCAGT
HPIV3_HN_Codon ATGGAATACTGGAAGCACACCAACCACGGCAAGGACGC 11
Optimized CGGCAACGAGCTGGAAACCAGCACAGCCACACACGGCA
ACAAGCTGACCAACAAGATCACCTACATCCTGTGGACC
ATCACCCCTGGTGCTGCTGAGCATCGTGTTTCATCATCGTG
CTGACCAATAGCATCAAGAGCGAGAAGGCCAGAGAGAG
CCTGCTGCAGGACATCAACAACGAGTTCATGGAAGTGA
CCGAGAAGATCCAGGTGGCCAGCGACAACACCAACGAC
CTGATCCAGAGCGGCGTGAACACCCGGCTGCTGACCATC
CAGAGCCACGTGCAGAACTACATCCCCATCAGCCTGACC
CAGCAGATCAGCGACCTGCGGAAGTTCATCAGCGAGAT
CACCATCCGGAACGACAACCAGGAAGTGCCCCCCCAGA

GAATCACCGCCATCAAGCCCCTGAACCCC
GACGATTTCTGGCGGTGTACAAGCGGCCTGCCCAGCCTG
ATGAAGACCCCCAAGATCCGGCTGATGCCTGGCCCTGG
ACTGCTGGCCATGCCTACCACAGTGGATGGCTGTGTGCG
GACCCCCAGCCTCGTGATCAACGATCTGATCTACGCCTA
CACCAGCAACCTGATCACCCGGGGCTGCCAGGATATCG
GCAAGAGCTACCAGGTGCTGCAGATCGGCATCATCACC
GTGAACTCCGACCTGGTGCCCGACCTGAACCCTCGGATC
AGCCACACCTTCAACATCAACGACAACAGAAAGAGCTG
CAGCCTGGCTCTGCTGAACACCGACGTGTACCAGCTGTG
CAGCACCCCCAAGGTGGACGAGAGAAGCGACTACGCCA
GCAGCGGCATCGAGGATATCGTGCTGGACATCGTGAAC
TACGACGGCAGCATCAGCACCAACCCGGTTCAAGAACAA
CAACATCAGCTTCGACCAGCCCTACGCCGCCCTGTACCC
TTCTGTGGGCCCTGGCATCTACTACAAGGGCAAGATCAT
CTTCCTGGGCTACGGCGGCCTGGAACACCCCATCAACGA
GAACGCCATCTGCAACACCACCGGCTGCCCTGGCAAGA
CCCAGAGAGACTGCAATCAGGCCAGCCACAGCCCCTGG
TTCAGCGACCGCAGAATGGTCAACTCTATCATCGTGGTG
GACAAGGGCCTGAACAGCGTGCCCAAGCTGAAAGTGTG
GACAATCAGCATGCGCCAGAACTACTGGGGCAGCGAGG
GCAGACTTCTGCTGCTGGGAAACAAGATCTACATCTACA
CCCGGTCCACCAGCTGGCACAGCAAATGCAGCTGGGA
ATCATCGACATCACCGACTACAGCGACATCCGGATCAA
GTGGACCTGGCACAACGTGCTGAGCAGACCCGGCAACA
ATGAGTGCCCTTGGGGCCACAGCTGCCCCGATGGATGTA
TCACCGGCGTGTACACCGACGCCTACCCCCTGAATCCTA
CCGGCTCCATCGTGTCCAGCGTGATCCTGGACAGCCAGA
AAAGCAGAGTGAACCCCGTGATCACATACAGCACCGCC
ACCGAGAGAGTGAACGAACCTGGCCATCAGAAACAAGAC
CCTGAGCGCCGGCTACACCACCACAAGCTGCATCACAC
ACTACAACAAGGGCTACTGCTTCCACATCGTGGAATCA
ACCACAAGTCCCTGAACACCTTCCAGCCCATGCTGTTCA
AGACCGAGATCCCCAAGAGCTGCTCC HPIV3_F_Codon
ATGCCCATCAGCATCCTGCTGATCATCACCACAATGATC 12 Optimized
ATGGCCAGCCACTGCCAGATCGACATACCAAGCTGCA
GCACGTGGGCGTGCTCGTGAACAGCCCCAAGGGCATGA
AGATCAGCCAGAACTTCGAGACACGCTACCTGATCCTGA
GCCTGATCCCCAAGATCGAGGACAGCAACAGCTGCGGC
GACCAGCAGATCAAGCAGTACAAGCGGCTGCTGGACAG
ACTGATCATCCCCCTGTACGACGGCCTGCGGCTGCAGAA
AGACGTGATCGTGACCAACCAGGAAAGCAACGAGAACAA
CCGACCCCCGGACCGAGAGATTCTTCGGCGGCGTGATCG
GCACAATCGCCCTGGGAGTGGCCACAAGCGCCCAGATT
ACAGCCGCTGTGGCCCTGGTGGAAGCCAAGCAGGCCAG
AAGCGACATCGAGAAGCTGAAAGAGGCCATCCGGGACA
CCAACAAGGCCGTGCAGAGCGTGCAGTCCAGCGTGGGC
AATCTGATCGTGGCCATCAAGTCCGTGCAGGACTACGTG
AACAAAGAAATCGTGCCCTCTATCGCCCGGCTGGGCTGT
GAAGCTGCCGGACTGCAGCTGGGCATTGCCCTGACACA

GCACTACAGTGTGACATCTTTCGGCGACAACA
TCGGCAGCCTGCAGGAAAAGGGCATTAGCTGCAGGGA
ATCGCCAGCCTGTACCGCACCAACATCACCGAGATCTTC
ACCACCAGCACCGTGGATAAGTACGACATCTACGACCT
GCTGTTACCGAGAGCATCAAAGTGCGCGTGATCGACGT
GGACCTGAACGACTACAGCATCACCTGCAAGTGCGGC
TGCCCCTGCTGACCAGACTGCTGAACACCCAGATCTACA
AGGTGGACAGCATCTCCTACAACATCCAGAACCGCGAG
TGGTACATCCCTCTGCCCAGCCACATTATGACCAAGGGC
GCCTTTCTGGGCGGAGCCGACGTGAAAGAGTGCATCGA
GGCCTTCAGCAGCTACATCTGCCCCAGCGACCCTGGCTT
CGTGCTGAACCACGAGATGGAAAGCTGCCTGAGCGGCA
ACATCAGCCAGTGCCCCAGAACCACCGTGACCTCCGAC
ATCGTGCCCAGATACGCCTTCGTGAATGGCGGCGTGGTG
GCCAACTGCATCACCAACACCTGTACCTGCAACGGGCATC
GGCAACCGGATCAACCAGCCTCCCGATCAGGGCGTGAA
GATTATCACCCACAAAGAGTGTAACACCATCGGCATCA
ACGGCATGCTGTTCAATAACCAACAAAGAGGGCACCTG
GCCTTCTACACCCCCGACGATATCACCTGAACAACCTCC
GTGGCTCTGGACCCCATCGACATCTCCATCGAGCTGAAC
AAGGCCAAGAGCGACCTGGAAGAGTCCAAAGAGTGGAT
CCGGCGGAGCAACCAGAAGCTGGACTCTATCGGCAGCT
GGCACCAGAGCAGCACCACCATCATCGTGATCCTGATTA
TGATGATTATCCTGTTTCATCATCAACATTACCATCATCAC
TATCGCCATTAAGTACTACCGGATCCAGAAACGGAACC
GGGTGGACCAGAATGACAAGCCCTACGTGCTGACAAAC AAG PIV3 mRNA
Sequences >gb|KJ672601.1|: AUGCCAAUUUCAAUACUGUUAUUUAUUAACAACCAUGA
61 4990-6609 Human UCAUGGCAUCACACUGCCAAAUAGACAUCACAAAACU
parainfluenza virus ACAGCAUGUAGGUGUAUUGGUCAACAGUCCCAAAGGG 3 strain
AUGAAGAUUAUCACAAAACUUCGAAACAAGAUUAUCUAA HPIV3/*Homo*
UCCUGAGUCUCAUACCAAAAAUAGAAGAUUCUAACUC *sapiens*/PER/FLA4
UUGUGGUGACCAACAGAUCAAGCAAUACAAGAGGUUA 815/2008 [fusion
UUGGAUAGACUGAUCAUUCCUUUAUAUGAUGGACUAA glycoprotein F0]
GAUUACAGAAGGAUGUGAUAGUGACUAAUCAAGAAUC
CAAUGAAAACACUGAUCCCAGAACAGAACGAUUCUUU
GGAGGGGUAAUUGGAACUAUUGCUCUAGGAGUAGCAA
CCUCAGCACAAAUUACAGCAGCAGUUGCUCUGGUUGA
AGCCAAGCAGGCAAGAUUCAGACAUUGAAAAACUCAAG
GAAGCAAUCAGGGACACAAAUAAGCAGUGCAGUCAG
UUCAGAGCUCUGUAGGAAAUUUGAUAGUAGCAAUUAA
AUCAGUCCAGGAUUAUGUCAACAAAGAAUCGUGCCA
UCGAUUGCGAGACUAGGUUGUGAAGCAGCAGGACUUC
AGUUAGGGAUUGCAUUAACACAGCAUUAUCAGAAUU
AACAAAUAUAUUUGGUGAUAAAUAGGAUCGUUACAA
GAAAAAGGAUAAAAUUAACAAGGUUAUAGCAUCAUUUAU
ACCGUACAAAUAUCACAGAAAUUAUUCACAACAUAAC
AGUUGACAAAUAUGAUUUUAUGAUCUAUUUUUACA
GAUCAAUAAAGGUGAGAGUUUAUGAUGUUGAUUUGA
AUGAUUACUCAAUAAACCCUCCAAGUCAGACUCCCUUU
AUUGACCAGACUGCUGAACACUCAAUUAACAAGUA

GAUUCUAAUUAUCCAAAUAGAGAAUGGU
AUAUCCCUCUUCCCAGCCAUAUCAUGACGAAAGGGGC
AUUUCUAGGUGGAGCAGAUGUCAAGAAUGCAUAGAA
GCAUUCAGCAGUUAUUAUGCCCUUCUGAUCCAGGAU
UUGUACUAAACCAUGAAAUGGAGAGCUGUCUAUCAGG
AAACAUAUCCCAAUGUCCAAGAACCACAGUCACAUCA
GACAUAGUUCCUAGGUAUGCAUUUGUCAUUGGAGGAG
UGGUUGCGAAUUGUAUAACAACUACAUGUACAUGCAA
UGGUAUCGGUAAUAGAAUCAACCAACCACCUGAUCAA
GGAGUCAAAAUUAUAACACAUAAGAAUGUAAUACAA
UAGGUAUCAACGGAAUGCUAUUCAACACAAACAAAGA
AGGAACUCUUGCAUUCUACACACCAGACGACAUAACA
UUAACAAUUCUGUUGCACUUGAUCCGAUUGACAUAU
CAAUCGAGCUCACAAGGCCAAAUCAGAUCUUGAGGA
AUCAAAAGAAUGGAUAAGAAGGUCAAAUCAAAGCUA
GAUUCUAUUGGAAGUUGGCAUCAUUCUAGCACUACAA
UCAUAGUUAUUUGAUAAUGAUGAUUAUUAUUGUUUAU
AAUUAUUAACAUAUUAUACAAUUGCAAUUAAGUAU
UACAGAAUUCAAAAGAGAAAUUCGAGUGGAUCAAUUG
AUAAGCCGU AUGUAUUAACAACAAG gi|612507167|gb|
AUGGAAUACUGGAAGCACACCAACCACGGAAAGGAUG 62 AHX22430.1|
CUGGUAAUGAGCUGGAGACAUCCACAGCCACUCAUGG hemagglutinin-
CAACAAGCUCACCAACAAGAUACAUAUAUUAUUGUGG neuraminidase
ACGAUAACCCUGGUGUUAUUAUCAAUAGUCUUCAUCA [Human
UAGUGCUAACUAAUUCCAUCAAAAGUGAAAAGGCCCG parainfluenza virus
CGAAUCAUUGC UACAAGACAUAUAAUUGAGUUUAUG 3]
GAAGUUACAGAAAAGAUCCAAGUGGCAUCGGAUAAUA
CUAAUGAUCUAAUACAGUCAGGAGUGAAUACAAGGCU
UCUUACAAUUCAGAGUCAUGUCCAGAAUUAUAUACCA
AUUAUCAUUGACACAACAAUUAUCGGAUCUUAAGGAAU
UCAUUAGUGAAAUAUCAAUUAAGAAUGAUAAUCAAGA
AGUGCCACCACAAAGAAUAACACAUGAUGUGGGUAUA
AAACCUUUAUUAUCCAGAUUAUUCUGGAGAUGCACGU
CUGGUCUCCAUCUUAUGAUGAAAACUCCAAAAUAAG
AUUAAUGCCGGGACCAGGAUUAUUAAGCUAUGCCAACG
ACUGUUGAUGGCUGUGUCAGAACCCCGUCCUUAUGUGA
UAAAUGAUCUGAUUUAUGCUUACACCUCAAAUCUAAU
UACUCGAGGUUGCCAGGAUAUAGGGAAUCAUAUCAA
GUAUUACAGAUAGGGAUAAUAACUGUAAACUCAGACU
UGGUACCUGACUAAAUCCUAGGAUCUCUCAUACCUU
CAACAUAAUUGACAAUAGAAAGUCAUGUUCUCUAGCA
CUCCUAAAUAUAGAUUAUUAUACUGUGUUAACCC
CAAAGUUGAUGAAAGAUCAUAUUAUGCAUCAUCAGG
CAUAGAAGAUUAUUGUACUUGAUUAUUGUCAAUUAUGAU
GGCUCAAUCUCGACAACAAGAUUUAAGAAUAAUAAUA
UAAGUUUUGAUCAACCAUAUGCGGCAUUAUACCCAUC
UGUUGGACCAGGGAUAUACUACAAAGGC AAAUAAUA
UUUCUCGGGU AUGGAGGUCUUGAACAUCCAAUAAAUG
AGAAUGCAAUCUGCAACACAACUGGGUGUCCUGGGAA
AACACAGAGAGACUGUAAUCAAGCAUCUCAUAGUCCA

UGGUUUUCAGAAUGGUCAACUCUAUAAUUG
UUGUUGACAAGGGCUUGAACUCAGUUCCAAAUUGAA
GGUAUGGACGAUAUCUAUGAGACAAAUACUGGGGG
UCAGAAGGAAGAUUACUUCUACUAGGUAACAAGAUCU
ACAUAUACACAAGAUCUACAAGUUGGCACAGCAAGUU
ACAAUUAGGAUAUAAUUGACAUUACUGACUACAGUGAU
AUAAGGAUAAAAUGGACAUGGCAUAAUGUGCUAUCAA
GACCAGGAAACAAUGAAUGUCCAUGGGGACAUUCAUG
UCCGGAUGGAUGUAUAAACGGGAGUAUUAUACCGAUGCA
UAUCCACUCAAUCCCACAGGAAGCAUUGUAUCAUCUG
UCAUAUUGGACUCACAAAAAUCGAGAGUCAACCCAGU
CAUAACUUAUCUACACAGCAACCGAAAGGGUAAACGAG
CUGGCUAUCCGAAACAAAACACUCUCAGCUGGGUACA
CAACAACAAGCUGCAUUAACACUAUAACAAAGGGUA
UUGUUUUCAUUAUAGUAGAAAUAAAUCAUAAAAGCUUA
AACACAUUUCAACCCAUGUUGUUCAAAACAGAGAUUC CAAAAAGCUGCAGU
HPIV3_HN_Codon AUGGAAUACUGGAAGCACACCAACCACGGCAAGGACG 63
Optimized CCGGCAACGAGCUGGAAACCAGCACAGCCACACACGGC
AACAAGCUGACCAACAAGAUCACCUACAUCCUGUGGA
CCAUCACCCUGGUGCUGCUGAGCAUCGUGUUCAUCAUC
GUGCUGACCAAUAGCAUCAAGAGCGAGAAGGCCAGAG
AGAGCCUGCUGCAGGACAUCAACAACGAGUUCAUGGA
AGUGACCGAGAAGAUAUCCAGGUGGCCAGCGACAACACC
AACGACCUGAUCCAGAGCGGCGUGAACACCCGGCUGCU
GACCAUCCAGAGCCACGUGCAGAACUACAUCCCAUCA
GCCUGACCCAGCAGAUCAAGCGACCUGCGGAAGUUCAUC
AGCGAGAUCAACCAUCCGGAACGACAACCAGGAAGUGC
CCCCCCAGAGAAUACCCACGACGUGGGCAUCAAGCCC
CUGAACCCCGACGAUUUCUGGCGGUGUACAAGCGGCC
UGCCCAGCCUGAUGAAGACCCCCAAGAUCCGGCUGAUG
CCUGGCCCUGGACUGCUGGCCAUGCCUACCACAGUGGA
UGGCUGUGUGCGGACCCCCAGCCUCGUGAUCAACGAUC
UGAUCUACGCCUACACCAGCAACCUGAUCACCCGGGGC
UGCCAGGAUAUCGGCAAGAGCUACCAGGUGCUGCAGA
UCGGCAUCAUACCGUGAACUCCGACCUGGUGCCCGAC
CUGAACCCUCGGAUCAGCCACACCUUCAACAUCAACGA
CAACAGAAAGAGCUGCAGCCUGGCUCUGCUGAACACC
GACGUGUACCAGCUGUGCAGCACCCCCAAGGUGGACG
AGAGAAGCGACUACGCCAGCAGCGGCAUCGAGGAUUAU
CGUGCUGGACAUCGUGAACUACGACGGCAGCAUCAGC
ACCACCCGGUUCAAGAACAACAUCAGCUUCGACCA
GCCCUCACGCCGCCUGUACCCUUCUGUGGGCCCUGGCA
UCUACUACAAGGGCAAGAUCAUUCUCCUGGGCUACGG
CGGCCUGGAACACCCCAUCAACGAGAACGCCAUCUGCA
ACACCACCGGCUGCCCUGGCAAGACCCAGAGAGACUGC
AAUCAGGCCAGCCACAGCCCCUGGUUCAGCGACCGCAG
AAUGGUCAACUCUAUCAUCGUGGUGGACAAGGGCCUG
AACAGCGUGCCCAAGCUGAAAGUGUGGACAAUCAGCA
UGCGCCAGAACUACUGGGGCAGCGAGGGCAGACUUCU
GCUGCUGGGAAACAAGAUCUACAUCUACACCCGGUCC

ACCAGCUGGACGUCUGGGAUUAUCCG
ACAUCACCGACUACAGCGACAUCCGGAUCAAGUGGACC
UGGCACAACGUGCUGAGCAGACCCGGCAACAAUGAGU
GCCCUUGGGGCCACAGCUGCCCCGAUGGAUGUAUCCACC
GGCGUGUACACCGACGCCUACCCCCUGAAUCCUACCGG
CUCCAUCGUGUCCAGCGUGAUCCUGGACAGCCAGAAA
AGCAGAGUGAACCCCGUGAUACAUAACAGCACCGCCAC
CGAGAGAGUGAACGAACUGGCCAUAGAAACAAGACC
CUGAGCGCCGGCUACACCACCACAAGCUGCAUCACACA
CUACAACAAGGGGCUACUGCUUCCACAUCGUGGAAAUC
AACCACAAGUCCCUGAACACCUUCCAGCCCAUGCUGUU
CAAGACCGAGAUCCCCAAGAGCUGCUCC HPIV3_F_Codon
AUGCCCAUCAGCAUCCUGCUGAUCAUACCACAAUGAU 64 Optimized mRNA
CAUGGCCAGCCACUGCCAGAUCGACAUCACCAAGCUGC sequence
AGCACGUGGGCGUGCUCGUGAACAGCCCCAAGGGCAU
GAAGAUAGCCAGAACUUCGAGACACGCUACCUGAUC
CUGAGCCUGAUCCCCAAGAUCGAGGACAGCAACAGCU
GCGGCGACCAGCAGAUCAAGCAGUACAAGCGGCUGCU
GGACAGACUGAUCAUCCCCCUGUACGACGGCCUGCGGC
UGCAGAAAGACGUGAUCGUGACCAACCAGGAAAGCAA
CGAGAACACCGACCCCCGGACCGAGAGAUUCUUCGGCG
GCGUGAUCGGCACAUCGCCCUGGGAGUGGCCACAAG
CGCCCAGAUUACAGCCGCUUGUGGCCCUUGGUGGAAGCCA
AGCAGGCCAGAAGCGACAUCGAGAAGCUGAAAGAGGC
CAUCCGGGACACCAACAAGGCCGUGCAGAGCGUGCAG
UCCAGCGUGGGCAAUCUGAUCGUGGCCAUCAAGUCCG
UGCAGGACUACGUGAACAAAGAAAUCGUGCCCUCUAU
CGCCCCGGCUGGGCUGUGAAGCUGCCGGACUGCAGCUG
GGCAUUGCCCUGACACAGCACUACAGCGAGCUGACCAA
CAUCUUCGGCGACAACAUCGGCAGCCUGCAGGAAAAG
GGCAUUAAGCUGCAGGGAAUCGCCAGCCUGUACCGCA
CCAACAUCACCGAGAUUCUACCAACCAGCACCGUGGAU
AAGUACGACAUCUACGACCUGCUGUUCACCGAGAGCA
UCAAAGUGCGCGUGAUCGACGUGGACCUGAACGACUA
CAGCAUCACCCUGCAAGUGCGGCUGCCCCUGCUGACCA
GACUGCUGAACACCCAGAUUAACAAGGUGGACAGCAU
CUCCUACAACAUCCAGAACCGCGAGUGGUACAUCCCUC
UGCCCAGCCACAUAUGACCAAGGGCGCCUUUCUGGGC
GGAGCCGACGUGAAAGAGUGCAUCGAGGCCUUCAGCA
GCUACAUCUGCCCCAGCGACCCUGGCUUCGUGCUGAAC
CACGAGAUGGAAAGCUGCCUGAGCGGCAACAUCAGCC
AGUGCCCCAGAACCACCGUGACCUCCGACAUCGUGCCC
AGAUACGCCUUCGUGAAUGGCGGGCGUGGUGGCCAACU
GCAUCACCACCACCUGUACCUUGCAACGGCAUCGGCAAC
CGGAUCAACCAGCCUCCC GAUCAGGGCGUGAAGAUUA
UCACCCACAAAGAGUGUAACACCAUCGGCAUCAACGGC
AUGCUGUCAAUACCAACAAAGAGGGCACCCUGGCCU
UCUACACCCCCGACGAUAUACCCUGAACAAACUCCGUG
GCUCUGGACCCCAUCGACAUCUCCAUCGAGCUGAACAA
GGCCAAGAGCGACCUGGAAGAGUCCAAAGAGUGGAUC

CGCGGAGCAACGACUGGACUCUAUCGGCAGCU
GGCACCAGAGCAGCACCACCAUCAUCGUGAUCCUGAUU
AUGAUGAUUAUCCUGUUCAUCAUCAACAUUACCAUCA
UCACUAUCGCCAUUAAGUACUACCGGAUCCAGAAACG
GAACCGGGUGGACCAGAAUGACAAGCCCUACGUGCUG ACAAACAAG
(577) TABLE-US-00007 TABLE 6 PIV3 Amino Acid Sequences SEQ ID Description
Sequence NO: >gi|612507166|gb| MPISILLIITTMIMASHCQIDITKLQHVGVLVNSPKGMKISQ
13 AHX22429.1| NFETRYLILSLIPKIEDSNSCGDQQIKQYKRLLDRLIIPLYDG fusion
glycoprotein LRLQKDVIVTNQESNENTDPRTERFFGGVIGTIALGVATSA F0 [Human
QITAAVALVEAKQARSDIEKLKEAIRDTNKAVQSVQSSVG parainfluenza virus
NLIVAIAKSVQDYVNKEIVPSIARLGCEAAGLQLGIALTQHYS 3]
ELTNIFGDNIGSLQEKGIKLQGIASLYRTNITEIFTTSTVDKY
DIYDLLFTESIKVRVIDVDLNDYSITLQVRLPLLTRLLNTQIY
KVDSISYNIQNREWIPLPSHIMTKGAFLGGADVKECIEAFS
SYICPSDPGFVLNHEMESCLSGNISQCPRTTVTSDIVPRYAF
VNGGVVANCITTTCTCNGIGNRINQPPDQGVKIITHKECNTI
GINGMLFNTNKEGTLAFYTPDDITLNNVALDPIDISIELNK
AKSDLEESKEWIRRSNQKLDSIGSWHQSSTTIIVILIMMILFI
INITIITIAIKYYRIQKRNRVDQNDKPYYVLTNK gi|612507167|gb|
MEYWKHTNHGKDAGNELETSTATHGNKLTNKITYILWTIT 14 AHX22430.1|
LVLLSIVFIIVLTNSIKSEKARESLLQDINNEFMEVTEKIQVA hemagglutinin-
SDNTNDLIQSGVNTRLTIQSHVQNYIPISLTQQISDLRKFI neuraminidase
EITIRNDNQEVPPQRITHDVGIKPLNPDDFWRCTSGLPSLMK [Human
TPKIRLMPGPGLLAMPTTVDGCVRTPSLVINDLIYAYTSNLI parainfluenza virus
TRGCQDIGKSYQVLQIGIITVNSDLVPDLNPRISHTFNINDN 3]
RKSCSLALLNTDVYQLCSTPKVDERSDYASSGIEDIVLDIV
NYDGSISTTRFKNNNISFDQPYAALYPSVGPGIYYKGKIIFL
GYGGLEHPINENAICNTTGCPGKTQRDCNQASHSPWFSDR
RMVNSIIVVDKGLNSVPKLKVWTISMQRQNYWGSEGRLLLL
GNKIYIYTRSTSWHSLQLGIIDITDYSIRIKWTWHNVLSR
PGNNECPWGHSCPDCITGVYTDAYPLNPTGSIVSSVILDS
QKSRVNPVITYSTATERVNELAIRNKTLISAGYTTTSCITHY
NKGYCFHIVEINHKS LNTFQPMLFKTEIPKSCS

(578) TABLE-US-00008 TABLE 7 PIV3 NCBI Accession Numbers (Nucleic Acid and Amino
Acid Sequences) GenBank Description Accession Fusion glycoprotein F0 [Human parainfluenza
virus 3] HPIV3/*Homo* KJ672601.1|:4990- *sapiens*/PER/FLA4815/2008 6609 AHX22429 (Fusion
protein) hemagglutinin-neuraminidase [Human parainfluenza virus 3] HPIV3/*Homo*
KJ672601.1|:6724- *sapiens*/PER/FLA4815/2008 8442 AHX22430 (HN protein) Recombinant
PIV3/PIV1 virus fusion glycoprotein (F) and hemagglutinin AF016281 (HN) genes, complete cds;
and RNA dependent RNA polymerase (L) gene, AAC23947 partial cds. (hemagglutinin)
Recombinant PIV3/PIV1 virus fusion glycoprotein (F) and hemagglutinin AF016281 (HN) genes,
complete cds; and RNA dependent RNA polymerase (L) gene, AAC23947 partial cds. (fusion
protein) hemagglutinin-neuraminidase [Human parainfluenza virus 3] BAO32044.1 hemagglutinin-
neuraminidase [Human parainfluenza virus 3] BAO32051.1 C protein [Human parainfluenza virus
3] NP_599251.1 C protein [Human parainfluenza virus 3] ABZ85670.1 C protein [Human
parainfluenza virus 3] AGT75164.1 C protein [Human parainfluenza virus 3] AAB48686.1 C
protein [Human parainfluenza virus 3] AHX22115.1 C protein [Human parainfluenza virus 3]
AGW51066.1 C protein [Human parainfluenza virus 3] AGW51162.1 C protein [Human
parainfluenza virus 3] AGT75252.1 C protein [Human parainfluenza virus 3] AGT75188.1 C
protein [Human parainfluenza virus 3] AGW51218.1 C protein [Human parainfluenza virus 3]

AGW51074.1 C protein [Human parainfluenza virus 3] AGT75323.1 C protein [Human parainfluenza virus 3] AGT75307.1 C protein [Human parainfluenza virus 3] AHX22131.1 C protein [Human parainfluenza virus 3] AGW51243.1 C protein [Human parainfluenza virus 3] AGT75180.1 C protein [Human parainfluenza virus 3] AGT75212.1 C protein [Human parainfluenza virus 3] AGW51186.1 C protein [Human parainfluenza virus 3] AHX22075.1 C protein [Human parainfluenza virus 3] AHX22163.1 C protein [Human parainfluenza virus 3] AGT75196.1 C protein [Human parainfluenza virus 3] AHX22491.1 C protein [Human parainfluenza virus 3] AHX22139.1 C protein [Human parainfluenza virus 3] AGW51138.1 C protein [Human parainfluenza virus 3] AGW51114.1 C protein [Human parainfluenza virus 3] AGT75220.1 C protein [Human parainfluenza virus 3] AHX22251.1 RecName: Full = Protein C; AltName: Full = VP18 protein P06165.1 C protein [Human parainfluenza virus 3] AHX22187.1 C protein [Human parainfluenza virus 3] AGT75228.1 C protein [Human parainfluenza virus 3] AHX22179.1 C protein [Human parainfluenza virus 3] AHX22427.1 C protein [Human parainfluenza virus 3] AGW51210.1 nonstructural protein C [Human parainfluenza virus 3] BAA00922.1 C protein [Human parainfluenza virus 3] AHX22315.1 C protein [Human parainfluenza virus 3] AGW51259.1 C protein [Human parainfluenza virus 3] AHX22435.1 C protein [Human parainfluenza virus 3] AHX22123.1 C protein [Human parainfluenza virus 3] AHX22299.1 C protein [Human parainfluenza virus 3] AGW51267.1 unnamed protein product [Human parainfluenza virus 3] CAA28430.1 C protein [Human parainfluenza virus 3] AGW51178.1 C protein [Human parainfluenza virus 3] AHX22411.1 RecName: Full = Protein C P06164.1 phosphoprotein [Human parainfluenza virus 3] NP_067149.1 phosphoprotein [Human parainfluenza virus 3] AAB48685.1 phosphoprotein [Human parainfluenza virus 3] AHX22498.1 phosphoprotein [Human parainfluenza virus 3] AHX22490.1 phosphoprotein [Human parainfluenza virus 3] AGT75259.1 phosphoprotein [Human parainfluenza virus 3] AGW51137.1 phosphoprotein [Human parainfluenza virus 3] AGW51145.1 phosphoprotein [Human parainfluenza virus 3] AGT75298.1 phosphoprotein [Human parainfluenza virus 3] AGW51113.1 phosphoprotein [Human parainfluenza virus 3] AGT75203.1 phosphoprotein [Human parainfluenza virus 3] AGT75163.1 phosphoprotein [Human parainfluenza virus 3] AHX22506.1 phosphoprotein [Human parainfluenza virus 3] AGW51129.1 phosphoprotein [Human parainfluenza virus 3] AHX22194.1 phosphoprotein [Human parainfluenza virus 3] AGT75211.1 phosphoprotein [Human parainfluenza virus 3] AHX22258.1 phosphoprotein [Human parainfluenza virus 3] AGW51121.1 phosphoprotein [Human parainfluenza virus 3] AGT75282.1 phosphoprotein [Human parainfluenza virus 3] AHX22146.1 phosphoprotein [Human parainfluenza virus 3] AHX22138.1 phosphoprotein [Human parainfluenza virus 3] AHX22322.1 phosphoprotein [Human parainfluenza virus 3] AHX22370.1 phosphoprotein [Human parainfluenza virus 3] AHX22098.1 phosphoprotein [Human parainfluenza virus 3] AHX22130.1 phosphoprotein [Human parainfluenza virus 3] AHX22418.1 phosphoprotein [Human parainfluenza virus 3] AHX22114.1 phosphoprotein [Human parainfluenza virus 3] AHX22410.1 phosphoprotein [Human parainfluenza virus 3] AGT75306.1 phosphoprotein [Human parainfluenza virus 3] AHX22170.1 phosphoprotein [Human parainfluenza virus 3] AHX22266.1 phosphoprotein [Human parainfluenza virus 3] AHX22090.1 phosphoprotein [Human parainfluenza virus 3] AGT75195.1 phosphoprotein [Human parainfluenza virus 3] AHX22226.1 phosphoprotein [Human parainfluenza virus 3] AHX22178.1 phosphoprotein [Human parainfluenza virus 3] AHX22122.1 phosphoprotein [Human parainfluenza virus 3] AHX22186.1 phosphoprotein [Human parainfluenza virus 3] AHX22066.1 phosphoprotein [Human parainfluenza virus 3] AHX22522.1 phosphoprotein [Human parainfluenza virus 3] AGW51225.1 phosphoprotein [Human parainfluenza virus 3] BAN29032.1 phosphoprotein [Human parainfluenza virus 3] ABZ85669.1 phosphoprotein [Human parainfluenza virus 3] AHX22426.1 phosphoprotein [Human parainfluenza virus 3] AHX22058.1 phosphoprotein [Simian Agent 10] ADR00400.1 phosphoprotein [Human parainfluenza virus 3] AHX22250.1 phosphoprotein

[Human parainfluenza virus 3] AHX22434.1 phosphoprotein [Human parainfluenza virus 3] AHX22298.1 phosphoprotein [Human parainfluenza virus 3] AHX22442.1 phosphoprotein [Human parainfluenza virus 3] AHX22074.1 phosphoprotein [Human parainfluenza virus 3] AGW51153.1 phosphoprotein [Human parainfluenza virus 3] AGW51241.1 phosphoprotein [Human parainfluenza virus 3] AHX22210.1 phosphoprotein [Human parainfluenza virus 3] AGW51105.1 phosphoprotein [Human parainfluenza virus 3] AGT75251.1 phosphoprotein [Human parainfluenza virus 3] AHX22362.1 phosphoprotein [Human parainfluenza virus 3] AHX22474.1 phosphoprotein [Human parainfluenza virus 3] AGW51217.1 phosphoprotein [Human parainfluenza virus 3] AIG60038.1 phosphoprotein [Human parainfluenza virus 3] AHX22378.1 phosphoprotein [Human parainfluenza virus 3] AGW51057.1 phosphoprotein [Human parainfluenza virus 3] AGT75187.1 phosphoprotein [Human parainfluenza virus 3] AGW51233.1 phosphoprotein [Human parainfluenza virus 3] AHX22482.1 phosphoprotein [Human parainfluenza virus 3] AGW51161.1 phosphoprotein [Human parainfluenza virus 3] AHX22306.1 phosphoprotein [Human parainfluenza virus 3] AHX22162.1 phosphoprotein [Human parainfluenza virus 3] ACJ70087.1 phosphoprotein [Human parainfluenza virus 3] AHX22466.1 phosphoprotein [Human parainfluenza virus 3] AHX22346.1 phosphoprotein [Human parainfluenza virus 3] AGW51089.1 phosphoprotein [Human parainfluenza virus 3] AGW51073.1 phosphoprotein [Human parainfluenza virus 3] AGW51185.1 phosphoprotein [Human parainfluenza virus 3] AGW51065.1 phosphoprotein [Human parainfluenza virus 3] ABY47603.1 phosphoprotein [Human parainfluenza virus 3] AGW51049.1 phosphoprotein [Human parainfluenza virus 3] AHX22330.1 phosphoprotein [Human parainfluenza virus 3] AGW51250.1 phosphoprotein [Human parainfluenza virus 3] AGT75227.1 phosphoprotein [Human parainfluenza virus 3] AGW51282.1 phosphoprotein [Human parainfluenza virus 3] AGW51209.1 phosphoprotein [Human parainfluenza virus 3] AGW51193.1 phosphoprotein [Human parainfluenza virus 3] AGT75322.1 phosphoprotein [Human parainfluenza virus 3] AGT75219.1 phosphoprotein [Human parainfluenza virus 3] AGW51258.1 phosphoprotein [Human parainfluenza virus 3] AGW51041.1 phosphoprotein [Human parainfluenza virus 3] ACD99698.1 phosphoprotein [Human parainfluenza virus 3] AGW51266.1 phosphoprotein [Human parainfluenza virus 3] AGT75179.1 phosphoprotein [Human parainfluenza virus 3] AHX22282.1 phosphoprotein [Human parainfluenza virus 3] AGW51169.1 phosphoprotein [Human parainfluenza virus 3] AGW51274.1 phosphoprotein [Human parainfluenza virus 3] AGW51201.1 phosphoprotein [Human parainfluenza virus 3] AGW51177.1 RecName: Full = Phosphoprotein; Short = Protein P P06162.1 P protein [Human parainfluenza virus 3] AAA66818.1 phosphoprotein [Human parainfluenza virus 3] AAA46866.1 phosphoprotein [Human parainfluenza virus 3] BAA00031.1 polymerase-associated nucleocapsid phosphoprotein (version 2) - RRNZP5 parainfluenza virus type 3 [Human parainfluenza virus 3] phosphoprotein [Human parainfluenza virus 3] AGT75171.1 phosphoprotein [Human parainfluenza virus 3] BAA00921.1 D protein [Human parainfluenza virus 3] NP_599250.1 D protein [Human parainfluenza virus 3] AHX22377.1 D protein [Human parainfluenza virus 3] AHX22121.1 D protein [Human parainfluenza virus 3] AGT75297.1 D protein [Human parainfluenza virus 3] AGW51136.1 D protein [Human parainfluenza virus 3] AGW51242.1 D protein [Human parainfluenza virus 3] AGW51112.1 D protein [Human parainfluenza virus 3] AHX22497.1 D protein [Human parainfluenza virus 3] AHX22145.1 D protein [Human parainfluenza virus 3] AGT75202.1 D protein [Human parainfluenza virus 3] AHX22385.1 D protein [Human parainfluenza virus 3] AGW51216.1 D protein [Human parainfluenza virus 3] AGT75281.1 D protein [Human parainfluenza virus 3] AGT75194.1 D protein [Human parainfluenza virus 3] AHX22521.1 D protein [Human parainfluenza virus 3] AGW51120.1 D protein [Human parainfluenza virus 3] AGT75313.1 D protein [Human parainfluenza virus 3] AHX22249.1 D protein [Human parainfluenza virus 3] AHX22097.1 D protein [Human parainfluenza virus 3] AGW51144.1 D protein [Human parainfluenza virus 3] AHX22089.1 D protein [Human parainfluenza virus 3]

AHX22225.1 D protein [Human parainfluenza virus 3] AHX22137.1 D protein [Human parainfluenza virus 3] AHX22065.1 D protein [Human parainfluenza virus 3] AGW51224.1 D protein [Human parainfluenza virus 3] AGT75210.1 D protein [Human parainfluenza virus 3] AHX22393.1 D protein [Human parainfluenza virus 3] AGT75258.1 D protein [Human parainfluenza virus 3] AHX22345.1 D protein [Human parainfluenza virus 3] AGT75250.1 D protein [Human parainfluenza virus 3] AHX22113.1 D protein [Human parainfluenza virus 3] AGW51232.1 D protein [Human parainfluenza virus 3] AHX22057.1 D protein [Human parainfluenza virus 3] AHX22209.1 D protein [Human parainfluenza virus 3] AGW51056.1 D protein [Human parainfluenza virus 3] AHX22161.1 D protein [Simian Agent 10] ADR00402.1 D protein [Human parainfluenza virus 3] AHX22361.1 D protein [Human parainfluenza virus 3] AGW51281.1 D protein [Human parainfluenza virus 3] AGW51184.1 D protein [Human parainfluenza virus 3] AGW51160.1 D protein [Human parainfluenza virus 3] AHX22465.1 D protein [Human parainfluenza virus 3] AHX22329.1 D protein [Human parainfluenza virus 3] AGW51064.1 D protein [Human parainfluenza virus 3] AGW51040.1 D protein [Human parainfluenza virus 3] AGT75226.1 D protein [Human parainfluenza virus 3] AHX22425.1 D protein [Human parainfluenza virus 3] AHX22305.1 D protein [Human parainfluenza virus 3] AGW51249.1 D protein [Human parainfluenza virus 3] AHX22481.1 D protein [Human parainfluenza virus 3] AHX22281.1 D protein [Human parainfluenza virus 3] AGW51048.1 D protein [Human parainfluenza virus 3] AHX22297.1 D protein [Human parainfluenza virus 3] AGW51088.1 D protein [Human parainfluenza virus 3] AGT75305.1 D protein [Human parainfluenza virus 3] AHX22185.1 D protein [Human parainfluenza virus 3] AGW51104.1 D protein [Human parainfluenza virus 3] AHX22081.1 D protein [Human parainfluenza virus 3] AGW51192.1 D protein [Human parainfluenza virus 3] AHX22489.1 D protein [Human parainfluenza virus 3] AHX22441.1 D protein [Human parainfluenza virus 3] AHX22409.1 D protein [Human parainfluenza virus 3] AHX22369.1 D protein [Human parainfluenza virus 3] AHX22321.1 D protein [Human parainfluenza virus 3] AHX22073.1 D protein [Human parainfluenza virus 3] AGW51152.1 D protein [Human parainfluenza virus 3] AGW51072.1 D protein [Human parainfluenza virus 3] AGT75321.1 D protein [Human parainfluenza virus 3] AHX22257.1 D protein [Human parainfluenza virus 3] AHX22129.1 D protein [Human parainfluenza virus 3] AHX22417.1 D protein [Human parainfluenza virus 3] AGT75218.1 D protein [Human parainfluenza virus 3] AHX22265.1 D protein [Human parainfluenza virus 3] AGT75178.1 D protein [Human parainfluenza virus 3] AHX22433.1 D protein [Human parainfluenza virus 3] AGW51273.1 D protein [Human parainfluenza virus 3] AGW51208.1 D protein [Human parainfluenza virus 3] AGT75170.1 D protein [Human parainfluenza virus 3] AGT75162.1 D protein [Human parainfluenza virus 3] AGW51257.1 D protein [Human parainfluenza virus 3] AGW51200.1 D protein [Human parainfluenza virus 3] AGW51176.1 D protein [Human parainfluenza virus 3] AGT75186.1 D protein [Human parainfluenza virus 3] AGW51265.1 D protein [Human parainfluenza virus 3] AGW51168.1

(579) TABLE-US-00009 TABLE 8 Signal Peptides SEQ ID Description Sequence NO:
HuIgG.sub.k signal METPAQLLFLLLWLPDTTG 15 peptide IgE heavy chain
MDWTWILFLVAAATRVHS 16 epsilon-1 signal peptide Japanese
MLGSNSGQRVVFTILLLLVAPAYS 17 encephalitis PRM signal sequence VSVg protein
MKCLLYLAFLFIGVNCA 18 signal sequence Japanese MWLVSLAIVTACAGA 19
encephalitis JEV signal sequence

(580) TABLE-US-00010 TABLE 9 hMPV/PIV Cotton Rat Challenge Study Design Group n Test Article [conc]/µg Route Challenge 1 5 Placebo n/a IM hMPV/A2 2 5 hMPV vaccine mRNA 30 IM hMPV/A2 3 5 hMPV vaccine mRNA 15 IM hMPV/A2 4 5 hMPV vaccine mRNA 10 IM hMPV/A2 5 5 hMPV/PIV3 vaccine 30 IM hMPV/A2 mRNA (15/15) 6 5 FI-hMPV n/a IM hMPV/A2 7 5 Placebo n/a IM PIV3 8 5 PIV3 vaccine mRNA 30 IM PIV3 9 5 PIV3 vaccine mRNA 15 IM PIV3 10 5 PIV3 vaccine mRNA 10 IM PIV3 11 5 hMPV/PIV3 vaccine 30 IM PIV3

mRNA (15/15) 12 5 FI-PIV3 n/a IM PIV3 60

(581) TABLE-US-00011 TABLE 10 SEQ ID Strain Nucleic Acid Sequence NO:

Betacoronavirus Nucleic Acid Sequence gb|KJ156934.1|:

ATGATACTCAGTGTCTTCTACTGATGTTCTTGTTAACACC 20 21405-25466 Middle

TACAGAAAGTTACGTTGATGTAGGGCCAGATTCTGTAAAG East respiratory

TCTGCTTGTATTGAGGTTGATATACAACAGACCTTCTTTGA syndrome

TAAAACTTGGCCTAGGCCAATTGATGTTTCTAAGGCTGAC coronavirus

GGTATTATATAACCCTCAAGGCCGTACATATTCTAACATAA isolate

CTATCACTTATCAAGGTCTTTTTCCCTATCAGGGAGACCAT Riyadh_14_2013,

GGTGATATGTATGTTTACTCTGCAGGACATGCTACAGGCA spike protein

CAACTCCACAAAAGTTGTTTGTAGCTAACTATTCTCAGGA (nucleotide)

CGTCAAACAGTTTGCTAATGGGTTTGTCGTCCGTATAGGA

GCAGCTGCCAATTCCACTGGCACTGTTATTATTAGCCCATC

TACCAGCGCTACTATACGAAAAATTTACCCTGCTTTTATGC

TGGGTTCTTCAGTTGGTAATTTCTCAGATGGTAAAATGGG

CCGCTTCTTCAATCATACTCTAGTTCTTTTGCCCGATGGAT

GTGGCACTTTACTTAGAGCTTTTTATTGTATTCTAGAGCCT

CGCTCTGGAAATCATTGTCCTGCTGGCAATTCCTATACTTC

TTTTGCCACTTATCACACTCCTGCAACAGATTGTTCTGATG

GCAATTACAATCGTAATGCCAGTCTGAACTCTTTTAAGGA

GTATTTTAATTTACGTAAGTGCACCTTTATGTACACTTATA

ACATTACCGAAGATGAGATTTTAGAGTGGTTTGGCATTAC

ACAAACTGCTCAAGGTGTTACCTCTTCTCATCTCGGTATG

TTGATTTGTACGGCGGCAATATGTTTCAATTTGCCACCTTG

CCTGTTTATGATACTATTAAGTATTATTCTATCATTCCTCA

CAGTATTCGTTCTATCCAAAGTGATAGAAAAGCTTGGGCT

GCCTTCTACGTATATAAACTTCAACCGTTAACTTTCCTGTT

GGATTTTTCTGTTGATGGTTATATACGCAGAGCTATAGACT

GTGGTTTTAATGATTTGTCACAACTCCACTGCTCATATGAA

TCCTTCGATGTTGAATCTGGAGTTTATTCAGTTTCGTCTTT

CGAAGCAAAACCTTCTGGCTCAGTTGTGGAACAGGCTGAA

GGTGTTGAATGTGATTTTTACCTCTTCTGTCTGGCACACC

TCCTCAGGTTTATAATTTCAAGCGTTTGGTTTTTACCAATT

GCAATTATAATCTTACCAAATTGCTTTCACTTTTTTCTGTG

AATGATTTTACTTGTAGTCAAATATCTCCAGCAGCAATTGC

TAGCAACTGTTATTCTTCACTGATTTTGGATTATTTTTTCAT

ACCCACTTAGTATGAAATCCGATCTCAGTGTTAGTTCTGCT

GGTCCAATATCCCAGTTTAATTATAAACAGTCCTTTTCTAA

TCCCACATGTTTGATCTTAGCGACTGTTCTCATAACCTTA

CTACTATTACTAAGCCTCTTAAGTACAGCTATATTAACAA

GTGCTCTCGTCTTCTTTCTGATGATCGTACTGAAGTACCTC

AGTTAGTGAACGCTAATCAATACTCACCTGTGTATCCATT

GTCCCATCCACTGTGTGGGAAGACGGTGATTATTATAGGA

AACAACCTATCTCCACTTGAAGGTGGTGGCTGGCTTGTTGC

TAGTGGCTCAACTGTTGCCATGACTGAGCAATTACAGATG

GGCTTTGGTATTACAGTTCAATATGGTACAGACACCAATA

GTGTTTGCCCCAAGCTTGAATTTGCTAATGACACAAAAT

TGCCTCTCAATTAGGCAATTGCGTGGAATATTCCCTCTATG

GTGTTTCGGGCCGTGGTGTTTTTCAGAATTGCACAGCTGTA

GGTGTTTCGACAGCAGCGCTTTGTTTATGATGCGTACCAGA

ATTAGTTAGTCTATTGATGGCAACTACTACTGT
CTGCGTGCTTGTGTTAGTGTTTCCTGTTTCTGTTCATCTATGA
TAAAGAACTAAAACCCACGCTACTCTATTTGGTAGTGTT
GCATGTGAACACATTTCTTCTACCATGTCTCAATACTCCCG
TTCTACGCGATCAATGCTTAAACGGCGAGATTCTACATAT
GGCCCCCTTCAGACACCTGTTGGTTGTGTCCTAGGACTTGT
TAATTCCTCTTTGTTTCGTAGAGGACTGCAAGTTGCCTCTCG
GTCAATCTCTCTGTGCTCTTCCTGACACACCTAGTACTCTC
ACACCTCGCAGTGTGCGCTCTGTGCCAGGTGAAATGCGCT
TGGCATCCATTGCTTTTAATCATCCCATTGAGGTTGATCAA
CTTAATAGTAGTTATTTTAAATTAAGTATAACCCACTAATTT
TTCCTTTGGTGTGACTCAGGAGTACATTCAGACAACCATTC
AGAAAGTTACTGTTGATTGTAAACAGTACGTTTGCAATGG
TTTCCAGAAGTGTGAGCAATTACTGCGCGAGTATGGCCAG
TTTTGTTCCAAAATAAACCAGGCTCTCCATGGTGCCAATTT
ACGCCAGGATGATTCTGTACGTAATTTGTTTGCGAGCGTG
AAAAGCTCTCAATCATCTCCTATCATAACCAGGTTTTGGAG
GTGACTTTAATTTGACACTTCTAGAACCTGTTTCTATATCT
ACTGGCAGTCGTAGTGCACGTAGTGCTATTGAGGATTTGC
TATTTGACAAAGTCACTATAGCTGATCCTGGTTATATGCA
AGGTTACGATGATTGTATGCAGCAAGGTCCAGCATCAGCT
CGTGATCTTATTTGTGCTCAATATGTGGCTGGTTATAAAGT
ATTACCTCCTCTTATGGATGTTAATATGGAAGCCGCGTATA
CTTCATCTTTGCTTGGCAGCATAGCAGGTGTTGGCTGGACT
GCTGGCTTATCCTCCTTTGCTGCTATTCCATTTGCACAGAG
TATYTTTTATAGGTAAACGGTGTTGGCATTACTCAACAG
GTTCTTTCAGAGAACC AAAAGCTTATTGCCAATAAGTTTA
ATCAGGCTCTGGGAGCTATGCAAACAGGCTTCACTACAAC
TAATGAAGCTTTTCGGAAGGTTGAGGATGCTGTGAACAAC
AATGCACAGGCTCTATCCAAATTAGCTAGCGAGCTATCTA
ATACTTTTGGTGCTATTTCCGCCTCTATTGGAGACATCATA
CAACGTCTTGATGTTCTCGAACAGGACGCCCAAATAGACA
GACTTATTAATGGCCGTTTGACAACACTAAATGCTTTTGTT
GCACAGCAGCTTGTTTCGTTCCGAATCAGCTGCTCTTTCCGC
TCAATTGGCTAAAGATAAAGTCAATGAGTGTGTCAAGGCA
CAATCCAAGCGTTCTGGATTTTGCGGTCAAGGCACACATA
TAGTGTCCTTTGTTGTAAATGCCCCTAATGGCCTTTACTTT
ATGCATGTTGGTTATTACCCTAGCAACCACATTGAGGTTGT
TTCTGCTTATGGTCTTTGCGATGCAGCTAACCCTACTAATT
GTATAGCCCCTGTTAATGGCTACTTTATTAAAATAATAAC
ACTAGGATTGTTGATGAGTGGTCATATACTGGCTCGTCCTT
CTATGCACCTGAGCCCATCACCTCTCTTAATACTAAGTATG
TTGCACCACAGGTGACATAACCAAACATTTCTACTAACCT
CCCTCCTCCTCTTCTCGGCAATTCACCGGGATTGACTTCC
AAGATGAGTTGGATGAGTTTTTCAAAAATGTTAGCACCAG
TATACCTAATTTTGGTTCTCTAACACAGATTAATACTACAT
TACTCGATCTTACCTACGAGATGTTGTCTCTTCAACAAGTT
GTTAAAGCCCTTAATGAGTCTTACATAGACCTTAAAGAGC
TTGGCAATTATACTTATTACAACAAATGGCCGTGGTACAT
TTGGCTTGGTTTCATTGCTGGGCTTGTTGCCTTAGCTCTAT

CGCTCTTCTTCTACATACTGTGCTGCACTGGTTGTGGGCACAAAC
TGTATGGGAAAACCTTAAGTGTAAATCGTTGTTGTGATAGAT
ACGAGGAATACGACCTCGAGCCGCATAAGGTTTCATGTTCA CTAA MERS S FL
ATGATACACTCAGTGTCTTCTACTGATGTTCTTGTTAACACC 21 SPIKE
TACAGAAAGTTACGTTGATGTAGGGCCAGATTCTGTAAAG 2cEMC/2012
TCTGCTTGTATTGAGGTTGATATACAACAGACTTTCTTTGA (XBal change(T to
TAAAACCTTGGCCTAGGCCAATTGATGTTTCTAAGGCTGAC G)) (nucleotide)
GGTATTATATAACCCTCAAGGCCGTACATATTCTAACATAA
CTATCACTTATCAAGGTCTTTTTCCCTATCAGGGAGACCAT
GGTGATATGTATGTTTACTCTGCAGGACATGCTACAGGCA
CAACTCCACAAAAGTTGTTTGTAGCTAACTATTCTCAGGA
CGTCAAACAGTTTGCTAATGGGTTTGTCTCGTCCGTATAGGA
GCAGCTGCCAATTCCACTGGCACTGTTATTATTAGCCCATC
TACCAGCGCTACTATACGAAAAATTTACCCTGCTTTTATGC
TGGGTTCTTCAGTTGGTAATTTCTCAGATGGTAAAATGGG
CCGCTTCTTCAATCATACTCTAGTTCTTTTGCCCGATGGAT
GTGGCACTTTACTTAGAGCTTTTTATTGTATTCTGGAGCCT
CGCTCTGGAAATCATTGTCCTGCTGGCAATTCCTATACTTC
TTTTGCCACTTATCACACTCCTGCAACAGATTGTTCTGATG
GCAATTACAATCGTAATGCCAGTCTGAACTCTTTTAAGGA
GTATTTTAATTTACGTAACTGCACCTTTATGTACACTTATA
ACATTACCGAAGATGAGATTTTAGAGTGGTTTGGCATTAC
ACAAACTGCTCAAGGTGTTACCTCTTCTCATCTCGGTATG
TTGATTTGTACGGCGGCAATATGTTTCAATTTGCCACCTTG
CCTGTTTATGATACTATTAAGTATTATTCTATCATTCTCA
CAGTATTTCGTTCTATCCAAAGTGATAGAAAAGCTTGGGCT
GCCTTCTACGTATATAAACTTCAACCGTTAACTTTCTCTGTT
GGATTTTTCTGTTGATGGTTATATACGCAGAGCTATAGACT
GTGGTTTTAATGATTTGTACAACTCCACTGCTCATATGAA
TCCTTCGATGTTGAATCTGGAGTTTATTCAGTTTCGTCTTT
CGAAGCAAAACCTTCTGGCTCAGTTGTGGAACAGGCTGAA
GGTGTTGAATGTGATTTTTACCTCTTCTGTCTGGCACACC
TCCTCAGGTTTATAATTTCAAGCGTTTGGTTTTTACCAATT
GCAATTATAATCTTACCAAATTGCTTTCACTTTTTTCTGTG
AATGATTTTACTTGTAGTCAAATATCTCCAGCAGCAATTGC
TAGCAACTGTTATTCTTCACTGATTTTGGATTACTTTTCAT
ACCCACTTAGTATGAAATCCGATCTCAGTGTTAGTTCTGCT
GGTCCAATATCCCAGTTTAATTATAAACAGTCCTTTTCTAA
TCCCACATGTTTGATTTTAGCGACTGTTCCCTCATAACCTTA
CTACTATTACTAAGCCTCTTAAGTACAGCTATATTAACAA
GTGCTCTCGTCTTCTTTCTGATGATCGTACTGAAGTACCTC
AGTTAGTGAACGCTAATCAATACTCACCTGTGTATCCATT
GTCCCATCCACTGTGTGGGAAGACGGTGATTATTATAGGA
AACAACATCTCCACTTGAAGGTGGTGGCTGGCTTGTTGC
TAGTGGCTCAACTGTTGCCATGACTGAGCAATTACAGATG
GGCTTTGGTATTACAGTTCAATATGGTACAGACACCAATA
GTGTTTGCCCCAAGCTTGAATTTGCTAATGACACAAAAT
TGCCTCTCAATTAGGCAATTGCGTGGAATATTCCCTCTATG
GTGTTTCGGGCCGTGGTGTTTTTCAGAATTGCACAGCTGTA
GGTGTTTCGACAGCAGCGCTTTGTTTATGATGCGTACCAGA

ATTTAGTTGATTGATGGTCAACTACTACTGT
TTGCGTGCTTGTGTTAGTGTTCTGTTTCTGTCTATGAT
AAAGAACTAAAACCCACGCTACTCTATTTGGTAGTGTTG
CATGTGAACACATTTCTTCTACCATGTCTCAATACTCCCGT
TCTACGCGATCAATGCTTAAACGGCGAGATTCTACATATG
GCCCCCTTCAGACACCTGTTGGTTGTGTCCTAGGACTTGTT
AATTCCTCTTTGTTCTGCTAGAGGACTGCAAGTTGCCTCTTGG
TCAATCTCTCTGTGCTCTTCCTGACACACCTAGTACTCTCA
CACCTCGCAGTGTGCGCTCTGTTCCAGGTGAAATGCGCTT
GGCATCCATTGCTTTTAATCATCCTATTCAGGTTGATCAAC
TTAATAGTAGTTATTTTAAATTAAGTATACCCACTAATTTT
TCCTTTGGTGTGACTCAGGAGTACATTCAGACAACCATTC
AGAAAGTTACTGTTGATTGTAAACAGTACGTTTGCAATGG
TTTCCAGAAGTGTGAGCAATTACTGCGCGAGTATGGCCAG
TTTTGTTCCAAAATAAACCAGGCTCTCCATGGTGCCAATTT
ACGCCAGGATGATTCTGTACGTAATTTGTTTGCGAGCGTG
AAAAGCTCTCAATCATCTCCTATCATAACCAGGTTTTGGAG
GTGACTTTAATTTGACACTTCTGGAACCTGTTTCTATATCT
ACTGGCAGTCGCTAGTGCACGTAGTGCTATTGAGGATTTGC
TATTTGACAAAGTCACTATAGCTGATCCTGGTTATATGCA
AGGTTACGATGATTGCATGCAGCAAGGTCCAGCATCAGCT
CGTGATCTTATTTGTGCTCAATATGTGGCTGGTTACAAAGT
ATTACCTCCTCTTATGGATGTTAATATGGAAGCCGCGTATA
CTTCATCTTTGCTTGGCAGCATAGCAGGTGTTGGCTGGACT
GCTGGCTTATCCTCCTTTGCTGCTATTCCATTTGCACAGAG
TATCTTTTATAGGTAAACGGTGTGTCATTACTCAACAGG
TTCTTTTCAGAGAACC AAAAGCTTATTGCCAATAAGTTTAA
TCAGGCTCTGGGAGCTATGCAAACAGGCTTCACTACA ACT
AATGAAGCTTTTCAGAAGGTTCAAGGATGCTGTGAACAACA
ATGCACAGGCTCTATCCAAATTAGCTAGCGAGCTATCTAA
TACTTTTGGTGCTATTTCCGCCTCTATTGGAGACATCATA
AACGTCTTGATGTTCTCGAACAGGACGCCCAAATAGACAG
ACTTATTAATGGCCGTTTGACAACACTAAATGCTTTTGTTG
CACAGCAGCTTGTTCTGTTCCGAATCAGCTGCTCTTTCCGCT
CAATTGGCTAAAGATAAAGTCAATGAGTGTGTCAAGGCAC
AATCCAAGCGTTCTGGATTTTGCGGTCAAGGCACACATAT
AGTGTCTTTGTTGTAAATGCCCTAATGGCCTTTACTTCA
TGCATGTTGGTTATTACCCTAGCAACCACATTGAGGTTGTT
TCTGCTTATGGTCTTTGCGATGCAGCTAACCCTACTAATTG
TATAGCCCCTGTTAATGGCTACTTTATTAAA ACTAATAACA
CTAGGATTGTTGATGAGTGGTCATATACTGGCTCGTCCTTC
TATGCACCTGAGCCCATACCTCCCTTAATACTAAGTATGT
TGCACCACAGGTGACATACCAAACATTTCTACTAACCTC
CCTCCTCCTCTTCTCGGCAATTCCACCGGGATTGACTTCCA
AGATGAGTTGGATGAGTTTTTCAAAAATGTTAGCACCAGT
ATACCTAATTTTGGTTCCCTAACACAGATTAATACTACATT
ACTCGATCTTACCTACGAGATGTTGTCTCTTCAACAAGTTG
TTAAAGCCCTTAATGAGTCTTACATAGACCTTAAAGAGCT
TGGCAATTATACTTATTACAACAAATGGCCGTGGTACATT
TGGCTTGGTTTTATTGCTGGGCTTGTGCTTAGCTCTATG

CGTCTTCTCATCTGCTTGTGTCACAACT
GTATGGGAAACTTAAGTGTAATCGTTGTTGTGATAGATA
CGAGGAATACGACCTCGAGCCGCATAAGGTTTCATGTTTAC TAA Novel_MERS_S2_
ATGATCCACTCCGTGTTCTCCTCATGTTCTGTTGACCCC 22 subunit_trimeric
CACTGAGTCAGACTGCAAGCTCCCGCTGGGACAGTCCCTG vaccine
TGTGCGCTGCCTGACACTCCTAGCACTCTGACCCACGCTC (nucleotide)
CGTGCGGTGCGGTGCCTGGCGAAATGCGGCTGGCCTCCATC
GCCTTCAATCACCCAATCCAAGTGGATCAGCTGAATAGCT
CGTATTTCAAGCTGTCCATCCCCACGAACTTCTCGTTTCGGG
GTCACCCAGGAGTACATCCAGACCACAATTCAGAAGGTCA
CCGTCGATTGCAAGCAATACGTGTGCAACGGCTTCCAGAA
GTGCGAGCAGCTGCTGAGAGAATACGGGCAGTTTTGCAGC
AAGATCAACCAGGCGCTGCATGGAGCTAACTTGCGCCAGG
ACGACTCCGTGCGCAACCTCTTTGCCTCTGTGAAGTCATCC
CAGTCCTCCCCAATCATCCCGGGATTTCGGAGGGGACTTCA
ACCTGACCCTCCTGGAGCCCGTGTCGATCAGCACCGGTAG
CAGATCGGCGCGCTCAGCCATTGAAGATCTTCTGTTTCGAC
AAGGTCACCATCGCCGATCCGGGCTACATGCAGGGATACG
ACGACTGTATGCAGCAGGGACCAGCCTCCGCGAGGGACCT
CATCTGCGCGCAATACGTGGCCGGGTACAAAGTGCTGCCT
CCTCTGATGGATGTGAACATGGAGGCCGCTTATACTTCGT
CCCTGCTCGGCTCTATCGCCGGCGTGGGGTGGACCGCCGG
CCTGTCTCCTTCGCCGCTATCCCCTTTGCACAATCCATTT
TCTACCGGCTCAACGGCGTGGGCATTACTCAACAAGTCCT
GTCGGAGAACCAGAAGTTGATCGCAAACAAGTTCAATCA
GGCCCTGGGGGGCCATGCAGACTGGATTCACTACGACTAAC
GAAGCGTTCCAGAAGGTCCAGGACGCTGTGAACAACAAC
GCCCAGGCGCTCTCAAAGCTGGCCTCCGAACTCAGCAACA
CCTTCGGAGCCATCAGCGCATCGATCGGTGACATAATTCA
GCGGCTGGACGTGCTGGAGCAGGACGCCCAGATCGACCG
CCTCATCAACGGACGGCTGACCACCTTGAATGCCTTCGTG
GCACAACAGCTGGTCCGGAGCGAATCAGCGGCACTTTCCG
CCCAACTCGCCAAGGACAAAGTCAACGAATGCGTGAAGG
CCCAGTCCAAGAGGTCCGGTTTCTGCGGTCAAGGAACCCA
TATTGTGTCCTTCGTGCTGAACGCGCCCAACGGTCTGTACT
TTATGCACGTCGGCTACTACCCGAGCAATCATATCGAAGT
GGTGTCCGCCTACGGCCTGTGCGATGCCGCTAACCCCACT
AACTGTATTGCCCCTGTGAACGGATATTTTATTAAGACCA
ACAACACCCGCAATTGTGGACGAATGGTCATACACCGGTTC
GTCCTTCTACGCGCCCGAGCCCATCACTTCACTGAACACC
AAATACGTGGCTCCGCAAGTGACCTACCAGAACATCTCCA
CCAATTTGCCGCCGCGCTGCTCGGAAACAGCACCGGAAT
TGATTTCCAAGATGAACTGGACGAATTCTTCAAGAACGTG
TCCACTTCCATTCCCAACTTCGGAAGCCTGACACAGATCA
ACACCACCCTTCTCGACCTGACCTACGAGATGCTGAGCCT
TCAACAAGTGGTCAAGGCCCTGAACGAGAGCTACATCGAC
CTGAAGGAGCTGGGCAACTATACCTACTACAACAAGTGGC
CGGACAAGATTGAGGAGATTCTGTGCGAAAATCTACCACAT
TGAAAACGAGATCGCCAGAATCAAGAAGCTTATCGGCGA AGCC MERS_S0_Full-
ATGGAAACCCCTGCCCAGCTGCTGTTCTGCTGCTGCTGTG 23 length Spike

GCTGCTGCTATACCTAGCTATGTGGACGTGGGCCCC protein
GATAGCGTGAAGTCCGCCTGTATCGAAGTGGACATCCAGC (nucleotide, codon
AGACCTTTTTTCGACAAGACCTGGCCCAGACCCATCGACGT optimized)
GTCCAAGGCCGACGGCATCATCTATCCACAAGGCCGGACC
TACAGCAACATCACCATTACCTACCAGGGCCTGTTCCCAT
ATCAAGGCGACCACGGCGATATGTACGTGTACTCTGCCGG
CCACGCCACCGGCACCACACCCCAGAACTGTTTCGTGGCC
AACTACAGCCAGGACGTGAAGCAGTTCGCCAACGGCTTCG
TCGTGCGGATTGGCGCCGCTGCCAATAGCACCGGCACAGT
GATCATCAGCCCCAGCACCGCGCCACCATCCGGAAGATC
TACCCCGCCTTCATGCTGGGCAGCTCCGTGGGCAATTTCA
GCGACGGCAAGATGGGCCGGTTCTTCAACCACACCCTGGT
GCTGCTGCCCCGATGGCTGTGGCACACTGCTGAGAGCCTTC
TACTGCATCCTGGAACCCAGAAGCGGCAACCACTGCCCTG
CCGGCAATAGCTACACCAGCTTCGCCACCTACCACACACC
CGCCACCGATTGCTCCGACGGCAACTACAACCGGAACGCC
AGCCTGAACAGCTTCAAAGAGTACTTCAACCTGCGGAACT
GCACCTTCATGTACACCTACAATATCACCGAGGACGAGAT
CCTGGAATGGTTCGGCATCACCCAGACCGCCCAGGGCGTG
CACCTGTTTCAGCAGCAGATACGTGGACCTGTACGGCGGCA
ACATGTTCCAGTTTGCCACCCTGCCCCGTGTACGACACCATC
AAGTACTACAGCATCATCCCCACAGCATCCGGTCCATCC
AGAGCGACAGAAAAGCCTGGGCGCGCCTTCTACGTGTACAA
GCTGCAGCCCCCTGACCTTCCTGCTGGACTTCAGCGTGGAC
GGCTACATCAGACGGGCCATCGACTGCGGCTTCAACGACC
TGAGCCAGCTGCACTGCTCCTACGAGAGCTTCGACGTGGA
AAGCGGCGTGTACAGCGTGTCCAGCTTCGAGGCCAAGCCT
AGCGGCAGCGTGGTGGAACAGGCTGAGGGCGTGGAATGC
GACTTCAGCCCTCTGCTGAGCGGCACCCCTCCCCAGGTGT
ACAACCTTCAAGCGGCTGGTGTTCAACCAACTGCAATTACAA
CCTGACCAAGCTGCTGAGCCTGTTCTCCGTGAACGACTTC
ACCTGTAGCCAGATCAGCCCTGCCGCCATTGCCAGCAACT
GCTACAGCAGCCTGATCCTGGACTACTTCAGCTACCCCCCT
GAGCATGAAGTCCGATCTGAGCGTGTCTCCGCCGGACCC
ATCAGCCAGTTCAACTACAAGCAGAGCTTCAGCAACCCTA
CCTGCCTGATTCTGGCCACCGTGCCCCACAATCTGACCAC
CATCACCAAGCCCCTGAAGTACAGCTACATCAACAAGTGC
AGCAGACTGCTGTCCGACGACCGGACCGAAGTGCCCCAGC
TCGTGAACGCCAACCAGTACAGCCCCTGCGTGTCCATCGT
GCCCAGCACCGTGTGGGAGGACGGCGACTACTACAGAAA
GCAGCTGAGCCCCCTGGAAGGCGGCGGATGGCTGGTGGCT
TCTGGAAGCACAGTGGCCATGACCGAGCAGCTGCAGATG
GGCTTTGGCATCACCGTGCAGTACGGCACCGACACCAACA
GCGTGTGCCCCAAGCTGGAATTCGCCAATGACACCAAGAT
CGCCAGCCAGCTGGGAAACTGCGTGGAATACTCCCTGTAT
GGCGTGTCCGGACGGGGCGTGTTCCAGAATTGCACAGCAG
TGGGAGTGCGGCAGCAGAGATTCGTGTACGATGCCTACCA
GAACCTCGTGGGCTACTACAGCGACGACGGCAATTACTAC
TGCCTGCGGGCCTGTGTGTCCGTGCCCCGTGTCCGTGATCTA
CGACAAAGAGACAAAGACCCACGCCACACTGTTTCGGCTCC

GTGGCCTGACCTCCAGCTCCACCATGAGCCAGTACT
CCCGCTCCACCCGGTCCATGCTGAAGCGGAGAGATAGCAC
CTACGGCCCCCTGCAGACACCTGTGGGATGTGTGCTGGGC
CTCGTGAACAGCTCCCTGTTTGTGGAAGATTGCAAGCTGC
CCCTGGGCCAGAGCCTGTGTGCCCTGCCAGATACCCCTAG
CACCTGACCCCTAGAAGCGTGCGCTCTGTGCCCCGGCGAA
ATGCGGCTGGCCTCTATCGCCTTCAATCACCCCATCCAGGT
GGACCAGCTGAACTCCAGCTACTTCAAGCTGAGCATCCCC
ACCAACTTCAGCTTCGGCGTGACCCAGGAGTACATCCAGA
CCACAATCCAGAAAGTGACCGTGGAAGTGAAGCAGTACGT
GTGCAACGGCTTTCAGAAGTGCGAACAGCTGCTGCGCGAG
TACGGCCAGTTCTGCAGCAAGATCAACCAGGCCCTGCACG
GCGCCAACCTGAGACAGGATGACAGCGTGCGGAACCTGTT
CGCCAGCGTGAAAAGCAGCCAGTCCAGCCCCATCATCCCT
GGCTTCGGCGGCGACTTTAACCTGACCCTGCTGGAACCTG
TGTCCATCAGCACCGGCTCCAGAAGCGCCAGATCCGCCAT
CGAGGACCTGCTGTTCGACAAAGTGACCATTGCCGACCCC
GGCTACATGCAGGGCTACGACGATTGCATGCAGCAGGGCC
CAGCCAGCGCCAGGGATCTGATCTGTGCCCAGTATGTGGC
CGGCTACAAGGTGCTGCCCCCCCCTGATGGACGTGAACATG
GAAGCCGCCTACACCTCCAGCCTGCTGGGCTCTATTGCTG
GCGTGGGATGGACAGCCGGCCTGTCTAGCTTTGCCGCCAT
CCCTTTCGCCCAGAGCATCTTCTACCGGCTGAACGGCGTG
GGCATCACACAACAGGTGCTGAGCGAGAACCAGAAGCTG
ATCGCCAACAAGTTTAACCAGGCACTGGGCGCCATGCAGA
CCGGCTTCACCACCACCAACGAGGCCTTCAGAAAGGTGCA
GGACGCCGTGAACAACAACGCCCAGGCTCTGAGCAAGCT
GGCCTCCGAGCTGAGCAATACCTTCGGCGCCATCAGCGCC
TCCATCGGCGACATCATCCAGCGGCTGGACGTGCTGGAAC
AGGACGCCCAGATCGACCGGCTGATCAACGGCAGACTGA
CCACCCTGAACGCCTTCGTGGCACAGCAGCTCGTGCGGAG
CGAATCTGCCGCTCTGTCTGCTCAGCTGGCCAAGGACAAA
GTGAACGAGTGCGTGAAGGCCCAGTCCAAGCGGAGCGGC
TTTTGTGGCCAGGGCACCCACATCGTGTCTTCGTCTGAA
TGCCCCCAACGGCCTGTACTTTATGCACGTGGGCTATTACC
CCAGCAACCACATCGAGGTGGTGTCCGCCTATGGCCTGTG
CGACGCCGCCAATCCTACCAACTGTATCGCCCCCGTGAAC
GGCTACTTCATCAAGACCAACAACACCCGGATCGTGGACG
AGTGGTCCTACACAGGCAGCAGCTTCTACGCCCCCGAGCC
CATCACCTCCCTGAACACCAAATACGTGGCCCCCACAAGTG
ACATACCAGAACATCTCCACCAACCTGCCCCCTCCACTGC
TGGGAAATTCCACCGGCATCGACTTCCAGGACGAGCTGGA
CGAGTTCTTCAAGAACGTGTCCACCTCCATCCCCAACTTCG
GCAGCCTGACCCAGATCAACACCACTCTGCTGGACCTGAC
CTACGAGATGCTGTCCCTGCAACAGGTTCGTGAAAGCCCTG
AACGAGAGCTACATCGACCTGAAAGAGCTGGGGAACTAC
ACCTACTACAACAAGTGGCCTTGGTACATTTGGCTGGGCT
TTATCGCCGGCCTGGTGGCCCTGGCCCTGTGCGTGTTCTTC
ATCCTGTGCTGCACCGGCTGCGGCACCAATTGCATGGGCA
AGCTGAAATGCAACCGGTGCTGCGACAGATACGAGGAAT

ACGACCTGGACCTAGTGCATGTGCAC Betacoronavirus mRNA Sequences
gb|KJ156934.1|: AUGAUACACUCAGUGUUUCUACUGAUGUUCUUGUUAAC 65 21405-
25466 Middle ACCUACAGAAAGUUACGUUGAUGUAGGGGCCAGAUUCUG East
respiratory UUAAGUCUGCUUGUAUUGAGGUUGAUUAACAACAGACC syndrome
UUCUUUGAUAAAACUUGGCCUAGGCCAAUUGAUGUUUC coronavirus
UAAGGCUGACGGUAUUUAUUAUACCCUCAAGGCCGUACAU isolate
AUUCUAACAUAACUAUCACUUAUCAAGGUCUUUUUCCCCU Riyadh_14_2013,
AUCAGGGAGACCAUGGUGAUUAUGUAUGUUUACUCUGCA spike protein
GGACAUGCUACAGGCACAACUCCACAAAAGUUGUUUGU (nucleotide)
AGCUAACUAUUCUCAGGACGUCAAACAGUUUUGC UAAUG
GGUUUGUCGUCCGUUAGGAGCAGCUGCCAAUUCCACUG
GCACUGUUAUUUAUUAGCCCAUCUACCAGCGCUACUUAUAC
GAAAAAUUUACCCUGCUUUUAUGCUGGGUUCUUCAGUU
GGUAAUUUCUCAGAUUGGUAAAAUUGGGCCGCUUCUUCAA
UCAUACUCUAGUUCUUUUGCCCGAUGGAUGUGGCACUU
UACUUAGAGCUUUUUUAUUGUAUUCUAGAGCCUCGCUCU
GGAAAUCAUUGUCCUGCUGGCAAUUCCUAUACUUCUUU
UGCCACUUAUCACACUCCUGCAACAGAUUGUUCUGAUGG
CAAUUACAAUCGUAAUGCCAGUCUGAACUCUUUUAAAGG
AGUAUUUUAAUUUACGUAAUCUGCACCUUUAUGUACACU
UAUAACAUAUACCGAAGAUGAGAUUUUAGAGUGGUUUGG
CAUUACACAAACUGCUCUAAAGGUGUUCACCUCUUCUCAUC
UCGGUAUGUUGAUUUUGUACGGCGGCAAUAUGUUUCAAU
UUGCCACCUUGCCUGUUUAUGAUACUAUUAAGUAUUAU
UCUAUCAUUCCUCACAGUAUUCGUUCUAUCCAAAGUGAU
AGAAAAGCUUGGGCUGCCUUCUACGUAAUUAUAAACUUCA
ACCGUUAACUUUCCUGUUGGAUUUUUCUGUUGAUGGUU
AUUAACGCAGAGCUAUAGACUGUGGUUUUAAUGAUUUG
UCACAACUCCACUGCUCUAUAUGAAUCCUUCGAUGUUGAA
UCUGGAGUUUAUUCAGUUUCGUCUUUCGAAGCAAACCC
UUCUGGCUCAGUUGUGGAACAGGCUGAAGGUGUUGAAU
GUGAUUUUUCACCUCUUCUGUCUGGCACACCUCUCAGG
UUUAUAAUUUCAAGCGUUUGGUUUUUACCAAUUGCAAU
UAUAAUCUUACCAAUUGCUUUCACUUUUUUCUGUGAA
UGAUUUUACUUGUAGUCAAAUAUCUCCAGCAGCAAUUG
CUAGCAACUGUUUAUUCUUCACUGAUUUUUGGAUUAUUUU
UCAUACCCACUAGUAUGAAAUCCGAUCUCAGUGUUAG
UUCUGCUGGUCCAAUAUCCCAGUUUAAUUAUAAACAGU
CCUUUUUCUAAUCCCACAUGUUUGAUCUUAGCGACUGUUC
CUCAUAACCUUACUACUAUUACUAAGCCUCUUAAGUACA
GCUAUUAUUAACAAGUGCUCUCGUCUUCUUCUGAUGAU
CGUACUGAAGUACCUCAGUUAGUGAACGCUAAUCAUA
CUCACCCUGUGUAUCCAUUGUCCCAUCCACUGUGUGGGA
AGACGGUGAUUAUUUAUAGGAAACAACUAUCUCCACUUG
AAGGUGGUGGCUGGCUUGUUGCUAGUGGCUCAACUGUU
GCCAUGACUGAGCAAUUACAGAUGGGCUUUGGUAAUUA
AGUUCAAUAUGGUACAGACACCAAUAGUGUUUGCCCCA
AGCUUGAAUUUGCUAAUGACACAAAAAUUGCCUCUCA
UUAGGCAAUUGCGUGGAAUAUUCCCUCUAUGGUGUUUC
GGGCCGUGGUGUUUUUCAGAAUUGCACAGCUGUAGGUG

UUCGACAGUAGCGUACCCAGAAU
UUAGUUGGCUAUUAUUCUGAUGAUGGCAACUACUACUG
UCUGCGUGCUUGUGUAGUGUCCUGUUUCUGUCAUCU
AUGAUAAAGAAACUAAAACCCACGCUACUCUAUUUGGU
AGUGUUGCAUGUGAACACAUAUUCUUCUACCAUGUCUCA
AUACUCCCGUUCUACGCGAUCAAUGCUUAAACGGCGAGA
UUCUACAUAUGGGCCCCCUUCAGACACCUGUUGGUUGUGU
CCUAGGACUUGUUAUUCUUCUUGUUCGUAGAGGACU
GCAAGUUGCCUCUCGGUCAUUCUCUCUGUGCUCUUCUG
ACACACCUAGUACUCUCACACCUCGCGAGUGUGCGCUCUG
UGCCAGGUGAAAUGCGCUUGGCAUCCAUUGCUUUUAAU
CAUCCCAUUCAGGUUGAUCAACUUAUAGUAGUUUUU
UAAAUUAAGUAUACCCACUAAUUUUUCCUUUGGUGUGA
CUCAGGAGUACAUUCAGACAACCAUUCAGAAAGUUACU
GUUGAUUGUAAACAGUACGUUUGCAAUGGUUUCAGAA
GUGUGAGCAAUACUGCGCGAGUAUGGCCAGUUUUGUU
CCAAAAUAAACCAGGCUCUCCAUGGUGCCAAUUUACGCC
AGGAUGAUUCUGUACGUAAUUUGUUUGCGAGCGUGAAA
AGCUCUCAAUCAUCUCCUAUCAUACCAGGUUUUGGAGGU
GACUUUAAUUUGACACUUCUAGAACCUGUUUCUAUAUC
UACUGGCAGUCGUAGUGCACGUAGUGCUAUUGAGGAUU
UGCUAUUUGACAAAGUCACUAUAGCUGAUCCUGGUUAU
AUGCAAGGUUACGAUGAUUGUAUGCAGCAAGGUCCAGC
AUCAGCUCGUGAUCUUAUUUGUGCUCAAUAUGUGGCUG
GUUAUAAAGUAUUACCUCUUAUGGAUGUUAAUAUG
GAAGCCGCGUAUACUUCAUCUUUGCUUGGCAGCAUAGCA
GGUGUUGGCUGGACUGCUGGCUUAUCCUCCUUUGCUGCU
AUUCCAUUUGCACAGAGUAUYUUUUUAUAGGUUAAACGG
UGUUGGCAUUACUCAACAGGUUCUUUCAGAGAACCAAA
AGCUUAUUGCCAUAUAGUUUAUUCAGGCUCUGGGAGCU
AUGCAAACAGGCUUCACUACAACUAAUGAAGCUUUUCG
GAAGGUUCAGGAUGCUGUGAACAACAAUGCACAGGCUC
UAUCCAAAUUAGCUAGCGAGCUAUCUAAUACUUUUGGU
GCUAUUUCCGCCUCUAUUGGAGACAUCAUACAACGUCUU
GAUGUUCUCGAACAGGACGCCCAAUAGACAGACUUUAU
UAAUGGCCGUUUGACAACACUAAAUGCUUUUUGUUGCAC
AGCAGCUUGUUCGUUCCGAAUCAGCUGCUCUUUCCGCUC
AAUUGGCUAAAGAUAAAGUCAUAGAGUGUGUCAAGGCA
CAAUCCAAGCGUUCUGGAUUUUGCGGUCAAGGCACACAU
AUAGUGUCCUUUGUUGUAAAUGCCCCUAAUGGCCUUUA
CUUUUAGCAUGUUGGUUAUUACCCUAGCAACCACAUUG
AGGUUGUUUCUGCUUAUGGUCUUUUGCGAUGCAGCUAAC
CCUACUAAUUGUAUAGCCCCUGUUAUUGGCUACUUUAU
UAAAACUAAUAACACUAGGAUUGUUGAUGAGUGGUCAU
AUACUGGCUCGUCCUUCUAUGCACCUGAGCCCAUCACCU
CUCUUAUACUAAGUAUGUUGCACCACAGGUGACAUACC
AAAACAUUUCUACUAACCUCCCUCCUCUUCUCGGCA
AUUCCACCGGGAUUGACUCCAAGAUGAGUUGGAUGAG
UUUUUCAAAAAUGUUAGCACCAGUAUACCUAAUUUUGG
UUCUCUAAACACAGAUUAUACUACAUAUCUCGAUCUAC

CUACGAGAUUUAACAAGUUGUUAAAGCCC
UUAUGAGUCUUACAAGACCUUAAAGAGCUUGGCAAU
UAUACUUAUUACAACAAUGGCCGUGGUACA UUUGGCU
UGGUUUCAUUGCUGGGCUUGUUGCCUUAAGCUCUAUGCG
UCUUCUUCAUACUGUGCUGCACUGGUUGUGGCACAAACU
GUAUGGGGAAACUUAAGUGUAAUCGUUGUUGUGAUAGA
UACGAGGAAUACGACCUCGAGCCGCAUAAGGUUCAUGU UCACUAA MERS S FL
AUGAUACACUCAGUGUUUCUACUGAUGUUCUUGUUAAC 66 SPIKE
ACCUACAGAAAGUUACGUUGAUGUAGGGGCCAGAUUCUG 2cEMC/2012
UUAAGUCUGCUUGUAUUGAGGUUGAUUAACAACAGACU (XBaI change(U
UUCUUUGAUAAAACUUGGCCUAGGCCAAUUGAUGUUUC to G)) (nucleotide)
UAAGGCUGACGGUAUUUAUUAACCCUCAAGGCCGUACAU
AUUCUAACAUAACUAUCACUUAUCAAGGUCUUUUUCCCU
AUCAGGGAGACCAUGGUGAUUAUGUAUGUUUACUCUGCA
GGACAUGCUACAGGCACAACUCCACAAAAGUUGUUUGU
AGCUAACUAUUCUCAGGACGUCAAACAGUUUGC UAAUG
GGUUUGUCGUCCGUUAUAGGAGCAGCUGCCAAUUCCACUG
GCACUGUUAUUUAUUAAGCCCAUCUACCAGCGCUACUAUAC
GAAAAAUUUACCCUGCUUUUAUGCUGGGUUCUUCAGUU
GGUAAUUUCUCAGAUUGGUAAAAUGGGCCGCUUCUUCAA
UCAUACUCUAGUUCUUUUGCCCGAUGGAUGUGGCACUU
UACUUAGAGCUUUUUUAUUGUAUUCUGGAGCCUCGCUCU
GGAAAUCAUUGUCCUGCUGGCAAUUCCUAUACUUCUUU
UGCCACUUAUCACACUCCUGCAACAGAUUGUUCUGAUGG
CAAUUACAAUCGUAAUGCCAGUCUGAACUCUUUUAAAGG
AGUAUUUUAAUUUACGUAAACUGCACCUUUUAUGUACACU
UAUAACAUAUACCGAAGAUGAGAUUUUAGAGUGGUUUGG
CAUUACACAAACUGCUCAAGGUGUUCACCUCUUCUCAUC
UCGGUAUGUUGAUUUUGUACGGCGGCAAUAUGUUUCAAU
UUGCCACCUUGCCUGUUUAUGAUACUAUUAAGUAUUUAU
UCUAUCAUUCUCACAGUAUUCGUUCUAUCCAAAGUGAU
AGAAAAGCUUGGGCUGCCUUCUACGUAAUAUAAACUUCA
ACCGUUAACUUUCCUGUUGGAUUUUUCUGUUGAUGGUU
AUUAACGCAGAGCUAUAGACUGUGGUUUUAAUGAUUUG
UCACAACUCCACUGCUCAUAUGAAUCCUUCGAUGUUGAA
UCUGGAGUUUAUUCAGUUUCGUCUUUCGAAGCAAACCC
UUCUGGCUCAGUUGUGGAACAGGCUGAAGGUGUUGAAU
GUGAUUUUUCACCUCUUCUGUCUGGCACACCUCUCAGG
UUUAUAAUUUCAAGCGUUUGGUUUUUACCAAUUGCAAU
UAUAAUCUUAACCAAUUGCUUUCACUUUUUUCUGUGAA
UGAUUUUACUUGUAGUCAAAUAUCUCCAGCAGCAAUUG
CUAGCAACUGUUAUUCUUCACUGAUUUUUGGAUUACUUU
UCAUACCCACUUAAGUAUGAAAUCCGAUCUCAGUGUUAG
UUCUGCUGGUCCAAUAUCCAGUUUAAUUAUAAACAGU
CCUUUUUCUAAUCCCAUGUUUGAUUUUAGCGACUGUUC
CUCAUAACCUUACUACUAUUACUAAGCCUCUUAAGUACA
GCUAUUAUAAACAAGUGCUCUCGUCUUCUUCUGAUGAU
CGUACUGAAGUACCUCAGUUAGUGAACGCUAAUCAUA
CUCACCCUGUGUAUCCAUUGUCCCAUCCACUGUGUGGGA
AGACGGUGAUUAUUAUAGGAAACAACUAUCUCCACUUG

[illegible]

UACUGGCUACGACUGGACCCCAUUAACCUCCUUAACUACUAAGUAUGUUGCACCACAGGUGACAUACCA
AAACAUUUCUACUAACCUCUCCUCCUCUUCUCGGCAA
UUCCACCGGGAUUGACUUCCAAGAUGAGUUGGAUGAGU
UUUUCAAAAAUGUUAAGCACCAGUAUACCUAUUUUUGGU
UCCCUAACACAGAUUAAUACUACAUUACUCGAUCUUACC
UACGAGAUGUUGUCUCUUAACAAGUUGUUAAGCCCU
UAAUGAGUCUACAUAGACCUUAAAGAGCUUGGCAAU
AUACUUAUUACAACAAAUGGCCGUGGUACAUUUGGCU
GGUUUCAUUGCUGGGCUUGUUGCCUAGCUCUAUGCGU
CUUCUUAUACUGUGCUGCACUGGUUGUGGCACAAACUG
UAUGGGAAAACUUAAGUGUAAUCGUUGUUGUGAUAGAU
ACGAGGAUACGACCUCGAGCCGCAUAAGGUUCAUGUUC ACUAA Novel_MERS_S2_
AUGAUCCACUCCGUGUUCCUCCUCAUGUUCCUGUUGACC 67 subunit_trimeric
CCCACUGAGUCAGACUGCAAGCUCCCGCUGGGACAGUCC vaccine
CUGUGUGCGCUGCCUGACACUCCUAGCACUCUGACCCCA (nucleotide)
CGCUCCGUGCGGUCGGUGCCUGGCGAAAUGCGGCUGGCC
UCCAUCGCCUUCAAUCACCCAUCCAAGUGGAUCAGCUG
AAUAGCUCGUAAUUAAGCUGUCCAUCCCCACGAACUUC
UCGUUCGGGGUCACCCAGGAGUACAUCCAGACCACAAU
CAGAAGGUCACCGUCGAUUGCAAGCAAUACGUGUGCAAC
GGCUUCCAGAAGUGCGAGCAGCUGCUGAGAGAAUACGG
GCAGUUUUGCAGCAAGAUAACCAGGCGCUGCAUGGAGC
UAACUUGCGCCAGGACGACUCCGUGCGCAACCUCUUUGC
CUCUGUGAAGUCAUCCAGUCCUCCCCAAUCAUCCCGGG
AUUCGGAGGGGACUUAACCUGACCCUCCUGGAGCCCGU
GUCGAUCAGCACCGGUAGCAGAUCCGGCGCGCUCAGCCAU
UGAAGAUCUUCUGUUCGACAAGGUCACCAUCGCCGAUCC
GGGCUACAUGCAGGGAUACGACGACUGUAUGCAGCAGG
GACCAGCCUCCGCGAGGGACCUCAUCUGCGCGCAAUACG
UGGCCGGGUACAAAGUGCUGCCUCCUCUGAUGGAUGUG
AACAUGGAGGCCGCUUAUACUUCGUCCUGCUCGGCUCU
AUCGCCGGCGUGGGGUGGACCGCCGGCCUGUCCUCCUUC
GCCGCUAUCCCCUUGCACAAUCCAUUUUCUACCGGCUC
AACGGCGUGGGCAUUAACUCAACAAGUCCUGUCGGAGAAC
CAGAAGUUGAUCGCAAACAAGUUAUACAGGCCCUUGG
GGCCAUGCAGACUGGAUUCACUACGACUAAACGAAGCGUU
CCAGAAGGUCCAGGACGCUGUGAACAACAACGCCAGGC
GCUCUCAAAAGCUGGCCUCCGAACUCAGCAACACCUUCGG
AGCCAUCAGCGCAUCGAUCGGUGACAUAUUCAGCGGCU
GGACGUGCUGGAGCAGGACGCCCAGAUCGACCGCCUCAU
CAACGGACGGCUGACCACCUUGAAUGCCUUCGUGGCACA
ACAGCUGGUCCGGAGCGAAUCAGCGGCACUUUCCGCCCA
ACUCGCCAAGGACAAAGUCAACGAAUGCGUGAAGGCCCA
GUCCAAGAGGUCCGGUUUCUGCGGUCAAGGAACCCAUAU
UGUGUCCUUCGUCGUGAACGCGCCCAACGGUCUGUACUU
UAUGCACGUCGGCUACUACCCGAGCAAUCAUAUCGAAGU
GGUGUCCGCCUACGGCCUGUGCGAUGCCGCUAACCCAC
UAACUGUAUUGCCCCUGUGAACGGAUAUUUUAUUAAGA
CCAACAACACCCGCAUUGUGGACGAAUGGUCAUACACCG

GUUCGUCUCCAGCGCCCAUCACUUCACUGA
ACACCAAUACGUGGCUCCGCAAGUGACCUACCAGAACA
UCUCCACCAAUUUGCCGCCGCCGUGCUCGGAAACAGCA
CCGGAAUUGAUUUCCAAGAUGAACUGGACGAAUUCUUC
AAGAACGUGUCCACUUCCAUUCCCAACUUCGGAAGCCUG
ACACAGAUCAACACCACCCUUCUCGACCUGACCUACGAG
AUGCUGAGCCUUCAACAAGUGGUCAAGGCCUGAACGAG
AGCUACAUCGACCUGAAGGAGCUGGGCAACUAUACCUAC
UACAACAAGUGGCCGGACAAGAUGAGGAGAUUCUGUC
GAAAUCUACCACAUUGAAAACGAGAUCGCCAGAAUCA
AGAAGCUUAUCGGCGAAGCC MERS_S0_Full-
AUGGAAACCCUGCCCAGCUGCUGUCCUGCUGCUGCUG 68 length Spike
UGGCUGCCUGAUACCACCGGCAGCUAUGUGGACGUGGGC protein
CCCGAUAGCGUGAAGUCCGCCUGUAUCGAAGUGGACAUC (nucleotide, codon
CAGCAGACCUUUUUCGACAAGACCUGGCCCAGACCCAUC optimized)
GACGUGUCCAAGGCCGACGGCAUCAUCUAUCCACAAGGC
CGGACCUACAGCAACAUCACCAUUACCUACCAGGGCCUG
UUCCCAUAUCAAGGCGACCACGGCGAUUGUACGUGUAC
UCUGCCCGCCACGCCACCGGCACCACACCCAGAAACUG
UUCGUGGCCAACUACAGCCAGGACGUGAAGCAGUUCGCC
AACGGCUUCGUCGUGCGGAUUGGCGCCGCUGCCAAUAGC
ACCGGCACAGUGAUCAUCAGCCCCAGCACCAGCGCCACC
AUCCGGAAGAUCUACCCCGCCUUCAUGCUGGGCAGCUCC
GUGGGCAAUUUCAGCGACGGCAAGAUGGGCCGGUUCUU
CAACCACACCCUGGUGCUGCUGCCCGAUGGCUGUGGCAC
ACUGCUGAGAGCCUUCUACUGCAUCCUGGAACCCAGAAG
CGGCAACCACUGCCCUGCCGGCAAUAGCUACACCAGCUU
CGCCACCUACCACACACCCGCCACCGAUUGCUCCGACGG
CAACUACAACCGGAACGCCAGCCUGAACAGCUUCAAGA
GUACUUAACCGUGCGGAACUGCACCUUCAUGUACACCUA
CAAUAUCACCGAGGACGAGAUCCUGGAAUGGUUCGGCA
UCACCCAGACCGCCCAGGGCGUGCACCUGUUCAGCAGCA
GAUACGUGGACCUGUACGGCGGCAACAUGUUCAGUUU
GCCACCCUGCCCGUGUACGACACCAUCAAGUACUACAGC
AUCAUCCCCCACAGCAUCCGGUCCAUCCAGAGCGACAGA
AAAGCCUGGGCCGCCUUCUACGUGUACAAGCUGCAGCCC
CUGACCUUCCUGCUGGACUUCAGCGUGGACGGCUACAUC
AGACGGGCCAUCGACUGCGGCUUCAACGACCUGAGCCAG
CUGCACUGCUCCUACGAGAGCUUCGACGUGGAAAGCGGC
GUGUACAGCGUGUCCAGCUUCGAGGCCAAGCCUAGCGGC
AGCGUGGUGGAACAGGCUGAGGGCGUGGAAUGCGACUU
CAGCCCUCUGCUGAGCGGCACCCCUCCCCAGGUGUACAA
CUUCAAGCGGCUGGUGUUCACCAACUGCAAUACAACCU
GACCAAGCUGCUGAGCCUGUUCUCCGUGAACGACUUCAC
CUGUAGCCAGAUCAGCCCUGCCGCCAUUGCCAGCAACUG
CUACAGCAGCCUGAUCCUGGACUACUUCAGCUACCCCU
GAGCAUGAAGUCCGAUCUGAGCGUGUCCUCCGCCGGACC
CAUCAGCCAGUUCAACUACAAGCAGAGCUUCAGCAACCC
UACCUGCCUGAUUCUGGCCACCGUGCCCCACAAUCUGAC
CACCAUCACCAAGCCCCUGAAGUACAGCUACAUCAACAA

GUGAGCAGACUGACACCGGACCGAAGUGCC
CCAGCUCGUGAACGCCAACCAGUACAGCCCCUGCGUGUC
CAUCGUGCCCAGCACCGUGUGGGAGGACGGCGACUACUA
CAGAAAGCAGCUGAGCCCCCUGGAAGGCGGCGGAUGGCU
GGUGGCUUCUGGAAGCACAGUGGCCAUGACCGAGCAGCU
GCAGAUGGGCUUUGGCAUCACCGUGCAGUACGGCACCGA
CACCAACAGCGUGUGCCCCAAGCUGGAAUUCGCCAAUGA
CACCAAGAUCGCCAGCCAGCUGGGAAACUGCGUGGAAUA
CUCCCUGUAUGGCGUGUCCGGACGGGGCGUGUUCCAGAA
UUGCACAGCAGUGGGAGUGCGGCAGCAGAGAUUCGUGU
ACGAUGCCUACCAGAACCUCGUGGGCUACUACAGCGACG
ACGGCAAUACUACUGCCUGCGGGCCUGUGUGUCCGUGC
CCGUGUCCGUGAUCUACGACAAAGAGACAAAGACCCACG
CCACACUGUUCGGCUCGUGGGCCUGCGAGCACAUACAGCU
CCACCAUGAGCCAGUACUCCCGCUCCACCCGGUCCAUGC
UGAAGCGGAGAGAUAGCACCUACGGCCCCCUGCAGACAC
CUGUGGGAUGUGUGCUGGGCCUCGUGAACAGCUCCCUGU
UUGUGGAAGAUUGCAAGCUGCCCCUGGGCCAGAGCCUGU
GUGCCCUGCCAGAUACCCCUAGCACCCUGACCCCUAGAA
GCGUGCGCUCUGUGCCCCGGCGAAAUGCGGCUGGGCCUCUA
UCGCCUUCAAUCACCCCAUCCAGGUGGACCAGCUGAACU
CCAGCUACUUCAAGCUGAGCAUCCCCACCAACUUCAGCU
UCGGCGUGACCCAGGAGUACAUCCAGACCACAAUCCAGA
AAGUGACCGUGGACUGCAAGCAGUACGUGUGCAACGGC
UUUCAGAAGUGCGAACAGCUGCUGCGCGAGUACGGCCAG
UUCUGCAGCAAGAUCAACCAGGCCCUGCACGGCGCCAAC
CUGAGACAGGAUGACAGCGUGCGGAACCUGUUCGCCAGC
GUGAAAAGCAGCCAGUCCAGCCCCAUCAUCCCUGGCUUC
GGCGGCGACUUUAACCUGACCCUGCUGGAACCUGUGUCC
AUCAGCACCGGCUCCAGAAGCGCCAGAUCCGCCAUUCGAG
GACCUGCUGUUCGACAAAGUGACCAUUGCCGACCCCGGC
UACAUGCAGGGCUACGACGAUUGCAUGCAGCAGGGCCCA
GCCAGCGCCAGGGAUCUGAUCUGUGCCCAGUAUGUGGCC
GGCUACAAGGUGCUGCCCCCCCUGAUGGACGUGAACAUG
GAAGCCGCCUACACCUCCAGCCUGCUGGGCUCUAUUGCU
GGCGUGGGAUGGACAGCCGGCCUGUCUAGCUUUGCCGCC
AUCCCUUUCGCCCAGAGCAUCUUCUACCGGCUGAACGGC
GUGGGCAUCACACAACAGGUGCUGAGCGAGAACCAGAA
GCUGAUCGCCAACAAGUUUAACCAGGCACUGGGCGCCAU
GCAGACCGGCUUCACCACCACCAACGAGGCCUUCAGAAA
GGUGCAGGACGCCGUGAACAACAACGCCCAGGCUCUGAG
CAAGCUGGCCUCCGAGCUGAGCAAUACCUUCGGCGCCAU
CAGCGCCUCCAUCGGCGACAUAUCCAGCGGCUGGACGU
GCUGGAACAGGACGCCCAGAUUCGACCGGCUGAUCAACGG
CAGACUGACCACCCUGAACGCCUUCGUGGCACAGCAGCU
CGUGCGGAGCGAAUCUGCCGCUCUGUCUGCUCAGCUGGC
CAAGGACAAAGUGAACGAGUGCGUGAAGGCCCAGUCCA
AGCGGAGCGGCUUUUGUGGGCCAGGGCACCCACAUCGUGU
CCUUCGUCGUGAAUGCCCCCAACGGCCUGUACUUUAUGC
ACGUGGGCUAUUACCCAGCAACCACAUCGAGGUGGUGU

CCGCCUACCGCCGCCGCGCCGCAUCCUACCCAACU
GUAUCGCCCCCGUGAACGGCUACUUCAAGACCAACA
ACACCCGGAUCGUGGACGAGUGGUCCUACACAGGCAGCA
GCUUCUACGCCCCCGAGCCCAUACCUCCCUGAACACCA
AAUACGUGGCCCCCACAAGUGACAUACCAGAACAUUCUCCA
CCAACCUGCCCCCUCCACUGCUGGGGAAAUUCCACCGGCA
UCGACUUCAGGACGAGCUGGACGAGUUCUUCAAGAACG
UGUCCACCUCCAUCCCCAACUUCGGCAGCCUGACCCAGA
UCAACACCACUCUGCUGGACCUGACCUACGAGAUGCUGU
CCCUGCAACAGGUCGUGAAAGCCCUGAACGAGAGCUACA
UCGACCUGAAAGAGCUGGGGAACUACACCUACUACAACA
AGUGGCCUUGGUACAUUUGGCUGGGCUUUUAUCGCCGGCC
UGGUGGCCUGGGCCUGUGCGUGUUCUUCAUCCUGUGCU
GCACCGGCUGCGGCACCAAUUGCAUGGGCAAGCUGAAAU
GCAACCGGUGCUGCGACAGAUACGAGGAUACGACCUGG
AACCUCACAAAGUGCAUGUGCAC

(582) TABLE-US-00012 TABLE 11 Betacoronavirus Amino Acid Sequences SEQ ID
Strain Amino Acid Sequence NO: gb|KJ156934.1|:

MIHSVFLLMFLLTPTESYVDVGPDSVKSACIEVDIQQTFFDKT 24 21405-25466 Middle
WPRPIDVSKADGIIYPQGRITYSNITITYQGLFPYQGDHGDY East respiratory
VYSAGHATGTTpQKLFVANYSQDVKQFANGFVVRIGAAANS syndrome
TGTVIIISPSTSATIRKIYPFMLGSSVGNFSDGKMGRFFNHTL coronavirus
VLLPDGCGTLLRAFYCILEPRSGNHCPAGNSYTSFATYHTPA isolate
TDCSDGNYNRNASLNSFKEYFNLRNCTFMYTYNITEDEILEW Riyadh_14_2013,
FGITQTAQGVHLFSSRYVDLYGGNMFQFATLPVYDTIKYYSII spike protein
PHSIRSIQSDRKAWAAFYVYKLQPLTFLDFSVDGYIRRAIDC (amino acid)
GFNDLSQLHCSYESFDVESGVYSVSSFEAKPSGSVVEQAEGV
ECDFSPLLSGTPPQVYNFKRLVFTNCNYNLTCLLSLFSVNDFt
CSQISPAAIASNCYSSLILDYFSYPLSMKSDLSVSSAGPISQFN
YKQSFSNPTCLILATVPHNLTTITKPLKYSYINKCSRLLSDDRT
EVPQLVNANQYSPCVSIVPSTVWEDGDYRKQLSPLEGGGW
LVASGSTVAMTEQLQMGGGITVQYGTDTNSVCPKLEFANDT
KIASQLGNCVEYSLYGVSGRGVFQNCTAVGVRQQRVYDA
YQNLVGYYSDDGNYCLRACVSPVSVIYDKETKTHATLFG
SVACEHISSTMSQYSRSTRSMLKRRDSTYGPLQTPVGCVLGL
VNSSLFVEDCKLPLGQSLCALPDTPTSTLTPRSVRSVPGEMRLA
SIAFNHPIQVDQLNSSYFKLSIPTNFSFGVTQEYIQTTIQKVTV
DCKQYVCNGFQKCEQLLREYQGFCSKINqALHGANLRQDDS
VRNLFASVKSSQSSPIPGFGGDFNLTLLEPVSISTGSRARS
EDLLFDKVTIADPGYMQGYDDCMQQGPASARDLICAQYVA
GYKVLPLMDVNMEAAYTSSLLGSIAGVGWTAGLSSFAAIPF
AQSFYRLNGVGITQQVLSENQKLIANKFNQALGAMQTGFTT
TNEAFrKVQDAVNNNAQALSKLASELSNTFGAISASIGDIIQR
LDVLEQDAQIDRLINGRLTTLNAFVAQQLVRSESAALSAQLA
KDKVNECVKAQSKRSGFCGQGTHIVSFVVNAPNGLYFMHV
GYYPNHIENVVSAYGLCDAANPTNCIAPVNGYFIKTNNTRIV
DEWSYTGSSFYAPEPITSLNTKYVAPQVTYQNISTNLPPPLLG
NSTGIDFQDELDEFFKNVSTSI PNFGSLTQINTTLLDLTYEMLS
LQQVVKALNESYIDLKELGNYTYYNKWPWYIWLGFIAGLVA
LALCVFFILCCTGCGTNCMGKCLKNRCCDRYEEYDLEPHKV HVH MERS S FL

MIHVSFLLMFLLTPTESDVGPDSVKACIEVDIQQTFFDKT 25 SPIKE
WPRPIDVSKADGHIYPQGRITYSNITITYQGLFPYQGDHGDYMY 2cEMC/2012
VYSAGHATGTTPQKLFVANYSQDVVKQFANGFVVRIGAAANS (XBal change(T to
TGTVIIISPSTSATIRKIYPAFMLGSSVGNFSDGKMGRFFNHTL G)) (amino acid)
VLLPDGCGTLLRAFYCILEPRSGNHCPAGNSYTSFATYHTPA
TDCSDGNYNRNASLNSFKEYFNLRNCTFMYTYNITEDEILEW
FGITQTAQGVHLFSSRYVDLYGGNMFQFATLPVYDTIKYYYSII
PHSIRSIQSDRKAWAAFYVYKLQPLTFLLDVSVDGYIRRAIDC
GFNDLSQLHCSYESFDVESGVYSVSSFEAKPSGSVVEQAEGV
ECDFSPLLSGTPPQVYNFKRLVFTNCNYNLTKLLSLFSVNDFT
CSQISPAAIASNCYSSLILDYFSYPLSMKSDLSVSSAGPISQFN
YKQSFNPTCLILATVPHNLTTITKPLKYSYINKCSRLLSDDRT
EVPQLVNANQYSPCVSIVPSTVWEDGDYRQKQLSPLEGGGW
LVASGSTVAMTEQLQMGGFQITVQYGTDTNSVCPKLEFANDT
KIASQLGNCVEYSLYGVSGRGVFQNCTAVGVRQQRFFVYDA
YQNLVGYYSDDGNYCLRACVSVPSVIYDKETKTHATLFG
SVACEHISSTMSQYSRSTRSMLKRRDSTYGPLQTPVGCVLGL
VNSSLFVEDCKLPLGQSLCALPDTPSTLTTPRSVRSVPGEMRLA
SIAFNHPIQVDQLNSSYFKLSIPTNFSFGVTQEYIQTTIQKVTV
DCKQYVCNGFQKCEQLLREYGGQFCSKINQALHGANLRQDDS
VRNLFASVKSSQSSPIIPGFGGDFNLTLLEPVSISTGSRARS
EDLLFDKVTIADPGYMQGYDDCMQQGPASARDLICAQYVA
GYKVLPLMDVNMEAAAYTSSLLGSIAGVGWTAGLSSFAAIPF
AQSIIFYRLNGVGITQQVLSENQKLIANKFNQALGAMQTGFTT
TNEAFQKVQDAVNNAQALSKLASELSNTFGAISASIGDIIQR
LDVLEQDAQIDRLINGRLTTLNAFVAQQLVRSESAALSAQLA
KDKVNECVKAQSKRSGFCGQGTHIVSFVVNAPNGLYFMHV
GYPSNHIEVVSAYGLCDAANPTNCIAPVNGYFIKTNNTRIV
DEWSYTGSSFYAPEPITSLNTKYVAPQVTYQNI
STNLPPPLLG
NSTGIDFQDELDEFFKNVSTSI
PNFGSLTQINTTL
LDLT
YEMLSLQQVVKALNESYIDLKELGNYTYYNKWPWYIWLGF
IAGLVALALCVFFILCCTGCGTNCMGLKLCNRCCDRYEEYDLEPHKV
HVVH Novel_MERS_S2_MIHVSFLLMFLLTPTESDCKLPLGQSLCALPDTPSTLT
TPRSVR 26 subunit_trimeric SVPGEMRLASIAFNHPIQVDQLNSSYFKLSIPTNFSFGVTQEYI
vaccine (amino QTTIQKVTV
DCKQYVCNGFQKCEQLLREYGGQFCSKINQALH acid)
GANLRQDDSVRNLFASVKSSQSSPIIPGFGGDFNLTLLEPVSIS
TGSRARS
AIEDLLFDKVTIADPGYMQGYDDCMQQGPASARDLICAQYVAGYKVLPLMDVNMEAAAYTSSLLGSIAGVGWTA
GLSSFAAIPFAQSIIFYRLNGVGITQQVLSENQKLIANKFNQALGAMQTGFTTTNEAFQKVQDAVNNAQALSKLASELSNTFG
AISASIGDIIQR
LDVLEQDAQIDRLINGRLTTLNAFVAQQLVRS
ESAALSAQLAKDKVNECVKAQSKRSGFCGQGTHIVSFVVNAPNGLYFMHVG
YPSNHIEVVSAYGLCDAANPTNCIAPVNGYFIKTNNTRIV
DEWSYTGSSFYAPEPITSLNTKYVAPQVTYQNI
STNLPPPLLG
NSTGIDFQDELDEFFKNVSTSI
PNFGSLTQINTTL
LDLT
YEMLSLQQVVKALNESYIDLKELGNYTYYNKWPDKIE
EILSKIYHIENEIARIKKLIGEA Isolate A1-MIHVSFLLMFLLTPTESYVDVGPDSVKACIEVDIQQTFFDKT 27 Hasa_1_2013
WPRPIDVSKADGHIYPQGRITYSNITITYQGLFPYQGDHGDYMY (NCBI accession
VYSAGHATGTTPQKLFVANYSQDVVKQFANGFVVRIGAAANS #AGN70962)

TGTVIIISPSTSATIRKIYP AFMLGSSVGNFSDGKMGRFFNHTL
VLLPDGCGTLLRAFYCILEPRSGNHCPAGNSYTSFATYHTPA
TDCSDGNYNRNASLNSFKEYFNLRNCTFMYTYNITEDEILEW
FGITQTAQGVHLFSSRYVDLYGGNMFQFATLPVYDTIKYYYSII
PHSIRSIQSDRKAWAAFYVYKLQPLTFLLDFSVDGYIRRAIDC
GFNDLSQLHCSYESFDVESGVYSVSSFEAKPSGSVVEQAEGV
ECDFSPLLSTPPQVYNFKRLVFTNCNYNLTKLLSLFSVNDFT
CSQISPAAIASNCYSSLILDYFSYPLSMKSDLSVSSAGPISQFN
YKQSFNPTCLILATVPHNLTTITKPLKYSYINKCSRLLSDDRT
EVPQLVNANQYSPCVSIVPSTVWEDGDYRQKQLSPLEGGGW
LVASGSTVAMTEQLQMGGFQITVQYGTDTNSVCPKLEFANDT
KIASQLGNCVEYSLYGVSGRGVFQNCTAVGVRQQRFFVYDA
YQNLVGYYSDDGNYCLRACVSPVSVIYDKETKTHATLFG
SVACEHISSTMSQYSRSTRSMLKRRDSTYGPLQTPVGCVLGL
VNSSLFVEDCKLPLGQSLCALPDTPSTLTPRSVRSVPGEMRLA
SIAFNHPIQVDQLNSSYFKLSIPTNFSFGVTQEYIQTTIQKVTV
DCKQYVCNGFQKCEQLLREYGQFCSKINQALHGANLRQDDS
VRNLFASVKSSQSSPIPGFGGDFNLTLLEPVSISTGSRARS
EDLLFDKVTIADPGYMQGYDDCMQQGPASARDLICAQYVA
GYKVLPLMDVNMEAAYTSSLLGSIAGVGWTAGLSSFAAIPF
AQSFYRLNGVGITQQVLSENQKLIANKFNQALGAMQTGFTT
TNEAFRKVQDAVNNAQAALSKLASELSNTFGAISASIGDIIQR
LDVLEQDAQIDRLINGRLTTLNAFVAQQLVRSESAALSAQLA
KDKVNECVKAQSKRSGFCGQGTHIVSFVVNAPNGLYFMHV
GYPSNHIEVVSAAYGLCDAANPTNCIAPVNGYFIKTNNTRIV
DEWSYTGSSFYAPEPITSLNTKYVAPHVTYQNISTNLPPPLLG
NSTGIDFQDELDEFFKNVSTSIPNFGSLTQINTTLLDLTYEMLS
LQQVVKALNESYIDLKELGNYTYYNKWPWYIWLGFIAGLVA
LALCVFFILCCTGCGTNCMGLKLCNRCCDRYEEYDLEPHKV HVH Middle East
MIHSVFLLMFLLTPTESYVDVGPDSVKSACIEVDIQQTFFDKT 28 respiratory
WPRPIDVSKADGHIYPQGRITYSNITITYQGLFPYQGDHGD MY syndrome
VYSAGHATGTTPQKLFVANYSQDVKQFANGFVVRIGAAANS coronavirus S
TGTVIIISPSTSATIRKIYP AFMLGSSVGNFSDGKMGRFFNHTL protein
VLLPDGCGTLLRAFYCILEPRSGNHCPAGNSYTSFATYHTPA UniProtKB -
TDCSDGNYNRNASLNSFKEYFNLRNCTFMYTYNITEDEILEW R9UQ53
FGITQTAQGVHLFSSRYVDLYGGNMFQFATLPVYDTIKYYYSII
PHSIRSIQSDRKAWAAFYVYKLQPLTFLLDFSVDGYIRRAIDC
GFNDLSQLHCSYESFDVESGVYSVSSFEAKPSGSVVEQAEGV
ECDFSPLLSTPPQVYNFKRLVFTNCNYNLTKLLSLFSVNDFT
CSQISPAAIASNCYSSLILDYFSYPLSMKSDLSVSSAGPISQFN
YKQSFNPTCLILATVPHNLTTITKPLKYSYINKCSRLLSDDRT
EVPQLVNANQYSPCVSIVPSTVWEDGDYRQKQLSPLEGGGW
LVASGSTVAMTEQLQMGGFQITVQYGTDTNSVCPKLEFANDT
KIASQLGNCVEYSLYGVSGRGVFQNCTAVGVRQQRFFVYDA
YQNLVGYYSDDGNYCLRACVSPVSVIYDKETKTHATLFG
SVACEHISSTMSQYSRSTRSMLKRRDSTYGPLQTPVGCVLGL
VNSSLFVEDCKLPLGQSLCALPDTPSTLTPRSVRSVPGEMRLA
SIAFNHPIQVDQLNSSYFKLSIPTNFSFGVTQEYIQTTIQKVTV
DCKQYVCNGFQKCEQLLREYGQFCSKINQALHGANLRQDDS
VRNLFASVKSSQSSPIPGFGGDFNLTLLEPVSISTGSRARS

EDLLFDKVDYCMQYQGPASARDLICAQYVA
GYKVLPLMDVNMEAAYTSSLLGSIAGVGWTAGLSSFAAIPF
AQSFYRLNGVGITQQVLSENQKLIANKFNQALGAMQTGFTT
TNEAFRKVQDAVNNNAQALSKLASELSNTFGAISASIGDIIQR
LDVLEQDAQIDRLINGRLTTLNAFVAQQLVRSESAALSAQLA
KDKVNECVKAQSKRSGFCGQGTHIVSFVVNAPNGLYFMHV
GYYP SNHIEVV SAYGLCDAANPTNCIAPVNGYFIKTNNTRIV
DEWSYTGSSFYAPEPITSLNTKYVAPHV TYQNISTNLPPPLLG
NSTGIDFQDELDEFFKNVSTSIPNFGSLTQINTTLLDLTYEMLS
LQQVVKALNESYIDLKELGNYTYYNKWPWYIWLGFIAGLVA
LALCVFFILCCTGCGTNCMGKLCNRCDDRYEEYDLEPHKV HVH Human SARS
MFIFLLFLTLTSGSDLDRCTTFDDVQAPNYTQHTSSMRGVYY 29 coronavirus
PDEIFRSDTLYLTQDLFLPFYSNVTGFHTINHTFGNPVIPFKDG (SARS-CoV)
IYFAATEKSNVVRGWVFGSTMNNKSQSVIINNSTNVVIRAC (Severe acute
NFELCDNPFFAVSKPMGTQTHTMIFDNAFNCTFEYISDAFSLD respiratory
VSEKSGNFKHLREFVFKNKDGLYVYKGYQPIDVVRDLPSGF syndrome
NTLKPIFKLPLGINITNFRAILTAFSPAQDIWGTSAAYFVGYL coronavirus)
KPTTFMLKYDENG TITDAVDCSQNPLAELKCSVK SFEIDKGI Spike
YQTSNFRVVP SGDVVRFPNITNLCPFGEVFNATKFPSVY AWE glycoprotein
RKKISNCVADYSVLYNSTFFSTFKCYGVSATKLNDLCFSNVY UniProtKB-
ADSFVVKGDDVRQIAPGQTGVIADYNYKLPDDFMGCVLAW P59594
NTRNIDATSTGNYNYKYRYLRH GKLRPFERDISNVPFSPDGK
PCTPPALNCYWPLNDYGFYTTTGIGYQPYRVVLSFELLNAP
ATVCGPKLSTD LIKNQCVNFNFNGLTGTGVLT PSSKRFQPFQ
QFGRDVSDFTDSVRDPKTSEILDISPCSFGGVSVITPGTNASSE
VAVLYQDVNCTDVSTAIHADQLTPAWRIYSTGNNVFQTQAG
CLIGAEHVDTSYECDIPIGAGICASYHTVSLLRSTSQKSIVAYT
MSLGADSSIAYSNNTIAIPTNFSISITTEVMPV SMAKTSVDCN
MYICGDSTECANLLLQYGSFCTQLNRALSGIAAEQDRNTREV
FAQVKQMYKTPTLKYFGGFNFSQILPDPLKPTKRSFIEDLLFN
KVTLADAGFMKQYGECLGDINARDLICAQKFNGLTVLPPLL
TDDMIAAYTAALVSGTATAGWTFGAGAALQIPFAMQMAYR
FNGIGVTQNVLYENQKQIANQFNKAISQIQESLTTTSTALGKL
QDVVNQNAQALNTLVKQLSSNFGAISSVLNDILSR LDKVEAE
VQIDRLITGRLQSLQTYVTQQLIRAAEIRASANLAATKMSEC
VLGQSKRVD FCGKGYHLMSPQAAPHGVVFLHVTYVPSQER
NFTTAPAICHEGKAYFPREGVFVFNGTSWFITQRNFFSPQIITT
DNTFVSGNCDVVIGIINNTVYDPLQPELDSFKEELDKYFKNH
TSPDVDLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDLQE
LGKYEQYIKWPWYVWLGFIAGLIAIVMVTILLCCMTSCCSCL
KGACSCGSCCKFDEDDSEPV LKGVKLHYT Human
MFLILLISLPTAFAVIGDLKCTSDNINDKDTGPPPISTDTVDVT 30 coronavirus OC43
NGLGTYYYVLDRVYLN TTFLNGYYPTSGSTYRNMALKGSVL (HCoV-OC43)
LSRLWFKPPFLSDFINGIFAKVKNTKVIKDRVMYSEFP AITIGS Spike
TFVNTSYSVVVQPR TINSTQDGDNKLQGLLEV SVCQYNMCE glycoprotein
YPQTICHPNLGNHRKELWHLDTGVVSCLYKRNFTYDVNAD UniProtKB-
YLYFHFYQEGGTFYAYFTDTGVVTKFLFNVYLG MALSHYYV P36334
MPLTCNSKLTLEYWVTP LTSRQYLLAFNQDGIIFNAEDCMSD
FMSEIKCKTQSIAPPTGVYELNGYTVQPIADVYRRKPNLPNC
NIEAWLNDKSVPSPLNWERKTFSNCNFMSSLM SFIQADSFT

CNNIDAAKIACMFSSSTIDFAIPNGRKNVDLQLGNLGYLQSF
NYRIDTTATSCQLYYNLPAAANVSVSRLFNPSTWNKRFGFIEDS
VFKPRPAGVLTNHDVVYAQHCFAKPNFCPCKLNGSCVGS
PGKNNIGITCPAGTNYLTCDNLCTPDPITFTGTYKCPQTKSL
VGIGEHCSEGLAVKSDYCGGNSCTCRPQAFLGWSADSCSQGD
KCNIFANFILHDVNSGLTCSTDLQKANTDILGVCVNYDLYGI
LGQGIFVEVNATYYNSWQNLLYDSNGNLYGFRDYIINRTFMI
RSCYSGRVSAAFHANSSEPALLFRNIKCNYVFNNSLTRQLQPI
NYFDSYLGCVVNAYNSTAISVQTCDLTVGSGYCVDYSKNRR
SRGAITTYRFTNFEPFTVNSVNDLSLEPVGGLYEIQIPSEFTIG
NMVEFIQTSSPKVTIDCAAFVCGDYAACKSQLVEYGSFCDNI
NAILTEVNELLDTTQLQVANSLMNGVTLSTKLKDGVNFNVD
DINFSPVLGCLGSECSKASSRSAIEDLLFDKVKLSDVGFVEAY
NNCTGGAEIRDLCVQSYKGIVLPPLLENQISGYTLAATSA
SLFPPWTAAAGVPFYLNVQYRINGLGVTMDVLSQNQKLIAN
AFNNALYAIQEGFDATNSALVKIQAVVNANAEALNNLLQQL
SNRFGAISASLQEILSRLEDAEAQIDRLINGRLTALNAYVS
QQLSDSTLVKFSAQAAMEKVNCEVKSQSSRINFCGNGNHIIS
LVQNAPYGLYFIHFSYVPTKYVTARVSPGLCIAGDRGIAPKS
GYFVNVNNTWMYTGSYYYPEPITENNVVVMSTCAVNYTK
APYVMLNTSIPNLPDFKEELDQWFKNQTSVAPDLSLDYINVT
FLDLQVEMNRLQEAIVLNQSYINLKDIGTYEYVVKWPWYV
WLLICLAGVAMLVLLFFICCTGCGTSCFKKCGGCCDDYTGYQELVIKTSHDD Human
MFLIIFILPTTLAVIGDFNCTNSFINDYNKTIPISEDVVDVSLG 31 coronavirus
LGTYYYVLNRVYLNNTLLFTGYFPKSGANFRDLALKGSYILST HKU1 (isolate
LWYKPPFLSDFNNGIFSKVKNTKLYVNNTLYSEFSTIVIGSVF N5) (HCoV-
VNTSYTIVVQPHNGILEITACQYTMCEYPHTVCKSKGSIRNES HKU1) Spike
WHIDSSEPLCLFKKNFTYNVSADWLYFHFYQERGVFYAYYA glycoprotein
DVGMPPTTFLFSLYLGTLSHYYVMPLTCNAISSNTDNETLEY UniProtKB-
WVTPLSRRQYLLNFDEHGVITNAVDCSSSFLSEIQCKTQSFAP Q0ZME7
NTGVYDLSGFTVKPVATVYRRIPNLPDCDIDNWLNNVSVPS
LNWERRIFSNCNFNLSLRLVHVDSEFCNNLDKSKIFGSCFN
SITVDKFAIPNRRRDDLQLGSSGFLQSSNYKIDISSSSCQLYYS
LPLVNVTINNFPSSWNRRYGFGSFNLSSYDVVYSDHCFSVN
SDFCPCADPSVVNSCAKSKPPSAICPAGTKYRHCDLDTTLYV
KNWCRCSCLPDPITYSPNTCPQKKVVVGIGEHCPLGINEE
KCGTQLNHSSCFCSPDAFLGWSFDSCISNNRCNIFSNFIFNGIN
SGTTCSNDLLYSNTEISTGVCVNYDLYGITGQGIFKEVSAAY
YNNWQNLLYDSNGNIIGFKDFLTNKTYTILPCYSGRVSAAFY
QNSSSPALLYRNLKCSYVLNNISFISQPFYFDSYLGCVLNAV
LTSYSVSSCDLRMGSGFCIDYALPSSRRKRRGISSPYRFVTFEP
FNVSFVNDVSVETVGGLEFIQIPTNFTIAGHEEFIQTSSPKVTIDC
SAFVCSNYAACHDLLSEYGTFCDNINSILNEVNDLLDITQLQV
ANALMQGVTLSSNLNTNLHSDVDNIDFKSLLGCLGSQCGSS
RSLLEDLLFNKVKLSDVGFVEAYNNCTGGSEIRDLLCVQSFN
GIKVLPPILSETQISGYTTAATVAAMFPPWSAAAGVPFSLNVQ
YRINGLGVTMDVLNKNQKLIANAFNKALLSIQNGFTATNSAL
AKIQSVVNANAQALNSLLQQLFNKFGAISSSLQEILSRLDNLE
AQVQIDRLINGRLTALNAYVSQQLSDITLIKAGASRAIEKVNE
CVKSQSPRINFCGNGNHILSLVQNAPYGLLFIHFSYKPTSFKT

VLVSPGLDGLSGDRGIAPKQGYFIKQNDGLQSWDMFTGSSYYYPEPIS
DKNVVFMMNSCSVNFTKAPFIYLNNSIPNLSDFEAEKSLWFKN
HTSIAPNLTFNHINATFLDLYYEMNVIQESIKSLNSSFINLKEI
GTYEMYVKWPWYIWLIVILFIIFLMILFFICCCTGCGSACFSK
CHNCCDEYGGHNDFVIKASHDD Novel_SARS_S2
MFIFLLFLTTLTSGSDLDRALSGIAAEQDRNTREVFAQVKQMY 32
KTPTLKYFGGFNFSQILPDPLKPTKRSFIEDLLFNKVTLADAG
FMKQYGECLGDINARDLICAQKFNGLTVLPPLLTDDMIAAYT
AALVSGTATAGWTFGAGAALQIPFAMQMAYRFNGIGVTQN
VLYENQKQIANQFNKAISQIQESLTTTSTALGKLQDVVNQNA
QALNTLVKQLSSNFGAISSVLNDILSRLDKVEAEVQIDRLITG
RLQSLQTYVTQQLIRAAEIRASANLAATKMSECVLGQSKRV
DFCGKGYHLMSPQAAPHGVVFLHVTYVPSQERNFTTAPAIC
HEGKAYFPREGVFVFNGTSWFITQRNFFSPQIITDNTFVSGN
CDVVIGIINNTVYDPLQPELDSFKEELDKYFKNHTSPDVDLG
DISGINASVVNIQKEIDRLNEVAKNLNESLIDLQELGKYEQYI
KWPWYVWLGFIAGLIAIVMVTILLCCMTSCCCLKGACSCGS
CCKFDEDDSEPVLKGVKLHYT Novel_MERS_S2
MIHSVFLLMFLLTPTESDCKLPLGQSLCALPDTPTSTLTPRSVR 33
SVPGEMRLASIAFNHPIQVDQLNSSYFKLSIPTNFSFGVTQEYI
QTTIQKVTVDCKQYVCNGFQKCEQLLREYGGQFCSKINQALH
GANLRQDDSVRNLFASVKSSQSSPIPGFGGDFNLTLLPVSIS
TGSRSARSAIEDLLFDKVTIADPGYMQGYDDCMQQGPASAR
DLICAQYVAGYKVLPLMDVNMEAAAYTSSLLGSIAGVGWTA
GLSSFAAIPFAQSIFYRLNGVGITQQVLSNQLIANKFNQAL
GAMQTGFTTTNEAFQKVQDAVNNNAQALSKLASELSNTFG
AISASIGDIIQRLDVLEQDAQIDRLINGRLTTLNAFVAQQLVRS
ESAALSAQLAKDKVNECVKAQSKRSGFCGQGTHIVSFVVNA
PNGLYFMHVGYYPSNHIEVVSAYGLCDAANPTNCIAPVNGY
FIKTNNTRIVDEWSYTGSSFYAPEPITSLNTKYVAPQVTYQNI
STNLPPPLLGNSTGIDFQDELDEFFKNVSTSIPNFGSLTQINTTL
LDLTYEMLSLQQVVKALNESYIDLKELGNYTYYNKWP Novel_Trimeric_
MFIFLLFLTTLTSGSDLDRALSGIAAEQDRNTREVFAQVKQMY 34 SARS_S2
KTPTLKYFGGFNFSQILPDPLKPTKRSFIEDLLFNKVTLADAG
FMKQYGECLGDINARDLICAQKFNGLTVLPPLLTDDMIAAYT
AALVSGTATAGWTFGAGAALQIPFAMQMAYRFNGIGVTQN
VLYENQKQIANQFNKAISQIQESLTTTSTALGKLQDVVNQNA
QALNTLVKQLSSNFGAISSVLNDILSRLDKVEAEVQIDRLITG
RLQSLQTYVTQQLIRAAEIRASANLAATKMSECVLGQSKRV
DFCGKGYHLMSPQAAPHGVVFLHVTYVPSQERNFTTAPAIC
HEGKAYFPREGVFVFNGTSWFITQRNFFSPQIITDNTFVSGN
CDVVIGIINNTVYDPLQPELDSFKEELDKYFKNHTSPDVDLG
DISGINASVVNIQKEIDRLNEVAKNLNESLIDLQELGKYEQYI
KWPWYVWLGFIAGLIAIVMVTILLCCMTSCCCLKGACSCGS
CCKFDEDDSEPVLKGVKLHYT

(583) TABLE-US-00013 TABLE 12 Full-length Spike Glycoprotein Amino Acid Sequences
(*Homo sapiens* strains) GenBank Collection Accession Country Date Release Date Virus Name
AFY13307 United Sep. 11, 2012 Dec. 5, 2012 Betacoronavirus England 1, Kingdom complete
genome AFS88936 Jun. 13, 2012 Sep. 27, 2012 Human betacoronavirus 2c EMC/2012, complete
genome AGG22542 United Sep. 19, 2012 Feb. 27, 2013 Human betacoronavirus 2c England-

Kingdom Qatar/2012, complete genome AHY21469 Jordan 2012 May 4, 2014 Human betacoronavirus 2c Jordan- N3/2012 isolate MG167, complete genome AGH58717 Jordan April 2012 Mar. 25, 2013 Human betacoronavirus 2c Jordan- N3/2012, complete genome AGV08444 Saudi May 7, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_12_2013, complete genome AGV08546 Saudi May 11, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_15_2013, complete genome AGV08535 Saudi May 12, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_16_2013, complete genome AGV08558 Saudi May 15, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_17_2013, complete genome AGV08573 Saudi May 23, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_18_2013, complete genome AGV08480 Saudi May 23, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_19_2013, complete genome AGN70962 Saudi May 9, 2013 Jun. 10, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_1_2013, complete genome AGV08492 Saudi May 30, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_21_2013, complete genome AHI48517 Saudi May 2, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_25_2013, complete genome AGN70951 Saudi Apr. 21, 2013 Jun. 10, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_2_2013, complete genome AGN70973 Saudi Apr. 22, 2013 Jun. 10, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_3_2013, complete genome AGN70929 Saudi May 1, 2013 Jun. 10, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Al-Hasa_4_2013, complete genome AGV08408 Saudi Jun. 19, 2012 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Bisha_1_2012, complete genome AGV08467 Saudi May 13, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Buraidah_1_2013, complete genome AID50418 United Feb. 10, 2013 Jun. 18, 2014 Middle East respiratory syndrome Kingdom coronavirus isolate England/2/2013, complete genome AJD81451 United Feb. 10, 2013 Jan. 18, 2015 Middle East respiratory syndrome Kingdom coronavirus isolate England/3/2013, complete genome AJD81440 United Feb. 13, 2013 Jan. 18, 2015 Middle East respiratory syndrome Kingdom coronavirus isolate England/4/2013, complete genome AHB33326 France May 7, 2013 Dec. 7, 2013 Middle East respiratory syndrome coronavirus isolate FRA/UAE, complete genome AIZ48760 USA June 2014 Dec. 14, 2014 Middle East respiratory syndrome coronavirus isolate Florida/USA- 2_Saudi Arabia_2014, complete genome AGV08455 Saudi Jun. 4, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Hafr-Al- Batin_1_2013, complete genome AHI48561 Saudi Aug. 5, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Hafr-Al- Batin_2_2013, complete genome AHI48539 Saudi Aug. 28, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Hafr-Al- Batin_6_2013, complete genome AIZ74417 France Apr. 26, 2013 Mar. 10, 2015 Middle East respiratory syndrome coronavirus isolate Hu-France (UAE) - FRA1_1627- 2013_BAL_Sanger, complete genome AIZ74433 France May 7, 2013 Mar. 10, 2015 Middle East respiratory syndrome coronavirus isolate Hu-France - FRA2_130569-2013_IS-HTS, complete genome AIZ74439 France May 7, 2013 Mar. 10, 2015 Middle East respiratory syndrome coronavirus isolate Hu-France - FRA2_130569-2013_InSpu_Sanger, complete genome AIZ74450 France May 7, 2013 Mar. 10, 2015 Middle East respiratory syndrome coronavirus isolate Hu-France - FRA2_130569-2013_Isolate_Sanger, complete genome AKK52602 Saudi Feb. 10, 2015 Jun. 8, 2015 Middle East respiratory syndrome Arabia coronavirus isolate Hu/Riyadh_KSA_2959_2015, complete genome AKK52612 Saudi Mar. 1, 2015 Jun. 8, 2015 Middle East respiratory syndrome Arabia coronavirus isolate Hu/Riyadh_KSA_4050_2015, complete genome AHN10812 Saudi Nov. 6, 2013 Mar. 24, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_1_2013, complete genome AID55071 Saudi Apr. 21, 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_C10306/KSA/2014-04-20, complete genome AID55066 Saudi 2014

Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_C7149/KSA/2014-04-05, complete genome AID55067 Saudi 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_C7569/KSA/2014-04-03, complete genome AID55068 Saudi Apr. 7, 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_C7770/KSA/2014-04-07, complete genome AID55069 Saudi Apr. 12, 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_C8826/KSA/2014-04-12, complete genome AID55070 Saudi Apr. 14, 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Jeddah_C9055/KSA/2014-04-14, complete genome AHE78108 Saudi Nov. 5, 2013 May 1, 2014 Middle East respiratory syndrome Arabia coronavirus isolate MERS-CoV- Jeddah-human-1, complete genome AKL59401 South May 20, 2015 Jun. 9, 2015 Middle East respiratory syndrome Korea coronavirus isolate MERS-CoV/KOR/KNIH/002_05_2015, complete genome ALD51904 Thailand Jun. 17, 2015 Jul. 7, 2015 Middle East respiratory syndrome coronavirus isolate MERS- CoV/THA/CU/17_06_2015, complete genome AID55072 Saudi Apr. 15, 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Makkah_C9355/KSA/Makkah/2014- 04-15, complete genome AHC74088 Qatar Oct. 13, 2013 Dec. 23, 2013 Middle East respiratory syndrome coronavirus isolate Qatar3, complete genome AHC74098 Qatar Oct. 17, 2013 Dec. 23, 2013 Middle East respiratory syndrome coronavirus isolate Qatar4, complete genome AHI48572 Saudi Aug. 15, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_14_2013, complete genome AGV08379 Saudi Oct. 23, 2012 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_1_2012, complete genome AID55073 Saudi Apr. 22, 2014 Nov. 12, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_2014KSA_683/KSA/2014, complete genome AGV08584 Saudi Oct. 30, 2012 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_2_2012, complete genome AGV08390 Saudi Feb. 5, 2013 Sep. 17, 2013 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_3_2013, complete genome AHI48605 Saudi Mar. 1, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_4_2013, complete genome AHI48583 Saudi Jul. 2, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_5_2013, complete genome AHI48528 Saudi Jul. 17, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Riyadh_9_2013, complete genome AHI48594 Saudi Jun. 12, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Taif_1_2013, complete genome AHI48550 Saudi Jun. 12, 2013 Feb. 6, 2014 Middle East respiratory syndrome Arabia coronavirus isolate Wadi-Ad- Dawasir_1_2013, complete genome AIY60558 United Mar. 7, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi/Gayathi_UAE_2_2014, complete genome AIY60538 United Apr. 10, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_16_2014, complete genome AIY60528 United Apr. 10, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_18_2014, complete genome AIY60588 United Apr. 13, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_26_2014, complete genome AIY60548 United Apr. 19, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_30_2014, complete genome AIY60568 United Apr. 17, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_33_2014, complete genome AIY60518 United Apr. 7, 2014 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_8_2014, complete genome AIY60578 United Nov. 15, 2013 Dec. 6, 2014 Middle East respiratory syndrome Arab coronavirus strain Abu Emirates Dhabi_UAE_9_2013, complete genome AKJ80137 China May 27, 2015 Jun. 5, 2015 Middle East respiratory syndrome coronavirus strain ChinaGD01, complete genome AHZ64057 USA May 10, 2014 May 14, 2014 Middle East respiratory syndrome coronavirus strain Florida/USA- 2_Saudi Arabia_2014, complete genome AKM76229 Oman Oct. 28, 2013 Jun. 23, 2015 Middle East

respiratory syndrome coronavirus strain Hu/Oman_2285_2013, complete genome AKM76239
Oman Dec. 28, 2013 Jun. 23, 2015 Middle East respiratory syndrome coronavirus strain
Hu/Oman_2874_2013, complete genome AKI29284 Saudi Jan. 6, 2015 May 27, 2015 Middle East
respiratory syndrome Arabia coronavirus strain Hu/Riyadh-KSA- 2049/2015, complete genome
AKI29265 Saudi Jan. 21, 2015 May 27, 2015 Middle East respiratory syndrome Arabia
coronavirus strain Hu/Riyadh-KSA- 2343/2015, complete genome AKI29255 Saudi Jan. 21, 2015
May 27, 2015 Middle East respiratory syndrome Arabia coronavirus strain Hu/Riyadh-KSA-
2345/2015, complete genome AKI29275 Saudi Jan. 26, 2015 May 27, 2015 Middle East
respiratory syndrome Arabia coronavirus strain Hu/Riyadh-KSA- 2466/2015, complete genome
AKK52582 Saudi Feb. 10, 2015 Jun. 8, 2015 Middle East respiratory syndrome Arabia coronavirus
strain Hu/Riyadh_KSA_2959_2015, complete genome AKK52592 Saudi Mar. 1, 2015 Jun. 8, 2015
Middle East respiratory syndrome Arabia coronavirus strain Hu/Riyadh_KSA_4050_2015,
complete genome AHZ58501 USA Apr. 30, 2014 May 13, 2014 Middle East respiratory syndrome
coronavirus strain Indiana/USA- 1_Saudi Arabia_2014, complete genome AGN52936 United 2013
Jun. 10, 2013 Middle East respiratory syndrome Arab coronavirus, complete genome Emirates
(584) TABLE-US-00014 TABLE 13 SEQ ID Description Sequence NO: MeV Nucleic
Acid Sequences GC_F_MEASLES_

TCAAGCTTTTGGACCCTCGTACAGAAGCTAATACGACT 35 B3.1
CACTATAGGGAAATAAGAGAGAGAAAAGAAGAGTAAGAA Sequence, NT (5'
GAAATATAAGAGCCACCATGGGTCTCAAGGTGAACGTC UTR, ORF, 3'
TCTGCCGTATTCATGGCAGTACTGTAACTCTCCAAACA UTR)
CCCGCCGGTCAAATTCATTGGGGCAATCTCTCTAAGAT Sequence Length:
AGGGGTAGTAGGAATAGGAAGTGCAAGCTACAAAGTT 1864
ATGACTCGTTCCAGCCATCAATCATTAGTCATAAAATT
AATGCCCAATATAACTCTCCTCAATAACTGCACGAGGG
TAGAGATTGCAGAATACAGGAGACTACTAAGAACAGTT
TTGGAACCAATTAGGGATGCACTTAATGCAATGACCCA
GAACATAAGGCCGGTTCAGAGCGTAGCTTCAAGTAGGA
GACACAAGAGATTTGCGGGAGTAGTCCTGGCAGGTGCG
GCCCTAGGTGTTGCCACAGCTGCTCAGATAACAGCCGG
CATTGCACTTCACCGGTCCATGCTGAACTCTCAGGCCAT
CGACAATCTGAGAGCGAGCCTGGAAACTACTAATCAGG
CAATTGAGGCAATCAGACAAGCAGGGCAGGAGATGAT
ATTGGCTGTTCAAGGTGTCCAAGACTACATCAATAATG
AGCTGATACCGTCTATGAACCAGCTATCTTGTGATCTA
ATCGGTCAGAAGCTCGGGCTCAAATTGCTTAGATACTA
TACAGAAATCCTGTCATTATTTGGCCCCAGCCTACGGG
ACCCCATATCTGCGGAGATATCTATCCAGGCTTTGAGTT
ATGCACTTGGAGGAGATATCAATAAGGTGTTAGAAAAG
CTCGGATACAGTGGAGGCGATTTACTAGGCATCTTAGA
GAGCAGAGGAATAAAGGCTCGGATAACTCACGTTCGAC
ACAGAGTCCTACTTCATAGTCCTCAGTATAGCCTATCCG
ACGCTGTCCGAGATTAAGGGGGTGATTGTCCACCGGCT
AGAGGGGGTCTCGTACAACATAGGCTCTCAAGAGTGGT
ATACCACTGTGCCCAAGTATGTTGCAACCCAAGGGTAC
CTTATCTCGAATTTTGATGAGTCATCATGTACTTTTCATG
CCAGAGGGGACTGTGTGCAGCCAAAATGCCTTGTACCC
GATGAGTCCTCTGCTCCAAGAATGCCTCCGGGGGTCCA
CCAAGTCCTGTGCTCGTACACTCGTATCCGGGTCTTTTG
GGAACCGGTTCATTTTATCACAAGGGAACCTAATAGCC

ATTGTGCTGAATTCATTGTAAGTGTACACAACAGGT
ACGATTATTAATCAAGACCCTGACAAGATCCTAACATA
CATTGCTGCCGATCGCTGCCCCGGTAGTCGAGGTGAACG
GCGTGACCATCCAAGTCGGGAGCAGGAGGTATCCAGA
CGCTGTGTACTTGCACAGAATTGACCTCGGTCCTCCCAT
ATCATTGGAGAGGTTGGACGTAGGGACAAATCTGGGG
AATGCAATTGCCAAATTGGAGGATGCCAAGGAATTGTT
GGAATCATCGGACCAGATATTGAGAAGTATGAAAGGTT
TATCGAGCACTAGCATAGTCTACATCCTGATTGCAGTG
TGTCTTGGAGGGTTGATAGGGATCCCCACTTTAATATGT
TGCTGCAGGGGGCGTTGTAACAAAAAGGGAGAACAAG
TTGGTATGTCAAGACCAGGCCTAAAGCCTGACCTTACA
GGAACATCAAAATCCTATGTAAGATCGCTTTGATGATA
ATAGGCTGGAGCCTCGGTGGCCAAGCTTCTTGCCCCCTT
GGGCCTCCCCCAGCCCCCTCCTCCCCTTCCTGCACCCGT
ACCCCCGTGGTCTTTGAATAAAGTCTGAGTGGGCGGC GC_F_MEASLES_
ATGGGTCTCAAGGTGAACGTCTCTGCCGTATTCATGGC 36 B3.1
AGTACTGTAACTCTCCAAACACCCGCCGGTCAAATTC ORF Sequence, NT
ATTGGGGCAATCTCTCTAAGATAGGGGTAGTAGGAATA
GGAAGTGCAAGCTACAAAGTTATGACTCGTTCCAGCCA
TCAATCATTAGTCATAAAATTAATGCCCAATATAACTCT
CCTCAATAACTGCACGAGGGTAGAGATTGCAGAATACA
GGAGACTACTAAGAACAGTTTTTGGAAACCAATTAGGGAT
GCACTTAATGCAATGACCCAGAACATAAGGCCGGTTCA
GAGCGTAGCTTCAAGTAGGAGACACAAGAGATTGCG
GGAGTAGTCCTGGCAGGTGCGGCCCTAGGTGTTGCCAC
AGCTGCTCAGATAACAGCCGGCATTGCACTTCACCGGT
CCATGCTGAACTCTCAGGCCATCGACAATCTGAGAGCG
AGCCTGGAACTACTAATCAGGCAATTGAGGCAATCAG
ACAAGCAGGGCAGGAGATGATATTGGCTGTTCAGGGTG
TCCAAGACTACATCAATAATGAGCTGATACCGTCTATG
AACCAGCTATCTTGTGATCTAATCGGTCAGAAGCTCGG
GCTCAAATTGCTTAGATACTATACAGAAATCCTGTCATT
ATTTGGCCCCAGCCTACGGGACCCCATATCTGCGGAGA
TATCTATCCAGGCTTTGAGTTATGCACTTGGAGGAGAT
ATCAATAAGGTGTTAGAAAAGCTCGGATACAGTGGAG
GCGATTTACTAGGCATCTTAGAGAGCAGAGGAATAAAG
GCTCGGATAACTCACGTCGACACAGAGTCCTACTTCAT
AGTCCTCAGTATAGCCTATCCGACGCTGTCCGAGATTA
AGGGGGTGATTGTCCACCGGCTAGAGGGGGTCTCGTAC
AACATAGGCTCTCAAGAGTGGTATAACCACTGTGCCCAA
GTATGTTGCAACCCAAGGGTACCTTATCTCGAATTTTGA
TGAGTCATCATGTACTTTCATGCCAGAGGGGACTGTGT
GCAGCCAAAATGCCTTGTACCCGATGAGTCCTCTGCTC
CAAGAATGCCTCCGGGGGGTCCACCAAGTCCTGTGCTCG
TACACTCGTATCCGGGTCTTTTGGGAACCGGTTCATTTT
ATCACAAGGGAACCTAATAGCCAATTGTGCATCAATTC
TTTGTAAGTGTTACACAACAGGTACGATTATTAATCAA
GACCCTGACAAGATCCTAACATACATTGCTGCCGATCG
CTGCCCCGGTAGTCGAGGTGAACGGCGTGACCATCCAAG

TCGGGACGACGATCCAGACGCTGTGTACTTGCCAC
AGAATTGACCTCGGTCCTCCCATATCATTGGAGAGGTT
GGACGTAGGGACAAATCTGGGGAATGCAATTGCCAAA
TTGGAGGATGCCAAGGAATTGTTGGAATCATCGGACCA
GATATTGAGAAGTATGAAAGGTTTATCGAGCACTAGCA
TAGTCTACATCCTGATTGCAGTGTGTCTTGGAGGGTTGA
TAGGGATCCCCACTTTAATATGTTGCTGCAGGGGGCGT
TGTAACAAAAAGGGAGAACAAGTTGGTATGTCAAGAC
CAGGCCTAAAGCCTGACCTTACAGGAACATCAAAATCC TATGTAAGATCGCTTTGA
GC_F_MEASLES_ G*GGGAAATAAGAGAGAAAAGAAGAGTAAGAAGAAAT 37 B3.1
ATAAGAGCCACCATGGGTCTCAAGGTGAACGTCTCTGC mRNA Sequence
CGTATTCATGGCAGTACTGTAACTCTCCAAACACCCG (assumes T100 tail)
CCGGTCAAATTCATTGGGGCAATCTCTCTAAGATAGGG mRNA Sequence
GTAGTAGGAATAGGAAGTGCAAGCTACAAAGTTATGA Length: 1925
CTCGTTCCAGCCATCAATCATTAGTCATAAAATTAATGC
CCAATATACTCTCCTCAATAACTGCACGAGGGTAGAG
ATTGCAGAATACAGGAGACTACTAAGAACAGTTTTTGA
ACCAATTAGGGATGCACTTAATGCAATGACCCAGAACA
TAAGGCCGGTTCAGAGCGTAGCTTCAAGTAGGAGACAC
AAGAGATTTGCGGGAGTAGTCCTGGCAGGTGCGGCCCT
AGGTGTTGCCACAGCTGCTCAGATAACAGCCGGCATTG
CACTTCACCGGTCCATGCTGAACTCTCAGGCCATCGAC
AATCTGAGAGCGAGCCTGGAACTACTAATCAGGCAAT
TGAGGCAATCAGACAAGCAGGGCAGGAGATGATATTG
GCTGTTCAAGGTGTCCAAGACTACATCAATAATGAGCT
GATACCGTCTATGAACCAGCTATCTTGTGATCTAATCG
GTCAGAAGCTCGGGCTCAAATTGCTTAGATACTATACA
GAAATCCTGTCATTATTTGGCCCCAGCCTACGGGACCC
CATATCTGCGGAGATATCTATCCAGGCTTTGAGTTATGC
ACTTGGAGGAGATATCAATAAGGTGTTAGAAAAGCTCG
GATACAGTGGAGGCGATTTACTAGGCATCTTAGAGAGC
AGAGGAATAAAGGCTCGGATAACTCACGTCGACACAG
AGTCCTACTTCATAGTCCTCAGTATAGCCTATCCGACGC
TGTCCGAGATTAAGGGGGTGATTGTCCACCGGCTAGAG
GGGGTCTCGTACAACATAGGCTCTCAAGAGTGGTATAC
CACTGTGCCCAAGTATGTTGCAACCCAAGGGTACCTTA
TCTCGAATTTTGATGAGTCATCATGTACTTTCATGCCAG
AGGGGACTGTGTGCAGCCAAAATGCCTTGTACCCGATG
AGTCCTCTGCTCCAAGAATGCCTCCGGGGGTCCACCAA
GTCCTGTGCTCGTACACTCGTATCCGGGTCTTTTGGGAA
CCGGTTCATTTTATCACAAGGGAACCTAATAGCCAATT
GTGCATCAATTCTTTGTAAGTGTTACACAACAGGTACG
ATTATTAATCAAGACCCTGACAAGATCCTAACATACAT
TGCTGCCGATCGCTGCCCCGTAGTCGAGGTGAACGGCG
TGACCATCCAAGTCGGGAGCAGGAGGTATCCAGACGCT
GTGTACTTGCACAGAATTGACCTCGGTCTCCCATATCA
TTGGAGAGGTTGGACGTAGGGACAAATCTGGGGAATG
CAATTGCCAAATTGGAGGATGCCAAGGAATTGTTGGAA
TCATCGGACCAGATATTGAGAAGTATGAAAGGTTTATC
GAGCACTAGCATAGTCTACATCCTGATTGCAGTGTGTC

TTGGAGTTGCTGAGTCCCTGACCTTTAATATGTTGCT
GCAGGGGGCGTTGTAACAAAAAGGGGAGAACAAGTTGG
TATGTCAAGACCAGGCCTAAAGCCTGACCTTACAGGAA
CATCAAAATCCTATGTAAGATCGCTTTGATGATAATAG
GCTGGAGCCTCGGTGGCCAAGCTTCTTGCCCCTTGGGC
CTCCCCCAGCCCCCTCCTCCCCCTTCTGACCCGTACCC
CCGTGGTCTTTGAATAAAGTCTGAGTGGGCGGCAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAATCTAG GC_F_MEASLES_
TCAAGCTTTTGGACCCTCGTACAGAAGCTAATACGACT 38 D8
CACTATAGGGAAATAAGAGAGAGAAAAGAAGAGTAAGAA Sequence, NT (5'
GAAATATAAGAGCCACCATGGGTCTCAAGGTGAACGTC UTR, ORF, 3'
TCTGTCATATTCATGGCAGTACTGTAACTCTTCAAACA UTR)
CCCACCGGTCAAATCCATTGGGGCAATCTCTCTAAGAT Sequence Length:
AGGGGTGGTAGGGGTAGGAAGTGCAAGCTACAAAGTT 1864
ATGACTCGTTCCAGCCATCAATCATTAGTCATAAAGTT
AATGCCCAATATAACTCTCCTCAACAATTGCACGAGGG
TAGGGATTGCAGAATACAGGAGACTACTGAGAACAGTT
CTGGAACCAATTAGAGATGCACTTAATGCAATGACCCA
GAATATAAGACCGGTTTCAGAGTGTAGCTTCAAGTAGGA
GACACAAGAGATTTGCGGGAGTTGTCCTGGCAGGTGCG
GCCCTAGGCGTTGCCACAGCTGCTCAAATAACAGCCGG
TATTGCACTTCACCAGTCCATGCTGAACTCTCAAGCCAT
CGACAATCTGAGAGCGAGCCTAGAACTACTAATCAGG
CAATTGAGGCAATCAGACAAGCAGGGCAGGAGATGAT
ATTGGCTGTTCAGGGTGTCCAAGACTACATCAATAATG
AGCTGATACCGTCTATGAATCAACTATCTTGTGATTTAA
TCGGCCAGAAGCTAGGGCTCAAATTGCTCAGATACTAT
ACAGAAATCCTGTCAATTATTTGGCCCCAGCTTACGGGA
CCCCATATCTGCGGAGATATCTATCCAGGCTTTGAGCT
ATGCGCTTGGAGGAGATATCAATAAGGTGTTGGAAAAG
CTCGGATACAGTGGAGGTGATCTACTGGGCATCTTAGA
GAGCAGAGGAATAAAGGCCCGGATAACTCACGTCGAC
ACAGAGTCCTACTTCATTGTACTCAGTATAGCCTATCCG
ACGCTATCCGAGATTAAGGGGGTGATTGTCCACCGGCT
AGAGGGGGTCTCGTACAACATAGGCTCTCAAGAGTGGT
ATACCACTGTGCCCAAGTATGTTGCAACCCAAGGGTAC
CTTATCTCGAATTTTGATGAGTCATCATGCACTTTCATG
CCAGAGGGGACTGTGTGCAGCCAGAATGCCTTGTACCC
GATGAGTCCTCTGCTCCAAGAATGCCTCCGGGGGTCCA
CTAAGTCCTGTGCTCGTACACTCGTATCCGGGTCTTTTCG
GGAACCGGTTCATTTTATCACAGGGGAACCTAATAGCC
AATTGTGCATCAATCCTTTGCAAGTGTTACACAACAGG
AACAATCATTAATCAAGACCCTGACAAGATCCTAACAT
ACATTGCTGCCGATCACTGCCCGGTGGTCGAGGTGAAT
GGCGTGACCATCCAAGTCGGGAGCAGGAGGTATCCGG
ACGCTGTGTACTTGCACAGGATTGACCTCGGTCCTCCC
ATATCTTTGGAGAGGTTGGACGTAGGGACAAATCTGGG
GAATGCAATTGCTAAGTTGGAGGATGCCAAGGAATTGT

TGGAGTGCATGATATTGAGGAGGTATGAAAGGT
TTATCGAGCACTAGTATAGTTTACATCCTGATTGCAGTG
TGTCTTGGAGGATTGATAGGGATCCCCGCTTTAATATGT
TGCTGCAGGGGGCGTTGTAACAAGAAGGGAGAACAAG
TTGGTATGTCAAGACCAGGCCTAAAGCCTGATCTTACA
GGAACATCAAAATCCTATGTAAGGTCACCTCTGATGATA
ATAGGCTGGAGCCTCGGTGGCCAAGCTTCTTGCCCCCTT
GGGCCTCCCCCAGCCCCCTCCTCCCCTTCCTGCACCCGT
ACCCCCGTGGTCTTTGAATAAAGTCTGAGTGGGCGGC GC_F_MEASLES_
ATGGGTCTCAAGGTGAACGTCTCTGTCATATTCATGGC 39 D8
AGTACTGTAACTCTTCAAACACCCACCGGTCAAATCC ORF Sequence, NT
ATTGGGGCAATCTCTCTAAGATAGGGGTGGTAGGGGTA
GGAAGTGCAAGCTACAAAGTTATGACTCGTTCCAGCCA
TCAATCATTAGTCATAAAGTTAATGCCCAATATAACTCT
CCTCAACAATTGCACGAGGGTAGGGATTGCAGAATACA
GGAGACTACTGAGAACAGTTCTGGAACCAATTAGAGAT
GCACTTAATGCAATGACCCAGAATATAAGACCGGTTCA
GAGTGTAGCTTCAAGTAGGAGACACAAGAGATTTGCGG
GAGTTGTCCTGGCAGGTGCGGCCCTAGGCGTTGCCACA
GCTGCTCAAATAACAGCCGGTATTGCACTTCACCAGTC
CATGCTGAACTCTCAAGCCATCGACAATCTGAGAGCGA
GCCTAGAACTACTAATCAGGCAATTGAGGCAATCAGA
CAAGCAGGGCAGGAGATGATATTGGCTGTTCAAGGTGT
CCAAGACTACATCAATAATGAGCTGATACCGTCTATGA
ATCAACTATCTTGTGATTTAATCGGCCAGAAGCTAGGG
CTCAAATTGCTCAGATACTATACAGAAATCCTGTCATT
ATTTGGCCCCAGCTTACGGGACCCCATATCTGCGGAGA
TATCTATCCAGGCTTTGAGCTATGCGCTTGGAGGAGAT
ATCAATAAGGTGTTGGAAAAGCTCGGATACAGTGGAG
GTGATCTACTGGGCATCTTAGAGAGCAGAGGAATAAAG
GCCCCGATAACTCACGTCGACACAGAGTCCTACTTCAT
TGTACTIONAGCCTATCCGACGCTATCCGAGATTA
AGGGGGTGATTGTCCACCGGCTAGAGGGGGTCTCGTAC
AACATAGGCTCTCAAGAGTGGTATAACCACTGTGCCCAA
GTATGTTGCAACCCAAGGGTACCTTATCTCGAATTTTGA
TGAGTCATCATGCACTTTCATGCCAGAGGGGACTGTGT
GCAGCCAGAATGCCTTGTACCCGATGAGTCCTCTGCTC
CAAGAATGCCTCCGGGGGTCCACTAAGTCCTGTGCTCG
TACACTCGTATCCGGGTCTTTCGGGAACCGGTTCATTTT
ATCACAGGGGAACCTAATAGCCAATTGTGCATCAATCC
TTTGCAAGTGTTACACAACAGGAACAATCATTAAATCAA
GACCCTGACAAGATCCTAACATAACATTGCTGCCGATCA
CTGCCCGGTGGTCGAGGTGAATGGCGTGACCATCCAAG
TCGGGAGCAGGAGGTATCCGGACGCTGTGTACTTGCAC
AGGATTGACCTCGGTCCTCCCATATCTTTGGAGAGGTT
GGACGTAGGGACAAATCTGGGGAATGCAATTGCTAAGT
TGGAGGATGCCAAGGAATTGTTGGAGTCATCGGACCAG
ATATTGAGGAGTATGAAAGGTTTATCGAGCACTAGTAT
AGTTTACATCCTGATTGCAGTGTGTCTTGGAGGATTGAT
AGGGATCCCCGCTTTAATATGTTGCTGCAGGGGGCGTT

GTAACAAGGAGCAAGATAAGTTGGTATGTCAAGACC
AGGCCTAAAGCCTGATCTTACAGGAACATCAAAATCCT ATGTAAGGTCACCTCTGA
GC_F_MEASLES_ G*GGGAAATAAGAGAGAAAAGAAGAGTAAGAAGAAAT 40 D8
ATAAGAGCCACCATGGGTCTCAAGGTGAACGTCTCTGT mRNA Sequence
CATATTCATGGCAGTACTGTAACTCTTCAAACACCCAC (assumes T100 tail)
CGGTCAAATCCATTGGGGCAATCTCTCTAAGATAGGGG Sequence Length:
TGGTAGGGGTAGGAAGTGCAAGCTACAAAGTTATGACT 1925
CGTTCCAGCCATCAATCATTAGTCATAAAGTTAATGCC
CAATATAACTCTCCTCAACAATTGCACGAGGGTAGGGA
TTGCAGAATACAGGAGACTACTGAGAACAGTTCTGGAA
CCAATTAGAGATGCACTTAATGCAATGACCCAGAATAT
AAGACCGGTTTCAAGAGTGTAGCTTCAAGTAGGAGACACA
AGAGATTTGCGGGAGTTGTCTTGGCAGGTGCGGCCCTA
GGCGTTGCCACAGCTGCTCAAATAACAGCCGGTATTGC
ACTTCACCAGTCCATGCTGAACTCTCAAGCCATCGACA
ATCTGAGAGCGAGCCTAGAACTACTAATCAGGCAATT
GAGGCAATCAGACAAGCAGGGCAGGAGATGATATTGG
CTGTTCAAGGTGTCCAAGACTACATCAATAATGAGCTG
ATACCGTCTATGAATCAACTATCTTGTGATTTAATCGGC
CAGAAGCTAGGGCTCAAATTGCTCAGATACTATACAGA
AATCCTGTCATTATTTGGCCCCAGCTTACGGGACCCCAT
ATCTGCGGAGATATCTATCCAGGCTTTGAGCTATGCGC
TTGGAGGAGATATCAATAAGGTGTTGGAAAAGCTCGGA
TACAGTGGAGGTGATCTACTGGGCATCTTAGAGAGCAG
AGGAATAAAGGCCCGGATAACTCACGTCGACACAGAG
TCCTACTTCATTGTACTCAGTATAGCCTATCCGACGCTA
TCCGAGATTAAGGGGGTGATTGTCCACCGGCTAGAGGG
GGTCTCGTACAACATAGGCTCTCAAGAGTGGTATACCA
CTGTGCCCAAGTATGTTGCAACCCAAGGGTACCTTATC
TCGAATTTTGATGAGTCATCATGCACTTTCATGCCAGAG
GGGACTGTGTGCAGCCAGAATGCCTTGTACCCGATGAG
TCCTCTGCTCCAAGAATGCCTCCGGGGGTCCACTAAGT
CCTGTGCTCGTACACTCGTATCCGGGTCTTTCGGGAACC
GGTTCATTTTATCACAGGGGAACCTAATAGCCAATTGT
GCATCAATCCTTTGCAAGTGTTACACAACAGGAACAAT
CATTAAATCAAGACCCTGACAAGATCCTAACATACATTG
CTGCCGATCACTGCCCCGGTGGTCGAGGTGAATGGCGTG
ACCATCCAAGTCGGGAGCAGGAGGTATCCGGACGCTGT
GTACTTGACAGGATTGACCTCGGTCTCCCATATCTTT
GGAGAGGTTGGACGTAGGGACAAATCTGGGGAATGCA
ATTGCTAAGTTGGAGGATGCCAAGGAATTGTTGGAGTC
ATCGGACCAGATATTGAGGAGTATGAAAGGTTTATCGA
GCACTAGTATAGTTTACATCCTGATTGCAGTGTGTCTTG
GAGGATTGATAGGGATCCCCGCTTTAATATGTTGCTGC
AGGGGGCGTTGTAACAAGAAGGGAGAACAAGTTGGTA
TGTCAGAGACCAGGCCTAAAGCCTGATCTTACAGGAACA
TCAAAATCCTATGTAAGGTCACCTCTGATGATAATAGGC
TGGAGCCTCGGTGGCCAAGCTTCTTGCCCCTTGGGCCTC
CCCCCAGCCCCTCCTCCCCTTCCCTGCACCCGTACCCCG
TGGTCTTTGAATAAAGTCTGAGTGGGCGGCCAAAAAAA

AA
AA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATCTAG GC_H_MEASLES_
TCAAGCTTTTGGACCCTCGTACAGAAGCTAATACGACT 41 B3
CACTATAGGGAAATAAGAGAGAGAAAAGAAGAGTAAGAA Sequence, NT (5'
GAAATATAAGAGCCACCATGTCACCGCAACGAGACCG UTR, ORF, 3'
GATAAATGCCTTCTACAAAGATAACCCTTATCCCAAGG UTR)
GAAGTAGGATAGTTATTAACAGAGAACATCTTATGATT Sequence Length:
GACAGACCCTATGTTCTGCTGGCTGTTCTGTTTCGTCATG 2065
TTTCTGAGCTTGATCGGATTGCTGGCAATTGCAGGCATT
AGACTTCATCGGGCAGCCATCTACACCGCGGAGATCCA
TAAAAGCCTCAGTACCAATCTGGATGTGACTAACTCCA
TCGAGCATCAGGTCAAGGACGTGCTGACACCACTCTTT
AAAATCATCGGGGATGAAGTGGGCCTGAGAACACCTC
AGAGATTCACTGACCTAGTGAAATTCATCTCGGACAAG
ATTAAATTCCTTAATCCGGATAGGGAGTACGACTTCAG
AGATCTCACTTGGTGCATCAACCCGCCAGAGAGGATCA
AACTAGATTATGATCAATACTGTGCAGATGTGGCTGCT
GAAGAGCTCATGAATGCATTGGTGAAC TCAACTCTACT
GGAGACCAGAACAACTCAGTTCCTAGCTGTCTCAA
AGGGAAACTGCTCAGGGCCCACTACAATCAGAGGTCA
ATTCTCAAACATGTCGCTGTCCTTGTTGGACTTGTACTT
AGGTCGAGGTTACAATGTGTCATCTATAGTCACTATGA
CATCCCAGGGAATGTATGGGGGAACCTACCTAGTTGAA
AAGCCTAATCTGAACAGCAAAGGGTCAGAGTTGTCACA
ACTGAGCATGTACCGAGTGTTTGAAGTAGGTGTGATCA
GAAACCCGGGTTTGGGGGGCTCCGGTGTTCCATATGACA
AACTATTTTGAGCAACCAGTCAGTAATGGTCTCGGCAA
CTGTATGGTGGCTTTGGGGGAGCTCAAAC TCGCAGCCC
TTTGTCACGGGGACGATTCTATCATAATTCCCTATCAGG
GATCAGGGAAAGGTGTCAGCTTCCAGCTCGTCAAGCTG
GGTGTCTGGAATCCCCAACCGACATGCAATCCTGGGT
CCCCTTATCAACGGATGATCCAGTGGTAGACAGGCTTT
ACCTCTCATCTCACAGAGGTGTCATCGCTGACAATCAA
GCAAATGGGCTGTCCCGACAACACGAACAGATGACA
AGTTGCGAATGGAGACATGCTTCCAGCAGGCGTGTA
GGTAAAATCCAAGCACTCTGCGAGAATCCCGAGTGGGT
ACCATTGAAGGATAACAGGATTCCTTCATACGGGGGTCC
TGTCTGTTGATCTGAGTCTGACGGTTGAGCTTAAAATCA
AAATTGCTTCGGGATTCGGGCCATTGATCACACACGGC
TCAGGGATGGACCTATACAAATCCAAC TGAACAATGT
GTATTGGCTGACTATTCCGCCAATGAGAAATCTAGCCT
TAGGCGTAATCAACACATTGGAGTGGATACCGAGATTC
AAGGTTAGTCCCAACCTCTTCACTGTCCCAATTAAGGA
AGCAGGCGAAGACTGCCATGCCCCAACATACCTACCTG
CGGAGGTGGACGGTGATGTCAAAC TCAAGTTCCAACCTG
GTGATTCTACCTGGTCAAGATCTCCAATATGTTTTGGCA
ACCTACGATACCTCCAGGGTTGAGCATGCTGTGGTTTA
TTACGTTTACAGCCCAAGCCGCTCATTTTCTTACTTTTA
TCCTTTTAGGTTGCCTATAAAGGGGGTCCCAATCGAAC

TACAAGTGGATGACATGGGATCAAAAACCTCTGG
TGCCGTCACCTTCTGTGTGCTTGCGGACTCAGAATCCGGT
GGACTTATCACTCACTCTGGGATGGTGGGCATGGGAGT
CAGCTGCACAGCTACCCGGGAAGATGGAACCAATCGC
AGATAATGATAATAGGCTGGAGCCTCGGTGGCCAAGCT
TCTTGCCCCCTTGGGCCTCCCCCCAGCCCCCTCCTCCCCTT
CCTGCACCCGTACCCCGTGGTCTTTGAATAAAGTCTG AGTGGGCGGC
GC_H_MEASLES_ ATGTCACCGCAACGAGACCGGATAAATGCCTTCTACAA 42 B3
AGATAACCTTATCCCAAGGGAAGTAGGATAGTTATTA ORF Sequence, NT
ACAGAGAACATCTTATGATTGACAGACCCTATGTTCTG
CTGGCTGTTCTGTTCGTCATGTTTCTGAGCTTGATCGGA
TTGCTGGCAATTGCAGGCATTAGACTTCATCGGGCAGC
CATCTACACCGCGGAGATCCATAAAAGCCTCAGTACCA
ATCTGGATGTGACTAACTCCATCGAGCATCAGGTCAAG
GACGTGCTGACACCACTCTTTAAAATCATCGGGGATGA
AGTGGGCCTGAGAACACCTCAGAGATTCACTGACCTAG
TGAAATTCATCTCGGACAAGATTAAATTCCTTAATCCG
GATAGGGAGTACGACTTCAGAGATCTCACTTGGTGCAT
CAACCCGCCAGAGAGGATCAAACCTAGATTATGATCAAT
ACTGTGCAGATGTGGCTGCTGAAGAGCTCATGAATGCA
TTGGTGAACCTCAACTCTACTGGAGACCAGAACAACCAC
TCAGTTCCTAGCTGTCTCAAAGGGGAACTGCTCAGGGC
CCACTACAATCAGAGGTCAATTCTCAAACATGTCGCTG
TCCTTGTTGGACTTGTACTTAGGTCGAGGTTACAATGTG
TCATCTATAGTCACTATGACATCCCAGGGAATGTATGG
GGGAACCTACCTAGTTGAAAAGCCTAATCTGAACAGCA
AAGGGTCAGAGTTGTCACAACCTGAGCATGTACCGAGTG
TTTGAAGTAGGTGTGATCAGAAACCCGGGTTTGGGGGC
TCCGGTGTTCCATATGACAAACTATTTTGAGCAACCAG
TCAGTAATGGTCTCGGCAACTGTATGGTGGCTTTGGGG
GAGCTCAAACCTCGCAGCCCTTTGTCACGGGGACGATTC
TATCATAATTCCCTATCAGGGATCAGGGAAAGGTGTCA
GCTTCCAGCTCGTCAAGCTGGGTGTCTGGAAATCCCCA
ACCGACATGCAATCCTGGGTCCCCTTATCAACGGATGA
TCCAGTGGTAGACAGGCTTTACCTCTCATCTCACAGAG
GTGTCATCGCTGACAATCAAGCAAAATGGGCTGTCCCG
ACAACACGAACAGATGACAAGTTGCGAATGGAGACAT
GCTTCCAGCAGGCGTGTAAGGTAAAATCCAAGCACTC
TGCGAGAATCCCGAGTGGGTACCATTGAAGGATAACAG
GATTCCTTCATACGGGGTCCTGTCTGTTGATCTGAGTCT
GACGGTTGAGCTTAAAATCAAAATTGCTTCGGGATTCG
GGCCATTGATCACACACGGCTCAGGGATGGACCTATAC
AAATCCAACCTGCAACAATGTGTATTGGCTGACTATTCC
GCCAATGAGAAATCTAGCCTTAGGCGTAATCAACACAT
TGGAGTGGATACCGAGATTCAAGGTTAGTCCCAACCTC
TTCACTGTCCCAATTAAGGAAGCAGGCGAAGACTGCCA
TGCCCCAACATACCTACCTGCGGAGGTGGACGGTGATG
TCAAACCTCAGTTCCAACCTGGTGATTCTACCTGGTCAA
GATCTCCAATATGTTTTGGCAACCTACGATACCTCCAG
GGTTGAGCATGCTGTGGTTTATTACGTTTACAGCCCAA

GCCGCTTCTTCTTCTTATCCTTTTAGGTTGCCTAT
AAAGGGGGTCCCAATCGAACTACAAGTGAATGCTTCA
CATGGGATCAAAAACCTCTGGTGCCGTCCTTCTGTGTG
CTTGCGGACTCAGAATCCGGTGGACTTATCACTCACTCT
GGGATGGTGGGCATGGGAGTCAGCTGCACAGCTACCCG
GGAAGATGGAACCAATCGCAGATAA GC_H_MEASLES_
G*GGGAAATAAGAGAGAAAAGAAGAGTAAGAAGAAAT 43 B3
ATAAGAGCCACCATGTCACCGCAACGAGACCGGATAA mRNA Sequence
ATGCCTTCTACAAAGATAACCCTTATCCCAAGGGAAGT (assumes T100 tail)
AGGATAGTTATTAACAGAGAACATCTTATGATTGACAG Sequence Length:
ACCCTATGTTCTGCTGGCTGTTCTGTTTCGTCATGTTTCT 2126
GAGCTTGATCGGATTGCTGGCAATTGCAGGCATTAGAC
TTCATCGGGCAGCCATCTACACCGCGGAGATCCATAAA
AGCCTCAGTACCAATCTGGATGTGACTAACTCCATCGA
GCATCAGGTCAAGGACGTGCTGACACCACTCTTTAAAA
TCATCGGGGATGAAGTGGGCCTGAGAACACCTCAGAG
ATTCCTGACCTAGTGAAATTCATCTCGGACAAGATTA
AATTCCTTAATCCGGATAGGGAGTACGACTTCAGAGAT
CTCACTTGGTGCATCAACCCGCCAGAGAGGATCAAAC
AGATTATGATCAATACTGTGCAGATGTGGCTGCTGAAG
AGCTCATGAATGCATTGGTGAACCTCAACTCTACTGGAG
ACCAGAACAACCACTCAGTTCCTAGCTGTCTCAAAGGG
AAACTGCTCAGGGCCCACTACAATCAGAGGTCAATTCT
CAAACATGTCGCTGTCCTTGTTGGACTTGTAAGGTC
GAGGTTACAATGTGTCATCTATAGTCACTATGACATCC
CAGGGAATGTATGGGGGAACCTACCTAGTTGAAAAGCC
TAATCTGAACAGCAAAGGGTCAGAGTTGTCACAACTGA
GCATGTACCGAGTGTTTGAAGTAGGTGTGATCAGAAAC
CCGGGTTTGGGGGCTCCGGTGTTCATATGACAAACTA
TTTTGAGCAACCAGTCAGTAATGGTCTCGGCAACTGTA
TGGTGGCTTTGGGGGAGCTCAAACCTCGCAGCCCTTTGT
CACGGGGACGATTCTATCATAATTCCCTATCAGGGATC
AGGGAAAGGTGTCAGCTTCCAGCTCGTCAAGCTGGGTG
TCTGGAAATCCCCAACCGACATGCAATCCTGGGTCCCC
TTATCAACGGATGATCCAGTGGTAGACAGGCTTTACCT
CTCATCTCACAGAGGTGTCATCGCTGACAATCAAGCAA
AATGGGCTGTCCCGACAACACGAACAGATGACAAGTTG
CGAATGGAGACATGCTTCCAGCAGGCGTGTAAGGTAA
AATCCAAGCACTCTGCGAGAATCCCGAGTGGGTACCAT
TGAAGGATAACAGGATTCCTTCATACGGGGTCCTGTCT
GTTGATCTGAGTCTGACGGTTGAGCTTAAAATCAAAT
TGCTTCGGGATTTCGGGCCATTGATCACACACGGCTCAG
GGATGGACCTATACAAATCCAACCTGCAACAATGTGTAT
TGGCTGACTATTCCGCCAATGAGAAATCTAGCCTTAGG
CGTAATCAACACATTGGAGTGGATAACCGAGATTCAAGG
TTAGTCCCAACCTCTTCACTGTCCCAATTAAGGAAGCA
GGCGAAGACTGCCATGCCCCAACATACTACCTGCGGA
GGTGGACGGTGATGTCAAACCTCAGTTCCAACCTGGTGA
TTCTACCTGGTCAAGATCTCCAATATGTTTTGGCAACCT
ACGATACCTCCAGGGTTGAGCATGCTGTGGTTTATTAC

GTTTACAGCCCAAGCCGCTCATTTTCTTACTTTTATCCT
TTTAGGTTGCCTATAAAGGGGGTCCCAATCGAACTACA
AGTGAATGCTTCACATGGGATCAAAAACCTCTGGTGCC
GTCATTCTGTGTGCTTGCGGACTCAGAATCCGGTGGA
CTTATCACTCACTCTGGGATGGTGGGCATGGGAGTCAG
CTGCACAGCTACCCGGGAAGATGGAACCAATCGCAGAT
AATGATAATAGGCTGGAGCCTCGGTGGCCAAGCTTCTT
GCCCCTTGGGCCTCCCCCAGCCCCTCCTCCCCTTCCTG
CACCCGTACCCCGTGGTCTTTGAATAAAGTCTGAGTG
GGCGGCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAATC TAG GC_H_MEASLES_
TCAAGCTTTTGGACCCTCGTACAGAAGCTAATACGACT 44 D8
CACTATAGGGAAATAAGAGAGAAAAGAAGAGTAAGAA Sequence, NT (5'
GAAATATAAGAGCCACCATGTCACCACAACGAGACCG UTR, ORF, 3'
GATAAATGCCTTCTACAAAGACAACCCCCATCCTAAGG UTR)
GAAGTAGGATAGTTATTAACAGAGAACATCTTATGATT Sequence Length:
GATAGACCTTATGTTTTGCTGGCTGTTCTATTCGTCATG 2065
TTTCTGAGCTTGATCGGGTTGCTAGCCATTGCAGGCATT
AGACTTCATCGGGCAGCCATCTACACCGCAGAGATCCA
TAAAAGCCTCAGCACCAATCTGGATGTAACATACTCAA
TCGAGCATCAGGTAAAGGACGTGCTGACACCACTCTTC
AAGATCATCGGTGATGAAGTGGGCTTGAGGACACCTCA
GAGATTCACTGACCTAGTGAAGTTCATCTCTGACAAGA
TTAAATTCCTTAATCCGGACAGGGAATACGACTTCAGA
GATCTCACTTGGTGTATCAACCCGCCAGAGAGAATCAA
ATTGGATTATGATCAATACTGTGCAGATGTGGCTGCTG
AAGAACTCATGAATGCATTGGTGAACCTCAACTCTACTG
GAGACCAGGGCAACCAATCAGTTCCTAGCTGTCTCAA
GGGAACTGCTCAGGGCCCACTACAATCAGAGGCCAAT
TCTCAAACATGTCGCTGTCCCTGTTGGACTTGTATTTAA
GTCGAGGTTACAATGTGTCATCTATAGTCACTATGACA
TCCCAGGGAATGTACGGGGGAACTTACCTAGTGGA
GCCTAATCTGAGCAGCAAAGGGTCAGAGTTGTCACAAC
TGAGCATGCACCGAGTGTTTGAAGTAGGTGTTATCAGA
AATCCGGGTTTGGGGGCTCCGGTATTCCATATGACAAA
CTATCTTGAGCAACCAGTCAGTAATGATTCAGCAACT
GCATGGTGGCTTTGGGGGAGCTCAAGTTCGCAGCCCTC
TGTCACAGGGAAGATTCTATCACAATTCCCTATCAGGG
ATCAGGGAAAGGTGTCAGCTTCCAGCTTGTCAAGCTAG
GTGTCTGGAAATCCCAACCGACATGCAATCCTGGGTC
CCCCTATCAACGGATGATCCAGTGATAGACAGGCTTTA
CCTCTCATCTCACAGAGGCGTTATCGCTGACAATCAAG
CAAAATGGGCTGTCCCGACAACACGGACAGATGACAA
GTTGCGAATGGAGACATGCTTCCAGCAGGCGTGTAAGG
GTAAAATCCAAGCACTTTGCGAGAATCCCGAGTGGACA
CCATTGAAGGATAACAGGATTCCTTCATACGGGGTCTT
GTCTGTTGATCTGAGTCTGACAGTTGAGCTTAAAATCA
AAATTGTTTCAGGATTCGGGCCATTGATCACACACGGT
TCAGGGATGGACCTATACAAATCCAACCACAACAATAT

GTATTGGCTTATCCCTCCCAATGAAGAACCCTGGCCT
TAGGTGTAATCAACACATTGGAGTGGATACCGAGATTC
AAGGTTAGTCCCAACCTCTTCACTGTTCCAATTAAGGA
AGCAGGCGAGGACTGCCATGCCCCAACATACCTACCTG
CGGAGGTGGATGGTGTGTCAAACCTCAGTTCCAATCTG
GTGATTCTACCTGGTCAAGATCTCCAATATGTTCTGGCA
ACCTACGATACTTCCAGAGTTGAACATGCTGTAGTTTAT
TACGTTTACAGCCCAAGCCGCTCATTTTCTTACTTTTAT
CCTTTTAGGTTGCCTGTAAGGGGGGTCCCCATTGAATTA
CAAGTGGAATGCTTCACATGGGACCAAAAACCTCTGGTG
CCGTCACTTCTGTGTGCTTGCGGACTCAGAATCTGGTG
ACATATCACTCACTCTGGGATGGTGGGCATGGGAGTCA
GCTGCACAGCCACTCGGGAAGATGGAACCAGCCGCAG
ATAGTGATAATAGGCTGGAGCCTCGGTGGCCAAGCTTC
TTGCCCCCTTGGGCCTCCCCCAGCCCCCTCCTCCCCCTTCC
TGCACCCCGTACCCCGTGGTCTTTGAATAAAGTCTGAG TGGGCGGC GC_H_MEASLES_
ATGTCACCACAACGAGACCGGATAAATGCCTTCTACAA 45 D8
AGACAACCCCCATCCTAAGGGAAGTAGGATAGTTATTA ORF Sequence, NT
ACAGAGAACATCTTATGATTGATAGACCTTATGTTTTGC
TGGCTGTTCTATTCGTCATGTTTCTGAGCTTGATCGGGT
TGCTAGCCATTGCAGGCATTAGACTTCATCGGGCAGCC
ATCTACACCGCAGAGATCCATAAAAGCCTCAGCACCAA
TCTGGATGTAATAACTCAATCGAGCATCAGGTTAAGG
ACGTGCTGACACCACTCTTCAAGATCATCGGTGATGAA
GTGGGCTTGAGGACACCTCAGAGATTCACTGACCTAGT
GAAGTTCATCTCTGACAAGATTAAATTCCTTAATCCGG
ACAGGGAATACGACTTCAGAGATCTCACTTGGTGTATC
AACCCGCCAGAGAGAATCAAATTGGATTATGATCAATA
CTGTGCAGATGTGGCTGCTGAAGAACTCATGAATGCAT
TGGTGAACCTCAACTCTACTGGAGACCAGGGCAACCAAT
CAGTTCCTAGCTGTCTCAAAGGGAACTGCTCAGGGCC
CACTACAATCAGAGGCCAATTCTCAAACATGTCGCTGT
CCCTGTTGGACTTGTAATTTAAGTCGAGGTTACAATGTGT
CATCTATAGTCACTATGACATCCCAGGGAATGTACGGG
GGAACCTACCTAGTGGAAGAGCCTAATCTGAGCAGCAA
AGGGTCAGAGTTGTCACAACTGAGCATGCACCGAGTGT
TTGAAGTAGGTGTTATCAGAAATCCGGGTTTGGGGGCT
CCGGTATTCCATATGACAACTATCTTGAGCAACCAGT
CAGTAATGATTTAGCAACTGCATGGTGGCTTTGGGGG
AGCTCAAGTTCGCAGCCCTCTGTCACAGGGAAGATTCT
ATCACAATTCCCTATCAGGGATCAGGGAAAGGTGTGAG
CTTCCAGCTTGTCAAGCTAGGTGTCTGGAAATCCCCAA
CCGACATGCAATCCTGGGTCCCCCTATCAACGGATGAT
CCAGTGATAGACAGGCTTTACCTCTCATCTCACAGAGG
CGTTATCGCTGACAATCAAGCAAAATGGGCTGTCCCGA
CAACACGGACAGATGACAAGTTGCGAATGGAGACATG
CTTCCAGCAGGCGTGTAAGGGTAAAATCCAAGCACTTT
GCGAGAATCCCGAGTGGACACCATTGAAGGATAACAG
GATTCCTTCATACGGGGTCTTGTCTGTTGATCTGAGTCT
GACAGTTGAGCTTAAAATCAAATTTGTTTCAGGATTCTG

GGCCATTGATCACACACGGTTTCAGGGATGGACCTATAC
AAATCCAACCACAACAATATGTATTGGCTGACTATCCC
GCCAATGAAGAACCTGGCCTTAGGTGTAATCAACACAT
TGGAGTGGATACCGAGATTCAAGGTTAGTCCCAACCTC
TTCACTGTTCCAATTAAGGAAGCAGGCGAGGACTGCCA
TGCCCCAACATACCTACCTGCGGAGGTGGATGGTGATG
TCAAACCTCAGTTCCAATCTGGTGATTCTACCTGGTCAAG
ATCTCCAATATGTTCTGGCAACCTACGATACTTCCAGA
GTTGAACATGCTGTAGTTTATTACGTTTACAGCCCAAGC
CGCTCATTTTCTTACTTTTATCCTTTTAGGTTGCCTGTAA
GGGGGGTCCCCATTGAATTACAAGTGGAATGCTTCACA
TGGGACCAAAAACCTCTGGTGCCGTCACCTTCTGTGTGCTT
GCGGACTCAGAATCTGGTGGACATATCACTCACTCTGG
GATGGTGGGCATGGGAGTCAGCTGCACAGCCACTCGGG
AAGATGGAACCAGCCGCAGATAG GC_H_MEASLES_
G*GGGAAATAAGAGAGAAAAGAAGAGTAAGAAGAAAT 46 D8
ATAAGAGCCACCATGTCACCACAACGAGACCGGATAA mRNA Sequence
ATGCCTTCTACAAAGACAACCCCCATCCTAAGGGAAGT (assumes T100 tail)
AGGATAGTTATTAACAGAGAACATCTTATGATTGATAG Sequence Length:
ACCTTATGTTTTGCTGGCTGTTCTATTCGTCATGTTTCTG 2126
AGCTTGATCGGGTTGCTAGCCATTGCAGGCATTAGACT
TCATCGGGCAGCCATCTACACCGCAGAGATCCATAAAA
GCCTCAGCACCAATCTGGATGTAACATACTCAATCGAG
CATCAGGTTAAGGACGTGCTGACACCACTCTTCAAGAT
CATCGGTGATGAAGTGGGCTTGAGGACACCTCAGAGAT
TCACTGACCTAGTGAAGTTCATCTCTGACAAGATTAAA
TTCCTTAATCCGGACAGGGAATACGACTTCAGAGATCT
CACTTGGTGTATCAACCCGCCAGAGAGAATCAAATTGG
ATTATGATCAATACTGTGCAGATGTGGCTGCTGAAGAA
CTCATGAATGCATTGGTGAACCTCAACTCTACTGGAGAC
CAGGGCAACCAATCAGTTCCTAGCTGTCTCAAAGGGAA
ACTGCTCAGGGCCCACTACAATCAGAGGCCAATTCTCA
AACATGTCGCTGTCCCTGTTGGACTTGTATTTAAGTCGA
GGTTACAATGTGTCATCTATAGTCACTATGACATCCCA
GGGAATGTACGGGGGAACCTTAGTGGAAGGCCT
AATCTGAGCAGCAAAGGGTCAGAGTTGTCACAACTGAG
CATGCACCGAGTGTTTGAAGTAGGTGTTATCAGAAATC
CGGGTTTGGGGGCTCCGGTATTCCATATGACAAAC TAT
CTTGAGCAACCAGTCAGTAATGATTTTCAGCAACTGCAT
GGTGGCTTTGGGGGAGCTCAAGTTCGCAGCCCTCTGTC
ACAGGGAAGATTCTATCACAATCCCTATCAGGGATCA
GGGAAAGGTGTCAGCTTCCAGCTTGTCAAGCTAGGTGT
CTGGAAATCCCCAACCGACATGCAATCCTGGGTCCCCC
TATCAACGGATGATCCAGTGATAGACAGGCTTTACCTC
TCATCTCACAGAGGCGTTATCGCTGACAATCAAGCAAA
ATGGGCTGTCCCGACAACACGGACAGATGACAAGTTGC
GAATGGAGACATGCTTCCAGCAGGCGTGTAAGGGTAA
AATCCAAGCACTTTGCGAGAATCCCGAGTGGACACCAT
TGAAGGATAACAGGATTCCCTTCATACGGGGTCTTGTCT
GTTGATCTGAGTCTGACAGTTGAGCTTAAAATCAAAT

TGTTTCAGGCTTGCATTCACACACGCTTCAG
GGATGGACCTATACAAATCCAACCACAACAATATGTAT
TGGCTGACTATCCCGCCAATGAAGAACCTGGCCTTAGG
TGTAATCAACACATTGGAGTGGATACCGAGATTCAAGG
TTAGTCCCAACCTCTTCACTGTTCCAATTAAGGAAGCA
GGCGAGGACTGCCATGCCCCAACATACCTACCTGCGGA
GGTGGATGGTGTGTCAAACCTCAGTTCCAATCTGGTGA
TTCTACCTGGTCAAGATCTCCAATATGTTCTGGCAACCT
ACGATACTTCCAGAGTTGAACATGCTGTAGTTTATTAC
GTTTACAGCCCAAGCCGCTCATTTTCTTACTTTTATCCT
TTTAGGTTGCCTGTAAGGGGGGTCCCCATTGAATTACA
AGTGGAATGCTTCACATGGGACCAAAAACCTCTGGTGCC
GTCACCTTCTGTGTGCTTGCGGACTCAGAATCTGGTGGA
CATATCACTCACTCTGGGATGGTGGGCATGGGAGTCAG
CTGCACAGCCACTCGGGAAGATGGAACCAGCCGCAGA
TAGTGATAATAGGCTGGAGCCTCGGTGGCCAAGCTTCT
TGCCCCTTGGGCCTCCCCCAGCCCCTCCTCCCCTTCCT
GCACCCGTACCCCGTGGTCTTTGAATAAAGTCTGAGT
GGGCGGCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAT CTAG MeV mRNA
Sequences GC_F_MEASLES_ UCAAGCUUUUGGACCCUCGUACAGAAGCUAAUACGAC
69 B3.1 UCACUAUAGGGAAAUUAGAGAGAAAAGAAGAGUAAG Sequence, NT (5'
AAGAAUAUAAGAGCCACCAUGGGUCUCAAGGUGAA UTR, ORF, 3'
CGUCUCUGCCGUUAUCAUGGCAGUACUGUUAACUCUC UTR)
CAAACACCCGCCGGUCAAAUUCAUUGGGGCAAUCUCU Sequence Length:
CUAAGAUAGGGGUAGUAGGAAUAGGAAGUGCAAGCU 1864
ACAAAGUUAUGACUCGUUCCAGCCAUCAAUCAUUAGU
CAUAAAAUUAUGCCCAAUAUAACUCUCCUCAUAAC
UGCACGAGGGUAGAGAUUGCAGAAUACAGGAGACUA
CUAAGAACAGUUUUGGAACCAAUUAGGGAUGCACUU
AAUGCAAUGACCCAGAACAUAAGGCCGGUUCAGAGCG
UAGCUUCAAGUAGGAGACACAAGAGAUUUGCGGGAG
UAGUCCUGGCAGGUGCGGCCCAGGUGUUGCCACAGC
UGCUCAGAUAAACAGCCGGCAUUGCACUUCACCGGUCC
AUGCUGAACUCUCAGGCCAUUCGACAAUCUGAGAGCGA
GCCUGGAAACUACUAAUCAGGCAAUUGAGGCAAUCAG
ACAAGCAGGGCAGGAGAUGAUUAUUGGCUGUUCAGGG
UGUCCAAGACUACAUAUAUAAUGAGCUGAUACCGUCU
AUGAACCAGCUAUCUUGUGAUCUAAUCGGUCAGAAGC
UCGGGCUCAAAUUGCUUAGAUACUAUACAGAAAUCCU
GUCAUUAUUUGGCCCCAGCCUACGGGACCCCAUAUCU
GCGGAGAUUAUCUAUCCAGGCUUUGAGUUAUGCACUU
GGAGGAGAUUAUAUAAGGUGUAGAAAAGCUCGGA
UACAGUGGAGGCGAUUUACUAGGCAUCUUAGAGAGC
AGAGGAAUAAAGGCUCGGAUAAACUCACGUCGACACAG
AGUCCUACUUCAUAGUCCUCAGUAUAGCCUAUCCGAC
GCUGUCCGAGAUUAAGGGGGUGAUUGUCCACCGGCUA
GAGGGGGUCUCGUACAACAUAGGCUCUCAAGAGUGG
UAUACCACUGUGCCCAAGUAUGUUGCAACCCAAGGGU

ACCUUAUAGCCAAUUGUGCAUCAAUUCUUUGUAAGU
UCAUGCCAGAGGGGACUGUGUGCAGCCAAAUGCCUU
GUACCCGAUGAGUCCUCUGCUCCAAGAAUGCCUCCGG
GGGUCCACCAAGUCCUGUGCUCGUACACUCGUAUCCG
GGUCUUUUGGGAACCGGUUCAUUUUAUCACAAGGGA
ACCUAAUAGCCAAUUGUGCAUCAAUUCUUUGUAAGU
GUUACACAACAGGUACGAUUAAUUAUCAAGACCCUGA
CAAGAUCCUAACAUAACAUUGCUGCCGAUCGCUGCCCG
GUAGUCGAGGUGAACGGCGUGACCAUCCAAGUCGGGA
GCAGGAGGUAUCCAGACGCUGUGUACUUGCACAGAAU
UGACCUCGGUCCUCCCAUAUCAUUGGAGAGGUUGGAC
GUAGGGACAAAUCUGGGGAAUGCAAUUGCCAAAUUG
GAGGAUGCCAAGGAAUUGUUGGAAUCAUCGGACCAG
AAUUGGAGAAGUAUGAAAGGUUUAUCGAGCACUAGC
AUAGUCUACAUCCUGAUUGCAGUGUGUCUUGGAGGG
UUGAUAGGGAUCCCCACUUUAAUAUGUUGCUGCAGG
GGGCGUUGUAACAAAAAGGGAGAACAAGUUGGUAUG
UCAAGACCAGGCCUAAAGCCUGACCUUACAGGAACAU
CAAAAUCCUAUGUAAGAUCGCUUUGAUGAUAAUAGG
CUGGAGCCUCGGUGGCCAAGCUUCUUGCCCCUUGGGC
CUCCCCCAGCCCCUCCUCCCCUUCUGCACCCGUACC
CCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC GC_F_MEASLES_
AUGGGUCUCAAGGUGAACGUCUCUGCCGUAUUCAUGG 70 B3.1
CAGUACUGUUAACUCUCCAAACACCCGCCGGUCAAAU ORF Sequence, NT
UCAUUGGGGCAAUCUCUCUAAGAUAAGGGGUAGUAGG
AAUAGGAAGUGCAAGCUACAAAGUUAUGACUCGUUC
CAGCCAUCAUCAUUAAGUCAUAAAAUUAUUGCCCAAU
AUAACUCUCCUCAAUAAACUGCACGAGGGUAGAGAUUG
CAGAAUACAGGAGACUACUAAGAACAGUUUUGGAAC
CAAUUAGGGAUGCACUUAUUGCAAUGACCCAGAACAU
AAGGCCGGUUCAGAGCGUAGCUUCAAGUAGGAGACAC
AAGAGAUUUGCGGGAGUAGUCCUGGCAGGUGCGGCCC
UAGGUGUUGCCACAGCUGCUCAGAUAAACAGCCGGCAU
UGCACUUCACCGGUCCAUGCUGAACUCUCAGGCCAUC
GACAAUCUGAGAGCGAGCCUGGAAACUACUAAUCAGG
CAAUUGAGGCAAUCAGACAAGCAGGGCAGGAGAUGA
UAUUGGCUGUUCAGGGUGUCCAAGACUACAUCAAUA
AUGAGCUGAUACCGUCUAUGAACCAGCUAUCUUGUGA
UCUAAUUCGGUCAGAAGCUCGGGCUCAAAUUGCUUAGA
UACUAUACAGAAAUCCUGUCAUUAUUUGGCCCCAGCC
UACGGGACCCCAUAUCUGCGGAGAUAUCAUUCAGGC
UUUGAGUUAUGCACUUGGAGGAGAUUCAUAAGGU
GUUAGAAAAGCUCGGAUACAGUGGAGGCGAUUUACU
AGGCAUCUUAGAGAGCAGAGGAAUAAAGGCUCGGAU
AACUCACGUCGACACAGAGUCCUACUUCAUAGUCCUC
AGUAUAGCCUAUCCGACGCUGUCCGAGAUUAAGGGGG
UGAUUGUCCACCGGCUAGAGGGGGUCUCGUACAACAU
AGGCUCUCAAGAGUGGUUAUACCACUGUGCCCAAGUAU
GUUGCAACCCAAGGGUACCUUAUCUCGAAUUUUGAUG
AGUCAUCAUGUACUUCAUGCCAGAGGGGACUGUGU

GCGAGCCAAAGUACCCGAGUCCUCUCUCU
CCAAGAAUGCCUCCGGGGGUCCACCAAGUCCUGUGCU
CGUACACUCGUAUCCGGGUCUUUUGGGAACCGGUUCA
UUUUUAUCACAAGGGAACCUAAUAGCCAAUUGUGCAUC
AAUUCUUUGUAAGUGUACACAACAGGUACGAUUAU
UAAUCAAGACCCUGACAAGAUCCUAACAUACAUUGCU
GCCGAUCGCUGCCCGGUAGUCGAGGUGAACGGCGUGA
CCAUCCAAGUCGGGAGCAGGAGGUAUCCAGACGCUGU
GUACUUGCACAGAAUUGACCUCGGUCCUCCCAUAUCA
UUGGAGAGGUUGGACGUAGGGACAAAUCUGGGGAAU
GCAAUUGCCAAAUUGGAGGAUGCCAAGGAAUUGUUG
GAAUCAUCGGACCAGAUAUUGAGAAGUAUGAAAGGU
UUAUCGAGCACUAGCAUAGUCUACAUCUGAUUUGCAG
UGUGUCUUGGAGGGUUGAUAGGGAUCCCCACUUUAA
UAUGUUGCUGCAGGGGGCGUUGUAACAAAAAGGGAG
AACAAGUUGGU AUGUCAAGACCAGGCCUAAAGCCUGA
CCUUACAGGAACAUCAAAAUCCUAUGUAAGAUCGCUU UGA GC_F_MEASLES_
G*GGGAAAU AAGAGAGAAAAGAAGAGUAAGAAGAAA 71 B3.1
UAUAAGAGCCACCAUGGGUCUCAAGGUGAACGUCUCU mRNA Sequence
GCCGUAUUCAUGGCAGUACUGUUAACUCUCCAAACAC (assumes T100 tail)
CCGCCGGUCAAAUUCAUUGGGGGCAAUCUCUCUAAGAU mRNA Sequence
AGGGGUAGUAGGAAUAGGAAGUGCAAGCUACAAAGU Length: 1925
UAUGACUCGUUCCAGCCAUCAAUCAUUAAGUCAUAAAA
UUA AUGCCCAUAUAACUCUCCUCAUAACUGCACGA
GGGUAGAGAUUGCAGAAUACAGGAGACUACUAAGAA
CAGUUUUGGAACCAAUUAAGGGAUGCACUUA AUGCAA
UGACCCAGAACAUAAAGGCCGGUUCAGAGCGUAGCUUC
AAGUAGGAGACACAAGAGAUUUGCGGGAGUAGUCCU
GGCAGGUGCGGCCCUAGGUGUUGCCACAGCUGCUCAG
AUAACAGCCGGCAUUGCACUUCACCGGUCCAUGCUGA
ACUCUCAGGCCAUUCGACAAUCUGAGAGCGAGCCUGGA
AACUACUAAUCAGGCAAUUGAGGCAAUCAGACAAGCA
GGGCAGGAGAUGAUAUUGGCUGUUCAGGGUGUCCAA
GACUACAUCAAUAAUGAGCUGAUACCGUCUAUGAACC
AGCUAUCUUGUGAUCUAAUCGGUCAGAAGCUCGGGCU
CAA AUUGCUUAGAUACUAUACAGAAAUCCUGUCAUU
AUUUGGCCCCAGCCUACGGGACCCCAUAUCUGCGGAG
AUAUCUAUCCAGGCUUUGAGUUAUGCACUUGGAGGA
GAUAUCAAUAAAGGUGUUAAGAAAAGCUCGGAUACAGU
GGAGGCGAUUUACUAGGCAUCUUAGAGAGCAGAGGA
AUAAAGGCUCGGAUAACUCACGUCGACACAGAGUCCU
ACUUCAUAGUCCUCAGUAUAGCCUAUCCGACGCUGUC
CGAGAUUAAGGGGGUGAUUGUCCACCGGCUAGAGGG
GGUCUCGUACAACAUAGGCUCUCAAGAGUGGUUAUACC
ACUGUGCCCAAGUAUGUUGCAACCCAAGGGUACCUUA
UCUCGAAUUUUGAUGAGUCAUCAUGUACUUUCAUGCC
AGAGGGGACUGUGUGCAGCCAAA AUGCCUUGUACCCG
AUGAGUCCUCUGCUCCAAGAAUGCCUCCGGGGGUCCA
CCAAGUCCUGUGCUCGUACACUCGUAUCCGGGUCUUU
UGGGAACCGGUUCAUUUUAUCACAAGGGAACCUAAU

AGCCAAUUCUUCUUGUAAGUGUUACAC
AACAGGUACGAUUAUUAUCAAGACCCUGACAAGAUC
CUAACAUACAUUGCUGCCGAUCGCGUGCCCGGUAGUCG
AGGUGAACGGCGUGACCAUCCAAGUCGGGAGCAGGAG
GUAUCCAGACGCUGUGUACUUGCACAGAAUUGACCUC
GGUCCUCCCAUAUCAUUGGAGAGGUUGGACGUAGGG
ACAAAUCUGGGGAAUGCAAUUGCCAAAUUGGAGGAU
GCCAAGGAAUUGUUGGAAUCAUCGGACCAGAUAUUG
AGAAGUAUGAAAGGUUUAUUCGAGCACUAGCAUAGUC
UACAUCCUGAUUUGCAGUGUGUCUUGGAGGGGUUGAUA
GGGAUCCCCACUUUAAUAUGUUGCUGCAGGGGGCGUU
GUAACAAAAAGGGGAGAACAGUUGGUAUGUCAAGAC
CAGGCCUAAAGCCUGACCUUACAGGAACAUCAAAAUC
CUAUGUAAGAUCGCUUUGAUGAUAAUAGGCUGGAGC
CUCGGUGGCCAAGCUUCUUGCCCCUUGGGCCUCCCC
CAGCCCCUCCUCCCCUUCUGCACCCGUACCCCCGUGG
UCUUUGAAUAAAGUCUGAGUGGGCGGCAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAUCUAG GC_F_MEASLES_
UCAAGCUUUUGGACCCUCGUACAGAAGCUAAUACGAC 72 D8
UCACUAUAGGGAAUAAGAGAGAAAAGAAGAGUAAG Sequence, NT (5'
AAGAAUAUAAGAGCCACCAUGGGUCUCAAGGUGAA UTR, ORF, 3'
CGUCUCUGUCAUAUUCAUGGCAGUACUGUUAACUCUU UTR)
CAAACACCCACCGGUCAAAUCCAUUGGGGCAAUCUCU Sequence Length:
CUAAGAUAGGGGUGGUAGGGGUAGGAAGUGCAAGCU 1864
ACAAAGUUAUGACUCGUUCCAGCCAUCAUCAUUAGU
CAUAAAGUUAUUGCCCAAUAUAACUCUCCUCAACAAU
UGCACGAGGGUAGGGAUUUGCAGAAUACAGGAGACUA
CUGAGAACAGUUCUGGAACCAAUUAGAGAUGCACUU
AAUGCAAUGACCCAGAAUAUAAGACCGGUUCAGAGU
GUAGCUUCAAGUAGGAGACACAAGAGAUUUGCGGGA
GUUGUCCUGGCAGGUGCGGCCCUAGGCGUUGCCACAG
CUGCUCAAAUAACAGCCGGUAUUGCACUUCACCAGUC
CAUGCUGAACUCUCAAGCCAUCGACAAUCUGAGAGCG
AGCCUAGAAACUACUAAUCAGGCAAUUGAGGCAAUCA
GACAAGCAGGGCAGGAGAUGAUAAUUGGCUGUUCAGG
GUGUCCAAGACUACAUCAAUAAUUGAGCUGAUACCGUC
UAUGAAUCAACUAUCUUGUGAUUUAAUCGGCCAGAA
GCUAGGGCUCAAAUUGCUCAGAUACUAUACAGAAAUC
CUGUCAUUAUUUGGCCCCAGCUUACGGGACCCCAUAU
CUGCGGAGAUAUCUAUCCAGGCUUUGAGCUAUGCGCU
UGGAGGAGAUUCAAUAAAGGUGUUGGAAAAGCUCGG
AUACAGUGGAGGUGAUUCUACUGGGCAUCUUAAGAGAG
CAGAGGAAUAAAGGCCCGGAUAACUCACGUCGACACA
GAGUCCUACUUCAUUGUACUCAGUAUAGCCUAUCCGA
CGCUAUCCGAGAUUAAGGGGGUGAUUGUCCACCGGCU
AGAGGGGGUCUCGUACAACAUAAGGCUCUCAAGAGUG
GUAUACCACUGUGCCCAAGUAUGUUGCAACCCAAGGG
UACCUUAUCUCGAAUUUUGAUGAGUCAUCAUGCACUU

UCAUGCCAGAGUGUGCAGGCCAGAAUGCCUU
GUACCCGAUGAGUCCUCUGCUCCAAGAAUGCCUCCGG
GGGUCCACUAAGUCCUGUGCUCGUACACUCGUAUCCG
GGUCUUUUCGGGAACCGGUUCAUUUUAUCACAGGGGA
ACCUAAUAGCCAAUUGUGCAUCAAUCCUUUGCAAGUG
UUACACAACAGGAACAAUCAUUAUAUCAAGACCCUGAC
AAGAUCCUAACAUAUAUUGCUGCCGAUCACUGCCCGG
UGGUCGAGGUGAAUGGGCGUGACCAUCCAAGUCGGGA
GCAGGAGGUAUCCGGACGCUGUGUACUUGCACAGGAU
UGACCUCGGUCCUCCCAUAUCUUUGGAGAGGUUGGAC
GUAGGGACAAAUUCUGGGGAAUGCAAUUGCUAAGUUG
GAGGAUGCCAAGGAAUUGUUGGAGUCAUCGGACCAG
AUAUUGAGGAGUAUGAAAGGUUUAUCGAGCACUAGU
AUAGUUUACAUCUGAUUUGCAGUGUGUCUUGGAGGA
UUGAUAGGGAUCCCCGCUUUAUAUGUUGCUGCAGG
GGGCGUUGUAACAAGAAGGGAGAACAAGUUGGUUUG
UCAAGACCAGGCCUAAAGCCUGAUCUUACAGGAACAU
CAAAAUCCUAUGUAAGGUCACUCUGAUGAUAAUAGG
CUGGAGCCUCGGUGGCCAAGCUUCUUGCCCCUUGGGC
CUCCCCCAGCCCCUCCUCCCCUUCUGCACCCGUACC
CCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC GC_F_MEASLES_
AUGGGUCUCAAGGUGAACGUCUCUGUCAUAUUCAUG 73 D8
GCAGUACUGUUAACUCUUCAAACACCCACCGGUCAAA ORF Sequence, NT
UCCAUUGGGGCAAUCUCUCUAAGAUAGGGGUGGUAG
GGGUAGGAAGUGCAAGCUACAAAGUUAUGACUCGUU
CCAGCCAUCAAUCAUUAUGUCAUAAAGUUAUUGCCCAA
UAUAACUCUCCUCAACAAUUGCACGAGGGUAGGGAUU
GCAGAAUACAGGAGACUACUGAGAACAGUUCUGGAA
CCAAUUAGAGAUGCACUUAUUGCAAUGACCCAGAAUA
UAAGACCGGUUCAGAGUGUAGCUUCAAGUAGGAGAC
ACAAGAGAUUUGCGGGAGUUGUCCUGGCAGGUGCGG
CCCUAGGCGUUGCCACAGCUGCUCAAUAAACAGCCGG
UAUUGCACUUCACCAGUCCAUGCUGAACUCUCAAGCC
AUCGACAAUCUGAGAGCGAGCCUAGAAACUACUAAUC
AGGCAAUUGAGGCAAUCAGACAAGCAGGGCAGGAGA
UGAUAUUUGGCUGUUCAGGGUGUCCAAGACUACAUCA
AUAAUGAGCUGAUACCGUCUAUGAAUCAACUAUCUU
GUGAUUUAAUCGGCCAGAAGCUAGGGCUCAAUUGC
UCAGAUACUUAACAGAAAUCCUGUCAUUAUUUGGCC
CAGCUUACGGGACCCCAUAUCUGCGGAGAUUAUCUAUC
CAGGCUUUGAGCUAUGCGCUUGGAGGAGAUUCAAU
AAGGUGUUGGAAAAGCUCGGAUACAGUGGAGGUGAU
CUACUGGGCAUCUUAGAGAGCAGAGGAAUAAAGGCC
GGAUAACUCACGUCGACACAGAGUCCUACUUCAUUGU
ACUCAGUAUAGCCUAUCCGACGCUAUCCGAGAUUAAG
GGGGUGAUUGUCCACCGGCUAGAGGGGGUCUCGUACA
ACAUAGGCUCUCAAGAGUGGUUAUACCACUGUGCCCAA
GUAUGUUGCAACCCAAGGGUACCUUAUCUCGAAUUUU
GAUGAGUCAUCAUGCACUUUCAUGCCAGAGGGGACUG
UGUGCAGCCAGAAUGCCUUGUACCCGAUGAGUCCUCU

GCUCCAACACUCGUAUCCGGGUGCCACUAAGUCCUGU
GCUCGUACACUCGUAUCCGGGUCUUCGGGAACCGGU
UCAUUUUUAUCACAGGGGAACCUAAUAGCCAAUUGUGC
AUCAAUCCUUUGCAAGUGUUACACAACAGGAACAAUC
AUUAAUCAAGACCCUGACAAGAUAUCCUAACAUAUUG
CUGCCGAUCACUGCCCGGUGGUCGAGGUGAAUGGCGU
GACCAUCCAAGUCGGGAGCAGGAGGUAUCCGGACGCU
GUGUACUUGCACAGGAUUGACCUCGGUCCUCCCAUUAU
CUUUGGAGAGGUUGGACGUAGGGACAAAUCUGGGGA
AUGCAAUUGCUAAGUUGGAGGAUGCCAAGGAAUUGU
UGGAGUCAUCGGACCAGAUUUGAGGAGUAUGAAAG
GUUUAUUCGAGCACUAGUAUAGUUUACAUCCUGAUUG
CAGUGUGUCUUGGAGGAUUGAUAGGGAUCCCCGCUU
UAAUAUGUUGCUGCAGGGGGCGUUGUAACAAGAAGG
GAGAACAAGUUGGUAUGUCAAGACCAGGCCUAAAGCC
UGAUCUUACAGGAACAUCAAAAUCCUAUGUAAGGUC ACUCUGA GC_F_MEASLES_
G*GGGAAAUAAGAGAGAGAAAAGAAGAGUAAGAAGAAA 74 D8
UAUAAGAGCCACCAUGGGUCUCAAGGUGAACGUCUCU mRNA Sequence
GUCAUAUUCAUGGCAGUACUGUUAAUCUCUUCAAACAC (assumes T100 tail)
CCACCGGUCAAUCCAUUGGGGCAAUCUCUCUAAGAU Sequence Length:
AGGGGUGGUAGGGGUAGGAAGUGCAAGCUACAAAGU 1925
UAUGACUCGUUCCAGCCAUCAAUCAUUAGUCAUAAAG
UUA AUGCCCAAUAUAACUCUCCUCAACAAUUGCACGA
GGGUAGGGAUUGCAGAAUACAGGAGACUACUGAGAA
CAGUUCUGGAACCAAUUAAGAGAUUGCACUUA AUGCAA
UGACCCAGAAUAUAAGACCGGUUCAGAGUGUAGCUUC
AAGUAGGAGACACAAGAGAUUUGCGGGAGUUGUCCU
GGCAGGUGCGGCCCCUAGGCGUUGCCACAGCUGCUCAA
AUAACAGCCGGUAUUGCACUUCACCAGUCCAUGCUGA
ACUCUCAAGCCAUCGACAAUCUGAGAGCGAGCCUAGA
AACUACUAAUCAGGCAAUUGAGGCAAUCAGACAAGCA
GGGCAGGAGAUGAUUUGGCUGUUCAGGGUGUCCAA
GACUACAUCAAUAAUGAGCUGAUACCGUCUAUGAAUC
AACUAUCUUGUGAUUUAAUCGGCCAGAAGCUAGGGC
UCAAAUUGCUCAGAUACUAUACAGAAAUCCUGUCAUU
AUUUGGCCCCAGCUUACGGGACCCCAUAUCUGCGGAG
AUAUCUAUCCAGGCUUUGAGCUAUGCGCUUGGAGGA
GAUAUCAUAAGGUGUUGGAAAAGCUCGGAUACAGU
GGAGGUGAUCUACUGGGCAUCUUAGAGAGCAGAGGA
AUAAGGCCCCGAUAACUCACGUCGACACAGAGUCCU
ACUUCAUUGUACUCAGUAUAGCCUAUCCGACGCUAUC
CGAGAUUAAGGGGGUGAUUGUCCACCGGCUAGAGGG
GGUCUCGUACAACAUAGGCUCUCAAGAGUGGUUAUACC
ACUGUGCCCAAGUAUGUUGCAACCCAAGGGUACCUUA
UCUCGAAUUUUGAUGAGUCAUCAUGCACUUUCAUGCC
AGAGGGGACUGUGUGCAGCCAGAAUGCCUUGUACCCG
AUGAGUCCUCUGCUCCAAGAAUGCCUCCGGGGGUCCA
CUAAGUCCUGUGCUCGUACACUCGUAUCCGGGUCUUU
CGGGAACCGGUUCAUUUUUAUCACAGGGGAACCUAAUA
GCCAAUUGUGCAUCAAUCCUUUGCAAGUGUUACACAA

CAGGAACCAUAGACCCUGACAAGAUCU
AACAUACAUUGCUGCCGAUCACUGCCCGGUGGUCGAG
GUGAAUGGCGUGACCAUCCAAGUCGGGAGCAGGAGG
UAUCCGGACGCUGUGUACUUGCACAGGAUUGACCUCG
GUCCUCCCAUAUCUUUGGAGAGGUUGGACGUAGGGAC
AAAUUCUGGGGAAUGCAAUUGCUAAGUUGGAGGAUGC
CAAGGAAUUGUUGGAGUCAUCGGACCAGAUAUUGAG
GAGUAUGAAAGGUUUAUCGAGCACUAGUAUAGUUUA
CAUCCUGAUUGCAGUGUGUCUUGGAGGAUUGAUAGG
GAUCCCCGCUUUAUAUGUUGCUGCAGGGGGGCGUUGU
ACAAGAAGGGAGAACAAGUUGGUAUGUCAAGACCA
GGCCUAAAGCCUGAUCUACAGGAACAUCAAAAUCCU
AUGUAAGGUCACUCUGAUGAUAAUAGGCUGGAGCCU
CGGUGGCCAAGCUUCUUGCCCCUUGGGCCUCCCCCA
GCCCCUCCUCCCCUUCUGCACCCGUACCCCCGUGGUC
UUUGAAUAAAGUCUGAGUGGGCGGCAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAUCUAG GC_H_MEASLES_
UCAAGCUUUUGGACCCUCGUACAGAAGCUAAUACGAC 75 B3
UCACUAUAGGGAAAUAAAGAGAGAAAAGAAGAGUAAG Sequence, NT (5'
AAGAAAUUAAGAGCCACCAUGUCACCGCAACGAGAC UTR, ORF, 3'
CGGAUAAAUGCCUUCUACAAAGAUAAACCCUUAUCCCA UTR)
AGGGAAGUAGGAUAGUUAUUAACAGAGAACAUUCUUA Sequence Length:
UGAUUGACAGACCCUAUGUUCUGCUGGCUGUUCUGUU 2065
CGUCAUGUUUCUGAGCUUGAUCGGAUUGCUGGCAAU
UGCAGGCAUUAAGACUUCAUCGGGCAGCCAUCUACACC
GCGGAGAUCCAUAAAAGCCUCAGUACCAAUUCUGGAUG
UGACUAACUCCAUCGAGCAUCAGGUCAAGGACGUGCU
GACACCACUCUUUAAAAUCAUCGGGGAUGAAGUGGGC
CUGAGAACACCUCAGAGAUUCACUGACCUAGUGAAAU
UCAUCUCGGACAAGAUUAAAUCCCUUAAUCCGGAUAG
GGAGUACGACUUCAGAGAUUCACUUGGUGCAUCAA
CCGCCAGAGAGGAUCAAAACUAGAUUAUGAUCAAUACU
GUGCAGAUGUGGCUGCUGAAGAGCUCAUGAAUGCAU
UGGUGAACUCAACUCUACUGGAGACCAGAACAACCAC
UCAGUUCCUAGCUGUCUCAAAGGGAAACUGCUCAGGG
CCCACUACAAUCAGAGGUCAAUUCUCAAACAUGUCGC
UGUCCUUGUUGGACUUGUACUUAAGGUCGAGGUUACA
AUGUGUCAUCUAUAGUCACUAUGACAUCCCAGGGAAU
GUAUGGGGGGAACCUACCUAGUUGAAAAGCCUAAUCU
GAACAGCAAAGGGUCAGAGUUGUCACAACUGAGCAU
GUACCGAGUGUUUGAAGUAGGUGUGAUCAGAAACCC
GGGUUUGGGGGCUCCGGUGUCCAUAUGACAAACUA
UUUUGAGCAACCAGUCAGUAAUGGUCUCGGCAACUGU
AUGGUGGCUUUGGGGGAGCUCAAACUCGCAGCCCUU
GUCACGGGGACGAUUCUAUCAUAAUUCCCUAUCAGGG
AUCAGGGAAAGGUGUCAGCUUCCAGCUCGUCAAGCUG
GGUGUCUGGAAAUCCCCAACCGACAUGCAAUCCUGGG
UCCCCUUAUCAACGGAUGAUCCAGUGGUAGACAGGCU

UUACCUUCUAGGUGUCUACUCGUCUGACAAU
CAAGCAAAAUGGGCUGUCCCCGACAACACGAACAGAUG
ACAAGUUGCGAAUGGAGACAUGCUUCCAGCAGGCGUG
UAAAGGUAAAAUCCAAGCACUCUGCGAGAAUCCCCGAG
UGGGUACCAUUGAAGGAUAACAGGAUUCCUUCAUAC
GGGUUCCUGUCUGUUGAUCUGAGUCUGACGGUUGAG
CUUAAAAUCAAAAUUGCUCGCGGAUUCGGGCCAUUG
AUCACACACGGCUCAGGGAUGGACCUAUACAAAUCCA
ACUGCAACAAUGUGUAUUGGCUGACUUAUCCGCCAAU
GAGAAAUCUAGCCUUAGGCGUAAUCAACACAUUGGA
GUGGAUACCGAGAUUCAAGGUUAGUCCCAACCUUCUUC
ACUGUCCCAAUUAAGGAAGCAGGCGAAGACUGCCAUG
CCCCAACAUACCUACCUGCGGAGGUGGACGGUGAUGU
CAAACUCAGUUCCAACCUGGUGAUUCUACCUGGUCAA
GAUCUCCAAUAUGUUUUGGCAACCUACGAUACCUCCA
GGGUUGAGCAUGCUGUGGUUUAUUACGUUUACAGCC
CAAGCCGCUCAUUUUCUUAUUUAUCCUUUUAGGUU
GCCUAUAAAGGGGGUCCCAAUCGAACUACAAGUGGAA
UGCUCACAUUGGGAUCAAAAACUCUGGUGCCGUCACU
UCUGUGUGCUUGCGGACUCAGAAUCCGGUGGACUUAU
CACUCACUCUGGGAUGGUGGGCAUGGGAGUCAGCUGC
ACAGCUACCCGGGAAGAUGGAACCAUUCGCAGAUAAU
GAUAAUAGGCUGGAGCCUCGGUGGCCAAGCUUCUUGC
CCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGC
ACCCGUACCCCGUGGUCUUUGAAUAAAGUCUGAGUG GGCGGC GC_H_MEASLES_
AUGUCACCGCAACGAGACCGGAUAAAUGCCUUCUACA 76 B3
AAGAUAACCCUUAUCCCAAGGGAAGUAGGAUAGUUA ORF Sequence, NT
UUAACAGAGAACAUUCUUAUGAUUGACAGACCCUAUG
UUCUGCUGGCUGUUCUGUUCGUCAUGUUUCUGAGCUU
GAUCGGAUUGCUGGCAAUUGCAGGCAUUAGACUUCA
UCGGGCAGCCAUCUACACCGCGGAGAUCCAUAAAAGC
CUCAGUACCAAUUCUGGAUGUGACUAAUCUCCAUCGAGC
AUCAGGUCAAGGACGUGCUGACACCACUCUUUAAAAU
CAUCGGGGAUGAAGUGGGCCUGAGAACACCUCAGAGA
UUCACUGACCUAGUGAAAUUCAUCUCGGACAAGAUUA
AAUUCCUUAUCCGGAUAGGGAGUACGACUUCAGAG
AUCUCACUUGGUGCAUCAACCCGCCAGAGAGGAUCAA
ACUAGAUAUUGAUCAAUACUGUGCAGAUGUGGCUGC
UGAAGAGCUCAUGAAUGCAUUGGUGAACUCAACUCU
ACUGGAGACCAGAACAACCACUCAGUUCCUAGCUGUC
UCAAGGGGAAACUGCUCAGGGCCCACUACAAUCAGAG
GUCAAUUCUCAAAACAUUGUCGUGUCCUUGUUGGACUU
GUACUUAGGUCGAGGUUACAAUGUGUCAUCUAUAGU
CACUAUGACAUCCCAGGGAAUGUAUGGGGGGAACCUAC
CUAGUUGAAAAGCCUAAUCUGAACAGCAAAGGGUCA
GAGUUGUCACAACUGAGCAUGUACCGAGUGUUUGAA
GUAGGUGUGAUCAGAAACCCGGGUUUGGGGGCUCCG
GUGUCCAUUAUGACAAACUAUUUUGAGCAACAGUCA
GUAAUGGUCUCGGCAACUGUAUGGUGGCUUUGGGGG
AGCUCAAACUCGCAGCCCUUUGUCACGGGGACGAUUC

UAUCAUAUACAGGGAUCCAGGUAAGGUGU
CAGCUUCCAGCUCGUCAAGCUGGGUGUCUGGAAAUCC
CCAACCGACAUGCAAUCCUGGGUCCCCUUAUCAACGG
AUGAUCCAGUGGUAGACAGGCUUUACCUCUCAUCUCA
CAGAGGUGUCAUCGCUGACAAUCAAGCAAAAUGGGCU
GUCCCGACAACACGAACAGAUGACAAGUUGCGAAUGG
AGACAUGCUUCCAGCAGGCGUGUAAAGGUAAAAUCCA
AGCACUCUGCGAGAAUCCCGAGUGGGUACCAUUGAAG
GAUAACAGGAUUCCUUCAUACGGGGUCCUGUCUGUUG
AUCUGAGUCUGACGGUUGAGCUUAAAAUCAAAAUUG
CUUCGGGAUUCGGGCCAUUGAUCACACACGGCUCAGG
GAUGGACCUAUACAAAUCCAACUGCAACAAUGUGUAU
UGGCUGACUAUUCCGCCAAUGAGAAAUCUAGCCUAG
GCGUAAUCAACACAUUGGAGUGGAUACCGAGAUUCA
AGGUUAGUCCCAACCUCUUCACUGUCCCAAUUAAGGA
AGCAGGCGAAGACUGCCAUGCCCCAACAUACCUACCU
GCGGAGGUGGACGGUGAUGUCAAAACUCAGUUCCAACC
UGGUGAUUCUACCUUGGUCAAGAUCUCCAUAUGUUU
UGGCAACCUACGAUACCUCCAGGGUUGAGCAUGCUGU
GGUUUAUUACGUUUACAGCCCAAGCCGCUCAUUUUCU
UACUUUUAUCCUUUUAGGUUGCCUAUAAAGGGGGUC
CCAAUCGAACUACAAGUGGAAUGCUUCACAUGGGAUC
AAAAACUCUGGUGCCGUCACUUCUGUGUGCUUGCGGA
CUCAGAAUCCGGUGGACUUAUCACUCACUCUGGGAUG
GUGGGCAUGGGGAGUCAGCUGCACAGCUACCCGGGAAG
AUGGAACCAAUCGCAGAUAA GC_H_MEASLES_
G*GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAA 77 B3
UAUAAGAGCCACCAUGUCACCGCAACGAGACCGGAUA mRNA Sequence
AAUGCCUUCUACAAAGAUAAACCCUUAUCCCAAGGGAA (assumes T100
GUAGGAUAGUUAUUAACAGAGAACAUUUUAUGAUUG Tail)
ACAGACCCUAUGUUCUGCUGGCUGUUCUGUUCGUCAU Sequence Length:
GUUUCUGAGCUUGAUUCGGAUUGCUGGCAAUUGCAGG 2126
CAUUAGACUUCAUCGGGCAGCCAUCUACACCGCGGAG
AUCCAUAAAAGCCUCAGUACCAAUCUGGAUGUGACUA
ACUCCAUCGAGCAUCAGGUCAAGGACGUGCUGACACC
ACUCUUUAAAAUCAUCGGGGGAUGAAGUGGGCCUGAG
AACACCUCAGAGAUUCACUGACCUAGUGAAAUUCAUC
UCGGACAAGAUUAAAUUCCUUAUCCGGAUAGGGAG
UACGACUUCAGAGAUUCACUUGGUGCAUCAACCCGC
CAGAGAGGAUCAAAACUAGAUUAUGAUCAAUACUGUG
CAGAUUGGCUGCUGAAGAGCUCAUGAAUGCAUUGG
UGAACUCAACUCUACUGGAGACCAGAACCAACACUCA
GUUCCUAGCUGUCUCAAGGGAAACUGCUCAGGGCCC
ACUACAAUCAGAGGUCAAUUCUCAAAACAUGUCGCUGU
CCUUGUUGGACUUGUACUUAAGGUCGAGGUACAAUG
UGUCAUCUAUAGUCACUAUGACAUCCCGAGGGAUGUA
UGGGGGAACCUACCUAGUUGAAAAGCCUAAUCUGAAC
AGCAAAGGGUCAGAGUUGUCACAACUGAGCAUGUACC
GAGUGUUUGAAGUAGGUGUGAUCAGAAACCCGGGUU
UGGGGGCUCCGGUGUCCAUAUGACAAACUAUUUUG

AGCAAACCAUGGUGUCUGCGGCAACUGGUAUGGUAUGGCUUUGGGGGAGCUCAAACUCGCAGCCCUUUGUCAC
GGGGACGAUUCUAUCAUAAUUCCCUAUCAGGGGAUCAG
GGAAAGGUGUCAGCUUCCAGCUCGUCAAGCUGGGUGU
CUGGAAAUCCCCAACCGACAUGCAAUCCUGGGUCCCC
UUAUCAACGGAUGAUCCAGUGGUAGACAGGCCUUUACC
UCUCAUCUCACAGAGGUGUCAUCGCUGACAAUCAAGC
AAAAUUGGGCUGUCCCGACAACACGAACAGAUGACAAG
UUGCGAAUGGAGACAUGCUUCCAGCAGGCGUGUAAA
GGUAAAAUCCAAGCACUCUGCGAGAAUCCCGAGUGGG
UACCAUUGAAGGAUAACAGGAUUCCUUCAUACGGGG
UCCUGUCUGUUGAUCUGAGUCUGACGGUUGAGCUUA
AAAUCAAAAUUGCUUCGGGAUUCGGGCCAUUGAUCAC
ACACGGCUCAGGGAUGGACCUAUACAAAUCCAACUGC
AACAAUGUGUAUUGGCUGACUAUUCGGCCAAUGAGA
AAUCUAGCCUUAAGGCGUAAUCAACACAUUGGAGUGG
AUACCGAGAUUCAAGGUUAGUCCCAACCUCUUCACUG
UCCCAAUUAAGGAAGCAGGCGAAGACUGCCAUGCCCC
AACAUACCUACCUGCGGAGGUGGACGGUGAUGUCAAA
CUCAGUUCCAACCUGGUGAUUCUACCUGGUCAAGAUC
UCCAAUAUGUUUUGGCAACCUACGAUACCUCCAGGGU
UGAGCAUGCUGUGGUUUUAUACGUUUACAGCCCAAGC
CGCUCAUUUUCUUAUUUUUAUCCUUUUAGGUUGCCUA
UAAAGGGGGUCCCAAUUCGAACUACAAGUGGAAUGCU
UCACAUGGGAUCAAAAACUCUGGUGCCGUCACUUCUG
UGUGCUUGCGGACUCAGAAUCCGGUGGACUUAUCACU
CACUCUGGGAUGGUGGGCAUGGGAGUCAGCUGCACAG
CUACCCGGGAAGAUGGAACCAAUCGCAGAUAAUGAUA
AUAGGCUGGAGCCUCGGUGGCCAAGCUUCUUGCCCCU
UGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACCC
GUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCG
GCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAUCUAG GC_H_MEASLES_
UCAAGCUUUUGGACCCUCGUACAGAAGCUAAUACGAC 78 D8
UCACUAUAGGGAAAUAAGAGAGAAAAGAAGAGUAAG Sequence, NT (5'
AAGAAAUUAAGAGCCACCAUGUCACCACAACGAGAC UTR, ORF, 3'
CGGAUAAAUGCCUUCUACAAAGACAACCCCCAUCCUA UTR)
AGGGAAGUAGGAUAGUUUAUUAACAGAGAACAUCUUA Sequence Length:
UGAUUGAUAGACCUUAUGUUUUGCUGGCUGUUCUUAU 2065
UCGUCAUGUUUCUGAGCUUGAUCGGGUUGCUAGCCAU
UGCAGGCAUUAGACUUCAUCGGGCAGCCAUCUACACC
GCAGAGAUCCAUAAAAGCCUCAGCACCAUUCUGGAUG
UAACUAACUCAUUCGAGCAUCAGGUUAAGGACGUGCU
GACACCACUCUUCAAGAUCAUCGGUGAUGAAGUGGGC
UUGAGGACACCUCAGAGAUUCACUGACCUAGUGAAGU
UCAUCUCUGACAAGAUUAAAUUCCUUAUUCGGACAG
GGAAUACGACUUCAGAGAUUCACUUGGUGUAUCAAC
CCGCCAGAGAGAAUCAAAUUGGAUUAUGAUCAAUAC
UGUGCAGAUUGGCUGCUGAAGAACUCAUGAAUGCA

UUGUGAGACACACAGGGCAACCA
AUCAGUUCCUAGCUGUCUCAAGGGAAACUGCUCAGG
GCCCACUACAAUCAGAGGCCAAUUCUCAAAACAUGUCG
CUGUCCCUGUUGGACUUGUAUUUAAGUCGAGGUUAC
AAUGUGUCAUCUAUAGUCACUAUGACAUCCCAGGGAA
UGUACGGGGGGAACUUACCUAGUGGAAAAGCCUAAUC
UGAGCAGCAAAGGGUCAGAGUUGUCACAACUGAGCA
UGCACCGAGUGUUUGAAGUAGGUGUUAUCAGAAAUC
CGGGUUUGGGGGCUCCGGUUUCCAUAUGACAAACUA
UCUUGAGCAACCAGUCAGUAAUGAUUUCAGCAACUGC
AUGGUGGCUUUGGGGGAGCUCAGUUCGCAGCCCUCU
GUCACAGGGAAGAUUCUAUCACAAUUCCCUAUCAGGG
AUCAGGGAAAGGUGUCAGCUUCCAGCUUGUCAAGCUA
GGUGUCUGGAAAUCCCCAACCGACAUGCAAUCCUGGG
UCCCCCUAUAACGGAUGAUCCAGUGAUAGACAGGCU
UUACCUCUCAUCUCACAGAGGCGUUAUCGCUGACAAU
CAAGCAAAAUGGGCUGUCCCGACAACACGGACAGAUG
ACAAGUUGCGAAUGGAGACAUGC UCCAGCAGGCGUG
UAAGGGUAAA AUCCAAGCACUUUGCGAGAAUCCCGAG
UGGACACCAUUGAAGGAUAACAGGAUUCUUAUACG
GGGUCUUGUCUGUUGAUCUGAGUCUGACAGUUGAGC
UUAAA AUCAAAAUUGUUUCAGGAUUCGGGCCAUUGA
UCACACACGGUUCAGGGAUUGGACCUAUACAAAUCCAA
CCACAACAAUAUGUAUUGGCUGACUAUCCCGCCAAUG
AAGAACCUGGCCUUAGGUGUAAUCAAACACAUUGGAG
UGGAUACCGAGAUUCAAGGUUAGUCCCAACCUCUUA
CUGUUCCAAUUAAGGAAGCAGGCGAGGACUGCCAUGC
CCCAACAUACCUACCUUGCGGAGGUGGAUGGUGAUGUC
AAACUCAGUUCCAAUCUGGUGAUUCUACCUUGGUCAAG
AUCUCCAAUAUGUUCUGGCAACCUACGAUACUUCAG
AGUUGAACAUUGCUGUAGUUUAUUACGUUUACAGCCC
AAGCCGCUCAUUUUCUUACUUUU AUCCUUUUAGGUUG
CCUGUAAGGGGGGUCCCCAUUGAAUUAACAAGUGGAA
UGC UUCACAUGGGACCAAAAACUCUGGUGCCGUCACU
UCUGUGUGCUUGCGGACUCAGAAUCUGGUGGACAU
UCACUCACUCUGGGAUGGUGGGCAUGGGAGUCAGCUG
CACAGCCACUCGGGAAGAUGGAACCAGCCGCAGAUAG
UGAUAAUAGGCUGGAGCCUCGGUGGCCAAGCUUCUUG
CCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUG
CACCCGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGU GGGCGGC GC_H_MEASLES_
AUGUCACCACAACGAGACCGGAUAAAUGCCUUCUACA 79 D8
AAGACAACCCCCAUCCUAAGGGAAGUAGGAUAGUUAU ORF Sequence, NT
UAACAGAGAACAUCUUAUGAUUGAUAGACCUUAUGU
UUUGCUGGCUGUUCUAUUCGUCAUGUUUCUGAGCUU
GAUCGGGUUGC UAGCCAUUGCAGGCAUUAGACUUCAU
CGGGCAGCCAUCUACACCGCAGAGAUCCAUAAAAGCC
UCAGCACCAAUUCUGGAUGUAACUAACUCAAUCGAGCA
UCAGGUUAAGGACGUGCUGACACCACUCUUCAAGAUC
AUCGGUGAUGAAGUGGGCUUGAGGACACCUCAGAGA
UUCACUGACCUAGUGAAGUUCAUCUCUGACAAGAUUA

AAUUCUUAUUCGGAUUAACGACUUCAGAGA
UCUCACUUGGUGUAUCAACCCGCCAGAGAGAAUCAA
UUGGAUUUAUGAUCAAUACUGUGCAGAUUGGCGUCU
GAAGAACUCAUGAAUGCAUUGGUGAACUCAACUCUAC
UGGAGACCAGGGCAACCAAUCAGUUCCUAGCUGUCUC
AAAGGGAAACUGCUCAGGGCCCACUACAAUCAGAGGC
CAAUUCUCAAAACAUGUCGCUGUCCCUGUUGGACUUGU
AUUUAAGUCGAGGUUACAAUGUGUCAUCUAUAGUCA
CUAUGACAUCCCAGGGAAUGUACGGGGGAACUUACCU
AGUGGAAAAGCCUAAUCUGAGCAGCAAAGGGUCAGA
GUUGUCACAACUGAGCAUGCACCGAGUGUUUGAAGU
AGGUGUUUAUCAGAAAUCCGGGUUUGGGGGCUCCGGU
AUUCCAUAUGACAAACUAUCUUGAGCAACCAGUCAGU
AAUGAUUUCAGCAACUGCAUGGUGGCUUUGGGGGAG
CUCAAGUUCGCAGCCCUCUGUCACAGGGAAGAUCUA
UCACAAUUCCCUAUCAGGGAUCAGGGAAGGUGUCAG
CUUCCAGCUUGUCAAGCUAGGUGUCUGGAAAUCCCCA
ACCGACAUGCAAUCCUGGGUCCCCCUAUC AACGGAUG
AUCCAGUGAUAGACAGGCUUUACCUUCUAUCUCACAG
AGGCGUUUAUCGCUGACAAUCAAGCAAAAUGGGCUGUC
CCGACAACACGGACAGAUACAAGUUGCGAAUGGAGA
CAUGCUUCCAGCAGGCGUGUAAGGGUAAAAUCCAAGC
ACUUUGCGAGAAUCCCGAGUGGACACCAUUGAAGGAU
AACAGGAUUCCUUCAUACGGGGUCUUGUCUGUUGAUC
UGAGUCUGACAGUUGAGCUUAAAAUCAAAAUUGUUU
CAGGAUUCGGGCCAUUGAUCACACACGGUUCAGGGAU
GGACCUAUACAAAUCCAACCACAACAAUAUGUAUUGG
CUGACUAUCCCGCCAAUGAAGAACCUGGCCUUAGGUG
UAAUCAACACAUUGGAGUGGAUACCGAGAUUCAAGG
UUAGUCCCAACCUCUUCACUGUUCCAAUUAAGGAAGC
AGGCGAGGACUGCCAUGCCCCAACAUACCUACCUGCG
GAGGUGGAUGGUGAUGUCAAACUCAGUUCCAAUCUG
GUGAUUCUACCUGGUCAAGAUCUCCAAUAUGUUCUGG
CAACCUACGAUACUUC CAGAGUUGAACAU GCUGUAGU
UUAUUACGUUUACAGCCCAAGCCGCUCAUUUUCUUAC
UUUUAUCCUUUUAGGUUGCCUGUAAGGGGGGUCCCCA
UUGAAUUACAAGUGGAAUGCUUCACAUGGGACCAA
AACUCUGGUGCCGUCACUUCUGUGUCUUGCGGACUC
AGAAUCUGGUGGACAUAUCACUCACUCUGGGAUGGU
GGGCAUGGGAGUCAGCUGCACAGCCACUCGGGAAGAU GGAACCAGCCGCAGAUAG
GC_H_MEASLES_ G*GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAA 80 D8
UAUAAGAGCCACCAUGUCACCACAACGAGACCGGAUA mRNA Sequence
AAUGCCUUCUACAAAGACAACCCCCAUCCUAAGGGAA (assumes T100 tail)
GUAGGAUAGUUAUUAACAGAGAACAUUCUUAUGAUUG Sequence Length:
AUAGACCUUAUGUUUUGCUGGCGUGUUCUAUUCGUCA 2126
UGUUUCUGAGCUUGAUCGGGUUGCUAGCCAUUGCAG
GCAUUAGACUUCAUCGGGCAGCCAUCUACACCGCAGA
GAUCCAUAAAAGCCUCAGCACCAAUCUGGAUGUAACU
AACUCAAU CGAGCAUCAGGUUAAGGACGUGCUGACAC
CACUCUUCAAGAUCAUCGGUGAUGAAGUGGGCUUGA

GGACACUCAGACCUAGUGAAGUUCAU
CUCUGACAAGAUUAAAUUCCUUAUCCGGACAGGGAA
UACGACUUCAGAGAUUCACUUGGUGUAUCAACCCGC
CAGAGAGAAUCAAUUGGAUUUAUGAUCAAUACUGUG
CAGAUGUGGCUGCUGAAGAACUCAUGAAUGCAUUGG
UGAACUCAACUCUACUGGAGACCAGGGCAACCAAUCA
GUUCCUAGCUGUCUCAAAGGGAAACUGCUCAGGGCCC
ACUACAAUCAGAGGCCAAUUCUCAAACAUGUCGCUGU
CCCUGUUGGACUUGUAUUUAAGUCGAGGUUACAAUG
UGUCAUCUAUAGUCACUAUGACAUCCCAGGGAAUGUA
CGGGGGAACUUAACCUAGUGGAAAAGCCUAAUCUGAGC
AGCAAAGGGUCAGAGUUGUCACAACUGAGCAUGCACC
GAGUGUUUGAAGUAGGUGUUAUCAGAAAUCCGGGUU
UGGGGGCUCCGGUUAUCCAUAUGACAAACUAUCUUGA
GCAACCAGUCAGUAAUGAUUUCAGCAACUGCAUGGUG
GCUUUGGGGGAGCUCAGUUCGCAGCCCUCUGUCACA
GGGAAGAUCUAUCACAAUUCCCUAUCAGGGAUCAGG
GAAAGGUGUCAGCUUCCAGCUUGUCAAGCUAGGUGUC
UGGAAAUCCCCAACCGACAUGCAAUCCUGGGUCCCCC
UAUCAACGGAUGAUCCAGUGAUAGACAGGCUUUACCU
CUCAUCUCACAGAGGCGUUAUCGCUGACAAUCAAGCA
AAUUGGGCUGUCCCGACAACACGGACAGAUACAAGU
UGCGAAUGGAGACAUGCUCUCCAGCAGGCGUGUAAGG
GUAAAUAUCCAAGCACUUUGCGAGAAUCCCGAGUGGAC
ACCAUUGAAGGAUAACAGGAUUCCUUCAUACGGGGUC
UUGUCUGUUGAUCUGAGUCUGACAGUUGAGCUUAAA
AUCAAAAUUGUUUCAGGAUUCGGGCCAUUGAUCACAC
ACGGUUCAGGGAUGGACCUAUACAAAUCCAACCACAA
CAAUAUGUAUUGGCUGACUAUCCCGCCAAUGAAGAAC
CUGGCCUUAAGGUGUAAUCAACACAUAUGGAGUGGAUA
CCGAGAUUCAAGGUUAGUCCCAACCUCUUCACUGUUC
CAAUUAAGGAAGCAGGCGAGGACUGCCAUGCCCCAAC
AUACCUACCUGCGGAGGUGGAUGGUGAUGUCAAACUC
AGUUCCAAUCUGGUGAUUCUACCUGGUCAAGAUCUCC
AAUAUGUUCUGGCAACCUACGAUACUCCAGAGUUGA
ACAUGCUGUAGUUUAUUACGUUUACAGCCCAAGCCGC
UCAUUUUCUUAUUUUUAUCCUUUUUAGGUUGCCUGUA
AGGGGGGUCCCCAUUGAAUUACAAGUGGAAUGCUUC
ACAUGGGACCAAAAACUCUGGUGCCGUCACUUCUGUG
UGCUGCGGACUCAGAAUCUGGUGGACAUAUCACUCA
CUCUGGGAUGGUGGGCAUGGGAGUCAGCUGCACAGCC
ACUCGGGAAGAUGGAACCAGCCGCAGAUAGUGAUAA
UAGGCUGGAGCCUCGGUGGCCAAGCUUCUUGCCCCU
GGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACCCG
UACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGG
CAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAUCUAG

MGLKVNVSVMFVAVLTLQTPAGQIHWGNLSKIGVV 47 ORF Sequence, AA

GIGSASYKVMTRSSHQSLVIKLMFNITLLNNCTRVEIA
EYRLLRTVLEPIRDALNAMTQNIRPVQSVASSRRHK
RFAGVVLAGAALGVATAAQITAGIALHRSMLNSQAID
NLRASLETTNQAIEAIRQAGQEMILAVQGVQDYINNE
LIPSMNQLSCDLIGQKLGLKLLRYYTEILSLFGPSLRDP
ISAEISIQALSALGGDINKVLEKLGYSGGDLLGILES
GIKARITHVDTESYFIVLSIAYPTLSEIKGVIVHRLEGVS
YNIGSQEWYTTVPKYVATQGYLISNFEDESSCTFMPEG
TVCSQNALYPMSPLLQECLRGSTKSCARTLVSGSFGN
RFILSQGNLIANCASILCKCYTTGTIINQDPDKILTYIAA
DRCPVVEVNGVTIQVGSRRYPDAVYLHRIDLGPISLE
RLDVGTNLGNAIAKLEDAKELLESSDQILRSMKGLSST
SIVYILIAVCLGGLIGIPTLICCCRGRCNKKGEQVGMSR PGLKPDLTGTSSKSYVRSL*

GC_F_MEASLES_D8 MGLKVNVSVMFVAVLTLQTPAGQIHWGNLSKIGVV 48 ORF
Sequence, AA VGSASYKVMTRSSHQSLVIKLMFNITLLNNCTRVGIAE

YRLLRTVLEPIRDALNAMTQNIRPVQSVASSRRHKKR
FAGVVLAGAALGVATAAQITAGIALHQSMLNSQAIDN
LRASLETTNQAIEAIRQAGQEMILAVQGVQDYINNLI
PSMNQLSCDLIGQKLGLKLLRYYTEILSLFGPSLRDPIS
AEISIQALSALGGDINKVLEKLGYSGGDLLGILESRI
KARITHVDTESYFIVLSIAYPTLSEIKGVIVHRLEGVSY
NIGSQEWYTTVPKYVATQGYLISNFEDESSCTFMPEGT
VCSQNALYPMSPLLQECLRGSTKSCARTLVSGSFGNR
FILSQGNLIANCASILCKCYTTGTIINQDPDKILTYIAAD
HCPVVEVNGVTIQVGSRRYPDAVYLHRIDLGPISLER
LDVGTNLGNAIAKLEDAKELLESSDQILRSMKGLSSTS
IVYILIAVCLGGLIGIPALICCCRGRCNKKGEQVGMSR PGLKPDLTGTSSKSYVRSL*

GC_H_MEASLES_B3 MSPQRDRINAFYKDNPPYKGSRVINREHLMIDRPYVL 49 ORF
Sequence, AA LAVLFVMFSLIGLLAIAGIRLHRAAIYTAIEIHKSLSTN

LDVTNSIEHQVKDVLTPLFKIIGDEVGLRTPQRFTDLV
KFISDKIKFLNPDREYDFRDLTWCINPPERIKLDYDQY
CADVAAEELMNALVNSTLLETTRTTTQFLAVSKGNCS
GPTTIRGQFSNMSLSLLDLYLGRGYNVSSIVTMTSQG
MYGGTYLVEKPNLNSKGESELSQLSMYRVFEVGVIRNP
GLGAPVFHMTNYFEQPVSNGLGNCMVALGELKLAAL
CHGDDSIIPYQGSKGVSFQLVKLG VWKSPTDMQSW
VPLSTDDPVVDRLYLSSHGVIADNQAQWAVPTTRT
DDKLRMETCFQQACKGKIQALCENPEWVPLKDNRIPS
YGVLSVDLSLTVELKIKIASGFGPLITHGSGMDLYKSN
CNNVYWLTIPPMRNALGVINTLEWIPRFKVSPNLFTV
PIKEAGEDCHAPTYLPAEVDGDVKLSSNLVILPGQDL
QYVLATYDTSRVEHAVVYVYVSPSRFSYFYFRLPIK
GVPIELQVECFDWDQKLWCRHFCVLADSESGGLITHS GMVGMGVSCATREDGTNRR*

GC_H_MEASLES_D8 MSPQRDRINAFYKDNPPYKGSRVINREHLMIDRPYVL 50 ORF
Sequence, AA LAVLFVMFSLIGLLAIAGIRLHRAAIYTAIEIHKSLSTN

LDVTNSIEHQVKDVLTPLFKIIGDEVGLRTPQRFTDLV
KFISDKIKFLNPDREYDFRDLTWCINPPERIKLDYDQY
CADVAAEELMNALVNSTLLETTRATNQFLAVSKGNCS
GPTTIRGQFSNMSLSLLDLYLSRGYNVSSIVTMTSQGM

YGGTYLVEKPLSSQSLSMHRVFEVGVIRNPG
LGAPVFHMTNYLEQPVSNDFSNCMVALGELKFAALC
HREDSITIPYQGSQKGVSFQLVKLGWKSPTDMQSW
VPLSTDDPVIDRLYLSSHRGVIADNQAKWAVPTTRTD
DKLRMETCFQQACKGKIQALCENPEWTPLKDNRI
PSYGVLSVDLSLTVELKIKIVSGFGPLITHGSGMDLYKSNH
NNMYWLTIPPMKNLALGVINTLEWIPRFKVSPNLFTV
PIKEAGEDCHAPTYLPAEVDGDVKLSSNLVILPGQDL
QYVLATYDTSRVEHAVVYYVYSPSRFSYFYFRLPV
RGVPIELQVECFTWDQKLWCRHFCVLADSESGGHITH
SGMVGMGVSCATREDGTSRR*
(586) TABLE-US-00016 TABLE 15 MeV NCBI Accession Numbers (Amino Acid Sequences)
Type Virus Name GenBank Accession hemagglutinin hemagglutinin [Measles virus strain
Moraten] AAF85673.1 hemagglutinin hemagglutinin [Measles virus strain Rubeovax] AAF85689.1
hemagglutinin hemagglutinin [Measles virus] AAF89824.1 hemagglutinin hemagglutinin protein
[Measles virus] CAA91369.1 hemagglutinin hemagglutinin [Measles virus] BAJ23068.1
hemagglutinin hemagglutinin protein [Measles virus] BAB39848.1 hemagglutinin hemagglutinin
[Measles virus] AAA50551.1 hemagglutinin RecName: Full = Hemagglutinin glycoprotein
P08362.1 hemagglutinin hemagglutinin [Measles virus] AAB63802.1 hemagglutinin hemagglutinin
[Measles virus] AAA56650.1 hemagglutinin hemagglutinin [Measles virus] AAA56642.1
hemagglutinin hemagglutinin [Measles virus] AAA74936.1 hemagglutinin hemagglutinin protein
[Measles virus] BAH56665.1 hemagglutinin hemagglutinin [Measles virus] ACC86105.1
hemagglutinin hemagglutinin [Measles virus strain Edmonston-Zagreb] AAF85697.1
hemagglutinin hemagglutinin [Measles virus] AAR89413.1 hemagglutinin hemagglutinin [Measles
virus] AAA56653.1 hemagglutinin RecName: Full = Hemagglutinin glycoprotein P35971.1
hemagglutinin Hemagglutinin [Measles virus] CAB94916.1 hemagglutinin hemagglutinin
[Measles virus] AAC03036.1 hemagglutinin hemagglutinin [Measles virus] AAF85681.1
hemagglutinin Hemagglutinin [Measles virus] CAB94927.1 hemagglutinin Hemagglutinin
[Measles virus] CAB94925.1 hemagglutinin hemagglutinin protein [Measles virus] BAB39835.1
hemagglutinin Hemagglutinin [Measles virus] CAB94931.1 hemagglutinin hemagglutinin
[Measles virus genotype A] AFO84712.1 hemagglutinin hemagglutinin [Measles virus]
AAA56639.1 hemagglutinin Hemagglutinin [Measles virus] CAB94926.1 hemagglutinin
hemagglutinin protein [Measles virus] BAB39836.1 hemagglutinin Hemagglutinin [Measles virus]
CAB94929.1 hemagglutinin RecName: Full = Hemagglutinin glycoprotein P06830.1
hemagglutinin Hemagglutinin [Measles virus] CAB94928.1 hemagglutinin hemagglutinin protein
[Measles virus] BAB39837.1 hemagglutinin hemagglutinin [Measles virus] AAA74935.1
hemagglutinin hemagglutinin protein [Measles virus] CAB43780.1 hemagglutinin hemagglutinin
[Measles virus] BAA09952.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43815.1
hemagglutinin hemagglutinin [Measles virus] AAF28390.1 hemagglutinin Hemagglutinin [Measles
virus] CAB94923.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43785.1
hemagglutinin hemagglutinin [Measles virus] ABD34001.1 hemagglutinin hemagglutinin protein
[Measles virus] CAB43782.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43781.1
hemagglutinin hemagglutinin [Measles virus] BAH22353.1 hemagglutinin hemagglutinin [Measles
virus] AAC35878.2 hemagglutinin hemagglutinin protein [Measles virus] AAL86996.1
hemagglutinin hemagglutinin [Measles virus] CAA76066.2 hemagglutinin hemagglutinin [Measles
virus] AAA46428.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43803.1
hemagglutinin Hemagglutinin [Measles virus] CAB94918.1 hemagglutinin hemagglutinin
[Measles virus] AAF72162.1 hemagglutinin hemagglutinin [Measles virus] AAM70154.1
hemagglutinin hemagglutinin protein [Measles virus] CAB43776.1 hemagglutinin hemagglutinin
[Measles virus genotype D4] ACT78395.1 hemagglutinin hemagglutinin [Measles virus genotype
D7] AAL02030.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43789.1

hemagglutinin hemagglutinin protein [Measles virus] CAB43774.1 hemagglutinin Hemagglutinin
[Measles virus] CAB94920.1 hemagglutinin Hemagglutinin [Measles virus] CAB94922.1
hemagglutinin hemagglutinin [Measles virus] ABB59491.1 hemagglutinin hemagglutinin protein
[Measles virus] BAB39843.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43804.1
hemagglutinin hemagglutinin [Measles virus] AAX52048.1 hemagglutinin Hemagglutinin
[Measles virus] CAB94930.1 hemagglutinin hemagglutinin [Measles virus] AAA74526.1
hemagglutinin hemagglutinin protein [Measles virus] CAB43814.1 hemagglutinin hemagglutinin
[Measles virus] ABB59493.1 hemagglutinin hemagglutinin [Measles virus genotype D4]
AAL02019.1 hemagglutinin Hemagglutinin [Measles virus] CAB94919.1 hemagglutinin
hemagglutinin protein [Measles virus] AAL86997.1 hemagglutinin hemagglutinin [Measles virus
genotype C2] AAL02017.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43769.1
hemagglutinin hemagglutinin protein [Measles virus] CAB43808.1 hemagglutinin hemagglutinin
[Measles virus] BAO97032.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43805.1
hemagglutinin hemagglutinin protein [Measles virus] CAB43777.1 hemagglutinin hemagglutinin
[Measles virus] AAL67793.1 hemagglutinin hemagglutinin [Measles virus] AAF89816.1
hemagglutinin hemagglutinin [Measles virus genotype D4] AAL02020.1 hemagglutinin
hemagglutinin protein [Measles virus] CAB43786.1 hemagglutinin hemagglutinin protein [Measles
virus strain MVi/New AEP40452.1 Jersey.USA/45.05] hemagglutinin hemagglutinin [Measles
virus] AAA74531.1 hemagglutinin hemagglutinin [Measles virus] AAB63800.1 hemagglutinin
hemagglutinin [Measles virus] AAO21711.1 hemagglutinin hemagglutinin [Measles virus genotype
D8] ALE27189.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43810.1
hemagglutinin hemagglutinin [Measles virus] AAF89817.1 hemagglutinin hemagglutinin [Measles
virus genotype D6] AAL02022.1 hemagglutinin hemagglutinin protein [Measles virus]
CAB43800.1 hemagglutinin hemagglutinin protein [Measles virus genotype B3] AGA17219.1
hemagglutinin hemagglutinin protein [Measles virus] CAB43770.1 hemagglutinin hemagglutinin
protein [Measles virus strain AEP40444.1 MVi/Texas.USA/4.07] hemagglutinin hemagglutinin
[Measles virus] AAX52047.1 hemagglutinin hemagglutinin [Measles virus] AAB63794.1
hemagglutinin hemagglutinin [Measles virus] AAB63796.1 hemagglutinin hemagglutinin [Measles
virus] AAA74528.1 hemagglutinin hemagglutinin [Measles virus] AAB63774.1 hemagglutinin
hemagglutinin [Measles virus] AAB63795.1 hemagglutinin hemagglutinin [Measles virus]
AAA74519.1 hemagglutinin hemagglutinin protein [Measles virus] CAB43778.1 fusion protein
fusion protein [Measles virus strain Moraten] AAF85672.1 fusion protein fusion protein [Measles
virus] AAA56645.1 fusion protein fusion protein [Measles virus strain Rubeovax] AAF85688.1
fusion protein fusion protein [Measles virus] AAF85680.1 fusion protein fusion protein [Measles
virus] AEF30359.1 fusion protein fusion protein [Measles virus] BAA09957.1 fusion protein
fusion protein [Measles virus] AAV84957.1 fusion protein fusion protein [Measles virus MeV-
eGFP_Edm-tag] AII16636.1 fusion protein fusion protein [Measles virus] ABY58018.1 fusion
protein fusion protein [Measles virus] BAA19838.1 fusion protein fusion protein [Measles virus]
AAA56641.1 fusion protein F protein [Measles virus] ABK40529.1 fusion protein fusion protein
[Measles virus] AAA56652.1 fusion protein fusion protein [Measles virus] ABY58017.1 fusion
protein fusion protein [Measles virus] ABB71645.1 fusion protein fusion protein [Measles virus]
NP_056922.1 fusion protein fusion protein [Measles virus strain AIK-C] AAF85664.1 fusion
protein fusion protein [Measles virus] BAB60865.1 fusion protein fusion protein [Measles virus]
BAA09950.1 fusion protein fusion protein [Measles virus strain MVi/New AEP40403.1
York.USA/26.09/3] fusion protein fusion protein [Measles virus] AAA74934.1 fusion protein
fusion protein [Measles virus] CAB38075.1 fusion protein fusion protein [Measles virus strain
AEP40443.1 MVi/Texas.USA/4.07] fusion protein fusion protein [Measles virus] AAF02695.1
fusion protein fusion protein [Measles virus] AAF02696.1 fusion protein fusion protein [Measles
virus] AAT99301.1 fusion protein fusion protein [Measles virus] ABB71661.1 fusion protein fusion
protein [Measles virus] BAK08874.1 fusion protein fusion protein [Measles virus] AAF02697.1

fusion protein fusion protein [Measles virus genotype D4] AFY12704.1 fusion protein fusion protein [Measles virus strain AEP40467.1 MVi/California.USA/16.03] fusion protein fusion protein [Measles virus genotype D8] AHN07989.1 fusion protein fusion protein [Measles virus] AAA46421.1 fusion protein fusion protein [Measles virus] AAA56638.1 fusion protein fusion protein [Measles virus strain AEP40419.1 MVi/Virginia.USA/15.09] fusion protein fusion protein [Measles virus genotype D8] ALE27200.1 fusion protein fusion protein [Measles virus genotype D8] AFY12695.1 fusion protein fusion protein [Measles virus genotype D8] ALE27248.1 fusion protein fusion protein [Measles virus genotype D8] ALE27224.1 fusion protein fusion protein [Measles virus] AAT99300.1 fusion protein fusion protein [Measles virus] BAH96592.1 fusion protein fusion protein [Measles virus strain AEP40459.1 MVi/California.USA/8.04] fusion protein fusion protein [Measles virus genotype D8] AIG94081.1 fusion protein fusion protein [Measles virus] BAA09951.1 fusion protein fusion protein [Measles virus genotype D8] ALE27194.1 fusion protein fusion protein [Measles virus] BAA33871.1 fusion protein fusion protein [Measles virus strain AEP40427.1 MVi/Washington.USA/18.08/1] fusion protein fusion protein [Measles virus] ABY21182.1 fusion protein fusion protein [Measles virus genotype D8] ALE27284.1 fusion protein fusion protein [Measles virus] ACA09725.1 fusion protein fusion protein [Measles virus genotype D8] ALE27314.1 fusion protein fusion protein [Measles virus genotype G3] AFY12712.1 fusion protein fusion protein [Measles virus genotype D8] ALE27368.1 fusion protein RecName: Full = Fusion glycoprotein F0; Contains: P35973.1 RecName: Full = Fusion glycoprotein F2; Contains: RecName: Full = Fusion glycoprotein F1; Flags: Precursor fusion protein fusion protein [Measles virus genotype H1] AIG53713.1 unnamed protein product [Measles virus] CAA34588.1 fusion protein fusion protein [Measles virus] CAA76888.1 fusion protein fusion protein [Measles virus genotype B3.1] AIY55563.1 fusion protein fusion protein [Measles virus] ADO17330.1 fusion protein fusion protein [Measles virus genotype H1] AIG53703.1 fusion protein fusion protein [Measles virus genotype B3] AGA17208.1 fusion protein fusion protein [Measles virus] AAL29688.1 fusion protein fusion protein [Measles virus genotype H1] AIG53706.1 fusion protein fusion protein [Measles virus genotype H1] AIG53701.1 fusion protein fusion protein [Measles virus genotype B3] ALE27092.1 fusion protein fusion protein [Measles virus genotype H1] AIG53714.1 fusion protein fusion protein [Measles virus genotype H1] AIG53694.1 fusion protein fusion protein [Measles virus genotype H1] AIG53668.1 fusion protein fusion protein [Measles virus] ACC86094.1 fusion protein fusion protein [Measles virus genotype H1] AIG53670.1 fusion protein fusion protein [Measles virus genotype H1] AIG53707.1 fusion protein fusion protein [Measles virus genotype B3] AGA17216.1 fusion protein fusion protein [Measles virus genotype H1] AIG53671.1 fusion protein fusion protein [Measles virus strain MVi/New AEP40451.1 Jersey.USA/45.05] fusion protein fusion protein [Measles virus genotype H1] AIG53684.1 fusion protein fusion protein [Measles virus genotype H1] AIG53688.1 fusion protein fusion protein [Measles virus genotype B3] AGA17214.1 fusion protein fusion protein [Measles virus genotype H1] AIG53683.1 fusion protein fusion protein [Measles virus genotype H1] AIG53667.1 fusion protein fusion protein [Measles virus genotype H1] AIG53686.1 fusion protein fusion protein [Measles virus genotype H1] AIG53685.1 fusion protein fusion protein [Measles virus genotype H1] AIG53681.1 unnamed protein product [Measles virus] CAA34589.1 fusion protein fusion protein [Measles virus genotype H1] AIG53678.1 fusion protein fusion protein [Measles virus genotype H1] AIG53710.1 fusion protein fusion protein [Measles virus genotype H1] AIG53669.1 fusion protein fusion protein [Measles virus genotype H1] AIG53664.1 fusion protein fusion protein [Measles virus] AAA50547.1 fusion protein fusion protein [Measles virus genotype H1] AIG53679.1 fusion protein fusion protein [Measles virus genotype H1] AIG53709.1 fusion protein fusion protein [Measles virus genotype H1] AIG53672.1 fusion protein fusion protein [Measles virus genotype H1] AIG53697.1 fusion protein fusion protein [Measles virus genotype H1] AIG53689.1 fusion protein fusion protein [Measles virus genotype H1] AIG53676.1 fusion protein fusion protein [Measles virus genotype H1] AIG53675.1 fusion protein fusion protein [Measles

virus genotype H1] AIG53663.1 fusion protein fusion protein [Measles virus] BAA19841.1 fusion protein fusion protein [Measles virus] AAF02701.1 fusion protein fusion protein [Measles virus genotype H1] AIG53680.1 fusion protein fusion protein [Measles virus genotype H1] AIG53674.1 C protein C protein [Measles virus strain Moraten] AAF85670.1 C protein RecName: Full = Protein C P03424.1 C protein C protein [Measles virus] ACN54404.1 C protein C protein [Measles virus] ACN54412.1 C protein RecName: Full = Protein C P35977.1 C protein C protein [Measles virus] AAF85678.1 C protein C protein [Measles virus] ABD33998.1 C protein unnamed protein product [Measles virus] CAA34586.1 C protein C protein [Measles virus] BAJ51786.1 C protein C protein [Measles virus] BAA33869.1 C protein virulence factor [Measles virus] ABO69700.1 C protein C protein [Measles virus] NP_056920.1 C protein C protein [Measles virus] ADO17333.1 C protein C protein [Measles virus] ACC86082.1 C protein C protein [Measles virus] BAA33875.1 C protein C protein [Measles virus] ABY21189.1 C protein C protein [Measles virus] BAE98296.1 C protein C protein [Measles virus] ADU17782.1 C protein C protein [Measles virus strain MVi/Virginia.USA/15.09] AEP40417.1 C protein C protein [Measles virus] ADU17814.1 C protein C protein [Measles virus] ADU17798.1 C protein C protein [Measles virus genotype D4] AFY12700.1 C protein C protein [Measles virus] ADU17784.1 C protein C protein [Measles virus strain AEP40465.1 MVi/California.USA/16.03] C protein C protein [Measles virus] ABB71643.1 C protein C protein [Measles virus] AEI91027.1 C protein C protein [Measles virus] ADU17874.1 C protein C protein [Measles virus] ADU17903.1 C protein C protein [Measles virus] CAA34579.1 C protein C protein [Measles virus] ADU17790.1 C protein C protein [Measles virus] ADU17800.1 C protein C protein [Measles virus] ABB71667.1 C protein unnamed protein product [Measles virus] CAA34572.1 C protein C protein [Measles virus strain AEP40433.1 MVi/Arizona.USA/11.08/2] C protein C protein [Measles virus] ADU17830.1 C protein C protein [Measles virus] ADU17947.1 C protein C protein [Measles virus] ADU17818.1 C protein C protein [Measles virus strain MVi/New AEP40449.1 Jersey.USA/45.05] C protein C protein [Measles virus strain MVi/Texas.USA/4.07 AEP40441.1 C protein C protein [Measles virus] ADU17864.1 C protein C protein [Measles virus] ADU17838.1 C protein C protein [Measles virus] ADU17881.1 C protein C protein [Measles virus strain AEP40425.1 MVi/Washington.USA/18.08/1] C protein C protein [Measles virus] ADU17927.1 C protein C protein [Measles virus] ADU17953.1 C protein C protein [Measles virus] ADU17889.1 C protein C protein [Measles virus] ADU17963.1 C protein C protein [Measles virus] ADU17893.1 C protein C protein [Measles virus] ADU17820.1 C protein C protein [Measles virus] ABB71651.1 C protein C protein [Measles virus] ADU17786.1 C protein C protein [Measles virus] ADU17862.1 C protein C protein [Measles virus] ADU17923.1 C protein C protein [Measles virus] ADU17959.1 C protein C protein [Measles virus] ADU17951.1 C protein C protein [Measles virus] ADU17916.1 C protein C protein [Measles virus] ADU17957.1 C protein C protein [Measles virus] ADU17925.1 C protein C protein [Measles virus] ADU17901.1 C protein C protein [Measles virus] ADU17887.1 C protein C protein [Measles virus] ADU17832.1 C protein C protein [Measles virus] ADU17891.1 C protein C protein [Measles virus] ADU17961.1 C protein C protein [Measles virus] ADU17872.1 C protein C protein [Measles virus] ADU17929.1 C protein C protein [Measles virus] ADU17908.1 C protein C protein [Measles virus] ADU17910.1 C protein C protein [Measles virus] ADU17921.1 C protein C protein [Measles virus] ADU17824.1 C protein C protein [Measles virus strain AEP40473.1 MVi/Pennsylvania.USA/20.09] C protein C protein [Measles virus] ADU17828.1 C protein C protein [Measles virus] ADU17812.1 C protein C protein [Measles virus genotype D8] AFY12692.1 C protein nonstructural C protein [Measles virus] ABA59559.1 C protein RecName: Full = Protein C Q00794.1 C protein nonstructural C protein [Measles virus] ADO17934.1 C protein nonstructural C protein [Measles virus] ACJ66773.1 C protein C protein [Measles virus genotype G3] AFY12708.1 C protein RecName: Full = Protein C P26035.1 C protein C protein [Measles virus] BAA84128.1 nucleoprotein RecName: Full = Nucleoprotein; AltName: Q77M43.1 Full = Nucleocapsid protein; Short = NP; Short = Protein N nucleoprotein nucleocapsid protein [Measles virus strain Rubeovax] AAF85683.1 nucleoprotein

RecName: Full = Nucleoprotein; AltName: Q89933.1 Full = Nucleocapsid protein; Short = NP; Short = Protein N nucleoprotein nucleocapsid protein [Measles virus strain AIK-C] AAF85659.1 nucleoprotein nucleoprotein [Measles virus] ABI54102.1 nucleoprotein nucleoprotein [Measles virus] AAA56643.1 nucleoprotein nucleoprotein [Measles virus] AAC03050.1 nucleoprotein nucleoprotein [Measles virus] AAA18990.1 nucleoprotein nucleoprotein [Measles virus] AAA56640.1 nucleoprotein RecName: Full = Nucleoprotein; AltName: P35972.1 Full = Nucleocapsid protein; Short = NP; Short = Protein N nucleoprotein RecName: Full = Nucleoprotein; AltName: P10050.1 Full = Nucleocapsid protein; Short = NP; Short = Protein N nucleoprotein N protein [Measles virus] BAB60956.1 nucleoprotein RecName: Full = Nucleoprotein; AltName: B1AAA7.1 Full = Nucleocapsid protein; Short = NP; Short = Protein N nucleoprotein nucleoprotein [Measles virus] AAA18991.1 nucleoprotein nucleoprotein [Measles virus] CAB46894.1 nucleoprotein nucleoprotein [Measles virus] CAB46871.1 nucleoprotein nucleoprotein [Measles virus] CAB46872.1 nucleoprotein nucleoprotein [Measles virus] ABU49606.1 nucleoprotein nucleocapsid protein [Measles virus] AAA75494.1 nucleoprotein nucleoprotein [Measles virus] CAB46883.1 nucleoprotein nucleoprotein [Measles virus] CAB46892.1 nucleoprotein unnamed protein product [Measles virus] CAA34584.1 nucleoprotein nucleoprotein [Measles virus] AAA18997.1 nucleoprotein nucleoprotein [Measles virus] CAB46863.1 nucleoprotein nucleoprotein [Measles virus] AEF30352.1 nucleoprotein nucleoprotein [Measles virus] ABI54103.1 nucleoprotein nucleocapsid protein [Measles virus] AAA46433.1 nucleoprotein nucleoprotein [Measles virus] CAB46902.1 nucleoprotein nucleoprotein [Measles virus] CAB46873.1 nucleoprotein nucleoprotein [Measles virus] CAB46906.1 nucleoprotein nucleoprotein [Measles virus] AAA74547.1 nucleoprotein nucleoprotein [Measles virus] AAA74537.1 nucleoprotein nucleoprotein [Measles virus] CAB46862.1 nucleoprotein nucleocapsid protein [Measles virus] BAA09961.1 nucleoprotein nucleoprotein [Measles virus] AAO15875.1 nucleoprotein nucleoprotein [Measles virus] AAO15871.1 nucleoprotein nucleoprotein [Measles virus] CAB46882.1 nucleoprotein nucleoprotein [Measles virus] CAB60124.1 nucleoprotein nucleoprotein [Measles virus] ABI54104.1 nucleoprotein nucleoprotein [Measles virus] CAB46869.1 nucleoprotein nucleoprotein [Measles virus] CAB46880.1 nucleoprotein nucleoprotein [Measles virus] AAA74541.1 nucleoprotein nucleocapsid protein [Measles virus strain MVi/New AEP40446.1 Jersey.USA/45.05] nucleoprotein nucleoprotein [Measles virus] ABI54110.1 nucleoprotein nucleoprotein [Measles virus] CAB46903.1 nucleoprotein nucleoprotein [Measles virus] CAB46899.1 nucleoprotein nucleoprotein [Measles virus] CAB46901.1 nucleoprotein nucleocapsid protein [Measles virus] ABB71640.1 nucleoprotein nucleoprotein [Measles virus] CAB60113.1 nucleoprotein nucleoprotein [Measles virus] CAB60114.1 nucleoprotein nucleoprotein [Measles virus] CAB60116.1 nucleoprotein nucleoprotein [Measles virus] CAB46895.1 nucleoprotein nucleoprotein [Measles virus] CAB60121.1 nucleoprotein nucleoprotein [Measles virus] ABI54111.1 nucleoprotein nucleoprotein [Measles virus] CAB46889.1 nucleoprotein nucleoprotein [Measles virus] CAB46898.1 nucleoprotein nucleoprotein [Measles virus genotype B3] ALE27083.1 nucleoprotein nucleoprotein [Measles virus] CAB60118.1 nucleoprotein nucleocapsid protein [Measles virus] CAA34570.1 nucleoprotein nucleoprotein [Measles virus] AAC29443.1 nucleoprotein nucleocapsid protein [Measles virus strain AEP40422.1 MVi/Washington.USA/18.08/1] nucleoprotein nucleoprotein [Measles virus] AAO15872.1 nucleoprotein nucleoprotein [Measles virus] CAB46874.1 nucleoprotein nucleoprotein [Measles virus] AAA74550.1 nucleoprotein nucleocapsid protein [Measles virus] ABB71648.1 nucleoprotein nucleoprotein [Measles virus] CAB46900.1 nucleoprotein nucleoprotein [Measles virus] BAH22440.1 nucleoprotein nucleocapsid protein [Measles virus] AAA46432.1 nucleoprotein nucleocapsid protein [Measles virus] BAA33867.1 nucleoprotein nucleoprotein [Measles virus] AAA74539.1 nucleoprotein nucleoprotein [Measles virus] CAB60115.1 nucleoprotein nucleoprotein [Measles virus] CAB60123.1 nucleoprotein

nucleocapsid protein [Measles virus] ABB71664.1 nucleoprotein nucleoprotein [Measles virus]
 CAB60125.1 nucleoprotein nucleoprotein [Measles virus] AAA74546.1 nucleoprotein
 nucleoprotein [Measles virus] CAB46886.1 nucleoprotein nucleoprotein [Measles virus]
 BAH22350.1 nucleoprotein nucleoprotein [Measles virus] CAB46867.1 nucleoprotein
 nucleocapsid protein [Measles virus] BAA09954.1 nucleoprotein nucleoprotein [Measles virus]
 AAO15873.1 nucleoprotein nucleocapsid protein [Measles virus] AEP95735.1 nucleoprotein
 nucleoprotein [Measles virus] AAL37726.1 nucleoprotein nucleoprotein [Measles virus]
 AAA74549.1 nucleoprotein RecName: Full = Nucleoprotein; AltName: P26030.1 Full =
 Nucleocapsid protein; Short = NP; Short = Protein N nucleoprotein nucleoprotein [Measles virus
 ETH55/99] AAK07777.1 nucleoprotein nucleoprotein [Measles virus genotype B3] AGA17238.1
 nucleoprotein nucleoprotein [Measles virus] AEF30351.1 nucleoprotein nucleoprotein [Measles
 virus genotype B3] AGA17242.1 nucleoprotein nucleoprotein [Measles virus ETH54/98]
 AAK07776.1 nucleoprotein nucleoprotein [Measles virus] AAA74548.1 nucleoprotein
 nucleoprotein [Measles virus] AAA19221.1 nucleoprotein nucleoprotein [Measles virus]
 AAC03039.1 nucleoprotein nucleoprotein [Measles virus] AAA19223.1 nucleoprotein
 nucleoprotein [Measles virus genotype B3] AGA17241.1 nucleoprotein nucleoprotein [Measles
 virus] CAB60122.1 nucleoprotein nucleoprotein [Measles virus] CAC34599.1 nucleoprotein
 nucleoprotein [Measles virus] AAC03042.1 nucleoprotein nucleoprotein [Measles virus]
 CAC34604.1 nucleoprotein nucleoprotein [Measles virus] AAA74544.1 nucleoprotein
 nucleocapsid protein [Measles virus] NP_056918.1 V Protein RecName: Full = Non-structural
 protein V Q9IC37.1 V Protein RecName: Full = Non-structural protein V Q9EMA9.1 V Protein V
 protein [Measles virus] ACN54411.1 V Protein V protein [Measles virus] ACN54403.1 V Protein V
 protein [Measles virus] AEP95742.1 V Protein V protein [Measles virus strain
 MVi/Virginia.USA/15.09] AEP40416.1 V Protein V protein [Measles virus] ADU17801.1 V
 Protein V protein [Measles virus] ADU17849.1 V Protein V protein [Measles virus] ABB71642.1 V
 Protein V protein [Measles virus genotype D8] AFY12693.1 V Protein V protein [Measles virus]
 YP_003873249.2 V Protein V protein [Measles virus strain AEP40432.1
 MVi/Arizona.USA/11.08/2] V Protein RecName: Full = Non-structural protein V P26036.1 V
 Protein V protein [Measles virus strain AEP40464.1 MVi/California.USA/16.03] V Protein V
 protein [Measles virus strain AEP40456.1 MVi/California.USA/8.04] V Protein V protein [Measles
 virus] ABY21188.1 V Protein V protein [Measles virus strain AEP40424.1
 MVi/Washington.USA/18.08/1] V Protein V protein [Measles virus] BAH96581.1 V Protein V
 protein [Measles virus] ABB71666.1 V Protein RecName: Full = Non-structural protein V
 P60168.1 V Protein V protein [Measles virus] BAH96589.1 V Protein V protein [Measles virus]
 ADU17954.1 V Protein V protein [Measles virus strain MVi/New AEP40400.1 York.USA/26.09/3]
 V Protein V protein [Measles virus] ABY21196.1 V Protein virulence factor [Measles virus]
 ABO69701.1 V Protein V protein [Measles virus] ABB71650.1 V Protein V protein [Measles virus]
 ACC86086.1 V Protein V protein [Measles virus genotype D4] AFY12702.1 V Protein V protein
 [Measles virus strain MVi/New AEP40448.1 Jersey.USA/45.05] V Protein V protein [Measles
 virus] BAE98295.1 V Protein V protein [Measles virus] ACC86083.1 V Protein V protein [Measles
 virus] ACU57139.1 V Protein V protein [Measles virus] ADO17334.1 V Protein V protein
 [Measles virus] ADU17930.1 V Protein V protein [Measles virus genotype G3] AFY12710.1 V
 Protein V protein [Measles virus strain AEP40472.1 MVi/Pennsylvania.USA/20.09] V Protein
 phosphoprotein [Measles virus] ADU17839.1 V Protein V protein [Measles virus] ADU17894.1 V
 Protein V protein [Measles virus] ACN50010.1 V Protein V protein [Measles virus] ADU17892.1
 unnamed protein product [Measles virus] CAA34585.1 V Protein V protein [Measles virus]
 ABD33997.1

(587) TABLE-US-00017 TABLE 16 SEQ ID Name Sequence NO: Flagellin Nucleic
 Acid Sequences NT (5'

TCAAGCTTTTGGACCCTCGTACAGAAGCTAATACGACTCACTAT 51 UTR, ORF,

AGGGAAATAAAGAGAGAGAAAAGAGAGTAAGAAGATAAGAAAGATAG 3' UTR)
AGCCACCATGGCACAAGTCATTAATACAAACAGCCTGTCGCTG
TTGACCCAGAATAACCTGAACAAATCCCAGTCCGCACTGGGCA
CTGCTATCGAGCGTTTGTCTTCCGGTCTGCGTATCAACAGCGCG
AAAGACGATGCGGCAGGACAGGCGATTGCTAACCGTTTTACCG
CGAACATCAAAGGTCTGACTCAGGCTTCCCGTAACGCTAACGA
CGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACGAA
ATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCAGT
CTGCGAATGGTACTAACTCCCAGTCTGACCTCGACTCCATCCAG
GCTGAAATCACCCAGCGCCTGAACGAAATCGACCGTGTATCCG
GCCAGACTCAGTTCAACGGCGTGAAAGTCCTGGCGCAGGACAA
CACCTTGACCATCCAGGTTGGTGCCAACGACGGTGAAACTATC
GATATTGATTTAAAAGAAATCAGCTCTAAAACACTGGGACTTG
ATAAGCTTAATGTCCAAGATGCCTACACCCCGAAAGAACTGC
TGTAACCGTTGATAAACTACCTATAAAAATGGTACAGATCCT
ATTACAGCCCAGAGCAATACTGATATCCAAACTGCAATTGGCG
GTGGTGCAACGGGGGTTACTGGGGCTGATATCAAATTTAAAGA
TGGTCAATACTATTTAGATGTTAAAGGCGGTGCTTCTGCTGGTG
TTTATAAAGCCACTTATGATGAAACTACAAAGAAAGTTAATAT
TGATACGACTGATAAACTCCGTTGGCAACTGCGGAAGCTACA
GCTATTCGGGGAACGGCCACTATAACCCACAACCAAATTGCTG
AAGTAACAAAAGAGGGTGTTGATACGACCACAGTTGCGGCTCA
ACTTGCTGCAGCAGGGGTTACTGGCGCCGATAAGGACAATACT
AGCCTTGTAAACTATCGTTTGAGGATAAAAACGGTAAGGTTA
TTGATGGTGGCTATGCAGTGAAAATGGGCGACGATTTCTATGC
CGCTACATATGATGAGAAAACAGGTGCAATTACTGCTAAAACC
ACTACTTATACAGATGGTACTGGCGTTGCTCAAACCTGGAGCTGT
GAAATTTGGTGGCGCAAATGGTAAATCTGAAGTTGTTACTGCT
ACCGATGGTAAGACTTACTTAGCAAGCGACCTTGACAAACATA
ACTTCAGAACAGGCGGTGAGCTTAAAGAGGTTAATACAGATAA
GACTGAAAACCCACTGCAGAAAATTGATGCTGCCTTGGCACAG
GTTGATACACTTCGTTCTGACCTGGGTGCGGTTTCAGAACCGTTT
CAACTCCGCTATCACCAACCTGGGCAATACCGTAAATAACCTG
TCTTCTGCCCCGTAGCCGTATCGAAGATTCCGACTACGCAACCGA
AGTCTCCAACATGTCTCGCGCGCAGATTCTGCAGCAGGCCGGT
ACCTCCGTTCTGGCGCAGGCGAACCAGGTTCCGCAAAACGTCC
TCTCTTTACTGCGTTGATAATAGGCTGGAGCCTCGGTGGCCATG
CTTCTTGCCCCCTTGGGCCTCCCCCAGCCCCCTCCTCCCCCTTCCTG
CACCCGTACCCCCGTGGTCTTTGAATAAAGTCTGAGTGGGCGG C ORF
ATGGCACAAGTCATTAATACAAACAGCCTGTGCTGTTGACCC 52 Sequence,
AGAATAACCTGAACAAATCCCAGTCCGCACTGGGCACTGCTAT NT
CGAGCGTTTGTCTTCCGGTCTGCGTATCAACAGCGCGAAAGAC
GATGCGGCAGGACAGGCGATTGCTAACCGTTTTACCGCGAACA
TCAAAGGTCTGACTCAGGCTTCCCGTAACGCTAACGACGGTAT
CTCCATTGCGCAGACCACTGAAGGCGCGCTGAACGAAATCAAC
AACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCAGTCTGCGA
ATGGTACTAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAA
ATCACCCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGA
CTCAGTTCAACGGCGTGAAAGTCCTGGCGCAGGACAACACCCT
GACCATCCAGGTTGGTGCCAACGACGGTGAAACTATCGATATT

GATTTAAACCTCTATAACAACTGGGACTTGATAAGC
TTAATGTCCAAGATGCCTACACCCCGAAAGAACTGCTGTAAC
CGTTGATAAACTACCTATAAAAATGGTACAGATCCTATTACA
GCCCAGAGCAATACTGATATCCAACTGCAATTGGCGGTGGTG
CAACGGGGGTACTGGGGCTGATATCAAATTTAAAGATGGTCA
ATACTATTTAGATGTTAAAGGCGGTGCTTCTGCTGGTGTTTATA
AAGCCACTTATGATGAACTACAAAGAAAGTTAATATTGATAC
GACTGATAAACTCCGTTGGCAACTGCGGAAGCTACAGCTATT
CGGGGAACGGCCACTATAACCCACAACCAAATTGCTGAAGTAA
CAAAAGAGGGTGTGATACGACCACAGTTGCGGCTCAACTTGC
TGCAGCAGGGGTACTGGCGCCGATAAGGACAATACTAGCCTT
GTAAAACTATCGTTTGAGGATAAAAACGGTAAGGTTATTGATG
GTGGCTATGCAGTGAAAATGGGCGACGATTTCTATGCCGCTAC
ATATGATGAGAAAACAGGTGCAATTACTGCTAAAACCACTACT
TATACAGATGGTACTGGCGTTGCTCAAACCTGGAGCTGTGAAAT
TTGGTGGCGCAAATGGTAAATCTGAAGTTGTTACTGCTACCGAT
GGTAAGACTTACTTAGCAAGCGACCTTGACAAACATAACTTCA
GAACAGGCGGTGAGCTTAAAGAGGTTAATACAGATAAGACTG
AAAACCCACTGCAGAAAATTGATGCTGCCTTGGCACAGGTTGA
TACTTTCGTTCTGACCTGGGTGCGGTTTCAGAACCGTTTCAACT
CCGCTATCACCAACCTGGGCAATACCGTAAATAACCTGTCTTCT
GCCCGTAGCCGTATCGAAGATTCCGACTACGCAACCGAAGTCT
CCAACATGTCTCGCGCGCAGATTCTGCAGCAGGCCGGTACCTC
CGTTCTGGCGCAGGCGAACCAGGTTCCGCAAAACGTCCTCTCTT TACTGCGT mRNA
G*GGGAAAUAAAGAGAGAAAAGAAGAGUAAAGAAGAAAUAAUAA 53 Sequence
GAGCCACCAUGGCACAAGUCAUUAUACAAACAGCCUGUCGC (assumes
UGUUGACCCAGAAUAACCUGAACAAAUCCCAGUCCGCACUGG T100 tail)
GCACUGCUAUCGAGCGUUUGUCUUCCGGUCUGCGUAUCAACA
GCGCGAAAGACGAUGCGGCAGGACAGGCGAUUGCUAACCGUU
UUACCGCGAACAUCAAAGGUCUGACUCAGGCUUCCCGUAACG
CUAACGACGGUAUCUCCAUUGCGCAGACCACUGAAGGCGCGC
UGAACGAAAUCAACAACAACCUGCAGCGUGUGCGUGAACUGG
CGGUUCAGUCUGCGAAUGGUACUAAUCCCAGUCUGACCUCG
ACUCCAUCCAGGCUGAAAUCACCCAGCGCCUGAACGAAAUUG
ACCGUGUAUCCGGCCAGACUCAGUUCAACGGCGUGAAAGUCC
UGGCGCAGGACAACACCCUGACCAUCCAGGUUGGUGCCAACG
ACGGUGAAACUAUCGAUAUUGAUUUAAAAGAAAUCAGCUCU
AAAACACUGGGACUUGAUAAAGCUUAAUGUCCAAGAUGCCUAC
ACCCCGAAAGAAACUGCUGUAACCGUUGAUAAAACUACCUAU
AAAAAUGGUACAGAUCCUAUUACAGCCCAGAGCAAUACUGAU
AUCCAAACUGCAAUUGGCGGUGGUGCAACGGGGGUUACUGG
GGCUGAUUAUCAAUUUAAAAGAUUGGUCAAUACUAAUUAGAUG
UUAAAGGCGGUGCUUCUGCUGGUGUUUAAUAAAGCCACUUAU
GAUGAAACUACAAAGAAAGUUAAUUAUUGAUACGACUGAUAA
AACUCCGUUGGCAACUGCGGAAGCUACAGCUAUUCGGGGAAC
GGCCACUAUAACCCACAACCAAUUGCUGAAGUAACAAAAGA
GGGUGUUGAUACGACCACAGUUGCGGCUCAACUUGCUGCAGC
AGGGGUUACUGGCGCCGAUAAGGACAAUACUAGCCUUGUAA
AACUAUCGUUUGAGGAUAAAAACGGUAAGGUUAUUGAUGGU
GGCUAUGCAGUGAAAAUGGGCGACGAUUUCUAUGCCGCUACA

[illegible]

AUGGCACAAGUCAUUAUACAAACAGCCUGUCGCUGUUGACC 82 Sequence,
CAGAAUAACCUGAACAAAUCCCAGUCCGCACUGGGGCACUGCU NT
AUCGAGCGUUUGUCUUCCGGUCUGCGUAUCAACAGCGCGAAA
GACGAUGCGGCAGGACAGGCGAUUGCUAACCGUUUUACCGCG
AACAUCAAAGGUCUGACUCAGGCUUCCCGUAACGCUAACGAC
GGUAUCUCCAUUGCGCAGACCACUGAAGGCGCGCUGAACGAA
AUCAACAACAACCUGCAGCGUGUGCGUGAACUGGCGGUUCAG
UCUGCGAAUGGUACUAACUCCCAGUCUGACCUCGACUCCAUC
CAGGCUGAAAUCACCCAGCGCCUGAACGAAAUCGACCGUGUA
UCCGGCCAGACUCAGUUCAACGGCGUGAAAGUCCUGGCGCAG
GACAACACCCUGACCAUCCAGGUUGGUGCCAACGACGGUGAA
ACUAUCGAUAUUGAUUUAAAAGAAAUCAGCUCUAAAACACU
GGGACUUGAUAAAGCUUAAUGUCCAAGAUGCCUACACCCCGAA
AGAAACUGCUGUAACCGUUGAUAAAACUACCUAUAAAAAUG
GUACAGAUCCUAUUCAGCCCAGAGCAAUACUGAUAUCCAAA
CUGCAAUUGGCGGUGGUGCAACGGGGGUUACUGGGGGCUGAU
AUCAAUUUUAAAGAUGGUCAAUACUAUUUAGAUGUUAAAGG
CGGUGCUUCUGCUGGUGUUUAUAAAGCCACUUAUGAUGAAA
CUACAAAGAAAGUUAUAUUGAUACGACUGAUAAAACUCCG
UUGGCAACUGCGGAAGCUACAGCUAUUCGGGGAACGGCCACU
AUAACCCACAACCAAUUGCUGAAGUAACAAAAGAGGGUGU
UGAUACGACCACAGUUGCGGCUCAACUUGCUGCAGCAGGGGU
UACUGGCGCCGAUAAGGACAAUACUAGCCUUGUAAAACUAUC
GUUUGAGGAUAAAAACGGUAAGGUUAUUGAUGGUGGCUAUG
CAGUGAAAAUGGGCGACGAUUUCUAUGCCGCUACAUAUGAU
GAGAAAACAGGUGCAAUACUGCUAAAACCACUACUUAUACA
GAUGGUACUGGCGUUGCUCAAACUGGAGCUGUGAAAUUUGG
UGGCGCAAUUGGUAAAUCUGAAGUUGUUAACUGCUACCGAUG
GUAAGACUUAACUAGCAAGCGACCUUGACAAACAUAACUUCA
GAACAGGCGGUGAGCUUAAAGAGGUUAAUACAGAUAAAGACU
GAAAACCCACUGCAGAAAAUUGAUGCUGCCUUGGCACAGGUU
GAUACACUUCGUUCUGACCUGGGUGCGGUUCAGAACCGUUUC
AACUCCGCUAUCACCAACCUGGGCAAUACCGUAAAUAACCG
UCUUCUGCCCGUAGCCGUAU CGAAGAUUCCGACUACGCAACC
GAAGUCUCCAACAUGUCUCGCGCGCAGAUUCUGCAGCAGGCC
GGUACCUC CGUUCUGGCGCAGGCGAACCAGGUUCCGCAAAC
GUCCUCUCUUUACUGCGU mRNA

G*GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAUAUAA 83 Sequence
GAGCCACCAUGGCACAAGUCAUUAUACAAACAGCCUGUCGC (assumes
UGUUGACCCAGAAUAACCUGAACAAAUCCCAGUCCGCACUGG T100 tail)
GCACUGCUAUCGAGCGUUUGUCUUCCGGUCUGCGUAUCAACA

GCGCGAAGGCGGAUUGCUAACCGUU
UUACCGCGAACAUCAAAGGUCUGACUCAGGCUUCCCGUAACG
CUAACGACGGUAUCUCCAUUGCGCAGACCACUGAAGGCGCGC
UGAACGAAAUCAACAACAACCUUGCAGCGUGUGCGUGAACUGG
CGGUUCAGUCUGCGAAUGGUACUAAACUCCCAGUCUGACCUCG
ACUCCAUCCAGGCUGAAAUCACCCAGCGCCUGAACGAAAUCG
ACCGUGUAUCCGGCCAGACUCAGUUCAACGGCGUGAAAGUCC
UGGCGCAGGACAACACCCUGACCAUCCAGGUUGGUGCCAACG
ACGGUGAAACUAUCGAUAUUGAUUUAAAAGAAAUCAGCUCU
AAAACACUGGGACUUGAUAAAGCUUAAUGUCCAAGAUGCCUAC
ACCCCGAAAGAAACUGCUGUAACCGUUGAUAAAACUACCUAU
AAAAAUGGUACAGAUCCUAUUACAGCCCAGAGCAAUACUGAU
AUCCAAACUGCAAUUGGCGGUGGUGCAACGGGGGUUACUGG
GGCUGAUUAUCAAUUUAAAAGAUGGUCAAUACUAAUUAGAUG
UAAAAGGCGGUGCUUCUGCUGGUGUUUAUAAAGCCACUUAU
GAUGAAACUACAAAGAAAGUUAAUAUUGAUACGACUGAUAA
AACUCCGUUGGCAACUGCGGAAGCUACAGCUAUUCGGGGAAC
GGCCACUAUAACCCACAACCAAUUGCUGAAGUAACAAAAGA
GGGUGUUGAUACGACCACAGUUGCGGCUCAACUUGCUGCAGC
AGGGGUUACUGGCGCCGAUAAGGACAAUACUAGCCUUGUAA
AACUAUCGUUUGAGGAUAAAAACGGUAAGGUUAUUGAUGGU
GGCUAUGCAGUGAAAAUGGGCGACGAUUUCUAUGCCGCUACA
UAUGAUGAGAAAACAGGUGCAAUUACUGCUAAAACCACUAC
UUAUACAGAUUGGUACUGGCGUUGCUCAAACUGGAGCUGUGA
AAUUUGGUGGCGCAAUUGGUAAAUCUGAAGUUGUUACUGCU
ACCGAUGGUAAAGACUUAUUAGCAAGCGACCUUGACAAACAU
AACUUCAGAACAGGCGGUGAGCUUAAAGAGGUUAAUACAGA
UAAGACUGAAAAACCCACUGCAGAAAAUUGAUGCUGCCUUGGC
ACAGGUUGAUACACUUCGUUCUGACCUGGGUGCGGUUCAGAA
CCGUUUAACUCCGCUAUCACCAACCUGGGCAAUACCGUAAA
UAACCUGUCUUCUGCCCGUAGCCGUUUCGAAGAUUCCGACUA
CGCAACCGAAGUCUCCAACAUGUCUCGCGCGCAGAUUCUGCA
GCAGGCCGGUACCUCGCUUCUGGCGCAGGCGAACCAGGUUCC
GCAAACGUCCUCUCUUUACUGCGUUGAUAAUAGGCUGGAGC
CUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCCCCCCAGCC
CCUCCUCCCCUUCUGCACCCGUACCCCGUGGUCUUUGAAU
AAAGUCUGAGUGGGCGGCAAAAAAAAAAAAAAAAAAAAAAAAAA
AA
AAAUCUAG

(588) TABLE-US-00018 TABLE 17 SEQ ID Name Sequence NO: Flagellin Amino Acid
Sequences ORF MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSAKDDEA 54
Sequence, GQAIANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQRV AA
RELAVQSANGTNSQSDLDISIQAEITQRLNEIDRVSGQTQFNGVKVL
AQDNTLTIQVGANDGETIDIDLKEISSKTLGLDKLNVQDAYTPKET
AVTVDKTTYKNGTDPITAQSNTDIQTAIGGGATGVTGADIKFKDG
QYYLDVKGGASAGVYKATYDETTKKVNIDTTDKTPLATAEATAI
RGTATITHNQIAEVTKEGVDTTVAQAALAAAGVTGADKDNTSLV
KLSFEDKNGKVIDGGYAVKMGDDFYAATYDEKTGAITAKTTTTYT
DGTGVAQTGAVKFGGANGKSEVVTATDGKTYLASDLDKHNFRT
GGELKEVNTDKTENPLQKIDAALAQVDTLRSDLGAVQNRFNSAIT

DNAGITPAISLDLMTDAELARAVPNMPTSAQIKLMLENRAMVRRKGFIL
CGVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWY
CQNAGSTVYYPNEKDCETRGRDHFCDTAAGINVAEQSKECNINISTTNYP
KVSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIIKQLNKGCSYITNQ
DADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEHQWHVALDQVFE
NIENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTG
APPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 87
DM_Krarup_ DVENLTCSDGPSLIKTELDLLKSALRELKTVSADQLAREEQIENPGSGSFVLG
T74LD185P AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIPDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 88
TM_Krarup_ DVENLTCSDGPSLIKTELDLLKSALRELKTVSADQLAREEQIENPGSGSFVLG
T74LD185PD454N
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIPDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPENNQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 89
4M_Krarup_ DVENLTCSDGPSLIKTELDLLKSALRELKTVSADQLAREEQIENPGSGSFVLG
T74LS170LD185P
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVLKNLTRAINKNKCDIPDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 90
5M_Krarup_ DVENLTCSDGPSLIKTELDLLKSALRELKTVSADQLAREEQIENPGSGSFVLG
T74LS170LD185PD454N
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVLKNLTRAINKNKCDIPDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHFCDTAAGINVAEQSKECNINISTTNYPCK

VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPENQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVL GSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTL_PVG 91
DM_Krarup_ DVENLTCSDGPSLIKTEL DL_KSALRELKTVSADQLAREEQIENPGSGSFVLG
E51PT74L AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETR GDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVL GSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTL_PVG 92
TM_Krarup_
DVENLTCSDGPSLIKTEL DL_KSALRELKTVSADQLAREEQIENPGSGSFVLG
E51PT74LD454N
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETR GDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPENQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVL GSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_SC_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV G 93
StabilizeAlpha_
DVENLTCSDGPSLIKTEL DL_KSALRELKTVSADQLAREEQIENPGSGSFVLG T74L
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETR GDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVL GSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV G 94 SC_
DL ENLTCSDGPSLIKTEL DTKSALRELKTVSADQLAREEQIENPGSGSFVLG
StabilizeAlpha_ AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
V55L LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFIL
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETR GDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVL GSSMILVSIFIIKKTKKPTGA

PPELSGVTNNGFIPHN HMPV_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 95 SC_
DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPSGSGSFVLG
StabilizeAlpha_ AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
S170L LATAVRELKDFVLKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGILI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVLGGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 96 SC_
DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPSGSGSFVLG
StabilizeAlpha_ AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
T174W LATAVRELKDFVSKNLWRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGILI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVLGGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_ SC_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 97 4M_
DLENLTCSDGPSLIKTELDLKSALRELKTVSADQLAREEQIENPSGSGSFVLG
StabilizeAlpha_ AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
V55LT74LS170LT174W
LATAVRELKDFVLKNLWRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGILI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVLGGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLPVG 98
ProlineStab_ DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPSGSGSFVLG
E51P AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGILI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILI AVLGGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 99
ProlineStab_ DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPSGSGSFVLG
D185P AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIPDLKMAVSFSQFNRRFLNVVRQFS

DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGLI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNYACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV G 100
ProlineStab_ DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPGSGSFVLG
D183P AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCPIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGLI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNYACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV G 101
ProlineStab_ DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPGSGSFVLG
E131P AIALGVAAAAAVTAGVAIAKTIRLPSEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGLI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNYACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV G 102
ProlineStab_ DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPGSGSFVLG
D447P AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGLI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNYACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFPPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV G 103
TrimerRepulsionD454N

DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPGSGSFVLG
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGLI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNYACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIKGRPVSSSFDPIKFPENQFQVALDQVFENI

ENSQALVDQSNRILSSAEKGNTGFIIVIIILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 104
TrimerRepulsionE453N
DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPGSGSFVLG
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQFNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGILI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIGRVPVSSSFDPIKFPQDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVIIILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN HMPV_
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVFTLEV 105
StabilizeAlphaF196W

DVENLTCSDGPSLIKTELDTKSALRELKTVSADQLAREEQIENPGSGSFVLG
AIALGVAAAAAVTAGVAIAKTIRLESEVTAINNALKKTNEAVSTLGNGVRV
LATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVSFSQWNNRRFLNVVRQFS
DNAGITPAISLDLMTDAELARAVPNMPTSAGQIKLMLENRAMVRRKGFGILI
GVYGSSVIYMVQLPIFGVIDTPCWIVKAAPSCSEKKGNACLLREDQGWYC
QNAGSTVYYPNEKDCETRGRDHVFCDTAAGINVAEQSKECNINISTTNYPCK
VSTGRHPISMVALSPLGALVACYKGVSCSIGSNRVGIKQLNKGCSYITNQD
ADTVTIDNTVYQLSKVEGEQHVIGRVPVSSSFDPIKFPEDQFQVALDQVFENI
ENSQALVDQSNRILSSAEKGNTGFIIVIIILIAVLGSSMILVSIFIIKKTKKPTGA
PPELSGVTNNGFIPHN

(590) TABLE-US-00020 TABLE 19 SEQ ID Strain Nucleic Acid Sequence NO: Human
Metapneumovirus Mutant Nucleic Acid Sequences HMPV_SC_

ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 106 DSCAV1_4MMV
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCTGCAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CTTTGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGGCCCTGAACAAGAACAAGTGCAGCATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGTGTGGCGTGACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG

TAGCAGAGAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGGATCAGTTCAACGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAAC HMPV_SC_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 107 DSTRIC_4MMV
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCTGCAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGTGTGGCGTGACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC

[illegible]

CAATGGCTTCATCCCTACAAC HMPV_SC_TM_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 109 Krarup_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT T74LD185PD454N
CCTGCAGCACCATCACAGAGGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG
CCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAGC
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAACC
TGACACGGGGCCATTAACAAGAACAAGTGCGACATCCCTGA
CCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCGG
TTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGAA
TCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTGA
GCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGGC
CAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGAC
GGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGCAG
CGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATCG
ACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTAG
CGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGGA
CCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTAC
TACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGACCAC
GTGTTCTGTGATAACGCCGCTGGAATCAATGTGGCCGAGC
AGAGCAAAGAGTGCAACATCAACATCAGCACCACT
ATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATACCAAC
CAGGACGCCGATAACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGAACCAGTTCCAGGTGGCCCTGGACCAGGTGTTGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTACAAC HMPV_SC_4M_

ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 110 Krarup_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT T74LS170LD185P
CCTGCAGCACCATCACAGAGGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG

CCATTGCTCTTGGAGTGGAGTGGCTGCTGCTGAGCTGTTACAGC
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGCTTAAGAACC
TGACACGGGCCATTAACAAGAACAAGTGCGACATCCCTGA
CCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCGG
TTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGAA
TCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTGA
GCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGGC
CAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGAC
GGAAAGGCTTCGGCATTCTGATTGGCGTGACGGCAGCAG
CGTGATCTATATGGTGACGCTGCCTATCTTCGGCGTGATCG
ACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTAG
CGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGGA
CCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTAC
TACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGACCAC
GTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCGAGC
AGAGCAAAGAGTGCAACATCAACATCAGCACCACTCAACT
ATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATAACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAC HMPV_SC_5M_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 111 Krarup_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT T74LS170LD185PD454N
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACC GGCTGGTACACCAACGTGTTACACTGGAAGTGGGC
GACGTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG
CCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAGC
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGCTTAAGAACC
TGACACGGGCCATTAACAAGAACAAGTGCGACATCCCTGA
CCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCGG
TTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGAA
TCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTGA

GCTTGGCTAGCCATGACATCTGCCGGC
CAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGAC
GGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGCAG
CGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATCG
ACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTAG
CGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGGA
CCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTAC
TACCCTAACGAGAAGGACTGCGAGACAAGAGGCGACCAC
GTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCGAGC
AGAGCAAAGAGTGCAACATCAACATCAGCACCACT
ATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATACCAAC
CAGGACGCCGATACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGAACCAGTTCCAGGTGGCCCTGGACCAGGTGTTGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAAC HMPV_SC_DM_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 112 Krarup_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT E51PT74L
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGCCTGTGGGC
GACGTGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG
CCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAGC
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAACC
TGACACGGGGCCATTAACAAGAACAAGTGCGACATCGACG
ACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGCA
GCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGACCA
CGTGTCTGTGATACCGCCGCTGGAATCAATGTGGCCGAG

CAGACGAAGAATCAACATCAGCACCAACCAAC
TATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAC HMPV_SC_TM_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 113 Krarup_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT E51PT74LD454N
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGCCTGTGGGC
GACGTGAGAAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG
CCATTGCTCTTGAGTGGCTGCTGCTGCAGCTGTTACAGC
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAACC
TGACACGGGGCCATTAACAAGAACAAGTGCGACATCGACG
ACCTGAAGATGGCCGTGTCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGACGGCAGCA
GCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGACCA
CGTGTCTGTGATACCGCCGCTGGAATCAATGTGGCCGAG
CAGAGCAAAGAGTGCAACATCAACATCAGCACCAACCAAC
TATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGAACCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG

AACATCATGACCAATCTGGTGGACCAAGTCCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAC HMPV_SC_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 114 StabilizeAlpha_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT T74L
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG
CCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAGC
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAACC
TGACACGGGGCCATTAACAAGAACAAGTGCGACATCGACG
ACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGACGGCAGCA
GCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGACCA
CGTGTCTGTGATACCGCCGCTGGAATCAATGTGGCCGAG
CAGAGCAAAGAGTGCAACATCAACATCAGCACCAACCAAC
TATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAC HMPV_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 115 SC_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT StabilizeAlpha_
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG V55L

AACCGGCTGGGCTGGTTCACACTGGAAGTGGGGC
GACCTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATAACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAC HMPV_SC_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 116 StabilizeAlpha_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT S170L
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC

CACAGCGCTGCGAGCTTGAAGGACTTCGTGCTTAAGAAC
CTGACACGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTT
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAC HMPV_SC_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 117 StabilizeAlpha_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT T174W
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTGAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGTGGCGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA

TCGACACACCATTCGATTGTAAGGCCGCTCCTCGATG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGAC
CACGTGTTCTGTGATAACGCCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATAACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CTGACACGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGAC
CACGTGTTCTGTGATAACGCCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATAACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAAC HMPV_SC_4M_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 118 StabilizeAlpha_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT V55LT74LS170LT174W
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACCTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGCTCAAGAGCGCCCTGAGAGA
ACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGAA
CAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGAG

CCATTGCTCTTGAGTGGAGTGGCTGCTGCTGCAGCTGTTACACG
AGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTGAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGCTTAAGAACC
TGTGGCGGGCCATTAACAAGAACAAGTGCGACATCGACG
ACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGACGGCAGCA
GCGTGATCTATATGGTGACGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGACCA
CGTGTTCTGTGATAACGCCGCTGGAATCAATGTGGCCGAG
CAGAGCAAAGAGTGCAACATCAACATCAGCACCAACCAAC
TATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATAACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAC HMPV_ProlineStab_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 119 E51P
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACACTGCCTGTGGGC
GACGTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTGAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT

GAGCTGGCTAGCGCTAGCCTAACATGGCTTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATAACGCCGCTGGAATCATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTTGG
GCATCATCAAGCAGCTGAACAAGGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATACCGTGACCATCGACAACACCGTGG
TATCAGCTGAGCAAGGTGGAAGGCCGAACAGCACGTGATC
AAGGGCAGACCRGRTGTCCAGCAGCTTCGACCCTATCAAGT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAAC HMPV_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 120 ProlineStab_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT D185P
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAACTCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTGAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGCCATTAACAAGAACAAGTGCGACATCCCTG
ACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGCA
GCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGACCA
CGTGTTCTGTGATAACGCCGCTGGAATCAATGTGGCCGAG
CAGAGCAAAGAGTGCAACATCAACATCAGCACCACCAAC

TATCCCTGCAAGGTGGCCAGGCACCCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATAACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAAC HMPV_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 121 ProlineStab_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT D183P
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTGAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGCCATTAACAAGAACAAGTGCCCTATCGACG
ACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGCA
GCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGACCA
CGTGTTCCTGTGATAACCGCCGCTGGAATCAATGTGGCCGAG
CAGAGCAAAGAGTGCAACATCAACATCAGCACCAACCAAC
TATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATAACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA

GAATGTCCTGCTGATGAGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAAC HMPV_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 122 ProlineStab_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT E131P
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGCCTAGCGA
AGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACGA
GGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGCC
ACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAACC
TGACACGGGGCCATTAACAAGAACAAGTGCGACATCGACG
ACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGCG
GTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGGA
ATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCTG
AGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCGG
CCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCGA
CGGAAAGGCTTCGGCATTCTGATTGGCGTGACGGCAGCA
GCGTGATCTATATGGTGACGCTGCCTATCTTCGGCGTGATC
GACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTGTA
GCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGAGG
ACCAAGGCTGGTATTGTGTCAGAACGCCGGCAGCACCGTGTA
CTACCCTAACGAGAAGGACTGCGAGACAAGAGGGCGACCA
CGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCGAG
CAGAGCAAAGAGTGCAACATCAACATCAGCACCAACCAAC
TATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTCTAT
GGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTTATA
AGGGCGTGTCCTGTAGCATCGGCAGCAACAGAGTGGGCAT
CATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAAC
CAGGACGCCGATAACCGTGACCATCGACAACACCGTGTATC
AGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATCAAGG
GCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGTTCCC
TGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTTTCGAG
AACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCCAACA
GAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGCTTCAT
CATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTCCATG
ATCCTGGTGTCCATCTTCATCATTATCAAGAAGACCAAGA
AGCCACCGGCGCTCCTCCAGAACTGAGCGGAGTGACCAA
CAATGGCTTCATCCCTCACAAAC HMPV_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 123 ProlineStab_
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT D447P
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC

GACGTCGATCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCCG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCCCACCTATCAAGT
TCCCTGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAC HMPV_

ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 124 TrimerRepulsionD454N

CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTCGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC

CTGACGACGGGCAAGCAAGTGGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGAACCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAC HMPV_
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 125 TrimerRepulsionE453N
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTTCAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG

TAGCAGAGAAGGCTGGCTTACGCCTGGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGATACCGTGACCATCGACAACACCGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTCAGGATCAGTTCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAC HMPV_Stabilize
ATGAGCTGGAAGGTGGTCATCATCTTCAGCCTGCTGATCA 126 AlphaF196W
CACCTCAGCACGGCCTGAAAGAGAGCTACCTGGAAGAGT
CCTGCAGCACCATCACAGAGGGCTACCTGTCTGTGCTGAG
AACCGGCTGGTACACCAACGTGTTCACTGGAAGTGGGC
GACGTGAGAAATCTGACATGCTCTGATGGCCCTAGCCTGA
TCAAGACCGAGCTGGATCTGACCAAGAGCGCCCTGAGAG
AACTCAAGACCGTGTCTGCCGATCAGCTGGCCAGAGAGGA
ACAGATCGAGAATCCTGGCAGCGGCAGCTTTGTGCTGGGA
GCCATTGCTCTTGGAGTGGCTGCTGCTGCAGCTGTTACAG
CAGGCGTGGCCATCGCTAAGACCATCAGACTGGAAAGCG
AAGTGACCGCCATCAACAACGCCCTGAAGAAGACAAACG
AGGCCGTCAGCACACTCGGCAATGGCGTTAGAGTGCTGGC
CACAGCCGTGCGCGAGCTGAAGGACTTCGTGTCCAAGAAC
CTGACACGGGGCCATTAACAAGAACAAGTGCGACATCGAC
GACCTGAAGATGGCCGTGTCCTTTAGCCAGTGGAACCGGC
GGTTTCTGAACGTCGTGCGGCAGTTTAGCGACAACGCCGG
AATCACACCAGCCATCAGCCTGGACCTGATGACAGATGCT
GAGCTGGCTAGAGCCGTGCCTAACATGCCTACATCTGCCG
GCCAGATCAAGCTGATGCTCGAGAATAGAGCCATGGTCCG
ACGGAAAGGCTTCGGCATTCTGATTGGCGTGTACGGCAGC
AGCGTGATCTATATGGTGCAGCTGCCTATCTTCGGCGTGA
TCGACACACCCTGCTGGATTGTGAAGGCCGCTCCTAGCTG
TAGCGAGAAGAAGGGCAATTACGCCTGCCTGCTGAGAGA
GGACCAAGGCTGGTATTGTCAGAACGCCGGCAGCACCGTG
TACTACCCTAACGAGAAGGACTGCGAGACAAGAGGCGAC
CACGTGTTCTGTGATACCGCCGCTGGAATCAATGTGGCCG
AGCAGAGCAAAGAGTGCAACATCAACATCAGCACCACCA
ACTATCCCTGCAAGGTGTCCACCGGCAGGCACCCTATTTC
TATGGTGGCTCTGTCTCCTCTGGGAGCCCTGGTGGCTTGTT
ATAAGGGCGTGTCTGTAGCATCGGCAGCAACAGAGTGG
GCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC

CAACCAAGGACGACGATCGACCATCGACCAACACCGGTG
TATCAGCTGAGCAAGGTGGAAGGCGAACAGCACGTGATC
AAGGGCAGACCTGTGTCCAGCAGCTTCGACCCTATCAAGT
TCCCTGAGGATCAGTTCCAGGTGGCCCTGGACCAGGTGTT
CGAGAACATCGAGAATTCCCAGGCTCTGGTGGACCAGTCC
AACAGAATCCTGTCTAGCGCCGAGAAGGGAAACACCGGC
TTCATCATCGTGATCATCCTGATCGCCGTGCTGGGCAGCTC
CATGATCCTGGTGTCCATCTTCATCATTATCAAGAAGACC
AAGAAGCCCACCGGCGCTCCTCCAGAACTGAGCGGAGTG
ACCAACAATGGCTTCATCCCTCACAAC Human Metapneumovirus mRNA Sequences
HMPV_SC_AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 127
DSCAV1_4MMV CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA
GUCCUGCAGCACCAUCACAGAGGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUACAGCAGGCGUGGCCAUUCUGCAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCUUUGCCGUGCGGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCUGAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUCUGAGCUGGCUAGAGCCGUGCCUAC
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GUGUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGCGACCACGUGUUCUGUG
AUACCGCCGCUUGGAAUCAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUCUAUUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGGAUCAGUUCAACGUGGCCCUUGGACCAGGUGUUCG
AGAACAUUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGA

CCAAGACCAAGCCUCCAGAACGUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 128 DSURIC_4MMV
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUCACAGCAGGCGUGGCCAUCUGCAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCUAA
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUCU
GUGUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAAACGA
GAAGGACUGCGAGACAAGAGGGCACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACCUAUUCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCAGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCC
UGAGCACCAUGUGGCAUGUGGCCCUUGGACCAGGUGUUCGA
GAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCAA
CAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCUU
CAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCUC
CAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGAC
CAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAGU
GACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_DM_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 129 Krarup_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA U74LD185P
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUG
CCUGAUCAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU

GAGAGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCACAUCCCUGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCUAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUCCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_UM_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 130 Krarup_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA U74LD185PD454N
GUCCUGCAGCACCAUCACAGAGGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG

UGCGACAUCGGAUAGGCGGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCUAAAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACCUAUUCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGAACCAGUUCAGGUGGCCCUUGGACCAGGUGUUCGA
GAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCAA
CAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCUU
CAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCUC
CAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGAC
CAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAGU
GACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_4M_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 131 Krarup_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA U74LS170LD185P
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGCUIAAGAACCUGACACGGGCCAUUAACAAGAACAA
GUGCGACAUCCCUGACCUGAAGAUGGCCGUGUCCUUUAG
CCAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUU
UAGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGA
CCUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCUAA
CAUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGA
GAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUC
UGAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUG
CAGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGG

AUUGUAGCGCGAGAAAGAAGGG
CAAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUCAGGUGGCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_5M_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 132 Krarup_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA U74LS170LD185PD454N
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGCUIAAGAACCUGACACGGGCCAUUAACAAGAACA
GUGCGACAUCCCUGACCUGAAGAUGGCCGUGUCCUUUAG
CCAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUU
UAGCGACAACGCCGGAUCACACCAGCCAUCAGCCUGGA
CCUGAUGACAGAUUCUGAGCUGGCUAGAGCCGUGCCUAA
CAUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGA
GAAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUC
UGAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUG
CAGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGG
AUUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGG
CAAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC

GUGUCUGUACGACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGAACCAGUUCCAGGUGGCCCUGGACCAGGUGUUCGA
GAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCAA
CAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCUU
CAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCUC
CAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGAC
CAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAGU
GACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_DM_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 133 Krarup_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA E51PU74L
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGCCUGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUG
CCUGAUAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCU AAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCU AACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGGAUCAGUUCCAGGUGGCCCUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU

UCAUCAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_UM_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 134 Krarup_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA E51PU74LD454N
GUCCUGCAGCACCAUCACAGAGGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGCCUGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUG
CCUGAUCAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAU CGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCUAAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCAGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGAACCAGUUCCAGGUGGCCCUUGGACCAGGUGUUCGA
GAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCAA
CAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCUU
CAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCUC
CAUGAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGAC
CAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAGU
GACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 135 SUabilizeAlpha_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA U74L
GUCCUGCAGCACCAUCACAGAGGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU

GGGCGGACGUCUGCAUUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUGCUGAGCUGGCCUAGAGCCGUGCCUAAAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGGCGAACAGCACGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 136 SC_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA SUabilizeAlpha_
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU V55L
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACCUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU

AGAGUGCUGCGCGAGCUGGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCUAAAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCAGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUCCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGUCCUCCAGAACUGAGCGGAG
UGACCAACAUAUGGCUUCAUCCCUCACAAC HMPV_SC_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 137 SUabilizeAlpha_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA S170L
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUAACAGCAGGCGUGGCCAUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGCUUAAGAACCUGACACGGGCCAUUAACAAGAACA
GUGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAG
CCAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUU
UAGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGA
CCUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCUAA
CAUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGA
GAAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUC

UGAGUUGGCGUGAGCGUGAUCUAUAUGGUG
CAGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGG
AUUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGG
CAAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SC_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 138 SUabilizeAlpha_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA U174W
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGUGGCGGGCCAUAACAAGAACAA
GUGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAG
CCAGUUAACCGGCGGUUUCUGAACGUCGUGCGGCAGUU
UAGCGACAACGCCGGAUCACACCAGCCAUCAGCCUGGA
CCUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCUAA
CAUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGA
GAAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUC
UGAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUG
CAGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGG
AUUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGG
CAAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA

AGGUGGUCUAGCCUACUAGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGGAUCAGUUCCAGGUGGCCCUGGACCAGGUGUUCG
AGAAACAUUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUAUCCUCACAAC HMPV_SC_4M_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 139 SUabilizeAlpha_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA V55LU74LS170LU174W
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACCUCGAGAAUCUGACAUGCUCUGAUGGCCCUG
CCUGAUAAGACCGAGCUGGAUCUGCUCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUAACAGCAGGCGUGGCCAUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGCUUAAGAACCUGUGGGCGGGCCAUAUAACAAGAACA
GUGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAG
CCAGUUAACCGGCGGUUUCUGAACGUCGUGCGGCAGUU
UAGCGACAACGCCGGAUCACACCAGCCAUCAGCCUGGA
CCUGAUGACAGAUUCUGAGCUGGCUAGAGCCGUGCCUAA
CAUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGA
GAAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUC
UGAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUG
CAGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGG
AUUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGG
CAAUUAACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGGAUCAGUUCCAGGUGGCCCUGGACCAGGUGUUCG

AGAACAUCGAGGCUUCAGGUGGACCGAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 140 ProlineSUab_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA E51P
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGCCUGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAAUACACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCUAC
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCGAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 141 ProlineSUab_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA D185P

GUCCUGCAGGCUACGAGCCUACCUUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCCCUGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCU AAC
AUGCCUACAUCUGCCGGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUC CAGGUGGCCCUUGGACCAGGUGUUCG
AGAAUCAUCGAGAAU UCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGGAAACACCGGCU
UCAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 142 ProlineSUab_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA D183P
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUCGCUAAGACCAUCAGA

CUGGAAAGCGAAGCAACACGCCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCCCUAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCU AAC
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCU AACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUC CAGGUGGCCCUGGACCAGGUGUUCG
AGAACAUCGAGAAU UCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGA
CCAAGAAGCCCACCGGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 143 ProlineSUab_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA E131P
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAU CGCUAAGACCAUCAGA
CUGCCUAGCGAAGUGACCGCCAUCAACAACGCCCU GAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAUUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCU AAC

AUGCCUAAGCUAAGCUAUGCCUCCGAG
AAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUCCCC
UGAGGAUCAGUUC CAGGUGGCCUGGACCAGGUGUUCG
AGAACAUUCGAGAAU UCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUAUCAUCUUCAGCCUGCUGAU 144 ProlineSUab_
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA D447P
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAU CGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGC GACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAAUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAU GCUGAGCUGGCUAGAGCCGUGCCU AAC
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGGCGACCACGUGUUCUGUG

AUACCGCGGAGCAUUGGCGGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG
CAGACCUGUGUCCAGCAGCUUCCCACCUAUAAGUCCC
UGAGGAUCAGUUCCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUAUCAUAUAUCAAGAAGA
CCAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 145 UrimerRepulsionD454N
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAU CGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAAUACACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCUAC
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCCUGCUGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCUAACGA
GAAGGACUGCGAGACAAGAGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCUUGUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUAAGGG

CAGAGCUGGACCCUAGUCAAGUUC
UGAGAACCAGUUCCAGGUGGCCUGGACCAGGUGUUCGA
GAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCAA
CAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCUU
CAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCUC
CAUGAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGAC
CAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAGU
GACCAACAAUGGCUUCAUCCCUCACAAC HMPV_
AUGAGCUGGAAGGUGGUCAUCAUCUUCAGCCUGCUGAU 146 UrimerRepulsionE453N
CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUGCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUAG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUACAGCAGGCGUGGCCAU CGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUUCAACCGGCGGUUUCUGAACGUCGUGCGGCAGUUU
AGCGACAACGCCGGAAUCACACCAGCCAUCAGCCUGGAC
CUGAUGACAGAUUGCUGAGCUGGCUAGAGCCGUGCCU AAC
AUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGAG
AAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUCU
GAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUGC
AGCUGCCUAUCUUCGGCGUGAUCGACACACCUCUGGGA
UUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGGC
AAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGCGACCACGUGUUCUGUG
AUACCGCCGCUGGAAUCA AUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCUUGGUGGCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUC
UCAGGAUCAGUUCCAGGUGGCCUGGACCAGGUGUUCGA
GAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCAA
CAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCUU
CAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCUC
CAUGAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGAC
CAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAGU
GACCAACAAUGGCUUCAUCCCUCACAAC HMPV_SUabilize

CACACCUCAGCACGGCCUGAAAGAGAGCUACCUGGAAGA
GUCCUGCAGCACCAUCACAGAGGGCUACCUGUCUGUCU
GAGAACCGGCUGGUACACCAACGUGUUCACACUGGAAGU
GGGCGACGUCGAGAAUCUGACAUGCUCUGAUGGCCCUG
CCUGAUCAAGACCGAGCUGGAUCUGACCAAGAGCGCCCU
GAGAGAACUCAAGACCGUGUCUGCCGAUCAGCUGGCCAG
AGAGGAACAGAUUCGAGAAUCCUGGCAGCGGCAGCUUUG
UGCUGGGAGCCAUUGCUCUUGGAGUGGCUGCUGCUGCA
GCUGUUACAGCAGGCGUGGCCAUUCGCUAAGACCAUCAGA
CUGGAAAGCGAAGUGACCGCCAUCAACAACGCCCUGAAG
AAGACAAACGAGGCCGUCAGCACACUCGGCAAUGGCGUU
AGAGUGCUGGCCACAGCCGUGCGCGAGCUGAAGGACUUC
GUGUCCAAGAACCUGACACGGGCCAUUAACAAGAACAAG
UGCGACAUCGACGACCUGAAGAUGGCCGUGUCCUUUAGC
CAGUGGAACCGGGCGGUUUCUGAACGUCGUGCGGCAGUU
UAGCGACAACGCCGGAAUCACACCAGCCAUCAGCCUGGA
CCUGAUGACAGAUGCUGAGCUGGCUAGAGCCGUGCCUAA
CAUGCCUACAUCUGCCGGCCAGAUCAAGCUGAUGCUCGA
GAAUAGAGCCAUGGUCCGACGGAAAGGCUUCGGCAUUC
UGAUUGGCGUGUACGGCAGCAGCGUGAUCUAUAUGGUG
CAGCUGCCUAUCUUCGGCGUGAUCGACACACCUGCUGG
AUUGUGAAGGCCGCUCCUAGCUGUAGCGAGAAGAAGGG
CAAUUACGCCUGCCUGCUGAGAGAGGACCAAGGCUGGUA
UUGUCAGAACGCCGGCAGCACCGUGUACUACCCUAACGA
GAAGGACUGCGAGACAAGAGGCGACCACGUGUUCUGUG
AUACCGCCGCUUGGAAUCAAUUGUGGCCGAGCAGAGCAAAG
AGUGCAACAUCAACAUCAGCACCAACUAUCCCUGCA
AGGUGUCCACCGGCAGGCACCCUAUUUCUAUGGUGGCUC
UGUCUCCUCUGGGAGCCCUGGUGGCCUUGUUUAUAAGGGC
GUGUCCUGUAGCAUCGGCAGCAACAGAGUGGGCAUCAUC
AAGCAGCUGAACAAGGGCUGCAGCUACAUCACCAACCCAG
GACGCCGAUACCGUGACCAUCGACAACACCGUGUAUCAG
CUGAGCAAGGUGGAAGGCGAACAGCACGUGAUCAAGGG
CAGACCUGUGUCCAGCAGCUUCGACCCUAUCAAGUUCCC
UGAGGAUCAGUUCAGGUGGCCCUUGGACCAGGUGUUCG
AGAACAUCGAGAAUUCCCAGGCUCUGGUGGACCAGUCCA
ACAGAAUCCUGUCUAGCGCCGAGAAGGGAAACACCGGCU
UCAUCAUCGUGAUCAUCCUGAUCGCCGUGCUGGGCAGCU
CCAUGAUCCUGGUGUCCAUCUUCAUCAUUAUCAAGAAGA
CCAAGAAGCCCACCGGCGCUCCUCCAGAACUGAGCGGAG
UGACCAACAAUGGCUUCAUCCCUCACAAC

EQUIVALENTS

(591) Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the disclosure described herein. Such equivalents are intended to be encompassed by the following claims.

(592) All references, including patent documents, disclosed herein are incorporated by reference in their entirety.

Claims

1. A method of vaccinating a human subject against a betacoronavirus, the method comprising administering to the human subject a dose of a vaccine comprising a messenger ribonucleic acid (mRNA) formulated in a lipid nanoparticle comprising 20-60 mol % ionizable cationic lipid, 5-25 mol % neutral lipid, 25-55 mol % cholesterol, and 0.5-15 mol % PEG-modified lipid, wherein the mRNA comprises an open reading frame encoding a full-length betacoronavirus spike protein, wherein the open reading frame comprises nucleosides consisting of N1-methylpseudouridine, adenosine, guanosine, and cytidine, wherein the dose comprises from 20-50 µg of the mRNA and is administered intramuscularly.
 2. The method of claim 1, wherein the dose comprises 20 µg of the mRNA.
 3. The method of claim 1, wherein the dose comprises 25 µg of the mRNA.
 4. The method of claim 1, wherein the dose comprises 30 µg of the mRNA.
 5. The method of claim 1, wherein the dose comprises 50 µg of the mRNA.
 6. A method of vaccinating a human subject against a betacoronavirus, the method comprising administering to the human subject a dose of a vaccine comprising a messenger ribonucleic acid (mRNA) formulated in a lipid nanoparticle comprising 20-60 mol % ionizable cationic lipid, 5-25 mol % neutral lipid, 25-55 mol % cholesterol, and 0.5-15 mol % PEG-modified lipid, wherein the mRNA comprises an open reading frame encoding a full-length betacoronavirus spike protein, wherein the open reading frame comprises nucleosides consisting of N1-methyl-pseudouridine, adenosine, guanosine, and cytidine, wherein the dose comprises from 10-15 µg of the mRNA and is administered intramuscularly.
 7. The method of claim 6, wherein the dose comprises 10 µg of the mRNA.
 8. A method of vaccinating a human subject against a betacoronavirus, the method comprising administering to the human subject a dose of a vaccine comprising a messenger ribonucleic acid (mRNA) formulated in a lipid nanoparticle comprising 20-60 mol % ionizable cationic lipid, 5-25 mol % neutral lipid, 25-55 mol % cholesterol, and 0.5-15 mol % PEG-modified lipid, wherein the mRNA comprises an open reading frame encoding a full-length betacoronavirus spike protein, wherein the open reading frame comprises nucleosides consisting of N1-methyl-pseudouridine, adenosine, guanosine, and cytidine, wherein the dose comprises from 10-100 µg of the mRNA and is administered intramuscularly.
-