



US012404492B2

(12) **United States Patent**
Campana et al.

(10) **Patent No.:** **US 12,404,492 B2**

(45) **Date of Patent:** **Sep. 2, 2025**

(54) **METHODS FOR ENHANCING EFFICACY OF THERAPEUTIC IMMUNE CELLS**

(71) Applicant: **National University of Singapore**, Singapore (SG)

(72) Inventors: **Dario Campana**, Singapore (SG);
Takahiro Kamiya, Singapore (SG)

(73) Assignee: **NATIONAL UNIVERSITY OF SINGAPORE**, Singapore (SG)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 467 days.

(21) Appl. No.: **17/862,797**

(22) Filed: **Jul. 12, 2022**

(65) **Prior Publication Data**

US 2022/0370501 A1 Nov. 24, 2022

Related U.S. Application Data

(63) Continuation of application No. 16/943,400, filed on Jul. 30, 2020, which is a continuation of application (Continued)

(51) **Int. Cl.**
C12N 5/0783 (2010.01)
A61K 40/11 (2025.01)
(Continued)

(52) **U.S. Cl.**
CPC **C12N 5/0646** (2013.01); **A61K 40/11** (2025.01); **A61K 40/15** (2025.01); **A61K 40/31** (2025.01);
(Continued)

(58) **Field of Classification Search**
CPC C12N 5/0646; C12N 5/0636; C12N 2510/00; A61K 39/4611; A61K 39/4613;
(Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,946,778 A 8/1990 Ladner et al.
5,260,203 A 11/1993 Ladner et al.
(Continued)

FOREIGN PATENT DOCUMENTS

CA 2937157 A1 1/2018
CN 107709548 A 2/2018
(Continued)

OTHER PUBLICATIONS

Janeway CA Jr, Travers P, Walport M, et al. , Immunobiology: The Immune System in Health and Disease. 5th edition New York: Garland Science; 2001. (Year: 2001).*

(Continued)

Primary Examiner — Tracy Vivlemore

Assistant Examiner — Thomas R. Amick


(74) *Attorney, Agent, or Firm* — Wilson Sonsini Goodrich & Rosati

(57) **ABSTRACT**

The present invention relates to a method of using a receptor (e.g., chimeric antigen receptor—CAR) that activates an immune response upon binding a cancer cell ligand in conjunction with a target-binding molecule that targets a protein or molecule for removal or neutralization to generate enhanced anti-cancer immune cells. The present invention also relates to engineered immune cells having enhanced therapeutic efficacy and uses thereof.

21 Claims, 10 Drawing Sheets

Specification includes a Sequence Listing.



Related U.S. Application Data

No. 15/548,577, filed as application No. PCT/SG2016/050063 on Feb. 5, 2016, now Pat. No. 10,765,699.

- (60) Provisional application No. 62/130,970, filed on Mar. 10, 2015, provisional application No. 62/112,765, filed on Feb. 6, 2015.

(51) Int. Cl.

A61K 40/15 (2025.01)
A61K 40/31 (2025.01)
A61K 40/42 (2025.01)
C07K 16/28 (2006.01)

(52) U.S. Cl.

CPC **A61K 40/4211** (2025.01); **A61K 40/4224** (2025.01); **C07K 16/2809** (2013.01); **C07K 16/2833** (2013.01); **C07K 16/289** (2013.01); **C12N 5/0636** (2013.01); **A61K 2239/48** (2023.05); **C07K 2319/03** (2013.01); **C12N 2510/00** (2013.01)

(58) Field of Classification Search

CPC A61K 39/4631; A61K 39/464412; A61K 39/464429; A61K 2239/48; C07K 16/2809; C07K 16/2833; C07K 16/289; C07K 2319/03; C07K 14/7051; C07K 16/2803; C07K 2317/56; C07K 2317/622; C07K 2319/04; A61P 35/00; A61P 37/02
 See application file for complete search history.

(56) References Cited**U.S. PATENT DOCUMENTS**

- | | | | |
|--------------|----|---------|-----------------|
| 7,887,805 | B2 | 2/2011 | Pedersen et al. |
| 8,119,775 | B2 | 2/2012 | Moretta et al. |
| 8,399,645 | B2 | 3/2013 | Campana et al. |
| 8,580,714 | B2 | 11/2013 | Almagro et al. |
| 8,614,307 | B2 | 12/2013 | Moretta et al. |
| 8,637,258 | B2 | 1/2014 | Padkjaer et al. |
| 8,796,427 | B2 | 8/2014 | Spee et al. |
| 8,981,065 | B2 | 3/2015 | Moretta et al. |
| 9,181,527 | B2 | 11/2015 | Sentman |
| 9,273,283 | B2 | 3/2016 | Sentman |
| 9,422,368 | B2 | 8/2016 | Spee et al. |
| 9,683,042 | B2 | 6/2017 | Lee et al. |
| 9,902,936 | B2 | 2/2018 | Moretta et al. |
| 10,550,183 | B2 | 2/2020 | Png et al. |
| 10,730,942 | B2 | 8/2020 | Pule et al. |
| 10,765,699 | B2 | 9/2020 | Campana et al. |
| 11,161,907 | B2 | 11/2021 | June et al. |
| 11,440,958 | B2 | 9/2022 | Png et al. |
| 11,648,269 | B2 | 5/2023 | Campana et al. |
| 11,679,132 | B2 | 6/2023 | Campana et al. |
| 11,945,865 | B2 | 4/2024 | Png et al. |
| 2005/0282181 | A1 | 12/2005 | Yan et al. |
| 2006/0034834 | A1 | 2/2006 | Marasco et al. |
| 2007/0036773 | A1 | 2/2007 | Cooper et al. |
| 2009/0196850 | A1 | 8/2009 | Romagne et al. |
| 2012/0282256 | A1 | 11/2012 | Campana et al. |
| 2013/0266551 | A1 | 10/2013 | Campana et al. |
| 2013/0287748 | A1 | 10/2013 | June et al. |
| 2014/0120622 | A1 | 5/2014 | Gregory et al. |
| 2014/0186387 | A1 | 7/2014 | Lauer et al. |
| 2016/0256488 | A1 | 9/2016 | Wu |
| 2016/0312182 | A1 | 10/2016 | Sentman |
| 2017/0335331 | A1 | 11/2017 | Zhao et al. |
| 2018/0008638 | A1 | 1/2018 | Campana et al. |
| 2018/0066034 | A1 | 3/2018 | Ma et al. |
| 2018/0086831 | A1 | 3/2018 | Pule et al. |
| 2019/0002912 | A1 | 1/2019 | Lu et al. |
| 2019/0038733 | A1 | 2/2019 | Campana et al. |
| 2019/0046571 | A1 | 2/2019 | Campana et al. |

2019/0345217	A1	11/2019	Ma et al.
2020/0087398	A1	3/2020	Qasim et al.
2021/0046112	A1	2/2021	Campana et al.
2022/0347219	A1	11/2022	Campana et al.

FOREIGN PATENT DOCUMENTS

CN	113713091	A	11/2021
CN	107709548	B	6/2022
CN	115029362	A	9/2022
EP	0404097	A2	12/1990
EP	2247619	A1	11/2010
EP	3169773	A2	5/2017
EP	3322801	A1	5/2018
EP	3359168	A1	8/2018
EP	3474867	A1	5/2019
EP	3568467	A1	11/2019
JP	H03219896	A	9/1991
JP	H09501824	A	2/1997
JP	2001516766	A	10/2001
JP	2004529610	A	9/2004
JP	2008506368	A	3/2008
JP	2008518021	A	5/2008
JP	2009511495	A	3/2009
JP	2010537671	A	12/2010
JP	2011510047	A	3/2011
JP	2014507118	A	3/2014
JP	6895380	B2	6/2021
JP	2021137024	A	9/2021
WO	WO-8801649	A1	3/1988
WO	WO-9311161	A1	6/1993
WO	WO-9914353	A2	3/1999
WO	WO-03051926	A2	6/2003
WO	WO-2005017163	A2	2/2005
WO	WO-2006003179	A2	1/2006
WO	WO-2009092805	A1	7/2009
WO	WO-2012079000	A1	6/2012
WO	WO-2013126712	A1	8/2013
WO	WO-2014011984	A1	1/2014
WO	WO-2014124143	A1	8/2014
WO	WO-2014134165	A1	9/2014
WO	WO-2014159940	A1	10/2014
WO	WO-2015075468	A1	5/2015
WO	WO-2015121454	A1	8/2015
WO	WO-2015150771	A1	10/2015
WO	WO-2016055551	A1	4/2016
WO	WO-2016102965	A1	6/2016
WO	WO-2016126213	A1	8/2016
WO	WO-2017213979	A1	12/2017
WO	WO-2018027036	A1	2/2018
WO	WO-2019032916	A1	2/2019

OTHER PUBLICATIONS

Khan et al., Protein disulfide isomerase a multifunctional protein with multiple physiological roles. *Front. Chem.*, Aug. 25, 2014 Sec. Cellular Biochemistry vol. 2—2014 (Year: 2014).*

Brown et al. Novel Treatments for Chronic Lymphocytic Leukemia and Moving Forward. American Society of Clinical Oncology Educational Book, pp. e317-e325 (2014). Retrieved at URL: https://ascopubs.org/doi/pdfdirect/10.14694/EdBook_AM.2014.34.e317.

Caruana et al. From Monoclonal Antibodies to Chimeric Antigen Receptors for the Treatment of Human Malignancies. *Semin Oncol.* Oct. 2014 ; 41(5): 661-666.

Certified copy of PCT/US2016/019953 priority document U.S. Appl. No. 62/121,842, MA, Yupo et al., inventors, filed Feb. 27, 2015.

Feng et al. Treatment of Aggressive T Cell Lymphoblastic Lymphoma/leukemia Using Anti-CD5 Car T Cells. *Stem Cell Reviews and Reports* (2021) 17:652-661. Published online Jan. 6, 2021.

Fujiwara, et al. Adoptive Immunotherapy for Hematological Malignancies Using T Cells Gene-Modified to Express Tumor Antigen-Specific Receptors. *Pharmaceuticals* (Basel). Dec. 2014; 7(12): 1049-1068.

Glienke et al. Advantages and applications of CAR-expressing natural killer cells. *Frontiers in Pharmacology*, vol. 6, Article 21 (Feb. 12, 2015). 7 pages.

(56)

References Cited**OTHER PUBLICATIONS**

- Guo et al. Efficiency and side effects of anti-CD38 CAR T cells in an adult patient with relapsed B-ALL after failure of bi-specific CD19/CD22 CAR T cell treatment. *Cell Mol Immunol* 17, 430-432 (2020).
- Hishima et al. CD5 Expression in Thymic Carcinoma. *American Journal of Pathology*, vol. 145, No. 2, pp. 268-275 (Aug. 1994).
- Imboden et al. Stimulation of CD5 enhances signal transduction by the T cell antigen receptor. *J Clin Invest.* 1990. 85:130-134.
- Jena et al., Redirecting T-cell specificity by introducing a tumor-specific chimeric antigen receptor. *Blood.* 116:1035-1044 (2010).
- Kapp et al. Chapter 1: Post-Targeting Functions of Signal Peptides. In *Protein Transport into the Endoplasmic Reticulum*, Zimmerman, ed., 2009, Landes Bioscience. 13 pages. Retrieved at URL: <https://www.ncbi.nlm.nih.gov/books/NBK6322/>.
- Lemaistre et al. Phase I Trial of H65-RTA Immunoconjugate in Patients With Cutaneous T-cell Lymphoma. *Blood*, vol. 78, No. 5 (Sep. 1), 1991: pp. 1173-1182.
- Lewis et al. The immunophenotype of pre-TALL/LBL revisited. *Experimental and Molecular Pathology* 81 (2006) 162-165. Available online Aug. 14, 2006.
- Li et al. Flow Cytometry in the Differential Diagnosis of Lymphocyte-Rich Thymoma From Precursor T-Cell Acute Lymphoblastic Leukemia/ Lymphoblastic Lymphoma. *Am J Clin Pathol* 2004;121:268-274.
- Litzow et al. How I treat T-cell acute lymphoblastic leukemia in adults. *Blood.* 2015; 126(7):833-841. Retrieved at URL: <https://www.academia.edu/download/47911039/j.yexmp.2006.06.00620160809-3503-cb5sgz.pdf>.
- Liu et al. [NK cell surface receptors and their research progress—review]. *Zhongguo shi yan xue ye xue za zhi.* Aug. 2012;20(4):1034-1038. English abstract only. Retrieved at URL: <https://pubmed.ncbi.nlm.nih.gov/22931679/>. One page.
- Morris et al. Antibody-based therapy of leukaemia. *Expert Rev Mol Med.* Sep. 30, 2009; 11: e29.
- Pinz et al. Preclinical targeting of human T cell malignancies using CD4-specific chimeric antigen receptor (CAR)-engineered T cells. *Leukemia*, accepted article preview (Nov. 3, 2015). Retrieved at URL: https://www.researchgate.net/profile/Alexander-Jares/publication/283493430_Preclinical_targeting_of_human_T_cell_malignancies_using_CD4-specific_chimeric_antigen_receptor_CAR-engineered_T_cells/links/5643662808aef646e6c6a549/Preclinical-targeting-of-human-T-cell-malignancies-using-CD4-specific-chimeric-antigen-receptor-CAR-engineered-T-cells.pdf. 30 pages.
- Rezvani et al. The Application of Natural Killer Cell immunotherapy for the Treatment of Cancer. *Frontiers in Immunology*, vol. 6, Article 578 (Nov. 17, 2015). 13 pages.
- Roitt et al. *Roitt's Essential Immunology*. 12th Edition, Wiley-Blackwell, Chapter 4 (2012).
- Sommermeier, et al., Chimeric antigen receptor-modified T cells derived from defined CD8+ and CD4+ subsets confer superior antitumor reactivity in vivo. *Leukemia* 30(2):492-500 (2016).
- U.S. Appl. No. 17/862,721 Notice of Allowance dated Jun. 9, 2023.
- Zhong-Fu et al. Clinical Applications of NK Cells in Tumor Immunotherapy. *Chinese Journal of Biochemistry and Molecular Biology* 27(12):1088-1093 (Dec. 2011). With English translation.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_143513_us-15-548-577-37.rag. Search run on Jan. 7, 2019. Copy retrieved Sep. 11, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b67>. . . . 4 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_143514_us-15-548-577-36.rapbm. Search run on Jan. 7, 2019. Copy retrieved Sep. 11, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 13 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_143514_us-15-548-577-37.rai. Search run on Jan. 7, 2019. Copy retrieved Jan. 8, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 11 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_145543_us-15-548-577-32.rag. Search run on Jan. 7, 2019. Copy retrieved Jan. 8, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 23 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_145544_us-15-548-577-32.rai. Search run on Jan. 7, 2019. Copy retrieved Jan. 8, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 4 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_145544_us-15-548-577-32.rapbm. Search run on Jan. 7, 2019. Copy retrieved Jan. 8, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 6 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_145544_us-15-548-577-33.rai. Search run on Jan. 7, 2019. Copy retrieved Jan. 8, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 4 pages.
- Score Search Results Details for U.S. Appl. No. 15/548,577 and Search Result 20190107_145544_us-15-548-577-33.rapbm. Search run on Jan. 7, 2019. Copy retrieved Jan. 8, 2019 at <http://score.uspto.gov/ScoreAccessWeb/getItem.htm?AppId=15548577&seqId=09323b678>. . . . 12 pages.
- Chen, Kevin H., et al. Novel Anti-cd3 Chimeric Antigen Receptor Targeting of Aggressive T Cell Malignancies. *Oncotarget*, vol. 7, 56219-56232 (2016).
- Kipriyanov, Sergey M., et al. Two amino acid mutations in an anti-human CD3 single chain Fv antibody fragment that affect the yield on bacterial secretion but not the affinity. *Protein Engineering*, vol. 10, 445-453 (1997).
- U.S. Appl. No. 18/295,226 Office Action dated Aug. 2, 2024.
- EP22180254.9 Extended European Search Report dated Jan. 16, 2023.
- U.S. Appl. No. 16/100,117 Notice of Allowance dated Feb. 17, 2023.
- U.S. Appl. No. 16/100,117 Notice of Allowance dated Jan. 5, 2023.
- U.S. Appl. No. 17/862,721 Notice of Allowance dated Feb. 15, 2023.
- U.S. Appl. No. 17/862,721 Office Action dated Jan. 5, 2023.
- Alanen et al., Beyond KDEL: the role of positions 5 and 6 in determining ER localization. *J. Mol. Biol.* (2011) 409, 291-297. Available online Apr. 6, 2011.
- Alarcon et al., Assembly of the human T cell receptor-CD3 complex takes place in the endoplasmic reticulum and involves intermediary complexes between the CD3-gamma.delta.epsilon core and single T cell receptor alpha or beta chains, *Journal of Biological Chemistry*, Feb. 25, 1988;263(6):2953-61.
- Almagro, Juan C., Fransson, Johan. Humanization of antibodies. *Frontiers in Bioscience* 13;1619-1633 (Jan. 1, 2008).
- Altschul et al. Basic local alignment search tool. *J Mol Biol* 215(3):403-410 (1990).
- Anti-CD3 epsilon [OKT-3 (muromonab)]. Absolute Antibody. Website. Copyright 2021. Retrieved Dec. 26, 2021 at URL: <https://absoluteantibody.com/product/anti-cd3-epsilon-okt-3>. . . . 4 pages.
- Anti-CD3D monoclonal antibody, clone PLU4 (DCABH-10124). Product Information. CD Creative Diagnostics. Publication date unknown. 2 pages.
- Anti-TCR [BMA031]. Absolute Antibody. Website. Copyright 2021. Retrieved Dec. 26, 2021 at URL: <https://absoluteantibody.com/product/anti-tcr-bma031/>. 4 pages.

(56)

References Cited**OTHER PUBLICATIONS**

- Anwer et al., Donor origin CAR T cells: graft versus malignancy effect without GVHD, a systematic review, *Immunotherapy*, Jan. 2017;9(2):123-130.
- APPELBAUM. Haematopoietic cell transplantation as immunotherapy. *Nature*, vol. 411, pp. 385-389 (2001).
- Arafat et al. Antineoplastic effect of anti-erbB-2 intrabody is not correlated with scFv affinity for its target. *Cancer Gene Therapy*, vol. 7, No. 9, 2000: pp. 1250-1256.
- Arase et al. Recognition of virus infected cells by NK cells (w/ English abstract). Department of Immunochemistry, Research Institute for Microbial Diseases, Osaka University, vol. 54, No. 2, pp. 153-160 (2004).
- Austyn et al. T cell activation by anti-CD3 antibodies: function of Fc receptors on B cell blasts, but not resting B cells, and CD18 on the responding T cells. *Eur J Immunol* 17:1329-1335 (1987).
- Böldicke et al., Blocking translocation of cell surface molecules from the ER to the cell surface by intracellular antibodies targeted to the Er, *J. Cell. Mol. Med.*, 11(1):54-70 (2007).
- Böldicke et al. Functional inhibition of transitory proteins by intrabody-mediated retention in the endoplasmatic reticulum. *Meth-ods* 56 (2012) 338-350. Available online Oct. 20, 2011.
- Bleakley et al., Molecules and mechanisms of the graft-versus-leukaemia effect, *Nature Reviews, Cancer*, May 2004;4(5):371-80.
- Boettcher et al., Choosing the Right Tool for the Job: RNAi, TALEN, or CRISPR, *Mol Cell*, May 21, 2015;58:575-85.
- Brentjens et al. CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia. *Sci Transl Med*, Mar. 20, 2013;5(177):177ra38. 9 pages.
- Brentjens et al. Eradication of systemic B-cell tumors by genetically targeted human T lymphocytes co-stimulated by CD80 and interleukin-15. *Nat Med*, Mar. 2003;9(3):279-86.
- Brentjens et al. Safety and persistence of adoptively transferred autologous CD19-targeted T cells in patients with relapsed or chemotherapy refractory B-cell leukemias. *Blood*, Nov. 3, 2011;118(18):4817-28. Prepublished online Aug. 17, 2011.
- Brudno et al. Allogeneic T cells that express an anti-CD19 chimeric antigen receptor induce remissions of B-cell malignancies that progress after allogeneic hematopoietic stem-cell transplantation without causing graft-versus-host disease. *J Clin Oncol* 34(10):1112-1121 (2016).
- Burns et al. Two monoclonal anti-human T lymphocyte antibodies have similar biologic effects and recognize the same cell surface antigen. *J Immunol* 1982; 129:1451-1457.
- Campana et al. 4-1BB chimeric antigen receptors. *Cancer J*, Mar-Apr. 2014;20(2):134-40.
- Ceuppens et al. Failure of OKT3 monoclonal antibody to induce lymphocyte mitogenesis: a familial defect in monocyte helper function. *J Immunol*, vol. 134, No. 3, pp. 1498-1502 (Mar. 1985).
- Chang et al. A chimeric receptor with NKG2D specificity enhances natural killer cell activation and killing of tumor cells. *Cancer Res* 73(6):1777-86 (Mar. 15, 2013). Published online Jan. 9, 2013.
- Chen et al., Donor-derived CD19-targeted T cell infusion induces minimal residual disease-negative remission in relapsed B-cell acute lymphoblastic leukaemia with No. response to donor lymphocyte infusions after haploidentical haematopoietic stem cell transplantation. *British Journal of Haematology*, 2017, 179, 598-605. Epub Oct. 26, 2017.
- Clevers et al., The T cell receptor/CD3 complex: a dynamic protein ensemble, *Annual Review of Immunology*, 1988;6:629-62.
- Clift, Dean, et al. "A Method for the Acute and Rapid Degradation of Endogenous Proteins", Elsevier Inc., Cell 172, Dec. 14, 2017, pp. 1692-1706.
- Cooley et al. Donor selection for natural killer cell receptor genes leads to superior survival after unrelated transplantation for acute myelogenous leukemia. *Blood* (2010) 116 (14): 2411-2419.
- Cooper, et al. T-cell clones can be rendered specific for CD19: toward the selective augmentation of the graft-versus-B-lineage leukemia effect. *Blood*, Feb. 15, 2003;101(4):1637- 44. Epub Oct. 10, 2002.
- Dai et al., Tolerance and efficacy of autologous or donor-derived T cells expression CD19 chimeric antigen receptors in adult B-ALL with extramedullary leukemia. *Oncol Immunology*, 4:11, e1027469 (2015). 12 pages.
- Davila. Efficacy and toxicity management of 19-28z Car T cell therapy in B cell acute lymphoblastic leukemia. *Sci Transl Med* 6(224):224ra25 (2014).
- Dotti et al. Design and Development of Therapies using Chimeric Antigen Receptor-Expressing T cells. *Immunol Rev*, 257(1):35 pgs (2014).
- EP16746922.0 Extended European Search Report dated Sep. 21, 2018.
- EP18844536.5 Extended European Search Report dated Mar. 18, 2021.
- Eshhar et al. Specific activation and targeting of cytotoxic lymphocytes through chimeric single chains consisting of antibody-binding domains and the gamma or zeta subunits of the immunoglobulin and T-cell receptors. *Proc Natl Acad Sci U S A*, vol. 90, pp. 720-724 (Jan. 1993).
- Eyquem et al., Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances tumour rejection, *Nature*, Mar. 2, 2017; 543(7643): 113-117.
- Gale. Cellular and Molecular Basis of Cancer. Merck Manual Professional Version. Last full review/revision Nov. 2020. Content last modified Nov. 2020. Retrieved Jan. 8, 2021 at URL: <https://www.merckmanuals.com/professional/hematology-and-oncology/overview-of-cancer/cellular-and-molecular-basis-of-cancer#8> pages.
- Gan et al. Molecular Mechanisms and Potential Therapeutic Reversal of Pancreatic Cancer-Induced Immune Evasion. *Cancers* 2020, 12, 1872. Published Jul. 11, 2020. 22 pages.
- Gao et al. Retention mechanisms for ER and Golgi membrane proteins. *Trends in Plant Science*, vol. 19, Issue 8, pp. 508-515 (Aug. 2014).
- Geiger et al. Integrated src kinase and costimulatory activity enhances signal transduction through single-chain chimeric receptors in T lymphocytes. *Blood* 98(8):2364-2371 (2001).
- Geiger et al. The TCR zeta-chain immunoreceptor tyrosine-based activation motifs are sufficient for the activation and differentiation of primary T lymphocytes. *The Journal of Immunology*, 1999, 162: 5931-5939.
- Ghosh et al., Donor CD19 Car T cells exert potent graft-versus-lymphoma activity with diminished graft-versus-host activity, *Nat Med*, Feb. 2017 ; 23(2): 242-249.
- Giebel et al. Survival advantage with KIR ligand incompatibility in hematopoietic stem cell transplantation from unrelated donors. *Blood* (2003) 102 (3): 814-819.
- Grimshaw, B.D. et al., BGST Abstract Mar. 9, 2012, abstract P023, "Creating a 'null' T cell for use in adoptive immunotherapy", British Society for Gene and Cell Therapy 2012, <http://www.bsqct.org>, Human Gene Therapy, 22 pages.
- Grimshaw. Developing a universal T cell for use in adoptive immunotherapy (thesis), University College London (2015). Retrieved Aug. 22, 2022 at URL: <http://discovery.ucl.ac.uk/1470207/1/Grimshaw%20Ben%20Thesis.pdf>. 267 pages.
- Grimshaw et al. Creating a Null T Cell for Adoptive Immunotherapy. Poster. Mar. 9, 2012. One page.
- Grovender et al. Single-chain antibody fragment-based adsorbent for the extracorporeal removal of β 2-microglobulin. *Kidney International*, vol. 65 (2004), pp. 310-322.
- Grupp et al. Chimeric Antigen Receptor-Modified T Cells for Acute Lymphoid Leukemia. *N Engl J Med* 368; 16, pp. 1509-1508 (Apr. 18, 2013). With correction published *N. Engl J. Med* (2016) 374(10) 998.
- Haegert et al. Co-expression of surface immunoglobulin and T3 on hairy cells. *Scand J Haematol* 1986;37:196-202.
- Haynes et al. Rejection of syngeneic colon carcinoma by CTLs expressing single-chain antibody receptors codelivering CD28 costimulation. *J Immunol* 2002; 169:5780-5786.
- Haynes et al. Single-chain antigen recognition receptors that costimulate potent rejection of established experimental tumors. *Blood*, Nov. 1, 2002;100(9):3155-63. Published online Jul. 5, 2002.

(56)

References Cited

OTHER PUBLICATIONS

- Hegde, M. et al., "Combinational targeting offsets antigen escape and enhances effector functions of adoptively transferred T cells in glioblastoma", *Molecular Therapy*, 2013, vol. 21, pp. 2087-2101.
- Hegde, M. et al., Supplementary Material for "Combinational targeting offsets antigen escape and enhances effector functions of adoptively transferred T cells in glioblastoma", *Molecular Therapy*, 2013, vol. 21, pp. 2087-2101. Retrieved Jan. 6, 2022 from URL: <https://ars.els-cdn.com/content/image/1-s2.0-S1525001616309315-mmcl.pdf>. 9 pages.
- Hexham et al. Influence of relative binding affinity on efficacy in a panel of anti-CD3 scFv immunotoxins. *Molecular Immunology* 38 (2001) 397-408.
- Holliger et al. "Diabodies": small bivalent and bispecific antibody fragments. *Proc Natl Acad Sci USA*, vol. 90, pp. 6444-6448 (Jul. 1993).
- Hombach et al. Tumor-Specific T Cell Activation by Recombinant Immunoreceptors: CD3 ζ Signaling and CD28 Costimulation Are Simultaneously Required for Efficient IL-2 Secretion and Can Be Integrated Into One Combined CD28/CD3 ζ Signaling Receptor Molecule. *J Immunol* 2001; 167:6123-6131.
- Imai et al. Chimeric receptors with 4-1BB signaling capacity provoke potent cytotoxicity against acute lymphoblastic leukemia. *Leukemia* (2004) 18, 676-684.
- Imai et al. Genetic modification of primary natural killer cells overcomes inhibitory signals and induces specific killing of leukemic cells. *Blood* (2005) 106 (1): 376-383.
- Imamura, M. et al., "Autonomous growth and increased cytotoxicity of natural killer cells expressing membrane-bound interleukin-15", *Blood*, 2014, vol. 124, pp. 1081-1088.
- Imamura, M. et al., Supplementary Material for "Autonomous growth and increased cytotoxicity of natural killer cells expressing membrane-bound interleukin-15", *Blood*, 2014, vol. 124, pp. 1081-1088. Retrieved Jan. 6, 2021 at URL: https://ash.silverchair-cdn.com/ash/content_public/journal/blood/124/7/10.1182_blood-2014-02-556837/4/blood-2014-02-556837-1.pdf?Expires=1644340511&Signature=uyyKt0KS8WtXpVUULTgok2RyymzBJgE2vCNIDD4xdwCv13vsg0gLOpQLU~KVPtITvIHmCLEX2MhA7mexzXy~ydDqj6rHeZEBNohY4NOkmjpH9529c9SCChMFBIn80TH-cM-MgQfrEtegs40K6vjiveJODaZP6TfW1gGK~5JUAn5LesZfPv9W28NmBfMoAOMVeX4Pz54V~9dWaBcCfXCR7vOrx1N8cpbxmLAumSziwKqXNcy79dwOL6ddz3joiyKtMiGnuY1c6l1f6b~MwbLxZ3jK16EE-giQZhSxFLm2ctwuCbOMj8RIHCM4cO5a2zIKMAM6dut-dafGbQaRQ_&Key-Pair-Id=APKAIE5G5CRDK6RD3PGA. 11 pages.
- Jackson et al., Identification of a consensus motif for retention of transmembrane proteins in the endoplasmic reticulum, *EMBO J.* Oct. 1990; 9(10): 3153-3162.
- Joshi et al. Fusion to a highly charged proteasomal retargeting sequence increases soluble cytoplasmic expression and efficacy of diverse anti-synuclein intrabodies. *mAbs*, vol. 4, Issue 6, pp. 686-693 (2012). Published online: Aug. 28, 2012.
- Kalos, et al., "T Cells with Chimeric Antigen Receptors Have Potent Antitumor Effects and Can Establish Memory in Patients with Advanced Leukemia", *Science Translation Medicine* 3.95 (2011):95ra73-95ra73.
- Kamiya, et al., "A novel method to generate T-cell receptor-deficient antigen receptor T cells," *Blood Advances*, vol. 2, No. 5, Mar. 13, 2018, pp. 517-528.
- Kamiya et al. Blocking expression of inhibitory receptor NKG2A overcomes tumor resistance to NK cells. *J Clin Invest.* 2019; 129(5):2094-2106.
- Kloss, C.C. et al., Supplementary Text and Figures for "Combinatorial antigen recognition with balanced signaling promotes selective tumor eradication by engineered T cells", *Nature Biotechnology*, 2013, vol. 31, pp. 71-75. Retrieved Jan. 6, 2021 at URL: https://static-content.springer.com/esm/art%3A10.1038%2Fnt.2459/MediaObjects/41587_2013_BFnt2459_MOESM2_ESM.pdf. 5 pages.
- Kloss. Combinatorial antigen recognition with balanced signaling promotes selective tumor eradication by engineered T cells. *Nature Technology*, vol. 31, No. 1, pp. 71-75 (Jan. 2013). Published online Dec. 16, 2012.
- Kochenderfer et al. B-cell depletion and remissions of malignancy along with cytokine-associated toxicity in a clinical trial of anti-CD19 chimeric-antigen-receptor-transduced T cells. *Blood*. 2012;119(12):2709-2720. Published online Dec. 8, 2011.
- Kochenderfer et al., Chemotherapy-refractory diffuse large B-cell lymphoma and indolent B-cell malignancies can be effectively treated with autologous T cells expressing an anti-CD19 chimeric antigen receptor, *Journal of Clinical Oncology*, Feb. 20, 2015; 33(6); pp. 540-549. Published online Aug. 25, 2014.
- Kochenderfer et al., Donor-derived CD19-targeted T cells cause regression of malignancy persisting after allogeneic hematopoietic stem cell transplantation, *Blood*. 2013;122(25):4129-4139.
- Kolb et al. Graft-versus-leukemia effect of donor lymphocyte transfusions in marrow grafted patients. *Blood*, vol. 86. No. 5 (Sep. 1), 1995: pp. 2041-2050.
- Kotteas et al. Immunotherapy for pancreatic cancer. *J Cancer Res Clin Oncol* (2016) 142:1795-1805. Published online Feb. 3, 2016.
- Kudo et al. T lymphocytes expressing a CD16 signaling receptor exert antibody-dependent cancer cell killing. *Cancer Res* 74(1):93-103 (2013).
- Lanitis et al. Chimeric antigen receptor T Cells with dissociated signaling domains exhibit focused anti-tumor activity with reduced potential for toxicity in vivo. *Cancer Immunol Res* 1(1):43-53 (2013).
- Lee, et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood*. Jul. 10, 2014;124(2):188-95. doi: 10.1182/blood-2014-05-552729. Epub May 29, 2014.
- Lee et al., T cells expressing CD 19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. *Lancet*, Feb. 7, 2015; 385 (9967); pp. 517-528. Epub Oct. 13, 2014.
- Lo et al. Harnessing the tumour-derived cytokine, CSF-1, to co-stimulate T-cell growth and activation. *Molecular Immunology* 45 (2008) 1276-1287. Available online Oct. 24, 2007.
- Lorentzen et al. CD19-Chimeric Antigen Receptor T Cells for Treatment of Chronic Lymphocytic Leukaemia and Acute Lymphoblastic Leukaemia. *Scandinavian Journal of Immunology*, 2015, 82, 307-319.
- Maciocia et al. A simple protein-based method for generation of 'off the shelf' allogeneic chimeric antigen receptor T-cells. *Database Ebase [Online]* 1-15, Elsevier Science Publishers, Amsterdam, NL (May 1, 2018). Abstract. XP002784541. Database Accession No. EMB-623339718.
- Maciocia et al. A simple protein-based method for generation of 'off the shelf' allogeneic chimeric antigen receptor T-cells. *Molecular Therapy*, vol. 26, No. 5, Supplement 1, pp. 296-297 (May 2018). Cell Press NLD. May 16, 2018 to May 19, 2018 Chicago, IL-297 Conf. ISSN: 1525-0024.
- MacLeod et al., Integration of a CD19 Car into the TCR Alpha Chain Locus Streamlines Production of Allogeneic Gene-Edited CAR T Cells, *Molecular Therapy*, vol. 25, No. 4, pp. 949-961, Apr. 2017.
- Maier et al. Human T-lymphocyte cytotoxicity and proliferation directed by a single chimeric TCRzeta /CD28 receptor. *Nat Biotech* 20(1):70-75 (2002).
- Marasco et al. Design, intracellular expression, and activity of a human anti-human immunodeficiency virus type 1 gp120 single-chain antibody. *PNAS USA* 90:7889-7893 (1993).
- Marschall, Andrea LJ, et al. "Specific in vivo knockdown of protein function by intrabodies", Taylor & Francis Group, LLC, Nov./Dec. 2015, vol. 7, Issue 6, pp. 1010-1035.
- Maude, et al., Chimeric Antigen Receptor T Cells for Sustained Remissions in Leukemia. *N. Engl J. Med* (2014) 371(16) 1507-1517. With correction published *N. Engl J. Med* 374(10): 998 (2016).
- Miller et al. Successful adoptive transfer and in vivo expansion of human haploidentical NK cells in patients with cancer. *Blood* 105:3051-3057 (2005).

(56)

References Cited**OTHER PUBLICATIONS**

- Miller. Therapeutic applications: natural killer cells in the clinic. Hematology Am Soc Hematol Educ Program (2013) 2013 (1): 247-253.
- Milone et al. Chimeric receptors containing CD137 signal transduction domains mediate enhanced survival of T cells and increased antileukemic efficacy in vivo. Mol. Ther. 17:1453-1464 (2009).
- Munro et al., A C-terminal signal prevents secretion of luminal ER proteins, Cell. Mar. 13, 1987;48:899-907. Retrieved Apr. 7, 2022 at URL: <https://bio.davidson.edu/molecular/MunPelham/mufixed.html>.
- Kochenderfer et al., Eradication of B-lineage cells and regression of lymphoma in a patient treated with autologous T cells genetically engineered to recognize CD19 (Blood, 2010, 116:4099-4102) (Year: 2010).
- Needleman et al. A general method applicable to the search for similarities in the amino acid sequence of two proteins. Journal of molecular biology 48(3):443-453 (1970).
- Neelapu et al., Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma, N Engl J Med 377:26 pp. 2531-2544 (Dec. 28, 2017).
- Pardoll. The blockade of immune checkpoints in cancer immunotherapy. Nat Rev Cancer. 12(4):252-264 (2012).
- Park et al. Are all chimeric antigen receptors created equal? J Clin Oncol. Feb. 20, 2015;33(6):651-3. Epub Jan. 20, 2015.
- Park et al., CD19-targeted CAR T-cell therapeutics for hematologic malignancies: interpreting clinical outcomes to date, Blood. 2016; 127(26):3312-3320.
- PCT/SG2016/050063 International Search Report and Written Opinion dated May 9, 2016.
- PCT/US2018/046137 International Search Report and Written Opinion dated Oct. 29, 2018.
- Pearson and Lipman, Improved tools for biological sequence comparison, Proc. Nat. Acad Sci USA., 85:2444-2448, (1988).
- Peipp et al. A Recombinant CD7-specific Single-Chain Immunotoxin Is a Potent Inducer of Apoptosis in Acute Leukemic T Cells. Cancer Research 62, pp. 2848-2855 (May 15, 2002).
- Png, et al., "Blockade of CD7 expression in T cells for effective chimeric antigen receptor targeting of T-cell malignancies", Blood Advances, vol. 1, No. 25, Nov. 28, 2017, pp. 2348-2360.
- Poirot et al. Multiplex Genome-Edited T-cell Manufacturing Platform for Off-the-Shelf Adoptive T-cell Immunotherapies. Cancer Research 75(18):3853-3864 (2015).
- Porter et al., Chimeric antigen receptor-modified T cells in chronic lymphoid leukemia. N Engl J Med 365:725-733 (2011). With correction published N. Engl J. Med (2016) 374(10) 998.
- Porter et al. Induction of Graft-versus-Host Disease as Immunotherapy for Relapsed Chronic Myeloid Leukemia. N Engl J Med 1994; 330:100-106.
- Qasim et al., Molecular remission of infant B-ALL after infusion of universal TALEN gene-edited CART cells. Sci. Transl. Med. 9, eaaj2013 (2017). With Erratum. Erratum retrieved Apr. 7, 2022 at URL: <https://www.science.org/doi/10.1126/scitranslmed.aam9292>. 3 pages.
- Reshef et al. Blockade of lymphocyte chemotaxis in visceral graft-versus-host disease. N Engl J Med 367;3 pp. 135-145 (Jul. 12, 2012).
- Rosenberg et al. Adoptive cell transfer as personalized immunotherapy for human cancer. Science 348(6230):62-68 (2015).
- Rossig, C. et al., "Genetic modification of T lymphocytes for adoptive immunotherapy", Molecular Therapy, 2004, vol. 10, pp. 5-18.
- Rowley, J. et al., "Expression of IL-15RA or an IL-15/IL-15RA fusion on CD8+ T cells modifies adoptively transferred T-cell function in cis", European Journal of Immunology, 2009, vol. 39, pp. 491-506.
- Rowley, J. et al., Supplementary Information for "Expression of IL-15RA or an IL-15/IL-15RA fusion on CD8+ T cells modifies adoptively transferred T-cell function in cis", European Journal of Immunology, 2009, vol. 39, pp. 491-506. Retrieved Jan. 6, 2022 at URL: https://onlinelibrary.wiley.com/action/downloadSupplement?doi=10.1002%2Feji.200838594&file=ejj_200838594_sm_SupplInfoFig.pdf. 6 pages.
- Rubnitz et al. NKAML: A Pilot Study to Determine the Safety and Feasibility of Haploidentical Natural Killer Cell Transplantation in Childhood Acute Myeloid Leukemia. J Clin Oncol. Feb. 20, 2010; 28(6): 955-959.
- Rudikoff, et al. Single amino acid substitution altering antigen-binding specificity. Proc Natl Acad Sci U S A. Mar. 1982;79(6):1979-83.
- Ruggeri et al. Effectiveness of donor natural killer cell alloreactivity in mismatched hematopoietic transplants. Science 295(5562):2097-2100 (2002).
- Sadelain et al. The Basic Principles of Chimeric Antigen Receptor Design. Cancer Discov. Apr. 2013 ; 3(4): 388-398.
- Sadelain et al. Therapeutic T cell engineering. Nature 545:423-431 (2017).
- Sanz, L. et al., "Antibodies and gene therapy: teaching old 'magic bullets' new tricks", Trends in Immunology, 2004, vol. 25, pp. 85-91.
- Sato et al. Single domain intrabodies against WASP inhibit TCR-induced immune responses in transgenic mice T cells. Sci Rep. Oct. 21, 2013;3:3003. 10 pages.
- Schumann, et al. Generation of knock-in primary human T cells using Cas9 ribonucleoproteins. Proc Natl Acad Sci U S A. Aug. 18, 2015;112(33):10437-42. doi: 10.1073/pnas. 1512503112. Epub Jul. 27, 2015.
- Schuster et al., Chimeric Antigen Receptor T Cells in Refractory B-Cell Lymphomas, Dec. 28, 2017,, the New England Journal of Medicine, 2017; 377; pp. 2545-2554.
- Schwartz. T cell anergy. Annu Rev Immunol. 2003;21:305-34. First published online as a Review in Advance on Dec. 5, 2002.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-16.rag. Search run on Aug. 29, 2022. Copy retrieved Aug. 30, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=bc544287-1664-44c7-9aff-7ec7ba8cb7ea&itemName=20220829_091720_... 14 pages.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-16.rai. Search run on Aug. 29, 2022. Copy retrieved Aug. 30, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=131a34dedcc5-448f-b75f-7ea24f699275&itemName=20220829_091720_... 8 pages.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-16.rapm. Search run on Aug. 29, 2022. Copy retrieved Aug. 30, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=e6f49583-c02c-4dee-a87c-e0ece396955c&itemName=20220829_091720_... 10 pages.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-17.rag. Search run on Aug. 29, 2022. Copy retrieved Aug. 30, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=ccb41ae5-1511-498f-ac7c-0b33abd2492f&itemName=20220829_091720_... 15 pages.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-17.rai. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=73e892a9-d59f-4aff-8df1-883c983c2966&itemName=20220829_091720_u... 8 pages.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-17.rapm. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=9851a8bc-7a77-4e69-985e-903a1f2622ae&itemName=20220829_09172_... 10 pages.
- Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-17.rpr. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/

(56)

References Cited**OTHER PUBLICATIONS**

ScoreAccessWeb/getItemDetail?appId=17862721&docId=e04b2be2-c5c8-4cba-b806-cd02a1efe868&itemName=20220829_091720_... 7 pages.

Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-20.rai. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=d17ad86f-3b93-47e9-a94e-a9258afaa3e3&itemName=20220829_091720_... 9 pages.

Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-20.rapm. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=5747899e-d2f6-4407-8197-6ae6e43ac551&itemName=20220829_09172_... 11 pages.

Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-20.rpr. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=8db4eb9f-566a-448b-8a1b-9244e2b8f0ee&itemName=20220829_091720_... 7 pages.

Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-21.rag. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=69e4190a-317d-4f26-9846-c30ab0601669&itemName=20220829_09172_... 14 pages.

Score Search Results Detail for Application 17862721 and Search Results 20220829_091720_us-17-862-721-21.rai. Search run on Aug. 29, 2022. Copy retrieved Aug. 31, 2022 at score.uspto.gov/ScoreAccessWeb/getItemDetail?appId=17862721&docId=a9c72e86-3699-4e4c-8110-8367d0fe6cf4&itemName=20220829_091720_... 8 pages.

Sharma et al., "Novel cancer immunotherapy agents with survival benefit: recent successes and next steps," *Nat Rev Cancer*, 11(11):805-812 (2011).

Shikano et al., Membrane receptor trafficking: Evidence of proximal and distal zones conferred by two independent endoplasmic reticulum localization signals, *PNAS* May 13, 2003 100 (10) 5783-5788.

Shimasaki et al. A clinically adaptable method to enhance the cytotoxicity of natural killer cells against B-cell malignancies. *Cytotherapy*, 2012; 14: 830-840.

Shimasaki et al., Natural killer cell reprogramming with chimeric immune receptors, *Methods Mol Biol*. 2013;969:203-20.

Slavin et al. Allogeneic cell therapy with donor peripheral blood cells and recombinant human interleukin-2 to treat leukemia relapse after allogeneic bone marrow transplantation. *Blood* (1996) 87 (6): 2195-2204.

Smith et al. Comparison of Biosequences. *Advances in Applied Mathematics*. 2:482-489 (1981).

Smith et al. T cell activation by anti-T3 antibodies: Comparison of IgG1 and IgG2b switch variants and direct evidence for accessory function of macrophage Fc receptors. *Eur J Immunol* 16:478-486 (1986).

Sommermeier et al., Fully human CD19-specific chimeric antigen receptors for T-cell therapy. *Leukemia*. Oct. 2017 ; 31(10): 2191-2199.

Strebe et al. Functional knockdown of VCAM-1 at the post-translational level with ER retained antibodies. *Journal of Immunological Methods* 341 (2009) 30-40. Available online Nov. 25, 2008.

Su et al., CRISPR-Cas9 mediated efficient PD-I disruption on human primary T cells from cancer patients, *Sci Rep* 6, 20070, Published: Jan. 28, 2016. 14 pages.

Till et al. CD20-specific adoptive immunotherapy for lymphoma using a chimeric antigen receptor with both CD28 and 4-1BB domains: pilot clinical trial results. *Blood* 119(17):3940-3950 (2012).

Todorovska et al. Design and application of diabodies, triabodies and tetrabodies for cancer targeting. *Journal of Immunological Methods* 248 (2001) 47-66.

Topp et al. Targeted therapy with the T-cell-engaging antibody blinatumomab of chemotherapy-refractory minimal residual disease in B-lineage acute lymphoblastic leukemia patients results in high response rate and prolonged leukemia-free survival. *J Clin Oncol*. Jun. 20, 2011;29(18):2493-8.

Torikai et al. A foundation for universal T-cell based immunotherapy: T cells engineered to express a CD19-specific chimeric-antigen-receptor and eliminate expression of endogenous TCR. *Blood* 119(24):5697-5705 (2012).

Turtle et al. CD19 Car-T cells of defined CD4+:CD8+ composition in adult B cell ALL patients. *J Clin Invest* 126(6):2123-2138 (2016).

U.S. Appl. No. 15/548,577 Notice of Allowance dated May 4, 2020.

U.S. Appl. No. 15/548,577 Office Action dated Jan. 14, 2019.

U.S. Appl. No. 15/548,577 Office Action dated Jul. 20, 2018.

U.S. Appl. No. 15/548,577 Office Action dated Sep. 17, 2019.

U.S. Appl. No. 16/100,117 Office Action dated Aug. 12, 2022.

U.S. Appl. No. 16/100,117 Office Action dated Dec. 9, 2021.

U.S. Appl. No. 16/100,117 Office Action dated May 3, 2021.

U.S. Appl. No. 16/100,120 Office Action dated Aug. 2, 2019.

U.S. Appl. No. 16/100,120 Office Action dated Feb. 24, 2020.

U.S. Appl. No. 16/100,120 Office Action dated Jan. 28, 2019.

U.S. Appl. No. 16/100,120 Office Action dated Oct. 6, 2020.

U.S. Appl. No. 17/862,721 Office Action dated Sep. 7, 2022.

Van Wauwe et al. Human T lymphocyte activation by monoclonal antibodies; OKT3, but not UCHL1, triggers mitogenesis via an interleukin 2-dependent mechanism. *J Immunol*, vol. 133, No. 1, pp. 129-132 (Jul. 1984).

Verneris et al. Natural Killer Cell Consolidation for Acute Myelogenous Leukemia: A Cell Therapy Ready for Prime Time? *J Clin Oncol*. Feb. 20, 2010; 28(6): 909-910.

Vervilghen et al. Differences in the stimulating capacity of immobilized anti-CD3 monoclonal antibodies: variable dependence on interleukin-1 as a helper signal for T-cell activation. *Immunology* 72:269-276 (1991).

Vivier, Eric et al., Innate or adaptive immunity? The example of natural killer cells, *Science* (New York, N.Y.) vol. 331,6013 (2011): 44-9. doi:10.1126/science.1198687.

Wang et al. Expression and characterization of recombinant soluble monkey CD3 molecules: mapping the FN18 polymorphic epitope. *Molecular Immunology* 40:1179-1188 (2004).

Weetall et al. T-cell depletion and graft survival induced by anti-human CD3 immunotoxins in human CD3 β transgenic mice. *Transplantation*, vol. 73, 1658-1666, No. 10 (May 27, 2002).

Wheeler et al. Intrabody and intrakine strategies for molecular therapy. *Molecular Therapy*, vol. 8, No. 3, pp. 355-366, Sep. 2003.

Wunderlich et al. OKT3 prevents xenogeneic GVHD and allows reliable xenograft initiation from unfractionated human hematopoietic tissues. *Blood*. 2014;123(24):e134-e144.

Yang et al., Challenges and opportunities of allogeneic donor-derived CAR T cells, *Current Opinion in Hematology*, Nov. 2015; 22 (6); pp. 509-515.

Zang et al. The B7 family and cancer therapy: costimulation and coinhibition. *Clin Cancer Res* 13(18) pp. 5271-5279 (Sep. 15, 2007).

Zhan et al. Modification of ricin A chain, by addition of endoplasmic reticulum (KDEL) or Golgi (YQRL) retention sequences, enhances its cytotoxicity and translocation. *Cancer Immunology, Immunotherapy*. Vol. 46, pp. 55-60 (1998).

Zhou, P. et al., "Cells transfected with a non-neutralizing antibody gene are resistant to HIV infection: targeting the endoplasmic reticulum and trans-Golgi network", *The Journal of Immunology*, 1998, vol. 160, pp. 1489-1496.

Arakawa et al. Cloning and Sequencing of the VH and YK Genes of an Anti-CD3 Monoclonal Antibody, and Construction of a Mouse/Human Chimeric Antibody. *J Biochem*, vol. 120, No. 3, pp. 657-662 (1996).

Certificate of Disclosure by Benjamin D. Grimshaw signed Dec. 15, 2021 for: Grimshaw et al. Creating a Null T Cell for Adoptive Immunotherapy. Poster. Mar. 9, 2012. One page.

(56)

References Cited**OTHER PUBLICATIONS**

Certificate of Disclosure by David Linch signed Dec. 16, 2021 for: Grimshaw et al. Creating a Null T Cell for Adoptive Immunotherapy. Poster. Mar. 9, 2012. One page.

Certificate of Disclosure by Martin Pule signed Dec. 15, 2021 for: Grimshaw et al. Creating a Null T Cell for Adoptive Immunotherapy. Poster. Mar. 9, 2012. One page.

Co-pending U.S. Appl. No. 18/295,226, inventors Campana; Dario et al., filed on Apr. 3, 2023.

Declaration by Benjamin D. Grimshaw signed Sep. 17, 2022. 2 pages.

Nilsson et al. Retention and retrieval in the endoplasmic reticulum and the Golgi apparatus. *Current Opinion in Cell Biology* 1994, 6:517-521.

Dong, et al. Modern Hematopoietic Stem Cell Transplant Therapeutics. Military Health Press. 1st Edition (2001):175-176. With English machine translation.

GenBank Accession No. AAA83267. Version No. AAA83267.1. sFv antibody, partial [Mus musculus]. Record created Dec. 14,

1995. Page Range: 1-4. Retrieved May 15, 2024 at URL: <https://www.ncbi.nlm.nih.gov/protein/AAA83267.1>.

GenBank Accession No. AAM12015. Version No. AAM12015.1. monoclonal anti-alpha-1,3-galactosyltransferase IgM heavy chain, partial [Mus musculus]. Record created Apr. 17, 2002. Page Range: 1-3. Retrieved May 15, 2024 at URL: <https://www.ncbi.nlm.nih.gov/protein/AAM12015.1>.

GenBank Accession No. ABK63997. Version No. ABK63997.1. Immunoglobulin light chain variable region, partial [Mus musculus]. Record created Nov. 25, 2006. Page Range: 1-3. Retrieved May 15, 2024 at URL: <https://www.ncbi.nlm.nih.gov/protein/ABK63997>.


Lanitis, Evripidis, et al., A Human ErbB2-Specific T-Cell Receptor Confers Potent Antitumor Effector Functions in Genetically Engineered Primary Cytotoxic Lymphocytes. *Human Gene Therapy* 25:730-739 (2014).

U.S. Appl. No. 16/943,400 Office Action dated Apr. 24, 2024.

Tragoolpua, Khajornsak et al. Generation of functional scFv intrabody to abate the expression of CD147 surface molecule of 293A cells. *BMC biotechnology* 8:5, 1-13 (2008).

U.S. Appl. No. 16/943,400 Office Action dated Feb. 5, 2025.

* cited by examiner



Name	Specificity tested	Signal peptide & ScFv			Localization domains
		CD8 SP	VL	Linker	
mb	CD3, CD45, β 2M, HLA I, KIR2DL1-3, NKG2A		VL	Linker	CD8 TM
PEST	CD3		VL	Linker	PEST
mb PEST	CD3, CD45, β 2M, KIR2DL1-3, NKG2A		VL	Linker	CD8 TM
KDEL	CD3, CD45		VL	Linker	PEST
SEKDEL	CD3		VL	Linker	KDEL
AEKDEL	CD3		VL	Linker	SEKDEL
myc KDEL	CD3, CD7, CD45, β 2M		VL	Linker	AEKDEL
PEST KDEL	CD3		VL	Linker	myc
link (20) KDEL	CD3		VL	Linker	PEST
link (10) KDEL	CD3		VL	Linker	KDEL
link (20) AEKDEL	CD3, CD7, CD45, β 2M, HLA I, KIR2DL1-3, NKG2A		VL	Linker	Linker
mb EEKMP	CD3, CD45, β 2M, HLA I, KIR2DL1-3, NKG2A		VL	Linker	Linker
mb OEKMP	CD3, CD7, CD45, β 2M, KIR2DL1-3, NKG2A		VL	Linker	Linker
mb HA-KKMP	CD3		VL	Linker	Linker
mb PEST KKMP	CD3		VL	Linker	Linker
mb KKTN	CD3		VL	Linker	Linker
mb YORL	CD3, CD45		VL	Linker	Linker
mb RNIKCD	CD3		VL	Linker	Linker
link (20) KDER1	CD3		VL	Linker	Linker

FIG. 2

FIG. 3B





FIG. 3A






FIG. 3C




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

1

METHODS FOR ENHANCING EFFICACY OF THERAPEUTIC IMMUNE CELLS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 16/943,400, filed Jul. 30, 2020, which is a continuation of U.S. patent application Ser. No. 15/548,577, filed Aug. 3, 2017, now U.S. Pat. No. 10,765,699, which is a 371 U.S. National Phase Application of International Patent Cooperation Treaty Application PCT/SG2016/050063, filed Feb. 5, 2016, which claims benefit to U.S. Provisional Application No. 62/112,765, filed Feb. 6, 2015, and U.S. Provisional Application No. 62/130,970, filed Mar. 10, 2015, the disclosures of which are incorporated herein by reference in their entirety for all purposes.

REFERENCE TO A SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on Jul. 11, 2022, is named 62190_701_302_SL.xml and is 87,725 bytes in size.

BACKGROUND OF THE INVENTION

Immune cells can be potent and specific “living drugs”. Immune cells have the potential to target tumor cells while sparing normal tissues; several clinical observations indicate that they can have major anti-cancer activity. Thus, in patients receiving allogeneic hematopoietic stem cell transplantation (HSCT), T-cell-mediated graft-versus-host disease (GvHD) (Weiden, P L et al., *N. Engl. J. Med.* 1979; 300(19):1068-1073; Appelbaum, FR *Nature*, 2001; 411 (6835):385-389; Porter, D L et al., *N. Engl. J. Med.* 1994; 330(2):100-106; Kolb, H J et al. *Blood*. 1995; 86(5):2041-2050; Slavin, S. et al., *Blood*. 1996; 87(6):2195-2204), and donor natural killer (NK) cell alloreactivity (Ruggeri L, et al. *Science*. 2002; 295(5562):2097-2100; Giebel S, et al. *Blood*. 2003; 102(3):814-819; Cooley S, et al. *Blood*. 2010; 116 (14):2411-2419) are inversely related to leukemia recurrence. Besides the HSCT context, administration of antibodies that release T cells from inhibitory signals (Sharma P, et al., *Nat Rev Cancer*. 2011; 11(11):805-812.; Pardoll D M., *Nat Rev Cancer*. 2012; 12(4):252-264), or bridge them to tumor cells (Topp M S, et al. *J Clin. Oncol.* 2011; 29(18):2493-2498) produced major clinical responses in patients with either solid tumors or leukemia. Finally, infusion of genetically-modified autologous T lymphocytes induced complete and durable remission in patients with refractory leukemia and lymphoma (Maude S L, et al. *N Engl J Med*. 2014; 371(16):1507-1517).

Nevertheless, there is a significant need for improving immune cell therapy by broadening its applicability and enhancing its efficacy.

SUMMARY OF THE INVENTION

The present invention relates to engineered immune cells having enhanced therapeutic efficacy for, e.g., cancer therapy. In certain embodiments, the present invention provides an engineered immune cell that comprises a nucleic acid comprising a nucleotide sequence encoding an immune

2

activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule linked to a localizing domain.

In other embodiments, the present invention provides the use of an engineered immune cell that comprises a gene encoding an immune activating receptor, and a gene encoding a target-binding molecule linked to a localizing domain for treating cancer, comprising administering a therapeutic amount of the engineered immune cell to a subject in need thereof.

In various embodiments, the present invention also provides a method for producing an engineered immune cell, the method comprising introducing into an immune cell a nucleic acid comprising a nucleotide sequence encoding an immune activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule linked to a localizing domain, thereby producing an engineered immune cell.

In some embodiments, the engineered immune cells possess enhanced therapeutic efficacy as a result of one or more of reduced graft-versus-host disease (GvHD) in a host, reduced or elimination of rejection by a host, extended survival in a host, reduced inhibition by the tumor in a host, reduced self-killing in a host, reduced inflammatory cascade in a host, or sustained natural/artificial receptor-mediated (e.g., CAR-mediated) signal transduction in a host.

BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing will be apparent from the following more particular description of example embodiments of the invention, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating embodiments of the present invention.

FIGS. 1A-1B is a schematic representation of a strategy employed in the present invention. FIG. 1A is an overall mechanism of CAR mediated killing of cancer cells. FIG. 1B shows the combined expression of CAR with different formats of compartment-directed scFv (an example of a target-binding molecule linked to a localizing domain) and examples of possible targets. The CAR can be replaced by other receptors that can enhance immune cell capacity.

FIG. 2 is a schematic diagram of constructs containing scFv together with domains that localize them to specific cellular compartments. Abbreviations: β 2M, β -2 microglobulin; SP, signal peptide; VL, variable light chain; VH, variable heavy chain; TM, transmembrane domain; HA, human influenza hemagglutinin. Additional constructs not listed in the figure include membrane-bound (mb) myc EEKKMP, mb myc KKTN, mb myc YQRL, mb TGN38 cytoplasmic domain, mb myc RNIKCD, linker(20-amino acid) mb EEKKMP, as well as variants of constructs without signaling peptide and with a varying number of amino acids in the CD8 transmembrane domain. The nucleotide sequence of the 10-amino acid linker is GGTGGTGGCGGCAGTGGTGGCGGTGGCTCA (SEQ ID NO: 61); the amino acid sequence is GGGGSGGGGS (SEQ ID NO: 62). The nucleotide sequence of the 20-amino acid linker is GGTGGTGGCGGCAGTGGTGGCGGTGGCTCAGGCGGTGGTGGCTCCGGTGGCGGT GGCTCT (SEQ ID NO: 63); the amino acid sequence is GGGGSGGGGS (SEQ ID NO: 41). Various localization domains are indicated under the heading “Localization domains,” and depicts linkers in some examples, as indicated. The constructs “myc KDEL” and

3

“PEST KDEL” show the use of more than one localization domains in a single construct.

FIGS. 3A-3C show downregulation of CD3/TCR in T cells by scFv targeting of CD3 ϵ . FIG. 3A shows expression of surface CD3 ϵ in Jurkat cells, transduced with either a retroviral vector containing green fluorescent protein (GFP) only (“mock”) or a vector containing GFP plus different constructs as indicated. Expression of CD3 ϵ on the cell membrane was compared to that of mock-transduced cells 1 week after transduction using an anti-CD3 antibody conjugated to allophycocyanin (BD Biosciences). All comparisons were performed after gating on GFP-positive cells. FIG. 3B depicts similar experiments performed with peripheral blood T lymphocytes expanded with anti-CD3/CD28 beads (Lifesciences). Staining was performed 1 week after transduction. FIG. 3C shows flow cytometry plots illustrating downregulation of membrane CD3 ϵ in Jurkat cells after transduction with the constructs indicated. Dashed rectangles on the upper right quadrant of each plot enclose GFP+ CD3+ cells.

FIG. 4 shows downregulation of CD3 ϵ and TC $\alpha\beta$ on the cell membrane in Jurkat T cells upon transduction with anti-CD3 ϵ scFv-KDEL or -PEST, or -mb EEKKMP. Membrane marker expression was measured 1 week after transduction using an anti-CD3 antibody conjugated to allophycocyanin (BD Biosciences) or an anti-TC $\alpha\beta$ conjugated to phycoerythrin (Biollegend). Lines labeled “Control” represent labelling of mock-transduced cells. Dashed vertical line represents the upper limit of staining obtained with an isotype-matched non-reactive antibody.

FIG. 5 shows that anti-scFv and CAR can be expressed simultaneously. Flow cytometric dot plots represent staining of Jurkat cells (top row) or peripheral blood lymphocytes (bottom row) with anti-CD3 allophycocyanin antibody and goat-anti-mouse Fab2 biotin plus streptavidin conjugated to phycoerythrin (to detect the CAR). Cells were transduced with the anti-CD3 scFv-myc KDEL construct, the anti-CD19-4-1BB-CD3 ζ construct, or both. After gating on GFP-positive cells, those transduced with anti-CD3 scFv-myc KDEL downregulated CD3 (left column, bottom left quadrants) and those transduced with the anti-CD19-4-1BB-CD3 ζ construct expressed the CAR (middle column, top right quadrants). A substantial proportion of cells transduced with both constructs were CD3-negative and CAR-positive (right column, top left quadrants).

FIG. 6 illustrates that anti-CD19 CAR triggers T cell activation and degranulation regardless of CD3/TCR downregulation. Jurkat cells were transduced with the anti-CD3 scFv-myc KDEL construct, the anti-CD19-4-1BB-CD3 ζ construct, or both. T cell activation and degranulation was compared to that of mock-transduced cells. Cells were co-cultured alone or with the CD19+ leukemia cell line OP-1 at a 1:1 ratio. After 18 hours, expression of CD69 and CD25 was tested by flow cytometry using specific antibodies (from BD Biosciences); expression of CD107a was tested after 6 hours (antibody from BD Biosciences). In the presence of OP-1 cells, CD69 and CD25 expression in CAR-expressing cells occurred regardless of whether cells were also transduced with anti-CD3 scFv-KDEL; no activation occurred in mock- or anti-CD3 scFv-myc KDEL transduced cells, or in the absence of OP-1 cells. CAR stimulation enhanced CD107 expression which was not affected by CD3/TCR downregulation.

FIG. 7 shows that anti-CD19 CAR expressed in T cells causes T cell proliferation regardless of CD3/TCR downregulation. Peripheral blood T lymphocytes were transduced with both the anti-CD3 scFv-myc KDEL construct and the

4

anti-CD19-4-1BB-CD3 ζ construct. Transduced T lymphocytes were co-cultured with OP-1 cells treated with Streck (Omaha, Nebr.) to inhibit their proliferation for the time indicated. Expansion of CD3-positive and CD3-negative T lymphocytes expressing the anti-CD19 CAR was compared to that of mock-transduced T cells. Each symbol shows the average cell count of two parallel cultures. CAR T cell expanded equally well regardless of CD3/TCR expression.

FIG. 8 shows expression of CD7 on the membrane of peripheral blood T lymphocytes transduced with either a retroviral vector containing GFP only (“mock”) or a vector containing GFP plus anti-CD7 scFv-myc KDEL construct. Expression of CD7 on the cell membrane was compared to that of mock-transduced cells 1 week after transduction using an anti-CD7 antibody conjugated to phycoerythrin (BD Biosciences). Dashed rectangles on the upper right quadrant of each plot enclose GFP+CD7+ cells.

FIG. 9 depicts the downregulation of HLA Class I in T cells by scFv targeting of β 2-microglobulin. Jurkat T cells were transduced with anti- β 2M scFv-myc KDEL. Expression of HLA-ABC on the cell membrane was compared to that of mock-transduced cells 1 week after transduction using an anti-HLA-ABC antibody conjugated to phycoerythrin (BD Biosciences). Staining with an isotype-matched control antibody is also shown. Analysis was performed after gating on GFP-positive cells.

FIG. 10 depicts the downregulation of Killer Immunoglobulin-like Receptor (KIR) 2DL1 and KIR2DL2/DL3 in human NK cells by scFv targeting of KIR2DL1 and KIR2DL2/DL3. NK cells, expanded ex vivo and selected for KIR2DL1 expression, were transduced with anti-KIR2DL1-KIR2DL2/DL3 scFv-linker (20) AEKDEL or -EEKKMP. Expression of the corresponding KIR on the cell membrane was compared to that of mock-transduced cells 8 days after transduction using an anti-KIR2DL1 antibody conjugated to allophycocyanin (R&D Systems) or an anti-KIR2DL2/DL3 antibody conjugated to phycoerythrin (BD Biosciences). Staining with an isotype-matched control antibody is also shown. Analysis was performed after gating on GFP-positive cells.

FIG. 11 depicts the downregulation of NKG2A in human NK cells by scFv targeting. NK cells, expanded ex vivo, were transduced with anti-NKG2A scFv-EEKKMP. Expression of NKG2A on the cell membrane was compared to that of mock-transduced cells 8 days after transduction using an NKG2A antibody conjugated to phycoerythrin (Beckman Coulter). Staining with an isotype-matched control antibody is also shown. Analysis was performed after gating on GFP-positive cells.

DETAILED DESCRIPTION OF THE INVENTION

A description of example embodiments of the invention follows.

In recent years, gains in knowledge about the molecular pathways that regulate immune cells have been paralleled by a remarkable evolution in the capacity to manipulate them ex vivo, including their expansion and genetic engineering. It is now possible to reliably prepare highly sophisticated clinical-grade immune cell products in a timely fashion. A prime example of how the anti-cancer activity of immune cells can be directed and magnified by ex vivo cell engineering is the development of chimeric antigen receptor (CAR) T cells (Eshhar, Z. et al., *PNAS*. 1993; 90(2):720-724).

CARs are artificial multi-molecular proteins, which have been previously described (Geiger T L, et al., *J Immunol.* 1999; 162(10):5931-5939; Brentjens R J, et al., *Nat Med.* 2003; 9(3):279-286; Cooper L J, et al., *Blood.* 2003; 101(4):1637-1644). CARs comprise an extracellular domain that binds to a specific target, a transmembrane domain, and a cytoplasmic domain. The extracellular domain and transmembrane domain can be derived from any desired source for such domains, as described in, e.g., U.S. Pat. No. 8,399,645, incorporated by reference herein in its entirety. Briefly, a CAR may be designed to contain a single-chain variable region (scFv) of an antibody that binds specifically to a target. The scFv may be linked to a T-cell receptor (TCR)-associated signaling molecule, such as CD3 ζ , via transmembrane and hinge domains. Ligation of scFv to the cognate antigen triggers signal transduction. Thus, CARs can instantaneously redirect cytotoxic T lymphocytes towards cancer cells and provoke tumor cell lysis (Eshhar, Z. et al., *PNAS.* 1993; 90(2):720-724; Geiger T L, et al., *J Immunol.* 1999; 162(10):5931-5939; Brentjens R J, et al., *Nat Med.* 2003; 9(3):279-286; Cooper L J, et al., *Blood.* 2003; 101(4):1637-1644; Imai C, et al., *Leukemia.* 2004; 18:676-684). Because CD3 ζ signaling alone is not sufficient to durably activate T cells (Schwartz R H. *Annu Rev Immunol.* 2003; 21:305-334; Zang X and Allison J P. *Clin Cancer Res.* 2007; 13(18 Pt 1):5271-5279), co-stimulatory molecules such as CD28 and 4-1BB (or CD137) have been incorporated into CAR constructs to boost signal transduction. This dual signaling design ("second generation CAR") is useful to elicit effective anti-tumor activity from T cells (Imai C, et al., *Leukemia.* 2004; 18:676-684; Campana D, et al., *Cancer J.* 2014; 20(2):134-140).

A specific CAR, anti-CD19 CAR, containing both 4-1BB and CD3 ζ has been described in U.S. Pat. No. 8,399,645. Infusion of autologous T cells expressing an anti-CD19-4-1BB-CD3 ζ CAR resulted in dramatic clinical responses in patients with chronic lymphocytic leukemia (CLL) (Porter D L, et al., Chimeric antigen receptor-modified T cells in chronic lymphoid leukemia; 2011: *N Engl J Med.* 2011; 365(8):725-733; Kalos M, et al., *Sci Transl Med.* 2011; 3(95):95ra73), and acute lymphoblastic leukemia (ALL) (Grupp S A, et al., *N Engl J Med.* 2013; 368(16):1509-1518; Maude S L, et al., *N Engl J Med.* 2014; 371(16):1507-1517). These studies, and studies with CARs bearing different signaling modules (Till B G, et al., *Blood.* 2012; 119(17):3940-3950; Kochenderfer J N, et al., *Blood.* 2012; 119(12):2709-2720; Brentjens R J, et al., *Blood.* 2011; 118(18):4817-4828; Brentjens R J, et al., *Sci Transl Med.* 2013; 5(177):177ra138), provide a convincing demonstration of the clinical potential of this technology, and of immunotherapy in general.

The methods described herein enable rapid removal or inactivation of specific proteins in immune cells redirected by a natural or artificial receptor, e.g., CARs, thus broadening the application potential and significantly improving the function of the engineered cells. The method relies, in part, on a single construct or multiple constructs containing an immune activating receptor, e.g., a CAR (which comprises an extracellular domain (e.g., an scFv) that binds to a specific target, a transmembrane domain, and a cytoplasmic domain) together with a target-binding molecule that binds a target (e.g., protein) to be removed or neutralized; the target-binding molecule is linked to a domain (i.e., localizing domain) that directs it to specific cellular compartments, such as the Golgi or endoplasmic reticulum, the proteasome, or the cell membrane, depending on the application. For

simplicity, a target-binding molecule linked to a localizing domain (LD) is sometimes referred to herein as "LD-linked target-binding molecule."

As will be apparent from the teachings herein, a variety of immune activating receptors may be suitable for the methods of the present invention. That is, any receptor that comprises a molecule that, upon binding (ligation) to a ligand (e.g., peptide or antigen) expressed on a cancer cell, is capable of activating an immune response may be used according to the present methods. For example, as described above, the immune activating receptor can be a chimeric antigen receptor (CAR); methods for designing and manipulating a CAR is known in the art (see, Geiger T L, et al., *J Immunol.* 1999; 162(10):5931-5939; Brentjens R J, et al., *Nat Med.* 2003; 9(3):279-286; Cooper L J, et al., *Blood.* 2003; 101(4):1637-1644). Additionally, receptors with antibody-binding capacity can be used (e.g., CD16-4-1BB-CD3 ζ receptor—Kudo K, et al. *Cancer Res.* 2014; 74(1):93-103), which are similar to CARs, but with the scFv replaced with an antibody-binding molecule (e.g., CD16, CD64, CD32). Further, T-cell receptors comprising T-cell receptor alpha and beta chains that bind to a peptide expressed on a tumor cell in the context of the tumor cell HLA can also be used according to the present methods. In addition, other receptors bearing molecules that activate an immune response by binding a ligand expressed on a cancer cell can also be used—e.g., NKG2D-DAP10-CD3 ζ receptor, which binds to NKG2D ligand expressed on tumor cells (see, e.g., Chang Y H, et al., *Cancer Res.* 2013; 73(6):1777-1786). All such suitable receptors collectively, as used herein, are referred to as an "immune activating receptor" or a "receptor that activates an immune response upon binding a cancer cell ligand." Therefore, an immune activating receptor having a molecule activated by a cancer cell ligand can be expressed together with a LD-linked target-binding molecule according to the present methods.

The present methods significantly expand the potential applications of immunotherapies based on the infusion of immune cells redirected by artificial receptors. The method described is practical and can be easily incorporated in a clinical-grade cell processing. For example, a single bicistronic construct containing, e.g., a CAR and a LD-linked target-binding molecule, e.g., scFv-myc KDEL (or PEST or transmembrane) can be prepared by inserting an internal ribosomal entry site (IRES) or a 2A peptide-coding region site between the 2 cDNAs encoding the CAR and the LD-linked target-binding molecule. The design of tricistronic delivery systems to delete more than one target should also be feasible. Alternatively, separate transductions of the 2 genes (simultaneously or sequentially) could be performed. In the context of cancer cell therapy, the CAR could be replaced by an antibody-binding signaling receptor (Kudo K, et al., *Cancer Res.* 2014; 74(1):93-103), a T-cell receptor directed against a specific HLA-peptide combination, or any receptor activated by contact with cancer cells (Chang Y H, et al., *Cancer Res.* 2013; 73(6):1777-1786). The results of the studies described herein with simultaneous anti-CD19-4-1BB-CD3 ζ CAR and anti-CD3 ϵ scFv-KDEL demonstrate that the signaling capacity of the CAR was not impaired.

Both the anti-CD3 ϵ scFv-KDEL (and -PEST) tested herein stably downregulate CD3 as well as TCR expression. Residual CD3+ T cells could be removed using CD3 beads, an approach that is also available in a clinical-grade format. The capacity to generate CD3/TCR-negative cells that respond to CAR signaling represents an important advance. Clinical studies with CAR T cells have generally been

performed using autologous T cells. Thus, the quality of the cell product varies from patient to patient and responses are heterogeneous. Infusion of allogeneic T cells is currently impossible as it has an unacceptably high risk of potentially fatal GvHD, due to the stimulation of the endogenous TCR by the recipient's tissue antigens. Downregulation of CD3/TCR opens the possibility of infusing allogeneic T cells because lack of endogenous TCR eliminates GvHD capacity. Allogeneic products could be prepared with the optimal cellular composition (e.g., enriched in highly cytotoxic T cells, depleted of regulatory T cells, etc.) and selected so that the cells infused have high CAR expression and functional potency. Moreover, fully standardized products could be cryopreserved and be available for use regardless of the patient immune cell status and his/her fitness to undergo apheresis or extensive blood draws. Removal of TCR expression has been addressed using gene editing tools, such as nucleases (Torikai H, et al. *Blood*, 2012; 119(24):5697-5705). Although this is an effective approach, it is difficult to implement in a clinical setting as it requires several rounds of cell selection and expansion, with prolonged culture. The methods described herein have considerable practical advantages.

Additionally, a LD-linked target-binding molecule (e.g., scFv-myc KDEL, scFv-EEKKMP or scFv-PEST, wherein scFv targets a specific protein/molecule) can be used according to the present invention to delete HLA Class I molecules, reducing the possibility of rejection of allogeneic cells. While infusion of allogeneic T cells is a future goal of CAR T cell therapy, infusion of allogeneic natural killer (NK) cells is already in use to treat patients with cancer. A key factor that determines the success of NK cell-based therapy is that NK cells must persist in sufficient numbers to achieve an effector:target ratio likely to produce tumor cyoreduction (Miller J S. *Hematology Am Soc Hematol Educ Program*. 2013; 2013:247-253). However, when allogeneic cells are infused, their persistence is limited. Immunosuppressive chemotherapy given to the patient allows transient engraftment of the infused NK cells but these are rejected within 2-4 weeks of infusion (Miller J S, et al. *Blood*. 2005; 105:3051-3057; Rubnitz J E, et al., *J Clin Oncol*. 2010; 28(6):955-959). Contrary to organ transplantation, continuing immunosuppression is not an option because immunosuppressive drugs also suppress NK cell function. Because rejection is primarily mediated by recognition of HLA Class I molecules by the recipient's CD8+ T lymphocytes, removing HLA Class I molecules from the infused NK cells (or T cells) will diminish or abrogate the rejection rate, extend the survival of allogeneic cells, and hence their anti-tumor capacity.

Furthermore, a LD-linked target-binding molecule can be used according to the present invention to target inhibitory receptors. Specifically, administration of antibodies that release T cells from inhibitory signals such as anti-PD1 or anti-CTLA-4 have produced dramatic clinical responses (Sharma P, et al., *Nat Rev Cancer*. 2011; 11(11):805-812; Pardoll D M. *Nat Rev Cancer*. 2012; 12(4):252-264). CAR-T cells, particularly those directed against solid tumors, might be inhibited by similar mechanisms. Thus, expression of a target-binding molecule (e.g., scFv or ligands) against PD1, CTLA-4, Tim3 or other inhibitory receptors would prevent the expression of these molecules (if linked to, e.g., KDEL (SEQ ID NO: 4), EEKKMP (SEQ ID NO: 64) or PEST motif SHGFPPVEEQDDGTLPMSCAQESGMDRHPAACASARINV (SEQ ID NO: 7)) or prevent binding of the receptors to their ligands (if linked to a transmembrane domain) and sustain CAR-mediated signal

transduction. In NK cells, examples of inhibitory receptors include killer immunoglobulin-like receptors (KIRs) and NKG2A (Vivier E, et al., *Science*, 2011; 331(6013):44-49).

The methods of the present invention also enable targeting of a greater number of targets amenable for CAR-directed T cell therapy. One of the main limitations of CAR-directed therapy is the paucity of specific antigens expressed by tumor cells. In the case of hematologic malignancies, such as leukemias and lymphomas, molecules which are not expressed in non-hematopoietic cells could be potential targets but cannot be used as CAR targets because they are also expressed on T cells and/or NK cells. Expressing such CARs on immune cells would likely lead to the demise of the immune cells themselves by a "fratricidal" mechanism, nullifying their anti-cancer capacity. If the target molecule can be removed from immune cells without adverse functional effects, then the CAR with the corresponding specificity can be expressed. This opens many new opportunities to target hematologic malignancies. Examples of the possible targets include CD38 expressed in multiple myeloma, CD7 expressed in T cell leukemia and lymphoma, Tim-3 expressed in acute leukemia, CD30 expressed in Hodgkin disease, CD45 and CD52 expressed in all hematologic malignancies. These molecules are also expressed in a substantial proportion of T cells and NK cells.

Moreover, it has been shown that secretion of cytokines by activated immune cells triggers cytokine release syndrome and macrophage activation syndrome, presenting serious adverse effects of immune cell therapy (Lee D W, et al., *Blood*. 2014; 124(2):188-195). Thus, the LD-linked target-binding molecule can be used according to the present invention to block cytokines such as IL-6, IL-2, IL-4, IL-7, IL-10, IL-12, IL-15, IL-18, IL-21, IL-27, IL-35, interferon (IFN)- γ , IFN- β , IFN- α , tumor necrosis factor (TNF)- α , and transforming growth factor (TGF)- β , which may contribute to such inflammatory cascade.

Accordingly, in one embodiment, the present invention relates to an engineered immune cell that comprises a nucleic acid comprising a nucleotide sequence encoding an immune activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule linked to a localizing domain.

As used herein, an "engineered" immune cell includes an immune cell that has been genetically modified as compared to a naturally-occurring immune cell. For example, an engineered T cell produced according to the present methods carries a nucleic acid comprising a nucleotide sequence that does not naturally occur in a T cell from which it was derived. In some embodiments, the engineered immune cell of the present invention includes a chimeric antigen receptor (CAR) and a target-binding molecule linked to a localizing domain (LD-linked target-binding molecule). In a particular embodiment, the engineered immune cell of the present invention includes an anti-CD19-4-1BB-CD3 ζ CAR and an anti-CD3 scFv linked to a localizing domain.

In certain embodiments, the engineered immune cell is an engineered T cell, an engineered natural killer (NK) cell, an engineered NK/T cell, an engineered monocyte, an engineered macrophage, or an engineered dendritic cell.

In certain embodiments, an "immune activating receptor" as used herein refers to a receptor that activates an immune response upon binding a cancer cell ligand. In some embodiments, the immune activating receptor comprises a molecule that, upon binding (ligation) to a ligand (e.g., peptide or antigen) expressed on a cancer cell, is capable of activating an immune response. In one embodiment, the immune activating receptor is a chimeric antigen receptor (CAR);

methods for designing and manipulating a CAR are known in the art. In other embodiments, the immune activating receptor is an antibody-binding receptor, which is similar to a CAR, but with the scFv replaced with an antibody-binding molecule (e.g., CD16, CD64, CD32) (see e.g., CD16-4-1BB-CD3zeta receptor—Kudo K, et al. *Cancer Res.* 2014; 74(1):93-103). In various embodiments, T-cell receptors comprising T-cell receptor alpha and beta chains that bind to a peptide expressed on a tumor cell in the context of the tumor cell HLA can also be used according to the present methods. In certain embodiments, other receptors bearing molecules that activate an immune response by binding a ligand expressed on a cancer cell can also be used—e.g., NKG2D-DAP10-CD3zeta receptor, which binds to NKG2D ligand expressed on tumor cells (see, e.g., Chang Y H, et al., *Cancer Res.* 2013; 73(6):1777-1786). All such suitable receptors capable of activating an immune response upon binding (ligation) to a ligand (e.g., peptide or antigen) expressed on a cancer cell are collectively referred to as an “immune activating receptor.” As would be appreciated by those of skill in the art, an immune activating receptor need not contain an antibody or antigen-binding fragment (e.g., scFv); rather the portion of the immune activating receptor that binds to a target molecule can be derived from, e.g., a receptor in a receptor-ligand pair, or a ligand in a receptor-ligand pair.

In certain aspects, the immune activating receptor binds to molecules expressed on the surface of tumor cells, including but not limited to, CD20, CD22, CD33, CD2, CD3, CD4, CD5, CD7, CD8, CD45, CD52, CD38, CS-1, TIM3, CD123, mesothelin, folate receptor, HER2-neu, epidermal-growth factor receptor, and epidermal growth factor receptor. In some embodiments, the immune activating receptor is a CAR (e.g., anti-CD19-4-1BB-CD3ζ CAR). In certain embodiments, the immune activating receptor comprises an antibody or antigen-binding fragment thereof (e.g., scFv) that binds to molecules expressed on the surface of tumor cells, including but not limited to, CD20, CD22, CD33, CD2, CD3, CD4, CD5, CD7, CD8, CD45, CD52, CD38, CS-1, TIM3, CD123, mesothelin, folate receptor, HER2-neu, epidermal-growth factor receptor, and epidermal growth factor receptor. Antibodies to such molecules expressed on the surface of tumor cells are known and available in the art. By way of example, antibodies to CD3 and CD7 are commercially available and known in the art. Such antibodies, as well as fragments of antibodies (e.g., scFv) derived therefrom, can be used in the present invention, as exemplified herein. Further, methods of producing antibodies and antibody fragments against a target protein are well-known and routine in the art.

The transmembrane domain of an immune activating receptor according to the present invention (e.g., CAR) can be derived from a single-pass membrane protein, including, but not limited to, CD8α, CD8β, 4-1BB, CD28, CD34, CD4, FcεRIγ, CD16 (e.g., CD16A or CD16B), OX40, CD3ζ, CD38, CD3γ, CD3δ, TCRα, CD32 (e.g., CD32A or CD32B), CD64 (e.g., CD64A, CD64B, or CD64C), VEGFR2, FAS, and FGFR2B. In some examples, the membrane protein is not CD8α. The transmembrane domain may also be a non-naturally occurring hydrophobic protein segment.

The hinge domain of the immune activating receptor (e.g., CAR) can be derived from a protein such as CD8α, or IgG. The hinge domain can be a fragment of the transmembrane or hinge domain of CD8α, or a non-naturally occurring peptide, such as a polypeptide consisting of hydrophilic

residues of varying length, or a (GGGGS)_n (SEQ ID NO: 8) polypeptide, in which n is an integer of, e.g., 3-12, inclusive.

The signaling domain of the immune activating receptor (e.g., CAR) can be derived from CD3ζ, FcεRIγ, DAP10, DAP12 or other molecules known to deliver activating signals in immune cells. At least one co-stimulatory signaling domain of the receptor can be a co-stimulatory molecule such as 4-1BB (also known as CD137), CD28, CD28_{LL→GG} variant, OX40, ICOS, CD27, GITR, HVEM, TIM1, LFA1, or CD2. Such molecules are readily available and known in the art.

As would be appreciated by those of skill in the art, the components of an immune activating receptor can be engineered to comprise a number of functional combinations, as described herein, to produce a desired result. Using the particular CAR anti-CD19-4-1BB-CD3ζ as an example, the antibody (e.g., or antigen-binding fragment thereof such as an scFv) that binds a molecule can be substituted for an antibody that binds different molecule, as described herein (e.g., anti-CD20, anti-CD33, anti-CD123, etc., instead of anti-CD19). In other embodiments, the co-stimulatory molecule (4-1BB in this specific example) can also be varied with a different co-stimulatory molecule, e.g., CD28. In some embodiments, the stimulatory molecule (CD3ζ in this specific example), can be substituted with another known stimulatory molecule. In various embodiments, the transmembrane domain of the receptor can also be varied as desired. The design, production, and testing for functionality of such immune activating receptors can be readily determined by those of skill in the art. Similarly, the design, delivery into cells and expression of nucleic acids encoding such immune activating receptors are readily known and available in the art.

As used herein, the term “nucleic acid” refers to a polymer comprising multiple nucleotide monomers (e.g., ribonucleotide monomers or deoxyribonucleotide monomers). “Nucleic acid” includes, for example, genomic DNA, cDNA, RNA, and DNA-RNA hybrid molecules. Nucleic acid molecules can be naturally occurring, recombinant, or synthetic. In addition, nucleic acid molecules can be single-stranded, double-stranded or triple-stranded. In some embodiments, nucleic acid molecules can be modified. In the case of a double-stranded polymer, “nucleic acid” can refer to either or both strands of the molecule.

The term “nucleotide sequence,” in reference to a nucleic acid, refers to a contiguous series of nucleotides that are joined by covalent linkages, such as phosphorus linkages (e.g., phosphodiester, alkyl and aryl-phosphonate, phosphorothioate, phosphotriester bonds), and/or non-phosphorus linkages (e.g., peptide and/or sulfamate bonds). In certain embodiments, the nucleotide sequence encoding, e.g., a target-binding molecule linked to a localizing domain is a heterologous sequence (e.g., a gene that is of a different species or cell type origin).

The terms “nucleotide” and “nucleotide monomer” refer to naturally occurring ribonucleotide or deoxyribonucleotide monomers, as well as non-naturally occurring derivatives and analogs thereof. Accordingly, nucleotides can include, for example, nucleotides comprising naturally occurring bases (e.g., adenosine, thymidine, guanosine, cytidine, uridine, inosine, deoxyadenosine, deoxythymidine, deoxyguanosine, or deoxycytidine) and nucleotides comprising modified bases known in the art.

As will be appreciated by those of skill in the art, in some aspects, the nucleic acid further comprises a plasmid sequence. The plasmid sequence can include, for example,

one or more sequences selected from the group consisting of a promoter sequence, a selection marker sequence, and a locus-targeting sequence.

As used herein, the gene encoding a target-binding molecule linked to a localizing domain is sometimes referred to as "LD-linked target-binding molecule."

In certain embodiments, the target-binding molecule is an antibody or antigen-binding fragment thereof. As used herein, "antibody" means an intact antibody or antigen-binding fragment of an antibody, including an intact antibody or antigen-binding fragment that has been modified or engineered, or that is a human antibody. Examples of antibodies that have been modified or engineered are chimeric antibodies, humanized antibodies, multiparatopic antibodies (e.g., biparatopic antibodies), and multispecific antibodies (e.g., bispecific antibodies). Examples of antigen-binding fragments include Fab, Fab', F(ab')₂, Fv, single chain antibodies (e.g., scFv), minibodies and diabodies.

A "Fab fragment" comprises one light chain and the C_H1 and variable regions of one heavy chain. The heavy chain of a Fab molecule cannot form a disulfide bond with another heavy chain molecule.

An "Fc" region contains two heavy chain fragments comprising the CH2 and CH3 domains of an antibody. The two heavy chain fragments are held together by two or more disulfide bonds and by hydrophobic interactions of the CH3 domains.

A "Fab' fragment" contains one light chain and a portion of one heavy chain that contains the VH domain and the CH1 domain and also the region between the CH1 and CH2 domains, such that an interchain disulfide bond can be formed between the two heavy chains of two Fab' fragments to form a F(ab')₂ molecule.

A "F(ab')₂ fragment" contains two light chains and two heavy chains containing a portion of the constant region between the C_H1 and C_H2 domains, such that an interchain disulfide bond is formed between the two heavy chains. A F(ab')₂ fragment thus is composed of two Fab' fragments that are held together by a disulfide bond between the two heavy chains.

The "Fv region" comprises the variable regions from both the heavy and light chains, but lacks the constant regions.

In a particular embodiment, the target-binding molecule is single-chain Fv antibody ("scFv antibody"). scFv refers to antibody fragments comprising the VH and VL domains of an antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv, see Pluckthun (1994) *The Pharmacology Of Monoclonal Antibodies*, vol. 113, Rosenburg and Moore eds. Springer-Verlag, New York, pp. 269-315. See also, PCT Publication No. WO 88/01649 and U.S. Pat. Nos. 4,946,778 and 5,260,203. By way of example, the linker between the VH and VL domains of the scFvs disclosed herein comprise, e.g., GGGGSGGGSGGGSGGGGS (SEQ ID NO: 41) or GGGGSGGGSGGGGS (SEQ ID NO: 43). As would be appreciated by those of skill in the art, various suitable linkers can be designed and tested for optimal function, as provided in the art, and as disclosed herein.

The scFv that is part of the LD-linked target-binding molecule is not necessarily the same as the scFv that occurs in the context of, e.g., a chimeric antigen receptor (CAR) or a similar antibody-binding signaling receptor. In some

context of, e.g., a chimeric antigen receptor (CAR) or a similar antibody-binding signaling receptor.

In some embodiments, the nucleic acid comprising a nucleotide sequence encoding a target-binding molecule (e.g., an scFv in the context of a LD-linked target-binding molecule) comprises one or more sequences that have at least 80%, at least 85%, at least 88%, at least 90%, at least 92%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to any one or more of SEQ ID NOs: 14, 15, 18, 19, 22, 23, 26, 27, 30, 31, 34, 35, 38, or 39.

The term "sequence identity" means that two nucleotide or amino acid sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least, e.g., 70% sequence identity, or at least 80% sequence identity, or at least 85% sequence identity, or at least 90% sequence identity, or at least 95% sequence identity or more. For sequence comparison, typically one sequence acts as a reference sequence (e.g., parent sequence), to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally Ausubel et al., *Current Protocols in Molecular Biology*). One example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (publicly accessible through the National Institutes of Health NCBI internet server). Typically, default program parameters can be used to perform the sequence comparison, although customized parameters can also be used. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)).

In certain embodiments, the antibody (e.g., scFv) comprises VH and VL having amino acid sequences set forth in SEQ ID NO: 12 and 13, respectively; SEQ ID NO: 16 and 17, respectively; SEQ ID NO: 20 and 21, respectively; SEQ ID NO: 24 and 25, respectively; SEQ ID NO: 28 and 29, respectively; SEQ ID NO: 32 and 33, respectively; or SEQ ID NO: 36 and 37, respectively. In some embodiments, the antibody (e.g., scFv) comprises VH and VL having sequence that each have at least 90% sequence identity, at least 91% sequence identity, at least 92% sequence identity, at least 93% sequence identity, at least 94% sequence identity, at least 95% sequence identity, at least 96% sequence identity, at least 97% sequence identity, at least 98% sequence identity, at least 99% sequence identity, or 100% sequence identity to the VH and VL sequences set forth in SEQ ID

13

NO: 12 and 13, respectively; SEQ ID NO: 16 and 17, respectively; SEQ ID NO: 20 and 21, respectively; SEQ ID NO: 24 and 25, respectively; SEQ ID NO: 28 and 29, respectively; SEQ ID NO: 32 and 33, respectively; or SEQ ID NO: 36 and 37, respectively.

A “diabody” is a small antibody fragment with two antigen-binding sites. The fragments comprise a heavy chain variable region (VH) connected to a light chain variable region (VL) in the same polypeptide chain (VH-VL or VL-VH). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described in, e.g., patent documents EP 404,097; WO 93/11161; and Holliger et al., (1993) *Proc. Nat. Acad. Sci. USA* 90: 6444-6448.

In certain embodiments, the antibody is a triabody or a tetrabody. Methods of designing and producing triabodies and tetrabodies are known in the art. See, e.g., Todorovska et al., *J. Immunol. Methods* 248(1-2):47-66, 2001.

A “domain antibody fragment” is an immunologically functional immunoglobulin fragment containing only the variable region of a heavy chain or the variable region of a light chain. In some instances, two or more VH regions are covalently joined with a peptide linker to create a bivalent domain antibody fragment. The two VH regions of a bivalent domain antibody fragment may target the same or different antigens.

In some embodiments, the antibody is modified or engineered. Examples of modified or engineered antibodies include chimeric antibodies, multiparatopic antibodies (e.g., biparatopic antibodies), and multispecific antibodies (e.g., bispecific antibodies).

As used herein, “multiparatopic antibody” means an antibody that comprises at least two single domain antibodies, in which at least one single domain antibody is directed against a first antigenic determinant on an antigen and at least one other single domain antibody is directed against a second antigenic determinant on the same antigen. Thus, for example, a “biparatopic” antibody comprises at least one single domain antibody directed against a first antigenic determinant on an antigen and at least one further single domain antibody directed against a second antigenic determinant on the same antigen.

As used herein, “multispecific antibody” means an antibody that comprises at least two single domain antibodies, in which at least one single domain antibody is directed against a first antigen and at least one other single domain antibody is directed against a second antigen (different from the first antigen). Thus, for example, a “bispecific” antibody is one that comprises at least one single domain antibody directed against a first antigen and at least one further single domain antibody directed against a second antigen, e.g., different from the first antigen.

In some embodiments, the antibodies disclosed herein are monoclonal antibodies, e.g., murine monoclonal antibodies. Methods of producing monoclonal antibodies are known in the art. See, for example, Pluckthun (1994) *The Pharmacology of Monoclonal Antibodies*, Vol. 113, Rosenberg and Moore eds. Springer-Verlag, New York, pp. 269-315.

In various embodiments, the target-binding molecule in the context of a LD-linked target-binding molecule is a receptor or a ligand that binds to a target molecule. For example, that target-binding molecule can be a ligand that binds PD-1 (e.g., PD-L1 or PD-L2). Thus, as would be

14

appreciated by those of skill in the art, the target-binding molecule can be an antibody, or a ligand/receptor that binds a target molecule.

As used herein, “linked” in the context of a LD-linked target-binding molecule refers to a gene encoding a target-binding molecule directly in frame (e.g., without a linker) adjacent to one or more genes encoding one or more localizing domains. Alternatively, the gene encoding a target-binding molecule may be connected to one or more gene encoding one or more localizing domains through a linker sequence, as described herein. Various suitable linkers known in the art can be used to tether the target-binding molecule to a localizing domain. For example, non-naturally occurring peptides, such as a polypeptide consisting of hydrophilic residues of varying length, or a (GGGGS)_n (SEQ ID NO: 8) polypeptide, in which n is an integer of, e.g., 3-12, inclusive, can be used according to the present invention. In particular embodiments, the linker comprises, e.g., GGGGSGGGGS (SEQ ID NO: 62). In some embodiments, the linker comprises, e.g., GGGGSGGGGS (SEQ ID NO: 41). In various embodiments, peptide linkers having lengths of about 5 to about 100 amino acids, inclusive, can be used in the present invention. In certain embodiments, peptide linkers having lengths of about 20 to about 40 amino acids, inclusive, can be used in the present invention. In some embodiments, peptide linkers having lengths of at least 5 amino acids, at least 10 amino acids, at least 15 amino acids, at least 20 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, or at least 40 amino acids can be used in the present invention. As would be appreciated by those of skill in the art, such linker sequences as well as variants of such linker sequences are known in the art. Methods of designing constructs that incorporate linker sequences as well as methods of assessing functionality are readily available to those of skill in the art.

In certain embodiments, the LD-linked target-binding molecule binds to a target expressed on the surface of an immune cell. In some embodiments, the LD-linked target-binding molecule inhibits the activity or function of the target molecule. By way of example, as disclosed herein, the LD-linked target-binding molecule can be designed to bind to, e.g., CD3, CD7, CD45, hB2MG, KIR2DLT, KIR2DL2/DL3, or NKG2A, thereby downregulating the cell surface expression of such molecules. Downregulation of such molecules can be achieved through, for example, localizing/targeting the molecules for degradation and/or internalization. In other embodiments, the LD-linked target-binding molecule renders the target inactive (e.g., the target can no longer interact and/or bind to its cognate ligand or receptor).

In some embodiments, the engineered immune cells of the present invention have enhanced therapeutic efficacy. As used herein, “enhanced therapeutic efficacy” refers to one or more of reduced graft-versus-host disease (GvHD) in a host, reduced or elimination of rejection by a host, extended survival in a host, reduced inhibition by the tumor in a host, reduced self-killing in a host, reduced inflammatory cascade in a host, or sustained CAR-mediated signal transduction in a host.

In certain embodiments of the present invention, the target-binding molecule in the context of a LD-linked target-binding molecule binds to a molecule in a CD3/T-cell receptor (TCR) complex, a cytokine, a human leukocyte antigen (HLA) Class I molecule, or a receptor that down-regulates immune response.

15

In certain embodiments, a molecule in a CD3/TCR complex can be CD3 ϵ , TCR α , TCR β , TCR γ , TCR δ , CD3 δ , CD3 γ , or CD3 ζ . In a particular embodiment, the molecule is CD3 ϵ .

In another embodiment, the HLA Class I molecule is beta-2 microglobulin, α 1-microglobulin, α 2-microglobulin, or α 3-microglobulin.

In other embodiments, a receptor that downregulates immune response is selected from, e.g., PD-1, CTLA-4, Tim3, killer immunoglobulin-like receptors (KIRs—e.g., KIR2DL1 (also known as CD158a), KIR2DL2/DL3 (also known as CD158b)), CD94 or NKG2A (also known as CD159a), protein tyrosine phosphatases such as Src homology region 2 domain-containing phosphatase (SHP)-1 and SHP-2. Thus, such receptors can be targeted by moiety LD-linked target-binding molecule, as described herein.

In various embodiments, examples of cytokines that can be targeted with moiety LD-linked target-binding molecule include, e.g., interleukin (IL)-6, IL-2, IL-4, IL-7, IL-10, IL-12, IL-15, IL-18, IL-21, IL-27, IL-35, interferon (IFN)- γ , IFN- β , IFN- α , tumor necrosis factor (TNF)- α , or transforming growth factor (TGF)- β .

In a further aspect, the LD-linked target-binding molecule binds to a molecule selected from, e.g., CD2, CD4, CD5, CD7, CD8, CD30, CD38, CD45, CD52, or CD127.

Methods of producing antibodies and antibody fragments thereof against any target protein are well-known and routine in the art. Moreover, as exemplified herein, commercially available antibodies to various targets, e.g., CD3 and CD7 can be used to generate a LD-linked target-binding molecule, as exemplified herein. Antibodies known in the art, as well as fragments of antibodies (e.g., scFv) derived therefrom, can be used in the present invention, as exemplified herein.

In other aspects, the localizing domain of the LD-linked target-binding molecule comprises an endoplasmic reticulum (ER) retention sequence KDEL (SEQ ID NO: 4), or other ER or Golgi retention sequences such as KKXX (SEQ ID NO: 9), KXD/E (SEQ ID NO: 10) (where X can be any amino acid—see Gao C, et al., *Trends in Plant Science* 19: 508-515, 2014) and YQRL (SEQ ID NO: 11) (see Zhan J, et al., *Cancer Immunol Immunother* 46:55-60, 1998); a proteosome targeting sequence that comprises, e.g., “PEST” motif—SHGFPPEVEEQDDGTLPMSCAQESGMDRH-PAACASARINV (SEQ ID NO: 7); and/or a sequence that targets the target-binding molecule to the cell membrane, such as the CD8a transmembrane domain, or the transmembrane of another single-pass membrane protein, as described herein (e.g., CD8 α , CD8 β , 4-1BB, CD28, CD34, CD4, Fc ϵ R1 γ , CD16 (such as CD16A or CD16B), OX40, CD3 ζ , CD38, CD3 γ , CD3 δ , TCR α , CD32 (such as CD32A or CD32B), CD64 (such as CD64A, CD64B, or CD64C), VEGFR2, FAS, or FGFR2B). Examples of particular localizing domains (sequences) exemplified herein are shown in FIG. 2. Various other localizing sequences are known and available in the art.

As shown in FIG. 2, the LD-linked target-binding molecules of the present invention can comprise one or more localizing domains. For example, the LD-linked target-binding molecule can have at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten localizing domains linked together. When more than one localizing domain is used in a given LD-linked target-binding molecule, each localizing domain can be linked with or without any intervening linker. By way of example, as shown in FIG. 2, localization domains CD8 TM, PEST motif, and EEKMP

16

can be used in a single LD-linked target-binding molecule. While this particular construct shows the localization domains without any intervening linkers, various intervening linkers can be incorporated between some or all of the localization domains. Other examples are shown in FIG. 2.

As would be appreciated by those of skill in the art, the immune activating receptor and/or the LD-linked target-binding molecule can be designed to bind to the targets disclosed herein, as well as variants of the targets disclosed herein. By way of example, an immune activating receptor and/or the LD-linked target-binding molecule can be designed to bind to a molecule in a CD3/TCR complex, or a naturally-occurring variant molecule thereof. Such naturally-occurring variants can have the same function as the wild-type form of the molecule. In other embodiments, the variant can have a function that is altered relative to the wild-type form of the molecule (e.g., confers a diseased state).

As would be appreciated by those of skill in the art, the various components of the LD-linked target-binding molecule constructs shown in FIG. 2 can be substituted in different combinations (e.g., to contain a different linker, different localizing sequence, different scFv, etc.), so long as the combination produces a functional LD-linked target-binding molecule. Methods of assessing functionality for a particular construct are within the ambit of those of skill in the art, as disclosed herein.

In further aspects, the present invention relates to the use of an engineered immune cell that comprises a nucleic acid comprising a nucleotide sequence encoding an immune activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule (e.g., scFv) linked to a localizing domain for treating cancer, comprising administering a therapeutic amount of the engineered immune cell to a subject in need thereof.

In another aspect, the present invention relates to the use of an engineered immune cell that comprises a nucleic acid comprising a nucleotide sequence encoding a chimeric antigen receptor (CAR) and a nucleic acid comprising a nucleotide sequence encoding a single-chain variable fragment (scFv) linked to a localizing domain for treating cancer, comprising administering a therapeutic amount of the engineered immune cell to a subject in need thereof.

In other aspects, the present invention relates to the use of an engineered immune cell that comprises a nucleic acid comprising a nucleotide sequence encoding an immune activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule (e.g., scFv) linked to a localizing domain for treating an autoimmune disorder, comprising administering a therapeutic amount of the engineered immune cell to a subject in need thereof.

In other aspects, the present invention also relates to the use of an engineered immune cell that comprises a nucleic acid comprising a nucleotide sequence encoding an immune activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule (e.g., scFv) linked to a localizing domain for treating an infectious disease, comprising administering a therapeutic amount of the engineered immune cell to a subject in need thereof.

In various embodiments, the immune activating receptor is a CAR (e.g., anti-CD19-4-1BB-CD3 ζ CAR).

In other embodiments, the single-chain variable fragment (scFv) linked to a localizing domain is selected from any one or more constructs shown in FIG. 2.

In some aspects, the engineered immune cell is administered by infusion into the subject. Methods of infusing

immune cells (e.g., allogeneic or autologous immune cells) are known in the art. A sufficient number of cells are administered to the recipient in order to ameliorate the symptoms of the disease. Typically, dosages of 10^7 to 10^{10} cells are infused in a single setting, e.g., dosages of 10^9 cells. Infusions are administered either as a single 10^9 cell dose or divided into several 10^9 cell dosages. The frequency of infusions can be every 3 to 30 days or even longer intervals if desired or indicated. The quantity of infusions is generally at least 1 infusion per subject and preferably at least 3 infusions, as tolerated, or until the disease symptoms have been ameliorated. The cells can be infused intravenously at a rate of 50-250 ml/hr. Other suitable modes of administration include intra-arterial infusion, direct injection into tumor and/or perfusion of tumor bed after surgery, implantation at the tumor site in an artificial scaffold, intrathecal administration, and intraocular administration. Methods of adapting the present invention to such modes of delivery are readily available to one skilled in the art.

In certain aspects, the cancer to be treated is a solid tumor or a hematologic malignancy. Examples of hematologic malignancies include acute myeloid leukemia, chronic myelogenous leukemia, myelodysplasia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, multiple myeloma, Hodgkin and non-Hodgkin lymphoma. Examples of solid tumors include lung cancer, melanoma, breast cancer, prostate cancer, colon cancer, renal cell carcinoma, ovarian cancer, pancreatic cancer, hepatocellular carcinoma, neuroblastoma, rhabdomyosarcoma, brain tumor.

In another embodiment, the present invention relates to a method for producing an engineered immune cell of the present invention, comprising introducing into an immune cell a nucleic acid comprising a nucleotide sequence encoding an immune activating receptor, and a nucleic acid comprising a nucleotide sequence encoding a target-binding molecule linked to a localizing domain, thereby producing an engineered immune cell.

In certain embodiments, the nucleic acid comprising a nucleotide sequence is introduced into an immune cell *ex vivo*. In other embodiments, the nucleic acid comprising a nucleotide sequence is introduced into an immune cell *in vivo*.

In some embodiments, an "immune cell" includes, e.g., a T cell, a natural killer (NK) cell, an NK/T cell, a monocyte, a macrophage, or a dendritic cell.

The nucleic acid comprising a nucleotide sequence to be introduced can be a single bicistronic construct containing an immune activating receptor described herein and a target-binding molecule (e.g., scFv) linked to a localizing domain. As described herein, a single bicistronic construct can be prepared by inserting an internal ribosomal entry site (IRES) or a 2A peptide-coding region site between the 2 cDNAs encoding the immune activating receptor as described herein (e.g., CAR) and the target-binding molecule (e.g., scFv). The design of tricistronic delivery systems to delete more than one target should also be feasible. Alternatively, separate transductions (simultaneously or sequentially) of the individual constructs (e.g., CAR and LD-linked target-binding molecule) could be performed. Methods of introducing exogenous nucleic acids are exemplified herein, and are well-known in the art.

As used herein, the indefinite articles "a" and "an" should be understood to mean "at least one" unless clearly indicated to the contrary.

EXEMPLIFICATION

Methods

Cloning of scFv from Mouse Anti-Human CD3 Hybridoma PLU4 hybridoma cells secreting an anti-human CD3 monoclonal antibody (IgG2a isotype; Creative Diagnostics,

Shirley, N.Y.) were cultured in IMDM plus GlutaMAX medium (Life Technologies, Carlsbad, Calif.) with 20% fetal bovine serum (Thermo Fisher Scientific, Waltham, Mass.) and antibiotics. Total RNA was extracted using TRIzol reagent (Life Technologies), and cDNA was synthesized by M-MLV reverse transcriptase (Promega, Madison, Wis.) and Oligo(dT)₁₅ primer (Promega). IgG Library Primer Set Mouse BioGenomics (US Biological, Salem, Mass.) was used to amplify the variable region of heavy chain (VH) and light chain (VL); PCR products were cloned into TOPO TA cloning kit for sequencing (Life Technologies). The VH and VL genes were assembled into scFv by a flexible linker sequence which encodes (Gly₄Ser)₄ using splicing by overlapping extension-PCR. Signal peptide domain of CD8 α was subcloned by PCR using cDNA derived from human activated T cell from healthy donor, and connected to 5' end of the VL fragment. The Myc tag (EQKLISEEDL; SEQ ID NO: 1) was added to C-terminus of VH by PCR using sense primer: 5'-ATATATGAATTCGGCTTCCACCATGGCCTTACCAGTGACC-3' (SEQ ID NO: 2) and reverse primer: 5'-CAGATCTTCTTCAGAAATAAGTTTTGTTCGGCTTGAGGAGACTGTGAGAG-3' (SEQ ID NO: 3). Also the KDEL (SEQ ID NO: 4) coding sequence was generated after Myc tag by sense primer: 5'-ATATATGAATTCGGCTTCCACCATGGCCTTACCAGTGACC-3' (SEQ ID NO: 5) and reverse primer: 5'-TATATACTCGAGTTACAACCTCGTCCTTCAGATCTTCTTCAGAAATAAG-3' (SEQ ID NO: 6). The synthesized gene consisting of CD8 signal peptide, scFv against human CD3, Myc tag and KDEL (SEQ ID NO: 4) sequence was subcloned into EcoRI and XhoI sites of the MSCV-IRES-GFP vector. Constructs in which myc-KDEL was replaced by other sequences were also made as listed in FIG. 2.

The sequence of "PEST"—SHGFPEVE-EQDDGTLPMSCAQESGMDRHPAACASARINV (SEQ ID NO: 7) motif corresponding to amino acids 422-461 of mouse ornithine decarboxylase was obtained from GenBank (accession number NM_013614.2). Codon optimization and gene synthesis was done by GenScript (Piscataway, N.J.), and subcloned into the 3' end of VH by PCR. The constructs were subcloned into EcoRI and XhoI sites of the MSCV-IRES-GFP vector.

Cloning of scFv Against Human CD7

The sequence scFv derived from murine TH69 (anti-CD7) antibody was obtained from literature (Peipp et al., *Cancer Res* 2002 (62): 2848-2855). After codon optimization, the synthesized gene consisting of CD8 signal peptide, scFv against human CD7, Myc tag and KDEL (SEQ ID NO: 4) sequence was subcloned into EcoRI and XhoI sites of the MSCV-IRES-GFP vector. Constructs in which myc-KDEL was replaced by other sequences were also made as listed in FIG. 2.

Cloning of scFv Against Human Beta-2 Microglobulin (hB2M G)

The sequence scFv derived from murine BBM.1 (anti-hB2M G) IgG2b antibody was obtained from literature (Grosvender, E. A. et al., *Kidney Int.* 2004; 65(1):310-322). After codon optimization, synthesized gene consists of CD8 signal peptide, scFv against human B2MG, Myc tag and KDEL (SEQ ID NO: 4) sequence was subcloned into EcoRI and XhoI sites of the MSCV-IRES-GFP vector.

Cloning of scFv Against Human KIR2DL1 and KIR2DL2/DL3

The amino acid sequence of human monoclonal antibody I-7F9 (anti-KIR2DL1, KIR2DL2, and KIR2DL3) was

19

derived from published International Patent Application WO2006003179 A2 by Moretta et al. After codon optimization, the sequence of scFv was designed by connecting variable light (VL) region and variable heavy (VH) region with linker sequence. The synthesized gene consisting of CD8 signal peptide, scFv against human KIRs (KIR2DL1, KIR2DL2 and KIR2DL3), CD8 hinge and transmembrane domain, and KKMP sequence was subcloned into EcoRI and XhoI sites of the MSCV-IRES-GFP vector. Constructs in which KKMP was replaced by other sequences were also made as listed in FIG. 2.

Cloning of scFv Against Human NKG2A

The sequence of murine antibody Z199 (anti-NKG2A) was derived from the published patent by Spee et al. (EP2247619 A1). After codon optimization, the sequence of

20

scFv was designed by connecting variable light (VL) region and variable heavy (VH) region with linker sequence. The synthesized gene consisting of CD8 signal peptide, scFv against human NKG2A, CD8 hinge and transmembrane, and KKMP sequence was subcloned into EcoRI and XhoI sites of the MSCV-IRES-GFP vector. Constructs in which KKMP was replaced by other sequences were also made as listed in FIG. 2. The sequence information for the scFvs generated herein is shown in Table 1. Sequence information for the various components depicted in FIG. 2 is shown in Table 2. Anti-CD19-4-1BB-CD3 ζ CAR

This CAR was generated as previously described (Imai, C. et al., *Leukemia*. 2004; 18:676-684; Imai, C. et al., *Blood*. 2005; 106:376-383).

TABLE 1

scFv sequence information				
Target	VH amino acid	VL amino acid	VH cDNA	VL cDNA
CD3	EVQLQQSGAELAR PGASVKMSCKAS GYTFTRYTMHWV KQRPQGQLEWIGY INPSRGYTNYNQK FKDKATLTDDKSS STAYMQLSSLTSE DSAVYYCARYYD DHYCLDYWGQGT TLTVSSA (SEQ ID NO: 12)	QIVLTQSPAIMSA SPGEKVTMTCSAS SSVSVMNWYQQ KSGTSPKRWIYDT SKLASGVPAHFR GSGSGTSYSLTIS GMEAEADAATYYC QQWSSNPFTFGSG TKLEINR (SEQ ID NO: 13)	GAGGTCCAGCTGCAGCAG TCTGGGGCTGAAGTGGCA AGACCTGGGGCCCTCAGTG AAGATGTCCTGCAAGGCTT CTGGCTACACCTTTACTAG GTACACGATGCACTGGGT AAAACAGAGGCTCGGACA GGGTCTGGAATGGATTGG ATACATTAATCCTAGCCGT GGTTATACTAATTACAATC AGAAGTTCAAGGACAAGG CCACATTGACTACAGACA AATCCTCCAGCACAGCCTA CATGCAACTGAGCAGCCT GACATCTGAGGACTCTGCA GTCTATTACTGTGCAAGAT ATTATGATGATCATTACTG CCTTGACTACTGGGCCCCA GGCACCACTCTCACAGTCT CCTCAGCC (SEQ ID NO: 14)	CAAATTGTTCTCACCCAG TCTCCAGCAATCATGTCT GCATCTCCAGGGGAGAA GGTCACCATGACCTGCA GTGCCAGCTCAAGTGTGA AGTTACATGAACCTGGTAC CAGCAGAAGTCAGGCAC CTCCCCAAAAGATGGA TTTATGACACATCCAAAC TGGCTTCTGGAGTCCCTG CTCACTTCAGGGGCACTG GGTCTGGGACCTCTTACT CTCTCACAATCAGCGGCA TGGAGGCTGAAGATGCT GCCACTTATTACTGCCAG CAGTGAGTAGTAACCC ATTACAGTTCTGGCTCGGG GACAAAGTTGGAATAAA ACCGG (SEQ ID NO: 15)
CD7 (TH69)	EVQLVESGGGLVK PGGSLKLSCKASG LTPSSYAMSWVR QTPEKRLWEVASI SSGGFTYYPDSVK GRFTISRDNARNIL YLQMSSLRSEDTA MYVCARDEVGRY LDVWGAGTTVT SS (SEQ ID NO: 16)	AAYKDIQMTQTT SSLSASLGDRVTIS CSASQGISNVLN WYQQKPDGTVKL LIYYTSSLHSGVP SRFSGSGSGTDYS LTISNLEPEDYATY YCCQYSKLPTFY GGGKLEIKR (SEQ ID NO: 17)	GAGGTGCAGCTGGTCGAA TCTGGAGGAGGACTGGTG AAGCCAGGAGGATCTCTG AAACTGAGTTGTGCCGCTT CAGGCCGTGACCTTCTCAAG CTACGCCATGAGCTGGGTG CGACAGACACCTGAGAAG CGGCTGGAATGGGTGCGT AGCATCTCCTCTGGCGGGT TCACATACTATCCAGACTC CGTGAAAGGCAGATTACT ATCTCTCGGGATAACGCAA GAAATATTCTGTACCTGCA GATGAGTTCACTGAGGAG CGAGGACACCGCAATGTA CTATTGTGCCAGGGACGA AGTGCGCGGCTATCTGGAT GTCTGGGAGCTGGCACT ACCGTCACCGCTCTCCAGC (SEQ ID NO: 18)	GCCGCATACAAGGATAT TCAGATGACTCAGACCA CAAGCTCCCTGAGCGCCT CCCTGGGAGACCGAGTG ACAATCTCTGCGAGTGCA TCACAGGGAATTAGCAA CTACCTGAATTGGTATCA GCAGAGCCAGATGGCA CTGTGAACCTGCTGATCT ACTATACCTCTAGTCTGC ACAGTGGGGTCCCTCAC GATTACGCGGATCCGGCT CTGGGACAGACTACAGC CTGACTATCTCCAACTG GAGCCCGAAGATATTGC CACCTACTATTGCCAGCA GTACTCCAAGCTGCCTTA TACCTTGGCGGGGGAA CAAAGCTGGAGATTAA AGG (SEQ ID NO: 19)
CD7 (3a1f)	QVQLQESGAELVK PGASVKLSCKASG YTFYSYWMHWVK QRPQGQLEWIGKI NPSNGRTNYNEKF KSKATLTVDKSSS TAYMQLSSLTSED SAVYYCARGGVY YDLYYYALDYWG QGTTVTVSS (SEQ ID NO: 20)	DIELTQSPATLSVT PGDSVLSLSCRASQ SISNNLHWYQQK SHESPRLLIKSASQ SISGIPSRFSGSGS GTDFTLSINSVETE DFGMYFCQQSNS WPYTFGGGTLKLEI KR (SEQ ID NO: 21)	CAGGTCCAGCTGCAGGAG TCAGGGGAGAGCTGGTG AAACCCGAGGAGCTGTC AAACTGTCCTGTAAGGCCA GCGGCTATACTTTCACCGA CTACTGGATGCACTGGGTG AAACAGAGGCCAGGACAG GGCCTGGAGTGGATCGGC AAGATTAAACCCAGCAAT GGGCGCACCACTACAAC GAAAAGTTTAAATCCAAG GCTACACTGACTGTGGACA AGAGCTCCTCTACCGCATA CATGCAGCTGAGTTCACTG ACATCTGAAGATAGTGCC	GACATCGAGCTGACACA GTCTCCAGCACTCTGAG CGTGACCCCTGGCGATT TGTCAGTCTGTGATGAG AGCTAGCCAGTCCATCTC TAACATCTGCATGGTA CCAGCAGAATCACATG AAAGCCCTCGGCTGCTG ATTAAGAGTGCTTCACAG AGCATCTCCGGGATTCCA AGCAGATTCTCTGGCAGT GGGTCAGGAACCGACTT TACACTGTCCATTAACTC TGTGGAGACCGAAGATT TCGGCATGTATTTTGGC

TABLE 1-continued

scFv sequence information				
Target	VH amino acid	VL amino acid	VH cDNA	VL cDNA
			GTGTACTATTGCGCCAGAG GCGGGGTCTACTATGACCT GTACTATTACGCACTGGAT TATTGGGGGAGGGAACC ACAGTGACTGTCAGCTCC (SEQ ID NO: 22)	AGCAGAGCAATTCTCTGG CCTTACACATTCCGGAGGC GGGACTAACTGGAGAT TAAGAGG (SEQ ID NO: 23)
CD45	QVQLVESGGGLV QPGGSLKLSCAAS GFDFSRYSWV RQAPGKLEWIGE INPTSSINFTPSLK DKVFI SRD NAKNT LYLQMSKVRSED ALYYCARGNYR YGDAMDYWGQG TSVTVS (SEQ ID NO: 24)	DIVLTQSPASLAV SLGQRATISCRAS KSVSTSGYSYLH WYQQKPGQPPKL LIYLASNLESQVP ARFSGSGSDTFT LNHPVEEEDAAT YYCQHSRELPTF SGSTKLEIK (SEQ ID NO: 25)	CAGGTGCAGCTGGTCGAG TCTGGAGGAGGACTGGTG CAGCCTGGAGGAAGTCTG AAGCTGTCATGTGCAGCCA GCGGGTTCGACTTTCTCG ATACTGGATGAGTTGGGTG CGGCAGGCACAGGAAAA GGACTGGAATGGATCGGC GAGATTAACCAACTAGCT CCACCATCAATTTACACC CAGCCTGAAGGACAAAGT GTTTATTTCCAGAGATAAC GCCAAGAATACTCTGTATC TGCAGATGTCCAAAGTCA GGTCTGAAGATACCGCCT GTACTATTGTGCTCGGGGC AACTACTATAGATACGGG GACGCTATGATTATTGGG GGCAGGGAAGTACCGTGA CCGTGAGT (SEQ ID NO: 26)	GACATTGTGCTGACCCAG TCCCCTGCTTCACTGGCA GTGAGCCTGGGACAGAG GGCAACCATCAGCTGCC GAGCCTCTAAGAGTGTCT CAACAAGCGGATACTCC TATCTGCACTGGTACCAG CAGAAGCCAGGACAGCC ACCTAAACTGCTGATCTA TCTGGCTTCCAACTTGA ATCTGGAGTGCCTGCACG CTTCTCCGGATCTGGAAG TGGAACCGACTTTACACT GAATATTCAACCACTCGA GGAAGAGGATGCCGCTA CCTACTATTGCCAGCACA GCCGGGAGCTGCCCTTCA CATTGCGAGCGGGACT AAGCTGGAGATCAAG (SEQ ID NO: 27)
B2MG	EVQLQQSGAELVK PGASVKLSCTPSG FMVKDTYIHWVK QRPKGLEWIGRI DPSDDIKYDPKF QGKATITADTSN TVSLQLSLTSED AVYYCARWFGDY GAMNYWGQTSV TVSS (SEQ ID NO: 28)	DIQMTQSPASQSA SLGESVTITCLAS QTIGTWLAWYQQ KPGKSPQLLIYAA TSLADGVPSRFSG SGSGTKFSLKIRT LQAEDFVSYYCQ QLYSKPYTFGGY TKLEIKRAD (SEQ ID NO: 29)	GAGGTGCAGCTGCAGCAG AGCGGAGCAGAACTGGTG AAACCTGGAGGCAGCGTC AAGCTGTCCTGTACTCCAT CTGGCTTCAACGTGAAGG ACACATACATCTACTGGGT CAAGCAGCGGCCCAAACA GGGACTGGAGTGGATCGG CAGAATTGACCCATCCGAC GGCGATATCAAGTATGATC CCAAATTCCAGGGGAAGG CTACTATTACCGCAGATAC CAGCTCCAACACAGTGAG TCTGCAGCTGTCTAGTCTG ACTAGCGAAGACACCGCC GTCTACTATTGTGCTAGAT GGTTTGCGGATTACGGGGC CATGAATTATTGGGGGCA GGGAACCAGCGTCACCGT GTCCAGC (SEQ ID NO: 30)	GATATTCAGATGACCCA GTCCCCTGCATCACAGAG CGCCTCCCTGGGCGAGTC AGTGACCATCACATGCCT GGCTAGCCAGACAAATG GCACCTGGCTGGCATGGT ACCAGCAGAAGCCCGGC AAATCCCCTCAGCTGCTG ATCTATGCAGTACTCTCT CTGGCAGACGGAGTGCC CAGTAGGTTCTCTGGGAG TGGATCAGGCACCAAGT TTTCTCTGAAAATTCGCA CACTGCAGGCTGAGGAT TTCGTCTCTACTATTGC CAGCAGCTGTACTCTAAA CCTTATACATTGGCGGG GGAATAAGCTGGAAAT CAAACGAGCAGAC (SEQ ID NO: 31)
NKG2A	EVQLVESGGGLVK PGGSLKLSCAASG FTFSSYAMSWVRQ SPEKRLWVAEISS GGSYTYYPDTVTG RFTISRDNKNTL YLEISLRSSEDTAM YYCTRHDYPRFF DVWGAGTTVTVS S (SEQ ID NO: 32)	QIVLTQSPALMSA SPGEKVTMTCSAS SSVSYYVYQQK PRSSPKPWYILTS NLASGVPARFSGS SGSTSYSLTISSM EAEDAATYYCQ WSGNPYTFGGT KLEIKR (SEQ ID NO: 33)	GAGGTGCAGCTGGTGGAG AGCGGAGGAGGACTGGTG AAGCCAGGAGGAAGCCTG AAGCTGTCCTGTGCCGCT CTGGCTTCAATTTCTCTC TTATGCAATGAGCTGGGTG CGGCAGTCCCCAGAGAAG AGACTGGAGTGGGTGGCA GAGATCAGCTCCGGAGGA TCCTACACCTACTATCCTG ACACAGTGACCGCGCGGT TCACAATCTCTAGAGATAA CGCCAAGAATACCCTGTAT CTGGAGATCTCTAGCCTGA GATCCGAGGATACAGCCA TGTAATATTGCACACAGGCA CGGCGACTACCCACGCTTC TTTGACGTGTGGGGAGCA GGAACCACAGTGACCGTG TCCTCT (SEQ ID NO: 34)	CAGATTGTCCTGACCCAG TCTCCAGCCTGATGAGC GCCTCCCTGGCGAGAA GGTGACAATGACCTGCTC TGCCAGCTCCTCTGTGAG CTACATCTATTGGTACCA GCAGAAGCCTCGGAGCT CCCCAAAGCCTGGATCT ATCTGACATCCAACCTGG CCTCTGGCGTGCCAGCCA GATTCTCTGGCAGCGGCT CCGGCACATCTTACAGCC TGACCATCTCTAGCATGG AGGCCGAGGACGCGCC ACCTACTATTGCCAGCAG TGGTCCGCAATCCATAT ACATTGCGCGCGGCAC CAAGCTGGAGATCAAGA GG (SEQ ID NO: 35)
KIR 2DL1 and 2/3	QVQLVQSGAEVK KPGSSVKVSCAS GGTFSFYAISWVR QAPGKLEWMMGG FIPFGAANYAQKP QGRVTITADESTST	EIVLTQSPVTLSSL PGERATLSCRASQ SVSSYLAWYQQK PGQAPRLLIYDAS NRATGI PARFSGS SGSDFTLTISSE	CAGGTCCAGCTGGTGCAGT CTGGAGCTGAAGTGAAGA AACCAGGGAGCTCCGTCA AGGTGTCATGC AAAGCAA GCGGCGGAGCTTCTCCTT TTATGCAATCTCTGGGTG	GAGATCGTGCTGACCCA GTCTCCTGTACACTGAG TCTGTACCAGGGGAAC GGGTACACTGTCTTGCA GAGCAAGCCAGTCCGTG AGCTCCTACCTGGCCTGG

TABLE 1-continued

scFv sequence information				
Target	VH amino acid	VL amino acid	VH cDNA	VL cDNA
	AYMELSSSLRSDDT AVYYCARIPSGSY YYDYDMVWQ GTTVTVSS (SEQ ID NO: 36)	PEDFAVYYCOQR SNWMYTFGQGTK LEIKRT (SEQ ID NO: 37)	AGACAGGCACCTGGACAG GGACTGGAGTGGATGGGA GGCTTC ATCCCAATTTTG GAGCCGCTAACTATGCCCA GAAGTTCAGGGCAGGGT GACCATCACAGCTGATGA GTCTACTAGTACCGCATAC ATGGAAGTGTCTAGTCTGA GGAGCGACGATAACGCCG TGTACTATTGTGCTCGCAT TCCATCAGGCAGCTACTAT TACGACTATGATATGGACG TGTGGGGCCAGGGGACCA CAGTCACCGTGAGCAGC (SEQ ID NO: 38)	TATCAGCAGAAGCCAGG CCAGGCTCCCAGGCTGCT GATCTACGATGCAAGCA ACAGGGCCACTGGGATT CCCCCCCCTTCTCTGGC AGTGGGTCAGGAACCGA CTTTACTCTGACCATTTO TAGTCTGGAGCCTGAAG ATTTCCCGGTGTACTATT GCCAGCAGCGATCCAAT TGGATGTATACTTTTGGC CAGGGGACCAAGCTGGA GATCAAACGGACA (SEQ ID NO: 39)

TABLE 2

Sequence information for components depicted in FIG. 2						
Target	CD8 SP amino acid	VH-VL linker amino acid	CD8 hinge and TM amino acid	CD8 SP cDNA	VH-VL linker cDNA	CD8 hinge and TM cDNA
CD3	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GGSGGGG SGGGGS (SEQ ID NO: 41)	KPTTTPAPRPPTP APTIASQPLSLRP EACRPAAGGAVH TRGLDFACDIYI WAPLAGTCGVLL LSLVITLY (SEQ ID NO: 42)	ATGGCCTTACC AGTGACCGCCT TGCTCCTGCCG CTGGCCTTGCT GCTCCACGCCG CCAGGCCG (SEQ ID NO: 44)	GGTGGTGGTG GTTCTGGTGG TGGTGGTTCT GGCGGCGGCG GCTCCGGTGG TGGTGGATCC (SEQ ID NO: 51)	AAGCCCACCACG ACGCCAGCGCCG CGACCACCAACA CCGGCGCCACC ATCGCGTCGCAG CCCCGTCCCTGC GCCAGAGGCGT GCCGGCCAGCGG CGGGGGGCGCAG TGCACACGAGGG GGCTGGACTTCG CCTGTGATATCTA CATCTGGGCGCC CTTGGCCGGGAC TTGTGGGGTCCTT CTCCTGTCACTGG TTATCACCCCTTA C (SEQ ID NO: 57)
CD7 (TH69)	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GGSGGGG SGGGGS (SEQ ID NO: 41)	TTTPAPRPPTPAP TIASQPLSLRPEA CRPAAGGAVHTR GLDFACDIYIWA PLAGTCGVLLLS LVITLY (SEQ ID NO: 50)	ATGGCTCTGCC TGTGACCGCAC TGCTGCTGCCC CTGGCTCTGCT GCTGCACGCCG CAAGACCT (SEQ ID NO: 45)	GGAGGAGGAG GAAGCGGAGG AGGAGGATCC GGAGGCGGGG GATCTGGAGG AGGAGGAAGT (SEQ ID NO: 52)	ACCACTACACCT GCACCAAGGCCT CCCACACCCGCTC CCACTATCGCTTC CCAGCCACTGTCC CTGAGGCCCGAG GCCTGCAGGCCA GCAGCTGGCGGA GCCGTGCATACT AGGGGGCTGGAC TTCGCTTGCGACA TCTACATCTGGGC CCCCTGGCAGG GACATGCGGAGT CCTGCTGCTGTCC CTGGTCATCACAC TTTAC (SEQ ID NO: 58)
CD7 (3a1f)	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GGSGGGG S (SEQ ID NO: 43)	TTTPAPRPPTPAP TIASQPLSLRPEA CRPAAGGAVHTR GLDFACDIYIWA PLAGTCGVLLLS LVITLY (SEQ ID NO: 50)	ATGGCTCTGCC CGTCACCGCTC TGCTGCTGCC CTGGCTCTGCT GCTGCACGCTG CTCGACCA (SEQ ID NO: 46)	GGAGGAGGAG GATCCGGCGG AGGAGGCTCT GGGGGAGGCG GGAGT (SEQ ID NO: 53)	ACTACCACACCA GCTCCAAGACCA CCTACCCCTGCAC CAACAATTGCTA GTACGCCACTGTG ACTGAGACCAGA AGCATGTAGGCC TGCAGCTGGAGG AGCTGTGCACAC CAGAGGCCTGGA CTTGCTGCGAT

TABLE 2-continued

Sequence information for components depicted in FIG. 2						
Target	CD8 SP amino acid	VH-VL linker amino acid	CD8 hinge and TM amino acid	CD8 SP cDNA	VH-VL linker cDNA	CD8 hinge and TM cDNA
						ATCTACATTGGG CTCCTCTGGCAGG AACCTGTGGCGT GCTGCTGCTGTCT CTGGTCATCACAC TTTAC (SEQ ID NO: 59)
CD45	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GSGGGGG SGGGGS (SEQ ID NO: 41)	KPTTTPAPRPPTP APTIASQPLSLRP EACRPAAGGAVH TRGLDFACDIYI WAPLAGTCGVLL LSLVITLY (SEQ ID NO: 42)	ATGGCTCTGCC CGTGACCGCTC TGCTGCTGCCT CTGGCTCTGCT GCTGCATGCTG CTCGACCT (SEQ ID NO: 47)	GGAGGAGGAG GAAGTGGAGG AGGAGGATCA GGAGGCGGGG GAAGCGGCGG GGGAGGCTCC (SEQ ID NO: 54)	AAGCCCACCACG ACGCCAGCGCCG CGACCACCAACA CCGGCGCCACC ATCGCTCGCAG CCCTGTCCCTGC GCCCAGAGGCGT GCCGGCCAGCGG CGGGGGCGCAG TGCACACGAGGG GGCTGGACTTCG CCTGTGATATCTA CATCTGGGCGCC CTTGGCCGGGAC TTGTGGGGTCCTT CTCCTGTCACTGG TTATCACCCCTTA C (SEQ ID NO: 57)
B2MG	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GSGGGGG SGGGGS (SEQ ID NO: 41)	KPTTTPAPRPPTP APTIASQPLSLRP EACRPAAGGAVH TRGLDFACDIYI WAPLAGTCGVLL LSLVITLY (SEQ ID NO: 42)	ATGGCCCTGCC CGTCACCGCCC TGCTGCTGCCC CTGGCTCTGCT GCTGCACGCCG CAAGACCC (SEQ ID NO: 48)	GGAGGAGGAG GAAGTGGAGG AGGAGGGTCA CGAGGCGGGG GAAGCGGCGG GGGAGGATCC (SEQ ID NO: 55)	AAGCCCACCACG ACGCCAGCGCCG CGACCACCAACA CCGGCGCCACC ATCGCTCGCAG CCCTGTCCCTGC GCCCAGAGGCGT GCCGGCCAGCGG CGGGGGCGCAG TGCACACGAGGG GGCTGGACTTCG CCTGTGATATCTA CATCTGGGCGCC CTTGGCCGGGAC TTGTGGGGTCCTT CTCCTGTCACTGG TTATCACCCCTTA C (SEQ ID NO: 57)
NKG2A	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GSGGGGG SGGGGS (SEQ ID NO: 41)	KPTTTPAPRPPTP APTIASQPLSLRP EACRPAAGGAVH TRGLDFACDIYI WAPLAGTCGVLL LSLVITLY (SEQ ID NO: 42)	ATGGCTCTGCC CGTGACCGCCC TGCTGCTGCCT CTGGCTCTGCT GCTGCACGCTG CCCGCCCA (SEQ ID NO: 49)	GGAGGAGGAG GATCTGGAGG AGGAGGCAGC GGCGGCGGCG GCTCCGGCGG CGGCGGCTCT (SEQ ID NO: 56)	AAGCCAACCACA ACCCCTGCACCA AGGCCACCTACA CCAGCACCTACC ATCGCAAGCCAG CCACTGTCCCTGA GGCCAGAGGCAT GTAGGCCTGCAG CAGGAGGCGCCG TGCACACACGCG GCCTGGACTTTGC CTGCGATATCTAC ATCTGGGCACCA CTGGCAGGAACC TGTGGCGTGCTGC TGCTGAGCCTGGT GATTACCCTGTAT (SEQ ID NO: 60)
KIR 2DL1 and 2/3	MALPVT ALLLPLA LLLHAAR P (SEQ ID NO: 40)	GGGSGGG GSGGGGG SGGGGS (SEQ ID NO: 41)	KPTTTPAPRPPTP APTIASQPLSLRP EACRPAAGGAVH TRGLDFACDIYI WAPLAGTCGVLL LSLVITLY (SEQ ID NO: 42)	ATGGCCTTACC AGTGACCGCCT TGCTCCTGCCG CTGGCCTTGCT GCTCCACGCCG CCAGGCCG (SEQ ID NO: 44)	GGTGGTGGTG GTTCTGGTGG TGGTGGTTCT GGCGGCGGCG GTCCTGGTGG TGGTGGATCC (SEQ ID NO: 51)	AAGCCCACCACG ACGCCAGCGCCG CGACCACCAACA CCGGCGCCACC ATCGCTCGCAG CCCTGTCCCTGC GCCCAGAGGCGT GCCGGCCAGCGG CGGGGGCGCAG

TABLE 2-continued

Sequence information for components depicted in FIG. 2						
Target	CD8 SP amino acid	VH-VL linker amino acid	CD8 hinge and TM amino acid	CD8 SP cDNA	VH-VL linker cDNA	CD8 hinge and TM cDNA
						TGCACACGAGGG GGCTGGACTTCG CCTGTGATATCTA CATCTGGGCGCC CTTGGCCGGGAC TTGTGGGGTCCTT CTCCTGTCACTGG TTATCACCCCTTA C (SEQ ID NO: 57)

Gene Transduction, Cell Expansion, Flow Cytometric Analysis and Functional Studies

These were performed as previously described (Kudo, K et al., *Cancer Res.* 2014; 74(1):93-103).

Results

Generation of scFv Constructs

A schematic of the technology is outlined in FIG. 1. A schematic representation of the inhibitory constructs that we generated is shown in FIG. 2. The scFv portion can be derived by cloning the cDNA encoding variable light (VL) and variable heavy (VH) immunoglobulin chain regions from an antibody-producing hybridoma cell line or from the corresponding published sequences. VL and VH are linked with a short peptide sequence ("linker") according to standard techniques to make a full scFv. To be expressed, the scFv is linked to a signal peptide at the N-terminus; the signal peptide is required for the scFv to be expressed, as confirmed in preliminary experiments. Proteins containing scFv plus signal peptide are generally released into the cells' milieu. For example, in preliminary experiments (not shown), an anti-CD3 ϵ scFv plus signal peptide expressed in Jurkat T cells was detected in the cells' culture supernatant. By directing the scFv to specific compartments and preventing its secretion, possible effects on other cells are prevented. To direct it to the endoplasmic reticulum (ER), the KDEL (SEQ ID NO: 4) motif (which retains proteins in the ER) was utilized (Strebe N. et al., *J Immunol Methods.* 2009; 341(1-2):30-40). To promote the degradation of the targeted protein, we linked it to a proteasome-targeting PEST motif (Joshi, S. N. et al., *MABs.* 2012; 4(6):686-693). The scFv can also be directed to the cell membrane by linking it to the transmembrane domain and hinge of CD8 α or another transmembrane protein.

Downregulation of T-Cell Receptor in T Lymphocytes Expressing Anti-CD19-BB- ζ CAR

To determine whether the proposed strategy could be applied to generate immune cells expressing CAR and lacking one or more markers, T-cell receptor (TCR) expression was downregulated in anti-CD19 CAR T-cells.

To be expressed on the cell membrane, the CD3/TCR complex requires assembly of all its components (TCR α , TCR β , CD3 δ , CD3 ϵ , CD3 γ , CD3 ζ). Lack of one component prevents CD3/TCR expression and, therefore, antigen recognition. In preliminary studies, the scFv from an anti-CD3 ϵ hybridoma (purchased from Creative Diagnostics, Shirley, N.Y.) was cloned and generated the constructs containing KDEL (SEQ ID NO: 4), PEST, CD8 α transmembrane domain or others as shown in FIG. 2.

The constructs disclosed herein were transduced in the CD3/TCR+ Jurkat cell line using a murine stem cell virus

(MSCV) retroviral vector containing green fluorescent protein (GFP). Percentage of GFP+ cells after transduction was >90% in all experiments. FIG. 3A shows results of staining with anti-CD3 ϵ antibody among GFP+ cells, as measured by flow cytometry. Antibody staining of CD3 ϵ was decreased to variable degree in cells transduced with the constructs listed. Similar downregulation of CD3 ϵ was obtained with human peripheral blood T lymphocytes (FIG. 3B). FIG. 3C shows illustrative flow cytometry dot plots of CD3 ϵ expression in GFP-positive Jurkat cells after transduction with different gene constructs in comparison with cells transduced with a vector containing GFP alone. Downregulation of CD3 did not affect growth of Jurkat cells or expression of all other cell markers tested, including CD2, CD4, CD8, CD45, CD25, CD69. Inhibition of CD3 expression persisted for over 3 months. Further enrichment of CD3-negative cells could be achieved by CD3+ T cell depletion with anti-CD3 magnetic beads (Dyna, Life Technologies, Carlsbad, Calif.).

Staining with anti-TCR $\alpha\beta$ antibody of Jurkat cells or human peripheral blood T lymphocytes showed that downregulation of CD3 ϵ expression was associated with downregulation of TCR $\alpha\beta$ expression (FIG. 4).

Next, it was determined whether the anti-CD3 scFv-myc KDEL could be expressed simultaneously with an anti-CD19-4-1BB-CD3 ζ CAR. As shown in FIG. 5, this resulted in T cells lacking CD3 expression while expressing the anti-CD19 CAR. TCR was also absent on these cells (not shown).

To assess whether CAR could signal in Jurkat cells with downregulated CD3/TCR, the expression of the activation markers CD69 and CD25 was tested, and exocytosis of lytic granules was measured by CD107a expression in Jurkat cells co-cultured with the CD19+ leukemia cell line OP-1. As shown in FIG. 6, downregulation of CD3/TCR with the anti-CD3 scFv-myc KDEL construct did not diminish the capacity of anti-CD19-4-1BB-CD3 ζ CAR to activate Jurkat cells. To further explore the effects of CD3/TCR deletion on CAR signaling, it was determined whether CD3-negative T lymphocytes expressing the CAR could be stimulated by its ligation. As shown in FIG. 7, co-culture of T lymphocytes expressing the anti-CD19 CAR with CD19+ leukemic cells led to T cell proliferation regardless of whether CD3 was downregulated or not, indicating that CD3/TCR downregulation did not diminish the CAR proliferative stimulus.

Accordingly, CD3/TCR can be effectively downregulated in CAR-T cells using the anti-CD3 scFv-myc KDEL construct without affecting T cell activation, degranulation and proliferation driven by the CAR.

Downregulation of CD7

It was determined whether the strategy that successfully modulated CD3/TCR expression could be applied to other surface molecules. For this purpose, CD7 expression was modulated. The scFv sequence was derived from that published by Peipp et al. (Cancer Res 2002 (62): 2848-2855), which was linked to the CD8 signal peptide and the myc-KDEL sequence as illustrated in FIG. 2. Using the MSCV retroviral vector, the anti-CD7-myc KDEL construct was transduced in peripheral blood lymphocytes, which have high expression of CD7 as detected by an anti-CD7 antibody conjugated to phycoerythrin (BD Bioscience). As shown in FIG. 8, CD7 in T lymphocytes transduced with the construct was virtually abrogated.

Downregulation of HLA-Class I

The strategy was then applied to downregulate another surface molecule, HLA class I.

HLA class I consists of polymorphic α chains and a non-polymorphic chain termed β 2-microglobulin. Knock-down of the latter subunit results in abrogation of HLA (MHC in the mouse) Class I expression (Koller, B H et al., *Science*, 1990; 248(4960):1227-1230). An scFv reacting with β 2-microglobulin was used to suppress expression of HLA Class I in immune cells.

The scFv sequence was derived from that published by Grovender et al. (*Kidney Int.* 2004; 65(1):310-322), which was linked to the CD8 signal peptide and the myc KDEL sequence as illustrated in FIG. 2. Using the MSCV retroviral vector, the anti- β 2M-myc KDEL construct was transduced in Jurkat cells, which have high expression of HLA Class I as detected by an anti-HLA-ABC antibody conjugated to phycoerythrin (BD Pharmingen). As shown in FIG. 9, Jurkat cells transduced with the construct had a substantial down-regulation of HLA-ABC expression. Cells maintained their morphology and growth capacity.

Downregulation of Inhibitory Receptors in NK Cells

To determine if the strategy outlined above would also apply to surface molecules expressed in other immune cells,

downregulation of function of the inhibitory receptor KIR2DL1, KIR2DL2/DL3 and NKG2A was tested in NK cells.

To downregulate KIR receptors, an scFv reacting with KIR2DL1 and KIR2DL2/DL3 was used to suppress their expression in NK cells. The scFv sequence was derived from that published by Moretta et al. (patent WO2006003179 A2), which was linked to the CD8 signal peptide and the ER retention sequences as illustrated in FIG. 2. Using the MSCV retroviral vector, the constructs were transduced in NK cells expanded from human peripheral blood and selected for KIR2DL1 expression. These cells had high KIR2DL1 expression as detected by an anti-KIR2DL1 antibody conjugated to allophycocyanin (R&D Systems) and also high KIR2DL2/DL3 expression as detected by an anti-KIR2DL2/DL3 antibody conjugated to phycoerythrin (BD Bioscience). FIG. 10 shows results obtained with scFv-linker(20) AEKEDL and scFv-EEKKMP, with substantial down regulation of the targeted KIRs.

To downregulate NKG2A, an scFv reacting with NKG2A was used to suppress its expression in NK cells. The scFv sequence, which was derived from published European Patent Application No. EP2247619 A1 by Spee et al. was linked to the CD8 signal peptide and the ER retention sequences as illustrated in FIG. 2. Using the MSCV retroviral vector, the constructs were transduced in NK cells expanded from human peripheral blood, which had high NKG2A expression as detected by an anti-NKG2A antibody conjugated to phycoerythrin (Beckman Coulter). FIG. 11 shows substantial downregulation of NKG2A obtained with scFv-EEKKMP.

The teachings of all patents, published applications and references cited herein are incorporated by reference in their entirety.

While this invention has been particularly shown and described with references to example embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

SEQUENCE LISTING

```

Sequence total quantity: 64
SEQ ID NO: 1          moltype = AA  length = 10
FEATURE              Location/Qualifiers
REGION               1..10
                     note = Myc tag
source               1..10
                     mol_type = protein
                     organism = synthetic construct

SEQUENCE: 1
EQKLISEEDL                                     10

SEQ ID NO: 2          moltype = DNA  length = 40
FEATURE              Location/Qualifiers
misc_feature          1..40
                     note = Sense primer
source               1..40
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 2
atatatgaat tcggcttcca ccatggcctt accagtgacc    40

SEQ ID NO: 3          moltype = DNA  length = 50
FEATURE              Location/Qualifiers
misc_feature          1..50
                     note = Reverse primer
source               1..50
                     mol_type = other DNA
                     organism = synthetic construct

```

-continued

SEQUENCE: 3
cagatcttct tcagaaataa gtttttggtc ggctgaggag actgtgagag 50

SEQ ID NO: 4 moltype = AA length = 4
FEATURE Location/Qualifiers
REGION 1..4
 note = KDEL coding sequence
source 1..4
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 4
KDEL 4

SEQ ID NO: 5 moltype = DNA length = 40
FEATURE Location/Qualifiers
misc_feature 1..40
 note = Sense primer
source 1..40
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 5
atatatgaat tcggttcca ccatggcctt accagtgacc 40

SEQ ID NO: 6 moltype = DNA length = 48
FEATURE Location/Qualifiers
misc_feature 1..48
 note = Reverse primer
source 1..48
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 6
tatatactcg agttacaact cgtccttcag atcttcttca gaaataag 48

SEQ ID NO: 7 moltype = AA length = 40
FEATURE Location/Qualifiers
REGION 1..40
 note = PEST motif
source 1..40
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 7
SHGFPPEVEE QDDGTLPMSC AQESGMDRHP AACASARINV 40

SEQ ID NO: 8 moltype = AA length = 5
FEATURE Location/Qualifiers
REGION 1..5
 note = Hydrophilic peptide
source 1..5
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 8
GGGS 5

SEQ ID NO: 9 moltype = length =
SEQUENCE: 9
000

SEQ ID NO: 10 moltype = length =
SEQUENCE: 10
000

SEQ ID NO: 11 moltype = AA length = 4
FEATURE Location/Qualifiers
REGION 1..4
 note = ER or Golgi retention sequence
source 1..4
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 11
YQRL 4

SEQ ID NO: 12 moltype = AA length = 120
FEATURE Location/Qualifiers
REGION 1..120
 note = Variable heavy region for scFv targeting CD3
source 1..120
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 12

-continued

```

EVQLQQSGAE LARPGASVKM SCKASGYTFT RYTMHWVKQR PGQGLEWIGY INPSRGYTNV 60
NQKFKDKATL TTDKSSSTAY MQLSSLTSED SAVYYCARYY DDHYCLDYWG QGTTLTVSSA 120

SEQ ID NO: 13      moltype = AA length = 107
FEATURE            Location/Qualifiers
REGION            1..107
                  note = Variable light region for scFv targeting CD3
source            1..107
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 13
QIVLTQSPAI MSASPGEKVT MTCASSSSVS YMNWYQQKSG TSPKRWIYDT SKLASGVPAH 60
FRGSGSGTSY SLTISGMEAE DAATYYCQQW SSNPFTFGSG TKLEINR 107

SEQ ID NO: 14      moltype = DNA length = 360
FEATURE            Location/Qualifiers
misc_feature       1..360
                  note = Variable heavy region for scFv targeting CD3
source            1..360
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 14
gagggtccagc tgcagcagtc tgggggctgaa ctgggcaagac ctgggggcctc agtgaagatg 60
tcctgcaagg cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
gatgatcatt actgcttga ctactggggc caaggcacca ctctcacagt ctctcagcc 360

SEQ ID NO: 15      moltype = DNA length = 321
FEATURE            Location/Qualifiers
misc_feature       1..321
                  note = Variable light region for scFv targeting CD3
source            1..321
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 15
caaatgttgc tcaccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
atgacctgca gtgcagctc aagtgttaagt tacatgaact ggtaccagca gaagtcaggc 120
acctcccca aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcac 180
ttcaggggca gtgggtctgg gacctcttac tctctcaca tcagcggcat ggaggtgtaa 240
gatgctgcca cttattactg ccagcagtggt agtagtaacc cattcacgtt cggctcgggg 300
acaaagttgg aaataaaccg g 321

SEQ ID NO: 16      moltype = AA length = 117
FEATURE            Location/Qualifiers
REGION            1..117
                  note = Variable heavy region for scFv targeting CD7(TH69)
source            1..117
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 16
EVQLVESGGG LVKPGGSLKL SCAASGLTFS SYAMSWVRQT PEKRLEWVAS ISSGGFTYYP 60
DSVKGRFTIS RDNARNILYL QMSSLRSED AMYYCARDEV RGYLDVWGAG TTVTVSS 117

SEQ ID NO: 17      moltype = AA length = 112
FEATURE            Location/Qualifiers
REGION            1..112
                  note = Variable light region for scFv targeting CD7(TH69)
source            1..112
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 17
AAYKDIQMTQ TTSSLASLG DRVTISCSAS QGISNYLNWY QQKPDGTVKL LIYTTSSLHS 60
GVPSRFGSGG SGTDSLITIS NLEPEDIATY YCQQYSKLPY TFGGGTKLEI KR 112

SEQ ID NO: 18      moltype = DNA length = 351
FEATURE            Location/Qualifiers
misc_feature       1..351
                  note = Variable heavy region for scFv targeting CD7(TH69)
source            1..351
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 18
gagggtgcagc tgggtogaatc tggaggagga ctgggtgaagc caggaggatc tctgaaactg 60
agttgtgccg cttcaggcct gacctctca agctacgcca tgagctgggt gcgacagaca 120
cctgagaagc ggctggaatg ggtcgctagc atctcctctg cgggggtcac atactatcca 180
gactccgtga aaggcagatt tactatctct cgggataacg caagaaatat tctgtacctg 240
cagatgagtt cactgaggag cgaggacacc gcaatgtact attgtgccag ggacgaagtg 300

```

-continued

cgcggtatc tggatgtctg gggagctggc actaccgtca ccgtctccag c 351

SEQ ID NO: 19 moltype = DNA length = 336
 FEATURE Location/Qualifiers
 misc_feature 1..336
 note = Variable light region for scFv targeting CD7(TH69)
 source 1..336
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 19
 gccgcataca aggatattca gatgactcag accacaagct ccctgagcgc ctccctggga 60
 gaccgagtgca caatctcttg cagtgcacat caggggaatta gcaactacct gaattgggtat 120
 cagcagaagc cagatggcac tgtgaaactg ctgatctact atacctctag tctgcacagt 180
 ggggtccct cagcattcag cggatccggc tctgggacag actacagcct gactatctcc 240
 aacctggagc ccgaagatat tgccacctac tattgccagc agtactccaa gctgccttat 300
 acctttggcg ggggaacaaa gctggagatt aaaagg 336

SEQ ID NO: 20 moltype = AA length = 123
 FEATURE Location/Qualifiers
 REGION 1..123
 note = Variable heavy region for scFv targeting CD7(3a1f)
 source 1..123
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 20
 QVQLQESGAE LVKPGASVKL SCKASGYTFT SYWMHWVKQR PGQGLEWIGK INPSNGRTNY 60
 NEKFKSKATL TVDKSSSTAY MQLSSLTSED SAVYYCARGG VYYDLYYAL DYWGQGTTVT 120
 VSS 123

SEQ ID NO: 21 moltype = AA length = 108
 FEATURE Location/Qualifiers
 REGION 1..108
 note = Variable light region for scFv targeting CD7(3a1f)
 source 1..108
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 21
 DIELTQSPAT LSVTPGDSVS LSCRASQSIG NNLHWYQQKS HESPRLLIKS ASQSISGIPS 60
 RFSGSGSGTD FTLSINSVET EDFGMYFCQQ SNSWPYTFGG GTKLEIKR 108

SEQ ID NO: 22 moltype = DNA length = 369
 FEATURE Location/Qualifiers
 misc_feature 1..369
 note = Variable heavy region for scFv targeting CD7(3a1f)
 source 1..369
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 22
 caggtccagc tgcaggagtc aggggcagag ctgggtgaaac ccggagccag tgtcaaaactg 60
 tctgtgaagg ccagcggcta tactttcacc agctactgga tgcactgggt gaaacagagg 120
 ccaggacagg gcctggagtg gatcggaagc attaacccca gcaatgggag caccacactac 180
 aacgaaaagt ttaaatccaa ggctacactg actgtggaca agagctcctc taccgcatac 240
 atgcagctga gttcactgac atctgaagat agtgccgtgt actattgcgc cagagggcgg 300
 gtctactatg acctgtacta ttacgcactg gattattggg ggcagggaac cacagtgact 360
 gtcagctcc 369

SEQ ID NO: 23 moltype = DNA length = 324
 FEATURE Location/Qualifiers
 misc_feature 1..324
 note = Variable light region for scFv targeting CD7(3a1f)
 source 1..324
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 23
 gacatcgagc tgacacagtc tccagccact ctgagcgtga cccctggcga ttctgtcagt 60
 ctgtcatgta gagctagcca gtccatctct aacaatctgc actggtagca gcagaaatca 120
 catgaaagcc ctccggtgct gattaagagt gcttcacaga gcatctccgg gattccaagc 180
 agattctctg gcagtgggtc aggaaccgac ttacactgt ccattaactc tgtggagacc 240
 gaagatttcg gcattgattt ttgccagcag agcaattcct ggccttacac attcggaggc 300
 gggactaaac tggagattaa gagg 324

SEQ ID NO: 24 moltype = AA length = 120
 FEATURE Location/Qualifiers
 REGION 1..120
 note = Variable heavy region for scFv targeting CD45
 source 1..120
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 24

-continued

QVQLVESGGG LVQPGGSLKL SCAASGFDFS RYWMSWVRQA PGKGLEWIGE INPTSSTINF 60
 TPSLKDVKFI SRDNAKNTLY LQMSKVRSED TALYYCARGN YYRYGDAMDY WQGTSVTVS 120

SEQ ID NO: 25 moltype = AA length = 111
 FEATURE Location/Qualifiers
 REGION 1..111
 note = Variable light region for scFv targeting CD45
 source 1..111
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 25
 DIVLTQSPAS LAVSLGQRAT ISCRASKSVS TSGYSYLHWY QOKPGQPPKL LIYLASNLES 60
 GVPARFSGSG SGTDFTLNIH PVEEEDAATY YCQHSRELFP TFGSGTKLEI K 111

SEQ ID NO: 26 moltype = DNA length = 360
 FEATURE Location/Qualifiers
 misc_feature 1..360
 note = Variable heavy region for scFv targeting CD45
 source 1..360
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 26
 cagggtgcagc tgggtcgagtc tggaggagga ctgggtgcagc ctggagggaag tctgaagctg 60
 tcatgtgcag ccagcgggtt cgacttttct cgatactgga tgagttgggt gcggcaggca 120
 ccaggaaaag gactggaatg gatcggcgag attaaccctaa ctagctccac catcaatttc 180
 acacccagcc tgaaggacaa agtggtttatt tccagagata acgccaagaa tactctgtat 240
 ctgcagatgt ccaaagtcag gtctgaagat accgccctgt actattgtgc tcggggcaac 300
 tactatagat acggggacgc tatggattat tgggggcagg gaactagcgt gaccgtgagt 360

SEQ ID NO: 27 moltype = DNA length = 333
 FEATURE Location/Qualifiers
 misc_feature 1..333
 note = Variable light region for scFv targeting CD45
 source 1..333
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 27
 gacattgtgc tgaccagtc cctgcttca ctggcagtga gcctgggaca gagggcaacc 60
 atcagctgcc gagcctctaa gagtgtctca acaagcggat actcctatct gcactggtag 120
 cagcagaagc caggacagcc acctaaactg ctgactctac tggcttccaa cctggaatct 180
 ggagtgctgc cagcgtcttc cggatctgga agtggaaacc actttacact gaattattac 240
 ccagtcgagg aagaggatgc cgctacctac tattgccagc acagccggga gctgcccttc 300
 acatttgga cggggactaa gctggagatc aag 333

SEQ ID NO: 28 moltype = AA length = 119
 FEATURE Location/Qualifiers
 REGION 1..119
 note = Variable heavy region for scFv targeting B2MG
 source 1..119
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 28
 EVQLQQSGAE LVKPGASVKL SCTPSGFNVK DTYIHVVKQR PKQGLEWIGR IDPSDGDIKY 60
 DPKFQKQATI TADTSSNTVS LQLSSLTSED TAVYYCARWF GDYGMNYWG QGTSVTVSS 119

SEQ ID NO: 29 moltype = AA length = 110
 FEATURE Location/Qualifiers
 REGION 1..110
 note = Variable light region for scFv targeting B2MG
 source 1..110
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 29
 DIQMTQSPAS QSASLGESVT ITCLASQTIG TWLAWYQKP GKSPQLLIYA ATSLADGVPS 60
 RFSGSGSGTK FSLKIRTQA EDFVSYQCQ LYSKPYTFGG GTKLEIKRAD 110

SEQ ID NO: 30 moltype = DNA length = 357
 FEATURE Location/Qualifiers
 misc_feature 1..357
 note = Variable heavy region for scFv targeting B2MG
 source 1..357
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 30
 gagggtgcagc tgcagcagag cggagcagaa ctgggtgaaac ctggagccag cgtcaagctg 60
 tcctgtactc catctgctt caacgtgaag gacacataca ttactgggt caagcagcgg 120
 cccaaacagg gactggagtg gatcggcaga attgacctat ccgacggcga tatcaagtat 180
 gatcccaaat tccaggggaa ggctactatt accgcagata ccagctccaa cacagtgaat 240
 ctgcagctgt ctagtctgac tagcgaagac accgccgtct actattgtgc tagatggttt 300

-continued

```

ggcgattacg gggccatgaa ttattggggg cagggaaacca gcgtaccgt gtccagc 357

SEQ ID NO: 31      moltype = DNA length = 330
FEATURE           Location/Qualifiers
misc_feature       1..330
                   note = Variable light region for scFv targeting B2MG
source            1..330
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 31
gatattcaga tgaccagtc cctgcatac cagagcgct cctggggcga gtcagtgacc 60
atcacatgcc tggctagcca gacaattggc acttggtgg catggtacca gcagaagccc 120
ggcaaatccc ctgagctgct gatctatgca gctacctctc tggcagacgg agtgcccagt 180
aggttctctg gtagtggatc aggcaccaag tttctctga aaattcgac actgcaggct 240
gaggatttcg tctcctacta ttgccagcag ctgtactcta aacctatac atttggcggg 300
ggaactaagc tggaaatcaa acgagcagac 330

SEQ ID NO: 32      moltype = AA length = 119
FEATURE           Location/Qualifiers
REGION           1..119
                   note = Variable heavy region for scFv targeting NKG2A
source            1..119
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 32
EVQLVESGGG LVKPGGSLKL SCAASGFTFS SYAMSWVRQS PEKRLIEWAE ISSGGSYTTY 60
PDTVTGRFTI SRDIAKNTLY LETSSLRSED TAMYCTRHG DYPRFFDVGW AGTTVTVSS 119

SEQ ID NO: 33      moltype = AA length = 107
FEATURE           Location/Qualifiers
REGION           1..107
                   note = Variable light region for scFv targeting NKG2A
source            1..107
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 33
QIVLTQSPAL MSASPGEKVT MTCASASSVS YIYWYQOKPR SSPKPWIYLT SNLASGVPAR 60
FSGSGSGTSY SLTISSMEAE DAATYYCQQW SGNPYTFGGG TKLEIKR 107

SEQ ID NO: 34      moltype = DNA length = 357
FEATURE           Location/Qualifiers
misc_feature       1..357
                   note = Variable heavy region for scFv targeting NKG2A
source            1..357
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 34
gagggtgcagc tgggtggagag cggaggagga ctggtgaagc caggaggaag cctgaagctg 60
tcctgtgccg cctctggctt cacattttcc tcttatgcaa tgagctgggt gcggcagtc 120
ccagagaaga gactggagtg ggtggcagag atcagctccg gaggatccta cacctactat 180
cctgacacag tgaccggccg gtccacaatc tctagagata acgccaagaa taccctgtat 240
ctggagatct ctgacctgag atccagagat acagccatgt actattgcac caggcacggc 300
gactaccacc gcttctttga cgtgtgggga gcaggaacca cagtgaccgt gtectct 357

SEQ ID NO: 35      moltype = DNA length = 321
FEATURE           Location/Qualifiers
misc_feature       1..321
                   note = Variable light region for scFv targeting NKG2A
source            1..321
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 35
cagattgtcc tgaccagtc tccagccctg atgagcgct cccctggcga gaagggtgaca 60
atgacctgct ctgccagctc ctctgtgagc tacatctatt ggtaccagca gaagcctcgg 120
agctcccaaa agccctggat ctatctgaca tccaacctgg cctctggcgt gccagccaga 180
ttctctggca gcggctccgg cacatcttac agcctgacca tctctagcat ggaggccgag 240
gacgcccga cctactattg ccagcagtg tccggcaatc catatacatt tggcggcggc 300
accaagctgg agatcaagag g 321

SEQ ID NO: 36      moltype = AA length = 122
FEATURE           Location/Qualifiers
REGION           1..122
                   note = Variable heavy region for scFv targeting KIR2DL1and
                   2/3
source            1..122
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 36
QVQLVQSGAE VKKPGSSVKV SCKASGGTFS FYAISWVRQA PGQGLEWMGG FIPFGAANY 60

```

-continued

AQKFQGRVTI TADESTSTAY MELSSLRSDD TAVYYCARPS GSYYYDYDMD VWGQGTVTVT 120
SS 122

SEQ ID NO: 37 moltype = AA length = 109
FEATURE Location/Qualifiers
REGION 1..109
note = Variable light region for scFv targeting KIR2DL1and
2/3
source 1..109
mol_type = protein
organism = synthetic construct

SEQUENCE: 37
EIVLTQSPVT LSLSPGERAT LSCRASQSVS SYLAWYQQKP GQAPRLLIYD ASNRATGIPA 60
RFGSGSGTDT FTLTISLEP EDFAVYYCQQ RSNWMYTFGQ GTKLEIKRT 109

SEQ ID NO: 38 moltype = DNA length = 369
FEATURE Location/Qualifiers
misc_feature 1..369
note = Variable heavy region for scFv targeting KIR2DL1and
2/3
source 1..369
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 38
caggtccagc tgggtcagtc tggagctgaa gtgaagaaac cagggagctc cgtcaagggtg 60
tcatgcaaaag caagcggcgg gactttctcc ttttatgcaa tctcttgggt gagacaggca 120
cctggacagg gactggagtg gatgggaggg ttcaccccaa tttttggagc cgctaactat 180
gcccagaagt tccagggcag ggtgaccatc acagctgatg agtctactag taccgcatac 240
atggaaactgt ctagtctgag gagcgacgat accgcccgtg actattgtgc tcgcattcca 300
tcaggcagct actattacga ctatgatatg gacgtgtggg gccagggggac cacagtcacc 360
gtgagcagc 369

SEQ ID NO: 39 moltype = DNA length = 327
FEATURE Location/Qualifiers
misc_feature 1..327
note = Variable light region for scFv targeting KIR2DL1and
2/3
source 1..327
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 39
gagatcgtgc tgacccagtc tctgttcaca ctgagctctgt caccagggga acgggctaca 60
ctgtcttgca gagcaagcca gtccgtgagc tcctacctgg cctggatca gcagaagcca 120
ggccaggctc ccaggctgct gatctacgat gcaagcaaca gggccactgg gattcccgc 180
cgcttctctg gcagtggtgc aggaaccgac tttactctga ccatttctag tctggagcct 240
gaagatttcg ccgtgtacta ttgccagcag cgatccaatt ggatgtatac ttttggccag 300
gggaccaagc tggagatcaa acggaca 327

SEQ ID NO: 40 moltype = AA length = 21
FEATURE Location/Qualifiers
REGION 1..21
note = CD8 signal peptide
source 1..21
mol_type = protein
organism = synthetic construct

SEQUENCE: 40
MALPVTALLL PLALLLHAAR P 21

SEQ ID NO: 41 moltype = AA length = 20
FEATURE Location/Qualifiers
REGION 1..20
note = Variable heavy-variable light linker
source 1..20
mol_type = protein
organism = synthetic construct

SEQUENCE: 41
GGGSGGGGS GGGSGGGGS 20

SEQ ID NO: 42 moltype = AA length = 70
FEATURE Location/Qualifiers
REGION 1..70
note = CD8 hinge and transmembrane
source 1..70
mol_type = protein
organism = synthetic construct

SEQUENCE: 42
KPTTTPAPRP PTPAPTASQ PLSLRPEACR PAAGGAVHTR GLDFACDIYI WAPLAGTCGV 60
LLLSLVITLY 70

-continued

SEQ ID NO: 43	moltype = AA length = 15	
FEATURE	Location/Qualifiers	
REGION	1..15	
	note = Variable heavy-variable light linker	
source	1..15	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 43		
GGGSGGGGS GGGGS		15
SEQ ID NO: 44	moltype = DNA length = 63	
FEATURE	Location/Qualifiers	
misc_feature	1..63	
	note = CD8 signal peptide	
source	1..63	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 44		
atggccttac cagtgaccgc cttgtctctg ccgctggcct tgctgtctca cgccgccagg		60
cctg		63
SEQ ID NO: 45	moltype = DNA length = 63	
FEATURE	Location/Qualifiers	
misc_feature	1..63	
	note = CD8 signal peptide	
source	1..63	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 45		
atggctctgc ctgtgaccgc actgctgctg cccctggctc tgctgtctga cgccgcaaga		60
cct		63
SEQ ID NO: 46	moltype = DNA length = 63	
FEATURE	Location/Qualifiers	
misc_feature	1..63	
	note = CD8 signal peptide	
source	1..63	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 46		
atggctctgc ccgtcaccgc tctgtgctg cctctggctc tgctgtctga cgtgtctga		60
cca		63
SEQ ID NO: 47	moltype = DNA length = 63	
FEATURE	Location/Qualifiers	
misc_feature	1..63	
	note = CD8 signal peptide	
source	1..63	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 47		
atggctctgc ccgtgaccgc tctgtgctg cctctggctc tgctgtctga tgctgtctga		60
cct		63
SEQ ID NO: 48	moltype = DNA length = 63	
FEATURE	Location/Qualifiers	
misc_feature	1..63	
	note = CD8 signal peptide	
source	1..63	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 48		
atggccctgc ccgtcaccgc cctgtgctg cccctggctc tgctgtctga cgccgcaaga		60
ccc		63
SEQ ID NO: 49	moltype = DNA length = 63	
FEATURE	Location/Qualifiers	
misc_feature	1..63	
	note = CD8 signal peptide	
source	1..63	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 49		
atggctctgc ccgtgaccgc cctgtgctg cctctggctc tgctgtctga cgctgccgc		60
cca		63
SEQ ID NO: 50	moltype = AA length = 68	
FEATURE	Location/Qualifiers	
REGION	1..68	

-continued

```

source          note = CD8 hinge and transmembrane
                1..68
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 50
TTTPAPRPPT PAPTIASQPL SLRPEACRPA AGGAVHTRGL DFACDIYIWA PLAGTCGVLL 60
LSLVITLY                                           68

SEQ ID NO: 51      moltype = DNA length = 60
FEATURE           Location/Qualifiers
misc_feature       1..60
                  note = Variable heavy-variable light linker
source            1..60
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 51
ggtggtggtg gttctggtg tgggtggttct ggcggcgccg gctccggtg tggtggtacc 60

SEQ ID NO: 52      moltype = DNA length = 60
FEATURE           Location/Qualifiers
misc_feature       1..60
                  note = Variable heavy-variable light linker
source            1..60
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 52
ggaggaggag gaagcggagg aggaggatcc ggaggcgggg gatctggagg aggaggaagt 60

SEQ ID NO: 53      moltype = DNA length = 45
FEATURE           Location/Qualifiers
misc_feature       1..45
                  note = Variable heavy-variable light linker
source            1..45
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 53
ggaggaggag gatccggcgg aggaggctct gggggaggcg ggagt 45

SEQ ID NO: 54      moltype = DNA length = 60
FEATURE           Location/Qualifiers
misc_feature       1..60
                  note = Variable heavy-variable light linker
source            1..60
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 54
ggaggaggag gaagtggagg aggaggatca ggaggcgggg gaagcggcgg gggaggctcc 60

SEQ ID NO: 55      moltype = DNA length = 60
FEATURE           Location/Qualifiers
misc_feature       1..60
                  note = Variable heavy-variable light linker
source            1..60
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 55
ggaggaggag gaagtggagg aggagggtca ggaggcgggg gaagcggcgg gggaggatcc 60

SEQ ID NO: 56      moltype = DNA length = 60
FEATURE           Location/Qualifiers
misc_feature       1..60
                  note = Variable heavy-variable light linker
source            1..60
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 56
ggaggaggag gatctggagg aggaggcagc ggcggcgccg gctccggcgg cggcggctct 60

SEQ ID NO: 57      moltype = DNA length = 210
FEATURE           Location/Qualifiers
misc_feature       1..210
                  note = CD8 hinge and transmembrane
source            1..210
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 57
aagcccacca cgacgccagc gccgcgacca ccaacaccgg cgcccaccat cgcgtcgcag 60
cccctgtccc tgcgcccaga gccgtgccgg ccagcggcgg ggggcgcagt gcacacgagg 120
gggctggact tcgcctgtga tatctacatc tgggcgcctt tggccgggac ttgtggggtc 180

```

-continued

```

cttctcctgt cactggttat caccctttac 210

SEQ ID NO: 58      moltype = DNA  length = 204
FEATURE           Location/Qualifiers
misc_feature       1..204
                   note = CD8 hinge and transmembrane
source            1..204
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 58
accactacac ctgcaccaag gctcccccaca cccgctccca ctatcgettc ccagccactg 60
tccttgaggc ccgaggcctg caggccagca gctggcggag ccgtgcatac tagggggctg 120
gacttcgctt gcgacatcta catctgggcc ccactggcag ggacatgcgg agtcctgctg 180
ctgtccctgg tcatcacact ttac 204

SEQ ID NO: 59      moltype = DNA  length = 204
FEATURE           Location/Qualifiers
misc_feature       1..204
                   note = CD8 hinge and transmembrane
source            1..204
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 59
actaccacac cagctccaag accacctacc cctgcaccaa caattgctag tcagccactg 60
tcactgagac cagaagcatg taggctgca gctggaggag ctgtgcacac cagaggcctg 120
gactttgctt gcgatatcta catctgggct cctctggcag gaacctgtgg cgtgtgctg 180
ctgtctctgg tcatcacact ttac 204

SEQ ID NO: 60      moltype = DNA  length = 210
FEATURE           Location/Qualifiers
misc_feature       1..210
                   note = CD8 hinge and transmembrane
source            1..210
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 60
aagccaacca caaccctgc accaaggcca cctacaccag cacctaccat cgcaagccag 60
ccactgtccc tgaggccaga ggcagttagg cctgcagcag gaggcgccgt gcacacacgc 120
ggcctggact ttgcctgcga tatctacatc tgggcaccac tggcaggaac ctgtggcgtg 180
ctgtctgtga gcctggtgat taccctgtat 210

SEQ ID NO: 61      moltype = DNA  length = 30
FEATURE           Location/Qualifiers
misc_feature       1..30
                   note = Variable heavy-variable light linker
source            1..30
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 61
ggtggtggcg gcagtgggtg cggtggtctca 30

SEQ ID NO: 62      moltype = AA  length = 10
FEATURE           Location/Qualifiers
REGION            1..10
                   note = Variable heavy-variable light linker
source            1..10
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 62
GGGSGGGGS 10

SEQ ID NO: 63      moltype = DNA  length = 60
FEATURE           Location/Qualifiers
misc_feature       1..60
                   note = Variable heavy-variable light linker
source            1..60
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 63
ggtggtggcg gcagtgggtg cggtggtctca ggcggtggtg gctccggtgg cggtggtctc 60

SEQ ID NO: 64      moltype = AA  length = 6
FEATURE           Location/Qualifiers
REGION            1..6
                   note = Variable heavy-variable light linker

```

-continued

```

source          1..6
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 64
EEKKMP

```

6

What is claimed is:

1. A composition comprising one or more recombinant polynucleotide molecules, the composition comprising

a first recombinant polynucleotide sequence encoding a chimeric antigen receptor (CAR), wherein the CAR comprises a first binding domain and a signaling domain; and

a second recombinant polynucleotide sequence encoding a target-binding molecule linked to a localizing domain, wherein the target-binding molecule linked to the localizing domain comprises a second binding domain and the localizing domain comprising an ER retention sequence, a Golgi retention sequence, or a PEST sequence,

wherein the second recombinant polynucleotide sequence comprises, from 5' terminus to 3' terminus, a first nucleotide sequence encoding the second binding domain, a second nucleotide sequence encoding a peptide linker, and a third nucleotide sequence encoding the localizing domain, and wherein the peptide linker comprises at least 5 amino acids, and

wherein the first binding domain is capable of binding to a molecule expressed on the surface of a cancer cell that is also expressed in an immune cell, and the second binding domain is capable of binding to the same molecule and suppressing its expression in the immune cell.

2. The composition of claim 1, wherein the first recombinant polynucleotide sequence and the second recombinant polynucleotide sequence are on the same polynucleotide molecule.

3. The composition of claim 2, further comprising an internal ribosomal entry site or a 2A peptide-coding region site between the first recombinant polynucleotide sequence encoding the CAR and the second recombinant polynucleotide sequence encoding the target-binding molecule linked to a localizing domain.

4. The composition of claim 1, wherein the localizing domain comprises an ER retention sequence.

5. The composition of claim 4, wherein the ER retention sequence comprises a KDEL amino acid sequence.

6. The composition of claim 4, wherein the ER retention sequence comprises a KKXX amino acid sequence, wherein X represents any amino acid.

7. The composition of claim 1, wherein the localizing domain comprises a Golgi retention sequence.

8. The composition of claim 7, wherein the Golgi retention sequence comprises a YQRL amino acid sequence.

9. The composition of claim 1, wherein the localizing domain comprises a PEST sequence.

10. The composition of claim 1, wherein the polynucleotide is a vector.

11. The composition of claim 1, wherein the polynucleotide is a lentiviral vector.

12. The composition of claim 1, wherein the first or the second binding domain comprises a single chain binding domain.

13. The composition of claim 12, wherein the single chain binding domain is a single chain variable fragment (scFv).

14. The composition of claim 1, wherein the target-binding molecule linked to a localizing domain further comprises a transmembrane domain.

15. The composition of claim 14, wherein the transmembrane domain is a CD8 transmembrane domain.

16. The composition of claim 1, wherein the second binding domain is identical to the first binding domain.

17. The composition of claim 1, wherein the second binding domain and the first binding domain are different.

18. The composition of claim 1, wherein the signaling domain comprises a CD3 ζ , a Fe, a DAP10, or a DAP12 domain.

19. The composition of claim 1, wherein the CAR further comprises a co-stimulatory domain.

20. The composition of claim 1, wherein the second recombinant polynucleotide sequence further comprises a nucleotide sequence encoding a signal peptide.

21. The composition of claim 1, wherein the first binding domain is a first single chain variable fragment (scFv) and the second binding domain is a second scFv.

* * * * *