

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250283154

Kind Code

A1

Publication Date

September 11, 2025

Inventor(s)

Ramachandran Iyer; Eswar Prasad et al.

PROFILING OF BIOLOGICAL ANALYTES WITH SPATIALLY BARCODED OLIGONUCLEOTIDE ARRAYS

Abstract

This disclosure relates to methods for spatial profiling of analytes present in a biological sample. Also provided are methods for using spatially barcoded substrates to detect a biological analyte in a cell culture, an organism, and organoid. Also provided are methods for using spatially barcoded substrates to detect the temporal profile of a biological analyte.

Inventors: Ramachandran Iyer; Eswar Prasad (Oakland, CA), Mikkelsen; Tarjei Sigurd (Hayward, CA), Tentori; Augusto Manuel (Dublin, CA), Bharadwaj; Rajiv (Pleasanton, CA), Stoeckius; Marlon (Stockholm, SE), Chell; James Michael (Stockholm, SE), Uytingco; Cedric (Milpitas, CA)

Applicant: 10x Genomics, Inc. (Pleasanton, CA)

Family ID: 1000008629075

Appl. No.: 18/746646

Filed: June 18, 2024

Related U.S. Application Data

parent US continuation 18471948 20230921 ABANDONED child US 18746646
parent US continuation 17869636 20220720 ABANDONED child US 18471948
parent US continuation 17459931 20210827 ABANDONED child US 17869636
parent US continuation PCT/US2020/020193 20200227 PENDING child US 17459931
us-provisional-application US 62812219 20190228
us-provisional-application US 62819458 20190315
us-provisional-application US 62819449 20190315
us-provisional-application US 62819478 20190315
us-provisional-application US 62819456 20190315
us-provisional-application US 62819448 20190315
us-provisional-application US 62819495 20190315

us-provisional-application US 62822722 20190322
us-provisional-application US 62822680 20190322
us-provisional-application US 62822606 20190322
us-provisional-application US 62822575 20190322
us-provisional-application US 62822554 20190322
us-provisional-application US 62822566 20190322
us-provisional-application US 62822649 20190322
us-provisional-application US 62822622 20190322
us-provisional-application US 62822627 20190322
us-provisional-application US 62822592 20190322
us-provisional-application US 62822565 20190322
us-provisional-application US 62839264 20190426
us-provisional-application US 62839575 20190426
us-provisional-application US 62839212 20190426
us-provisional-application US 62839320 20190426
us-provisional-application US 62839223 20190426
us-provisional-application US 62839294 20190426
us-provisional-application US 62858331 20190607
us-provisional-application US 62924241 20191022
us-provisional-application US 62925578 20191024
us-provisional-application US 62931587 20191106
us-provisional-application US 62931779 20191106
us-provisional-application US 62933299 20191108
us-provisional-application US 62933318 20191108
us-provisional-application US 62933878 20191111
us-provisional-application US 62934356 20191112
us-provisional-application US 62934883 20191113
us-provisional-application US 62935043 20191113
us-provisional-application US 62939488 20191122
us-provisional-application US 62941581 20191127
us-provisional-application US 62959526 20200110

Publication Classification

Int. Cl.: **C12Q1/6841** (20180101); **C12Q1/6874** (20180101); **C12Q1/6886** (20180101);
G16B30/10 (20190101)

U.S. Cl.:

CPC **C12Q1/6841** (20130101); **C12Q1/6874** (20130101); **C12Q1/6886** (20130101);
G16B30/10 (20190201);

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS [0001] This application is a continuation of a U.S. application Ser. No. 18/471,948, filed Sep. 21, 2023, which is a continuation of a U.S. application Ser. No. 17/869,636, filed Jul. 20, 2022, which is a continuation of a U.S. application Ser. No. 17/459,931, filed on Aug. 27, 2021, which claims priority to International Application PCT/US2020/020193, with an international filing date of Feb. 27, 2020, which claims the benefit

of U.S. Provisional Patent Application No. 62/812,219, filed Feb. 28, 2019, U.S. Provisional Patent Application No. 62/819,495, filed Mar. 15, 2019, U.S. Provisional Patent Application No. 62/819,448, filed Mar. 15, 2019, U.S. Provisional Patent Application No. 62/819,456, filed Mar. 15, 2019, U.S. Provisional Patent Application No. 62/819,478, filed Mar. 15, 2019, U.S. Provisional Patent Application No. 62/819,449, filed Mar. 15, 2019, U.S. Provisional Patent Application No. 62/819,458, filed Mar. 15, 2019, U.S. Provisional Patent Application No. 62/822,610, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,565, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,592, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,627, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,622, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,649, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,566, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,554, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,575, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,605, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,606, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,680, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/822,722, filed Mar. 22, 2019, U.S. Provisional Patent Application No. 62/839,294, filed Apr. 26, 2019, U.S. Provisional Patent Application No. 62/839,223, filed Apr. 26, 2019, U.S. Provisional Patent Application No. 62/839,320, filed Apr. 26, 2019, U.S. Provisional Patent Application No. 62/839,212, filed Apr. 26, 2019, U.S. Provisional Patent Application No. 62/839,575, filed Apr. 26, 2019, U.S. Provisional Patent Application No. 62/839,264, filed Apr. 26, 2019, U.S. Provisional Patent Application No. 62/858,331, filed Jun. 7, 2019, U.S. Provisional Patent Application No. 62/924,241, filed Oct. 22, 2019, U.S. Provisional Patent Application No. 62/925,578, filed Oct. 24, 2019, U.S. Provisional Patent Application No. 62/931,587, filed Nov. 6, 2019, U.S. Provisional Patent Application No. 62/931,779, filed Nov. 6, 2019, U.S. Provisional Patent Application No. 62/933,318, filed Nov. 8, 2019, U.S. Provisional Patent Application No. 62/933,299, filed Nov. 8, 2019, U.S. Provisional Patent Application No. 62/933,878, filed Nov. 11, 2019, U.S. Provisional Patent Application No. 62/934,356, filed Nov. 12, 2019, U.S. Provisional Patent Application No. 62/935,043, filed Nov. 13, 2019, U.S. Provisional Patent Application No. 62/934,883, filed Nov. 13, 2019, U.S. Provisional Patent Application No. 62/939,488, filed Nov. 22, 2019, U.S. Provisional Patent Application No. 62/941,581, filed Nov. 27, 2019, and U.S. Provisional Patent Application No. 62/959,526, filed Jan. 10, 2020. The contents of these applications are incorporated by reference in their entireties.

SEQUENCE LISTING

[0002] This application contains a Sequence Listing that has been submitted electronically as an XML file named “47706-0074005_SL_ST26.XML.” The XML file, created on Oct. 9, 2024, is 7,167 bytes in size. The material in the XML file is hereby incorporated by reference in its entirety.

BACKGROUND

[0003] Cells within a tissue of a subject have differences in cell morphology and/or function due to varied analyte levels (e.g., gene and/or protein expression) within the different cells. The specific position of a cell within a tissue (e.g., the cell's position relative to neighboring cells or the cell's position relative to the tissue microenvironment) can affect, e.g., the cell's morphology, differentiation, fate, viability, proliferation, behavior, and signaling and cross-talk with other cells in the tissue.

[0004] Spatial heterogeneity has been previously studied using techniques that only provide data for a small handful of analytes in the context of an intact tissue or a portion of a tissue, or provide a lot of analyte data for single cells, but fail to provide information regarding the position of the single cell in a parent biological sample (e.g., tissue sample).

[0005] Genetic material, and related gene and protein expression, influences cellular fate and behavior. The spatial heterogeneity in developing systems has typically been studied via RNA

hybridization, immunohistochemistry, fluorescent reporters, or purification or induction of pre-defined subpopulations and subsequent genomic profiling (e.g., RNA-seq). Such approaches, however, rely on a small set of pre-defined markers, therefore introducing selection bias that limits discovery and increases the cost and labor required to localize RNA on transcriptome-wide basis.

SUMMARY

[0006] Provided herein are methods for identifying a location of a biological analyte in a biological sample, the method comprising: (a) providing a biological sample comprising live cells; (b) contacting the biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises (i) a spatial barcode and (ii) a capture domain that binds specifically to a biological analyte in the biological sample; and (c) determining (i) all or a part of the sequence corresponding to the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample.

[0007] Provided herein are methods of identifying a location of a biological analyte in a biological sample, the method comprising: (a) providing a biological sample comprising live cells; (b) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises (i) a spatial barcode and (ii) a capture domain that binds specifically to the analyte capture sequence; (c) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample.

[0008] Provided herein are methods for identifying a location of a biological analyte in a biological sample, the method comprising: (a) contacting the biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises (i) a spatial barcode and (ii) a capture domain that binds specifically to a biological analyte in the biological sample; and (b) determining (i) all or a part of the sequence corresponding to the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample.

[0009] In some embodiments, the biological sample is an organism. In some embodiments, the biological sample is an organoid. In some embodiments, the biological sample is an in vitro cell culture comprising a plurality of live cells.

[0010] In some embodiments, the biological analyte is selected from the group consisting of: CD45, CD3, CD4, CD8, CD56, CD19, CD20, CD11c, CD14, CD33, CD66b, CD34, CD41, CD61, CD235a, CD146, and epithelial cellular adhesion molecule (EpCAM).

[0011] In some embodiments, a live cell of the plurality of live cells is distributed onto the substrate. In some embodiments, a live cell of the plurality of live cells is distributed onto a second substrate. In some embodiments, the second substrate is contacted with the substrate. In some embodiments, a live cell of the plurality of live cells is immobilized by adding a hydrogel. In some embodiments, a live cell of the plurality of live cells is immobilized by applying an electric field.

[0012] Provided herein are methods for determining presence of a biological analyte in a live cell at a location on a substrate, comprising: (a) allowing a live cell of a plurality of live cells to migrate on a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises (i) a spatial barcode and (ii) a capture domain that specifically binds to the biological analyte; (b) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the

sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to determine the biological analyte was present in the live cell at a location on the substrate. [0013] Provided herein are methods for determining presence of a biological analyte in a live cell at a location on a substrate, comprising: (a) contacting a plurality of analyte capture agents to a live cell of a plurality of live cells disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; (b) allowing the live cell of the plurality of live cells to migrate on a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain that binds specifically to the analyte capture sequence; (c) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to determine that the biological analyte was present in the live cell at a location on the substrate.

[0014] In some embodiments, the migration is passive migration. In some embodiments, the migration is active migration. In some embodiments, the active migration is induced by a chemoattractant gradient. In some embodiments, the substrate comprises one or more channels. In some embodiments, the one or more channels confine cell migration along one axis of the substrate. In some embodiments, the one or more channels confine cell migration along more than one axis of the substrate. In some embodiments, the one or more channels confine cell migration along two axes of the substrate.

[0015] Provided herein are methods for identifying a location of a biological analyte in a biological sample, comprising: (a) contacting a biological sample comprising a cell with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from a subcellular region of the cell, wherein the biological analyte is specifically bound by the capture probe; and (c) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample.

[0016] Provided herein are methods for identifying a location of a biological analyte in a biological sample, comprising: (a) contacting a plurality of analyte capture agents to a biological sample comprising a live cell, wherein the biological sample is disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture proteins comprises (i) a spatial barcode, and (ii) a capture domain; (b) releasing a biological analyte from a subcellular region of the cell, wherein the analyte capture sequence is specifically bound by the capture domain; and (c) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample.

[0017] In some embodiments, the biological analyte is released from the subcellular region of the cell wherein release of other biological analytes from other subcellular regions of the cell is substantially prevented. In some embodiments, the subcellular region is cytosol. In some embodiments, the subcellular region is a nucleus. In some embodiments, the subcellular region is a mitochondria. In some embodiments, the subcellular region is a microsome.

[0018] Provided herein are methods for temporal and spatial profiling of a biological analyte present in a biological sample, comprising: (a) contacting a biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes

comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from the biological sample, wherein the capture domain binds specifically to the released biological analyte; (c) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (a)-(c) after a period of time. Further provided herein are methods for temporal and spatial profiling of a biological analyte present in a biological sample, comprising: (a) contacting a plurality of analyte capture agents to a biological sample comprising a live cell, wherein the biological sample is disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture proteins comprises (i) a spatial barcode, and (ii) a capture domain; (b) releasing a biological analyte from the live cell, wherein the analyte capture sequence is specifically bound by the capture domain; (c) determining (i) all or a part of the sequence of the analyte binding moiety barcode specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (a)-(c) after a period of time.

[0019] Provided herein are methods for temporal and spatial profiling of a biological analyte present in a biological sample, comprising: (a) contacting a biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from the biological sample, wherein the capture domain binds specifically to the released biological analyte; (c) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (b)-(c) after a period of time. Further provided herein are methods for temporal and spatial profiling of a biological analyte present in a biological sample, comprising: (a) contacting a plurality of analyte capture agents to a biological sample comprising a live cell, wherein the biological sample is disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture proteins comprises (i) a spatial barcode, and (ii) a capture domain; (b) releasing a biological analyte from the live cell, wherein the analyte capture sequence is specifically bound by the capture domain; (c) determining (i) all or a part of the sequence of the analyte binding moiety barcode specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (b)-(c) after a period of time.

[0020] In some embodiments, the biological sample comprises a tissue, an organ, an organism, an organoid, or a cell culture sample. In some embodiments, prior to the step of repeating, the biological sample is subjected to one or more conditions. In some embodiments, the one or more conditions include application of one or more reagents, application of light, a change in temperature, a change in pressure, application of an electric field, application of a magnetic field, and combinations thereof. In some embodiments, the one or more reagents comprises a DNA modification system. In some embodiments, the DNA modification system is a CRISPR system.

[0021] In some embodiments, steps (a) through (c) are repeated one, two, three, or four times. In some embodiments, the substrate comprises a cell-permissive coating. In some embodiments, the releasing step comprises permeabilizing one or more cells of the biological sample. In some embodiments, the permeabilizing comprises contacting the biological sample with a hydrogel that comprises the permeabilization reagent.

[0022] In some embodiments, the biological analyte is DNA. In some embodiments, the biological analyte is RNA. In some embodiments, the capture domain comprises a poly-dT sequence. In some embodiments, the capture probe further comprises a unique molecular identifier (UMI). In some embodiments, the capture probe further comprises a functional domain. In some embodiments, the capture probe further comprises a cleavage domain. In some embodiments, any one of the methods described herein further comprises washing the sample. In some embodiments, any one of the methods described herein, further comprises imaging the biological sample.

[0023] Provided herein are methods of identifying the location of a biological analyte in a biological sample comprising: (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain that binds specifically to the analyte capture sequence; (b) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample; and (c) performing immunofluorescence microscopy on the biological sample to detect the location of an additional biological analyte in the biological sample.

[0024] Provided herein are methods of identifying a location of a biological analyte in a biological sample comprising: (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence, wherein the analyte capture sequence is hybridized to a blocking probe; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; (b) releasing the blocking probe from the analyte capture sequence, thereby allowing the analyte capture sequence to bind specifically to the capture agent; and (c) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample.

[0025] In some embodiments, step (b) comprises heating the analyte capture agent to release the blocking probe. In some embodiments, step (b) comprises contacting the analyte capture agent with an enzyme to release the blocking probe. In some embodiments, the enzyme to release the blocking probe is RNase H. In some embodiments, the method further comprises facilitating movement or migration of the analyte capture agent to the substrate. In some embodiments, the method further comprises imaging the biological sample.

[0026] In some embodiments, the substrate is a bead array. In some embodiments, the analyte capture sequence comprises a nucleic acid sequence. In some embodiments, the nucleic acid sequence of the analyte capture sequence comprises a sequence that is substantially complementary to the blocking probe. In some embodiments, the nucleic acid sequence of the analyte capture sequence comprises a sequence that is completely complementary to the blocking probe. In some embodiments, the nucleic acid sequence of the analyte capture sequence comprises a poly(A) sequence. In some embodiments, the nucleic acid sequence of the analyte capture sequence has a

GC content of at least 30%. In some embodiments, the analyte capture agent comprises a label. In some embodiments, the label is a fluorophore.

[0027] In some embodiments, the biological analyte is a protein. In some embodiments, the analyte binding moiety comprises an antibody or an antigen-binding domain thereof. In some embodiments, the method further comprises performing immunofluorescence microscopy on the biological sample to detect the location of an additional biological analyte in the biological sample. In some embodiments, the additional biological analyte is a protein. In some embodiments, the method further comprises generating an image of the location of the biological analyte in the biological sample, and generating an image of the location of the additional biological analyte in the biological sample. In some embodiments, the method further comprises comparing or overlying the image of the location of the biological analyte in the biological sample and the image of the location of the additional biological analyte in the biological sample. In some embodiments, the biological analyte is a protein and the method further comprises performing immunofluorescence microscopy on the biological sample to detect the location of an additional biological analyte in the biological sample, wherein the additional biological analyte is a nucleic acid. In some embodiments, the additional biological analyte is a nucleic acid encoding the biological analyte.

[0028] In some embodiments, the method further comprises generating an image of the location of the biological analyte and an image of the location of the additional biological analyte in the biological sample. In some embodiments, the method further comprises comparing or overlying the image of the location of the biological analyte and the image of the location of the additional biological analyte in the biological sample. In some embodiments, the analyte capture agent comprises an analyte binding moiety linked to the capture agent barcode domain by a cleavable domain. In some embodiments, the method further comprises, before step (a), permeabilizing the biological sample. In some embodiments, any one of the methods described herein further comprises imaging the substrate. In some embodiments, any one of the methods described herein further comprises performing gene expression analysis. In some embodiments, the gene expression analysis comprises performing high-throughput sequencing. In some embodiments, the imaging comprises the use of fiducial markers.

[0029] In some embodiments, the biological sample comprises a tissue, a tissue section, an organ, an organism, an organoid, or a cell culture sample. In some embodiments, the biological sample is fixed prior to step (a). In some embodiments, the biological sample is fixed with formaldehyde. In some embodiments, the biological sample is fixed with methanol.

[0030] All publications, patents, patent applications, and information available on the internet and mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, patent application, or item of information was specifically and individually indicated to be incorporated by reference. To the extent publications, patents, patent applications, and items of information incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

[0031] Where values are described in terms of ranges, it should be understood that the description includes the disclosure of all possible sub-ranges within such ranges, as well as specific numerical values that fall within such ranges irrespective of whether a specific numerical value or specific sub-range is expressly stated.

[0032] The term “each,” when used in reference to a collection of items, is intended to identify an individual item in the collection but does not necessarily refer to every item in the collection, unless expressly stated otherwise, or unless the context of the usage clearly indicates otherwise.

[0033] Various embodiments of the features of this disclosure are described herein. However, it should be understood that such embodiments are provided merely by way of example, and numerous variations, changes, and substitutions can occur to those skilled in the art without

departing from the scope of this disclosure. It should also be understood that various alternatives to the specific embodiments described herein are also within the scope of this disclosure.

Description

DESCRIPTION OF DRAWINGS

[0034] The following drawings illustrate certain embodiments of the features and advantages of this disclosure. These embodiments are not intended to limit the scope of the appended claims in any manner. Like reference symbols in the drawings indicate like elements.

[0035] FIG. **1** shows an exemplary spatial analysis workflow.

[0036] FIG. **2** shows an exemplary spatial analysis workflow.

[0037] FIG. **3** shows an exemplary spatial analysis workflow.

[0038] FIG. **4** shows an exemplary spatial analysis workflow.

[0039] FIG. **5** shows an exemplary spatial analysis workflow.

[0040] FIG. **6** is a schematic diagram showing an example of a barcoded capture probe, as described herein.

[0041] FIG. **7** is a schematic illustrating a cleavable capture probe, wherein the cleaved capture probe can enter into a non-permeabilized cell and bind to target analytes within the sample.

[0042] FIG. **8** is a schematic diagram of an exemplary multiplexed spatially-barcoded feature.

[0043] FIG. **9** is a schematic diagram of an exemplary analyte capture agent.

[0044] FIG. **10** is a schematic diagram depicting an exemplary interaction between a feature-immobilized capture probe **1024** and an analyte capture agent **1026**.

[0045] FIGS. **11A**, **11B**, and **11C** are schematics illustrating how streptavidin cell tags can be utilized in an array-based system to produce a spatially-barcoded cells or cellular contents.

[0046] FIG. **12** is a schematic showing the arrangement of barcoded features within an array.

[0047] FIG. **13** is a schematic illustrating a side view of a diffusion-resistant medium, e.g., a lid.

[0048] FIGS. **14A** and **14B** are schematics illustrating expanded FIG. **14A** and side views FIG. **14B** of an electrophoretic transfer system configured to direct transcript analytes toward a spatially-barcoded capture probe array.

[0049] FIG. **15** is a schematic illustrating an exemplary workflow protocol utilizing an electrophoretic transfer system.

[0050] FIG. **16** shows an example of a microfluidic channel structure **1600** for partitioning dissociated sample (e.g., biological particles or individual cells from a sample).

[0051] FIG. **17A** shows an example of a microfluidic channel structure **1700** for delivering spatial barcode carrying beads to droplets.

[0052] FIG. **17B** shows a cross-section view of another example of a microfluidic channel structure **1750** with a geometric feature for controlled partitioning.

[0053] FIG. **17C** shows an example of a workflow schematic.

[0054] FIG. **18** is a schematic depicting cell tagging using either covalent conjugation of the analyte binding moiety to the cell surface or non-covalent interactions with cell membrane elements.

[0055] FIG. **19** is a schematic depicting cell tagging using either cell-penetrating peptides or delivery systems.

[0056] FIG. **20A** is a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for “pixelating” a sample, wherein the sample is cut, stamped, microdissected, or transferred by hollow-needle or microneedle, moving a small portion of the sample into an individual partition or well.

[0057] FIG. **20B** is a schematic depicting multi-needle pixilation, wherein an array of needles punched through a sample on a scaffold and into nanowells containing gel beads and reagents

below. Once the needle is in the nanowell, the cell(s) are ejected.

[0058] FIG. **21** shows a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for dissociating a spatially-barcoded sample for analysis via droplet or flow cell analysis methods.

[0059] FIG. **22A** is a schematic diagram showing an example sample handling apparatus that can be used to implement various steps and methods described herein.

[0060] FIG. **22B** is a schematic diagram showing an example imaging apparatus that can be used to obtain images of biological samples, analytes, and arrays of features.

[0061] FIG. **22C** is a schematic diagram of an example of a control unit of the apparatus of FIGS. **22A** and **22B**.

[0062] FIG. **23A** shows a histological section of an invasive ductal carcinoma annotated by a pathologist.

[0063] FIG. **23B** shows a tissue plot with spots colored by unsupervised clustering.

[0064] FIG. **23C** is a tSNE plot of spots colored by unsupervised clustering.

[0065] FIG. **23D** shows a gene expression heat map of the most variable genes between 9 clusters.

[0066] FIG. **23E** shows the expression levels of genes corresponding to human epidermal growth factor receptor 2 (ERBB2), estrogen receptor (ESR1), and progesterone receptor (PGR) in the tissue section.

[0067] FIG. **23F** shows the expression levels of genes of top differentially expressed genes from each of the 9 clusters on individual plots.

[0068] FIG. **23G** shows the expression levels of genes of top differentially expressed genes from each of the 5 clusters on a single plot.

[0069] FIG. **23H** is a plot of the expression levels of the top differentially expressed genes from each of the 8 clusters in invasive ductal cell carcinoma (IDC) and normal breast tissue.

[0070] FIG. **23I** shows the expression of KRT14 in IDC and match normal tissue.

[0071] FIG. **23J** is a plot of the expression levels of extracellular matrix genes in IDC and normal tissue.

[0072] FIG. **24A** shows a schematic of an example analytical workflow in which electrophoretic migration of analytes is performed after permeabilization.

[0073] FIG. **24B** shows a schematic of an example analytical workflow in which electrophoretic migration of analytes and permeabilization are performed simultaneously.

[0074] FIG. **25A** shows an example perpendicular, single slide configuration for use during electrophoresis.

[0075] FIG. **25B** shows an example parallel, single slide configuration for use during electrophoresis.

[0076] FIG. **25C** shows an example multi-slide configuration for use during electrophoresis.

[0077] FIGS. **26A** and **26B** is a schematic depicting exemplary steps for adding a biological sample to an array.

[0078] FIG. **27** is a schematic showing exemplary biological samples that can be used in the methods described herein.

[0079] FIG. **28** is a schematic depicting a cell migration assay combined with an array such as those described herein.

[0080] FIG. **29A** is a schematic showing a high density array with cells bound to spots having barcoded capture probes and cell capture molecules (e.g., a capture antibody).

[0081] FIG. **29B** is a schematic showing a portion of a spot on the array of FIG. **29A** with barcoded capture probes and a capture antibody.

[0082] FIG. **30** is a schematic illustrating exemplary steps involved in subcellular fractionation.

[0083] FIG. **31** is a schematic illustrating exemplary steps involved in subcellular fractionation.

[0084] FIG. **32A** is a schematic showing an exemplary substrate and capture probe for detecting a selectively releasing a biological analyte using, for example, second strand synthesis.

[0085] FIG. **32B** is a schematic depicting steps for second strand synthesis.

[0086] FIG. **33A** is a schematic showing an exemplary substrate and capture probes for detecting a selectively-released biological analyte using, for example, selective probe cleavage.

[0087] FIG. **33B** is a schematic depicting steps for using selective probe cleavage in methods involving detecting a selectively-released biological analyte.

[0088] FIG. **34** is a schematic depicting steps for using multiple substrates in methods involving detecting a selectively-released biological sample.

[0089] FIG. **35** is a schematic depicting steps for temporal analysis of a biological analyte.

[0090] FIG. **36** shows an exemplary workflow for temporal and spatial profiling of biological analytes at n time-points.

[0091] FIG. **37** shows a schematic representation of spatio-temporal profiling of biological analytes.

[0092] FIG. **38** is a schematic illustrating fluorescently-labeled antibodies interacting with analytes within a biological sample, wherein the biological sample is contacting a capture probe array.

[0093] FIG. **39** is a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for immunofluorescently detecting analytes on a capture probe array, wherein the biological sample is fixed, stained with fluorescent antibodies, and imaged to detect the spatial location of target analytes within the biological sample.

[0094] FIG. **40** shows exemplary antibody staining wherein tissue sections of mouse spleen are fixed in either 2% formaldehyde or methanol. In both the 2% formaldehyde and methanol fixation conditions, CD3 can be seen in the pockets of white pulp, and CD29 in the red pulp portion of the spleen.

[0095] FIGS. **41A** and **41B** shows exemplary tissue plot of spatially-resolved information, wherein a mouse spleen tissue section is analyzed on a barcoded feature array. Here, analytes are captured on a barcoded feature array as described herein and captured analytes are clustered using a nonlinear dimensionality reduction technique. The clusters can identify cellular subpopulations within a tissue sample.

[0096] FIG. **42** shows exemplary immunofluorescence staining of 2% formaldehyde of mouse spleen on the left, identifying the presence and location of CD3, CD29, and DAPI. This immunofluorescence staining is overlaid with CD3 UMI counts (right), obtained by performing the spatial workflows described herein. The CD3 UMI counts (right) demonstrates that specific analytes can be captured and barcoded according to the spatial workflows described herein, wherein the spatial workflows are compatible with and complementary to other staining/imaging techniques (e.g., immunofluorescence).

[0097] FIG. **43** shows exemplary immunofluorescence staining of methanol of mouse spleen on the left, identifying the presence and location of CD3, CD29, and DAPI. This immunofluorescence staining is overlaid with CD3 UMI counts (right), obtained by performing the spatial workflows described herein. The CD3 UMI counts (right) demonstrates that specific analytes can be captured and barcoded according to the spatial workflows described herein, wherein the spatial workflows are compatible with and complementary to other staining/imaging techniques (e.g., immunofluorescence).

[0098] FIG. **44** shows exemplary immunofluorescence staining of 2% formaldehyde and methanol fixation of mouse brain in three channels (NeuN; GFAP; DAPI).

[0099] FIGS. **45A** and **45B** shows exemplary UMI and cluster data of mouse brain for 2% formaldehyde (FIG. **43A**) and methanol (FIG. **43B**) fixation conditions, wherein clusters were determined using t-SNE to identify different cell types.

[0100] FIG. **46** shows exemplary immunofluorescence staining of 2% formaldehyde fixation of mouse brain on the left. Right panels show immunofluorescence staining overlaid with NeuN UMI counts (top), and GFAP UMI counts (bottom), obtained by performing the spatial workflows described herein. The NeuN and GFAP UMI counts demonstrates that specific analytes can be

captured and barcoded according to the spatial workflows described herein, wherein the spatial workflows are compatible with and complementary to other staining/imaging techniques (e.g., immunofluorescence).

[0101] FIG. **47** shows exemplary immunofluorescence staining of methanol fixation of mouse brain on the left. Right panels show immunofluorescence staining overlaid with NeuN UMI counts (top), and GFAP UMI counts (bottom), obtained by performing the spatial workflows described herein. The NeuN and GFAP UMI counts demonstrates that specific analytes can be captured and barcoded according to the spatial workflows described herein, wherein the spatial workflows are compatible with and complementary to other staining/imaging techniques (e.g., immunofluorescence).

[0102] FIG. **48** is an exemplary schematic illustrating spatially-tagged analyte capture agents wherein the analyte capture sequence of the capture agent barcode domain (poly-A sequence) can be blocked with a blocking probe (poly-T oligonucleotide in upper left corner). Once the blocking probe is removed, the spatially-tagged analyte capture agent binds to the capture probe on the substrate and is labeled with fluorescent antibodies.

[0103] FIG. **49** is an exemplary schematic illustrating a spatially-tagged analyte capture agent wherein the analyte capture sequence is blocked via a blocking probe. The blocking probe can be removed, e.g., with an RNase treatment.

[0104] FIG. **50** shows exemplary images showing the performance of an oligo (dT) blocking probe under various conditions. The staining was performed with oligo (dT) blocking probes of varying lengths (15-30 nucleotides) at 4° C. and 37° C. and results indicate analyte capture agents were blocked from binding to the capture probes at both temperatures, although lower temperatures appeared more effective when using shorter blocking probes.

[0105] FIG. **51** is a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for spatial analyte identification after antibody staining in a biological sample, wherein the sample is fixed, stained with fluorescent antibodies and spatially-tagged analyte capture agents, and imaged to detect the spatial location of target analytes within the biological sample.

[0106] FIG. **52** shows an exemplary image showing immunofluorescence staining for CD29 (left panel) and images showing spatially-resolved information after analyzing the capture agent barcode domains for CD29, CD3, CD169, and F4/80 (right panels) in tissue sections of mouse spleen.

[0107] FIG. **53** shows exemplary multiplex imaging results, wherein the immunofluorescent image shows immunofluorescent staining for CD29 and CD4 in tissue sections of mouse spleen (far left). The images in the series of right panels show the results of a multiplexed, spatially-tagged analyte capture agent workflow, wherein the spatial location of target proteins, CD29, CD3, CD4, CD8, CD19, B220, F4/80, and CD169 are visualized by sequencing the analyte-corresponding analyte binding moiety barcodes.

DETAILED DESCRIPTION

I. Introduction

[0108] This disclosure describes apparatus, systems, methods, and compositions for spatial analysis of biological samples. This section describes certain general terminology, analytes, sample types, and preparative steps that are referred to in later sections of the disclosure.

(a) Spatial Analysis

[0109] Tissues and cells can be obtained from any source. For example, tissues and cells can be obtained from single-cell or multicellular organisms (e.g., a mammal). Tissues and cells obtained from a mammal, e.g., a human, often have varied analyte levels (e.g., gene and/or protein expression) which can result in differences in cell morphology and/or function. The position of a cell or a subset of cells (e.g., neighboring cells and/or non-neighboring cells) within a tissue can affect, e.g., the cell's fate, behavior, morphology, and signaling and cross-talk with other cells in the tissue. Information regarding the differences in analyte levels (gene and/or protein expression)

within different cells in a tissue of a mammal can also help physicians select or administer a treatment that will be effective and can allow researchers to identify and elucidate differences in cell morphology and/or cell function in the single-cell or multicellular organisms (e.g., a mammal) based on the detected differences in analyte levels within different cells in the tissue. Differences in analyte levels within different cells in a tissue of a mammal can also provide information on how tissues (e.g., healthy and diseased tissues) function and/or develop. Differences in analyte levels within different cells in a tissue of a mammal can also provide information of different mechanisms of disease pathogenesis in a tissue and mechanism of action of a therapeutic treatment within a tissue. Differences in analyte levels within different cells in a tissue of a mammal can also provide information on drug resistance mechanisms and the development of the same in a tissue of a mammal. Differences in the presence or absence of analytes within different cells in a tissue of a multicellular organism (e.g., a mammal) can provide information on drug resistance mechanisms and the development of the same in a tissue of a multicellular organism.

[0110] The spatial analysis methodologies herein provide for the detection of differences in an analyte level (e.g., gene and/or protein expression) within different cells in a tissue of a mammal or within a single cell from a mammal. For example, spatial analysis methodologies can be used to detect the differences in analyte levels (e.g., gene and/or protein expression) within different cells in histological slide samples, the data from which can be reassembled to generate a three-dimensional map of analyte levels (e.g., gene and/or protein expression) of a tissue sample obtained from a mammal, e.g., with a degree of spatial resolution (e.g., single-cell resolution).

[0111] Spatial heterogeneity in developing systems has typically been studied via RNA hybridization, immunohistochemistry, fluorescent reporters, or purification or induction of pre-defined subpopulations and subsequent genomic profiling (e.g., RNA-seq). Such approaches, however, rely on a relatively small set of pre-defined markers, therefore introducing selection bias that limits discovery. These prior approaches also rely on a priori knowledge. Spatial RNA assays traditionally relied on staining for a limited number of RNA species. In contrast, single-cell RNA-sequencing allows for deep profiling of cellular gene expression (including non-coding RNA), but the established methods separate cells from their native spatial context.

[0112] Spatial analysis methodologies described herein provide a vast amount of analyte level and/or expression data for a variety of multiple analytes within a sample at high spatial resolution, e.g., while retaining the native spatial context. Spatial analysis methods include, e.g., the use of a capture probe including a spatial barcode (e.g., a nucleic acid sequence that provides information as to the position of the capture probe within a cell or a tissue sample (e.g., mammalian cell or a mammalian tissue sample) and a capture domain that is capable of binding to an analyte (e.g., a protein and/or nucleic acid) produced by and/or present in a cell. As described herein, the spatial barcode can be a nucleic acid that has a unique sequence, a unique fluorophore or a unique combination of fluorophores, a unique amino acid sequence, a unique heavy metal or a unique combination of heavy metals, or any other unique detectable agent. The capture domain can be any agent that is capable of binding to an analyte produced by and/or present in a cell (e.g., a nucleic acid that is capable of hybridizing to a nucleic acid from a cell (e.g., an mRNA, genomic DNA, mitochondrial DNA, or miRNA), a substrate including an analyte, a binding partner of an analyte, or an antibody that binds specifically to an analyte). A capture probe can also include a nucleic acid sequence that is complementary to a sequence of a universal forward and/or universal reverse primer. A capture probe can also include a cleavage site (e.g., a cleavage recognition site of a restriction endonuclease), a photolabile bond, a thermosensitive bond, or a chemical-sensitive bond.

[0113] The binding of an analyte to a capture probe can be detected using a number of different methods, e.g., nucleic acid sequencing, fluorophore detection, nucleic acid amplification, detection of nucleic acid ligation, and/or detection of nucleic acid cleavage products. In some examples, the detection is used to associate a specific spatial barcode with a specific analyte produced by and/or

present in a cell (e.g., a mammalian cell).

[0114] Capture probes can be, e.g., attached to a surface, e.g., a solid array, a bead, or a coverslip. In some examples, capture probes are not attached to a surface. In some examples, capture probes can be encapsulated within, embedded within, or layered on a surface of a permeable composition (e.g., any of the substrates described herein). For example, capture probes can be encapsulated or disposed within a permeable bead (e.g., a gel bead). In some examples, capture probes can be encapsulated within, embedded within, or layered on a surface of a substrate (e.g., any of the exemplary substrates described herein, such as a hydrogel or a porous membrane).

[0115] In some examples, a cell or a tissue sample including a cell are contacted with capture probes attached to a substrate (e.g., a surface of a substrate), and the cell or tissue sample is permeabilized to allow analytes to be released from the cell and bind to the capture probes attached to the substrate. In some examples, analytes released from a cell can be actively directed to the capture probes attached to a substrate using a variety of methods, e.g., electrophoresis, chemical gradient, pressure gradient, fluid flow, or magnetic field.

[0116] In other examples, a capture probe can be directed to interact with a cell or a tissue sample using a variety of methods, e.g., inclusion of a lipid anchoring agent in the capture probe, inclusion of an agent that binds specifically to, or forms a covalent bond with a membrane protein in the capture probe, fluid flow, pressure gradient, chemical gradient, or magnetic field.

[0117] Non-limiting aspects of spatial analysis methodologies are described in WO 2011/127099, WO 2014/210233, WO 2014/210225, WO 2016/162309, WO 2018/091676, WO 2012/140224, WO 2014/060483, U.S. Pat. Nos. 10,002,316, 9,727,810, U.S. Patent Application Publication No. 2017/0016053, Rodriques et al., *Science* 363 (6434): 1463-1467, 2019; WO 2018/045186, Lee et al., *Nat. Protoc.* 10 (3): 442-458, 2015; WO 2016/007839, WO 2018/045181, WO 2014/163886, Trejo et al., *PLOS ONE* 14 (2): e0212031, 2019, U.S. Patent Application Publication No. 2018/0245142, Chen et al., *Science* 348 (6233): aaa6090, 2015, Gao et al., *BMC Biol.* 15:50, 2017, WO 2017/144338, WO 2018/107054, WO 2017/222453, WO 2019/068880, WO 2011/094669, U.S. Pat. Nos. 7,709,198, 8,604,182, 8,951,726, 9,783,841, 10,041,949, WO 2016/057552, WO 2017/147483, WO 2018/022809, WO 2016/166128, WO 2017/027367, WO 2017/027368, WO 2018/136856, WO 2019/075091, U.S. Pat. No. 10,059,990, WO 2018/057999, WO 2015/161173, and Gupta et al., *Nature Biotechnol.* 36:1197-1202, 2018, and can be used herein in any combination. Further non-limiting aspects of spatial analysis methodologies are described herein.

(b) General Terminology

[0118] Specific terminology is used throughout this disclosure to explain various aspects of the apparatus, systems, methods, and compositions that are described. This sub-section includes explanations of certain terms that appear in later sections of the disclosure. To the extent that the descriptions in this section are in apparent conflict with usage in other sections of this disclosure, the definitions in this section will control.

(i) Barcode

[0119] A “barcode” is a label, or identifier, that conveys or is capable of conveying information (e.g., information about an analyte in a sample, a bead, and/or a capture probe). A barcode can be part of an analyte, or independent of an analyte. A barcode can be attached to an analyte. A particular barcode can be unique relative to other barcodes.

[0120] Barcodes can have a variety of different formats. For example, barcodes can include non-random, semi-random, and/or random nucleic acid and/or amino acid sequences, and synthetic nucleic acid and/or amino acid sequences. A barcode can be attached to an analyte or to another moiety or structure in a reversible or irreversible manner. A barcode can be added to, for example, a fragment of a deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) sample before or during sequencing of the sample. Barcodes can allow for identification and/or quantification of individual sequencing-reads (e.g., a barcode can be or can include a unique molecular identifier or “UMI”).

[0121] Barcodes can spatially-resolve molecular components found in biological samples, for

example, at single-cell resolution (e.g., a barcode can be or can include a “spatial barcode”). In some embodiments, a barcode includes both a UMI and a spatial barcode. In some embodiments, a barcode includes two or more sub-barcode that together function as a single barcode (e.g., a polynucleotide barcode). For example, a polynucleotide barcode can include two or more polynucleotide sequences (e.g., sub-barcode) that may be separated by one or more non-barcode sequences.

(ii) Nucleic Acid and Nucleotide

[0122] The terms “nucleic acid” and “nucleotide” are intended to be consistent with their use in the art and to include naturally-occurring species or functional analogs thereof. Particularly useful functional analogs of nucleic acids are capable of hybridizing to a nucleic acid in a sequence-specific fashion (e.g., capable of hybridizing to two nucleic acids such that ligation can occur between the two hybridized nucleic acids) or are capable of being used as a template for replication of a particular nucleotide sequence. Naturally-occurring nucleic acids generally have a backbone containing phosphodiester bonds. An analog structure can have an alternate backbone linkage including any of a variety of those known in the art. Naturally-occurring nucleic acids generally have a deoxyribose sugar (e.g., found in deoxyribonucleic acid (DNA)) or a ribose sugar (e.g., found in ribonucleic acid (RNA)).

[0123] A nucleic acid can contain nucleotides having any of a variety of analogs of these sugar moieties that are known in the art. A nucleic acid can include native or non-native nucleotides. In this regard, a native deoxyribonucleic acid can have one or more bases selected from the group consisting of adenine (A), thymine (T), cytosine (C), or guanine (G), and a ribonucleic acid can have one or more bases selected from the group consisting of uracil (U), adenine (A), cytosine (C), or guanine (G). Useful non-native bases that can be included in a nucleic acid or nucleotide are known in the art.

(iii) Probe and Target

[0124] A “probe” or a “target,” when used in reference to a nucleic acid or sequence of a nucleic acids, is intended as a semantic identifier for the nucleic acid or sequence in the context of a method or composition, and does not limit the structure or function of the nucleic acid or sequence beyond what is expressly indicated.

(iv) Oligonucleotide and Polynucleotide

[0125] The terms “oligonucleotide” and “polynucleotide” are used interchangeably to refer to a single-stranded multimer of nucleotides from about 2 to about 500 nucleotides in length. Oligonucleotides can be synthetic, made enzymatically (e.g., via polymerization), or using a “split-pool” method. Oligonucleotides can include ribonucleotide monomers (i.e., can be oligoribonucleotides) and/or deoxyribonucleotide monomers (i.e., oligodeoxyribonucleotides). In some examples, oligonucleotides can include a combination of both deoxyribonucleotide monomers and ribonucleotide monomers in the oligonucleotide (e.g., random or ordered combination of deoxyribonucleotide monomers and ribonucleotide monomers). An oligonucleotide can be 4 to 10, 10 to 20, 21 to 30, 31 to 40, 41 to 50, 51 to 60, 61 to 70, 71 to 80, 80 to 100, 100 to 150, 150 to 200, 200 to 250, 250 to 300, 300 to 350, 350 to 400, or 400-500 nucleotides in length, for example. Oligonucleotides can include one or more functional moieties that are attached (e.g., covalently or non-covalently) to the multimer structure. For example, an oligonucleotide can include one or more detectable labels (e.g., a radioisotope or fluorophore).

(v) Subject

[0126] A “subject” is an animal, such as a mammal (e.g., human or a non-human simian), or avian (e.g., bird), or other organism, such as a plant. Examples of subjects include, but are not limited to, a mammal such as a rodent, mouse, rat, rabbit, guinea pig, ungulate, horse, sheep, pig, goat, cow, cat, dog, primate (i.e. human or non-human primate); a plant such as *Arabidopsis thaliana*, corn, sorghum, oat, wheat, rice, canola, or soybean; an algae such as *Chlamydomonas reinhardtii*; a nematode such as *Caenorhabditis elegans*; an insect such as *Drosophila melanogaster*, mosquito,

fruit fly, or honey bee; an arachnid such as a spider; a fish such as zebrafish; a reptile; an amphibian such as a frog or *Xenopus laevis*; a *Dictyostelium discoideum*; a fungi such as *Pneumocystis carinii*, *Takifugu rubripes*, yeast, *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*; or a *Plasmodium falciparum*.

(vi) Genome

[0127] A “genome” generally refers to genomic information from a subject, which can be, for example, at least a portion of, or the entirety of, the subject's gene-encoded hereditary information. A genome can include coding regions (e.g., that code for proteins) as well as non-coding regions. A genome can include the sequences of some or all of the subject's chromosomes. For example, the human genome ordinarily has a total of 46 chromosomes. The sequences of some or all of these can constitute the genome.

(vii) Adaptor, Adapter, and Tag

[0128] An “adaptor,” an “adapter,” and a “tag” are terms that are used interchangeably in this disclosure, and refer to species that can be coupled to a polynucleotide sequence (in a process referred to as “tagging”) using any one of many different techniques including (but not limited to) ligation, hybridization, and tagmentation. Adaptors can also be nucleic acid sequences that add a function, e.g., spacer sequences, primer sequences/sites, barcode sequences, unique molecular identifier sequences.

(viii) Hybridizing, Hybridize, Annealing, and Anneal

[0129] The terms “hybridizing,” “hybridize,” “annealing,” and “anneal” are used interchangeably in this disclosure, and refer to the pairing of substantially complementary or complementary nucleic acid sequences within two different molecules. Pairing can be achieved by any process in which a nucleic acid sequence joins with a substantially or fully complementary sequence through base pairing to form a hybridization complex. For purposes of hybridization, two nucleic acid sequences are “substantially complementary” if at least 60% (e.g., at least 70%, at least 80%, or at least 90%) of their individual bases are complementary to one another.

(ix) Primer

[0130] A “primer” is a single-stranded nucleic acid sequence having a 3' end that can be used as a chemical substrate for a nucleic acid polymerase in a nucleic acid extension reaction. RNA primers are formed of RNA nucleotides, and are used in RNA synthesis, while DNA primers are formed of DNA nucleotides and used in DNA synthesis. Primers can also include both RNA nucleotides and DNA nucleotides (e.g., in a random or designed pattern). Primers can also include other natural or synthetic nucleotides described herein that can have additional functionality. In some examples, DNA primers can be used to prime RNA synthesis and vice versa (e.g., RNA primers can be used to prime DNA synthesis). Primers can vary in length. For example, primers can be about 6 bases to about 120 bases. For example, primers can include up to about 25 bases.

(x) Primer Extension

[0131] A “primer extension” refers to any method where two nucleic acid sequences (e.g., a constant region from each of two distinct capture probes) become linked (e.g., hybridized) by an overlap of their respective terminal complementary nucleic acid sequences (i.e., for example, 3' termini). Such linking can be followed by nucleic acid extension (e.g., an enzymatic extension) of one, or both termini using the other nucleic acid sequence as a template for extension. Enzymatic extension can be performed by an enzyme including, but not limited to, a polymerase and/or a reverse transcriptase.

(xi) Proximity Ligation

[0132] A “proximity ligation” is a method of ligating two (or more) nucleic acid sequences that are in proximity with each other through enzymatic means (e.g., a ligase). In some embodiments, proximity ligation can include a “gap-filling” step that involves incorporation of one or more nucleic acids by a polymerase, based on the nucleic acid sequence of a template nucleic acid molecule, spanning a distance between the two nucleic acid molecules of interest (see, e.g., U.S.

Pat. No. 7,264,929, the entire contents of which are incorporated herein by reference).

[0133] A wide variety of different methods can be used for proximity ligating nucleic acid molecules, including (but not limited to) “sticky-end” and “blunt-end” ligations. Additionally, single-stranded ligation can be used to perform proximity ligation on a single-stranded nucleic acid molecule. Sticky-end proximity ligations involve the hybridization of complementary single-stranded sequences between the two nucleic acid molecules to be joined, prior to the ligation event itself. Blunt-end proximity ligations generally do not include hybridization of complementary regions from each nucleic acid molecule because both nucleic acid molecules lack a single-stranded overhang at the site of ligation.

(xii) Nucleic Acid Extension

[0134] A “nucleic acid extension” generally involves incorporation of one or more nucleic acids (e.g., A, G, C, T, U, nucleotide analogs, or derivatives thereof) into a molecule (such as, but not limited to, a nucleic acid sequence) in a template-dependent manner, such that consecutive nucleic acids are incorporated by an enzyme (such as a polymerase or reverse transcriptase), thereby generating a newly synthesized nucleic acid molecule. For example, a primer that hybridizes to a complementary nucleic acid sequence can be used to synthesize a new nucleic acid molecule by using the complementary nucleic acid sequence as a template for nucleic acid synthesis. Similarly, a 3' polyadenylated tail of an mRNA transcript that hybridizes to a poly(dT) sequence (e.g., capture domain) can be used as a template for single-strand synthesis of a corresponding cDNA molecule.

(xiii) PCR Amplification

[0135] A “PCR amplification” refers to the use of a polymerase chain reaction (PCR) to generate copies of genetic material, including DNA and RNA sequences. Suitable reagents and conditions for implementing PCR are described, for example, in U.S. Pat. Nos. 4,683,202, 4,683,195, 4,800,159, 4,965,188, and 5,512,462, the entire contents of each of which are incorporated herein by reference. In a typical PCR amplification, the reaction mixture includes the genetic material to be amplified, an enzyme, one or more primers that are employed in a primer extension reaction, and reagents for the reaction. The oligonucleotide primers are of sufficient length to provide for hybridization to complementary genetic material under annealing conditions. The length of the primers generally depends on the length of the amplification domains, but will typically be at least 4 bases, at least 5 bases, at least 6 bases, at least 8 bases, at least 9 bases, at least 10 base pairs (bp), at least 11 bp, at least 12 bp, at least 13 bp, at least 14 bp, at least 15 bp, at least 16 bp, at least 17 bp, at least 18 bp, at least 19 bp, at least 20 bp, at least 25 bp, at least 30 bp, at least 35 bp, and can be as long as 40 bp or longer, where the length of the primers will generally range from 18 to 50 bp. The genetic material can be contacted with a single primer or a set of two primers (forward and reverse primers), depending upon whether primer extension, linear or exponential amplification of the genetic material is desired.

[0136] In some embodiments, the PCR amplification process uses a DNA polymerase enzyme. The DNA polymerase activity can be provided by one or more distinct DNA polymerase enzymes. In certain embodiments, the DNA polymerase enzyme is from a bacterium, e.g., the DNA polymerase enzyme is a bacterial DNA polymerase enzyme. For instance, the DNA polymerase can be from a bacterium of the genus *Escherichia*, *Bacillus*, *Thermus*, or *Pyrococcus*.

[0137] Suitable examples of DNA polymerases that can be used include, but are not limited to: *E. coli* DNA polymerase I, Bsu DNA polymerase, Bst DNA polymerase, Taq DNA polymerase, VENT™ DNA polymerase, DEEPVENT™ DNA polymerase, LongAmp® Taq DNA polymerase, LongAmp® Hot Start Taq DNA polymerase, Crimson LongAmp® Taq DNA polymerase, Crimson Taq DNA polymerase, OneTaq® DNA polymerase, OneTaq® Quick-Load® DNA polymerase, Hemo KlenTaq® DNA polymerase, REDTaq® DNA polymerase, Phusion® DNA polymerase, Phusion® High-Fidelity DNA polymerase, Platinum Pfx DNA polymerase, AccuPrime Pfx DNA polymerase, Phi29 DNA polymerase, Klenow fragment, Pwo DNA polymerase, Pfu DNA polymerase, T4 DNA polymerase and T7 DNA polymerase enzymes.

[0138] The term “DNA polymerase” includes not only naturally-occurring enzymes but also all modified derivatives thereof, including derivatives of naturally-occurring DNA polymerase enzymes. For instance, in some embodiments, the DNA polymerase is modified to remove 5'-3' exonuclease activity. Sequence-modified derivatives or mutants of DNA polymerase enzymes that can be used include, but are not limited to, mutants that retain at least some of the functional, e.g., DNA polymerase activity of the wild-type sequence. Mutations can affect the activity profile of the enzymes, e.g., enhance or reduce the rate of polymerization, under different reaction conditions, e.g., temperature, template concentration, primer concentration, etc. Mutations or sequence-modifications can also affect the exonuclease activity and/or thermostability of the enzyme.

[0139] In some embodiments, PCR amplification can include reactions such as, but not limited to, a strand-displacement amplification reaction, a rolling circle amplification reaction, a ligase chain reaction, a transcription-mediated amplification reaction, an isothermal amplification reaction, and/or a loop-mediated amplification reaction.

[0140] In some embodiments, PCR amplification uses a single primer that is complementary to the 3' tag of target DNA fragments. In some embodiments, PCR amplification uses a first and a second primer, where at least a 3' end portion of the first primer is complementary to at least a portion of the 3' tag of the target nucleic acid fragments, and where at least a 3' end portion of the second primer exhibits the sequence of at least a portion of the 5' tag of the target nucleic acid fragments. In some embodiments, a 5' end portion of the first primer is non-complementary to the 3' tag of the target nucleic acid fragments, and a 5' end portion of the second primer does not exhibit the sequence of at least a portion of the 5' tag of the target nucleic acid fragments. In some embodiments, the first primer includes a first universal sequence and/or the second primer includes a second universal sequence.

[0141] In some embodiments (e.g., when the PCR amplification amplifies captured DNA), the PCR amplification products can be ligated to additional sequences using a DNA ligase enzyme. The DNA ligase activity can be provided by one or more distinct DNA ligase enzymes. In some embodiments, the DNA ligase enzyme is from a bacterium, e.g., the DNA ligase enzyme is a bacterial DNA ligase enzyme. In some embodiments, the DNA ligase enzyme is from a virus (e.g., a bacteriophage). For instance, the DNA ligase can be T4 DNA ligase. Other enzymes appropriate for the ligation step include, but are not limited to, Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9oN) DNA ligase (9oN™ DNA ligase, available from New England Biolabs, Ipswich, MA), and Ampligase® (available from Lucigen, Middleton, WI). Derivatives, e.g., sequence-modified derivatives, and/or mutants thereof, can also be used.

[0142] In some embodiments, genetic material is amplified by reverse transcription polymerase chain reaction (RT-PCR). The desired reverse transcriptase activity can be provided by one or more distinct reverse transcriptase enzymes (i.e., RNA dependent DNA polymerases), suitable examples of which include, but are not limited to: M-MLV, MuLV, AMV, HIV, ArrayScript™, MultiScribe™, ThermoScript™, and SuperScript® I, II, III, and IV enzymes. “Reverse transcriptase” includes not only naturally occurring enzymes, but all such modified derivatives thereof, including also derivatives of naturally-occurring reverse transcriptase enzymes.

[0143] In addition, reverse transcription can be performed using sequence-modified derivatives or mutants of M-MLV, MuLV, AMV, and HIV reverse transcriptase enzymes, including mutants that retain at least some of the functional, e.g., reverse transcriptase, activity of the wild-type sequence. The reverse transcriptase enzyme can be provided as part of a composition that includes other components, e.g., stabilizing components that enhance or improve the activity of the reverse transcriptase enzyme, such as RNase inhibitor(s), inhibitors of DNA-dependent DNA synthesis, e.g., actinomycin D. Many sequence-modified derivative or mutants of reverse transcriptase enzymes, e.g., M-MLV, and compositions including unmodified and modified enzymes are commercially available, e.g., ArrayScript™, MultiScribe™, ThermoScript™, and SuperScript® I, II, III, and IV enzymes.

[0144] Certain reverse transcriptase enzymes (e.g., Avian Myeloblastosis Virus (AMV) Reverse Transcriptase and Moloney Murine Leukemia Virus (M-MuLV, MMLV) Reverse Transcriptase) can synthesize a complementary DNA strand using both RNA (cDNA synthesis) and single-stranded DNA (ssDNA) as a template. Thus, in some embodiments, the reverse transcription reaction can use an enzyme (reverse transcriptase) that is capable of using both RNA and ssDNA as the template for an extension reaction, e.g., an AMV or MMLV reverse transcriptase.

[0145] In some embodiments, the quantification of RNA and/or DNA is carried out by real-time PCR (also known as quantitative PCR or qPCR), using techniques well known in the art, such as but not limited to “TAQMAN™”, or dyes such as “SYBR®”, or on capillaries (“LightCycler® Capillaries”). In some embodiments, the quantification of genetic material is determined by optical absorbance and with real-time PCR. In some embodiments, the quantification of genetic material is determined by digital PCR. In some embodiments, the genes analyzed can be compared to a reference nucleic acid extract (DNA and RNA) corresponding to the expression (mRNA) and quantity (DNA) in order to compare expression levels of the target nucleic acids.

(xiv) Antibody

[0146] An “antibody” is a polypeptide molecule that recognizes and binds to a complementary target antigen. Antibodies typically have a molecular structure shape that resembles a Y shape, or polymers thereof. Naturally-occurring antibodies, referred to as immunoglobulins, belong to one of the immunoglobulin classes IgG, IgM, IgA, IgD, and IgE. Antibodies can also be produced synthetically. For example, recombinant antibodies, which are monoclonal antibodies, can be synthesized using synthetic genes by recovering the antibody genes from source cells, amplifying into an appropriate vector, and introducing the vector into a host to cause the host to express the recombinant antibody. In general, recombinant antibodies can be cloned from any species of antibody-producing animal using suitable oligonucleotide primers and/or hybridization probes. Recombinant techniques can be used to generate antibodies and antibody fragments, including non-endogenous species.

[0147] Synthetic antibodies can be derived from non-immunoglobulin sources. For example, antibodies can be generated from nucleic acids (e.g., aptamers), and from non-immunoglobulin protein scaffolds (such as peptide aptamers) into which hypervariable loops are inserted to form antigen binding sites. Synthetic antibodies based on nucleic acids or peptide structures can be smaller than immunoglobulin-derived antibodies, leading to greater tissue penetration.

[0148] Antibodies can also include affimer proteins, which are affinity reagents that typically have a molecular weight of about 12-14 kDa. Affimer proteins generally bind to a target (e.g., a target protein) with both high affinity and specificity. Examples of such targets include, but are not limited to, ubiquitin chains, immunoglobulins, and C-reactive protein. In some embodiments, affimer proteins are derived from cysteine protease inhibitors, and include peptide loops and a variable N-terminal sequence that provides the binding site.

[0149] Antibodies can also include single domain antibodies (VHH domains and VNAR domains), scFvs, and Fab fragments.

(xv) Affinity Group

[0150] An “affinity group” is a molecule or molecular moiety which has a high affinity or preference for associating or binding with another specific or particular molecule or moiety. The association or binding with another specific or particular molecule or moiety can be via a non-covalent interaction, such as hydrogen bonding, ionic forces, and van der Waals interactions. An affinity group can, for example, be biotin, which has a high affinity or preference to associate or bind to the protein avidin or streptavidin. An affinity group, for example, can also refer to avidin or streptavidin which has an affinity to biotin. Other examples of an affinity group and specific or particular molecule or moiety to which it binds or associates with include, but are not limited to, antibodies or antibody fragments and their respective antigens, such as digoxigenin and anti-digoxigenin antibodies, lectin, and carbohydrates (e.g., a sugar, a monosaccharide, a disaccharide,

or a polysaccharide), and receptors and receptor ligands.

[0151] Any pair of affinity group and its specific or particular molecule or moiety to which it binds or associates with can have their roles reversed, for example, such that between a first molecule and a second molecule, in a first instance the first molecule is characterized as an affinity group for the second molecule, and in a second instance the second molecule is characterized as an affinity group for the first molecule.

(xvi) Label, Detectable Label, and Optical Label

[0152] The terms “detectable label,” “optical label,” and “label” are used interchangeably herein to refer to a directly or indirectly detectable moiety that is associated with (e.g., conjugated to) a molecule to be detected, e.g., a capture probe or analyte. The detectable label can be directly detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, can be indirectly detectable, e.g., by catalyzing chemical alterations of a chemical substrate compound or composition, which chemical substrate compound or composition is directly detectable. Detectable labels can be suitable for small scale detection and/or suitable for high-throughput screening. As such, suitable detectable labels include, but are not limited to, radioisotopes, fluorophores, chemiluminescent compounds, bioluminescent compounds, and dyes.

[0153] The detectable label can be qualitatively detected (e.g., optically or spectrally), or it can be quantified. Qualitative detection generally includes a detection method in which the existence or presence of the detectable label is confirmed, whereas quantifiable detection generally includes a detection method having a quantifiable (e.g., numerically reportable) value such as an intensity, duration, polarization, and/or other properties. In some embodiments, the detectable label is bound to a feature or to a capture probe associated with a feature. For example, detectably labeled features can include a fluorescent, a colorimetric, or a chemiluminescent label attached to a bead (see, for example, Rajeswari et al., *J. Microbiol Methods* 139:22-28, 2017, and Forcucci et al., *J. Biomed Opt.* 10:105010, 2015, the entire contents of each of which are incorporated herein by reference).

[0154] In some embodiments, a plurality of detectable labels can be attached to a feature, capture probe, or composition to be detected. For example, detectable labels can be incorporated during nucleic acid polymerization or amplification (e.g., Cy5®-labelled nucleotides, such as Cy5®-dCTP). Any suitable detectable label can be used. In some embodiments, the detectable label is a fluorophore. For example, the fluorophore can be from a group that includes: 7-AAD (7-Aminoactinomycin D), Acridine Orange (+DNA), Acridine Orange (+RNA), Alexa Fluor® 350, Alexa Fluor® 430, Alexa Fluor® 488, Alexa Fluor® 532, Alexa Fluor® 546, Alexa Fluor® 555, Alexa Fluor® 568, Alexa Fluor® 594, Alexa Fluor® 633, Alexa Fluor® 647, Alexa Fluor® 660, Alexa Fluor® 680, Alexa Fluor® 700, Alexa Fluor® 750, Allophycocyanin (APC), AMCA/AMCA-X, 7-Aminoactinomycin D (7-AAD), 7-Amino-4-methylcoumarin, 6-Aminoquinoline, Aniline Blue, ANS, APC-Cy7, ATTO-TAG™ CBQCA, ATTO-TAG™ FQ, Auramine O-Feulgen, BCECF (high pH), BFP (Blue Fluorescent Protein), BFP/GFP FRET, BOBO™-1/BO-PRO™-1, BOBO™-3/BO-PRO™-3, BODIPY® FL, BODIPY® TMR, BODIPY® TR-X, BODIPY® 530/550, BODIPY® 558/568, BODIPY® 564/570, BODIPY® 581/591, BODIPY® 630/650-X, BODIPY® 650-665-X, BTC, Calcein, Calcein Blue, Calcium Crimson™, Calcium Green-1™, Calcium Orange™, Calcofluor® White, 5-Carboxyfluorescein (5-FAM), 5-Carboxynaphthofluorescein, 6-Carboxyrhodamine 6G, 5-Carboxytetramethylrhodamine (5-TAMRA), Carboxy-X-rhodamine (5-ROX), Cascade Blue®, Cascade Yellow™, CCF2 (GeneBLAzer™), CFP (Cyan Fluorescent Protein), CFP/YFP FRET, Chromomycin A3, C1-NERF (low pH), CPM, 6-CR 6G, CTC Formazan, Cy2®, Cy3®, Cy3.5®, Cy5®, Cy5.5®, Cy7®, Cychrome (PE-Cy5), Dansylamine, Dansyl cadaverine, Dansylchloride, DAPI, Dapoxyl, DCFH, DHR, DiA (4-Di-16-ASP), DiD (DiIc18 (5)), DIDS, DiI (DiIc18 (3)), DiO (DiOC18 (3)), DiR (DiIc18 (7)), Di-4 ANEPPS, Di-8 ANEPPS, DM-NERF (4.5-6.5 pH), DsRed (Red Fluorescent Protein), EBFP, ECFP, EGFP, ELF®-97 alcohol, Eosin, Erythrosin, Ethidium bromide, Ethidium homodimer-1 (EthD-1), Europium (III) Chloride, 5-FAM (5-

Carboxyfluorescein), Fast Blue, Fluorescein-dT phosphoramidite, FITC, Fluo-3, Fluo-4, FluorX®, Fluoro-Gold™ (high pH), Fluoro-Gold™ (low pH), Fluoro-Jade, FM® 1-43, Fura-2 (high calcium), Fura-2/BCECF, Fura Red™ (high calcium), Fura Red™/Fluo-3, GeneBLAzer™ (CCF2), GFP Red Shifted (rsGFP), GFP Wild Type, GFP/BFP FRET, GFP/DsRed FRET, Hoechst 33342 & 33258, 7-Hydroxy-4-methylcoumarin (pH 9), 1,5 IAEDANS, Indo-1 (high calcium), Indo-1 (low calcium), Indodicarbocyanine, Indotricarbocyanine, JC-1, 6-JOE, JOJO™-1/JO-PRO™-1, LDS 751 (+DNA), LDS 751 (+RNA), LOLO™-1/LO-PRO™-1, Lucifer Yellow, LysoSensor™ Blue (pH 5), LysoSensor™ Green (pH 5), LysoSensor™ Yellow/Blue (pH 4.2), LysoTracker® Green, Lyso Tracker® Red, LysoTracker® Yellow, Mag-Fura-2, Mag-Indo-1, Magnesium Green™ Marina Blue®, 4-Methylumbelliferone, Mithramycin, MitoTracker® Green, MitoTracker® Orange, MitoTracker® Red, NBD (amine), Nile Red, Oregon Green® 488, Oregon Green® 500, Oregon Green® 514, Pacific Blue, PBF1, PE (R-phycoerythrin), PE-Cy5, PE-Cy7, PE-Texas Red, PerCP (Peridinin chlorophyll protein), PerCP-Cy5.5 (TruRed), PharRed (APC-Cy7), C-phycoerythrin, R-phycoerythrin, R-phycoerythrin (PE), PI (Propidium Iodide), PKH26, PKH67, POPO™-1/P0-PRO™-1, POPO™-3/P0-PRO™-3, Propidium Iodide (PI), PyMPO, Pyrene, Pyronin Y, Quantam Red (PE-Cy5), Quinacrine Mustard, R670 (PE-Cy5), Red 613 (PE-Texas Red), Red Fluorescent Protein (DsRed), Resorufin, RH 414, Rhod-2, Rhodamine B, Rhodamine Green™, Rhodamine Red™, Rhodamine Phalloidin, Rhodamine 110, Rhodamine 123, 5-ROX (carboxy-X-rhodamine), S65A, S65C, S65L, S65T, SBFI, SITS, SNAFL®-1 (high pH), SNAFL®-2, SNARFR-1 (high pH), SNARF®-1 (low pH), Sodium Green™, SpectrumAqua®, SpectrumGreen® #1, SpectrumGreen® #2, SpectrumOrange®, SpectrumRed®, SYTOR 11, SYTOR 13, SYTOR 17, SYTOR 45, SYTOX® Blue, SYTOX® Green, SYTOX® Orange, 5-TAMRA (5-Carboxytetramethylrhodamine), Tetramethylrhodamine (TRITC), Texas Red®/Texas Red®-X, Texas Red®-X (NHS Ester), Thiadicarbocyanine, Thiazole Orange, TOTO®-1/TO-PRO®-1, TOTOR-3/TO-PRO®-3, TO-PRO®-5, Tri-color (PE-Cy5), TRITC (Tetramethylrhodamine), TruRed (PerCP-Cy5.5), WW 781, X-Rhodamine (XRITC), Y66F, Y66H, Y66 W, YFP (Yellow Fluorescent Protein), YOYO®-1/YO-PRO®-1, YOYO®-3/YO-PRO®-3, 6-FAM (Fluorescein), 6-FAM (NHS Ester), 6-FAM (Azide), HEX, TAMRA (NHS Ester), Yakima Yellow, MAX, TET, TEX615, ATTO 488, ATTO 532, ATTO 550, ATTO 565, ATTO Rho101, ATTO 590, ATTO 633, ATTO 647N, TYE 563, TYE 665, TYE 705, 5' IRDye® 700, 5' IRDye® 800, 5' IRDye® 800CW (NHS Ester), WellRED D4 Dye, WellRED D3 Dye, WellRED D2 Dye, Lightcycler® 640 (NHS Ester), and Dy 750 (NHS Ester).

[0155] As mentioned above, in some embodiments, a detectable label is or includes a luminescent or chemiluminescent moiety. Common luminescent/chemiluminescent moieties include, but are not limited to, peroxidases such as horseradish peroxidase (HRP), soybean peroxidase (SP), alkaline phosphatase, and luciferase. These protein moieties can catalyze chemiluminescent reactions given the appropriate chemical substrates (e.g., an oxidizing reagent plus a chemiluminescent compound). A number of compound families are known to provide chemiluminescence under a variety of conditions. Non-limiting examples of chemiluminescent compound families include 2,3-dihydro-1,4-phthalazinedione luminol, 5-amino-6,7,8-trimethoxy- and the dimethylamino[ca]benz analog. These compounds can luminesce in the presence of alkaline hydrogen peroxide or calcium hypochlorite and base. Other examples of chemiluminescent compound families include, e.g., 2,4,5-triphenylimidazoles, para-dimethylamino and -methoxy substituents, oxalates such as oxalyl active esters, p-nitrophenyl, N-alkyl acridinium esters, luciferins, lucigenins, or acridinium esters.

(xvii) Template Switching Oligonucleotide

[0156] A “template switching oligonucleotide” is an oligonucleotide that hybridizes to untemplated nucleotides added by a reverse transcriptase (e.g., enzyme with terminal transferase activity) during reverse transcription. In some embodiments, a template switching oligonucleotide hybridizes to untemplated poly(C) nucleotides added by a reverse transcriptase. In some embodiments, the template switching oligonucleotide adds a common 5' sequence to full-length cDNA that is used for

cDNA amplification.

[0157] In some embodiments, the template switching oligonucleotide adds a common sequence onto the 5' end of the RNA being reverse transcribed. For example, a template switching oligonucleotide can hybridize to untemplated poly(C) nucleotides added onto the end of a cDNA molecule and provide a template for the reverse transcriptase to continue replication to the 5' end of the template switching oligonucleotide, thereby generating full-length cDNA ready for further amplification. In some embodiments, once a full-length cDNA molecule is generated, the template switching oligonucleotide can serve as a primer in a cDNA amplification reaction.

[0158] In some embodiments, a template switching oligonucleotide is added before, contemporaneously with, or after a reverse transcription, or other terminal transferase-based reaction. In some embodiments, a template switching oligonucleotide is included in the capture probe. In certain embodiments, methods of sample analysis using template switching oligonucleotides can involve the generation of nucleic acid products from analytes of the tissue sample, followed by further processing of the nucleic acid products with the template switching oligonucleotide.

[0159] Template switching oligonucleotides can include a hybridization region and a template region. The hybridization region can include any sequence capable of hybridizing to the target. In some embodiments, the hybridization region can, e.g., include a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases can include 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases, or more than 5 G bases. The template sequence can include any sequence to be incorporated into the cDNA. In other embodiments, the hybridization region can include at least one base in addition to at least one G base. In other embodiments, the hybridization can include bases that are not a G base. In some embodiments, the template region includes at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. In some embodiments, the template region and hybridization region are separated by a spacer.

[0160] In some embodiments, the template regions include a barcode sequence. The barcode sequence can act as a spatial barcode and/or as a unique molecular identifier. Template switching oligonucleotides can include deoxyribonucleic acids; ribonucleic acids; modified nucleic acids including 2-aminopurine, 2,6-diaminopurine (2-amino-dA), inverted dT, 5-methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine), locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), or any combination of the foregoing.

[0161] In some embodiments, the length of a template switching oligonucleotide can be at least about 1, 2, 10, 20, 50, 75, 100, 150, 200, or 250 nucleotides or longer. In some embodiments, the length of a template switching oligonucleotide can be at most about 2, 10, 20, 50, 100, 150, 200, or 250 nucleotides or longer.

(xviii) Splint Oligonucleotide

[0162] A "splint oligonucleotide" is an oligonucleotide that, when hybridized to other polynucleotides, acts as a "splint" to position the polynucleotides next to one another so that they can be ligated together. In some embodiments, the splint oligonucleotide is DNA or RNA. The splint oligonucleotide can include a nucleotide sequence that is partially complimentary to nucleotide sequences from two or more different oligonucleotides. In some embodiments, the splint oligonucleotide assists in ligating a "donor" oligonucleotide and an "acceptor" oligonucleotide. In general, an RNA ligase, a DNA ligase, or another other variety of ligase is used to ligate two nucleotide sequences together

[0163] In some embodiments, the splint oligonucleotide is between 10 and 50 oligonucleotides in length, e.g., between 10 and 45, 10 and 40, 10 and 35, 10 and 30, 10 and 25, or 10 and 20 oligonucleotides in length. In some embodiments, the splint oligonucleotide is between 15 and 50,

15 and 45, 15 and 40, 15 and 35, 15 and 30, 15 and 25, or 15 and 25 nucleotides in length.

(c) Analytes

[0164] The apparatus, systems, methods, and compositions described in this disclosure can be used to detect and analyze a wide variety of different analytes. For the purpose of this disclosure, an “analyte” can include any biological substance, structure, moiety, or component to be analyzed. The term “target” can similarly refer to an analyte of interest.

[0165] Analytes can be broadly classified into one of two groups: nucleic acid analytes, and non-nucleic acid analytes. Examples of non-nucleic acid analytes include, but are not limited to, lipids, carbohydrates, peptides, proteins, glycoproteins (N-linked or O-linked), lipoproteins, phosphoproteins, specific phosphorylated or acetylated variants of proteins, amidation variants of proteins, hydroxylation variants of proteins, methylation variants of proteins, ubiquitylation variants of proteins, sulfation variants of proteins, viral coat proteins, extracellular and intracellular proteins, antibodies, and antigen binding fragments. In some embodiments, the analyte can be an organelle (e.g., nuclei or mitochondria).

[0166] Cell surface features corresponding to analytes can include, but are not limited to, a receptor, an antigen, a surface protein, a transmembrane protein, a cluster of differentiation protein, a protein channel, a protein pump, a carrier protein, a phospholipid, a glycoprotein, a glycolipid, a cell-cell interaction protein complex, an antigen-presenting complex, a major histocompatibility complex, an engineered T-cell receptor, a T-cell receptor, a B-cell receptor, a chimeric antigen receptor, an extracellular matrix protein, a posttranslational modification (e.g., phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation) state of a cell surface protein, a gap junction, and an adherens junction.

[0167] Analytes can be derived from a specific type of cell and/or a specific sub-cellular region. For example, analytes can be derived from cytosol, from cell nuclei, from mitochondria, from microsomes, and more generally, from any other compartment, organelle, or portion of a cell. Permeabilizing agents that specifically target certain cell compartments and organelles can be used to selectively release analytes from cells for analysis.

[0168] Examples of nucleic acid analytes include DNA analytes such as genomic DNA, methylated DNA, specific methylated DNA sequences, fragmented DNA, mitochondrial DNA, in situ synthesized PCR products, and RNA/DNA hybrids.

[0169] Examples of nucleic acid analytes also include RNA analytes such as various types of coding and non-coding RNA. Examples of the different types of RNA analytes include messenger RNA (mRNA), ribosomal RNA (rRNA), transfer RNA (tRNA), microRNA (miRNA), and viral RNA. The RNA can be a transcript (e.g., present in a tissue section). The RNA can be small (e.g., less than 200 nucleic acid bases in length) or large (e.g., RNA greater than 200 nucleic acid bases in length). Small RNAs mainly include 5.8S ribosomal RNA (5.8S rRNA), 5S rRNA, transfer RNA (tRNA), microRNA (miRNA), small interfering RNA (siRNA), small nucleolar RNA (snoRNAs), Piwi-interacting RNA (piRNA), tRNA-derived small RNA (tsRNA), and small rDNA-derived RNA (srRNA). The RNA can be double-stranded RNA or single-stranded RNA. The RNA can be circular RNA. The RNA can be a bacterial rRNA (e.g., 16S rRNA or 23S rRNA).

[0170] Additional examples of analytes include mRNA and cell surface features (e.g., using the labelling agents described herein), mRNA and intracellular proteins (e.g., transcription factors), mRNA and cell methylation status, mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), mRNA and metabolites (e.g., using the labelling agents described herein), a barcoded labelling agent (e.g., the oligonucleotide tagged antibodies described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein). In some embodiments, a perturbation agent can be a small molecule, an antibody, a drug, an aptamer, a miRNA, a physical environmental (e.g., temperature change), or any other known perturbation agents.

[0171] Analytes can include a nucleic acid molecule with a nucleic acid sequence encoding at least a portion of a V(D)J sequence of an immune cell receptor (e.g., a TCR or BCR). In some embodiments, the nucleic acid molecule is cDNA first generated from reverse transcription of the corresponding mRNA, using a poly(T) containing primer. The generated cDNA can then be barcoded using a capture probe, featuring a barcode sequence (and optionally, a UMI sequence) that hybridizes with at least a portion of the generated cDNA. In some embodiments, a template switching oligonucleotide hybridizes to a poly(C) tail added to a 3' end of the cDNA by a reverse transcriptase enzyme. The original mRNA template and template switching oligonucleotide can then be denatured from the cDNA and the barcoded capture probe can then hybridize with the cDNA and a complement of the cDNA generated. Additional methods and compositions suitable for barcoding cDNA generated from mRNA transcripts including those encoding V(D)J regions of an immune cell receptor and/or barcoding methods and composition including a template switch oligonucleotide are described in PCT Patent Application PCT/US2017/057269, filed Oct. 18, 2017, and U.S. patent application Ser. No. 15/825,740, filed Nov. 29, 2017, both of which are incorporated herein by reference in their entireties. V(D)J analysis can also be completed with the use of one or more labelling agents that bind to particular surface features of immune cells and associated with barcode sequences. The one or more labelling agents can include an MHC or MHC multimer.

[0172] As described above, the analyte can include a nucleic acid capable of functioning as a component of a gene editing reaction, such as, for example, clustered regularly interspaced short palindromic repeats (CRISPR)-based gene editing. Accordingly, the capture probe can include a nucleic acid sequence that is complementary to the analyte (e.g., a sequence that can hybridize to the CRISPR RNA (crRNA), single guide RNA (sgRNA), or an adapter sequence engineered into a crRNA or sgRNA).

[0173] In certain embodiments, an analyte can be extracted from a live cell. Processing conditions can be adjusted to ensure that a biological sample remains live during analysis, and analytes are extracted from (or released from) live cells of the sample. Live cell-derived analytes can be obtained only once from the sample, or can be obtained at intervals from a sample that continues to remain in viable condition.

[0174] In general, the systems, apparatus, methods, and compositions can be used to analyze any number of analytes. For example, the number of analytes that are analyzed can be at least about 2, at least about 3, at least about 4, at least about 5, at least about 6, at least about 7, at least about 8, at least about 9, at least about 10, at least about 11, at least about 12, at least about 13, at least about 14, at least about 15, at least about 20, at least about 25, at least about 30, at least about 40, at least about 50, at least about 100, at least about 1,000, at least about 10,000, at least about 100,000 or more different analytes present in a region of the sample or within an individual feature of the substrate. Methods for performing multiplexed assays to analyze two or more different analytes will be discussed in a subsequent section of this disclosure.

(d) Biological Samples

(i) Types of Biological Samples

[0175] A “biological sample” is obtained from the subject for analysis using any of a variety of techniques including, but not limited to, biopsy, surgery, and laser capture microscopy (LCM), and generally includes cells and/or other biological material from the subject. In addition to the subjects described above, a biological sample can be obtained from non-mammalian organisms (e.g., a plants, an insect, an arachnid, a nematode (e.g., *Caenorhabditis elegans*), a fungi, an amphibian, or a fish (e.g., zebrafish)). A biological sample can be obtained from a prokaryote such as a bacterium, e.g., *Escherichia coli*, Staphylococci or *Mycoplasma pneumoniae*; an archaea; a virus such as Hepatitis C virus or human immunodeficiency virus; or a viroid. A biological sample can be obtained from a eukaryote, such as a patient derived organoid (PDO) or patient derived xenograft (PDX). The biological sample can include organoids, a miniaturized and simplified version of an

organ produced in vitro in three dimensions that shows realistic micro-anatomy. Organoids can be generated from one or more cells from a tissue, embryonic stem cells, and/or induced pluripotent stem cells, which can self-organize in three-dimensional culture owing to their self-renewal and differentiation capacities. In some embodiments, an organoid is a cerebral organoid, an intestinal organoid, a stomach organoid, a lingual organoid, a thyroid organoid, a thymic organoid, a testicular organoid, a hepatic organoid, a pancreatic organoid, an epithelial organoid, a lung organoid, a kidney organoid, a gastruloid, a cardiac organoid, or a retinal organoid. Subjects from which biological samples can be obtained can be healthy or asymptomatic individuals, individuals that have or are suspected of having a disease (e.g., cancer) or a pre-disposition to a disease, and/or individuals that are in need of therapy or suspected of needing therapy.

[0176] Biological samples can be derived from a homogeneous culture or population of the subjects or organisms mentioned herein or alternatively from a collection of several different organisms, for example, in a community or ecosystem.

[0177] Biological samples can include one or more diseased cells. A diseased cell can have altered metabolic properties, gene expression, protein expression, and/or morphologic features. Examples of diseases include inflammatory disorders, metabolic disorders, nervous system disorders, and cancer. Cancer cells can be derived from solid tumors, hematological malignancies, cell lines, or obtained as circulating tumor cells.

[0178] Biological samples can also include fetal cells. For example, a procedure such as amniocentesis can be performed to obtain a fetal cell sample from maternal circulation. Sequencing of fetal cells can be used to identify any of a number of genetic disorders, including, e.g., aneuploidy such as Down's syndrome, Edwards syndrome, and Patau syndrome. Further, cell surface features of fetal cells can be used to identify any of a number of disorders or diseases.

[0179] Biological samples can also include immune cells. Sequence analysis of the immune repertoire of such cells, including genomic, proteomic, and cell surface features, can provide a wealth of information to facilitate an understanding the status and function of the immune system. By way of example, determining the status (e.g., negative or positive) of minimal residue disease (MRD) in a multiple myeloma (MM) patient following autologous stem cell transplantation is considered a predictor of MRD in the MM patient (see, e.g., U.S. Patent Application Publication No. 2018/0156784, the entire contents of which are incorporated herein by reference).

[0180] Examples of immune cells in a biological sample include, but are not limited to, B cells, T cells (e.g., cytotoxic T cells, natural killer T cells, regulatory T cells, and T helper cells), natural killer cells, cytokine induced killer (CIK) cells, myeloid cells, such as granulocytes (basophil granulocytes, eosinophil granulocytes, neutrophil granulocytes/hypersegmented neutrophils), monocytes/macrophages, mast cells, thrombocytes/megakaryocytes, and dendritic cells.

[0181] The biological sample can include any number of macromolecules, for example, cellular macromolecules and organelles (e.g., mitochondria and nuclei). The biological sample can be a nucleic acid sample and/or protein sample. The biological sample can be a carbohydrate sample or a lipid sample. The biological sample can be obtained as a tissue sample, such as a tissue section, biopsy, a core biopsy, needle aspirate, or fine needle aspirate. The sample can be a fluid sample, such as a blood sample, urine sample, or saliva sample. The sample can be a skin sample, a colon sample, a cheek swab, a histology sample, a histopathology sample, a plasma or serum sample, a tumor sample, living cells, cultured cells, a clinical sample such as, for example, whole blood or blood-derived products, blood cells, or cultured tissues or cells, including cell suspensions.

[0182] Cell-free biological samples can include extracellular polynucleotides. Extracellular polynucleotides can be isolated from a bodily sample, e.g., blood, plasma, serum, urine, saliva, mucosal excretions, sputum, stool, and tears.

[0183] As discussed above, a biological sample can include a single analyte of interest, or more than one analyte of interest. Methods for performing multiplexed assays to analyze two or more different analytes in a single biological sample is discussed in a subsequent section of this

disclosure.

(ii) Preparation of Biological Samples

[0184] A variety of steps can be performed to prepare a biological sample for analysis. Except where indicated otherwise, the preparative steps described below can generally be combined in any manner to appropriately prepare a particular sample for analysis.

(1) Tissue Sectioning

[0185] A biological sample can be harvested from a subject (e.g., via surgical biopsy, whole subject sectioning), grown in vitro on a growth substrate or culture dish as a population of cells, or prepared as a tissue slice or tissue section. Grown samples may be sufficiently thin for analysis without further processing steps. Alternatively, grown samples, and samples obtained via biopsy or sectioning, can be prepared as thin tissue sections using a mechanical cutting apparatus such as a vibrating blade microtome. As another alternative, in some embodiments, a thin tissue section can be prepared by applying a touch imprint of a biological sample to a suitable substrate material.

[0186] The thickness of the tissue section can be a fraction of (e.g., less than 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, or 0.1) the maximum cross-sectional dimension of a cell. However, tissue sections having a thickness that is larger than the maximum cross-section cell dimension can also be used. For example, cryostat sections can be used, which can be, e.g., 10-20 micrometers thick.

[0187] More generally, the thickness of a tissue section typically depends on the method used to prepare the section and the physical characteristics of the tissue, and therefore sections having a wide variety of different thicknesses can be prepared and used. For example, the thickness of the tissue section can be at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, 1.0, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 20, 30, 40, or 50 micrometers. Thicker sections can also be used if desired or convenient, e.g., at least 70, 80, 90, or 100 micrometers or more. Typically, the thickness of a tissue section is between 1-100 micrometers, 1-50 micrometers, 1-30 micrometers, 1-25 micrometers, 1-20 micrometers, 1-15 micrometers, 1-10 micrometers, 2-8 micrometers, 3-7 micrometers, or 4-6 micrometers, but as mentioned above, sections with thicknesses larger or smaller than these ranges can also be analysed.

[0188] Multiple sections can also be obtained from a single biological sample. For example, multiple tissue sections can be obtained from a surgical biopsy sample by performing serial sectioning of the biopsy sample using a sectioning blade. Spatial information among the serial sections can be preserved in this manner, and the sections can be analysed successively to obtain three-dimensional information about the biological sample.

(2) Freezing

[0189] In some embodiments, the biological sample (e.g., a tissue section as described above) can be prepared by deep freezing at a temperature suitable to maintain or preserve the integrity (e.g., the physical characteristics) of the tissue structure. Such a temperature can be, e.g., less than -20°C ., or less than -25°C ., -30°C ., -40°C ., -50°C ., -60°C ., -70°C ., 80°C .- 90°C ., -100°C ., -110°C ., -120°C ., -130°C ., -140°C ., -150°C ., -160°C ., -170°C ., -180°C ., -190°C ., or -200°C . The frozen tissue sample can be sectioned, e.g., thinly sliced, onto a substrate surface using any number of suitable methods. For example, a tissue sample can be prepared using a chilled microtome (e.g., a cryostat) set at a temperature suitable to maintain both the structural integrity of the tissue sample and the chemical properties of the nucleic acids in the sample. Such a temperature can be, e.g., less than -15°C ., less than -20°C ., or less than -25°C . A sample can be snap frozen in isopentane and liquid nitrogen. Frozen samples can be stored in a sealed container prior to embedding.

(3) Formalin Fixation and Paraffin Embedding

[0190] In some embodiments, the biological sample can be prepared using formalin-fixation and paraffin-embedding (FFPE), which are established methods. In some embodiments, cell suspensions and other non-tissue samples can be prepared using formalin-fixation and paraffin-embedding. Following fixation of the sample and embedding in a paraffin or resin block, the

sample can be sectioned as described above. Prior to analysis, the paraffin-embedding material can be removed from the tissue section (e.g., deparaffinization) by incubating the tissue section in an appropriate solvent (e.g., xylene) followed by a rinse (e.g., 99.5% ethanol for 2 minutes, 96% ethanol for 2 minutes, and 70% ethanol for 2 minutes).

(4) Fixation

[0191] As an alternative to formalin fixation described above, a biological sample can be fixed in any of a variety of other fixatives to preserve the biological structure of the sample prior to analysis. For example, a sample can be fixed via immersion in ethanol, methanol, acetone, formaldehyde (e.g., 2% formaldehyde), paraformaldehyde-Triton, glutaraldehyde, or combinations thereof.

[0192] In some embodiments, acetone fixation is used with fresh frozen samples, which can include, but are not limited to, cortex tissue, mouse olfactory bulb, human brain tumor, human post-mortem brain, and breast cancer samples. In some embodiments, a compatible fixation method is chosen and/or optimized based on a desired workflow. For example, formaldehyde fixation may be chosen as compatible for workflows using IHC/IF protocols for protein visualization. As another example, methanol fixation may be chosen for workflows emphasizing RNA/DNA library quality. Acetone fixation may be chosen in some applications to permeabilize the tissue. When acetone fixation is performed, pre-permeabilization steps (described below) may not be performed. Alternatively, acetone fixation can be performed in conjunction with permeabilization steps.

(5) Embedding

[0193] As an alternative to paraffin embedding described above, a biological sample can be embedded in any of a variety of other embedding materials to provide a substrate to the sample prior to sectioning and other handling steps. In general, the embedding material is removed prior to analysis of tissue sections obtained from the sample. Suitable embedding materials include, but are not limited to, waxes, resins (e.g., methacrylate resins), epoxies, and agar.

(6) Staining

[0194] To facilitate visualization, biological samples can be stained using a wide variety of stains and staining techniques. In some embodiments, a sample can be stained using any number of biological stains, including but not limited to, acridine orange, Bismarck brown, carmine, coomassie blue, cresyl violet, DAPI, eosin, ethidium bromide, acid fuchsine, hematoxylin, Hoechst stains, iodine, methyl green, methylene blue, neutral red, Nile blue, Nile red, osmium tetroxide, propidium iodide, rhodamine, or safranin.

[0195] The sample can be stained using known staining techniques, including Can-Grunwald, Giemsa, hematoxylin and eosin (H&E), Jenner's, Leishman, Masson's trichrome, Papanicolaou, Romanowsky, silver, Sudan, Wright's, and/or Periodic Acid Schiff (PAS) staining techniques. PAS staining is typically performed after formalin or acetone fixation.

[0196] In some embodiments, the biological sample can be stained using a detectable label (e.g., radioisotopes, fluorophores, chemiluminescent compounds, bioluminescent compounds, and dyes) as described elsewhere herein. In some embodiments, a biological sample is stained using only one type of stain or one technique. In some embodiments, staining includes biological staining techniques such as H&E staining. In some embodiments, staining includes identifying analytes using fluorescently-conjugated antibodies. In some embodiments, a biological sample is stained using two or more different types of stains, or two or more different staining techniques. For example, a biological sample can be prepared by staining and imaging using one technique (e.g., H&E staining and brightfield imaging), followed by staining and imaging using another technique (e.g., IHC/IF staining and fluorescence microscopy) for the same biological sample.

[0197] In some embodiments, biological samples can be destained. Methods of destaining or discoloring a biological sample are known in the art, and generally depend on the nature of the stain(s) applied to the sample. For example, H&E staining can be destained by washing the sample in HCl, or any other low pH acid (e.g., selenic acid, sulfuric acid, hydroiodic acid, benzoic acid,

carbonic acid, malic acid, phosphoric acid, oxalic acid, succinic acid, salicylic acid, tartaric acid, sulfurous acid, trichloroacetic acid, hydrobromic acid, hydrochloric acid, nitric acid, orthophosphoric acid, arsenic acid, selenous acid, chromic acid, citric acid, hydrofluoric acid, nitrous acid, isocyanic acid, formic acid, hydrogen selenide, molybdic acid, lactic acid, acetic acid, carbonic acid, hydrogen sulfide, or combinations thereof). In some embodiments, destaining can include 1, 2, 3, 4, 5, or more washes in a low pH acid (e.g., HCl). In some embodiments, destaining can include adding HCl to a downstream solution (e.g., permeabilization solution). In some embodiments, destaining can include dissolving an enzyme used in the disclosed methods (e.g., pepsin) in a low pH acid (e.g., HCl) solution. In some embodiments, after destaining hematoxylin with a low pH acid, other reagents can be added to the destaining solution to raise the pH for use in other applications. For example, SDS can be added to a low pH acid destaining solution in order to raise the pH as compared to the low pH acid destaining solution alone. As another example, in some embodiments, one or more immunofluorescence stains are applied to the sample via antibody coupling. Such stains can be removed using techniques such as cleavage of disulfide linkages via treatment with a reducing agent and detergent washing, chaotropic salt treatment, treatment with antigen retrieval solution, and treatment with an acidic glycine buffer. Methods for multiplexed staining and destaining are described, for example, in Bolognesi et al., *J. Histochem. Cytochem.* 2017; 65 (8): 431-444, Lin et al., *Nat Commun.* 2015; 6:8390, Pirici et al., *J. Histochem. Cytochem.* 2009; 57:567-75, and Glass et al., *J. Histochem. Cytochem.* 2009; 57:899-905, the entire contents of each of which are incorporated herein by reference.

(7) Hydrogel Embedding

[0198] In some embodiments, hydrogel formation occurs within a biological sample. In some embodiments, a biological sample (e.g., tissue section) is embedded in a hydrogel. In some embodiments, hydrogel subunits are infused into the biological sample, and polymerization of the hydrogel is initiated by an external or internal stimulus. A “hydrogel” as described herein can include a cross-linked 3D network of hydrophilic polymer chains. A “hydrogel subunit” can be a hydrophilic monomer, a molecular precursor, or a polymer that can be polymerized (e.g., cross-linked) to form a three-dimensional (3D) hydrogel network.

[0199] A hydrogel can swell in the presence of water. In some embodiments, a hydrogel comprises a natural material. In some embodiments, a hydrogel includes a synthetic material. In some embodiments, a hydrogel includes a hybrid material, e.g., the hydrogel material comprises elements of both synthetic and natural polymers. Any of the materials used in hydrogels or hydrogels comprising a polypeptide-based material described herein can be used. Embedding the sample in this manner typically involves contacting the biological sample with a hydrogel such that the biological sample becomes surrounded by the hydrogel. For example, the sample can be embedded by contacting the sample with a suitable polymer material, and activating the polymer material to form a hydrogel. In some embodiments, the hydrogel is formed such that the hydrogel is internalized within the biological sample.

[0200] In some embodiments, the biological sample is immobilized in the hydrogel via cross-linking of the polymer material that forms the hydrogel. Cross-linking can be performed chemically and/or photochemically, or alternatively by any other hydrogel-formation method known in the art. For example, the biological sample can be immobilized in the hydrogel by polyacrylamide crosslinking. Further, analytes of a biological sample can be immobilized in a hydrogel by crosslinking (e.g., polyacrylamide crosslinking).

[0201] The composition and application of the hydrogel to a biological sample typically depends on the nature and preparation of the biological sample (e.g., sectioned, non-sectioned, fresh-frozen tissue, type of fixation). A hydrogel can be any appropriate hydrogel where upon formation of the hydrogel on the biological sample the biological sample becomes anchored to or embedded in the hydrogel. Non-limiting examples of hydrogels are described herein or are known in the art. As one example, where the biological sample is a tissue section, the hydrogel can include a monomer

solution and an ammonium persulfate (APS) initiator/tetramethylethylenediamine (TEMED) accelerator solution. As another example, where the biological sample consists of cells (e.g., cultured cells or cells disassociated from a tissue sample), the cells can be incubated with the monomer solution and APS/TEMED solutions. For cells, hydrogel are formed in compartments, including but not limited to devices used to culture, maintain, or transport the cells. For example, hydrogels can be formed with monomer solution plus APS/TEMED added to the compartment to a depth ranging from about 0.1 μm to about 5 mm.

[0202] In some embodiments, a hydrogel includes a linker that allows anchoring of the biological sample to the hydrogel. In some embodiments, a hydrogel includes linkers that allow anchoring of biological analytes to the hydrogel. In such cases, the linker can be added to the hydrogel before, contemporaneously with, or after hydrogel formation. Non-limiting examples of linkers that anchor nucleic acids to the hydrogel can include 6-((Acryloyl)amino) hexanoic acid (Acryloyl-X SE) (available from ThermoFisher, Waltham, MA), Label-IT Amine (available from MirusBio, Madison, WI) and Label X (Chen et al., Nat. Methods 13:679-684, (2016)).

[0203] In some embodiments, functionalization chemistry can be used. In some embodiments, functionalization chemistry includes hydrogel-tissue chemistry (HTC). Any hydrogel-tissue backbone (e.g., synthetic or native) suitable for HTC can be used for anchoring biological macromolecules and modulating functionalization. Non-limiting examples of methods using HTC backbone variants include CLARITY, PACT, ExM, SWITCH and ePACT. In some embodiments, hydrogel formation within a biological sample is permanent. For example, biological macromolecules can permanently adhere to the hydrogel allowing multiple rounds of interrogation. In some embodiments, hydrogel formation within a biological sample is reversible.

[0204] In some embodiments, additional reagents are added to the hydrogel subunits before, contemporaneously with, and/or after polymerization. For example, additional reagents can include but are not limited to oligonucleotides (e.g., capture probes), endonucleases to fragment DNA, fragmentation buffer for DNA, DNA polymerase enzymes, dNTPs used to amplify the nucleic acid and to attach the barcode to the amplified fragments. Other enzymes can be used, including without limitation, RNA polymerase, transposase, ligase, proteinase K, and DNase. Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers, and switch oligonucleotides. In some embodiments, optical labels are added to the hydrogel subunits before, contemporaneously with, and/or after polymerization.

[0205] In some embodiments, HTC reagents are added to the hydrogel before, contemporaneously with, and/or after polymerization. In some embodiments, a cell tagging agent is added to the hydrogel before, contemporaneously with, and/or after polymerization. In some embodiments, a cell-penetrating agent is added to the hydrogel before, contemporaneously with, and/or after polymerization.

[0206] In some embodiments, a biological sample is embedded in a hydrogel to facilitate sample transfer to another location (e.g., to an array). For example, archived biological samples (e.g., FFPE tissue sections) can be transferred from storage to a spatial array to perform spatial analysis. In some embodiments, a biological sample on a substrate can be covered with any of the prepolymer solutions described herein. In some embodiments, the prepolymer solution can be polymerized such that a hydrogel is formed on top of and/or around the biological sample. Hydrogel formation can occur in a manner sufficient to anchor (e.g., embed) the biological sample to the hydrogel. After hydrogel formation, the biological sample is anchored to (e.g., embedded in) the hydrogel wherein separating the hydrogel from the substrate (e.g., glass slide) results in the biological sample separating from the substrate along with the hydrogel. The biological sample contained in the hydrogel can then be contacted with a spatial array, and spatial analysis can be performed on the biological sample.

[0207] Any variety of characteristics can determine the transfer conditions required for a given biological sample. Non-limiting examples of characteristics likely to impact transfer conditions

include the sample (e.g., thickness, fixation, and cross-linking) and/or the analyte of interest (different conditions to preserve and/or transfer different analytes (e.g., DNA, RNA, and protein)). [0208] In some embodiments, the hydrogel is removed after contacting the biological sample with the spatial array. For example, methods described herein can include an event-dependent (e.g., light or chemical) depolymerizing hydrogel, wherein upon application of the event (e.g., external stimuli) the hydrogel depolymerizes. In one example, a biological sample can be anchored to a DTT-sensitive hydrogel, where addition of DTT can cause the hydrogel to depolymerize and release the anchored biological sample.

[0209] Hydrogels embedded within biological samples can be cleared using any suitable method. For example, electrophoretic tissue clearing methods can be used to remove biological macromolecules from the hydrogel-embedded sample. In some embodiments, a hydrogel-embedded sample is stored in a medium before or after clearing of hydrogel (e.g., a mounting medium, methylcellulose, or other semi-solid mediums).

[0210] In some embodiments, the hydrogel chemistry can be tuned to specifically bind (e.g., retain) particular species of analytes (e.g., RNA, DNA, protein, etc.). In some embodiments, a hydrogel includes a linker that allows anchoring of the biological sample to the hydrogel. In some embodiments, a hydrogel includes linkers that allow anchoring of biological analytes to the hydrogel. In such cases, the linker can be added to the hydrogel before, contemporaneously with, or after hydrogel formation. Non-limiting examples of linkers that anchor nucleic acids to the hydrogel can include 6-((Acryloyl)amino) hexanoic acid (Acryloyl-X SE), Label-IT Amine and Label X (Chen et al., Nat. Methods 13:679-684, (2016)). Non-limiting examples of characteristics likely to impact transfer conditions include the sample (e.g., thickness, fixation, and cross-linking) and/or the analyte of interest (different conditions to preserve and/or transfer different analytes (e.g., DNA, RNA, and protein)).

[0211] Additional methods and aspects of hydrogel embedding of biological samples are described for example in Chen et al., Science 347 (6221): 543-548, 2015, the entire contents of which are incorporated herein by reference.

(8) Biological Sample Transfer

[0212] In some embodiments, a biological sample immobilized on a substrate (e.g., a biological sample prepared using methanol fixation or formalin-fixation and paraffin-embedding (FFPE)) is transferred to a spatial array using a hydrogel. In some embodiments, a hydrogel is formed on top of a biological sample on a substrate (e.g., glass slide). For example, hydrogel formation can occur in a manner sufficient to anchor (e.g., embed) the biological sample to the hydrogel. After hydrogel formation, the biological sample is anchored to (e.g., embedded in) the hydrogel wherein separating the hydrogel from the substrate results in the biological sample separating from the substrate along with the hydrogel. The biological sample can then be contacted with a spatial array, thereby allowing spatial profiling of the biological sample. In some embodiments, the hydrogel is removed after contacting the biological sample with the spatial array. For example, methods described herein can include an event-dependent (e.g., light or chemical) depolymerizing hydrogel, wherein upon application of the event (e.g., external stimuli) the hydrogel depolymerizes. In one example, a biological sample can be anchored to a DTT-sensitive hydrogel, where addition of DTT can cause the hydrogel to depolymerize and release the anchored biological sample. A hydrogel can be any appropriate hydrogel where upon formation of the hydrogel on the biological sample the biological sample becomes anchored to or embedded in the hydrogel. Non-limiting examples of hydrogels are described herein or are known in the art. In some embodiments, a hydrogel includes a linker that allows anchoring of the biological sample to the hydrogel. In some embodiments, a hydrogel includes linkers that allow anchoring of biological analytes to the hydrogel. In such cases, the linker can be added to the hydrogel before, contemporaneously with, or after hydrogel formation. Non-limiting examples of linkers that anchor nucleic acids to the hydrogel can include 6-((Acryloyl)amino) hexanoic acid (Acryloyl-X SE) (available from ThermoFisher, Waltham,

MA), Label-IT Amine (available from MirusBio, Madison, WI) and Label X (Chen et al., *Nat. Methods* 13:679-684, 2016). Any variety of characteristics can determine the transfer conditions required for a given biological sample. Non-limiting examples of characteristics likely to impact transfer conditions include the sample (e.g., thickness, fixation, and cross-linking) and/or the analyte of interest (different conditions to preserve and/or transfer different analytes (e.g., DNA, RNA, and protein)). In some embodiments, hydrogel formation can occur in a manner sufficient to anchor the analytes (e.g., embed) in the biological sample to the hydrogel. In some embodiments, the hydrogel can be imploded (e.g., shrunk) with the anchored analytes (e.g., embedded in the hydrogel) present in the biological sample. In some embodiments, the hydrogel can be expanded (e.g., isometric expansion) with the anchored analytes (e.g., embedded in the hydrogel) present in the biological sample. In some embodiments, the hydrogel can be imploded (e.g., shrunk) and subsequently expanded with anchored analytes (e.g., embedded in the hydrogel) present in the biological sample.

(9) Isometric Expansion

[0213] In some embodiments, a biological sample embedded in a hydrogel can be isometrically expanded. Isometric expansion methods that can be used include hydration, a preparative step in expansion microscopy, as described in Chen et al., *Science* 347 (6221): 543-548, 2015; Asano et al. *Current Protocols*. 2018, 80:1, doi: 10.1002/cpcb.56 and Gao et al. *BMC Biology*. 2017, 15:50, doi: 10.1186/s12915-017-0393-3, Wassie, A. T., et al, Expansion microscopy: principles and uses in biological research, *Nature Methods*, 16 (1): 33-41 (2018), each of which is incorporated by reference in its entirety.

[0214] In general, the steps used to perform isometric expansion of the biological sample can depend on the characteristics of the sample (e.g., thickness of tissue section, fixation, cross-linking), and/or the analyte of interest (e.g., different conditions to anchor RNA, DNA, and protein to a gel).

[0215] Isometric expansion can be performed by anchoring one or more components of a biological sample to a gel, followed by gel formation, proteolysis, and swelling. Isometric expansion of the biological sample can occur prior to immobilization of the biological sample on a substrate, or after the biological sample is immobilized to a substrate. In some embodiments, the isometrically expanded biological sample can be removed from the substrate prior to contacting the expanded biological sample with a spatially barcoded array (e.g., spatially barcoded capture probes on a substrate).

[0216] In some embodiments, proteins in the biological sample are anchored to a swellable gel such as a polyelectrolyte gel. An antibody can be directed to the protein before, after, or in conjunction with being anchored to the swellable gel. DNA and/or RNA in a biological sample can also be anchored to the swellable gel via a suitable linker. Examples of such linkers include, but are not limited to, 6-((Acryloyl)amino) hexanoic acid (Acryloyl-X SE) (available from ThermoFisher, Waltham, MA), Label-IT Amine (available from MirusBio, Madison, WI) and Label X (described for example in Chen et al., *Nat. Methods* 13:679-684, 2016, the entire contents of which are incorporated herein by reference).

[0217] Isometric expansion of the sample can increase the spatial resolution of the subsequent analysis of the sample. For example, isometric expansion of the biological sample can result in increased resolution in spatial profiling (e.g., single-cell profiling). The increased resolution in spatial profiling can be determined by comparison of an isometrically expanded sample with a sample that has not been isometrically expanded.

[0218] Isometric expansion can enable three-dimensional spatial resolution of the subsequent analysis of the sample. In some embodiments, isometric expansion of the biological sample can occur in the presence of spatial profiling reagents (e.g., analyte capture agents or capture probes). For example, the swellable gel can include analyte capture agents or capture probes anchored to the swellable gel via a suitable linker. In some embodiments, spatial profiling reagents can be delivered

to particular locations in an isometrically expanded biological sample.

[0219] In some embodiments, a biological sample is isometrically expanded to a volume at least $2\times$, $2.1\times$, $2.2\times$, $2.3\times$, $2.4\times$, $2.5\times$, $2.6\times$, $2.7\times$, $2.8\times$, $2.9\times$, $3\times$, $3.1\times$, $3.2\times$, $3.3\times$, $3.4\times$, $3.5\times$, $3.6\times$, $3.7\times$, $3.8\times$, $3.9\times$, $4\times$, $4.1\times$, $4.2\times$, $4.3\times$, $4.4\times$, $4.5\times$, $4.6\times$, $4.7\times$, $4.8\times$, or $4.9\times$ its non-expanded volume. In some embodiments, the sample is isometrically expanded to at least $2\times$ and less than $20\times$ of its non-expanded volume.

[0220] In some embodiments, a biological sample embedded in a hydrogel is isometrically expanded to a volume at least $2\times$, $2.1\times$, $2.2\times$, $2.3\times$, $2.4\times$, $2.5\times$, $2.6\times$, $2.7\times$, $2.8\times$, $2.9\times$, $3\times$, $3.1\times$, $3.2\times$, $3.3\times$, $3.4\times$, $3.5\times$, $3.6\times$, $3.7\times$, $3.8\times$, $3.9\times$, $4\times$, $4.1\times$, $4.2\times$, $4.3\times$, $4.4\times$, $4.5\times$, $4.6\times$, $4.7\times$, $4.8\times$, or $4.9\times$ its non-expanded volume. In some embodiments, the biological sample embedded in a hydrogel is isometrically expanded to at least $2\times$ and less than $20\times$ of its non-expanded volume.

(10) Substrate Attachment

[0221] In some embodiments, the biological sample can be attached to a substrate. Examples of substrates suitable for this purpose are described in detail below. Attachment of the biological sample can be irreversible or reversible, depending upon the nature of the sample and subsequent steps in the analytical method.

[0222] In certain embodiments, the sample can be attached to the substrate reversibly by applying a suitable polymer coating to the substrate, and contacting the sample to the polymer coating. The sample can then be detached from the substrate using an organic solvent that at least partially dissolves the polymer coating. Hydrogels are examples of polymers that are suitable for this purpose.

[0223] More generally, in some embodiments, the substrate can be coated or functionalized with one or more substances to facilitate attachment of the sample to the substrate. Suitable substances that can be used to coat or functionalize the substrate include, but are not limited to, lectins, polylysine, antibodies, and polysaccharides.

(11) Unaggregated Cells

[0224] In some embodiments, the biological sample corresponds to cells (e.g., derived from a cell culture or a tissue sample). In a cell sample with a plurality of cells, individual cells can be naturally unaggregated. For example, cells can be derived from a suspension of cells and/or disassociated or disaggregated cells from a tissue or tissue section.

[0225] Alternatively, the cells in the sample may be aggregated, and may be disaggregated into individual cells using, for example, enzymatic or mechanical techniques. Examples of enzymes used in enzymatic disaggregation include, but are not limited to, dispase, collagenase, trypsin, or combinations thereof. Mechanical disaggregation can be performed, for example, using a tissue homogenizer.

[0226] In some embodiments of unaggregated cells or disaggregated cells, the cells are distributed onto the substrate such that at least one cell occupies a distinct spatial feature on the substrate. The cells can be immobilized on the substrate (e.g., to prevent lateral diffusion of the cells). In some embodiments, a cell immobilization agent can be used to immobilize a non-aggregated or disaggregated sample on a spatially-barcoded array prior to analyte capture. A “cell immobilization agent” can refer to an antibody, attached to a substrate, which can bind to a cell surface marker. In some embodiments, the distribution of the plurality of cells on the substrate follows Poisson statistics.

[0227] In some embodiments, cells from a plurality of cells are immobilized on a substrate. In some embodiments, the cells are immobilized to prevent lateral diffusion, for example, by adding a hydrogel and/or by the application of an electric field.

(12) Suspended and Adherent Cells

[0228] In some embodiments, the biological sample can be derived from a cell culture grown in vitro. Samples derived from a cell culture can include one or more suspension cells which are anchorage-independent within the cell culture. Examples of such cells include, but are not limited

to, cell lines derived from hematopoietic cells, and from the following cell lines: Colo205, CCRF-CEM, HL-60, K562, MOLT-4, RPMI-8226, SR, HOP-92, NCI-H322M, and MALME-3M.

[0229] Samples derived from a cell culture can include one or more adherent cells which grow on the surface of the vessel that contains the culture medium. Non-limiting examples of adherent cells include DU145 (prostate cancer) cells, H295R (adrenocortical cancer) cells, HeLa (cervical cancer) cells, KBM-7 (chronic myelogenous leukemia) cells, LNCaP (prostate cancer) cells, MCF-7 (breast cancer) cells, MDA-MB-468 (breast cancer) cells, PC3 (prostate cancer) cells, SaOS-2 (bone cancer) cells, SH-SY5Y (neuroblastoma, cloned from a myeloma) cells, T-47D (breast cancer) cells, THP-1 (acute myeloid leukemia) cells, U87 (glioblastoma) cells, National Cancer Institute's 60 cancer cell line panel (NCI60), vero (African green monkey *Chlorocebus* kidney epithelial cell line) cells, MC3T3 (embryonic calvarium) cells, GH3 (pituitary tumor) cells, PC12 (pheochromocytoma) cells, dog MDCK kidney epithelial cells, *Xenopus* A6 kidney epithelial cells, zebrafish AB9 cells, and Sf9 insect epithelial cells.

[0230] Additional examples of adherent cells are shown in Table 1 and catalogued, for example, in “A Catalog of in Vitro Cell Lines, Transplantable Animal and Human Tumors and Yeast,” The Division of Cancer Treatment and Diagnosis (DCTD), National Cancer Institute (2013), and in Abaan et al., “The exomes of the NCI-60 panel: a genomic resource for cancer biology and systems pharmacology,” *Cancer Research* 73 (14): 4372-82, 2013, the entire contents of each of which are incorporated by reference herein.

TABLE-US-00001

TABLE 1	Examples of adherent cells	Organ	of Cell Line	Species	Origin
Disease	BT549	Human	Breast Ductal Carcinoma	HS 578T	Human Breast Carcinoma
	MCF7	Human Breast Adenocarcinoma	MDA-MB-231	Human Breast Adenocarcinoma	MDA-MB-468
	T-47D	Human Breast Ductal Carcinoma	SF268	Human CNS Anaplastic Astrocytoma	SF295
	SF295	Human CNS Glioblastoma-Multiforme	SF539	Human CNS Glioblastoma	SNB-19
	SNB-19	Human CNS Glioblastoma	SNB-75	Human CNS Astrocytoma	U251
	U251	Human CNS Glioblastoma	Colo205	Human Colon Dukes' type D, Colorectal adenocarcinoma	HCC 2998
	HCC 2998	Human Colon Carcinoma	HCT-116	Human Colon Carcinoma	HCT-15
	HCT-15	Human Colon Dukes' type C, Colorectal adenocarcinoma	HT29	Human Colon Colorectal adenocarcinoma	KM12
	KM12	Human Colon Adenocarcinoma, Grade III	SW620	Human Colon Adenocarcinoma	786-O
	786-O	Human Kidney renal cell adenocarcinoma	A498	Human Kidney Adenocarcinoma	ACHN
	ACHN	Human Kidney renal cell adenocarcinoma	CAKI	Human Kidney clear cell carcinoma	RXF 393
	RXF 393	Human Kidney Poorly Differentiated Hypernephroma	SN12C	Human Kidney Carcinoma	TK-10
	TK-10	Human Kidney Spindle Cell carcinoma	UO-31	Human Kidney Carcinoma	A549
	A549	Human Lung Adenocarcinoma	EKVX	Human Lung Adenocarcinoma	HOP-62
	HOP-62	Human Lung Adenocarcinoma	HOP-92	Human Lung Large Cell, Undifferentiated	NCI-H226
	NCI-H226	Human Lung squamous cell carcinoma; mesothelioma	NCI-H23	Human Lung adenocarcinoma; non-small cell lung cancer	NCI-H460
	NCI-H460	Human Lung carcinoma; large cell lung cancer	NCI-H522	Human Lung adenocarcinoma; non-small cell lung cancer	LOX IMVI
	LOX IMVI	Human Melanoma Malignant Amelanotic melanoma	M14	Human Melanoma malignant melanoma	MALME-3M
	MALME-3M	Human Melanoma malignant melanoma	MDA-MB-435	Human Melanoma Adenocarcinoma	SK-MEL-2
	SK-MEL-2	Human Melanoma malignant melanoma	SK-MEL-28	Human Melanoma malignant melanoma	SK-MEL-5
	SK-MEL-5	Human Melanoma malignant melanoma	UACC-257	Human Melanoma malignant melanoma	UACC-62
	UACC-62	Human Melanoma malignant melanoma	IGROV1	Human Ovary Cystadenocarcinoma	OVCAR-3
	OVCAR-3	Human Ovary Adenocarcinoma	OVCAR-4	Human Ovary Adenocarcinoma	OVCAR-5
	OVCAR-5	Human Ovary Adenocarcinoma	OVCAR-8	Human Ovary Adenocarcinoma	SK-OV-3
	SK-OV-3	Human Ovary Adenocarcinoma	NCI-ADR-RES	Human Ovary Adenocarcinoma	DU145
	DU145	Human Prostate Carcinoma	PC-3	Human Prostate grade IV, adenocarcinoma	

[0231] In some embodiments, the adherent cells are cells that correspond to one or more of the following cell lines: BT549, HS 578T, MCF7, MDA-MB-231, MDA-MB-468, T-47D, SF268, SF295, SF539, SNB-19, SNB-75, U251, Colo205, HCC 2998, HCT-116, HCT-15, HT29, KM12,

SW620, 786-O, A498, CAKI, RXF 393, SN12C, TK-10, UO-31, A549, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H460, NCI-H522, LOX IMVI, M14, MALME-3M, MDA-MB-435, SK-, EL-2, SK-MEL-28, SK-MEL-5, UACC-257, UACC-62, IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3, NCI-ADR-RES, DU145, PC-3, DU145, H295R, HeLa, KBM-7, LNCaP, MCF-7, MDA-MB-468, PC3, SaOS-2, SH-SY5Y, T-47D, THP-1, U87, vero, MC3T3, GH3, PC12, dog MDCK kidney epithelial, *Xenopus* A6 kidney epithelial, zebrafish AB9, and Sf9 insect epithelial cell lines.

(13) Tissue Permeabilization

[0232] In some embodiments, a biological sample can be permeabilized to facilitate transfer of analytes out of the sample, and/or to facilitate transfer of species (such as capture probes) into the sample. If a sample is not permeabilized sufficiently, the amount of analyte captured from the sample may be too low to enable adequate analysis. Conversely, if the tissue sample is too permeable, the relative spatial relationship of the analytes within the tissue sample can be lost.

Hence, a balance between permeabilizing the tissue sample enough to obtain good signal intensity while still maintaining the spatial resolution of the analyte distribution in the sample is desirable.

[0233] In general, a biological sample can be permeabilized by exposing the sample to one or more permeabilizing agents. Suitable agents for this purpose include, but are not limited to, organic solvents (e.g., acetone, ethanol, and methanol), cross-linking agents (e.g., paraformaldehyde), detergents (e.g., saponin, Triton X-100™, Tween-20™, or sodium dodecyl sulfate (SDS)), and enzymes (e.g., trypsin, proteases (e.g., proteinase K). In some embodiments, the detergent is an anionic detergent (e.g., SDS or N-lauroylsarcosine sodium salt solution). In some embodiments, the biological sample can be permeabilized using any of the methods described herein (e.g., using any of the detergents described herein, e.g., SDS and/or N-lauroylsarcosine sodium salt solution) before or after enzymatic treatment (e.g., treatment with any of the enzymes described herein, e.g., trypsin, proteases (e.g., pepsin and/or proteinase K)).

[0234] In some embodiments, a biological sample can be permeabilized by exposing the sample to greater than about 1.0 w/v % (e.g., greater than about 2.0 w/v %, greater than about 3.0 w/v %, greater than about 4.0 w/v %, greater than about 5.0 w/v %, greater than about 6.0 w/v %, greater than about 7.0 w/v %, greater than about 8.0 w/v %, greater than about 9.0 w/v %, greater than about 10.0 w/v %, greater than about 11.0 w/v %, greater than about 12.0 w/v %, or greater than about 13.0 w/v %) sodium dodecyl sulfate (SDS) and/or N-lauroylsarcosine or N-lauroylsarcosine sodium salt. In some embodiments, a biological sample can be permeabilized by exposing the sample (e.g., for about 5 minutes to about 1 hour, about 5 minutes to about 40 minutes, about 5 minutes to about 30 minutes, about 5 minutes to about 20 minutes, or about 5 minutes to about 10 minutes) to about 1.0 w/v % to about 14.0 w/v % (e.g., about 2.0 w/v % to about 14.0 w/v %, about 2.0 w/v % to about 12.0 w/v %, about 2.0 w/v % to about 10.0 w/v %, about 4.0 w/v % to about 14.0 w/v %, about 4.0 w/v % to about 12.0 w/v %, about 4.0 w/v % to about 10.0 w/v %, about 6.0 w/v % to about 14.0 w/v %, about 6.0 w/v % to about 12.0 w/v %, about 6.0 w/v % to about 10.0 w/v %, about 8.0 w/v % to about 14.0 w/v %, about 8.0 w/v % to about 12.0 w/v %, about 8.0 w/v % to about 10.0 w/v %, about 10.0 w/v % to about 14.0 w/v %, about 10.0 w/v % to about 12.0 w/v %, or about 12.0 w/v % to about 14.0 w/v %) SDS and/or N-lauroylsarcosine salt solution and/or proteinase K (e.g., at a temperature of about 4° C. to about 35° C., about 4° C. to about 25° C., about 4° C. to about 20° C., about 4° C. to about 10° C., about 10° C. to about 25° C., about 10° C. to about 20° C., about 10° C. to about 15° C., about 35° C. to about 50° C., about 35° C. to about 45° C., about 35° C. to about 40° C., about 40° C. to about 50° C., about 40° C. to about 45° C., or about 45° C. to about 50° C.).

[0235] In some embodiments, the biological sample can be incubated with a permeabilizing agent to facilitate permeabilization of the sample. Additional methods for sample permeabilization are described, for example, in Jamur et al., *Method Mol. Biol.* 588:63-66, 2010, the entire contents of which are incorporated herein by reference.

Lysis Reagents

[0236] In some embodiments, the biological sample can be permeabilized by adding one or more lysis reagents to the sample. Examples of suitable lysis agents include, but are not limited to, bioactive reagents such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other commercially available lysis enzymes.

[0237] Other lysis agents can additionally or alternatively be added to the biological sample to facilitate permeabilization. For example, surfactant-based lysis solutions can be used to lyse sample cells. Lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). More generally, chemical lysis agents can include, without limitation, organic solvents, chelating agents, detergents, surfactants, and chaotropic agents.

[0238] In some embodiments, the biological sample can be permeabilized by non-chemical permeabilization methods. Non-chemical permeabilization methods are known in the art. For example, non-chemical permeabilization methods that can be used include, but are not limited to, physical lysis techniques such as electroporation, mechanical permeabilization methods (e.g., bead beating using a homogenizer and grinding balls to mechanically disrupt sample tissue structures), acoustic permeabilization (e.g., sonication), and thermal lysis techniques such as heating to induce thermal permeabilization of the sample.

Proteases

[0239] In some embodiments, a medium, solution, or permeabilization solution may contain one or more proteases. In some embodiments, a biological sample treated with a protease capable of degrading histone proteins can result in the generation of fragmented genomic DNA. The fragmented genomic DNA can be captured using the same capture domain (e.g., capture domain having a poly(T) sequence) used to capture mRNA. In some embodiments, a biological sample is treated with a protease capable of degrading histone proteins and an RNA protectant prior to spatial profiling in order to facilitate the capture of both genomic DNA and mRNA.

[0240] In some embodiments, a biological sample is permeabilized by exposing the sample to a protease capable of degrading histone proteins. As used herein, the term “histone protein” typically refers to a linker histone protein (e.g., H1) and/or a core histone protein (e.g., H2A, H2B, H3, and H4). In some embodiments, a protease degrades linker histone proteins, core histone proteins, or linker histone proteins and core histone proteins. Any suitable protease capable of degrading histone proteins in a biological sample can be used. Non-limiting examples of proteases capable of degrading histone proteins include proteases inhibited by leupeptin and TLCK (Tosyl-L-lysyl-chloromethane hydrochloride), a protease encoded by the EUO gene from *Chlamydia trachomatis* serovar A, granzyme A, a serine protease (e.g., trypsin or trypsin-like protease, neutral serine protease, elastase, cathepsin G), an aspartyl protease (e.g., cathepsin D), a peptidase family C1 enzyme (e.g., cathepsin L), pepsin, proteinase K, a protease that is inhibited by the diazomethane inhibitor Z-Phe-Phe-CHN (2) or the epoxide inhibitor E-64, a lysosomal protease, or an azurophilic enzyme (e.g., cathepsin G, elastase, proteinase 3, neutral serine protease). In some embodiments, a serine protease is a trypsin enzyme, trypsin-like enzyme or a functional variant or derivative thereof (e.g., P00761; COHK48; Q8IYP2; Q8BW11; Q6IE06; P35035; P00760; P06871; Q90627; P16049; P07477; P00762; P35031; P19799; P35036; Q29463; P06872; Q90628; P07478; P07146; P00763; P35032; P70059; P29786; P35037; Q90629; P35030; P08426; P35033; P35038; P12788; P29787; P35039; P35040; Q8NHM4; P35041; P35043; P35044; P54624; P04814; P35045; P32821; P54625; P35004; P35046; P32822; P35047; C0HKA5; C0HKA2; P54627; P35005; C0HKA6; C0HKA3; P52905; P83348; P00765; P35042; P81071; P35049; P51588; P35050; P35034; P35051; P24664; P35048; P00764; P00775; P54628; P42278; P54629; P42279; Q91041; P54630; P42280; C0HKA4) or a combination thereof. In some embodiments, a trypsin enzyme is P00761, P00760, Q29463, or a combination thereof. In some embodiments, a protease capable of degrading one or more histone proteins comprises an amino acid sequence with at least 80%

sequence identity to P00761, P00760, or Q29463. In some embodiments, a protease capable of degrading one or more histone proteins comprises an amino acid sequence with at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to P00761, P00760, or Q29463. A protease may be considered a functional variant if it has at least 50% e.g., at least 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the activity relative to the activity of the protease in condition optimum for the enzyme. In some embodiments, the enzymatic treatment with pepsin enzyme, or pepsin like enzyme, can include: P03954/PEPA1_MACFU; P28712/PEPA1_RABIT; P27677/PEPA2_MACFU; P27821/PEPA2_RABIT; P0DJ8/PEPA3_HUMAN; P27822/PEPA3_RABIT; P0DJ7/PEPA4_HUMAN; P27678/PEPA4_MACFU; P28713/PEPA4_RABIT; P0DJ9/PEPA5_HUMAN; Q9D106/PEPA5_MOUSE; P27823/PEPAF_RABIT; P00792/PEPA_BOVIN; Q9N2D4/PEPA_CALJA; Q9GMY6/PEPA_CANLF; P00793/PEPA_CHICK; P11489/PEPA_MACMU; P00791/PEPA_PIG; Q9GMY7/PEPA_RHIFE; Q9GMY8/PEPA_SORUN; P81497/PEPA_SUNMU; P13636/PEPA_URSTH and functional variants and derivatives thereof, or a combination thereof.

In some embodiments, the pepsin enzyme can include: P00791/PEPA_PIG; P00792/PEPA_BOVIN, functional variants, derivatives, or combinations thereof.

[0241] Additionally, the protease may be contained in a reaction mixture (solution), which also includes other components (e.g., buffer, salt, chelator (e.g., EDTA), and/or detergent (e.g., SDS, N-Lauroylsarcosine sodium salt solution)). The reaction mixture may be buffered, having a pH of about 6.5-8.5, e.g., about 7.0-8.0. Additionally, the reaction mixture may be used at any suitable temperature, such as about 10-50° C., e.g., about 10-44° C., 11-43° C., 12-42° C., 13-41° C., 14-40° C., 15-39° C., 16-38° C., 17-37° C., e.g., about 10° C., 12° C., 15° C., 18° C., 20° C., 22° C., 25° C., 28° C., 30° C., 33° C., 35° C. or 37° C., preferably about 35-45° C., e.g., about 37° C.

Other Reagents

[0242] In some embodiments, a permeabilization solution can contain additional reagents or a biological sample may be treated with additional reagents in order to optimize biological sample permeabilization. In some embodiments, an additional reagent is an RNA protectant. As used herein, the term “RNA protectant” typically refers to a reagent that protects RNA from RNA nucleases (e.g., RNases). Any appropriate RNA protectant that protects RNA from degradation can be used. A non-limiting example of a RNA protectant includes organic solvents (e.g., at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% v/v organic solvent), which include, without limitation, ethanol, methanol, propan-2-ol, acetone, trichloroacetic acid, propanol, polyethylene glycol, acetic acid, or a combination thereof. In some embodiments, a RNA protectant includes ethanol, methanol and/or propan-2-ol, or a combination thereof. In some embodiments, a RNA protectant includes RNAlater ICE (ThermoFisher Scientific). In some embodiments, the RNA protectant comprises at least about 60% ethanol. In some embodiments, the RNA protectant comprises about 60-95% ethanol, about 0-35% methanol and about 0-35% propan-2-ol, wherein the total amount of organic solvent in the medium is not more than about 95%. In some embodiments, the RNA protectant comprises about 60-95% ethanol, about 5-20% methanol and about 5-20% propan-2-ol, wherein the total amount of organic solvent in the medium is not more than about 95%.

[0243] In some embodiments, the RNA protectant includes a salt. The salt may include ammonium sulfate, ammonium bisulfate, ammonium chloride, ammonium acetate, cesium sulfate, cadmium sulfate, cesium iron (II) sulfate, chromium (III) sulfate, cobalt (II) sulfate, copper (II) sulfate, lithium chloride, lithium acetate, lithium sulfate, magnesium sulfate, magnesium chloride, manganese sulfate, manganese chloride, potassium chloride, potassium sulfate, sodium chloride, sodium acetate, sodium sulfate, zinc chloride, zinc acetate and zinc sulfate. In some embodiments, the salt is a sulfate salt, for example, ammonium sulfate, ammonium bisulfate, cesium sulfate, cadmium sulfate, cesium iron (II) sulfate, chromium (III) sulfate, cobalt (II) sulfate, copper (II) sulfate, lithium sulfate, magnesium sulfate, manganese sulfate, potassium sulfate, sodium sulfate, or zinc sulfate. In some embodiments, the salt is ammonium sulfate. The salt may be present at a

concentration of about 20 g/100 ml of medium or less, such as about 15 g/100 ml, 10 g/100 ml, 9 g/100 ml, 8 g/100 ml, 7 g/100 ml, 6 g/100 ml, 5 g/100 ml or less, e.g., about 4 g, 3 g, 2 g or 1 g/100 ml.

[0244] Additionally, the RNA protectant may be contained in a medium that further includes a chelator (e.g., EDTA), a buffer (e.g., sodium citrate, sodium acetate, potassium citrate, or potassium acetate, preferably sodium acetate), and/or buffered to a pH between about 4-8 (e.g., about 5).

[0245] In some embodiments, the biological sample is treated with one or more RNA protectants before, contemporaneously with, or after permeabilization. For example, a biological sample is treated with one or more RNA protectants prior to treatment with one or more permeabilization reagents (e.g., one or more proteases). In another example, a biological sample is treated with a solution including one or more RNA protectants and one or more permeabilization reagents (e.g., one or more proteases). In yet another example, a biological sample is treated with one or more RNA protectants after the biological sample has been treated with one or more permeabilization reagents (e.g., one or more proteases). In some embodiments, a biological sample is treated with one or more RNA protectants prior to fixation.

[0246] In some embodiments, identifying the location of the captured analyte in the biological sample includes a nucleic acid extension reaction. In some embodiments where a capture probe captures a fragmented genomic DNA molecule, a nucleic acid extension reaction includes DNA polymerase. For example, a nucleic acid extension reaction includes using a DNA polymerase to extend the capture probe that is hybridized to the captured analyte (e.g., fragmented genomic DNA) using the captured analyte (e.g., fragmented genomic DNA) as a template. The product of the extension reaction includes a spatially-barcoded analyte (e.g., spatially-barcoded fragmented genomic DNA). The spatially-barcoded analyte (e.g., spatially-barcoded fragmented genomic DNA) can be used to identify the spatial location of the analyte in the biological sample. Any DNA polymerase that is capable of extending the capture probe using the captured analyte as a template can be used for the methods described herein. Non-limiting examples of DNA polymerases include T7 DNA polymerase; Bsu DNA polymerase; and *E. coli* DNA Polymerase pol I.

Diffusion-Resistant Media

[0247] In some embodiments, a diffusion-resistant medium, typically used to limit diffusion of analytes, can include at least one permeabilization reagent. For example, the diffusion-resistant medium (e.g., a hydrogel) can include wells (e.g., micro-, nano-, or picowells or pores) containing a permeabilization buffer or reagents. In some embodiments, the diffusion-resistant medium (e.g., a hydrogel) is soaked in permeabilization buffer prior to contacting the hydrogel with a sample. In some embodiments, the hydrogel or other diffusion-resistant medium can contain dried reagents or monomers to deliver permeabilization reagents when the diffusion-resistant medium is applied to a biological sample. In some embodiments, the diffusion-resistant medium, (e.g., hydrogel) is covalently attached to a solid substrate (e.g., an acrylated glass slide).

[0248] In some embodiments, the hydrogel can be modified to both deliver permeabilization reagents and contain capture probes. For example, a hydrogel film can be modified to include spatially-barcoded capture probes. The spatially-barcoded hydrogel film is then soaked in permeabilization buffer before contacting the spatially-barcoded hydrogel film to the sample. In another example, a hydrogel can be modified to include spatially-barcoded capture probes and designed to serve as a porous membrane (e.g., a permeable hydrogel) when exposed to permeabilization buffer or any other biological sample preparation reagent. The permeabilization reagent diffuses through the spatially-barcoded permeable hydrogel and permeabilizes the biological sample on the other side of the hydrogel. The analytes then diffuse into the spatially-barcoded hydrogel after exposure to permeabilization reagents. In such cases, the spatially-barcoded hydrogel (e.g., porous membrane) is facilitating the diffusion of the biological analytes in the biological sample into the hydrogel. In some embodiments, biological analytes diffuse into the hydrogel before exposure to permeabilization reagents (e.g., when secreted analytes are present

outside of the biological sample or in instances where a biological sample is lysed or permeabilized by other means prior to addition of permeabilization reagents). In some embodiments, the permeabilization reagent is flowed over the hydrogel at a variable flow rate (e.g., any flow rate that facilitates diffusion of the permeabilization reagent across the spatially-barcoded hydrogel). In some embodiments, the permeabilization reagents are flowed through a microfluidic chamber or channel over the spatially-barcoded hydrogel. In some embodiments, after using flow to introduce permeabilization reagents to the biological sample, biological sample preparation reagents can be flowed over the hydrogel to further facilitate diffusion of the biological analytes into the spatially-barcoded hydrogel. The spatially-barcoded hydrogel film thus delivers permeabilization reagents to a sample surface in contact with the spatially-barcoded hydrogel, enhancing analyte migration and capture. In some embodiments, the spatially-barcoded hydrogel is applied to a sample and placed in a permeabilization bulk solution. In some embodiments, the hydrogel film soaked in permeabilization reagents is sandwiched between a sample and a spatially-barcoded array. In some embodiments, target analytes are able to diffuse through the permeabilizing reagent soaked hydrogel and hybridize or bind the capture probes on the other side of the hydrogel. In some embodiments, the thickness of the hydrogel is proportional to the resolution loss. In some embodiments, wells (e.g., micro-, nano-, or picowells) can contain spatially-barcoded capture probes and permeabilization reagents and/or buffer. In some embodiments, spatially-barcoded capture probes and permeabilization reagents are held between spacers. In some embodiments, the sample is punch, cut, or transferred into the well, wherein a target analyte diffuses through the permeabilization reagent/buffer and to the spatially-barcoded capture probes. In some embodiments, resolution loss may be proportional to gap thickness (e.g., the amount of permeabilization buffer between the sample and the capture probes). In some embodiments, the diffusion-resistant medium (e.g., hydrogel) is between approximately 50-500 micrometers thick including 500, 450, 400, 350, 300, 250, 200, 150, 100, or 50 micrometers thick, or any thickness within 50 and 500 micrometers.

[0249] In some embodiments, a biological sample is exposed to a porous membrane (e.g., a permeable hydrogel) to aid in permeabilization and limit diffusive analyte losses, while allowing permeabilization reagents to reach a sample. Membrane chemistry and pore volume can be manipulated to minimize analyte loss. In some embodiments, the porous membrane may be made of glass, silicon, paper, hydrogel, polymer monoliths, or other material. In some embodiments, the material may be naturally porous. In some embodiments, the material may have pores or wells etched into solid material. In some embodiments, the permeabilization reagents are flowed through a microfluidic chamber or channel over the porous membrane. In some embodiments, the flow controls the sample's access to the permeabilization reagents. In some embodiments, the porous membrane is a permeable hydrogel. For example, a hydrogel is permeable when permeabilization reagents and/or biological sample preparation reagents can pass through the hydrogel using diffusion. Any suitable permeabilization reagents and/or biological sample preparation reagents described herein can be used under conditions sufficient to release analytes (e.g., nucleic acid, protein, metabolites, lipids, etc.) from the biological sample. In some embodiments, a hydrogel is exposed to the biological sample on one side and permeabilization reagent on the other side. The permeabilization reagent diffuses through the permeable hydrogel and permeabilizes the biological sample on the other side of the hydrogel. In some embodiments, permeabilization reagents are flowed over the hydrogel at a variable flow rate (e.g., any flow rate that facilitates diffusion of the permeabilization reagent across the hydrogel). In some embodiments, the permeabilization reagents are flowed through a microfluidic chamber or channel over the hydrogel. Flowing permeabilization reagents across the hydrogel enables control of the concentration of reagents. In some embodiments, hydrogel chemistry and pore volume can be tuned to enhance permeabilization and limit diffusive analyte losses.

[0250] In some embodiments, a porous membrane is sandwiched between a spatially-barcoded

array and the sample, wherein permeabilization solution is applied over the porous membrane. The permeabilization reagents diffuse through the pores of the membrane and into the biological sample. In some embodiments, the biological sample can be placed on a substrate (e.g., a glass slide). Biological analytes then diffuse through the porous membrane and into the space containing the capture probes. In some embodiments, the porous membrane is modified to include capture probes. For example, the capture probes can be attached to a surface of the porous membrane using any of the methods described herein. In another example, the capture probes can be embedded in the porous membrane at any depth that allows interaction with a biological analyte. In some embodiments, the porous membrane is placed onto a biological sample in a configuration that allows interaction between the capture probes on the porous membrane and the biological analytes from the biological sample. For example, the capture probes are located on the side of the porous membrane that is proximal to the biological sample. In such cases, permeabilization reagents on the other side of the porous membrane diffuse through the porous membrane into the location containing the biological sample and the capture probes in order to facilitate permeabilization of the biological sample (e.g., also facilitating capture of the biological analytes by the capture probes). In some embodiments, the porous membrane is located between the sample and the capture probes. In some embodiments, the permeabilization reagents are flowed through a microfluidic chamber or channel over the porous membrane.

Selective Permeabilization/Selective Lysis

[0251] In some embodiments, biological samples can be processed to selectively release an analyte from a subcellular region of a cell according to established methods. In some embodiments, a method provided herein can include detecting at least one biological analyte present in a subcellular region of a cell in a biological sample. As used herein, a “subcellular region” can refer to any subcellular region. For example, a subcellular region can refer to cytosol, a mitochondria, a nucleus, a nucleolus, an endoplasmic reticulum, a lysosome, a vesicle, a Golgi apparatus, a plastid, a vacuole, a ribosome, cytoskeleton, or combinations thereof. In some embodiments, the subcellular region comprises at least one of cytosol, a nucleus, a mitochondria, and a microsome. In some embodiments, the subcellular region is cytosol. In some embodiments, the subcellular region is a nucleus. In some embodiments, the subcellular region is a mitochondria. In some embodiments, the subcellular region is a microsome.

[0252] For example, a biological analyte can be selectively released from a subcellular region of a cell by selective permeabilization or selective lysing. In some embodiments, “selective permeabilization” can refer to a permeabilization method that can permeabilize a membrane of a subcellular region while leaving a different subcellular region substantially intact (e.g., biological analytes are not released from subcellular region due to the applied permeabilization method). Non-limiting examples of selective permeabilization methods include using electrophoresis and/or applying a permeabilization reagent. In some embodiments, “selective lysing” can refer to a lysis method that can lyse a membrane of a subcellular region while leaving a different subcellular region substantially intact (e.g., biological analytes are not released from subcellular region due to the applied lysis method). Several methods for selective permeabilization or lysis are known to one of skill in the art including the methods described in Lu et al. *Lab Chip*. 2005 January; 5 (1): 23-9; Niklas et al. *Anal Biochem*. 2011 Sep. 15; 416 (2): 218-27; Cox and Emili. *Nat Protoc*. 2006; 1 (4): 1872-8; Chiang et al. *J Biochem. Biophys. Methods*. 2000 Nov. 20; 46 (1-2): 53-68; and Yamauchi and Herr et al. *Microsyst. Nanoeng*. 2017; 3. pii: 16079; each of which is incorporated herein by reference in its entirety.

[0253] In some embodiments, “selective permeabilization” or “selective lysis” refer to the selective permeabilization or selective lysis of a specific cell type. For example, “selective permeabilization” or “selective lysis” can refer to lysing one cell type while leaving a different cell type substantially intact (e.g., biological analytes are not released from the cell due to the applied permeabilization or lysis method). A cell that is a “different cell type” than another cell can refer to a cell from a

different taxonomic kingdom, a prokaryotic cell versus a eukaryotic cell, a cell from a different tissue type, etc. Many methods are known to one of skill in the art for selectively permeabilizing or lysing different cell types. Non-limiting examples include applying a permeabilization reagent, electroporation, and/or sonication. See, e.g., International Application No. WO 2012/168003; Han et al. *Microsyst Nanoeng.* 2019 Jun. 17; 5:30; Gould et al. *Oncotarget.* 2018 Mar. 20; 9 (21): 15606-15615; Oren and Shai. *Biochemistry.* 1997 Feb. 18; 36 (7): 1826-35; Algayer et al. *Molecules.* 2019 May 31; 24 (11). pii: E2079; Hipp et al. *Leukemia.* 2017 October; 31 (10): 2278; International Application No. WO 2012/168003; and U.S. Pat. No. 7,785,869; all of which are incorporated by reference herein in their entireties.

[0254] In some embodiments, applying a selective permeabilization or lysis reagent comprises contacting the biological sample with a hydrogel comprising the permeabilization or lysis reagent.

[0255] In some embodiments, the biological sample is contacted with two or more arrays (e.g., flexible arrays, as described herein). For example, after a subcellular region is permeabilized and a biological analyte from the subcellular region is captured on a first array, the first array can be removed, and a biological analyte from a different subcellular region can be captured on a second array.

(14) Selective Enrichment of RNA Species

[0256] In some embodiments, where RNA is the analyte, one or more RNA analyte species of interest can be selectively enriched (e.g., Adiconis, et. al., Comparative analysis of RNA sequencing methods for degraded and low-input samples, *Nature*, vol. 10, July 2013, 623-632, herein incorporated by reference in its entirety). For example, one or more species of RNA can be selected by addition of one or more oligonucleotides to the sample. In some embodiments, the additional oligonucleotide is a sequence used for priming a reaction by a polymerase. For example, one or more primer sequences with sequence complementarity to one or more RNAs of interest can be used to amplify the one or more RNAs of interest, thereby selectively enriching these RNAs. In some embodiments, an oligonucleotide with sequence complementarity to the complementary strand of captured RNA (e.g., cDNA) can bind to the cDNA. For example, biotinylated oligonucleotides with sequence complementary to one or more cDNAs of interest binds to the cDNA and can be selected using biotinylation-streptavidin affinity using any of a variety of methods known to the field (e.g., streptavidin beads).

[0257] Alternatively, one or more species of RNA (e.g., ribosomal and/or mitochondrial RNA) can be down-selected (e.g., removed, depleted) using any of a variety of methods. Non-limiting examples of a hybridization and capture method of ribosomal RNA depletion include RiboMinus™, RiboCop™, and Ribo-Zero™. Another non-limiting RNA depletion method involves hybridization of complementary DNA oligonucleotides to unwanted RNA followed by degradation of the RNA/DNA hybrids using RNase H. Non-limiting examples of a hybridization and degradation method include NEBNext® rRNA depletion, NuGEN AnyDeplete, or RiboZero Plus. Another non-limiting ribosomal RNA depletion method includes ZapR™ digestion, for example SMARTer. In the SMARTer method, random nucleic acid adapters are hybridized to RNA for first-strand synthesis and tailing by reverse transcriptase, followed by template switching and extension by reverse transcriptase. Additionally, first round PCR amplification adds full-length Illumina sequencing adapters (e.g., Illumina indexes). Ribosomal RNA is cleaved by ZapR v2 and R probes v2. A second round of PCR is performed, amplifying non-rRNA molecules (e.g., cDNA). Parts or steps of these ribosomal depletion protocols/kits can be further combined with the methods described herein to optimize protocols for a specific biological sample.

[0258] In depletion protocols, probes can be administered to a sample that selectively hybridize to ribosomal RNA (rRNA), thereby reducing the pool and concentration of rRNA in the sample. Probes can be administered to a biological sample that selectively hybridize to mitochondria RNA (mtRNA), thereby reducing the pool and concentration of mtRNA in the sample. In some embodiments, probes complementary to mitochondrial RNA can be added during cDNA synthesis,

or probes complementary to both ribosomal and mitochondrial RNA can be added during cDNA synthesis. Subsequent application of capture probes to the sample can result in improved capture of other types of RNA due to a reduction in non-specific RNA (e.g., down-selected RNA) present in the sample. Additionally and alternatively, duplex-specific nuclease (DSN) treatment can remove rRNA (see, e.g., Archer, et al, Selective and flexible depletion of problematic sequences from RNA-seq libraries at the cDNA stage, *BMC Genomics*, 15 401, (2014), the entire contents of which are incorporated herein by reference). Furthermore, hydroxyapatite chromatography can remove abundant species (e.g., rRNA) (see, e.g., Vandernoot, V. A., cDNA normalization by hydroxyapatite chromatography to enrich transcriptome diversity in RNA-seq applications, *Biotechniques*, 53 (6) 373-80, (2012), the entire contents of which are incorporated herein by reference).

(15) Other Reagents

[0259] Additional reagents can be added to a biological sample to perform various functions prior to analysis of the biological sample. In some embodiments, nuclease inhibitors such as DNase and RNase inactivating agents or protease inhibitors, and/or chelating agents such as EDTA, can be added to the biological sample. In other embodiments nucleases, such as DNase or RNase, or proteases, such as pepsin or proteinase K, are added to the sample. In some embodiments, additional reagents may be dissolved in a solution or applied as a medium to the sample. In some embodiments, additional reagents (e.g., pepsin) may be dissolved in HCl prior to applying to the sample. For example, hematoxylin, from an H&E stain, can be optionally removed from the biological sample by washing in dilute HCl (0.001M to 0.1M) prior to further processing. In some embodiments, pepsin can be dissolved in dilute HCl (0.001M to 0.1M) prior to further processing. In some embodiments, biological samples can be washed additional times (e.g., 2, 3, 4, 5, or more times) in dilute HCl prior to incubation with a protease (e.g., pepsin), but after proteinase K treatment.

[0260] In some embodiments, the biological sample can be treated with one or more enzymes. For example, one or more endonucleases to fragment DNA, DNA polymerase enzymes, and dNTPs used to amplify nucleic acids can be added. Other enzymes that can also be added to the biological sample include, but are not limited to, polymerase, transposase, ligase, and DNase, and RNase.

[0261] In some embodiments, reverse transcriptase enzymes can be added to the sample, including enzymes with terminal transferase activity, primers, and template switch oligonucleotides (TSOs). Template switching can be used to increase the length of a cDNA, e.g., by appending a predefined nucleic acid sequence to the cDNA. In some embodiments, the appended nucleic acid sequence comprises one or more ribonucleotides.

[0262] In some embodiments, additional reagents can be added to improve the recovery of one or more target molecules (e.g., cDNA molecules, mRNA transcripts). For example, addition of carrier RNA to a RNA sample workflow process can increase the yield of extracted RNA/DNA hybrids from the biological sample. In some embodiments, carrier molecules are useful when the concentration of input or target molecules is low as compared to remaining molecules. Generally, single target molecules cannot form a precipitate, and addition of the carrier molecules can help in forming a precipitate. Some target molecule recovery protocols use carrier RNA to prevent small amounts of target nucleic acids present in the sample from being irretrievably bound. In some embodiments, carrier RNA can be added immediately prior to a second strand synthesis step. In some embodiments, carrier RNA can be added immediately prior to a second strand cDNA synthesis on oligonucleotides released from an array. In some embodiments, carrier RNA can be added immediately prior to a post in vitro transcription clean-up step. In some embodiments, carrier RNA can be added prior to amplified RNA purification and quantification. In some embodiments, carrier RNA can be added before RNA quantification. In some embodiments, carrier RNA can be added immediately prior to both a second strand cDNA synthesis and a post in vitro transcription clean-up step.

(16) Pre-Processing for Capture Probe Interaction

[0263] In some embodiments, analytes in a biological sample can be pre-processed prior to interaction with a capture probe. For example, prior to interaction with capture probes, polymerization reactions catalyzed by a polymerase (e.g., DNA polymerase or reverse transcriptase) are performed in the biological sample. In some embodiments, a primer for the polymerization reaction includes a functional group that enhances hybridization with the capture probe. The capture probes can include appropriate capture domains to capture biological analytes of interest (e.g., poly(dT) sequence to capture poly(A) mRNA).

[0264] In some embodiments, biological analytes are pre-processed for library generation via next generation sequencing. For example, analytes can be pre-processed by addition of a modification (e.g., ligation of sequences that allow interaction with capture probes). In some embodiments, analytes (e.g., DNA or RNA) are fragmented using fragmentation techniques (e.g., using transposases and/or fragmentation buffers).

[0265] Fragmentation can be followed by a modification of the analyte. For example, a modification can be the addition through ligation of an adapter sequence that allows hybridization with the capture probe. In some embodiments, where the analyte of interest is RNA, poly(A) tailing is performed. Addition of a poly(A) tail to RNA that does not contain a poly(A) tail can facilitate hybridization with a capture probe that includes a capture domain with a functional amount of poly(dT) sequence.

[0266] In some embodiments, prior to interaction with capture probes, ligation reactions catalyzed by a ligase are performed in the biological sample. In some embodiments, ligation can be performed by chemical ligation. In some embodiments, the ligation can be performed using click chemistry as further described below. In some embodiments, the capture domain includes a DNA sequence that has complementarity to a RNA molecule, where the RNA molecule has complementarity to a second DNA sequence, and where the RNA-DNA sequence complementarity is used to ligate the second DNA sequence to the DNA sequence in the capture domain. In these embodiments, direct detection of RNA molecules is possible.

[0267] In some embodiments, prior to interaction with capture probes, target-specific reactions are performed in the biological sample. Examples of target specific reactions include, but are not limited to, ligation of target specific adaptors, probes and/or other oligonucleotides, target specific amplification using primers specific to one or more analytes, and target-specific detection using in situ hybridization, DNA microscopy, and/or antibody detection. In some embodiments, a capture probe includes capture domains targeted to target-specific products (e.g., amplification or ligation).

II. General Spatial Array-Based Analytical Methodology

[0268] Provided herein are methods, apparatus, systems, and compositions for spatial array-based analysis of biological samples.

(a) Spatial Analysis Methods

[0269] Array-based spatial analysis methods involve the transfer of one or more analytes from a biological sample to an array of features on a substrate, where each feature is associated with a unique spatial location on the array. Subsequent analysis of the transferred analytes includes determining the identity of the analytes and the spatial location of each analyte within the biological sample. The spatial location of each analyte within the biological sample is determined based on the feature to which each analyte is bound on the array, and the feature's relative spatial location within the array.

[0270] There are at least two general methods to associate a spatial barcode with one or more neighboring cells, such that the spatial barcode identifies the one or more cells, and/or contents of the one or more cells, as associated with a particular spatial location. One general method is to promote analytes out of a cell and towards the spatially-barcode array. FIG. 1 depicts an exemplary embodiment of this general method. In FIG. 1, the spatially-barcode array populated with capture probes (as described further herein) is contacted with a biological sample **101**, and biological sample is permeabilized, allowing the analyte to migrate away from the sample and

toward the array. The analyte interacts with a capture probe on the spatially-barcoded array **102**. Once the analyte hybridizes/is bound to the capture probe, the sample is optionally removed from the array and the capture probes are analyzed in order to obtain spatially-resolved analyte information **103**.

[0271] Another general method is to cleave the spatially-barcoded capture probes from an array, and promote the spatially-barcoded capture probes towards and/or into or onto the biological sample. FIG. 2 depicts an exemplary embodiment of this general method, the spatially-barcoded array populated with capture probes (as described further herein) can be contacted with a sample **201**. The spatially-barcoded capture probes are cleaved and then interact with cells within the provided biological sample **202**. The interaction can be a covalent or non-covalent cell-surface interaction. The interaction can be an intracellular interaction facilitated by a delivery system or a cell penetration peptide. Once the spatially-barcoded capture probe is associated with a particular cell, the sample can be optionally removed for analysis. The sample can be optionally dissociated before analysis. Once the tagged cell is associated with the spatially-barcoded capture probe, the capture probes can be analyzed to obtain spatially-resolved information about the tagged cell **203**.

[0272] FIG. 3 shows an exemplary workflow that includes preparing a biological sample on a spatially-barcoded array **301**. Sample preparation may include placing the sample on a slide, fixing the sample, and/or staining the biological sample for imaging. The stained sample can be then imaged on the array **302** using both brightfield (to image the sample hematoxylin and eosin stain) and/or fluorescence (to image features) modalities. Optionally, the sample can be destained prior to permeabilization. In some embodiments, analytes are then released from the sample and capture probes forming the spatially-barcoded array hybridize or bind the released analytes **303**. The sample is then removed from the array **304** and the capture probes cleaved from the array **305**. The biological sample and array are then optionally imaged a second time in one or both modalities **305B** while the analytes are reverse transcribed into cDNA, and an amplicon library is prepared **306** and sequenced **307**. Images are then spatially-overlaid in order to correlate spatially-identified biological sample information **308**. When the sample and array are not imaged a second time, **305B**, a spot coordinate file is supplied instead. The spot coordinate file replaces the second imaging step **305B**. Further, amplicon library preparation **306** can be performed with a unique PCR adapter and sequenced **307**.

[0273] FIG. 4 shows another exemplary workflow that utilizes a spatially-barcoded array on a substrate, where spatially-barcoded capture probes are clustered at areas called features. The spatially-barcoded capture probes can include a cleavage domain, one or more functional domains, a spatial barcode, a unique molecular identifier, and a capture domain. The spatially-barcoded capture probes can also include a 5' end modification for reversible attachment to the substrate. The spatially-barcoded array is contacted with a biological sample **401**, and the sample is permeabilized through application of permeabilization reagents **402**. Permeabilization reagents may be administered by placing the array/sample assembly within a bulk solution. Alternatively, permeabilization reagents may be administered to the sample via a diffusion-resistant medium and/or a physical barrier such as a lid, wherein the sample is sandwiched between the diffusion-resistant medium and/or barrier and the array-containing substrate. The analytes are migrated toward the spatially-barcoded capture array using any number of techniques disclosed herein. For example, analyte migration can occur using a diffusion-resistant medium lid and passive migration. As another example, analyte migration can be active migration, using an electrophoretic transfer system, for example. Once the analytes are in close proximity to the spatially-barcoded capture probes, the capture probes can hybridize or otherwise bind a target analyte **403**. The biological sample can be optionally removed from the array **404**.

[0274] The capture probes can be optionally cleaved from the array **405**, and the captured analytes can be spatially-barcoded by performing a reverse transcriptase first strand cDNA reaction. A first strand cDNA reaction can be optionally performed using template switching oligonucleotides. For

example, a template switching oligonucleotide can hybridize to a poly(C) tail added to a 3' end of the cDNA by a reverse transcriptase enzyme in a template independent manner. The original mRNA template and template switching oligonucleotide can then be denatured from the cDNA and the spatially-barcoded capture probe can then hybridize with the cDNA and a complement of the cDNA can be generated. The first strand cDNA can then be purified and collected for downstream amplification steps. The first strand cDNA can be amplified using PCR **406**, where the forward and reverse primers flank the spatial barcode and analyte regions of interest, generating a library associated with a particular spatial barcode **407**. In some embodiments, the library preparation can be quantitated and/or quality controlled to verify the success of the library preparation steps **408**. In some embodiments, the cDNA comprises a sequencing by synthesis (SBS) primer sequence. The library amplicons are sequenced and analyzed to decode spatial information **407**.

[0275] Using the methods, compositions, systems, kits, and devices described herein, RNA transcripts present in biological samples (e.g., tissue samples) can be used for spatial transcriptome analysis. In particular, in some cases, the barcoded oligonucleotides may be configured to prime, replicate, and consequently yield barcoded extension products from an RNA template, or derivatives thereof. For example, in some cases, the barcoded oligonucleotides may include mRNA specific priming sequences, e.g., poly-T primer segments that allow priming and replication of mRNA in a reverse transcription reaction or other targeted priming sequences. Alternatively or additionally, random RNA priming may be carried out using random N-mer primer segments of the barcoded oligonucleotides. Reverse transcriptases (RTs) can use an RNA template and a primer complementary to the 3' end of the RNA template to direct the synthesis of the first strand complementary DNA (cDNA). Many RTs can be used in this reverse transcription reactions, including, for example, avian myeloblastosis virus (AMV) reverse transcriptase, moloney murine leukemia virus (M-MuLV or MMLV), and other variants thereof. Some recombinant M-MuLV reverse transcriptase, such as, for example, PROTOSCRIPT® II reverse transcriptase, can have reduced RNase H activity and increased thermostability when compared to its wild type counterpart, and provide higher specificity, higher yield of cDNA and more full-length cDNA products with up to 12 kilobase (kb) in length. In some embodiments, the reverse transcriptase enzyme is a mutant reverse transcriptase enzyme such as, but not limited to, mutant MMLV reverse transcriptase. In another embodiment, the reverse transcriptase is a mutant MMLV reverse transcriptase such as, but not limited to, one or more variants described in US Patent Publication No. 20180312822 and U.S. Provisional Patent Application No. 62/946,885 filed on Dec. 11, 2019, both of which are incorporated herein by reference in their entireties.

[0276] FIG. 5 depicts an exemplary workflow where the biological sample is removed from the spatially-barcoded array and the spatially-barcoded capture probes are removed from the array for barcoded analyte amplification and library preparation. Another embodiment includes performing first strand synthesis using template switching oligonucleotides on the spatially-barcoded array without cleaving the capture probes. In this embodiment, sample preparation **501** and permeabilization **502** are performed as described elsewhere herein. Once the capture probes capture the analyte(s), first strand cDNA created by template switching and reverse transcriptase **503** is then denatured and the second strand is then extended **504**. The second strand cDNA is then denatured from the first strand cDNA, neutralized, and transferred to a tube **505**. cDNA quantification and amplification can be performed using standard techniques discussed herein. The cDNA can then be subjected to library preparation **506** and indexing **507**, including fragmentation, end-repair, and a-tailing, and indexing PCR steps. The library preparation can optionally be quality controlled to verify the success of the library preparation methods **508**.

[0277] In a non-limiting example of the workflows described above, a biological sample (e.g., tissue section), can be fixed with methanol, stained with hematoxylin and eosin, and imaged. Optionally, the sample can be destained prior to permeabilization. The images can be used to map spatial gene expression patterns back to the biological sample. A permeabilization enzyme can be

used to permeabilize the biological sample directly on the slide. Analytes (e.g., polyadenylated mRNA) released from the overlying cells of the biological sample can be captured by capture probes within a capture area on a substrate. Reverse transcription (RT) reagents can be added to permeabilized biological samples. Incubation with the RT reagents can produce spatially-barcoded full-length cDNA from the captured analytes (e.g., polyadenylated mRNA). Second strand reagents (e.g., second strand primers, enzymes) can be added to the biological sample on the slide to initiate second strand synthesis. The resulting cDNA can be denatured from the capture probe template and transferred (e.g., to a clean tube) for amplification, and/or library construction. The spatially-barcoded, full-length cDNA can be amplified via PCR prior to library construction. The amplicons can then be enzymatically fragmented and/or size-selected in order to provide for desired amplicon size. In some embodiments, when utilizing an Illumina® library preparation methodology, P5 and P7 sequences can be added to the amplicons thereby allowing for capture of the library preparation on a sequencing flowcell (e.g., on Illumina sequencing instruments). Additionally, i7 and i5 can index sequences be added as sample indexes if multiple libraries are to be pooled and sequenced together. Further, Read 1 and Read 2 sequences can be added to the library preparation for sequencing purposes. The aforementioned sequences can be added to a library preparation sample, for example, via End Repair, A-tailing, Adaptor Ligation, and/or PCR. The cDNA fragments can then be sequenced using, for example, paired-end sequencing using TruSeq Read 1 and TruSeq Read 2 as sequencing primer sites.

[0278] In some embodiments, performing correlative analysis of data produced by this workflow, and other workflows described herein, can yield over 95% correlation of genes expressed across two capture areas (e.g., 95% or greater, 96% or greater, 97% or greater, 98% or greater, or 99% or greater). When performing the described workflows using single cell RNA sequencing of nuclei, in some embodiments, correlative analysis of the data can yield over 90% (e.g., over 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) correlation of genes expressed across two capture areas.

[0279] In some embodiments, after cDNA is generated (e.g., by reverse transcription) the cDNA can be amplified directly on the substrate surface. Generating multiple copies of the cDNA (e.g., cDNA synthesized from captured analytes) via amplification directly on the substrate surface can improve final sequencing library complexity. Thus, in some embodiments, cDNA can be amplified directly on the substrate surface by isothermal nucleic acid amplification. In some embodiments, isothermal nucleic acid amplification can amplify RNA or DNA.

[0280] In some embodiments, isothermal amplification can be faster than a standard PCR reaction. In some embodiments, isothermal amplification can be linear amplification (e.g., asymmetrical with a single primer), or exponential amplification (e.g., with two primers). In some embodiments, isothermal nucleic acid amplification can be performed by a template-switching oligonucleotide primer. In some embodiments, the template switching oligonucleotide adds a common sequence onto the 5' end of the RNA being reverse transcribed. For example, after a capture probe interacts with an analyte (e.g., mRNA) and reverse transcription is performed such that additional nucleotides are added to the end of the cDNA creating a 3' overhang as described herein. In some embodiments, a template switching oligonucleotide hybridizes to untemplated poly(C) nucleotides added by a reverse transcriptase to continue replication to the 5' end of the template switching oligonucleotide, thereby generating full-length cDNA ready for further amplification. In some embodiments, the template switching oligonucleotide adds a common 5' sequence to full-length cDNA that is used for cDNA amplification (e.g., a reverse complement of the template switching oligonucleotide).

[0281] In some embodiments, once a full-length cDNA molecule is generated, the template switching oligonucleotide can serve as a primer in a cDNA amplification reaction (e.g., with a DNA polymerase). In some embodiments, double stranded cDNA (e.g., first strand cDNA and second strand reverse complement cDNA) can be amplified via isothermal amplification with either a helicase or recombinase, followed by a strand displacing DNA polymerase. The strand

displacing DNA polymerase can generate a displaced second strand resulting in an amplified product.

[0282] In any of isothermal amplification methods described herein, barcode exchange (e.g., spatial barcode) can occur after the first amplification cycle where there are unused capture probes on the substrate surface. In some embodiments, the free 3'OH end of the unused capture probes can be blocked by any suitable 3'OH blocking method. In some embodiments, the 3'OH can be blocked by hairpin ligation.

[0283] Isothermal nucleic acid amplification can be used in addition to, or as an alternative to standard PCR reactions (e.g., a PCR reaction that requires heating to about 95° C. to denature double stranded DNA). Isothermal nucleic acid amplification generally does not require the use of a thermocycler, however in some embodiments, isothermal amplification can be performed in a thermocycler. In some embodiments, isothermal amplification can be performed from about 35° C. to about 75° C. In some embodiments, isothermal amplification can be performed from about 40° C., about 45° C., about 50° C., about 55° C., about 60° C., about 65° C., or about 70° C. or anywhere in between depending on the polymerase and auxiliary enzymes used.

[0284] Isothermal nucleic acid amplification techniques are known in the art, and can be used alone or in combination with any of the spatial methods described herein. For example, non-limiting examples of suitable isothermal nucleic acid amplification techniques include transcription mediated amplification, nucleic acid sequence-based amplification, signal mediated amplification of RNA technology, strand displacement amplification, rolling circle amplification, loop-mediated isothermal amplification of DNA (LAMP), isothermal multiple displacement amplification, recombinase polymerase amplification, helicase-dependent amplification, single primer isothermal amplification, and circular helicase-dependent amplification (See, e.g., Gill and Ghaemi, Nucleic acid isothermal amplification technologies: a review, *Nucleosides, Nucleotides, & Nucleic Acids*, 27 (3), 224-43, doi: 10.1080/15257770701845204 (2008), which is incorporated herein by reference in its entirety).

[0285] In some embodiments, the isothermal nucleic acid amplification is helicase-dependent nucleic acid amplification. Helicase-dependent isothermal nucleic acid amplification is described in Vincent, et. al., Helicase-dependent isothermal DNA amplification, *EMBO Rep.*, 795-800 (2004) and U.S. Pat. No. 7,282,328, which are both incorporated herein by reference in their entireties. Further, helicase-dependent nucleic acid amplification on a substrate (e.g., on-chip) is described in Andresen, et. al., Helicase-dependent amplification: use in OnChip amplification and potential for point-of-care diagnostics, *Expert Rev Mol Diagn.*, 9, 645-650, doi: 10.1586/erm.09.46 (2009), which is incorporated herein by reference in its entirety. In some embodiments, the isothermal nucleic acid amplification is recombinase polymerase nucleic acid amplification. Recombinase polymerase nucleic acid amplification is described in Piepenburg, et al., DNA Detection Using Recombinant Proteins, *PLOS Biol.*, 4, 7 e204 (2006) and Li, et. al., Review: a comprehensive summary of a decade development of the recombinase polymerase amplification, *Analyst*, 144, 31-67, doi: 10.1039/C8AN01621F (2019), both of which are incorporated herein by reference in their entireties.

[0286] Generally, isothermal amplification techniques use standard PCR reagents (e.g., buffer, dNTPs etc.) known in the art. Some isothermal amplification techniques can require additional reagents. For example, helicase dependent nucleic acid amplification uses a single-strand binding protein and an accessory protein. In another example, recombinase polymerase nucleic acid amplification uses recombinase (e.g., T4 UvsX), recombinase loading factor (e.g., TF UvsY), single-strand binding protein (e.g., T4 gp32), crowding agent (e.g., PEG-35K), and ATP.

[0287] After isothermal nucleic acid amplification of the full-length cDNA described by any of the methods herein, the isothermally amplified cDNAs (e.g., single-stranded or double-stranded) can be recovered from the substrate, and optionally followed by amplification with typical cDNA PCR in microcentrifuge tubes. The sample can then be used with any of the spatial methods described

herein.

(i) Immunohistochemistry and Immunofluorescence

[0288] In some embodiments, immunofluorescence or immunohistochemistry protocols (direct and indirect staining techniques) can be performed as a part of, or in addition to, the exemplary spatial workflows presented herein. For example, tissue sections can be fixed according to methods described herein. The biological sample can be transferred to an array (e.g., capture probe array), wherein analytes (e.g., proteins) are probed using immunofluorescence protocols. For example, the sample can be rehydrated, blocked, and permeabilized (3×SSC, 2% BSA, 0.1% Triton X, 1 U/μl RNase inhibitor for 10 min at 4° C.) before being stained with fluorescent primary antibodies (1:100 in 3×SSC, 2% BSA, 0.1% Triton X, 1 U/μl RNase inhibitor for 30 min at 4° C.). The biological sample can be washed, coverslipped (in glycerol+1 U/μl RNase inhibitor), imaged (e.g., using a confocal microscope or other apparatus capable of fluorescent detection), washed, and processed according to analyte capture or spatial workflows described herein.

[0289] As used herein, an “antigen retrieval buffer” can improve antibody capture in IF/IHC protocols. An exemplary protocol for antigen retrieval can be preheating the antigen retrieval buffer (e.g., to 95° C.), immersing the biological sample in the heated antigen retrieval buffer for a predetermined time, and then removing the biological sample from the antigen retrieval buffer and washing the biological sample.

[0290] In some embodiments, optimizing permeabilization can be useful for identifying intracellular analytes. Permeabilization optimization can include selection of permeabilization agents, concentration of permeabilization agents, and permeabilization duration. Tissue permeabilization is discussed elsewhere herein.

[0291] In some embodiments, blocking an array and/or a biological sample in preparation of labeling the biological sample decreases unspecific binding of the antibodies to the array and/or biological sample (decreases background). Some embodiments provide for blocking buffers/blocking solutions that can be applied before and/or during application of the label, wherein the blocking buffer can include a blocking agent, and optionally a surfactant and/or a salt solution. In some embodiments, a blocking agent can be bovine serum albumin (BSA), serum, gelatin (e.g., fish gelatin), milk (e.g., non-fat dry milk), casein, polyethylene glycol (PEG), polyvinyl alcohol (PVA), or polyvinylpyrrolidone (PVP), biotin blocking reagent, a peroxidase blocking reagent, levamisole, Carnoy's solution, glycine, lysine, sodium borohydride, pontamine sky blue, Sudan Black, trypan blue, FITC blocking agent, and/or acetic acid. The blocking buffer/blocking solution can be applied to the array and/or biological sample prior to and/or during labeling (e.g., application of fluorophore-conjugated antibodies) to the biological sample.

[0292] In some embodiments, additional steps or optimizations can be included in performing IF/IHC protocols in conjunction with spatial arrays. Additional steps or optimizations can be included in performing spatially-tagged analyte capture agent workflows discussed herein.

[0293] In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample (e.g., an analyte present in a biological sample, such as a tissue section) that include: (a) providing a biological sample on a substrate; (b) staining the biological sample on the substrate, imaging the stained biological sample, and selecting the biological sample or subsection of the biological sample (e.g., region of interest) to subject to analysis; (c) providing an array comprising one or more pluralities of capture probes on a substrate; (d) contacting the biological sample with the array, thereby allowing a capture probe of the one or more pluralities of capture probes to capture the analyte of interest; and (e) analyzing the captured analyte, thereby spatially detecting the analyte of interest. Any variety of staining and imaging techniques as described herein or known in the art can be used in accordance with methods described herein. In some embodiments, the staining includes optical labels as described herein, including, but not limited to, fluorescent, radioactive, chemiluminescent, calorimetric, or colorimetric detectable labels. In some

embodiments, the staining includes a fluorescent antibody directed to a target analyte (e.g., cell surface or intracellular proteins) in the biological sample. In some embodiments, the staining includes an immunohistochemistry stain directed to a target analyte (e.g., cell surface or intracellular proteins) in the biological sample. In some embodiments, the staining includes a chemical stain such as hematoxylin and eosin (H&E) or periodic acid-schiff (PAS). In some embodiments, significant time (e.g., days, months, or years) can elapse between staining and/or imaging the biological sample and performing analysis. In some embodiments, reagents for performing analysis are added to the biological sample before, contemporaneously with, or after the array is contacted to the biological sample. In some embodiments, step (d) includes placing the array onto the biological sample. In some embodiments, the array is a flexible array where the plurality of spatially-barcoded features (e.g., a substrate with capture probes, a bead with capture probes) are attached to a flexible substrate. In some embodiments, measures are taken to slow down a reaction (e.g., cooling the temperature of the biological sample or using enzymes that preferentially perform their primary function at lower or higher temperature as compared to their optimal functional temperature) before the array is contacted with the biological sample. In some embodiments, step (e) is performed without bringing the biological sample out of contact with the array. In some embodiments, step (e) is performed after the biological sample is no longer in contact with the array. In some embodiments, the biological sample is tagged with an analyte capture agent before, contemporaneously with, or after staining and/or imaging of the biological sample. In such cases, significant time (e.g., days, months, or years) can elapse between staining and/or imaging and performing analysis. In some embodiments, the array is adapted to facilitate biological analyte migration from the stained and/or imaged biological sample onto the array (e.g., using any of the materials or methods described herein). In some embodiments, a biological sample is permeabilized before being contacted with an array. In some embodiments, the rate of permeabilization is slowed prior to contacting a biological sample with an array (e.g., to limit diffusion of analytes away from their original locations in the biological sample). In some embodiments, modulating the rate of permeabilization (e.g., modulating the activity of a permeabilization reagent) can occur by modulating a condition that the biological sample is exposed to (e.g., modulating temperature, pH, and/or light). In some embodiments, modulating the rate of permeabilization includes use of external stimuli (e.g., small molecules, enzymes, and/or activating reagents) to modulate the rate of permeabilization. For example, a permeabilization reagent can be provided to a biological sample prior to contact with an array, which permeabilization reagent is inactive until a condition (e.g., temperature, pH, and/or light) is changed or an external stimulus (e.g., a small molecule, an enzyme, and/or an activating reagent) is provided.

[0294] In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample (e.g., present in a biological sample such as a tissue section) that include: (a) providing a biological sample on a substrate; (b) staining the biological sample on the substrate, imaging the stained biological sample, and selecting the biological sample or subsection of the biological sample (e.g., a region of interest) to subject to spatial transcriptomic analysis; (c) providing an array comprising one or more pluralities of capture probes on a substrate; (d) contacting the biological sample with the array, thereby allowing a capture probe of the one or more pluralities of capture probes to capture the biological analyte of interest; and (e) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest.

(b) Capture Probes

[0295] A “capture probe” refers to any molecule capable of capturing (directly or indirectly) and/or labelling an analyte (e.g., an analyte of interest) in a biological sample. In some embodiments, the capture probe is a nucleic acid or a polypeptide. In some embodiments, the capture probe is a conjugate (e.g., an oligonucleotide-antibody conjugate). In some embodiments, the capture probe

includes a barcode (e.g., a spatial barcode and/or a unique molecular identifier (UMI)) and a capture domain.

[0296] FIG. 6 is a schematic diagram showing an example of a capture probe, as described herein. As shown, the capture probe **602** is optionally coupled to a feature **601** by a cleavage domain **603**, such as a disulfide linker. The capture probe can include functional sequences that are useful for subsequent processing, such as functional sequence **604**, which can include a sequencer specific flow cell attachment sequence, e.g., a P5 or P7 sequence, as well as functional sequence **606**, which can include sequencing primer sequences, e.g., a R1 primer binding site, a R2 primer binding site. In some embodiments, sequence **604** is a P7 sequence and sequence **606** is a R2 primer binding site. A spatial barcode **605** can be included within the capture probe for use in barcoding the target analyte. The functional sequences can generally be selected for compatibility with any of a variety of different sequencing systems, e.g., Ion Torrent Proton or PGM, Illumina sequencing instruments, PacBio, Oxford Nanopore, etc., and the requirements thereof. In some embodiments, functional sequences can be selected for compatibility with non-commercialized sequencing systems. Examples of such sequencing systems and techniques, for which suitable functional sequences can be used, include (but are not limited to) Ion Torrent Proton or PGM sequencing, Illumina sequencing, PacBio SMRT sequencing, and Oxford Nanopore sequencing. Further, in some embodiments, functional sequences can be selected for compatibility with other sequencing systems, including non-commercialized sequencing systems.

[0297] In some embodiments, the spatial barcode **605**, functional sequences **604** (e.g., flow cell attachment sequence) and **606** (e.g., sequencing primer sequences) can be common to all of the probes attached to a given feature. The spatial barcode can also include a capture domain **607** to facilitate capture of a target analyte.

(i) Capture Domain

[0298] As discussed above, each capture probe includes at least one capture domain. The “capture domain” can be an oligonucleotide, a polypeptide, a small molecule, or any combination thereof, that binds specifically to a desired analyte. In some embodiments, a capture domain can be used to capture or detect a desired analyte.

[0299] In some embodiments, the capture domain is a functional nucleic acid sequence configured to interact with one or more analytes, such as one or more different types of nucleic acids (e.g., RNA molecules and DNA molecules). In some embodiments, the functional nucleic acid sequence can include an N-mer sequence (e.g., a random N-mer sequence), which N-mer sequences are configured to interact with a plurality of DNA molecules. In some embodiments, the functional sequence can include a poly(T) sequence, which poly(T) sequences are configured to interact with messenger RNA (mRNA) molecules via the poly(A) tail of an mRNA transcript. In some embodiments, the functional nucleic acid sequence is the binding target of a protein (e.g., a transcription factor, a DNA binding protein, or a RNA binding protein), where the analyte of interest is a protein.

[0300] Capture probes can include ribonucleotides and/or deoxyribonucleotides as well as synthetic nucleotide residues that are capable of participating in Watson-Crick type or analogous base pair interactions. In some embodiments, the capture domain is capable of priming a reverse transcription reaction to generate cDNA that is complementary to the captured RNA molecules. In some embodiments, the capture domain of the capture probe can prime a DNA extension (polymerase) reaction to generate DNA that is complementary to the captured DNA molecules. In some embodiments, the capture domain can template a ligation reaction between the captured DNA molecules and a surface probe that is directly or indirectly immobilized on the substrate. In some embodiments, the capture domain can be ligated to one strand of the captured DNA molecules. For example, SplintR ligase along with RNA or DNA sequences (e.g., degenerate RNA) can be used to ligate a single-stranded DNA or RNA to the capture domain. In some embodiments, ligases with RNA-templated ligase activity, e.g., SplintR ligase, T4 RNA ligase 2 or KOD ligase, can be used to

ligate a single-stranded DNA or RNA to the capture domain. In some embodiments, a capture domain includes a splint oligonucleotide. In some embodiments, a capture domain captures a splint oligonucleotide.

[0301] In some embodiments, the capture domain is located at the 3' end of the capture probe and includes a free 3' end that can be extended, e.g., by template dependent polymerization, to form an extended capture probe as described herein. In some embodiments, the capture domain includes a nucleotide sequence that is capable of hybridizing to nucleic acid, e.g., RNA or other analyte, present in the cells of the biological sample contacted with the array. In some embodiments, the capture domain can be selected or designed to bind selectively or specifically to a target nucleic acid. For example, the capture domain can be selected or designed to capture mRNA by way of hybridization to the mRNA poly(A) tail. Thus, in some embodiments, the capture domain includes a poly(T) DNA oligonucleotide, e.g., a series of consecutive deoxythymidine residues linked by phosphodiester bonds, which is capable of hybridizing to the poly(A) tail of mRNA. In some embodiments, the capture domain can include nucleotides that are functionally or structurally analogous to a poly(T) tail. For example, a poly(U) oligonucleotide or an oligonucleotide included of deoxythymidine analogues. In some embodiments, the capture domain includes at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides. In some embodiments, the capture domain includes at least 25, 30, or 35 nucleotides.

[0302] In some embodiments, a capture probe includes a capture domain having a sequence that is capable of binding to mRNA and/or genomic DNA. For example, the capture probe can include a capture domain that includes a nucleic acid sequence (e.g., a poly(T) sequence) capable of binding to a poly(A) tail of an mRNA and/or to a poly(A) homopolymeric sequence present in genomic DNA. In some embodiments, a homopolymeric sequence is added to an mRNA molecule or a genomic DNA molecule using a terminal transferase enzyme in order to produce an analyte that has a poly(A) or poly(T) sequence. For example, a poly(A) sequence can be added to an analyte (e.g., a fragment of genomic DNA) thereby making the analyte capable of capture by a poly(T) capture domain.

[0303] In some embodiments, random sequences, e.g., random hexamers or similar sequences, can be used to form all or a part of the capture domain. For example, random sequences can be used in conjunction with poly(T) (or poly(T) analogue) sequences. Thus, where a capture domain includes a poly(T) (or a "poly(T)-like") oligonucleotide, it can also include a random oligonucleotide sequence (e.g., "poly(T)-random sequence" probe). This can, for example, be located 5' or 3' of the poly(T) sequence, e.g., at the 3' end of the capture domain. The poly(T)-random sequence probe can facilitate the capture of the mRNA poly(A) tail. In some embodiments, the capture domain can be an entirely random sequence. In some embodiments, degenerate capture domains can be used.

[0304] In some embodiments, a pool of two or more capture probes form a mixture, where the capture domain of one or more capture probes includes a poly(T) sequence and the capture domain of one or more capture probes includes random sequences. In some embodiments, a pool of two or more capture probes form a mixture where the capture domain of one or more capture probes includes poly(T)-like sequence and the capture domain of one or more capture probes includes random sequences. In some embodiments, a pool of two or more capture probes form a mixture where the capture domain of one or more capture probes includes a poly(T)-random sequences and the capture domain of one or more capture probes includes random sequences. In some embodiments, probes with degenerate capture domains can be added to any of the preceding combinations listed herein. In some embodiments, probes with degenerate capture domains can be substituted for one of the probes in each of the pairs described herein.

[0305] The capture domain can be based on a particular gene sequence or particular motif sequence or common/conserved sequence, that it is designed to capture (i.e., a sequence-specific capture domain). Thus, in some embodiments, the capture domain is capable of binding selectively to a desired sub-type or subset of nucleic acid, for example a particular type of RNA, such as mRNA,

rRNA, tRNA, SRP RNA, tmRNA, snRNA, snoRNA, SmY RNA, scaRNA, gRNA, RNase P, RNase MRP, TERC, SL RNA, aRNA, cis-NAT, crRNA, lncRNA, miRNA, piRNA, siRNA, shRNA, tasiRNA, rasiRNA, 7SK, eRNA, ncRNA or other types of RNA. In a non-limiting example, the capture domain can be capable of binding selectively to a desired subset of ribonucleic acids, for example, microbiome RNA, such as 16S rRNA.

[0306] In some embodiments, a capture domain includes an “anchor” or “anchoring sequence”, which is a sequence of nucleotides that is designed to ensure that the capture domain hybridizes to the intended analyte. In some embodiments, an anchor sequence includes a sequence of nucleotides, including a 1-mer, 2-mer, 3-mer or longer sequence. In some embodiments, the short sequence is random. For example, a capture domain including a poly(T) sequence can be designed to capture an mRNA. In such embodiments, an anchoring sequence can include a random 3-mer (e.g., GGG) that helps ensure that the poly(T) capture domain hybridizes to an mRNA. In some embodiments, an anchoring sequence can be VN, N, or NN. Alternatively, the sequence can be designed using a specific sequence of nucleotides. In some embodiments, the anchor sequence is at the 3' end of the capture domain. In some embodiments, the anchor sequence is at the 5' end of the capture domain.

[0307] In some embodiments, capture domains of capture probes are blocked prior to contacting the biological sample with the array, and blocking probes are used when the nucleic acid in the biological sample is modified prior to its capture on the array. In some embodiments, the blocking probe is used to block or modify the free 3' end of the capture domain. In some embodiments, blocking probes can be hybridized to the capture probes to mask the free 3' end of the capture domain, e.g., hairpin probes, partially double stranded probes, or complementary sequences. In some embodiments, the free 3' end of the capture domain can be blocked by chemical modification, e.g., addition of an azidomethyl group as a chemically reversible capping moiety such that the capture probes do not include a free 3' end. Blocking or modifying the capture probes, particularly at the free 3' end of the capture domain, prior to contacting the biological sample with the array, prevents modification of the capture probes, e.g., prevents the addition of a poly(A) tail to the free 3' end of the capture probes.

[0308] Non-limiting examples of 3' modifications include dideoxy C-3' (3'-ddC), 3' inverted dT, 3' C3 spacer, 3'Amino, and 3' phosphorylation. In some embodiments, the nucleic acid in the biological sample can be modified such that it can be captured by the capture domain. For example, an adaptor sequence (including a binding domain capable of binding to the capture domain of the capture probe) can be added to the end of the nucleic acid, e.g., fragmented genomic DNA. In some embodiments, this is achieved by ligation of the adaptor sequence or extension of the nucleic acid. In some embodiments, an enzyme is used to incorporate additional nucleotides at the end of the nucleic acid sequence, e.g., a poly(A) tail. In some embodiments, the capture probes can be reversibly masked or modified such that the capture domain of the capture probe does not include a free 3' end. In some embodiments, the 3' end is removed, modified, or made inaccessible so that the capture domain is not susceptible to the process used to modify the nucleic acid of the biological sample, e.g., ligation or extension.

[0309] In some embodiments, the capture domain of the capture probe is modified to allow the removal of any modifications of the capture probe that occur during modification of the nucleic acid molecules of the biological sample. In some embodiments, the capture probes can include an additional sequence downstream of the capture domain, e.g., 3' to the capture domain, namely a blocking domain.

[0310] In some embodiments, the capture domain of the capture probe can be a non-nucleic acid domain. Examples of suitable capture domains that are not exclusively nucleic-acid based include, but are not limited to, proteins, peptides, aptamers, antigens, antibodies, and molecular analogs that mimic the functionality of any of the capture domains described herein.

(ii) Cleavage Domain

[0311] Each capture probe can optionally include at least one cleavage domain. The cleavage domain represents the portion of the probe that is used to reversibly attach the probe to an array feature, as will be described further herein. Further, one or more segments or regions of the capture probe can optionally be released from the array feature by cleavage of the cleavage domain. As an example, spatial barcodes and/or universal molecular identifiers (UMIs) can be released by cleavage of the cleavage domain.

[0312] FIG. 7 is a schematic illustrating a cleavable capture probe, wherein the cleaved capture probe can enter into a non-permeabilized cell and bind to analytes within the sample. The capture probe **701** contains a cleavage domain **702**, a cell penetrating peptide **703**, a reporter molecule **704**, and a disulfide bond (—S—S—). **705** represents all other parts of a capture probe, for example a spatial barcode and a capture domain.

[0313] In some embodiments, the cleavage domain linking the capture probe to a feature is a bond capable of cleavage by an enzyme. An enzyme can be added to cleave the cleavage domain, resulting in release of the capture probe from the feature. As another example, heating can also result in degradation of the cleavage domain and release of the attached capture probe from the array feature. In some embodiments, laser radiation is used to heat and degrade cleavage domains of capture probes at specific locations. In some embodiments, the cleavage domain is a photo-sensitive chemical bond (e.g., a chemical bond that dissociates when exposed to light such as ultraviolet light). In some embodiments, the cleavage domain can be an ultrasonic cleavage domain. For example, ultrasonic cleavage can depend on nucleotide sequence, length, pH, ionic strength, temperature, and the ultrasonic frequency (e.g., 22 kHz, 44 kHz) (Grokhovsky, S. L., Specificity of DNA cleavage by ultrasound, *Molecular Biology*, 40 (2), 276-283 (2006)).

[0314] Oligonucleotides with photo-sensitive chemical bonds (e.g., photo-cleavable linkers) have various advantages. They can be cleaved efficiently and rapidly (e.g., in nanoseconds and milliseconds). In some cases, photo-masks can be used such that only specific regions of the array are exposed to cleavable stimuli (e.g., exposure to UV light, exposure to light, exposure to heat induced by laser). When a photo-cleavable linker is used, the cleavable reaction is triggered by light, and can be highly selective to the linker and consequently biorthogonal. Typically, wavelength absorption for the photocleavable linker is located in the near-UV range of the spectrum. In some embodiments, λ_{max} of the photocleavable linker is from about 300 nm to about 400 nm, or from about 310 nm to about 365 nm. In some embodiments, λ_{max} of the photocleavable linker is about 300 nm, about 312 nm, about 325 nm, about 330 nm, about 340 nm, about 345 nm, about 355 nm, about 365 nm, or about 400 nm.

[0315] Non-limiting examples of a photo-sensitive chemical bond that can be used in a cleavage domain include those described in Leriche et al. *Bioorg Med Chem.* 2012 Jan. 15; 20 (2): 571-82 and U.S. Publication No. 2017/0275669, both of which are incorporated by reference herein in their entireties. For example, linkers that comprise photo-sensitive chemical bonds include 3-amino-3-(2-nitrophenyl) propionic acid (ANP), phenacyl ester derivatives, 8-quinolinyl benzenesulfonate, dicoumarin, 6-bromo-7-alkoxycoumarin-4-ylmethoxycarbonyl, a bimane-based linker, and a bis-arylhydrazone based linker. In some embodiments, the photo-sensitive bond is part of a cleavable linker such as an ortho-nitrobenzyl (ONB) linker below:

##STR00001##

wherein:

[0316] X is selected from O and NH; [0317] R_{sup.1} is selected from H and C_{sub.1-3} alkyl; [0318] R_{sup.2} is selected from H and C_{sub.1-3} alkoxy; [0319] n is 1, 2, or 3; and [0320] a and b each represent either the point of attachment of the linker to the substrate, or the point of attachment of the linker to the capture probe.

[0321] In some embodiments, at least one spacer is included between the substrate and the ortho-nitrobenzyl (ONB) linker, and at least one spacer is included between the ortho-nitrobenzyl (ONB) linker and the capture probe. In some aspects of these embodiments, the spacer comprises at least

one group selected from C.sub.1-6 alkylene, C.sub.2-6 alkenylene, C.sub.2-6 alkynylene, C=O, O, S, NH, —(C=O)O—, —(C—O) NH—, —S—S—, ethylene glycol, polyethyleneglycol, propylene glycol, and polypropyleneglycol, or any combination thereof. In some embodiments, X is O. In some embodiments, X is NH. In some embodiments, R.sup.1 is H. In some embodiments, R.sup.1 is C.sub.1-3 alkyl. In some embodiments, R.sup.1 is methyl. In some embodiments, R.sup.2 is H. In some embodiments, R.sup.2 is C.sub.1-3 alkoxy. In some embodiments, R.sup.2 is methoxy. In some embodiments, R.sup.1 is H and R.sup.2 is H. In some embodiments, R.sup.1 is H and R.sup.2 is methoxy. In some embodiments, R.sup.1 is methyl and R.sup.2 is H. In some embodiments, R.sup.1 is methyl and R.sup.2 is methoxy.

In some embodiments, the photocleavable linker has formula:

##STR00002##

In some embodiments, the photocleavable linker has formula:

##STR00003##

In some embodiments, the photocleavable linker has formula:

##STR00004##

In some embodiments, the photocleavable linker has formula:

##STR00005##

In some embodiments, the photocleavable linker has formula:

##STR00006##

Without being bound to any particular theory, it is believed that excitation of the ortho-nitrobenzyl (ONB) linker leads to Norrish-type hydrogen abstraction in the γ -position, followed by formation of azinic acid, which is highly reactive and rearranges into nitroso compound, resulting in the complete cleavage of the linker, as shown on the following scheme:

##STR00007##

In some embodiments, the photocleavable linker is 3-amino-3-(2-nitrophenyl) propionic acid (ANP) linker:

##STR00008## [0322] wherein X, R.sup.2, n, a, and b are as described herein for the ortho-nitrobenzyl (ONB) linker.

In some embodiments, the photocleavable linker has formula:

##STR00009##

In some embodiments, the photocleavable linker is phenacyl ester linker:

##STR00010##

wherein a and b are as described herein for the ortho-nitrobenzyl (ONB) linker.

[0323] Other examples of photo-sensitive chemical bonds that can be used in a cleavage domain include halogenated nucleosides such as bromodeoxyuridine (BrdU). BrdU is an analog of thymidine that can be readily incorporated into oligonucleotides (e.g., in the cleavage domain of a capture probe), and is sensitive to UVB light (280-320 nm range). Upon exposure to UVB light, a photo-cleavage reaction occurs (e.g., at a nucleoside immediately 5' to the site of BrdU incorporation (Doddridge et al. Chem. Comm., 1998, 18:1997-1998 and Cook et al. Chemistry and Biology. 1999, 6:451-459)) that results in release of the capture probe from the feature.

[0324] Other examples of cleavage domains include labile chemical bonds such as, but not limited to, ester linkages (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), an abasic or apurinic/apyrimidinic (AP) site (e.g., cleavable with an alkali or an AP endonuclease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)).

[0325] In some embodiments, the cleavage domain includes a sequence that is recognized by one or more enzymes capable of cleaving a nucleic acid molecule, e.g., capable of breaking the phosphodiester linkage between two or more nucleotides. A bond can be cleavable via other nucleic

acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases). For example, the cleavage domain can include a restriction endonuclease (restriction enzyme) recognition sequence. Restriction enzymes cut double-stranded or single stranded DNA at specific recognition nucleotide sequences known as restriction sites. In some embodiments, a rare-cutting restriction enzyme, e.g., enzymes with a long recognition site (at least 8 base pairs in length), is used to reduce the possibility of cleaving elsewhere in the capture probe.

[0326] In some embodiments, the cleavage domain includes a poly(U) sequence which can be cleaved by a mixture of Uracil DNA glycosylase (UDG) and the DNA glycosylase-lyase Endonuclease VIII, commercially known as the USER[™] enzyme. Releasable capture probes can be available for reaction once released. Thus, for example, an activatable capture probe can be activated by releasing the capture probes from a feature.

[0327] In some embodiments, where the capture probe is attached indirectly to a substrate, e.g., via a surface probe, the cleavage domain includes one or more mismatch nucleotides, so that the complementary parts of the surface probe and the capture probe are not 100% complementary (for example, the number of mismatched base pairs can be one, two, or three base pairs). Such a mismatch is recognized, e.g., by the MutY and T7 endonuclease I enzymes, which results in cleavage of the nucleic acid molecule at the position of the mismatch. As described herein a “surface probe” can be any moiety present on the surface of the substrate capable of attaching to an agent (e.g., a capture probe). In some embodiments, the surface probe is an oligonucleotide. In some embodiments, the surface probe is part of the capture probe.

[0328] In some embodiments, where the capture probe is attached (e.g., immobilized) to a feature indirectly, e.g., via a surface probe, the cleavage domain includes a nickase recognition site or sequence. Nickases are endonucleases which cleave only a single strand of a DNA duplex. Thus, the cleavage domain can include a nickase recognition site close to the 5' end of the surface probe (and/or the 5' end of the capture probe) such that cleavage of the surface probe or capture probe destabilizes the duplex between the surface probe and capture probe thereby releasing the capture probe) from the feature.

[0329] Nickase enzymes can also be used in some embodiments where the capture probe is attached (e.g., immobilized) to the feature directly. For example, the substrate can be contacted with a nucleic acid molecule that hybridizes to the cleavage domain of the capture probe to provide or reconstitute a nickase recognition site, e.g., a cleavage helper probe. Thus, contact with a nickase enzyme will result in cleavage of the cleavage domain thereby releasing the capture probe from the feature. Such cleavage helper probes can also be used to provide or reconstitute cleavage recognition sites for other cleavage enzymes, e.g., restriction enzymes.

[0330] Some nickases introduce single-stranded nicks only at particular sites on a DNA molecule, by binding to and recognizing a particular nucleotide recognition sequence. A number of naturally-occurring nickases have been discovered, of which at present the sequence recognition properties have been determined for at least four. Nickases are described in U.S. Pat. No. 6,867,028, which is incorporated herein by reference in its entirety. In general, any suitable nickase can be used to bind to a complementary nickase recognition site of a cleavage domain. Following use, the nickase enzyme can be removed from the assay or inactivated following release of the capture probes to prevent unwanted cleavage of the capture probes.

[0331] Examples of suitable capture domains that are not exclusively nucleic-acid based include, but are not limited to, proteins, peptides, aptamers, antigens, antibodies, and molecular analogs that mimic the functionality of any of the capture domains described herein.

[0332] In some embodiments, a cleavage domain is absent from the capture probe. Examples of substrates with attached capture probes lacking a cleavage domain are described for example in Macosko et al., (2015) Cell 161, 1202-1214, the entire contents of which are incorporated herein by reference.

[0333] In some embodiments, the region of the capture probe corresponding to the cleavage

domain can be used for some other function. For example, an additional region for nucleic acid extension or amplification can be included where the cleavage domain would normally be positioned. In such embodiments, the region can supplement the functional domain or even exist as an additional functional domain. In some embodiments, the cleavage domain is present but its use is optional.

(iii) Functional Domain

[0334] Each capture probe can optionally include at least one functional domain. Each functional domain typically includes a functional nucleotide sequence for a downstream analytical step in the overall analysis procedure.

[0335] In some embodiments, the capture probe can include a functional domain for attachment to a sequencing flow cell, such as, for example, a P5 sequence for Illumina® sequencing. In some embodiments, the capture probe or derivative thereof can include another functional domain, such as, for example, a P7 sequence for attachment to a sequencing flow cell for Illumina® sequencing. The functional domains can be selected for compatibility with a variety of different sequencing systems, e.g., 454 Sequencing, Ion Torrent Proton or PGM, Illumina X10, etc., and the requirements thereof.

[0336] In some embodiments, the functional domain includes a primer. The primer can include an R1 primer sequence for Illumina® sequencing, and in some embodiments, an R2 primer sequence for Illumina® sequencing. Examples of such capture probes and uses thereof are described in U.S. Patent Publication Nos. 2014/0378345 and 2015/0376609, the entire contents of each of which are incorporated herein by reference.

(iv) Spatial Barcode

[0337] As discussed above, the capture probe can include one or more spatial barcodes (e.g., two or more, three or more, four or more, five or more) spatial barcodes. A “spatial barcode” is a contiguous nucleic acid segment or two or more non-contiguous nucleic acid segments that function as a label or identifier that conveys or is capable of conveying spatial information. In some embodiments, a capture probe includes a spatial barcode that possesses a spatial aspect, where the barcode is associated with a particular location within an array or a particular location on a substrate.

[0338] A spatial barcode can be part of an analyte, or independent from an analyte (e.g., part of the capture probe). A spatial barcode can be a tag attached to an analyte (e.g., a nucleic acid molecule) or a combination of a tag in addition to an endogenous characteristic of the analyte (e.g., size of the analyte or end sequence(s)). A spatial barcode can be unique. In some embodiments where the spatial barcode is unique, the spatial barcode functions both as a spatial barcode and as a unique molecular identifier (UMI), associated with one particular capture probe.

[0339] Spatial barcodes can have a variety of different formats. For example, spatial barcodes can include polynucleotide spatial barcodes; random nucleic acid and/or amino acid sequences; and synthetic nucleic acid and/or amino acid sequences. In some embodiments, a spatial barcode is attached to an analyte in a reversible or irreversible manner. In some embodiments, a spatial barcode is added to, for example, a fragment of a DNA or RNA sample before, during, and/or after sequencing of the sample. In some embodiments, a spatial barcode allows for identification and/or quantification of individual sequencing-reads. In some embodiments, a spatial barcode is used as a fluorescent barcode for which fluorescently labeled oligonucleotide probes hybridize to the spatial barcode.

[0340] In some embodiments, the spatial barcode is a nucleic acid sequence that does not substantially hybridize to analyte nucleic acid molecules in a biological sample. In some embodiments, the spatial barcode has less than 80% sequence identity (e.g., less than 70%, 60%, 50%, or less than 40% sequence identity) to the nucleic acid sequences across a substantial part (e.g., 80% or more) of the nucleic acid molecules in the biological sample.

[0341] The spatial barcode sequences can include from about 6 to about 20 or more nucleotides

within the sequence of the capture probes. In some embodiments, the length of a spatial barcode sequence can be about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a spatial barcode sequence can be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a spatial barcode sequence is at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or shorter.

[0342] These nucleotides can be completely contiguous, e.g., in a single stretch of adjacent nucleotides, or they can be separated into two or more separate subsequences that are separated by 1 or more nucleotides. Separated spatial barcode subsequences can be from about 4 to about 16 nucleotides in length. In some embodiments, the spatial barcode subsequence can be about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the spatial barcode subsequence can be at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the spatial barcode subsequence can be at most about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or shorter.

[0343] For multiple capture probes that are attached to a common array feature, the one or more spatial barcode sequences of the multiple capture probes can include sequences that are the same for all capture probes coupled to the feature, and/or sequences that are different across all capture probes coupled to the feature.

[0344] FIG. 8 is a schematic diagram of an exemplary multiplexed spatially-barcoded feature. In FIG. 8, the feature **801** can be coupled to spatially-barcoded capture probes, wherein the spatially-barcoded probes of a particular feature can possess the same spatial barcode, but have different capture domains designed to associate the spatial barcode of the feature with more than one target analyte. For example, a feature may be coupled to four different types of spatially-barcoded capture probes, each type of spatially-barcoded capture probe possessing the spatial barcode **802**. One type of capture probe associated with the feature includes the spatial barcode **802** in combination with a poly(T) capture domain **803**, designed to capture mRNA target analytes. A second type of capture probe associated with the feature includes the spatial barcode **802** in combination with a random N-mer capture domain **804** for gDNA analysis. A third type of capture probe associated with the feature includes the spatial barcode **802** in combination with a capture domain complementary to the analyte capture agent of interest **805**. A fourth type of capture probe associated with the feature includes the spatial barcode **802** in combination with a capture probe that can specifically bind a nucleic acid molecule **806** that can function in a CRISPR assay (e.g., CRISPR/Cas9). While only four different capture probe-barcoded constructs are shown in FIG. 8, capture-probe barcoded constructs can be tailored for analyses of any given analyte associated with a nucleic acid and capable of binding with such a construct. For example, the schemes shown in FIG. 8 can also be used for concurrent analysis of other analytes disclosed herein, including, but not limited to: (a) mRNA, a lineage tracing construct, cell surface or intracellular proteins and metabolites, and gDNA; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq) cell surface or intracellular proteins and metabolites, and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (c) mRNA, cell surface or intracellular proteins and/or metabolites, a barcoded labelling agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor). In some embodiments, a perturbation agent can be a small molecule, an antibody, a drug, an aptamer, a miRNA, a physical environmental (e.g., temperature change), or any other known perturbation agents.

[0345] Capture probes attached to a single array feature can include identical (or common) spatial barcode sequences, different spatial barcode sequences, or a combination of both. Capture probes attached to a feature can include multiple sets of capture probes. Capture probes of a given set can include identical spatial barcode sequences. The identical spatial barcode sequences can be different from spatial barcode sequences of capture probes of another set.

[0346] The plurality of capture probes can include spatial barcode sequences (e.g., nucleic acid barcode sequences) that are associated with specific locations on a spatial array. For example, a first plurality of capture probes can be associated with a first region, based on a spatial barcode sequence common to the capture probes within the first region, and a second plurality of capture probes can be associated with a second region, based on a spatial barcode sequence common to the capture probes within the second region. The second region may or may not be associated with the first region. Additional pluralities of capture probes can be associated with spatial barcode sequences common to the capture probes within other regions. In some embodiments, the spatial barcode sequences can be the same across a plurality of capture probe molecules.

[0347] In some embodiments, multiple different spatial barcodes are incorporated into a single arrayed capture probe. For example, a mixed but known set of spatial barcode sequences can provide a stronger address or attribution of the spatial barcodes to a given spot or location, by providing duplicate or independent confirmation of the identity of the location. In some embodiments, the multiple spatial barcodes represent increasing specificity of the location of the particular array point.

(v) Unique Molecular Identifier

[0348] The capture probe can include one or more (e.g., two or more, three or more, four or more, five or more) Unique Molecular Identifiers (UMIs). A unique molecular identifier is a contiguous nucleic acid segment or two or more non-contiguous nucleic acid segments that function as a label or identifier for a particular analyte, or for a capture probe that binds a particular analyte (e.g., via the capture domain).

[0349] A UMI can be unique. A UMI can include one or more specific polynucleotide sequences, one or more random nucleic acid and/or amino acid sequences, and/or one or more synthetic nucleic acid and/or amino acid sequences, or combinations thereof.

[0350] In some embodiments, the UMI is a nucleic acid sequence that does not substantially hybridize to analyte nucleic acid molecules in a biological sample. In some embodiments, the UMI has less than 80% sequence identity (e.g., less than 70%, 60%, 50%, or less than 40% sequence identity) to the nucleic acid sequences across a substantial part (e.g., 80% or more) of the nucleic acid molecules in the biological sample.

[0351] The UMI can include from about 6 to about 20 or more nucleotides within the sequence of the capture probes. In some embodiments, the length of a UMI sequence can be about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a UMI sequence can be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a UMI sequence is at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or shorter.

[0352] These nucleotides can be completely contiguous, i.e., in a single stretch of adjacent nucleotides, or they can be separated into two or more separate subsequences that are separated by 1 or more nucleotides. Separated UMI subsequences can be from about 4 to about 16 nucleotides in length. In some embodiments, the UMI subsequence can be about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the UMI subsequence can be at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the UMI subsequence can be at most about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or shorter.

[0353] In some embodiments, a UMI is attached to an analyte in a reversible or irreversible manner. In some embodiments, a UMI is added to, for example, a fragment of a DNA or RNA sample before, during, and/or after sequencing of the analyte. In some embodiments, a UMI allows for identification and/or quantification of individual sequencing-reads. In some embodiments, a UMI is used as a fluorescent barcode for which fluorescently labeled oligonucleotide probes hybridize to the UMI.

(vi) Other Aspects of Capture Probes

[0354] For capture probes that are attached to an array feature, an individual array feature can

include one or more capture probes. In some embodiments, an individual array feature includes hundreds or thousands of capture probes. In some embodiments, the capture probes are associated with a particular individual feature, where the individual feature contains a capture probe including a spatial barcode unique to a defined region or location on the array.

[0355] In some embodiments, a particular feature can contain capture probes including more than one spatial barcode (e.g., one capture probe at a particular feature can include a spatial barcode that is different than the spatial barcode included in another capture probe at the same particular feature, while both capture probes include a second, common spatial barcode), where each spatial barcode corresponds to a particular defined region or location on the array. For example, multiple spatial barcode sequences associated with one particular feature on an array can provide a stronger address or attribution to a given location by providing duplicate or independent confirmation of the location. In some embodiments, the multiple spatial barcodes represent increasing specificity of the location of the particular array point. In a non-limiting example, a particular array point can be coded with two different spatial barcodes, where each spatial barcode identifies a particular defined region within the array, and an array point possessing both spatial barcodes identifies the sub-region where two defined regions overlap, e.g., such as the overlapping portion of a Venn diagram.

[0356] In another non-limiting example, a particular array point can be coded with three different spatial barcodes, where the first spatial barcode identifies a first region within the array, the second spatial barcode identifies a second region, where the second region is a subregion entirely within the first region, and the third spatial barcode identifies a third region, where the third region is a subregion entirely within the first and second subregions.

[0357] In some embodiments, capture probes attached to array features are released from the array features for sequencing. Alternatively, in some embodiments, capture probes remain attached to the array features, and the probes are sequenced while remaining attached to the array features (e.g., via in situ sequencing). Further aspects of the sequencing of capture probes are described in subsequent sections of this disclosure.

[0358] In some embodiments, an array feature can include different types of capture probes attached to the feature. For example, the array feature can include a first type of capture probe with a capture domain designed to bind to one type of analyte, and a second type of capture probe with a capture domain designed to bind to a second type of analyte. In general, array features can include one or more (e.g., two or more, three or more, four or more, five or more, six or more, eight or more, ten or more, 12 or more, 15 or more, 20 or more, 30 or more, 50 or more) different types of capture probes attached to a single array feature.

[0359] In some embodiments, the capture probe is nucleic acid. In some embodiments, the capture probe is attached to the array feature via its 5' end. In some embodiments, the capture probe includes from the 5' to 3' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe includes from the 5' to 3' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain. In some embodiments, the capture probe includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), a second functional domain, and a capture domain. In some embodiments, the capture probe includes from the 5' to 3' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain. In some embodiments, the capture probe does not include a spatial barcode. In some embodiments, the capture probe does not include a UMI. In some embodiments, the capture probe includes a sequence for initiating a sequencing reaction.

[0360] In some embodiments, the capture probe is immobilized on a feature via its 3' end. In some embodiments, the capture probe includes from the 3' to 5' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe

includes from the 3' to 5' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe includes from the 3' to 5' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain. In some embodiments, the capture probe includes from the 3' to 5' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain.

[0361] In some embodiments, a capture probe includes an in situ synthesized oligonucleotide. The in situ synthesized oligonucleotide can be attached to a substrate, or to a feature on a substrate. In some embodiments, the in situ synthesized oligonucleotide includes one or more constant sequences, one or more of which serves as a priming sequence (e.g., a primer for amplifying target nucleic acids). The in situ synthesized oligonucleotide can, for example, include a constant sequence at the 3' end that is attached to a substrate, or attached to a feature on a substrate. Additionally or alternatively, the in situ synthesized oligonucleotide can include a constant sequence at the free 5' end. In some embodiments, the one or more constant sequences can be a cleavable sequence. In some embodiments, the in situ synthesized oligonucleotide includes a barcode sequence, e.g., a variable barcode sequence. The barcode can be any of the barcodes described herein. The length of the barcode can be approximately 8 to 16 nucleotides (e.g., 8, 9, 10, 11, 12, 13, 14, 15, or 16 nucleotides). The length of the in situ synthesized oligonucleotide can be less than 100 nucleotides (e.g., less than 90, 80, 75, 70, 60, 50, 45, 40, 35, 30, 25 or 20 nucleotides). In some instances, the length of the in situ synthesized oligonucleotide is about 20 to about 40 nucleotides. Exemplary in situ synthesized oligonucleotides are produced by Affymetrix. In some embodiments, the in situ synthesized oligonucleotide is attached to a feature of an array.

[0362] Additional oligonucleotides can be ligated to an in situ synthesized oligonucleotide to generate a capture probe. For example, a primer complementary to a portion of the in situ synthesized oligonucleotide (e.g., a constant sequence in the oligonucleotide) can be used to hybridize an additional oligonucleotide and extend (using the in situ synthesized oligonucleotide as a template e.g., a primer extension reaction) to form a double stranded oligonucleotide and to further create a 3' overhang. In some embodiments, the 3' overhang can be created by template-independent ligases (e.g., terminal deoxynucleotidyl transferase (TdT) or poly(A) polymerase). An additional oligonucleotide comprising one or more capture domains can be ligated to the 3' overhang using a suitable enzyme (e.g., a ligase) and a splint oligonucleotide, to generate a capture probe. Thus, in some embodiments, a capture probe is a product of two or more oligonucleotide sequences, (e.g., the in situ synthesized oligonucleotide and the additional oligonucleotide) that are ligated together. In some embodiments, one of the oligonucleotide sequences is an in situ synthesized oligonucleotide.

[0363] In some embodiments, the capture probe can be prepared using a splint oligonucleotide (e.g., any of the splint oligonucleotides described herein). Two or more oligonucleotides can be ligated together using a splint oligonucleotide and any variety of ligases known in the art or described herein (e.g., SplintR ligase).

[0364] One of the oligonucleotides can include, for example, a constant sequence (e.g., a sequence complementary to a portion of a splint oligonucleotide), a degenerate sequence, and/or a capture domain (e.g., as described herein). One of the oligonucleotides can also include a sequence compatible for ligating or hybridizing to an analyte of interest in the biological sample. An analyte of interest (e.g., an mRNA) can also be used as a splint oligonucleotide to ligate further oligonucleotides onto the capture probe. In some embodiments, the capture probe is generated by having an enzyme add polynucleotides at the end of an oligonucleotide sequence. The capture probe can include a degenerate sequence, which can function as a unique molecular identifier.

[0365] A degenerate sequence, which is a sequence in which some positions of a nucleotide sequence contain a number of possible bases. A degenerate sequence can be a degenerate nucleotide sequence including about or at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, or 50 nucleotides. In some embodiments, a nucleotide sequence contains 1,

2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, or more degenerate positions within the nucleotide sequence. In some embodiments, the degenerate sequence is used as a UMI.

[0366] In some embodiments, a capture probe includes a restriction endonuclease recognition sequence or a sequence of nucleotides cleavable by specific enzyme activities. For example, uracil sequences can be enzymatically cleaved from a nucleotide sequence using uracil DNA glycosylase (UDG) or Uracil Specific Excision Reagent (USER). As another example, other modified bases (e.g., modified by methylation) can be recognized and cleaved by specific endonucleases. The capture probes can be subjected to an enzymatic cleavage, which removes the blocking domain and any of the additional nucleotides that are added to the 3' end of the capture probe during the modification process. Removal of the blocking domain reveals and/or restores the free 3' end of the capture domain of the capture probe. In some embodiments, additional nucleotides can be removed to reveal and/or restore the 3' end of the capture domain of the capture probe.

[0367] In some embodiments, a blocking domain can be incorporated into the capture probe when it is synthesized, or after its synthesis. The terminal nucleotide of the capture domain is a reversible terminator nucleotide (e.g., 3'-O-blocked reversible terminator and 3'-unblocked reversible terminator), and can be included in the capture probe during or after probe synthesis.

(vii) Extended Capture Probes

[0368] An “extended capture probe” is a capture probe with an enlarged nucleic acid sequence. For example, where the capture probe includes nucleic acid, an “extended 3' end” indicates that further nucleotides were added to the most 3' nucleotide of the capture probe to extend the length of the capture probe, for example, by standard polymerization reactions utilized to extend nucleic acid molecules including templated polymerization catalyzed by a polymerase (e.g., a DNA polymerase or reverse transcriptase).

[0369] In some embodiments, extending the capture probe includes generating cDNA from the captured (hybridized) RNA. This process involves synthesis of a complementary strand of the hybridized nucleic acid, e.g., generating cDNA based on the captured RNA template (the RNA hybridized to the capture domain of the capture probe). Thus, in an initial step of extending the capture probe, e.g., the cDNA generation, the captured (hybridized) nucleic acid, e.g., RNA, acts as a template for the extension, e.g., reverse transcription, step.

[0370] In some embodiments, the capture probe is extended using reverse transcription. For example, reverse transcription includes synthesizing cDNA (complementary or copy DNA) from RNA, e.g., (messenger RNA), using a reverse transcriptase. In some embodiments, reverse transcription is performed while the tissue is still in place, generating an analyte library, where the analyte library includes the spatial barcodes from the adjacent capture probes. In some embodiments, the capture probe is extended using one or more DNA polymerases.

[0371] In some embodiments, the capture domain of the capture probe includes a primer for producing the complementary strand of the nucleic acid hybridized to the capture probe, e.g., a primer for DNA polymerase and/or reverse transcription. The nucleic acid, e.g., DNA and/or cDNA, molecules generated by the extension reaction incorporate the sequence of the capture probe. The extension of the capture probe, e.g., a DNA polymerase and/or reverse transcription reaction, can be performed using a variety of suitable enzymes and protocols.

[0372] In some embodiments, a full-length DNA, e.g., cDNA, molecule is generated. In some embodiments, a “full-length” DNA molecule refers to the whole of the captured nucleic acid molecule. However, if the nucleic acid, e.g., RNA, was partially degraded in the tissue sample, then the captured nucleic acid molecules will not be the same length as the initial RNA in the tissue sample. In some embodiments, the 3' end of the extended probes, e.g., first strand cDNA molecules, is modified. For example, a linker or adaptor can be ligated to the 3' end of the extended probes. This can be achieved using single stranded ligation enzymes such as T4 RNA ligase or Circligase™ (available from Lucigen, Middleton, WI). In some embodiments, template switching oligonucleotides are used to extend cDNA in order to generate a full-length cDNA (or as close to a

full-length cDNA as possible). In some embodiments, a second strand synthesis helper probe (a partially double stranded DNA molecule capable of hybridizing to the 3' end of the extended capture probe), can be ligated to the 3' end of the extended probe, e.g., first strand cDNA, molecule using a double stranded ligation enzyme such as T4 DNA ligase. Other enzymes appropriate for the ligation step are known in the art and include, e.g., Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9°N) DNA ligase (9° NTM DNA ligase, New England Biolabs), AmpligaseTM (available from Lucigen, Middleton, WI), and SplintR (available from New England Biolabs, Ipswich, MA). In some embodiments, a polynucleotide tail, e.g., a poly(A) tail, is incorporated at the 3' end of the extended probe molecules. In some embodiments, the polynucleotide tail is incorporated using a terminal transferase active enzyme.

[0373] In some embodiments, double-stranded extended capture probes are treated to remove any unextended capture probes prior to amplification and/or analysis, e.g., sequence analysis. This can be achieved by a variety of methods, e.g., using an enzyme to degrade the unextended probes, such as an exonuclease enzyme, or purification columns.

[0374] In some embodiments, extended capture probes are amplified to yield quantities that are sufficient for analysis, e.g., via DNA sequencing. In some embodiments, the first strand of the extended capture probes (e.g., DNA and/or cDNA molecules) acts as a template for the amplification reaction (e.g., a polymerase chain reaction).

[0375] In some embodiments, the amplification reaction incorporates an affinity group onto the extended capture probe (e.g., RNA-cDNA hybrid) using a primer including the affinity group. In some embodiments, the primer includes an affinity group and the extended capture probes includes the affinity group. The affinity group can correspond to any of the affinity groups described previously.

[0376] In some embodiments, the extended capture probes including the affinity group can be coupled to an array feature specific for the affinity group. In some embodiments, the substrate can include an antibody or antibody fragment. In some embodiments, the array feature includes avidin or streptavidin and the affinity group includes biotin. In some embodiments, the array feature includes maltose and the affinity group includes maltose-binding protein. In some embodiments, the array feature includes maltose-binding protein and the affinity group includes maltose. In some embodiments, amplifying the extended capture probes can function to release the extended probes from the array feature, insofar as copies of the extended probes are not attached to the array feature.

[0377] In some embodiments, the extended capture probe or complement or amplicon thereof is released from an array feature. The step of releasing the extended capture probe or complement or amplicon thereof from an array feature can be achieved in a number of ways. In some embodiments, an extended capture probe or a complement thereof is released from the feature by nucleic acid cleavage and/or by denaturation (e.g., by heating to denature a double-stranded molecule).

[0378] In some embodiments, the extended capture probe or complement or amplicon thereof is released from the array feature by physical means. For example, methods for inducing physical release include denaturing double stranded nucleic acid molecules. Another method for releasing the extended capture probes is to use a solution that interferes with the hydrogen bonds of the double stranded molecules. In some embodiments, the extended capture probe is released by applying heated water such as water or buffer of at least 85° C., e.g., at least 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99° C. In some embodiments, a solution including salts, surfactants, etc. that can further destabilize the interaction between the nucleic acid molecules is added to release the extended capture probe from the array feature. In some embodiments, a formamide solution can be used to destabilize the interaction between nucleic acid molecules to release the extended capture probe from the array feature.

(viii) Amplification of Capture Probes

[0379] In some embodiments, methods are provided herein for amplifying a capture probe affixed

to a spatial array, where amplification of the capture probe increases the number of capture domains and spatial barcodes on the spatial array. In some embodiments where a capture probe is amplified, the amplification is performed by rolling circle amplification. In some embodiments, the capture probe to be amplified includes sequences (e.g., docking sequences, functional sequences, and/or primer sequences) that enable rolling circle amplification. In one example, the capture probe can include a functional sequence that is capable of binding to a primer used for amplification. In another example, the capture probe can include one or more docking sequences (e.g., a first docking sequence and a second docking sequence) that can hybridize to one or more oligonucleotides (e.g., a padlock probe(s)) used for rolling circle amplification. In some embodiments, additional probes are affixed to the substrate, where the additional probes include sequences (e.g., a docking sequence(s), a functional sequence(s), and/or a primer sequence(s)) that enable rolling circle amplification. In some embodiments, the spatial array is contacted with an oligonucleotide (e.g., a padlock probe). As used herein, a “padlock probe” refers to an oligonucleotide that has, at its 5′ and 3′ ends, sequences that are complementary to adjacent or nearby target sequences (e.g., docking sequences) on a capture probe. Upon hybridization to the target sequences (e.g., docking sequences), the two ends of the padlock probe are either brought into contact or an end is extended until the two ends are brought into contact, allowing circularization of the padlock probe by ligation (e.g., ligation using any of the methods described herein). In some embodiments, after circularization of the oligonucleotide, rolling circle amplification can be used to amplify the ligation product, which includes at least a capture domain and a spatial barcode from the capture probe. In some embodiments, amplification of the capture probe using a padlock oligonucleotide and rolling circle amplification increases the number of capture domains and the number of spatial barcodes on the spatial array.

[0380] In some embodiments, a method of increasing capture efficiency of a spatial array includes amplifying all or part of a capture probe affixed to a substrate. For example, amplification of all or part of the capture probes affixed to the substrate can increase the capture efficiency of the spatial array by increasing the number of capture domains and spatial barcodes. In some embodiments, a method of determining a location of an analyte in a biological sample includes using a spatial array having increased capture efficiency (e.g., a spatial array where a capture probe has been amplified as described herein). For example, the capture efficiency of a spatial array can be increased by amplification of all or part of the capture probe prior to contact with a biological sample. The amplification results in an increased number of capture domains that enable capture of more analytes as compared to a spatial array where the capture probe was not amplified prior to contacting the biological sample. In some embodiments, a method of producing a spatial array that has increased capture efficiency includes amplifying all or part of a capture probe. In some embodiments where a spatial array having increased capture efficiency is produced by amplifying all or part of a capture probe, the amplification increases the number of capture domains and the number of spatial barcodes on the spatial array. In some embodiments, a method of determining the location of a capture probe (i.e., a capture probe on a feature) on a spatial array includes amplifying all or part of a capture probe. For example, amplification of the capture probe affixed to the substrate can increase the number of spatial barcodes used for direct decoding (e.g., direct decoding using any of the methods described herein including, without limitation, in situ sequencing) of the location of the capture probe.

(ix) Analyte Capture Agents

[0381] This disclosure also provides methods and materials for using analyte capture agents for spatial profiling of biological analytes (e.g., mRNA, genomic DNA, accessible chromatin, and cell surface or intracellular proteins and/or metabolites). As used herein, an “analyte capture agent” (also referred to previously at times as a “cell labelling” agent”) refers to an agent that interacts with an analyte (e.g., an analyte in a sample) and with a capture probe (e.g., a capture probe attached to a substrate) to identify the analyte. In some embodiments, the analyte capture agent

includes an analyte binding moiety and a capture agent barcode domain.

[0382] FIG. 9 is a schematic diagram of an exemplary analyte capture agent **902** comprised of an analyte binding moiety **904** and a capture agent barcode domain **908**. An analyte binding moiety **904** is a molecule capable of binding to an analyte **906** and interacting with a spatially-barcoded capture probe. The analyte binding moiety can bind to the analyte **906** with high affinity and/or with high specificity. The analyte capture agent can include a capture agent barcode domain **908**, a nucleotide sequence (e.g., an oligonucleotide), which can hybridize to at least a portion or an entirety of a capture domain of a capture probe. The analyte binding moiety **904** can include a polypeptide and/or an aptamer (e.g., an oligonucleotide or peptide molecule that binds to a specific target analyte). The analyte binding moiety **904** can include an antibody or antibody fragment (e.g., an antigen-binding fragment).

[0383] As used herein, the term “analyte binding moiety” refers to a molecule or moiety capable of binding to a macromolecular constituent (e.g., an analyte, e.g., a biological analyte). In some embodiments of any of the spatial profiling methods described herein, the analyte binding moiety of the analyte capture agent that binds to a biological analyte can include, but is not limited to, an antibody, or an epitope binding fragment thereof, a cell surface receptor binding molecule, a receptor ligand, a small molecule, a bi-specific antibody, a bi-specific T-cell engager, a T-cell receptor engager, a B-cell receptor engager, a pro-body, an aptamer, a monobody, an affimer, a darpin, and a protein scaffold, or any combination thereof. The analyte binding moiety can bind to the macromolecular constituent (e.g., analyte) with high affinity and/or with high specificity. The analyte binding moiety can include a nucleotide sequence (e.g., an oligonucleotide), which can correspond to at least a portion or an entirety of the analyte binding moiety. The analyte binding moiety can include a polypeptide and/or an aptamer (e.g., a polypeptide and/or an aptamer that binds to a specific target molecule, e.g., an analyte). The analyte binding moiety can include an antibody or antibody fragment (e.g., an antigen-binding fragment) that binds to a specific analyte (e.g., a polypeptide).

[0384] In some embodiments, an analyte binding moiety of an analyte capture agent includes one or more antibodies or antigen binding fragments thereof. The antibodies or antigen binding fragments including the analyte binding moiety can specifically bind to a target analyte. In some embodiments, the analyte is a protein (e.g., a protein on a surface of the biological sample (e.g., a cell) or an intracellular protein). In some embodiments, a plurality of analyte capture agents comprising a plurality of analyte binding moieties bind a plurality of analytes present in a biological sample. In some embodiments, the plurality of analytes includes a single species of analyte (e.g., a single species of polypeptide). In some embodiments in which the plurality of analytes includes a single species of analyte, the analyte binding moieties of the plurality of analyte capture agents are the same. In some embodiments in which the plurality of analytes includes a single species of analyte, the analyte binding moieties of the plurality of analyte capture agents are the different (e.g., members of the plurality of analyte capture agents can have two or more species of analyte binding moieties, wherein each of the two or more species of analyte binding moieties binds a single species of analyte, e.g., at different binding sites). In some embodiments, the plurality of analytes includes multiple different species of analyte (e.g., multiple different species of polypeptides).

[0385] An analyte capture agent can include an analyte binding moiety. The analyte binding moiety can be an antibody. Exemplary, non-limiting antibodies that can be used as analyte binding moieties in an analyte capture agent or that can be used in the IHC/IF applications disclosed herein include any of the following including variations thereof: A-ACT, A-AT, ACTH, Actin-Muscle-specific, Actin-Smooth Muscle (SMA), AE1, AE1/AE3, AE3, AFP, AKT Phosphate, ALK-1, Amyloid A, Androgen Receptor, Annexin A1, B72.3, BCA-225, BCL-1 (Cyclin D1), BCL-1/CD20, BCL-2, BCL-2/BCL-6, BCL-6, Ber-EP4, Beta-amyloid, Beta-catenin, BG8 (Lewis Y), BOB-1, CA 19.9, CA 125, CAIX, Calcitonin, Caldesmon, Calponin, Calretinin, CAM 5.2, CAM 5.2/AE1,

CD1a, CD2, CD3 (M), CD3 (P), CD3/CD20, CD4, CD5, CD7, CD8, CD10, CD14, CD15, CD20, CD21, CD22, CD 23, CD25, CD30, CD31, CD33, CD34, CD35, CD43, CD45 (LCA), CD45RA, CD56, CD57, CD61, CD68, CD71, CD74, CD79a, CD99, CD117 (c-KIT), CD123, CD138, CD163, CDX-2, CDX-2/CK-7, CEA (M), CEA (P), Chromogranin A, Chymotrypsin, CK-5, CK-5/6, CK-7, CK-7/TTF-1, CK-14, CK-17, CK-18, CK-19, CK-20, CK-HMW, CK-LMW, CMV-IH, COLL-IV, COX-2, D2-40, DBA44, Desmin, DOG1, EBER-ISH, EBV (LMP1), E-Cadherin, EGFR, EMA, ER, ERCC1, Factor VIII (vWF), Factor XIIIa, Fascin, FLI-1, FHS, Galectin-3, Gastrin, GCDFP-15, GFAP, Glucagon, Glycophorin A, Glypican-3, Granzyme B, Growth Hormone (GH), GST, HAM 56, HMBE-1, HBP, HCAG, HCG, Hemoglobin A, HEP B CORE (HBcAg), HEP B SURF, (HBsAg), HepPar1, HER2, Herpes I, Herpes II, HHV-8, HLA-DR, HMB 45, HPL, HPV-IHC, HPV (6/11)-ISH, HPV (16/18)-ISH, HPV (31/33)-ISH, HPV WSS-ISH, HPV High-ISH, HPV Low-ISH, HPV High & Low-ISH, IgA, IgD, IgG, IgG4, IgM, Inhibin, Insulin, JC Virus-ISH, Kappa-ISH, KER PAN, Ki-67, Lambda-IHC, Lambda-ISH, LH, Lipase, Lysozyme (MURA), Mammaglobin, MART-1, MBP, M-Cell Tryptase, MEL-5, Melan-A, Melan-A/Ki-67, Mesothelin, MiTF, MLH-1, MOC-31, MPO, MSH-2, MSH-6, MUC1, MUC2, MUC4, MUC5AC, MUM-1, MYO D1, Myogenin, Myoglobin, Myoin Heavy Chain, Napsin A, NB84a, NEW-N, NF, NK1-C.sub.3, NPM, NSE, OCT-2, OCT-3/4, OSCAR, p16, p21, p27/Kip1, p53, p57, p63, p120, P504S, Pan Melanoma, PANC.POLY, Parvovirus B19, PAX-2, PAX-5, PAX-5/CD43, PAX=5/CD5, PAX-8, PC, PD1, Perforin, PGP 9.5, PLAP, PMS-2, PR, Prolactin, PSA, PSAP, PSMA, PTEN, PTH, PTS, RB, RCC, S6, S100, Serotonin, Somatostatin, Surfactant (SP-A), Synaptophysin, Synuclein, TAU, TCL-1, TCR beta, TdT, Thrombomodulin, Thyroglobulin, TIA-1, TOXO, TRAP, TriView™ breast, TriView™ prostate, Trypsin, TS, TSH, TTF-1, Tyrosinase, Ubiquitin, Uroplakin, VEGF, Villin, Vimentin (VIM), VIP, VZV, WT1 (M) N-Terminus, WT1 (P) C-Terminus, ZAP-70.

[0386] Further, exemplary, non-limiting antibodies that can be used as analyte binding moieties in an analyte capture agent or that can be used in the IHC/IF applications disclosed herein include any of the following antibodies (and variations thereof) to: cell surface proteins, intracellular proteins, kinases (e.g., AGC kinase family (e.g., AKT1, AKT2, PDK1, Protein Kinase C, ROCK1, ROCK2, SGK3), CAMK kinase family (e.g., AMPK1, AMPK2, CAMK, Chk1, Chk2, Zip), CK1 kinase family, TK kinase family (e.g., Ab12, AXL, CD167, CD246/ALK, c-Met, CSK, c-Src, EGFR, ErbB2 (HER2/neu), ErbB3, ErbB4, FAK, Fyn, LCK, Lyn, PKT7, Syk, Zap70), STE kinase family (e.g., ASK1, MAPK, MEK1, MEK2, MEK3 MEK4, MEK5, PAK1, PAK2, PAK4, PAK6), CMGC kinase family (e.g., Cdk2, Cdk4, Cdk5, Cdk6, Cdk7, Cdk9, Erk1, GSK3, Jnk/MAPK8, Jnk2/MAPK9, JNK3/MAPK10, p38/MAPK), and TKL kinase family (e.g., ALK1, ILK1, IRAK1, IRAK2, IRAK3, IRAK4, LIMK1, LIMK2, M3K11, RAF1, RIP1, RIP3, VEGFR1, VEGFR2, VEGFR3), Aurora A kinase, Aurora B kinase, IKK, Nemo-like kinase, PINK, PLK3, ULK2, WEE1, transcription factors (e.g., FOXP3, ATF3, BACH1, EGR, ELF3, FOXA1, FOXA2, FOXO1, GATA), growth factor receptors, tumor suppressors (e.g., anti-p53, anti-BLM, anti-Cdk2, anti-Chk2, anti-BRCA-1, anti-NBS1, anti-BRCA-2, anti-WRN, anti-PTEN, anti-WT1, anti-p38).

[0387] In some embodiments, analyte capture agents are capable of binding to analytes present inside a cell. In some embodiments, analyte capture agents are capable of binding to cell surface analytes that can include, without limitation, a receptor, an antigen, a surface protein, a transmembrane protein, a cluster of differentiation protein, a protein channel, a protein pump, a carrier protein, a phospholipid, a glycoprotein, a glycolipid, a cell-cell interaction protein complex, an antigen-presenting complex, a major histocompatibility complex, an engineered T-cell receptor, a T-cell receptor, a B-cell receptor, a chimeric antigen receptor, an extracellular matrix protein, a posttranslational modification (e.g., phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation) state of a cell surface protein, a gap junction, and an adherens junction. In some embodiments, the analyte capture agents are capable of binding to cell surface analytes that are post-translationally modified. In such embodiments, analyte capture agents can be specific for cell surface analytes based on a given state of posttranslational modification

(e.g., phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation), such that a cell surface analyte profile can include posttranslational modification information of one or more analytes.

[0388] In some embodiments, the analyte capture agent includes a capture agent barcode domain that is conjugated or otherwise attached to the analyte binding moiety. In some embodiments, the capture agent barcode domain is covalently-linked to the analyte binding moiety. In some embodiments, a capture agent barcode domain is a nucleic acid sequence. In some embodiments, a capture agent barcode domain includes an analyte binding moiety barcode and an analyte capture sequence.

[0389] As used herein, the term “analyte binding moiety barcode” refers to a barcode that is associated with or otherwise identifies the analyte binding moiety. In some embodiments, by identifying an analyte binding moiety and its associated analyte binding moiety barcode, the analyte to which the analyte binding moiety binds can also be identified. An analyte binding moiety barcode can be a nucleic acid sequence of a given length and/or sequence that is associated with the analyte binding moiety. An analyte binding moiety barcode can generally include any of the variety of aspects of barcodes described herein. For example, an analyte capture agent that is specific to one type of analyte can have coupled thereto a first capture agent barcode domain (e.g., that includes a first analyte binding moiety barcode), while an analyte capture agent that is specific to a different analyte can have a different capture agent barcode domain (e.g., that includes a second barcode analyte binding moiety barcode) coupled thereto. In some aspects, such a capture agent barcode domain can include an analyte binding moiety barcode that permits identification of the analyte binding moiety to which the capture agent barcode domain is coupled. The selection of the capture agent barcode domain can allow significant diversity in terms of sequence, while also being readily attachable to most analyte binding moieties (e.g., antibodies or aptamers) as well as being readily detected, (e.g., using sequencing or array technologies).

[0390] In some embodiments, the capture agent barcode domain of an analyte capture agent includes an analyte capture sequence. As used herein, the term “analyte capture sequence” refers to a region or moiety configured to hybridize to, bind to, couple to, or otherwise interact with a capture domain of a capture probe. In some embodiments, an analyte capture sequence includes a nucleic acid sequence that is complementary to or substantially complementary to the capture domain of a capture probe such that the analyte capture sequence hybridizes to the capture domain of the capture probe. In some embodiments, an analyte capture sequence comprises a poly(A) nucleic acid sequence that hybridizes to a capture domain that comprises a poly(T) nucleic acid sequence. In some embodiments, an analyte capture sequence comprises a poly(T) nucleic acid sequence that hybridizes to a capture domain that comprises a poly(A) nucleic acid sequence. In some embodiments, an analyte capture sequence comprises a non-homopolymeric nucleic acid sequence that hybridizes to a capture domain that comprises a non-homopolymeric nucleic acid sequence that is complementary (or substantially complementary) to the non-homopolymeric nucleic acid sequence of the analyte capture region.

[0391] In some embodiments of any of the spatial analysis methods described herein that employ an analyte capture agent, the capture agent barcode domain can be directly coupled to the analyte binding moiety, or they can be attached to a bead, molecular lattice, e.g., a linear, globular, cross-linked, or other polymer, or other framework that is attached or otherwise associated with the analyte binding moiety, which allows attachment of multiple capture agent barcode domains to a single analyte binding moiety. Attachment (coupling) of the capture agent barcode domains to the analyte binding moieties can be achieved through any of a variety of direct or indirect, covalent or non-covalent associations or attachments. For example, in the case of a capture agent barcode domain coupled to an analyte binding moiety that includes an antibody or antigen-binding fragment, such capture agent barcode domains can be covalently attached to a portion of the antibody or antigen-binding fragment using chemical conjugation techniques (e.g., Lightning-

Link® antibody labelling kits available from Innova Biosciences). In some embodiments, a capture agent barcode domain can be coupled to an antibody or antigen-binding fragment using non-covalent attachment mechanisms (e.g., using biotinylated antibodies and oligonucleotides or beads that include one or more biotinylated linker(s), coupled to oligonucleotides with an avidin or streptavidin linker.) Antibody and oligonucleotide biotinylation techniques can be used, and are described for example in Fang et al., *Nucleic Acids Res.* (2003), 31 (2): 708-715, the entire contents of which are incorporated by reference herein. Likewise, protein and peptide biotinylation techniques have been developed and can be used, and are described for example in U.S. Pat. No. 6,265,552, the entire contents of which are incorporated by reference herein. Furthermore, click reaction chemistry such as a methyltetrazine-PEG5-NHS ester reaction, a TCO-PEG4-NHS ester reaction, or the like, can be used to couple capture agent barcode domains to analyte binding moieties. The reactive moiety on the analyte binding moiety can also include amine for targeting aldehydes, amine for targeting maleimide (e.g., free thiols), azide for targeting click chemistry compounds (e.g., alkynes), biotin for targeting streptavidin, phosphates for targeting EDC, which in turn targets active ester (e.g., NH₂). The reactive moiety on the analyte binding moiety can be a chemical compound or group that binds to the reactive moiety on the analyte binding moiety. Exemplary strategies to conjugate the analyte binding moiety to the capture agent barcode domain include the use of commercial kits (e.g., Solulink, Thunder link), conjugation of mild reduction of hinge region and maleimide labelling, stain-promoted click chemistry reaction to labeled amides (e.g., copper-free), and conjugation of periodate oxidation of sugar chain and amine conjugation. In the cases where the analyte binding moiety is an antibody, the antibody can be modified prior to or contemporaneously with conjugation of the oligonucleotide. For example, the antibody can be glycosylated with a chemical substrate-permissive mutant of β -1,4-galactosyltransferase, GalT (Y289L) and azide-bearing uridine diphosphate-N-acetylgalactosamine analog uridine diphosphate-GalNAz. The modified antibody can be conjugated to an oligonucleotide with a dibenzocyclooctyne-PEG4-NHS group. In some embodiments, certain steps (e.g., COOH activation (e.g., EDC) and homobifunctional cross linkers) can be avoided to prevent the analyte binding moieties from conjugating to themselves. In some embodiments of any of the spatial profiling methods described herein, the analyte capture agent (e.g., analyte binding moiety coupled to an oligonucleotide) can be delivered into the cell, e.g., by transfection (e.g., using transfectamine, cationic polymers, calcium phosphate or electroporation), by transduction (e.g., using a bacteriophage or recombinant viral vector), by mechanical delivery (e.g., magnetic beads), by lipid (e.g., 1,2-Dioleoyl-sn-glycero-3-phosphocholine (DOPC)), or by transporter proteins. An analyte capture agent can be delivered into a cell using exosomes. For example, a first cell can be generated that releases exosomes comprising an analyte capture agent. An analyte capture agent can be attached to an exosome membrane. An analyte capture agent can be contained within the cytosol of an exosome. Released exosomes can be harvested and provided to a second cell, thereby delivering the analyte capture agent into the second cell. An analyte capture agent can be releasable from an exosome membrane before, during, or after delivery into a cell. In some embodiments, the cell is permeabilized to allow the analyte capture agent to couple with intracellular constituents (such as, without limitation, intracellular proteins, metabolites, and nuclear membrane proteins). Following intracellular delivery, analyte capture agents can be used to analyze intracellular constituents as described herein.

[0392] In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domain coupled to an analyte capture agent can include modifications that render it non-extendable by a polymerase. In some embodiments, when binding to a capture domain of a capture probe or nucleic acid in a sample for a primer extension reaction, the capture agent barcode domain can serve as a template, not a primer. When the capture agent barcode domain also includes a barcode (e.g., an analyte binding moiety barcode), such a design can increase the efficiency of molecular barcoding by increasing the affinity between the capture agent barcode domain and

unbarcoded sample nucleic acids, and eliminate the potential formation of adaptor artifacts. In some embodiments, the capture agent barcode domain can include a random N-mer sequence that is capped with modifications that render it non-extendable by a polymerase. In some cases, the composition of the random N-mer sequence can be designed to maximize the binding efficiency to free, unbarcoded ssDNA molecules. The design can include a random sequence composition with a higher GC content, a partial random sequence with fixed G or C at specific positions, the use of guanosines, the use of locked nucleic acids, or any combination thereof.

[0393] A modification for blocking primer extension by a polymerase can be a carbon spacer group of different lengths or a dideoxynucleotide. In some embodiments, the modification can be an abasic site that has an apurine or apyrimidine structure, a base analog, or an analogue of a phosphate backbone, such as a backbone of N-(2-aminoethyl)-glycine linked by amide bonds, tetrahydrofuran, or 1',2'-Dideoxyribose. The modification can also be a uracil base, 2'OMe modified RNA, C3-18 spacers (e.g., structures with 3-18 consecutive carbon atoms, such as C3 spacer), ethylene glycol multimer spacers (e.g., spacer 18 (hexa-ethyleneglycol spacer), biotin, dideoxynucleotide triphosphate, ethylene glycol, amine, or phosphate.

[0394] In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domain coupled to the analyte binding moiety includes a cleavable domain. For example, after the analyte capture agent binds to an analyte (e.g., a cell surface analyte), the capture agent barcode domain can be cleaved and collected for downstream analysis according to the methods as described herein. In some embodiments, the cleavable domain of the capture agent barcode domain includes a U-excising element that allows the species to release from the bead. In some embodiments, the U-excising element can include a single-stranded DNA (ssDNA) sequence that contains at least one uracil. The species can be attached to a bead via the ssDNA sequence. The species can be released by a combination of uracil-DNA glycosylase (e.g., to remove the uracil) and an endonuclease (e.g., to induce an ssDNA break). If the endonuclease generates a 5' phosphate group from the cleavage, then additional enzyme treatment can be included in downstream processing to eliminate the phosphate group, e.g., prior to ligation of additional sequencing handle elements, e.g., Illumina full P5 sequence, partial P5 sequence, full R1 sequence, and/or partial R1 sequence.

[0395] In some embodiments, multiple different species of analytes (e.g., polypeptides) from the biological sample can be subsequently associated with the one or more physical properties of the biological sample. For example, the multiple different species of analytes can be associated with locations of the analytes in the biological sample. Such information (e.g., proteomic information when the analyte binding moiety (ies) recognizes a polypeptide(s)) can be used in association with other spatial information (e.g., genetic information from the biological sample, such as DNA sequence information, transcriptome information (i.e., sequences of transcripts), or both). For example, a cell surface protein of a cell can be associated with one or more physical properties of the cell (e.g., a shape, size, activity, or a type of the cell). The one or more physical properties can be characterized by imaging the cell. The cell can be bound by an analyte capture agent comprising an analyte binding moiety that binds to the cell surface protein and an analyte binding moiety barcode that identifies that analyte binding moiety, and the cell can be subjected to spatial analysis (e.g., any of the variety of spatial analysis methods described herein). For example, the analyte capture agent bound to the cell surface protein can be bound to a capture probe (e.g., a capture probe on an array), which capture probe includes a capture domain that interacts with an analyte capture sequence present on the capture agent barcode domain of the analyte capture agent. All or part of the capture agent barcode domain (including the analyte binding moiety barcode) can be copied with a polymerase using a 3' end of the capture domain as a priming site, generating an extended capture probe that includes the all or part of complementary sequence that corresponds to the capture probe (including a spatial barcode present on the capture probe) and a copy of the analyte binding moiety barcode. In some embodiments, an analyte capture agent with an extended

capture agent barcode domain that includes a sequence complementary to a spatial barcode of a capture probe is called a “spatially-tagged analyte capture agent.”

[0396] In some embodiments, the spatial array with spatially-tagged analyte capture agents can be contacted with a sample, where the analyte capture agent(s) associated with the spatial array capture the target analyte(s). The analyte capture agent(s) containing the extended capture probe(s), which includes a sequence complementary to the spatial barcode(s) of the capture probe(s) and the analyte binding moiety barcode(s), can then be denatured from the capture probe(s) of the spatial array. This allows the spatial array to be reused. The sample can be dissociated into non-aggregated cells (e.g., single cells) and analyzed by the single cell/droplet methods described herein. The spatially-tagged analyte capture agent can be sequenced to obtain the nucleic acid sequence of the spatial barcode of the capture probe and the analyte binding moiety barcode of the analyte capture agent. The nucleic acid sequence of the extended capture probe can thus be associated with an analyte (e.g., cell surface protein), and in turn, with the one or more physical properties of the cell (e.g., a shape or cell type). In some embodiments, the nucleic acid sequence of the extended capture probe can be associated with an intracellular analyte of a nearby cell, where the intracellular analyte was released using any of the cell permeabilization or analyte migration techniques described herein.

[0397] In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domains released from the analyte capture agents can then be subjected to sequence analysis to identify which analyte capture agents were bound to analytes. Based upon the capture agent barcode domains that are associated with a feature (e.g., a feature at a particular location) on a spatial array and the presence of the analyte binding moiety barcode sequence, an analyte profile can be created for a biological sample. Profiles of individual cells or populations of cells can be compared to profiles from other cells, e.g., ‘normal’ cells, to identify variations in analytes, which can provide diagnostically relevant information. In some embodiments, these profiles can be useful in the diagnosis of a variety of disorders that are characterized by variations in cell surface receptors, such as cancer and other disorders.

[0398] FIG. **10** is a schematic diagram depicting an exemplary interaction between a feature-immobilized capture probe **1024** and an analyte capture agent **1026**. The feature-immobilized capture probe **1024** can include a spatial barcode **1008** as well as one or more functional sequences **1006** and **1010**, as described elsewhere herein. The capture probe can also include a capture domain **1012** that is capable of binding to an analyte capture agent **1026**. The analyte capture agent **1026** can include a functional sequence **1018**, capture agent barcode domain **1016**, and an analyte capture sequence **1014** that is capable of binding to the capture domain **1012** of the capture probe **1024**. The analyte capture agent can also include a linker **1020** that allows the capture agent barcode domain **1016** to couple to the analyte binding moiety **1022**.

[0399] In some embodiments of any of the spatial profiling methods described herein, the methods are used to identify immune cell profiles. Immune cells express various adaptive immunological receptors relating to immune function, such as T cell receptors (TCRs) and B cell receptors (BCRs). T cell receptors and B cell receptors play a part in the immune response by specifically recognizing and binding to antigens and aiding in their destruction.

[0400] The T cell receptor, or TCR, is a molecule found on the surface of T cells that is generally responsible for recognizing fragments of antigen as peptides bound to major histocompatibility complex (MHC) molecules. The TCR is generally a heterodimer of two chains, each of which is a member of the immunoglobulin superfamily, possessing an N-terminal variable (V) domain, and a C terminal constant domain. In humans, in 95% of T cells, the TCR consists of an alpha (α) and beta (β) chain, whereas in 5% of T cells, the TCR consists of gamma and delta (γ/δ) chains. This ratio can change during ontogeny and in diseased states as well as in different species. When the TCR engages with antigenic peptide and MHC (peptide/MHC or pMHC), the T lymphocyte is activated through signal transduction.

[0401] Each of the two chains of a TCR contains multiple copies of gene segments—a variable ‘V’ gene segment, a diversity ‘D’ gene segment, and a joining ‘J’ gene segment. The TCR alpha chain (TCRa) is generated by recombination of V and J segments, while the beta chain (TCRb) is generated by recombination of V, D, and J segments. Similarly, generation of the TCR gamma chain involves recombination of V and J gene segments, while generation of the TCR delta chain occurs by recombination of V, D, and J gene segments. The intersection of these specific regions (V and J for the alpha or gamma chain, or V, D and J for the beta or delta chain) corresponds to the CDR3 region that is important for antigen-MHC recognition. Complementarity determining regions (e.g., CDR1, CDR2, and CDR3), or hypervariable regions, are sequences in the variable domains of antigen receptors (e.g., T cell receptor and immunoglobulin) that can complement an antigen. Most of the diversity of CDRs is found in CDR3, with the diversity being generated by somatic recombination events during the development of T lymphocytes. A unique nucleotide sequence that arises during the gene arrangement process can be referred to as a clonotype.

[0402] The B cell receptor, or BCR, is a molecule found on the surface of B cells. The antigen binding portion of a BCR is composed of a membrane-bound antibody that, like most antibodies (e.g., immunoglobulins), has a unique and randomly determined antigen-binding site. The antigen binding portion of a BCR includes membrane-bound immunoglobulin molecule of one isotype (e.g., IgD, IgM, IgA, IgG, or IgE). When a B cell is activated by its first encounter with a cognate antigen, the cell proliferates and differentiates to generate a population of antibody-secreting plasma B cells and memory B cells. The various immunoglobulin isotypes differ in their biological features, structure, target specificity, and distribution. A variety of molecular mechanisms exist to generate initial diversity, including genetic recombination at multiple sites.

[0403] The BCR is composed of two genes IgH and IgK (or IgL) coding for antibody heavy and light chains. Immunoglobulins are formed by recombination among gene segments, sequence diversification at the junctions of these segments, and point mutations throughout the gene. Each heavy chain gene contains multiple copies of three different gene segments—a variable ‘V’ gene segment, a diversity ‘D’ gene segment, and a joining ‘J’ gene segment. Each light chain gene contains multiple copies of two different gene segments for the variable region of the protein—a variable ‘V’ gene segment and a joining ‘J’ gene segment.

[0404] The recombination can generate a molecule with one of each of the V, D, and J segments. Furthermore, several bases can be deleted and others added (called N and P nucleotides) at each of the two junctions, thereby generating further diversity. After B cell activation, a process of affinity maturation through somatic hypermutation occurs. In this process, progeny cells of the activated B cells accumulate distinct somatic mutations throughout the gene with higher mutation concentration in the CDR regions leading to the generation of antibodies with higher affinity to the antigens.

[0405] In addition to somatic hypermutation, activated B cells undergo the process of isotype switching. Antibodies with the same variable segments can have different forms (isotypes) depending on the constant segment. Whereas all naïve B cells express IgM (or IgD), activated B cells mostly express IgG but also IgM, IgA, and IgE. This expression switching from IgM (and/or IgD) to IgG, IgA, or IgE occurs through a recombination event causing one cell to specialize in producing a specific isotype. A unique nucleotide sequence that arises during the gene arrangement process can similarly be referred to as a clonotype.

[0406] Certain methods described herein are utilized to analyze the various sequences of TCRs and BCRs from immune cells, for example, various clonotypes. In some embodiments, the methods are used to analyze the sequence of a TCR alpha chain, a TCR beta chain, a TCR delta chain, a TCR gamma chain, or any fragment thereof (e.g., variable regions including V(D)J or VJ regions, constant regions, transmembrane regions, fragments thereof, combinations thereof, and combinations of fragments thereof). In some embodiments, the methods described herein can be used to analyze the sequence of a B cell receptor heavy chain, B cell receptor light chain, or any

fragment thereof (e.g., variable regions including V(D)J or VJ regions, constant regions, transmembrane regions, fragments thereof, combinations thereof, and combinations of fragments thereof).

[0407] Where immune cells are to be analyzed, primer sequences useful in any of the various operations for attaching barcode sequences and/or amplification reactions can include gene specific sequences which target genes or regions of genes of immune cell proteins, for example immune receptors. Such gene sequences include, but are not limited to, sequences of various T cell receptor alpha variable genes (TRAV genes), T cell receptor alpha joining genes (TRAJ genes), T cell receptor alpha constant genes (TRAC genes), T cell receptor beta variable genes (TRBV genes), T cell receptor beta diversity genes (TRBD genes), T cell receptor beta joining genes (TRBJ genes), T cell receptor beta constant genes (TRBC genes), T cell receptor gamma variable genes (TRGV genes), T cell receptor gamma joining genes (TRGJ genes), T cell receptor gamma constant genes (TRGC genes), T cell receptor delta variable genes (TRDV genes), T cell receptor delta diversity genes (TRDD genes), T cell receptor delta joining genes (TRDJ genes), and T cell receptor delta constant genes (TRDC genes).

[0408] In some embodiments, the analyte binding moiety is based on the Major Histocompatibility Complex (MHC) class I or class II. In some embodiments, the analyte binding moiety is an MHC multimer including, without limitation, MHC dextramers, MHC tetramers, and MHC pentamers (see, for example, U.S. Patent Application Publication Nos. US 2018/0180601 and US 2017/0343545, the entire contents of each of which are incorporated herein by reference. MHCs (e.g., a soluble MHC monomer molecule), including full or partial MHC-peptides, can be used as analyte binding moieties of analyte capture agents that are coupled to capture agent barcode domains that include an analyte binding moiety barcode that identifies its associated MHC (and, thus, for example, the MHC's TCR binding partner). In some embodiments, MHCs are used to analyze one or more cell-surface features of a T-cell, such as a TCR. In some cases, multiple MHCs are associated together in a larger complex (MHC multi-mer) to improve binding affinity of MHCs to TCRs via multiple ligand binding synergies.

[0409] FIGS. **11A**, **11B**, and **11C** are schematics illustrating how streptavidin cell tags can be utilized in an array-based system to produce a spatially-barcoded cell or cellular contents. For example, as shown in FIG. **11A**, peptide-bound major histocompatibility complex (MHC) can be individually associated with biotin (β 2m) and bound to a streptavidin moiety such that the streptavidin moiety comprises multiple pMHC moieties. Each of these moieties can bind to a TCR such that the streptavidin binds to a target T-cell via multiple MCH/TCR binding interactions. Multiple interactions synergize and can substantially improve binding affinity. Such improved affinity can improve labelling of T-cells and also reduce the likelihood that labels will dissociate from T-cell surfaces. As shown in FIG. **11B**, a capture agent barcode domain **1101** can be modified with streptavidin **1102** and contacted with multiple molecules of biotinylated MHC **1103** such that the biotinylated MHC **1103** molecules are coupled with the streptavidin conjugated capture agent barcode domain **1101**. The result is a barcoded MHC multimer complex **1105**. As shown in FIG. **11B**, the capture agent barcode domain sequence **1101** can identify the MHC as its associated label and also includes optional functional sequences such as sequences for hybridization with other oligonucleotides. As shown in FIG. **11C**, one example oligonucleotide is capture probe **1106** that comprises a complementary sequence (e.g., rGrGrG corresponding to C C C), a barcode sequence and other functional sequences, such as, for example, a UMI, an adapter sequence (e.g., comprising a sequencing primer sequence (e.g., R1 or a partial R1 (“pR1”), R2), a flow cell attachment sequence (e.g., P5 or P7 or partial sequences thereof)), etc. In some cases, capture probe **1106** may at first be associated with a feature (e.g., a gel bead) and released from the feature. In other embodiments, capture probe **1106** can hybridize with a capture agent barcode domain **1101** of the MHC-oligonucleotide complex **1105**. The hybridized oligonucleotides (Spacer C C C and Spacer rGrGrG) can then be extended in primer extension reactions such that constructs comprising

sequences that correspond to each of the two spatial barcode sequences (the spatial barcode associated with the capture probe, and the barcode associated with the MHC-oligonucleotide complex) are generated. In some cases, one or both of these corresponding sequences may be a complement of the original sequence in capture probe **1106** or capture agent barcode domain **1101**. In other embodiments, the capture probe and the capture agent barcode domain are ligated together. The resulting constructs can be optionally further processed (e.g., to add any additional sequences and/or for clean-up) and subjected to sequencing. As described elsewhere herein, a sequence derived from the capture probe **1106** spatial barcode sequence may be used to identify a feature and the sequence derived from spatial barcode sequence on the capture agent barcode domain **1101** may be used to identify the particular peptide MHC complex **1104** bound on the surface of the cell (e.g., when using MHC-peptide libraries for screening immune cells or immune cell populations).

(c) Substrates

[0410] For the spatial array-based analytical methods described herein, a substrate functions as a support for direct or indirect attachment of capture probes to features of the array. In addition, in some embodiments, a substrate (e.g., the same substrate or a different substrate) can be used to provide support to a biological sample, particularly, for example, a thin tissue section. Accordingly, a “substrate” is a support that is insoluble in aqueous liquid and which allows for positioning of biological samples, analytes, features, and/or capture probes on the substrate.

[0411] Further, a “substrate” as used herein, and when not preceded by the modifier “chemical”, refers to a member with at least one surface that generally functions to provide physical support for biological samples, analytes, and/or any of the other chemical and/or physical moieties, agents, and structures described herein. Substrates can be formed from a variety of solid materials, gel-based materials, colloidal materials, semi-solid materials (e.g., materials that are at least partially cross-linked), materials that are fully or partially cured, and materials that undergo a phase change or transition to provide physical support. Examples of substrates that can be used in the methods and systems described herein include, but are not limited to, slides (e.g., slides formed from various glasses, slides formed from various polymers), hydrogels, layers and/or films, membranes (e.g., porous membranes), flow cells, cuvettes, wafers, plates, or combinations thereof. In some embodiments, substrates can optionally include functional elements such as recesses, protruding structures, microfluidic elements (e.g., channels, reservoirs, electrodes, valves, seals), and various markings, as will be discussed in further detail below.

(i) Substrate Attributes

[0412] A substrate can generally have any suitable form or format. For example, a substrate can be flat, curved, e.g., convexly or concavely curved towards the area where the interaction between a biological sample, e.g., tissue sample, and a substrate takes place. In some embodiments, a substrate is flat, e.g., planar, chip, or slide. A substrate can contain one or more patterned surfaces within the substrate (e.g., channels, wells, projections, ridges, divots, etc.).

[0413] A substrate can be of any desired shape. For example, a substrate can be typically a thin, flat shape (e.g., a square or a rectangle). In some embodiments, a substrate structure has rounded corners (e.g., for increased safety or robustness). In some embodiments, a substrate structure has one or more cut-off corners (e.g., for use with a slide clamp or cross-table). In some embodiments, where a substrate structure is flat, the substrate structure can be any appropriate type of support having a flat surface (e.g., a chip or a slide such as a microscope slide).

[0414] Substrates can optionally include various structures such as, but not limited to, projections, ridges, and channels. A substrate can be micropatterned to limit lateral diffusion (e.g., to prevent overlap of spatial barcodes). A substrate modified with such structures can be modified to allow association of analytes, features (e.g., beads), or probes at individual sites. For example, the sites where a substrate is modified with various structures can be contiguous or non-contiguous with other sites.

[0415] In some embodiments, the surface of a substrate can be modified so that discrete sites are

formed that can only have or accommodate a single feature. In some embodiments, the surface of a substrate can be modified so that features adhere to random sites.

[0416] In some embodiments, the surface of a substrate is modified to contain one or more wells, using techniques such as (but not limited to) stamping, microetching, or molding techniques. In some embodiments in which a substrate includes one or more wells, the substrate can be a concavity slide or cavity slide. For example, wells can be formed by one or more shallow depressions on the surface of the substrate. In some embodiments, where a substrate includes one or more wells, the wells can be formed by attaching a cassette (e.g., a cassette containing one or more chambers) to a surface of the substrate structure.

[0417] In some embodiments, the structures of a substrate (e.g., wells or features) can each bear a different capture probe. Different capture probes attached to each structure can be identified according to the locations of the structures in or on the surface of the substrate. Exemplary substrates include arrays in which separate structures are located on the substrate including, for example, those having wells that accommodate features.

[0418] In some embodiments where the substrate is modified to contain one or more structures, including but not limited to, wells, projections, ridges, features, or markings, the structures can include physically altered sites. For example, a substrate modified with various structures can include physical properties, including, but not limited to, physical configurations, magnetic or compressive forces, chemically functionalized sites, chemically altered sites, and/or electrostatically altered sites. In some embodiments where the substrate is modified to contain various structures, including but not limited to wells, projections, ridges, features, or markings, the structures are applied in a pattern. Alternatively, the structures can be randomly distributed.

[0419] The substrate (e.g., or a bead or a feature on an array) can include tens to hundreds of thousands or millions of individual oligonucleotide molecules (e.g., at least about 10,000, 50,000, 100,000, 500,000, 1,000,000, 10,000,000, 100,000,000, 1,000,000,000, or 10,000,000,000 oligonucleotide molecules).

[0420] In some embodiments, a substrate includes one or more markings on a surface of a substrate, e.g., to provide guidance for correlating spatial information with the characterization of the analyte of interest. For example, a substrate can be marked with a grid of lines (e.g., to allow the size of objects seen under magnification to be easily estimated and/or to provide reference areas for counting objects). In some embodiments, fiducial markers can be included on a substrate. Such markings can be made using techniques including, but not limited to, printing, sand-blasting, and depositing on the surface.

[0421] In some embodiments, imaging can be performed using one or more fiducial markers, i.e., objects placed in the field of view of an imaging system which appear in the image produced. Fiducial markers are typically used as a point of reference or measurement scale. Fiducial markers can include, but are not limited to, detectable labels such as fluorescent, radioactive, chemiluminescent, and colorimetric labels. The use of fiducial markers to stabilize and orient biological samples is described, for example, in Carter et al., *Applied Optics* 46:421-427, 2007), the entire contents of which are incorporated herein by reference. In some embodiments, a fiducial marker can be a physical particle (e.g., a nanoparticle, a microsphere, a nanosphere, a bead, a post, or any of the other exemplary physical particles described herein or known in the art).

[0422] In some embodiments, a fiducial marker can be present on a substrate to provide orientation of the biological sample. In some embodiments, a microsphere can be coupled to a substrate to aid in orientation of the biological sample. In some examples, a microsphere coupled to a substrate can produce an optical signal (e.g., fluorescence). In another example, a microsphere can be attached to a portion (e.g., corner) of an array in a specific pattern or design (e.g., hexagonal design) to aid in orientation of a biological sample on an array of features on the substrate. In some embodiments, a quantum dot can be coupled to the substrate to aid in the orientation of the biological sample. In some examples, a quantum dot coupled to a substrate can produce an optical signal.

[0423] In some embodiments, a fiducial marker can be an immobilized molecule with which a detectable signal molecule can interact to generate a signal. For example, a marker nucleic acid can be linked or coupled to a chemical moiety capable of fluorescing when subjected to light of a specific wavelength (or range of wavelengths). Such a marker nucleic acid molecule can be contacted with an array before, contemporaneously with, or after the tissue sample is stained to visualize or image the tissue section. Although not required, it can be advantageous to use a marker that can be detected using the same conditions (e.g., imaging conditions) used to detect a labelled cDNA.

[0424] In some embodiments, fiducial markers are included to facilitate the orientation of a tissue sample or an image thereof in relation to an immobilized capture probes on a substrate. Any number of methods for marking an array can be used such that a marker is detectable only when a tissue section is imaged. For instance, a molecule, e.g., a fluorescent molecule that generates a signal, can be immobilized directly or indirectly on the surface of a substrate. Markers can be provided on a substrate in a pattern (e.g., an edge, one or more rows, one or more lines, etc.).

[0425] In some embodiments, a fiducial marker can be randomly placed in the field of view. For example, an oligonucleotide containing a fluorophore can be randomly printed, stamped, synthesized, or attached to a substrate (e.g., a glass slide) at a random position on the substrate. A tissue section can be contacted with the substrate such that the oligonucleotide containing the fluorophore contacts, or is in proximity to, a cell from the tissue section or a component of the cell (e.g., an mRNA or DNA molecule). An image of the substrate and the tissue section can be obtained, and the position of the fluorophore within the tissue section image can be determined (e.g., by reviewing an optical image of the tissue section overlaid with the fluorophore detection). In some embodiments, fiducial markers can be precisely placed in the field of view (e.g., at known locations on a substrate). In this instance, a fiducial marker can be stamped, attached, or synthesized on the substrate and contacted with a biological sample. Typically, an image of the sample and the fiducial marker is taken, and the position of the fiducial marker on the substrate can be confirmed by viewing the image.

[0426] In some embodiments, a fiducial marker can be an immobilized molecule (e.g., a physical particle) attached to the substrate. For example, a fiducial marker can be a nanoparticle, e.g., a nanorod, a nanowire, a nanocube, a nanopyramid, or a spherical nanoparticle. In some examples, the nanoparticle can be made of a heavy metal (e.g., gold). In some embodiments, the nanoparticle can be made from diamond. In some embodiments, the fiducial marker can be visible by eye.

[0427] As noted herein, any of the fiducial markers described herein (e.g., microspheres, beads, or any of the other physical particles described herein) can be located at a portion (e.g., corner) of an array in a specific pattern or design (e.g., hexagonal design) to aid in orientation of a biological sample on an array of features on the substrate. In some embodiments, the fiducial markers located at a portion (e.g., corner) of an array (e.g., an array on a substrate) can be patterned or designed in at least 1, at least 2, at least 3, or at least 4 unique patterns. In some examples, the fiducial markers located at the corners of the array (e.g., an array on a substrate) can have four unique patterns of fiducial markers.

[0428] In some examples, fiducial markers can surround the array. In some embodiments the fiducial markers allow for detection of, e.g., mirroring. In some embodiments, the fiducial markers may completely surround the array. In some embodiments, the fiducial markers may not completely surround the array. In some embodiments, the fiducial markers identify the corners of the array. In some embodiments, one or more fiducial markers identify the center of the array. In some embodiments, the fiducial markers comprise patterned spots, wherein the diameter of one or more patterned spot fiducial markers is approximately 100 micrometers. The diameter of the fiducial markers can be any useful diameter including, but not limited to, 50 micrometers to 500 micrometers in diameter. The fiducial markers may be arranged in such a way that the center of one fiducial marker is between 100 micrometers and 200 micrometers from the center of one or more

other fiducial markers surrounding the array. In some embodiments, the array with the surrounding fiducial markers is approximately 8 mm by 8 mm. In some embodiments, the array without the surrounding fiducial markers is smaller than 8 mm by 50 mm.

[0429] In some embodiments, an array can be enclosed within a frame. Put another way, the perimeter of an array can have fiducial markers such that the array is enclosed, or substantially enclosed. In some embodiments, the perimeter of an array can be fiducial markers (e.g., any fiducial marker described herein). In some embodiments, the perimeter of an array can be uniform. For example, the fiducial markings can connect, or substantially connect, consecutive corners of an array in such a fashion that the non-corner portion of the array perimeter is the same on all sides (e.g., four sides) of the array. In some embodiments, the fiducial markers attached to the non-corner portions of the perimeter can be patterned or designed to aid in the orientation of the biological sample on the array. In some embodiments, the particles attached to the non-corner portions of the perimeter can be patterned or designed in at least 1, at least 2, at least 3, or at least 4 patterns. In some embodiments, the patterns can have at least 2, at least 3, or at least 4 unique patterns of fiducial markings on the non-corner portion of the array perimeter.

[0430] In some embodiments, an array can include at least two fiducial markers (e.g., at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 12, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100 fiducial markers or more (e.g., several hundred, several thousand, or tens of thousands of fiducial markers)) in distinct positions on the surface of a substrate. Fiducial markers can be provided on a substrate in a pattern (e.g., an edge, one or more rows, one or more lines, etc.).

[0431] A wide variety of different substrates can be used for the foregoing purposes. In general, a substrate can be any suitable support material. Exemplary substrates include, but are not limited to, glass, modified and/or functionalized glass, hydrogels, films, membranes, plastics (including e.g., acrylics, polystyrene, copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon™, cyclic olefins, polyimides etc.), nylon, ceramics, resins, Zeonor, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, optical fiber bundles, and polymers, such as polystyrene, cyclic olefin copolymers (COCs), cyclic olefin polymers (COPs), polypropylene, polyethylene polycarbonate, or combinations thereof.

[0432] Among the examples of substrate materials discussed above, polystyrene is a hydrophobic material suitable for binding negatively charged macromolecules because it normally contains few hydrophilic groups. For nucleic acids immobilized on glass slides, by increasing the hydrophobicity of the glass surface the nucleic acid immobilization can be increased. Such an enhancement can permit a relatively more densely packed formation (e.g., provide improved specificity and resolution).

[0433] In another example, a substrate can be a flow cell. Flow cells can be formed of any of the foregoing materials, and can include channels that permit reagents, solvents, features, and analytes to pass through the flow cell. In some embodiments, a hydrogel embedded biological sample is assembled in a flow cell (e.g., the flow cell is utilized to introduce the hydrogel to the biological sample). In some embodiments, a hydrogel embedded biological sample is not assembled in a flow cell. In some embodiments, the hydrogel embedded biological sample can then be prepared and/or isometrically expanded as described herein.

(ii) Conductive Substrates

[0434] Conductive substrates (e.g., electrophoretic compatible arrays) generated as described herein can be used in the spatial detection of analytes. For example, an electrophoretic field can be applied to facilitate migration of analytes towards the barcoded oligonucleotides (e.g., capture probes) on the array (e.g., capture probes immobilized on paper, capture probes immobilized in a hydrogel film, or capture probes immobilized on a glass slide having a conductive coating). In some embodiments, an electrophoresis assembly can be arranged. For example, an anode and a

cathode can be arranged such that an array of capture probes (e.g., capture probes immobilized on paper, capture probes immobilized in a hydrogel film, or capture probes immobilized on a glass slide having a conductive coating) and a biological sample are positioned between the anode and the cathode. In such embodiments, analytes in the biological sample are actively migrated toward the capture probes on the conductive substrate and captured. The biological sample can be prepared (e.g., permeabilized) according to any method described herein. In some embodiments, after electrophoretic-assisted capture of the analytes, the barcoded oligonucleotides (e.g., capture probes) and captured analytes can be collected, processed, and/or analyzed (e.g., sequenced) using any of the methods described herein.

[0435] In some embodiments, a conductive substrate can include glass (e.g., a glass slide) that has been coated with a substance or otherwise modified to confer conductive properties to the glass. In some embodiments, a glass slide can be coated with a conductive coating. In some embodiments, a conductive coating includes tin oxide (TO) or indium tin oxide (ITO). In some embodiments, a conductive coating includes a transparent conductive oxide (TCO). In some embodiments, a conductive coating includes aluminum doped zinc oxide (AZO). In some embodiments, a conductive coating includes fluorine doped tin oxide (FTO).

[0436] In some embodiments, arrays that are spotted or printed with oligonucleotides (e.g., capture probes, e.g., any of the variety of capture probes described herein) can be generated on a conductive substrate (e.g., any of the conductive substrates described herein). For example, the arrays described herein can be compatible with active analyte capture methods (e.g., any of the analyte capture methods described herein, including without limitation, electrophoretic capture methods). In some embodiments, a conductive substrate is a porous medium. Non-limiting examples of porous media that can be used in methods described herein that employ active analyte capture include a nitrocellulose or nylon membrane. In some embodiments, a porous medium that can be used in methods described herein that employ active analyte capture includes paper. In some embodiments, the oligonucleotides can be printed on a paper substrate. In some embodiments, the printed oligonucleotides can interact with the substrate (e.g., interact with fibers of the paper). In some embodiments, printed oligonucleotides can covalently bind the substrate (e.g., to fibers of the paper). In some embodiments, oligonucleotides in a molecular precursor solution can be printed on a conductive substrate (e.g., paper). In some embodiments, a molecular precursor solution can polymerize, thereby generating gel pads on the conductive substrate (e.g., paper). In some embodiments, a molecular precursor solution can be polymerized by light (e.g., photocured). In some embodiments, gel beads (e.g., any of the variety of gel beads described herein) containing oligonucleotides (e.g., barcoded oligonucleotides such as capture probes) can be printed on a conductive substrate (e.g., paper). In some embodiments, the printed oligonucleotides can be covalently attached into the gel matrix.

(iii) Coatings

[0437] In some embodiments, a surface of a substrate can be coated with a cell-permissive coating to allow adherence of live cells. A “cell-permissive coating” is a coating that allows or helps cells to maintain cell viability (e.g., remain viable) on the substrate. For example, a cell-permissive coating can enhance cell attachment, cell growth, and/or cell differentiation, e.g., a cell-permissive coating can provide nutrients to the live cells. A cell-permissive coating can include a biological material and/or a synthetic material. Non-limiting examples of a cell-permissive coating include coatings that feature one or more extracellular matrix (ECM) components (e.g., proteoglycans and fibrous proteins such as collagen, elastin, fibronectin and laminin), poly-lysine, poly(L)-ornithine, and/or a biocompatible silicone (e.g., CYTOSOFT®). For example, a cell-permissive coating that includes one or more extracellular matrix components can include collagen Type I, collagen Type II, collagen Type IV, elastin, fibronectin, laminin, and/or vitronectin. In some embodiments, the cell-permissive coating includes a solubilized basement membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma (e.g., MATRIGEL®). In some embodiments, the

cell-permissive coating includes collagen. A cell-permissive coating can be used to culture adherent cells on a spatially-barcoded array, or to maintain cell viability of a tissue sample or section while in contact with a spatially-barcoded array.

[0438] In some embodiments, a substrate is coated with a surface treatment such as poly(L)-lysine. Additionally or alternatively, the substrate can be treated by silanation, e.g., with epoxy-silane, amino-silane, and/or by a treatment with polyacrylamide.

[0439] In some embodiments, a substrate is treated in order to minimize or reduce non-specific analyte hybridization within or between features. For example, treatment can include coating the substrate with a hydrogel, film, and/or membrane that creates a physical barrier to non-specific hybridization. Any suitable hydrogel can be used. For example, hydrogel matrices prepared according to the methods set forth in U.S. Pat. Nos. 6,391,937, 9,512,422, and 9,889,422, and U.S. Patent Application Publication Nos. U.S. 2017/0253918 and U.S. 2018/0052081, can be used. The entire contents of each of the foregoing documents is incorporated herein by reference.

[0440] Treatment can include adding a functional group that is reactive or capable of being activated such that it becomes reactive after application of a stimulus (e.g., photoreactive functional groups). Treatment can include treating with polymers having one or more physical properties (e.g., mechanical, electrical, magnetic, and/or thermal) that minimize non-specific binding (e.g., that activate a substrate at certain locations to allow analyte hybridization at those locations).

[0441] A “conditionally removable coating” is a coating that can be removed from the surface of a substrate upon application of a releasing agent. In some embodiments, a conditionally removable coating includes a hydrogel as described herein, e.g., a hydrogel including a polypeptide-based material. Non-limiting examples of a hydrogel featuring a polypeptide-based material include a synthetic peptide-based material featuring a combination of spider silk and a trans-membrane segment of human muscle L-type calcium channel (e.g., PEPGEL®), an amphiphilic 16 residue peptide containing a repeating arginine-alanine-aspartate-alanine sequence (RADARADARADARADA) (e.g., PURAMATRIX®), EAK16 (AEAEAKAKAEAEAKAK), KLD12 (KLDLKLKLKL), and PGMATRIX™.

[0442] In some embodiments, the hydrogel in the conditionally removable coating is a stimulus-responsive hydrogel. A stimulus-responsive hydrogel can undergo a gel-to-solution and/or gel-to-solid transition upon application of one or more external triggers (e.g., a releasing agent). See, e.g., Willner, *Acc. Chem. Res.* 50:657-658, 2017, which is incorporated herein by reference in its entirety. Non-limiting examples of a stimulus-responsive hydrogel include a thermoresponsive hydrogel, a pH-responsive hydrogel, a light-responsive hydrogel, a redox-responsive hydrogel, an analyte-responsive hydrogel, or a combination thereof. In some embodiments, a stimulus-responsive hydrogel can be a multi-stimuli-responsive hydrogel.

[0443] A “releasing agent” or “external trigger” is an agent that allows for the removal of a conditionally removable coating from a substrate when the releasing agent is applied to the conditionally removable coating. An external trigger or releasing agent can include physical triggers such as thermal, magnetic, ultrasonic, electrochemical, and/or light stimuli as well as chemical triggers such as pH, redox reactions, supramolecular complexes, and/or biocatalytically driven reactions. See e.g., Echeverria, et al., *Gels* (2018), 4, 54; doi: 10.3390/gels4020054, which is incorporated herein by reference in its entirety. The type of “releasing agent” or “external trigger” can depend on the type of conditionally removable coating. For example, a conditionally removable coating featuring a redox-responsive hydrogel can be removed upon application of a releasing agent that includes a reducing agent such as dithiothreitol (DTT). As another example, a pH-responsive hydrogel can be removed upon the application of a releasing agent that changes the pH.

(iv) Gel Substrates

[0444] In some embodiments, a hydrogel can form a substrate. The term “hydrogel” herein refers to a macromolecular polymer gel including a network. Within the network, some polymer chains

can optionally be cross-linked, although cross-linking does not always occur. In some embodiments, the substrate includes a hydrogel and one or more second materials. In some embodiments, the hydrogel is placed on top of one or more second materials. For example, the hydrogel can be pre-formed and then placed on top of, underneath, or in any other configuration with one or more second materials. In some embodiments, hydrogel formation occurs after contacting one or more second materials during formation of the substrate. Hydrogel formation can also occur within a structure (e.g., wells, ridges, features, projections, and/or markings) located on a substrate. Where the substrate includes a gel (e.g., a hydrogel or gel matrix), oligonucleotides within the gel can attach to the substrate.

[0445] In some embodiments, a hydrogel can include hydrogel subunits. The hydrogel subunits can include any convenient hydrogel subunits, such as, but not limited to, acrylamide, bis-acrylamide, polyacrylamide and derivatives thereof, poly(ethylene glycol) and derivatives thereof (e.g., PEG-acrylate (PEG-DA), PEG-RGD), gelatin-methacryloyl (GelMA), methacrylated hyaluronic acid (MeHA), polyaliphatic polyurethanes, polyether polyurethanes, polyester polyurethanes, polyethylene copolymers, polyamides, polyvinyl alcohols, polypropylene glycol, polytetramethylene oxide, polyvinyl pyrrolidone, polyacrylamide, poly(hydroxyethyl acrylate), and poly(hydroxyethyl methacrylate), collagen, hyaluronic acid, chitosan, dextran, agarose, gelatin, alginate, protein polymers, methylcellulose, and the like, or combinations thereof.

[0446] In some embodiments, a hydrogel includes a hybrid material, e.g., the hydrogel material includes elements of both synthetic and natural polymers. Examples of suitable hydrogels are described, for example, in U.S. Pat. Nos. 6,391,937, 9,512,422, and 9,889,422, and in U.S. Patent Application Publication Nos. 2017/0253918, 2018/0052081 and 2010/0055733, the entire contents of each of which is incorporated herein by reference.

[0447] In some embodiments, cross-linkers and/or initiators are added to hydrogel subunits. Examples of cross-linkers include, without limitation, bis-acrylamide and diazirine. Examples of initiators include, without limitation, azobisisobutyronitrile (AIBN), riboflavin, and L-arginine. Inclusion of cross-linkers and/or initiators can lead to increased covalent bonding between interacting biological macromolecules in later polymerization steps.

[0448] In some embodiments, hydrogels can have a colloidal structure, such as agarose, or a polymer mesh structure, such as gelatin. In some embodiments, the hydrogel is a homopolymeric hydrogel. In some embodiments, the hydrogel is a copolymeric hydrogel. In some embodiments, the hydrogel is a multipolymer interpenetrating polymeric hydrogel.

[0449] In some embodiments, some hydrogel subunits are polymerized (e.g., undergo “formation”) covalently or physically cross-linked, to form a hydrogel network. For example, hydrogel subunits can be polymerized by any method including, but not limited to, thermal crosslinking, chemical crosslinking, physical crosslinking, ionic crosslinking, photo-crosslinking, free-radical initiation crosslinking, an addition reaction, condensation reaction, water-soluble crosslinking reactions, irradiative crosslinking (e.g., x-ray, electron beam), or combinations thereof. Techniques such as lithographic photopolymerization can also be used to form hydrogels.

[0450] In some embodiments, gel beads containing oligonucleotides (e.g., barcoded oligonucleotides such as capture probes) can be deposited on a substrate (e.g., a glass slide). In some embodiments, gel pads can be deposited on a substrate (e.g., a glass slide). In some embodiments, gel pads or gel beads are deposited on a substrate in an arrayed format. In some embodiments in which gel pads or gel beads are deposited on a substrate in an arrayed format, a hydrogel molecular precursor solution can be applied on top of the array (e.g., the array of gel pads or gel beads on a glass slide). In some embodiments, a hydrogel molecular precursor solution can be polymerized such that the deposited gel pads or gel beads are immobilized within the polymerized hydrogel. Any suitable method of polymerization can be used or (e.g., any of the variety of methods described herein). In some embodiments, a polymerized hydrogel that includes the gel pads or gel beads can be removed (e.g., peeled) from the substrate (e.g., glass slide) such

that the gel beads or gel pads are secured in the hydrogel. In some embodiments, a polymerized hydrogel that includes the gel pads or gel beads is a conductive substrate (as described herein) that can be used in accordance with any of the variety of analyte capture methods described herein (e.g., electrophoretic migration of analytes for capture).

[0451] Arrays can be prepared by depositing features (e.g., droplets, beads) on a substrate surface to produce a spatially-barcoded array. Methods of depositing (e.g., droplet manipulation) features are known in the art (see, U.S. Patent Application Publication No. 2008/0132429, Rubina, A. Y., et al., *Biotechniques*. 2003 May; 34 (5): 1008-14, 1016-20, 1022 and Vasiliskov et al. *Biotechniques*. 1999 September; 27 (3): 592-4, 596-8, 600 passim. each herein incorporated by reference in its entirety). A feature can be printed or deposited at a specific location on the substrate (e.g., inkjet printing). In some embodiments, each feature can have a unique oligonucleotide that functions as a spatial barcode. In some embodiments, each feature can have capture probes for multiplexing (e.g., capturing multiple analytes or multiple types of analytes, e.g., proteins and nucleic acids). In some embodiments, a feature can be printed or deposited at the specific location using an electric field. A feature can contain a photo-crosslinkable polymer precursor and an oligonucleotide. In some embodiments, the photo-crosslinkable polymer precursor can be deposited into a patterned feature on the substrate (e.g., well).

[0452] A “photo-crosslinkable polymer precursor” refers to a compound that cross-links and/or polymerizes upon exposure to light. In some embodiments, one or more photoinitiators may also be included to induce and/or promote polymerization and/or cross-linking. See, e.g., Choi et al. *Biotechniques*. 2019 January; 66 (1): 40-53, which is incorporated herein by reference in its entirety.

[0453] Non-limiting examples of photo-crosslinkable polymer precursors include polyethylene (glycol) diacrylate (PEGDA), gelatin-methacryloyl (GelMA), and methacrylated hyaluronic acid (MeHA). In some embodiments, a photo-crosslinkable polymer precursor comprises polyethylene (glycol) diacrylate (PEGDA), gelatin-methacryloyl (GelMA), methacrylated hyaluronic acid (MeHA), or a combination thereof. In some embodiments, a photo-crosslinkable polymer precursor (e.g., PAZAM) can be covalently linked (e.g., cross-linked) to a substrate. In some embodiments, a photo-crosslinkable polymer precursor is not covalently linked to a substrate surface. For example, a silane-free acrylamide can be used (See U.S. Patent Application Publication No. 2011/0059865, herein incorporated by reference in its entirety). The photo-crosslinkable polymer precursor in a feature (e.g., droplet or bead) can be polymerized by any known method. The oligonucleotides can be polymerized in a cross-linked gel matrix (e.g., copolymerized or simultaneously polymerized). In some embodiments, the features containing the photo-crosslinkable polymer precursor deposited on the substrate surface can be exposed to UV light. The UV light can induce polymerization of the photo-crosslinkable polymer precursor and result in the features becoming a gel matrix (e.g., gel pads) on the substrate surface (e.g., array).

[0454] Polymerization methods for hydrogel subunits can be selected to form hydrogels with different properties (e.g., pore volume, swelling properties, biodegradability, conduction, transparency, and/or permeability of the hydrogel). For example, a hydrogel can include pores of sufficient volume to allow the passage of macromolecules, (e.g., nucleic acids, proteins, chromatin, metabolites, gRNA, antibodies, carbohydrates, peptides, metabolites, and/or small molecules) to/from the sample (e.g., tissue section). It is known that pore volume generally decreases with increasing concentration of hydrogel subunits and generally increases with an increasing ratio of hydrogel subunits to cross-linker. Therefore, a hydrogel composition can be prepared that includes a concentration of hydrogel subunits that allows the passage of such biological macromolecules.

[0455] In some embodiments, hydrogel formation on a substrate occurs before, contemporaneously with, or after features (e.g., beads) are attached to the substrate. For example, when a capture probe is attached (e.g., directly or indirectly) to a substrate, hydrogel formation can be performed on the substrate already containing the capture probes.

(d) Arrays

[0456] In many of the methods described herein, features (as described further below) are collectively positioned on a substrate. An “array” is a specific arrangement of a plurality of features that is either irregular or forms a regular pattern. Individual features in the array differ from one another based on their relative spatial locations. In general, at least two of the plurality of features in the array include a distinct capture probe (e.g., any of the examples of capture probes described herein).

[0457] Arrays can be used to measure large numbers of analytes simultaneously. In some embodiments, oligonucleotides are used, at least in part, to create an array. For example, one or more copies of a single species of oligonucleotide (e.g., capture probe) can correspond to or be directly or indirectly attached to a given feature in the array. In some embodiments, a given feature in the array includes two or more species of oligonucleotides (e.g., capture probes). In some embodiments, the two or more species of oligonucleotides (e.g., capture probes) attached directly or indirectly to a given feature on the array include a common (e.g., identical) spatial barcode.

(i) Arrays for Analyte Capture

[0458] In some embodiments, an array can include a capture probe attached directly or indirectly to the substrate. The capture probe can include a capture domain (e.g., a nucleotide sequence) that can specifically bind (e.g., hybridize) to a target analyte (e.g., mRNA, DNA, or protein) within a sample. In some embodiments, the binding of the capture probe to the target (e.g., hybridization) can be detected and quantified by detection of a visual signal, e.g., a fluorophore, a heavy metal (e.g., silver ion), or chemiluminescent label, which has been incorporated into the target. In some embodiments, the intensity of the visual signal correlates with the relative abundance of each analyte in the biological sample. Since an array can contain thousands or millions of capture probes (or more), an array can interrogate many analytes in parallel.

[0459] In some embodiments, a substrate includes one or more capture probes that are designed to capture analytes from one or more organisms. In a non-limiting example, a substrate can contain one or more capture probes designed to capture mRNA from one organism (e.g., a human) and one or more capture probes designed to capture DNA from a second organism (e.g., a bacterium).

[0460] The capture probes can be attached to a substrate or feature using a variety of techniques. In some embodiments, the capture probe is directly attached to a feature that is fixed on an array. In some embodiments, the capture probes are immobilized to a substrate by chemical immobilization. For example, a chemical immobilization can take place between functional groups on the substrate and corresponding functional elements on the capture probes. Exemplary corresponding functional elements in the capture probes can either be an inherent chemical group of the capture probe, e.g., a hydroxyl group, or a functional element can be introduced on to the capture probe. An example of a functional group on the substrate is an amine group. In some embodiments, the capture probe to be immobilized includes a functional amine group or is chemically modified in order to include a functional amine group. Means and methods for such a chemical modification are well known in the art.

[0461] In some embodiments, the capture probe is a nucleic acid. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and includes from the 5' to 3' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and includes from the 5' to 3' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain.

[0462] In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more

barcodes (e.g., a spatial barcode and/or a UMI), a second functional domain, and a capture domain. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and includes from the 5' to 3' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and does not include a spatial barcode. In some embodiments, the capture probe is immobilized on a substrate or feature via its 5' end and does not include a UMI. In some embodiments, the capture probe includes a sequence for initiating a sequencing reaction.

[0463] In some embodiments, the capture probe is immobilized on a substrate or feature via its 3' end. In some embodiments, the capture probe is immobilized on a substrate or feature via its 3' end and includes from the 3' to 5' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe is immobilized on a substrate or feature via its 3' end and includes from the 3' to 5' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe is immobilized on a substrate or feature via its 3' end and includes from the 3' to 5' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain. In some embodiments, the capture probe is immobilized on a substrate or feature via its 3' end and includes from the 3' to 5' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain.

[0464] The localization of the functional group within the capture probe to be immobilized can be used to control and shape the binding behavior and/or orientation of the capture probe, e.g., the functional group can be placed at the 5' or 3' end of the capture probe or within the sequence of the capture probe. In some embodiments, a capture probe can further include a substrate. A typical substrate for a capture probe to be immobilized includes moieties which are capable of binding to such capture probes, e.g., to amine-functionalized nucleic acids. Examples of such substrates are carboxy, aldehyde, or epoxy substrates.

[0465] In some embodiments, the substrates on which capture probes can be immobilized can be chemically activated, e.g., by the activation of functional groups available on the substrate. The term "activated substrate" relates to a material in which interacting or reactive chemical functional groups are established or enabled by chemical modification procedures. For example, a substrate including carboxyl groups can be activated before use. Furthermore, certain substrates contain functional groups that can react with specific moieties already present in the capture probes.

[0466] In some embodiments, a covalent linkage is used to directly couple a capture probe to a substrate. In some embodiments a capture probe is indirectly coupled to a substrate through a linker separating the "first" nucleotide of the capture probe from the substrate, e.g., a chemical linker. In some embodiments, a capture probe does not bind directly to the substrate, but interacts indirectly, for example by binding to a molecule which itself binds directly or indirectly to the substrate. In some embodiments, the capture probe is indirectly attached to a substrate (e.g., attached to a substrate via a solution including a polymer).

[0467] In some embodiments where the capture probe is immobilized on a feature of the array indirectly, e.g., via hybridization to a surface probe capable of binding the capture probe, the capture probe can further include an upstream sequence (5' to the sequence that hybridizes to the nucleic acid, e.g., RNA of the tissue sample) that is capable of hybridizing to 5' end of a surface probe. Alone, the capture domain of the capture probe can be seen as a capture domain oligonucleotide, which can be used in the synthesis of the capture probe in embodiments where the capture probe is immobilized on the array indirectly.

[0468] In some embodiments, a substrate is comprised of an inert material or matrix (e.g., glass slides) that has been functionalized by, for example, treating the substrate with a material comprising reactive groups which enable immobilization of capture probes. See, for example, WO 2017/019456, the entire contents of which is herein incorporated by reference. Non-limiting examples include polyacrylamide hydrogels supported on an inert substrate (e.g., glass slide; see

WO 2005/065814 and U.S. Patent Application No. 2008/0280773, the entire contents of which is incorporated herein by reference).

[0469] In some embodiments, functionalized biomolecules (e.g., capture probes) are immobilized on a functionalized substrate using covalent methods. Methods for covalent attachment include, for example, condensation of amines and activated carboxylic esters (e.g., N-hydroxysuccinimide esters); condensation of amine and aldehydes under reductive amination conditions; and cycloaddition reactions such as the Diels-Alder [4+2] reaction, 1,3-dipolar cycloaddition reactions, and [2+2] cycloaddition reactions. Methods for covalent attachment also include, for example, click chemistry reactions, including [3+2] cycloaddition reactions (e.g., Huisgen 1,3-dipolar cycloaddition reaction and copper (I)-catalyzed azide-alkyne cycloaddition (CuAAC)); thiol-ene reactions; the Diels-Alder reaction and inverse electron demand Diels-Alder reaction; [4+1] cycloaddition of isonitriles and tetrazines; and nucleophilic ring-opening of small carbocycles (e.g., epoxide opening with amino oligonucleotides). Methods for covalent attachment also include, for example, maleimides and thiols; and para-nitrophenyl ester-functionalized oligonucleotides and polylysine-functionalized substrate. Methods for covalent attachment also include, for example, disulfide reactions; radical reactions (see, e.g., U.S. Pat. No. 5,919,626, the entire contents of which are herein incorporated by reference); and hydrazide-functionalized substrate (e.g., wherein the hydrazide functional group is directly or indirectly attached to the substrate) and aldehyde-functionalized oligonucleotides (see, e.g., Yershov et al. (1996) *Proc. Natl. Acad. Sci. USA* 93, 4913-4918, the entire contents of which are herein incorporated by reference).

[0470] In some embodiments, functionalized biomolecules (e.g., capture probes) are immobilized on a functionalized substrate using photochemical covalent methods. Methods for photochemical covalent attachment include, for example, immobilization of anthraquinone-conjugated oligonucleotides (see, e.g., Koch et al. (2000) *Bioconjugate Chem.* 11, 474-483, the entire contents of which is herein incorporated by reference).

[0471] In some embodiments, functionalized biomolecules (e.g., capture probes) are immobilized on a functionalized substrate using non-covalent methods. Methods for non-covalent attachment include, for example, biotin-functionalized oligonucleotides and streptavidin-treated substrates (see, e.g., Holmström et al. (1993) *Analytical Biochemistry* 209, 278-283 and Gilles et al. (1999) *Nature Biotechnology* 17, 365-370, the entire contents of which are herein incorporated by reference).

[0472] In some embodiments, an oligonucleotide (e.g., a capture probe) can be attached to a substrate or feature according to the methods set forth in U.S. Pat. Nos. 6,737,236, 7,259,258, 7,375,234, 7,427,678, 5,610,287, 5,807,522, 5,837,860, and 5,472,881; U.S. Patent Application Publication Nos. 2008/0280773 and 2011/0059865; Shalon et al. (1996) *Genome Research*, 639-645; Rogers et al. (1999) *Analytical Biochemistry* 266, 23-30; Stimpson et al. (1995) *Proc. Natl. Acad. Sci. USA* 92, 6379-6383; Beattie et al. (1995) *Clin. Chem.* 45, 700-706; Lamture et al. (1994) *Nucleic Acids Research* 22, 2121-2125; Beier et al. (1999) *Nucleic Acids Research* 27, 1970-1977; Joos et al. (1997) *Analytical Biochemistry* 247, 96-101; Nikiforov et al. (1995) *Analytical Biochemistry* 227, 201-209; Timofeev et al. (1996) *Nucleic Acids Research* 24, 3142-3148; Chrissey et al. (1996) *Nucleic Acids Research* 24, 3031-3039; Guo et al. (1994) *Nucleic Acids Research* 22, 5456-5465; Running and Urdea (1990) *BioTechniques* 8, 276-279; Fahy et al. (1993) *Nucleic Acids Research* 21, 1819-1826; Zhang et al. (1991) 19, 3929-3933; and Rogers et al. (1997) *Gene Therapy* 4, 1387-1392. The entire contents of each of the foregoing documents is incorporated herein by reference.

(ii) Generation of Capture Probes in an Array Format

[0473] Arrays can be prepared by a variety of methods. In some embodiments, arrays are prepared through the synthesis (e.g., in situ synthesis) of oligonucleotides on the array, or by jet printing or lithography. For example, light-directed synthesis of high-density DNA oligonucleotides can be achieved by photolithography or solid-phase DNA synthesis. To implement photolithographic

synthesis, synthetic linkers modified with photochemical protecting groups can be attached to a substrate and the photochemical protecting groups can be modified using a photolithographic mask (applied to specific areas of the substrate) and light, thereby producing an array having localized photo-deprotection. Many of these methods are known in the art, and are described e.g., in Miller et al., “Basic concepts of microarrays and potential applications in clinical microbiology.” *Clinical Microbiology Reviews* 22.4 (2009): 611-633; US201314111482A; U.S. Pat. No. 9,593,365B2; US2019203275; and WO2018091676, which are each incorporated herein by reference in its entirety.

(1) Spotting or Printing

[0474] In some embodiments, oligonucleotides (e.g., capture probes) can be “spotted” or “printed” onto a substrate to form an array. The oligonucleotides can be applied by either noncontact or contact printing. A noncontact printer can use the same method as computer printers (e.g., bubble jet or inkjet) to expel small droplets of probe solution onto the substrate. The specialized inkjet-like printer can expel nanoliter to picoliter volume droplets of oligonucleotide solution, instead of ink, onto the substrate. In contact printing, each print pin directly applies the oligonucleotide solution onto a specific location on the surface. The oligonucleotides can be attached to the substrate surface by the electrostatic interaction of the negative charge of the phosphate backbone of the DNA with a positively charged coating of the substrate surface or by UV-cross-linked covalent bonds between the thymidine bases in the DNA and amine groups on the treated substrate surface. In some embodiments, the substrate is a glass slide. In some embodiments, the oligonucleotides (e.g., capture probes) are attached to a substrate by a covalent bond to a chemical matrix, e.g., epoxy-silane, amino-silane, lysine, polyacrylamide, etc.

(2) In Situ Synthesis

[0475] Capture probes arrays can be prepared by in situ synthesis. In some embodiments, capture probe arrays can be prepared using photolithography. Photolithography typically relies on UV masking and light-directed combinatorial chemical synthesis on a substrate to selectively synthesize probes directly on the surface of an array, one nucleotide at a time per spot, for many spots simultaneously. In some embodiments, a substrate contains covalent linker molecules that have a protecting group on the free end that can be removed by light. UV light is directed through a photolithographic mask to deprotect and activate selected sites with hydroxyl groups that initiate coupling with incoming protected nucleotides that attach to the activated sites. The mask is designed in such a way that the exposure sites can be selected, and thus specify the coordinates on the array where each nucleotide can be attached. The process can be repeated, a new mask is applied activating different sets of sites and coupling different bases, allowing different oligonucleotides to be constructed at each site. This process can be used to synthesize hundreds of thousands of different oligonucleotides. In some embodiments, maskless array synthesizer technology can be used. Programmable micromirrors can create digital masks that reflect the desired pattern of UV light to deprotect the features.

[0476] In some embodiments, the inkjet spotting process can also be used for in situ oligonucleotide synthesis. The different nucleotide precursors plus catalyst can be printed on the substrate, and are then combined with coupling and deprotection steps. This method relies on printing picoliter volumes of nucleotides on the array surface in repeated rounds of base-by-base printing that extends the length of the oligonucleotide probes on the array.

(3) Electric Fields

[0477] Arrays can also be prepared by active hybridization via electric fields to control nucleic acid transport. Negatively charged nucleic acids can be transported to specific sites, or features, when a positive current is applied to one or more test sites on the array. The surface of the array can contain a binding molecule, e.g., streptavidin, which allows for the formation of bonds (e.g., streptavidin-biotin bonds) once electrically addressed biotinylated probes reach their targeted location. The positive current is then removed from the active features, and new test sites can be

activated by the targeted application of a positive current. The process are repeated until all sites on the array are covered.

(4) Ligation

[0478] In some embodiments, an array comprising barcoded probes can be generated through ligation of a plurality of oligonucleotides. In some instances, an oligonucleotide of the plurality contains a portion of a barcode, and the complete barcode is generated upon ligation of the plurality of oligonucleotides. For example, a first oligonucleotide containing a first portion of a barcode can be attached to a substrate (e.g., using any of the methods of attaching an oligonucleotide to a substrate described herein), and a second oligonucleotide containing a second portion of the barcode can then be ligated onto the first oligonucleotide to generate a complete barcode. Different combinations of the first, second and any additional portions of a barcode can be used to increase the diversity of the barcodes. In instances where the second oligonucleotide is also attached to the substrate prior to ligation, the first and/or the second oligonucleotide can be attached to the substrate via a surface linker which contains a cleavage site. Upon ligation, the ligated oligonucleotide can be linearized by cleaving at the cleavage site.

[0479] To increase the diversity of the barcodes, a plurality of second oligonucleotides comprising two or more different barcode sequences can be ligated onto a plurality of first oligonucleotides that comprise the same barcode sequence, thereby generating two or more different species of barcodes. To achieve selective ligation, a first oligonucleotide attached to a substrate containing a first portion of a barcode can initially be protected with a protective group (e.g., a photocleavable protective group), and the protective group can be removed prior to ligation between the first and second oligonucleotide. In instances where the barcoded probes on an array are generated through ligation of two or more oligonucleotides, a concentration gradient of the oligonucleotides can be applied to a substrate such that different combinations of the oligonucleotides are incorporated into a barcoded probe depending on its location on the substrate.

[0480] Probes can be generated by directly ligating additional oligonucleotides onto existing oligonucleotides via a splint oligonucleotide. In some embodiments, oligonucleotides on an existing array can include a recognition sequence that can hybridize with a splint oligonucleotide. The recognition sequence can be at the free 5' end or the free 3' end of an oligonucleotide on the existing array. Recognition sequences useful for the methods of the present disclosure may not contain restriction enzyme recognition sites or secondary structures (e.g., hairpins), and may include high contents of Guanine and Cytosine nucleotides.

(5) Polymerases

[0481] Barcoded probes on an array can also be generated by adding single nucleotides to existing oligonucleotides on an array, for example, using polymerases that function in a template-independent manner. Single nucleotides can be added to existing oligonucleotides in a concentration gradient, thereby generating probes with varying length, depending on the location of the probes on the array.

(6) Modification of Existing Capture Probes

[0482] Arrays can also be prepared by modifying existing arrays, for example, by modifying oligonucleotides already attached to an arrays. For instance, capture probes can be generated on an array that already comprises oligonucleotides that are attached to the array (or features on the array) at the 3' end and have a free 5' end. In some instances, an array is any commercially available array (e.g., any of the arrays available commercially as described herein). The oligonucleotides can be in situ synthesized using any of the in situ synthesis methods described herein. The oligonucleotide can include a barcode and one or more constant sequences. In some instances, the constant sequences are cleavable sequences. The length of the oligonucleotides attached to the substrate (e.g., array) can be less than 100 nucleotides (e.g., less than 90, 80, 75, 70, 60, 50, 45, 40, 35, 30, 25, 20, 15, or 10 nucleotides). To generate probes using oligonucleotides, a primer complementary to a portion of an oligonucleotide (e.g., a constant sequence shared by the

oligonucleotides) can hybridize to the oligonucleotide and extend the oligonucleotide (using the oligonucleotide as a template) to form a duplex and to create a 3' overhang. The 3' overhang can be created by template-independent ligases (e.g., terminal deoxynucleotidyl transferase (TdT) or poly(A) polymerase). The 3' overhang allows additional nucleotides or oligonucleotides to be added to the duplex, for example, by an enzyme. For instance, a capture probe can be generated by adding additional oligonucleotides to the end of the 3' overhang (e.g., via splint oligonucleotide mediated ligation), where the additional oligonucleotides can include a sequence or a portion of sequence of one or more capture domains, or a complement thereof.

[0483] The additional oligonucleotide (e.g., a sequence or a portion of sequence of a capture domain) can include a degenerate sequence (e.g., any of the degenerate sequences as described herein). The additional oligonucleotide (e.g., a sequence or a portion of sequence of a capture domain) can include a sequence compatible for hybridizing or ligating with an analyte of interest in a biological sample. An analyte of interest can also be used as a splint oligonucleotide to ligate additional oligonucleotides onto a probe. When using a splint oligonucleotide to assist in the ligation of additional oligonucleotides, an additional oligonucleotide can include a sequence that is complementary to the sequence of the splint oligonucleotide. Ligation of the oligonucleotides can involve the use of an enzyme, such as, but not limited to, a ligase. Non-limiting examples of suitable ligases include Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9oN) DNA ligase (9oN™ DNA ligase, New England Biolabs), Ampligase™ (available from Lucigen, Middleton, WI), and SplintR (available from New England Biolabs, Ipswich, MA). An array generated as described above is useful for spatial analysis of a biological sample. For example, the one or more capture domains can be used to hybridize with the poly(A) tail of an mRNA molecule. Reverse transcription can be carried out using a reverse transcriptase to generate cDNA complementary to the captured mRNA. The sequence and location of the captured mRNA can then be determined (e.g., by sequencing the capture probe that contains the barcode as well as the complementary cDNA).

[0484] An array for spatial analysis can be generated by various methods as described herein. In some embodiments, the array has a plurality of capture probes comprising spatial barcodes. These spatial barcodes and their relationship to the locations on the array can be determined. In some cases, such information is readily available, because the oligonucleotides are spotted, printed, or synthesized on the array with a pre-determined pattern. In some cases, the spatial barcode can be decoded by methods described herein, e.g., by in situ sequencing, by various labels associated with the spatial barcodes etc. In some embodiments, an array can be used as a template to generate a daughter array. Thus, the spatial barcode can be transferred to the daughter array with a known pattern.

(iii) Features

[0485] A “feature” is an entity that acts as a support or repository for various molecular entities used in sample analysis. In some embodiments, some or all of the features in an array are functionalized for analyte capture. In some embodiments, functionalized features include one or more capture probe(s). Examples of features include, but are not limited to, a bead, a spot of any two- or three-dimensional geometry (e.g., an ink jet spot, a masked spot, a square on a grid), a well, and a hydrogel pad. In some embodiments, features are directly or indirectly attached or fixed to a substrate. In some embodiments, the features are not directly or indirectly attached or fixed to a substrate, but instead, for example, are disposed within an enclosed or partially enclosed three dimensional space (e.g., wells or divots).

[0486] In addition to those above, a wide variety of other features can be used to form the arrays described herein. For example, in some embodiments, features that are formed from polymers and/or biopolymers that are jet printed, screen printed, or electrostatically deposited on a substrate can be used to form arrays. Jet printing of biopolymers is described, for example, in PCT Patent Application Publication No. WO 2014/085725. Jet printing of polymers is described, for example,

in de Gans et al., *Adv Mater.* 16 (3): 203-213 (2004). Methods for electrostatic deposition of polymers and biopolymers are described, for example, in Hoyer et al., *Anal. Chem.* 68 (21): 3840-3844 (1996). The entire contents of each of the foregoing references are incorporated herein by reference.

[0487] As another example, in some embodiments, features are formed by metallic micro- or nanoparticles. Suitable methods for depositing such particles to form arrays are described, for example, in Lee et al., *Beilstein J. Nanotechnol.* 8:1049-1055 (2017), the entire contents of which are incorporated herein by reference.

[0488] As a further example, in some embodiments, features are formed by magnetic particles that are assembled on a substrate. Examples of such particles and methods for assembling arrays are described in Ye et al., *Scientific Reports* 6:23145 (2016), the entire contents of which are incorporated herein by reference.

[0489] As another example, in some embodiments, features correspond to regions of a substrate in which one or more optical labels have been incorporated, and/or which have been altered by a process such as permanent photobleaching. Suitable substrates to implement features in this manner include a wide variety of polymers, for example. Methods for forming such features are described, for example, in Moshrefzadeh et al., *Appl. Phys. Lett.* 62:16 (1993), the entire contents of which are incorporated herein by reference.

[0490] As yet another example, in some embodiments, features can correspond to colloidal particles assembled (e.g., via self-assembly) to form an array. Suitable colloidal particles are described for example in Sharma, *Resonance* 23 (3): 263-275 (2018), the entire contents of which are incorporated herein by reference.

[0491] As a further example, in some embodiments, features can be formed via spot-array photopolymerization of a monomer solution on a substrate. In particular, two-photon and three-photon polymerization can be used to fabricate features of relatively small (e.g., sub-micron) dimensions. Suitable methods for preparing features on a substrate in this manner are described for example in Nguyen et al., *Materials Today* 20 (6): 314-322 (2017), the entire contents of which are incorporated herein by reference.

[0492] In some embodiments, features are directly or indirectly attached or fixed to a substrate that is liquid permeable. In some embodiments, features are directly or indirectly attached or fixed to a substrate that is biocompatible. In some embodiments, features are directly or indirectly attached or fixed to a substrate that is a hydrogel.

[0493] FIG. 12 depicts an exemplary arrangement of barcoded features within an array. From left to right, FIG. 12 shows (L) a slide including six spatially-barcoded arrays, (C) an enlarged schematic of one of the six spatially-barcoded arrays, showing a grid of barcoded features in relation to a biological sample, and (R) an enlarged schematic of one section of an array, showing the specific identification of multiple features within the array (labelled as ID578, ID579, ID560, etc.).

(1) Beads

[0494] A “bead” can be a particle. A bead can be porous, non-porous, solid, semi-solid, and/or a combination thereof. In some embodiments, a bead can be dissolvable, disruptable, and/or degradable, whereas in certain embodiments, a bead is not degradable. A semi-solid bead can be a liposomal bead. Solid beads can include metals including, without limitation, iron oxide, gold, and silver. In some embodiments, the bead can be a silica bead. In some embodiments, the bead can be rigid. In some embodiments, the bead can be flexible and/or compressible.

[0495] The bead can be a macromolecule. The bead can be formed of nucleic acid molecules bound together. The bead can be formed via covalent or non-covalent assembly of molecules (e.g., macromolecules), such as monomers or polymers. Polymers or monomers can be natural or synthetic. Polymers or monomers can be or include, for example, nucleic acid molecules (e.g., DNA or RNA).

[0496] A bead can be rigid, or flexible and/or compressible. A bead can include a coating including one or more polymers. Such a coating can be disruptable or dissolvable. In some embodiments, a bead includes a spectral or optical label (e.g., dye) attached directly or indirectly (e.g., through a linker) to the bead. For example, a bead can be prepared as a colored preparation (e.g., a bead exhibiting a distinct color within the visible spectrum) that can change color (e.g., colorimetric beads) upon application of a desired stimulus (e.g., heat and/or chemical reaction) to form differently colored beads (e.g., opaque and/or clear beads).

[0497] A bead can include natural and/or synthetic materials. For example, a bead can include a natural polymer, a synthetic polymer or both natural and synthetic polymers.

[0498] Examples of natural polymers include, without limitation, proteins, sugars such as deoxyribonucleic acid, rubber, cellulose, starch (e.g., amylose, amylopectin), enzymes, polysaccharides, silks, polyhydroxyalkanoates, chitosan, dextran, collagen, carrageenan, ispaghula, acacia, agar, gelatin, shellac, sterculia gum, xanthan gum, corn sugar gum, guar gum, gum karaya, agarose, alginic acid, alginate, or natural polymers thereof. Examples of synthetic polymers include, without limitation, acrylics, nylons, silicones, spandex, viscose rayon, polycarboxylic acids, polyvinyl acetate, polyacrylamide, polyacrylate, polyethylene glycol, polyurethanes, polylactic acid, silica, polystyrene, polyacrylonitrile, polybutadiene, polycarbonate, polyethylene, polyethylene terephthalate, poly(chlorotrifluoroethylene), poly(ethylene oxide), poly(ethylene terephthalate), polyethylene, polyisobutylene, poly(methyl methacrylate), poly(oxymethylene), polyformaldehyde, polypropylene, polystyrene, poly(tetrafluoroethylene), poly(vinyl acetate), poly(vinyl alcohol), poly(vinyl chloride), poly(vinylidene dichloride), poly(vinylidene difluoride), poly(vinyl fluoride) and/or combinations (e.g., co-polymers) thereof. Beads can also be formed from materials other than polymers, including for example, lipids, micelles, ceramics, glass-ceramics, material composites, metals, and/or other inorganic materials.

[0499] In some embodiments, a bead is a degradable bead. A degradable bead can include one or more species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) with a labile bond such that, when the bead/species is exposed to the appropriate stimuli, the labile bond is broken and the bead degrades. The labile bond can be a chemical bond (e.g., covalent bond, ionic bond) or can be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.). In some embodiments, a cross-linker used to generate a bead can include a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead including cystamine cross-linkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

[0500] Degradation can refer to the disassociation of a bound or entrained species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species can be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore volumes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some embodiments, an increase in pore volume due to osmotic swelling of a bead can permit the release of entrained species within the bead. In some embodiments, osmotic shrinking of a bead can cause a bead to better retain an entrained species due to pore volume contraction.

[0501] Any suitable agent that can degrade beads can be used. In some embodiments, changes in temperature or pH can be used to degrade thermo-sensitive or pH-sensitive bonds within beads. In some embodiments, chemical degrading agents can be used to degrade chemical bonds within beads by oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as DTT, where DTT can degrade the disulfide bonds formed between a cross-linker and gel precursors, thus degrading the bead. In some embodiments, a reducing agent can be added to degrade the bead, which can cause the bead to release its contents. Examples of reducing agents can include, without limitation, dithiothreitol (DTT), β -mercaptoethanol, (2S)-2-

amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris (2-carboxyethyl) phosphine (TCEP), or combinations thereof.

[0502] Any of a variety of chemical agents can be used to trigger the degradation of beads. Examples of chemical agents include, but are not limited to, pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

[0503] In some embodiments, a bead can be formed from materials that include degradable chemical cross-linkers, such as N,N'-bis-(acryloyl) cystamine (BAC) or cystamine. Degradation of such degradable cross-linkers can be accomplished through any variety of mechanisms. In some examples, a bead can be contacted with a chemical degrading agent that can induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as dithiothreitol (DTT). Additional examples of reducing agents can include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris (2-carboxyethyl) phosphine (TCEP), or combinations thereof.

[0504] In some embodiments, exposure to an aqueous solution, such as water, can trigger hydrolytic degradation, and thus degradation of the bead. Beads can also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat can cause melting of a bead such that a portion of the bead degrades. In some embodiments, heat can increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat can also act upon heat-sensitive polymers used as materials to construct beads.

[0505] Where degradable beads are used, it can be beneficial to avoid exposing such beads to the stimulus or stimuli that cause such degradation prior to a given time, in order to, for example, avoid premature bead degradation and issues that arise from such degradation, including for example poor flow characteristics and aggregation. By way of example, where beads include reducible cross-linking groups, such as disulfide groups, it will be desirable to avoid contacting such beads with reducing agents, e.g., DTT or other disulfide cleaving reagents. In such embodiments, treatment of the beads described herein will, in some embodiments be provided free of reducing agents, such as DTT. Because reducing agents are often provided in commercial enzyme preparations, it can be desirable to provide reducing agent free (or DTT free) enzyme preparations in treating the beads described herein. Examples of such enzymes include, e.g., polymerase enzyme preparations, reverse transcriptase enzyme preparations, ligase enzyme preparations, as well as many other enzyme preparations that can be used to treat the beads described herein. The terms "reducing agent free" or "DTT free" preparations refer to a preparation having less than about 1/10th, less than about 1/50th, or less than about 1/100th of the lower ranges for such materials used in degrading the beads. For example, for DTT, the reducing agent free preparation can have less than about 0.01 millimolar (mM), 0.005 mM, 0.001 mM DTT, 0.0005 mM DTT, or less than about 0.0001 mM DTT. In some embodiments, the amount of DTT can be undetectable.

[0506] A degradable bead can be useful to more quickly release an attached capture probe (e.g., a nucleic acid molecule, a spatial barcode sequence, and/or a primer) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species can have greater mobility and accessibility to other species in solution upon degradation of the bead. In some embodiments, a species can also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker can respond to the same stimuli as the degradable bead or the two degradable species can respond to different stimuli. For example, a capture probe having one or more spatial barcodes can be attached, via a disulfide bond, to a polyacrylamide bead including cystamine. Upon exposure of the spatially-barcoded bead to a reducing agent, the bead degrades and the capture probe having the one or more spatial barcode sequences is released upon breakage of both the disulfide linkage between the capture probe and

the bead and the disulfide linkages of the cystamine in the bead.

[0507] The addition of multiple types of labile bonds to a bead can result in the generation of a bead capable of responding to varied stimuli. Each type of labile bond can be sensitive to an associated stimulus (e.g., chemical stimulus, light, temperature, pH, enzymes, etc.) such that release of reagents attached to a bead via each labile bond can be controlled by the application of the appropriate stimulus. Some non-limiting examples of labile bonds that can be coupled to a precursor or bead include an ester linkage (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)). A bond can be cleavable via other nucleic acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases). Such functionality can be useful in controlled release of reagents from a bead. In some embodiments, another reagent including a labile bond can be linked to a bead after gel bead formation via, for example, an activated functional group of the bead as described above. In some embodiments, a gel bead including a labile bond is reversible. In some embodiments, a gel bead with a reversible labile bond is used to capture one or more regions of interest of a biological sample. For example, without limitation, a bead including a thermolabile bond can be heated by a light source (e.g., a laser) that causes a change in the gel bead that facilitates capture of a biological sample in contact with the gel bead. Capture probes having one or more spatial barcodes that are releasably, cleavably, or reversibly attached to the beads described herein include capture probes that are released or releasable through cleavage of a linkage between the capture probe and the bead, or that are released through degradation of the underlying bead itself, allowing the capture probes having the one or more spatial barcodes to be accessed or become accessible by other reagents, or both.

[0508] Beads can have different physical properties. Physical properties of beads can be used to characterize the beads. Non-limiting examples of physical properties of beads that can differ include volume, shape, circularity, density, symmetry, and hardness. For example, beads can be of different volumes. Beads of different diameters can be obtained by using microfluidic channel networks configured to provide beads of a specific volume (e.g., based on channel sizes, flow rates, etc.). In some embodiments, beads have different hardness values that can be obtained by varying the concentration of polymer used to generate the beads. In some embodiments, a spatial barcode attached to a bead can be made optically detectable using a physical property of the capture probe. For example, a nucleic acid origami, such as a deoxyribonucleic acid (DNA) origami, can be used to generate an optically detectable spatial barcode. To do so, a nucleic acid molecule, or a plurality of nucleic acid molecules, can be folded to create two- and/or three-dimensional geometric shapes. The different geometric shapes can be optically detected.

[0509] In some embodiments, special types of nanoparticles with more than one distinct physical property can be used to make the beads physically distinguishable. For example, Janus particles with both hydrophilic and hydrophobic surfaces can be used to provide unique physical properties.

[0510] A bead can generally be of any suitable shape. Examples of bead shapes include, but are not limited to, spherical, non-spherical, oval, oblong, amorphous, circular, cylindrical, cuboidal, hexagonal, and variations thereof. In some embodiments, non-spherical (e.g., hexagonal, cuboidal, shaped beads can assemble more closely (e.g., tighter) than spherical shaped beads. In some embodiments, beads can self-assemble into a monolayer. A cross section (e.g., a first cross-section) can correspond to a diameter or maximum cross-sectional dimension of the bead. In some embodiments, the bead can be approximately spherical. In such embodiments, the first cross-section can correspond to the diameter of the bead. In some embodiments, the bead can be approximately cylindrical. In such embodiments, the first cross-section can correspond to a diameter, length, or width along the approximately cylindrical bead.

[0511] Beads can be of uniform size or heterogeneous size. “Polydispersity” generally refers to heterogeneity of sizes of molecules or particles. The polydispersity index (PDI) of a bead can be calculated using the equation $PDI = M_w/M_n$, where M_w is the weight-average molar mass and M_n is the number-average molar mass. In certain embodiments, beads can be provided as a population or plurality of beads having a relatively monodisperse size distribution. Where it can be desirable to provide relatively consistent amounts of reagents, maintaining relatively consistent bead characteristics, such as size, can contribute to the overall consistency.

[0512] In some embodiments, the beads provided herein can have size distributions that have a coefficient of variation in their cross-sectional dimensions of less than 50%, less than 40%, less than 30%, less than 20%, less than 15%, less than 10%, less than 5%, or lower. In some embodiments, a plurality of beads provided herein has a polydispersity index of less than 50%, less than 45%, less than 40%, less than 35%, less than 30%, less than 25%, less than 20%, less than 15%, less than 10%, less than 5%, or lower.

[0513] In some embodiments, the bead can have a diameter or maximum dimension no larger than 100 μm (e.g., no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm .)

[0514] In some embodiments, a plurality of beads has an average diameter no larger than 100 μm . In some embodiments, a plurality of beads has an average diameter or maximum dimension no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm .

[0515] In some embodiments, the volume of the bead can be at least about 1 μm^3 , e.g., at least 1 μm^3 , 2 μm^3 , 3 μm^3 , 4 μm^3 , 5 μm^3 , 6 μm^3 , 7 μm^3 , 8 μm^3 , 9 μm^3 , 10 μm^3 , 12 μm^3 , 14 μm^3 , 16 μm^3 , 18 μm^3 , 20 μm^3 , 25 μm^3 , 30 μm^3 , 35 μm^3 , 40 μm^3 , 45 μm^3 , 50 μm^3 , 55 μm^3 , 60 μm^3 , 65 μm^3 , 70 μm^3 , 75 μm^3 , 80 μm^3 , 85 μm^3 , 90 μm^3 , 95 μm^3 , 100 μm^3 , 125 μm^3 , 150 μm^3 , 175 μm^3 , 200 μm^3 , 250 μm^3 , 300 μm^3 , 350 μm^3 , 400 μm^3 , 450 μm^3 , μm^3 , 500 μm^3 , 550 μm^3 , 600 μm^3 , 650 μm^3 , 700 μm^3 , 750 μm^3 , 800 μm^3 , 850 μm^3 , 900 μm^3 , 950 μm^3 , 1000 μm^3 , 1200 μm^3 , 1400 μm^3 , 1600 μm^3 , 1800 μm^3 , 2000 μm^3 , 2200 μm^3 , 2400 μm^3 , 2600 μm^3 , 2800 μm^3 , 3000 μm^3 , or greater.

[0516] In some embodiments, the bead can have a volume of between about 1 μm^3 and 100 μm^3 , such as between about 1 μm^3 and 10 μm^3 , between about 10 μm^3 and 50 μm^3 , or between about 50 μm^3 and 100 μm^3 . In some embodiments, the bead can include a volume of between about 100 μm^3 and 1000 μm^3 , such as between about 100 μm^3 and 500 μm^3 or between about 500 μm^3 and 1000 μm^3 . In some embodiments, the bead can include a volume between about 1000 μm^3 and 3000 μm^3 , such as between about 1000 μm^3 and 2000 μm^3 or between about 2000 μm^3 and 3000 μm^3 . In some embodiments, the bead can include a volume between about 1 μm^3 and 3000 μm^3 , such as between about 1 μm^3 and 2000 μm^3 , between about 1 μm^3 and 1000 μm^3 , between about 1 μm^3 and 500 μm^3 , or between about 1 μm^3 and 250 μm^3 .

[0517] The bead can include one or more cross-sections that can be the same or different. In some embodiments, the bead can have a first cross-section that is different from a second cross-section. The bead can have a first cross-section that is at least about 0.0001 micrometer, 0.001 micrometer, 0.01 micrometer, 0.1 micrometer, or 1 micrometer. In some embodiments, the bead can include a cross-section (e.g., a first cross-section) of at least about 1 micrometer (μm), 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , 7 μm , 8 μm , 9 μm , 10 μm , 11 μm , 12 μm , 13 μm , 14 μm , 15 μm , 16 μm , 17 μm , 18 μm ,

19 μm , 20 μm , 25 μm , 30 μm , 35 μm , 40 μm , 45 μm , 50 μm , 55 μm , 60 μm , 65 μm , 70 μm , 75 μm , 80 μm , 85 μm , 90 μm , 100 μm , 120 μm , 140 μm , 160 μm , 180 μm , 200 μm , 250 μm , 300 μm , 350 μm , 400 μm , 450 μm , 500 μm , 550 μm , 600 μm , 650 μm , 700 μm , 750 μm , 800 μm , 850 μm , 900 μm , 950 μm , 1 millimeter (mm), or greater. In some embodiments, the bead can include a cross-section (e.g., a first cross-section) of between about 1 μm and 500 μm , such as between about 1 μm and 100 μm , between about 100 μm and 200 μm , between about 200 μm and 300 μm , between about 300 μm and 400 μm , or between about 400 μm and 500 μm . For example, the bead can include a cross-section (e.g., a first cross-section) of between about 1 μm and 100 μm . In some embodiments, the bead can have a second cross-section that is at least about 1 μm . For example, the bead can include a second cross-section of at least about 1 micrometer (μm), 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , 7 μm , 8 μm , 9 μm , 10 μm , 11 μm , 12 μm , 13 μm , 14 μm , 15 μm , 16 μm , 17 μm , 18 μm , 19 μm , 20 μm , 25 μm , 30 μm , 35 μm , 40 μm , 45 μm , 50 μm , 55 μm , 60 μm , 65 μm , 70 μm , 75 μm , 80 μm , 85 μm , 90 μm , 100 μm , 120 μm , 140 μm , 160 μm , 180 μm , 200 μm , 250 μm , 300 μm , 350 μm , 400 μm , 450 μm , 500 μm , 550 μm , 600 μm , 650 μm , 700 μm , 750 μm , 800 μm , 850 μm , 900 μm , 950 μm , 1 millimeter (mm), or greater. In some embodiments, the bead can include a second cross-section of between about 1 μm and 500 μm , such as between about 1 μm and 100 μm , between about 100 μm and 200 μm , between about 200 μm and 300 μm , between about 300 μm and 400 μm , or between about 400 μm and 500 μm . For example, the bead can include a second cross-section of between about 1 μm and 100 μm .

[0518] In some embodiments, beads can be of a nanometer scale (e.g., beads can have a diameter or maximum cross-sectional dimension of about 100 nanometers (nm) to about 900 nanometers (nm) (e.g., 850 nm or less, 800 nm or less, 750 nm or less, 700 nm or less, 650 nm or less, 600 nm or less, 550 nm or less, 500 nm or less, 450 nm or less, 400 nm or less, 350 nm or less, 300 nm or less, 250 nm or less, 200 nm or less, 150 nm or less). A plurality of beads can have an average diameter or average maximum cross-sectional dimension of about 100 nanometers (nm) to about 900 nanometers (nm) (e.g., 850 nm or less, 800 nm or less, 750 nm or less, 700 nm or less, 650 nm or less, 600 nm or less, 550 nm or less, 500 nm or less, 450 nm or less, 400 nm or less, 350 nm or less, 300 nm or less, 250 nm or less, 200 nm or less, 150 nm or less). In some embodiments, a bead has a diameter or volume that is about the diameter of a single cell (e.g., a single cell under evaluation).

[0519] In some embodiments, a bead is able to identify multiple analytes (e.g., nucleic acids, proteins, chromatin, metabolites, drugs, gRNA, and lipids) from a single cell. In some embodiments, a bead is able to identify a single analyte from a single cell (e.g., mRNA).

[0520] A bead can have a tunable pore volume. The pore volume can be chosen to, for instance, retain denatured nucleic acids. The pore volume can be chosen to maintain diffusive permeability to exogenous chemicals such as sodium hydroxide (NaOH) and/or endogenous chemicals such as inhibitors. A bead can be formed of a biocompatible and/or biochemically compatible material, and/or a material that maintains or enhances cell viability. A bead can be formed from a material that can be depolymerized thermally, chemically, enzymatically, and/or optically.

[0521] In some embodiments, beads can be non-covalently loaded with one or more reagents. The beads can be non-covalently loaded by, for instance, subjecting the beads to conditions sufficient to swell the beads, allowing sufficient time for the reagents to diffuse into the interiors of the beads, and subjecting the beads to conditions sufficient to de-swell the beads. Swelling of the beads can be accomplished, for instance, by placing the beads in a thermodynamically favorable solvent, subjecting the beads to a higher or lower temperature, subjecting the beads to a higher or lower ion concentration, and/or subjecting the beads to an electric field.

[0522] The swelling of the beads can be accomplished by various swelling methods. In some embodiments, swelling is reversible (e.g., by subjecting beads to conditions that promote de-swelling). In some embodiments, the de-swelling of the beads is accomplished, for instance, by transferring the beads in a thermodynamically unfavorable solvent, subjecting the beads to lower or

higher temperatures, subjecting the beads to a lower or higher ion concentration, and/or adding or removing an electric field. The de-swelling of the beads can be accomplished by various de-swelling methods. In some embodiments, de-swelling is reversible (e.g., subject beads to conditions that promote swelling). In some embodiments, the de-swelling of beads can include transferring the beads to cause pores in the bead to shrink. The shrinking can then hinder reagents within the beads from diffusing out of the interiors of the beads. The hindrance created can be due to steric interactions between the reagents and the interiors of the beads. The transfer can be accomplished microfluidically. For instance, the transfer can be achieved by moving the beads from one co-flowing solvent stream to a different co-flowing solvent stream. The swellability and/or pore volume of the beads can be adjusted by changing the polymer composition of the bead. [0523] A bead can include a polymer that is responsive to temperature so that when the bead is heated or cooled, the characteristics or dimensions of the bead can change. For example, a polymer can include poly(N-isopropylacrylamide). A gel bead can include poly(N-isopropylacrylamide) and when heated the gel bead can decrease in one or more dimensions (e.g., a cross-sectional diameter, multiple cross-sectional diameters). A temperature sufficient for changing one or more characteristics of the gel bead can be, for example, at least about 0 degrees Celsius ($^{\circ}$ C.), 1° C., 2° C., 3° C., 4° C., 5° C., 10° C., or higher. For example, the temperature can be about 4° C. In some embodiments, a temperature sufficient for changing one or more characteristics of the gel bead can be, for example, at least about 25° C., 30° C., 35° C., 37° C., 40° C., 45° C., 50° C., or higher. For example, the temperature can be about 37° C.

[0524] Functionalization of beads for attachment of capture probes can be achieved through a wide range of different approaches, including, without limitation, activation of chemical groups within a polymer, incorporation of active or activatable functional groups in the polymer structure, or attachment at the pre-polymer or monomer stage in bead production. The bead can be functionalized to bind to targeted analytes, such as nucleic acids, proteins, carbohydrates, lipids, metabolites, peptides, or other analytes.

[0525] In some embodiments, a bead can contain molecular precursors (e.g., monomers or polymers), which can form a polymer network via polymerization of the molecular precursors. In some embodiments, a precursor can be an already polymerized species capable of undergoing further polymerization via, for example, a chemical cross-linkage. In some embodiments, a precursor can include one or more of an acrylamide or a methacrylamide monomer, oligomer, or polymer. In some embodiments, the bead can include prepolymers, which are oligomers capable of further polymerization. For example, polyurethane beads can be prepared using prepolymers. In some embodiments, a bead can contain individual polymers that can be further polymerized together (e.g., to form a co-polymer). In some embodiments, a bead can be generated via polymerization of different precursors, such that they include mixed polymers, co-polymers, and/or block co-polymers. In some embodiments, a bead can include covalent or ionic bonds between polymeric precursors (e.g., monomers, oligomers, and linear polymers), nucleic acid molecules (e.g., oligonucleotides), primers, and other entities. In some embodiments, covalent bonds can be carbon-carbon bonds or thioether bonds.

[0526] Cross-linking of polymers can be permanent or reversible, depending upon the particular cross-linker used. Reversible cross-linking can allow the polymer to linearize or dissociate under appropriate conditions. In some embodiments, reversible cross-linking can also allow for reversible attachment of a material bound to the surface of a bead. In some embodiments, a cross-linker can form a disulfide linkage. In some embodiments, a chemical cross-linker forming a disulfide linkage can be cystamine or a modified cystamine.

[0527] For example, where the polymer precursor material includes a linear polymer material, such as a linear polyacrylamide, PEG, or other linear polymeric material, the activation agent can include a cross-linking agent, or a chemical that activates a cross-linking agent within formed droplets. Likewise, for polymer precursors that include polymerizable monomers, the activation

agent can include a polymerization initiator. For example, in certain embodiments, where the polymer precursor includes a mixture of acrylamide monomer with a N,N'-bis-(acryloyl) cystamine (BAC) comonomer, an agent such as tetraethylmethylenediamine (TEMED) can be provided, which can initiate the copolymerization of the acrylamide and BAC into a cross-linked polymer network, or other conditions sufficient to polymerize or gel the precursors. The conditions sufficient to polymerize or gel the precursors can include exposure to heating, cooling, electromagnetic radiation, and/or light.

[0528] Following polymerization or gelling, a polymer or gel can be formed. The polymer or gel can be diffusively permeable to chemical or biochemical reagents. The polymer or gel can be diffusively impermeable to macromolecular constituents. The polymer or gel can include one or more of disulfide cross-linked polyacrylamide, agarose, alginate, polyvinyl alcohol, polyethylene glycol (PEG)-diacrylate, PEG-acrylate, PEG-thiol, PEG-azide, PEG-alkyne, other acrylates, chitosan, hyaluronic acid, collagen, fibrin, gelatin, or elastin. The polymer or gel can include any other polymer or gel.

[0529] In some embodiments, disulfide linkages can be formed between molecular precursor units (e.g., monomers, oligomers, or linear polymers) or precursors incorporated into a bead and nucleic acid molecules (e.g., oligonucleotides, capture probes). Cystamine (including modified cystamines), for example, is an organic agent including a disulfide bond that can be used as a cross-linker agent between individual monomeric or polymeric precursors of a bead. Polyacrylamide can be polymerized in the presence of cystamine or a species including cystamine (e.g., a modified cystamine) to generate polyacrylamide gel beads including disulfide linkages (e.g., chemically degradable beads including chemically-reducible cross-linkers). The disulfide linkages can permit the bead to be degraded (or dissolved) upon exposure of the bead to a reducing agent.

[0530] In some embodiments, chitosan, a linear polysaccharide polymer, can be cross-linked with glutaraldehyde via hydrophilic chains to form a bead. Crosslinking of chitosan polymers can be achieved by chemical reactions that are initiated by heat, pressure, change in pH, and/or radiation.

[0531] In some embodiments, a bead can include an acrydite moiety, which in certain aspects can be used to attach one or more capture probes to the bead. In some embodiments, an acrydite moiety can refer to an acrydite analogue generated from the reaction of acrydite with one or more species (e.g., disulfide linkers, primers, other oligonucleotides, etc.), such as, without limitation, the reaction of acrydite with other monomers and cross-linkers during a polymerization reaction.

Acrydite moieties can be modified to form chemical bonds with a species to be attached, such as a capture probe. Acrydite moieties can be modified with thiol groups capable of forming a disulfide bond or can be modified with groups already including a disulfide bond. The thiol or disulfide (via disulfide exchange) can be used as an anchor point for a species to be attached or another part of the acrydite moiety can be used for attachment. In some embodiments, attachment can be reversible, such that when the disulfide bond is broken (e.g., in the presence of a reducing agent), the attached species is released from the bead. In some embodiments, an acrydite moiety can include a reactive hydroxyl group that can be used for attachment of species.

[0532] In some embodiments, precursors (e.g., monomers or cross-linkers) that are polymerized to form a bead can include acrydite moieties, such that when a bead is generated, the bead also includes acrydite moieties. The acrydite moieties can be attached to a nucleic acid molecule (e.g., an oligonucleotide), which can include a priming sequence (e.g., a primer for amplifying target nucleic acids, random primer, primer sequence for messenger RNA) and/or one or more capture probes. The one or more capture probes can include sequences that are the same for all capture probes coupled to a given bead and/or sequences that are different across all capture probes coupled to the given bead. The capture probe can be incorporated into the bead. In some embodiments, the capture probe can be incorporated or attached to the bead such that the capture probe retains a free 3' end. In some embodiments, the capture probe can be incorporated or attached to the bead such that the capture probe retains a free 5' end. In some embodiments, beads can be

functionalized such that each bead contains a plurality of different capture probes. For example, a bead can include a plurality of capture probes e.g., Capture Probe 1, Capture Probe 2, and Capture Probe 3, and each of Capture Probes 1, Capture Probes 2, and Capture Probes 3 contain a distinct capture domain (e.g., capture domain of Capture Probe 1 includes a poly(dT) capture domain, capture domain of Capture Probe 2 includes a gene-specific capture domain, and capture domain of Capture Probe 3 includes a CRISPR-specific capture domain). By functionalizing beads to contain a plurality of different capture domains per bead, the level of multiplex capability for analyte detection can be improved.

[0533] In some embodiments, precursors (e.g., monomers or cross-linkers) that are polymerized to form a bead can include a functional group that is reactive or capable of being activated such that when it becomes reactive it can be polymerized with other precursors to generate beads including the activated or activatable functional group. The functional group can then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the beads. For example, some precursors including a carboxylic acid (COOH) group can co-polymerize with other precursors to form a bead that also includes a COOH functional group. In some embodiments, acrylic acid (a species including free COOH groups), acrylamide, and bis(acryloyl) cystamine can be co-polymerized together to generate a bead including free COOH groups. The COOH groups of the bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species including an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) as a functional group on a moiety to be linked to the bead.

[0534] Beads including disulfide linkages in their polymeric network can be functionalized with additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) via reduction of some of the disulfide linkages to free thiols. The disulfide linkages can be reduced via, for example, the action of a reducing agent (e.g., DTT, TCEP, etc.) to generate free thiol groups, without dissolution of the bead. Free thiols of the beads can then react with free thiols of a species or a species including another disulfide bond (e.g., via thiol-disulfide exchange) such that the species can be linked to the beads (e.g., via a generated disulfide bond). In some embodiments, free thiols of the beads can react with any other suitable group. For example, free thiols of the beads can react with species including an acrydite moiety. The free thiol groups of the beads can react with the acrydite via Michael addition chemistry, such that the species including the acrydite is linked to the bead. In some embodiments, uncontrolled reactions can be prevented by inclusion of a thiol capping agent such as N-ethylmaleimide or iodoacetate.

[0535] Activation of disulfide linkages within a bead can be controlled such that only a small number of disulfide linkages are activated. Control can be exerted, for example, by controlling the concentration of a reducing agent used to generate free thiol groups and/or concentration of reagents used to form disulfide bonds in bead polymerization. In some embodiments, a low concentration of reducing agent (e.g., molecules of reducing agent: gel bead ratios) of less than or equal to about 1:100,000,000,000, less than or equal to about 1:10,000,000,000, less than or equal to about 1:1,000,000,000, less than or equal to about 1:100,000,000, less than or equal to about 1:10,000,000, less than or equal to about 1:1,000,000, less than or equal to about 1:100,000, or less than or equal to about 1:10,000) can be used for reduction. Controlling the number of disulfide linkages that are reduced to free thiols can be useful in ensuring bead structural integrity during functionalization. In some embodiments, optically-active agents, such as fluorescent dyes can be coupled to beads via free thiol groups of the beads and used to quantify the number of free thiols present in a bead and/or track a bead.

[0536] In some embodiments, addition of moieties to a bead after bead formation can be

advantageous. For example, addition of a capture probe after bead formation can avoid loss of the species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) during chain transfer termination that can occur during polymerization. In some embodiments, smaller precursors (e.g., monomers or cross linkers that do not include side chain groups and linked moieties) can be used for polymerization and can be minimally hindered from growing chain ends due to viscous effects. In some embodiments, functionalization after bead synthesis can minimize exposure of species (e.g., oligonucleotides) to be loaded with potentially damaging agents (e.g., free radicals) and/or chemical environments. In some embodiments, the generated hydrogel can possess an upper critical solution temperature (UCST) that can permit temperature driven swelling and collapse of a bead. Such functionality can aid in oligonucleotide (e.g., a primer) infiltration into the bead during subsequent functionalization of the bead with the oligonucleotide. Post-production functionalization can also be useful in controlling loading ratios of species in beads, such that, for example, the variability in loading ratio is minimized. Species loading can also be performed in a batch process such that a plurality of beads can be functionalized with the species in a single batch. [0537] Reagents can be encapsulated in beads during bead generation (e.g., during polymerization of precursors). Such reagents can or cannot participate in polymerization. Such reagents can be entered into polymerization reaction mixtures such that generated beads include the reagents upon bead formation. In some embodiments, such reagents can be added to the beads after formation. Such reagents can include, for example, capture probes (e.g., oligonucleotides), reagents for a nucleic acid amplification reaction (e.g., primers, polymerases, dNTPs, co-factors (e.g., ionic co-factors), buffers) including those described herein, reagents for enzymatic reactions (e.g., enzymes, co-factors, chemical substrates, buffers), reagents for nucleic acid modification reactions such as polymerization, ligation, or digestion, and/or reagents for template preparation (e.g., tagmentation) for one or more sequencing platforms (e.g., Nextera® for Illumina®). Such reagents can include one or more enzymes described herein, including without limitation, polymerase, reverse transcriptase, restriction enzymes (e.g., endonuclease), transposase, ligase, proteinase K, DNase, etc. Such reagents can also or alternatively include one or more reagents such as lysis agents, inhibitors, inactivating agents, chelating agents, stimulus agents. Trapping of such reagents can be controlled by the polymer network density generated during polymerization of precursors, control of ionic charge within the bead (e.g., via ionic species linked to polymerized species), or by the release of other species. Encapsulated reagents can be released from a bead upon bead degradation and/or by application of a stimulus capable of releasing the reagents from the bead. In some embodiments, the beads or bead arrangements can be incubated in permeabilization reagents as described herein.

[0538] In some embodiments, the beads can also include (e.g., encapsulate or have attached thereto) a plurality of capture probes that include spatial barcodes, and the optical properties of the spatial barcodes can be used for optical detection of the beads. For example, the absorbance of light by the spatial barcodes can be used to distinguish the beads from one another. In some embodiments, a detectable label can directly or indirectly attach to a spatial barcode and provide optical detection of the bead. In some embodiments, each bead in a group of one or more beads has a unique detectable label, and detection of the unique detectable label determines the location of the spatial barcode sequence associated with the bead.

[0539] Optical properties giving rise to optical detection of beads can be due to optical properties of the bead surface (e.g., a detectable label attached to the bead), or optical properties from the bulk region of the bead (e.g., a detectable label incorporated during bead formation or an optical property of the bead itself). In some embodiments, a detectable label can be associated with a bead or one or more moieties coupled to the bead.

[0540] In some embodiments, the beads include a plurality of detectable labels. For example, a fluorescent dye can be attached to the surface of the beads and/or can be incorporated into the beads. Different intensities of the different fluorescent dyes can be used to increase the number of

optical combinations that can be used to differentiate between beads. For example, if N is the number of fluorescent dyes (e.g., between 2 and 10 fluorescent dyes, such as 4 fluorescent dyes) and M is the possible intensities for the dyes (e.g., between 2 and 50 intensities, such as 20 intensities), then $M \times N$ are the possible distinct optical combinations. In one example, 4 fluorescent dyes with 20 possible intensities can be used to generate 160,000 distinct optical combinations.

[0541] One or more optical properties of the beads or biological contents, such as cells or nuclei, can be used to distinguish the individual beads or biological contents from other beads or biological contents. In some embodiments, the beads are made optically detectable by including a detectable label having optical properties to distinguish the beads from one another.

[0542] In some embodiments, optical properties of the beads can be used for optical detection of the beads. For example, without limitation, optical properties can include absorbance, birefringence, color, fluorescence, luminosity, photosensitivity, reflectivity, refractive index, scattering, or transmittance. For example, beads can have different birefringence values based on degree of polymerization, chain length, or monomer chemistry.

[0543] In some embodiments, nanobeads, such as quantum dots or Janus beads, can be used as optical labels or components thereof. For example, a quantum dot can be attached to a spatial barcode of a bead.

[0544] Optical labels of beads can provide enhanced spectral resolution to distinguish (e.g., identify) between beads with unique spatial barcodes (e.g., beads including unique spatial barcode sequences). That is, the beads are manufactured in a way that the optical labels and the barcodes on the beads (e.g., spatial barcodes) are correlated with each other. In some aspects, the beads can be loaded into a flowcell such that beads are arrayed in a closely packed manner (e.g., single-cell resolution). Imaging can be performed, and the spatial location of the barcodes can be determined (e.g., based on information from a look-up table (LUT)). The optical labels for spatial profiling allow for quick deconvolution of bead-barcode (e.g., spatial barcode) identify.

[0545] In some examples, a lookup table (LUT) can be used to associate a property (e.g., an optical label, such as a color and/or intensity) of the bead with the barcode sequence. The property may derive from the particle (e.g., bead) or an optical label associated with the bead. The beads can be imaged to obtain optical information of the bead, including, for example, the property (e.g., color and/or intensity) of the bead or the optical label associated with the bead, and optical information of the biological sample. For example, an image can include optical information in the visible spectrum, non-visible spectrum, or both. In some embodiments, multiple images can be obtained across various optical frequencies.

[0546] In some embodiments, a first bead includes a first optical label and spatial barcodes each having a first spatial barcode sequence. A second bead includes a second optical label and spatial barcodes each having a second spatial barcode sequence. The first optical label and second optical label can be different (e.g., provided by two different fluorescent dyes or the same fluorescent dye at two different intensities). The first and second spatial barcode sequences can be different nucleic acid sequences. In some embodiments, the beads can be imaged to identify the first and second optical labels, and the first and second optical labels can then be used to associate the first and second optical labels with the first and second spatial barcode sequences, respectively. In some embodiments, the nucleic acid containing the spatial barcode can further have a capture domain for analytes (e.g., mRNA). In some embodiments, the nucleic acid (e.g., nucleic acid containing the spatial barcode) can have a unique molecular identifier, a cleavage domain, a functional domain, or combinations thereof.

[0547] In some embodiments, the optical label has a characteristic electromagnetic spectrum. As used herein, the “electromagnetic spectrum” refers to the range of frequencies of electromagnetic radiation. In some embodiments, the optical label has a characteristic absorption spectrum. As used herein, the “absorption spectrum” refers to the range of frequencies of electromagnetic radiation

that are absorbed. The “electromagnetic spectrum” or “absorption spectrum” can lead to different characteristic spectrum. In some embodiments, the peak radiation or the peak absorption occurs at 380-450 nm (Violet), 450-485 nm (Blue), 485-500 nm (Cyan), 500-565 nm (Green), 565-590 nm (Yellow), 590-625 nm (Orange), or 625-740 nm (Red). In some embodiments, the peak radiation or the peak absorption occurs around 400 nm, 460 nm, or 520 nm.

[0548] Optical labels included on the beads can identify the associated spatial barcode on the bead. Due to the relative limited diversity of optical labels it can be advantageous to limit the size of the spatial array for deconvolution. For example, the substrate can be partitioned into two or more partitions (e.g., bins). In some embodiments, the substrate can be partitioned into three or more partitions. In some embodiments, the substrate can be partitioned into four or more partitions (e.g., bins). In some embodiments, a set of beads are deposited to the partition. Within each set of beads, one or more beads (e.g., equal to or more than 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, or 5000 beads) can have an unique optical label.

[0549] In some cases, beads within the same partition can have different coordinates on the substrate. These coordinates can be determined e.g., by various imaging techniques, such as observation through microscope under an appropriate condition. In some embodiments, the beads within the same partition can share the same spatial barcode. In some embodiments, the beads (e.g., beads having capture probes with barcodes, e.g., spatial barcodes or UMI) are different from each other for different partition bins. In some embodiments, the beads having capture probes with barcodes (e.g., spatial barcodes or UMI) can have different barcodes. For example, in some cases, within each set of beads, which beads are associated with a capture probe, the capture probes on individual beads can have a unique barcode. In some cases, among all beads (e.g., within two or more sets of beads), individual beads can have capture probes with a unique barcode.

[0550] In some aspects, the present disclosure provides a substrate. The substrate can have 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900, 1000, or more than 1000 partitions (e.g., bins, or pre-defined area). The partitions can have the same shape or different shapes. In some embodiments, the substrate has only one partition (e.g., bin or pre-defined area).

[0551] In some embodiments, the first partition (e.g., the first pre-defined area, or the only bin on the substrate) can have a first set of beads. In some embodiments, at least one bead from the first set of beads comprises an optical label, and a capture probe (e.g., an oligonucleotide capture probe) comprising a barcode and a capture domain. At least one of the beads can have a unique optical label among the first set of beads. In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% of the beads in the first set of beads have a unique optical label. In some embodiments, each bead in the first set of beads has a unique optical label.

[0552] In some embodiments, the substrate can have a second partition (e.g., the second pre-defined area, or the second bin). The second partition can have a second set of beads. In some embodiments, at least one bead from the second set of beads comprises an optical label, and a capture probe (e.g., an oligonucleotide capture probe) comprising a barcode and a capture domain. At least one of the beads can have a unique optical label among the second set of beads. In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% of the beads in the second set of beads have a unique optical label. In some embodiments, each bead in the second set of beads has a unique optical label.

[0553] In some embodiments, the substrate can have a third partition, a fourth partition, a fifth partition, a sixth partition, a seventh partition, an eighth partition, a ninth partition, or a tenth partition, etc. In some embodiments, the substrate can have multiple partitions. In some cases, each of these partitions has properties that are similar to the first or the second partitions described

herein. For example, at least one bead from each set of beads comprises an optical label, and a capture probe (e.g., an oligonucleotide capture probe) comprising a barcode and a capture domain. At least one of these beads can have a unique optical label among each set of beads. In some embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% of the beads in each set of beads have a unique optical label. In some embodiments, each bead in each set of beads has a unique optical label.

[0554] In some embodiments, the beads are deposited on the substrate. In some embodiments, the beads can be deposited directly on or into a biological sample. Thus, in some cases, the biological sample can be fixed or attached on the substrate before beads are deposited onto the substrate.

[0555] In some embodiments, the beads are only deposited to areas of interest (e.g., specific locations on the substrate, specific cell types, and specific tissue structures). Thus, the deposited beads do not necessarily cover the entire biological sample. In some embodiments, one or more regions of a substrate can be masked or modified (e.g., capped capture domains) such that the masked regions do not interact with a corresponding region of the biological sample.

[0556] In some embodiments, two or more than two sets of beads (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more than 10 sets) are deposited at two or more than two partitions (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, or more than 10 partitions). These partitions do not need to be adjacent to each other. As long as the location of the partitions on the substrate is recorded, the identity of the beads can be determined from the optical labels.

[0557] In some embodiments, a set of beads can have equal to or more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, or 5000 beads. In some embodiments, a set of beads can have less than 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, or 5000 beads.

[0558] Optical labels can be included while generating the beads. For example, optical labels can be included in the polymer structure of a gel bead, or attached at the pre-polymer or monomer stage in bead production. In some embodiments, the beads include moieties that attach to one or more optical labels (e.g., at a surface of a bead and/or within a bead). In some embodiments, optical labels can be loaded into the beads with one or more reagents. For example, reagents and optical labels can be loaded into the beads by diffusion of the reagents (e.g., a solution of reagents including the optical labels). In some embodiments, optical labels can be included while preparing spatial barcodes. For example, spatial barcodes can be prepared by synthesizing molecules including barcode sequences (e.g., using a split pool or combinatorial approach). Optical labels can be attached to spatial barcodes prior to attaching the spatial barcodes to a bead. In some embodiments, optical labels can be included after attaching spatial barcodes to a bead. For example, optical labels can be attached to spatial barcodes coupled to the bead. In some embodiments, spatial barcodes or sequences thereof can be releasably or cleavably attached to the bead. Optical labels can be releasably or non-releasably attached to the bead. In some embodiments, a first bead (e.g., a bead including a plurality of spatial barcodes) can be coupled to a second bead including one or more optical labels. For example, the first bead can be covalently coupled to the second bead via a chemical bond. In some embodiments, the first bead can be non-covalently associated with the second bead.

[0559] The first and/or second bead can include a plurality of spatial barcodes. The plurality of spatial barcodes coupled to a given bead can include the same barcode sequences. Where both the first and second beads include spatial barcodes, the first and second beads can include spatial barcodes including the same barcode sequences or different barcode sequences.

[0560] Bead arrays containing captured analytes can be processed in bulk or partitioned into droplet emulsions for preparing sequencing libraries. In some embodiments, next generation sequencing reads are clustered and correlated to the spatial position of the spatial barcode on the

bead array. For example, the information can be computationally superimposed over a high-resolution image of the tissue section to identify the location(s), where the analytes were detected. [0561] In some embodiments, de-cross linking can be performed to account for de-crosslinking chemistries that may be incompatible with certain barcoding/library prep biochemistry (e.g., presence of proteases). For example, a two-step process is possible. In the first step, beads can be provided in droplets such that DNA binds to the beads after the conventional de-crosslinking chemistry is performed. In the second step, the emulsion is broken and beads collected and then re-encapsulated after washing for further processing.

[0562] In some embodiments, beads can be affixed or attached to a substrate using photochemical methods. For example, a bead can be functionalized with perfluorophenylazide silane (PFPA silane), contacted with a substrate, and then exposed to irradiation (see, e.g., Liu et al. (2006) *Journal of the American Chemical Society* 128, 14067-14072). For example, immobilization of anthraquinone-functionalized substrates (see, e.g., Koch et al. (2000) *Bioconjugate Chem.* 11, 474-483, the entire contents of which are herein incorporated by reference).

[0563] The arrays can also be prepared by bead self-assembly. Each bead can be covered with hundreds of thousands of copies of a specific oligonucleotide. In some embodiments, each bead can be covered with about 1,000 to about 1,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 1,000,000 to about 10,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 2,000,000 to about 3,000,000, about 3,000,000 to about 4,000,000, about 4,000,000 to about 5,000,000, about 5,000,000 to about 6,000,000, about 6,000,000 to about 7,000,000, about 7,000,000 to about 8,000,000, about 8,000,000 to about 9,000,000, or about 9,000,000 to about 10,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 10,000,000 to about 100,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 100,000,000 to about 1,000,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 1,000,000,000 to about 10,000,000,000 oligonucleotides. The beads can be irregularly distributed across etched substrates during the array production process. During this process, the beads can be self-assembled into arrays (e.g., on a fiber-optic bundle substrate or a silica slide substrate). In some embodiments, the beads irregularly arrive at their final location on the array. Thus, the bead location may need to be mapped or the oligonucleotides may need to be synthesized based on a predetermined pattern.

[0564] Beads can be affixed or attached to a substrate covalently, non-covalently, with adhesive, or a combination thereof. The attached beads can be, for example, layered in a monolayer, a bilayer, a trilayer, or as a cluster. As defined herein, a “monolayer” generally refers to an arrayed series of probes, beads, spots, dots, features, micro-locations, or islands that are affixed or attached to a substrate, such that the beads are arranged as one layer of single beads. In some embodiments, the beads are closely packed.

[0565] As defined herein, the phrase “substantial monolayer” or “substantially form(s) a monolayer” generally refers to (the formation of) an arrayed series of probes, beads, microspheres, spots, dots, features, micro-locations, or islands that are affixed or attached to a substrate, such that about 50% to about 99% (e.g., about 50% to about 98%) of the beads are arranged as one layer of single beads. This arrangement can be determined using a variety of methods, including microscopic imaging.

[0566] In some embodiments, the monolayer of beads is located in a predefined area on the substrate. For example, the predefined area can be partitioned with physical barriers, a photomask, divots in the substrate, or wells in the substrate.

[0567] As used herein, the term “reactive element” generally refers to a molecule or molecular moiety that can react with another molecule or molecular moiety to form a covalent bond. Reactive elements include, for example, amines, aldehydes, alkynes, azides, thiols, haloacetyls, pyridyl disulfides, hydrazides, carboxylic acids, alkoxyamines, sulfhydryls, maleimides, Michael

acceptors, hydroxyls, and active esters. Some reactive elements, for example, carboxylic acids, can be treated with one or more activating agents (e.g., acylating agents, isourea-forming agents) to increase susceptibility of the reactive element to nucleophilic attack. Non-limiting examples of activating agents include N-hydroxysuccinimide, N-hydroxysulfosuccinimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide, dicyclohexylcarbodiimide, diisopropylcarbodiimide, 1-hydroxybenzotriazole, (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate, (benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, 4-(N,N-dimethylamino)pyridine, and carbonyldiimidazole.

[0568] In some embodiments, the reactive element is bound directly to a bead. For example, hydrogel beads can be treated with an acrylic acid monomer to form acrylic acid-functionalized hydrogel beads. In some cases, the reactive element is bound indirectly to the bead via one or more linkers. As used herein, a “linker” generally refers to a multifunctional (e.g., bifunctional, trifunctional) reagent used for conjugating two or more chemical moieties. A linker can be a cleavable linker that can undergo induced dissociation. For example, the dissociation can be induced by a solvent (e.g., hydrolysis and solvolysis); by irradiation (e.g., photolysis); by an enzyme (e.g., enzymolysis); or by treatment with a solution of specific pH (e.g., pH 4, 5, 6, 7, or 8).

[0569] In some embodiments, the reactive element is bound directly to a substrate. For example, a glass slide can be coated with (3-aminopropyl)triethoxysilane. In some embodiments, the reactive element is bound indirectly to a substrate via one or more linkers.

Gel/Hydrogel Beads

[0570] In some embodiments, the bead can be a gel bead. A “gel” is a semi-rigid material permeable to liquids and gases. Exemplary gels include, but are not limited to, those having a colloidal structure, such as agarose; polymer mesh structures, such as gelatin; hydrogels; and cross-linked polymer structures, such as polyacrylamide, SFA (see, for example, U.S. Patent Application Publication No. 2011/0059865, which is incorporated herein by reference in its entirety) and PAZAM (see, for example, U.S. Patent Application Publication No. 2014/0079923, which is incorporated herein by reference in its entirety).

[0571] A gel can be formulated into various shapes and dimensions depending on the context of intended use. In some embodiments, a gel is prepared and formulated as a gel bead (e.g., a gel bead including capture probes attached or associated with the gel bead). A gel bead can be a hydrogel bead. A hydrogel bead can be formed from molecular precursors, such as a polymeric or monomeric species.

[0572] In some cases, a bead comprises a polymer or hydrogel. The polymer or hydrogel may determine one or more characteristics of the hydrogel bead, such as the volume, fluidity, porosity, rigidity, organization, or one or more other features of the hydrogel bead. In some embodiments, a hydrogel bead can include a polymer matrix (e.g., a matrix formed by polymerization or cross-linking). A polymer matrix can include one or more polymers (e.g., polymers having different functional groups or repeat units). Cross-linking can be via covalent, ionic, and/or inductive interactions, and/or physical entanglement.

[0573] A polymer or hydrogel may be formed, for example, upon cross-linking one or more cross-linkable molecules within the hydrogel bead. For example, a hydrogel may be formed upon cross-linking one or more molecules within the hydrogel bead. The hydrogel may be formed upon polymerizing a plurality of monomers within the hydrogel bead. The hydrogel may be formed upon polymerizing a plurality of polymers within the hydrogel bead. Polymeric or hydrogel precursors may be provided to the hydrogel bead and may not form a polymer or hydrogel without application of a stimulus (e.g., as described herein). In some cases, the hydrogel bead may be encapsulated within the polymer or hydrogel. Formation of a hydrogel bead may take place following one or more other changes to the cell that may be brought about by one or more other conditions.

[0574] The methods described herein may be applied to a single hydrogel bead or a plurality of hydrogel beads. A method of processing a plurality of hydrogel beads may comprise providing the

plurality of hydrogel beads within a vessel and subjecting the plurality of hydrogel beads to conditions sufficient to change one or more characteristics of the hydrogel bead. For example, plurality of hydrogel beads may be subjected to a first condition or set of conditions comprising a chemical species, and a cross-section of the hydrogel beads of the plurality of hydrogel beads may change from a first cross-section to a second cross-section, which second cross-section is less than the first cross-section. The chemical species may comprise, for example, an organic solvent such as ethanol, methanol, or acetone. The plurality of hydrogel beads may then be subjected to a second condition or set of conditions comprising a chemical species, and crosslinks may form within each of the hydrogel beads. The chemical species may comprise, for example, a cross-linking agent. The plurality of processed hydrogel beads may be provided in an aqueous fluid. In some instances, the second cross-section of the plurality of hydrogel beads is substantially maintained in the aqueous fluid. The plurality of processed hydrogel beads may be partitioned within a plurality of partitions. The partitions may be, for example, aqueous droplets included in a water-in-oil emulsion. The partitions may be, for example, a plurality of wells. The plurality of fixed hydrogel beads may be co-partitioned with one or more reagents. In some cases, the plurality of fixed hydrogel beads may be co-partitioned with one or more beads, where each bead comprises a plurality of nucleic acid barcode molecules attached thereto. The nucleic acid barcode molecules attached to a given bead may comprise a common barcode sequence, and the nucleic acid barcode molecules attached to each different bead may comprise a sequence comprising a different common barcode sequence. The nucleic acid barcode molecules, or portions thereof, may then be used in reactions with target molecules associated with hydrogel beads of the plurality of hydrogel beads.

Core/Shell Beads

[0575] In some embodiments, the bead is a core/shell bead that comprises an inner core (e.g., a nanosphere or microsphere) and an outer shell (e.g., a hydrogel coating the nanosphere or microsphere). In some embodiments, the inner core can be a solid nanoparticle or solid microparticle. In some embodiments, the inner core can be a silica inner core (e.g., a silica nanoparticle or silica microparticle). In some embodiments, the inner core of the core/shell bead can have an average diameter of about 1 micron. In some embodiments, the inner core can have an average diameter of about 2 microns. In some embodiments, the inner core can have an average diameter of about 3 microns. In some embodiments, the inner core can have an average diameter of about 4 microns. In some embodiments, the inner core can have an average diameter of about 5 microns. In some embodiments, the inner core can have an average diameter of about 6 microns. In some embodiments, the inner core can have an average diameter of about 7 microns. In some embodiments, the inner core can have an average diameter of about 8 microns. In some embodiments, the inner core can have an average diameter of about 9 microns. In some embodiments, the inner core can have an average diameter of about 10 microns. In some embodiments, the inner core can have an average diameter of about 100 nanometers to about 10 microns.

[0576] In some embodiments, the core/shell bead can decrease its outer shell volume by removing solvents, salts, or water (e.g., dehydrated, desiccated, dried, exsiccated) from the outer shell to form a shrunken core/shell bead. In another example, the core/shell bead can decrease its outer shell volume by adjusting temperature or pH, as described above. In some embodiments, the core/shell bead can expand its outer shell volume, for example by the addition of solvents, salts, or water (e.g., rehydration) to form an expanded core/shell bead. In some embodiments, the outer shell (e.g., coating the inner core) can have an average thickness of about 1 micron. In some embodiments, the outer shell can have an average thickness of about 2 microns. In some embodiments, the outer shell can have an average thickness of about 3 microns. In some embodiments, the outer shell can have an average thickness of about 4 microns. In some embodiments, the outer shell can have an average thickness of about 5 microns.

[0577] In some embodiments, the core/shell bead can have an average diameter of about 1 micron

to about 10 microns. In some embodiments, the core/shell bead can have an average diameter of about 1 micron. In some embodiments, the core/shell bead can have an average diameter of about 2 microns. In some embodiments, the core/shell bead can have an average diameter of about 3 microns. In some embodiments, the core/shell bead can have an average diameter of about 4 microns. In some embodiments, the core/shell bead can have an average diameter of about 5 microns. In some embodiments, the core/shell bead can have an average diameter of about 6 microns. In some embodiments, the core/shell bead can have an average diameter of about 7 microns. In some embodiments, the core/shell bead can have an average diameter of about 8 microns. In some embodiments, the core/shell bead can have an average diameter of about 9 microns. In some embodiments, the core/shell bead can have an average diameter of about 10 microns.

(2) Methods for Covalently Bonding Features to a Substrate

[0578] Provided herein are methods for the covalent bonding of features (e.g., optically labeled beads, hydrogel beads, microsphere beads) to a substrate.

[0579] In some embodiments, the features (e.g., beads) are coupled to a substrate via a covalent bond between a first reactive element and a second reactive element. In some embodiments, the covalently-bound beads substantially form a monolayer of features (e.g., hydrogel beads, microsphere beads) on the substrate.

[0580] In some embodiments, the features (e.g., beads) are functionalized with a first reactive element, which is directly bound to the features. In some embodiments, the features are functionalized with a first reactive element, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.

[0581] In some embodiments, the substrate is functionalized with a second reactive element, which is directly bound to the substrate. In some embodiments, the substrate is functionalized with a second reactive element, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. For example, the linker can be benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.

[0582] In some embodiments, the substrate is a glass slide. In some embodiments, the substrate is a pre-functionalized glass slide.

[0583] In some embodiments, about 99% of the covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 50% to about 98% form a monolayer of beads on the substrate. For example, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 65%, about 50% to about 60%, or about 50% to about 55% of the covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 98%, about 60% to about 98%, about 65% to about 98%, about 70% to about 98%, about 75% to about 98%, about 80% to about 98%, about 85% to about 98%, about 90% to about 95%, or about 95% to about 98% of the covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 95%, about 60% to about 90%, about 65% to about 95%, about 70% to about 95%, about 75% to about 90%, about 75% to about 95%, about 80% to about 90%, about 80% to about 95%, about 85% to about 90%, or about 85% to about 95% of the covalently-bound beads form a monolayer of beads on the substrate.

[0584] In some embodiments, at least one of the first reactive element and the second reactive element is selected from the group consisting of:

##STR00011##

wherein [0585] R_{sup.1} is selected from H, C_{sub.1}-C_{sub.6} alkyl, or —SO_{sub.3}; [0586] R_{sup.2} is

C.sub.1-C.sub.6 alkyl; and [0587] X is a halo moiety.

[0588] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00012## [0589] wherein the  custom-character indicates the point of attachment of the first reactive element or the second reactive element to the bead (e.g., hydrogel bead or microsphere bead) or to the substrate.

[0590] In some embodiments, at least one of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00013##

wherein [0591] R.sup.1 is selected from H, C.sub.1-C.sub.6 alkyl, or —SO.sub.3; [0592] R.sup.2 is C.sub.1-C.sub.6 alkyl; and [0593] X is a halo moiety.

[0594] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00014##

wherein R.sup.1 is selected from H, C.sub.1-C.sub.6 alkyl, or —SO.sub.3. In some embodiments, R.sup.1 is H. In some embodiments, R.sup.1 is C.sub.1-C.sub.6 alkyl. In some embodiments, R.sup.1 is —SO.sub.3.

[0595] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00015##

wherein R.sup.2 is C.sub.1-C.sub.6 alkyl. In some embodiments, R.sup.2 is methyl.

[0596] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00016##

In some embodiments,

##STR00017##

can be reacted with an activating agent to form an active ester. In some embodiments, the active ester is

##STR00018##

In some embodiments, the activating agent is an acylating agent (e.g., N-hydroxysuccinimide and N-hydroxysulfosuccinimide). In some embodiments, the activating agent is an O-acylisourea-forming agent (e.g., 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), dicyclohexylcarbodiimide, and diisopropylcarbodiimide). In some embodiments, the activating agent is a combination of at least one acylating agent and at least one O-isourea-forming agents (e.g., N-hydroxysuccinimide (NHS), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), N-hydroxysulfosuccinimide (sulfo-NHS), and a combination thereof).

[0597] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00019##

[0598] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00020##

wherein X is a halo moiety. For example, X is chloro, bromo, or iodo.

[0599] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00021##

[0600] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00022##

[0601] In some embodiments, at least one of the first reactive element or the second reactive

element comprises

##STR00023##

[0602] In some embodiments, at least one of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00024##

wherein [0603] R^{sup.3} is H or C_{sub.1}-C_{sub.6} alkyl; and [0604] R^{sup.4} is H or trimethylsilyl.

[0605] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00025##

wherein R^{sup.4} is H or trimethylsilyl. In some embodiments, R^{sup.4} is H.

[0606] In some embodiments, at least one of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00026##

wherein R^{sup.3} is H or C_{sub.1}-C_{sub.6} alkyl. In some embodiments, R^{sup.3} is H. In some embodiments, R^{sup.3} is C_{sub.1}-C_{sub.6} alkyl.

[0607] In some embodiments, at least one of the first reactive element or the second reactive element comprises

##STR00027##

wherein R^{sup.3} is H or C_{sub.1}-C_{sub.6} alkyl. In some embodiments, R^{sup.3} is H. In some embodiments, R^{sup.3} is C_{sub.1}-C_{sub.6} alkyl.

[0608] In some embodiments, at least one of the first reactive elements or the second reactive elements comprises

##STR00028##

In some embodiments, at least one of the first reactive elements or the second reactive elements comprises

##STR00029##

[0609] In some embodiments, one of the first reactive elements or the second reactive elements is selected from the group consisting of:

##STR00030##

wherein [0610] R^{sup.1} is selected from H, C_{sub.1}-C_{sub.6} alkyl, or —SO_{sub.3}; [0611] R^{sup.2} is C_{sub.1}-C_{sub.6} alkyl; [0612] X is a halo moiety;

and the other of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00031##

wherein [0613] R^{sup.3} is H or C_{sub.1}-C_{sub.6} alkyl; and [0614] R^{sup.4} is H or trimethylsilyl.

[0615] In some embodiments, one of the first reactive elements or the second reactive elements is selected from the group consisting of

##STR00032##

wherein R^{sup.3} is H or C_{sub.1}-C_{sub.6} alkyl;

##STR00033##

and the other of the first reactive element or the second reactive element is wherein R^{sup.4} is H or trimethylsilyl. In some embodiments, R^{sup.3} is H. In some embodiments, R^{sup.3} is C_{sub.1}-C_{sub.6} alkyl. In some embodiments, R^{sup.4} is H. In some embodiments, R^{sup.4} is trimethylsilyl.

[0616] In some embodiments, one of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00034##

wherein [0617] R^{sup.1} is selected from H, C_{sub.1}-C_{sub.6} alkyl, or —SO_{sub.3}; [0618] R^{sup.2} is C_{sub.1}-C_{sub.6} alkyl; [0619] X is a halo moiety;

and the other of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00035##

[0620] wherein R.sup.3 is H or C.sub.1-C.sub.6 alkyl. In some embodiments, R.sup.1 is H. In some embodiments, R.sup.1 is C.sub.1-C.sub.6 alkyl. In some embodiments, R.sup.1 is —SO.sub.3. In some embodiments, R.sup.2 is methyl. In some embodiments, X is iodo. In some embodiments, R.sup.3 is H. In some embodiments, R.sup.3 is C.sub.1-C.sub.6 alkyl.

[0621] In some embodiments, one of the first reactive elements or the second reactive elements is selected from the group consisting of:

##STR00036##

wherein

[0622] R.sup.1 is selected from H, C.sub.1-C.sub.6 alkyl, or —SO.sub.3; [0623] R.sup.2 is C.sub.1-C.sub.6 alkyl;

and the other of the first reactive elements or the second reactive elements comprises

##STR00037##

wherein R.sup.3 is H or C.sub.1-C.sub.6 alkyl. In some embodiments, R.sup.1 is H. In some embodiments, R.sup.1 is C.sub.1-C.sub.6 alkyl. In some embodiments, R.sup.1 is —SO.sub.3. In some embodiments, R.sup.2 is methyl. In some embodiments, R.sup.3 is H. In some embodiments, R.sup.3 is C.sub.1-C.sub.6 alkyl.

[0624] In some embodiments, one of the first reactive element or the second reactive element is selected from the group consisting of:

##STR00038## [0625] wherein X is a halo moiety; [0626] and the other of the first reactive element or the second reactive element comprises

##STR00039##

In some embodiments, X is bromo. In some embodiments, X is iodo.

[0627] In some embodiments, one of the first reactive element or the second reactive element is selected from the group consisting of

##STR00040##

and the other of the first reactive element or the second reactive element comprises

##STR00041##

[0628] The term “halo” refers to fluoro (F), chloro (Cl), bromo (Br), or iodo (I).

[0629] The term “alkyl” refers to a hydrocarbon chain that may be a straight chain or branched chain, containing the indicated number of carbon atoms. For example, C.sub.1-10 indicates that the group may have from 1 to 10 (inclusive) carbon atoms in it. Non-limiting examples include methyl, ethyl, iso-propyl, tert-butyl, n-hexyl.

[0630] The term “haloalkyl” refers to an alkyl, in which one or more hydrogen atoms is/are replaced with an independently selected halo.

[0631] The term “alkoxy” refers to an —O-alkyl radical (e.g., —OCH.sub.3).

[0632] The term “alkylene” refers to a divalent alkyl (e.g., —CH.sub.2—).

[0633] The term “alkenyl” refers to a hydrocarbon chain that may be a straight chain or branched chain having one or more carbon-carbon double bonds. The alkenyl moiety contains the indicated number of carbon atoms. For example, C.sub.2-6 indicates that the group may have from 2 to 6 (inclusive) carbon atoms in it.

[0634] The term “alkynyl” refers to a hydrocarbon chain that may be a straight chain or branched chain having one or more carbon-carbon triple bonds. The alkynyl moiety contains the indicated number of carbon atoms. For example, C.sub.2-6 indicates that the group may have from 2 to 6 (inclusive) carbon atoms in it.

[0635] The term “aryl” refers to a 6-20 carbon mono-, bi-, tri- or polycyclic group wherein at least one ring in the system is aromatic (e.g., 6-carbon monocyclic, 10-carbon bicyclic, or 14-carbon tricyclic aromatic ring system); and wherein 0, 1, 2, 3, or 4 atoms of each ring may be substituted by a substituent. Examples of aryl groups include phenyl, naphthyl, tetrahydronaphthyl, and the like.

(3) Methods for Non-Covalently Bonding Features to a Substrate

[0636] Provided herein are methods for the non-covalent bonding of features (e.g., optically-labeled beads, hydrogel beads, or microsphere beads) to a substrate.

[0637] In some embodiments, features (e.g., beads) are coupled to a substrate via a non-covalent bond between a first affinity group and a second affinity group. In some embodiments, the non-covalently-bound features (e.g., beads) substantially form a monolayer of beads (e.g., hydrogel beads, microsphere beads) on the substrate.

[0638] In some embodiments, the features are functionalized with a first affinity group, which is directly bound to the features. In some embodiments, the features are functionalized with a first affinity group, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.

[0639] In some embodiments, the substrate is functionalized with a second affinity group, which is directly bound to the substrate. In some embodiments, the substrate is functionalized with a second affinity group, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.

[0640] In some embodiments the first affinity group or the second affinity group is biotin, and the other of the first affinity group or the second affinity group is streptavidin.

[0641] In some embodiments, about 99% of the non-covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 50% to about 98% form a monolayer of beads on the substrate. For example, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 65%, about 50% to about 60%, or about 50% to about 55% of the non-covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 98%, about 60% to about 98%, about 65% to about 98%, about 70% to about 98%, about 75% to about 98%, about 80% to about 98%, about 85% to about 98%, about 90% to about 95%, or about 95% to about 98% of the non-covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 95%, about 60% to about 90%, about 65% to about 95%, about 70% to about 95%, about 75% to about 90%, about 75% to about 95%, about 80% to about 90%, about 80% to about 95%, about 85% to about 90%, or about 85% to about 95% of the non-covalently-bound beads form a monolayer of beads on the substrate.

[0642] In some embodiments, the monolayer of beads is formed in a predefined area on the substrate. In some embodiments, the predefined area is partitioned with physical barriers. For example, divots or wells in the substrate. In some embodiments, the predefined area is partitioned using a photomask. For example, the substrate is coated with a photo-activated solution, dried, and then irradiated under a photomask. In some embodiments, the photo-activated solution is UV-activated.

[0643] As used herein, an “adhesive” generally refers to a substance used for sticking objects or materials together. Adhesives include, for example, glues, pastes, liquid tapes, epoxy, bioadhesives, gels, and mucilage. In some embodiments, an adhesive is liquid tape. In some embodiments, the adhesive is glue.

[0644] In some embodiments, beads are adhered to a substrate using an adhesive (e.g., liquid tape, glue, paste). In some embodiments, the adhered beads substantially form a monolayer of beads on the substrate (e.g., a glass slide). In some embodiments, the beads are hydrogel beads. In some embodiments, the beads are microsphere beads. In some embodiments, the beads are coated with the adhesive, and then the beads are contacted with the substrate. In some embodiments, the substrate is coated with the adhesive, and then the substrate is contacted with the beads. In some

embodiments, both the substrate is coated with the adhesive and the beads are coated with the adhesive, and then the beads and substrate are contacted with one another.

[0645] In some embodiments, about 99% of the adhered beads form a monolayer of beads on the substrate. In some embodiments, about 50% to about 98% form a monolayer of beads on the substrate. For example, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 65%, about 50% to about 60%, or about 50% to about 55% of the adhered beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 98%, about 60% to about 98%, about 65% to about 98%, about 70% to about 98%, about 75% to about 98%, about 80% to about 98%, about 85% to about 98%, about 90% to about 95%, or about 95% to about 98% of the adhered beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 95%, about 60% to about 90%, about 65% to about 95%, about 70% to about 95%, about 75% to about 90%, about 75% to about 95%, about 80% to about 90%, about 80% to about 95%, about 85% to about 90%, or about 85% to about 95% of the adhered beads form a monolayer of beads on the substrate.

[0646] In some embodiments, beads can be deposited onto a biological sample such that the deposited beads form a monolayer of beads on the biological sample (e.g., over or under the biological sample). In some embodiments, beads deposited on the substrate can self-assemble into a monolayer of beads that saturate the intended surface area of the biological sample under investigation. In this approach, bead arrays can be designed, formulated, and prepared to evaluate a plurality of analytes from a biological sample of any size or dimension. In some embodiments, the concentration or density of beads (e.g., gel beads) applied to the biological sample is such that the area as a whole, or one or more regions of interest in the biological sample, is saturated with a monolayer of beads. In some embodiments, the beads are contacted with the biological sample by pouring, pipetting, spraying, and the like, onto the biological sample. Any suitable form of bead deposition can be used.

[0647] In some embodiments, the biological sample can be confined to a specific region or area of the array. For example, a biological sample can be affixed to a glass slide and a chamber, gasket, or cage positioned over the biological sample to act as a containment region or frame within which the beads are deposited. As will be apparent, the density or concentration of beads needed to saturate an area or biological sample can be readily determined by one of ordinary skill in the art (e.g., through microscopic visualization of the beads on the biological sample). In some embodiments, the bead array contains microfluidic channels to direct reagents to the spots or beads of the array.

(4) Feature Geometric Attributes

[0648] Features on an array can have a variety of sizes. In some embodiments, a feature of an array can have an average diameter or maximum dimension between 500 nm to 100 μ m. For example, between 500 nm to 2 μ m, 1 μ m to 3 μ m, 1 μ m to 5 μ m, 1 μ m to 10 μ m, 1 μ m to 20 μ m, 1 μ m to 30 μ m, 1 μ m to 40 μ m, 1 μ m to 50 μ m, 1 μ m to 60 μ m, 1 μ m to 70 μ m, 1 μ m to 80 μ m, 1 μ m to 90 μ m, 90 μ m to 100 μ m, 80 μ m to 100 μ m, 70 μ m to 100 μ m, 60 μ m to 100 μ m, 50 μ m to 100 μ m, 40 μ m to 100 μ m, 30 μ m to 100 μ m, 20 μ m to 100 μ m, 10 μ m to 100 μ m, about 40 μ m to about 70 μ m, or about 50 μ m to about 60 μ m. In some embodiments, the feature has an average diameter or maximum dimension between 30 μ m to 100 μ m, 40 μ m to 90 μ m, 50 μ m to 80 μ m, 60 μ m to 70 μ m, or any range within the disclosed sub-ranges. In some embodiments, the feature has an average diameter or maximum dimension no larger than 95 μ m, 90 μ m, 85 μ m, 80 μ m, 75 μ m, 70 μ m, 65 μ m, 60 μ m, 55 μ m, 50 μ m, 45 μ m, 40 μ m, 35 μ m, 30 μ m, 25 μ m, 20 μ m, 15 μ m, 14 μ m, 13 μ m, 12 μ m, 11 μ m, 10 μ m, 9 μ m, 8 μ m, 7 μ m, 6 μ m, 5 μ m, 4 μ m, 3 μ m, 2 μ m, or 1 μ m. In some embodiments, the feature has an average diameter or maximum dimension of approximately 65 μ m. In some embodiments, the feature has an average diameter or maximum distance of approximately 55 μ m.

[0649] In some embodiments, the size and/or shape of a plurality of features of an array are

approximately uniform. In some embodiments, the size and/or shape of a plurality of features of an array is not uniform. For example, in some embodiments, features in an array can have an average cross-sectional dimension, and a distribution of cross-sectional dimensions among the features can have a full-width and half-maximum value of 0% or more (e.g., 5% or more, 10% or more, 20% or more, 30% or more, 40% or more, 50% or more, 70% or more, or 100% or more) of the average cross-sectional dimension for the distribution.

[0650] In certain embodiments, features in an array can have an average cross-sectional dimension of between about 1 μm and about 10 μm . This range in average feature cross-sectional dimension corresponds to the approximate diameter of a single mammalian cell. Thus, an array of such features can be used to detect analytes at, or below, mammalian single-cell resolution.

[0651] In some embodiments, a plurality of features has a mean diameter or mean maximum dimension of about 0.1 μm to about 100 μm (e.g., about 0.1 μm to about 5 μm , about 1 μm to about 10 μm , about 1 μm to about 20 μm , about 1 μm to about 30 μm , about 1 μm to about 40 μm , about 1 μm to about 50 μm , about 1 μm to about 60 μm , about 1 μm to about 70 μm , about 1 μm to about 80 μm , about 1 μm to about 90 μm , about 90 μm to about 100 μm , about 80 μm to about 100 μm , about 70 μm to about 100 μm , about 60 μm to about 100 μm , about 50 μm to about 100 μm , about 40 μm to about 100 μm , about 30 μm to about 100 μm , about 20 μm to about 100 μm , or about 10 μm to about 100 μm). In some embodiments, the plurality of features has a mean diameter or mean maximum dimension between 30 μm to 100 μm , 40 μm to 90 μm , 50 μm to 80 μm , 60 μm to 70 μm , or any range within the disclosed sub-ranges. In some embodiments, the plurality of features has a mean diameter or a mean maximum dimension no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm . In some embodiments, the plurality of features has a mean average diameter or a mean maximum dimension of approximately 65 μm , approximately 60 μm , approximately 55 μm , approximately 50 μm , approximately 45 μm , approximately 40 μm , approximately 35 μm , approximately 30 μm , approximately 25 μm , approximately 20 μm , approximately 15 μm , approximately 10 μm , approximately 5 μm , approximately 4 μm , approximately 3 μm , approximately 2 μm , or approximately 1 μm .

(iv) Array Geometric Attributes

[0652] In some embodiments, an array includes a plurality of features. For example, an array includes between 4,000 and 50,000 features, or any range within 4,000 to 40,000 features. For example, an array includes between 4,000 to 35,000 features, 4,000 to 30,000 features, 4,000 to 25,000 features, 4,000 to 20,000 features, 4,000 to 15,000 features, 4,000 to 10,000 features, 4,000 to 6,000 features, or 4,400 to 6,000 features. In some embodiments, the array includes between 4,100 and 5,900 features, between 4,200 and 5,800 features, between 4,300 and 5,700 features, between 4,400 and 5,600 features, between 4,500 and 5,500 features, between 4,600 and 5,400 features, between 4,700 and 5,300 features, between 4,800 and 5,200 features, between 4,900 and 5,100 features, or any range within the disclosed sub-ranges. For example, the array can include about 4,000 features, about 4,200 features, about 4,400 features, about 4,800 features, about 5,000 features, about 5,200 features, about 5,400 features, about 5,600 features, or about 6,000 features, about 10,000 features, about 20,000 features, about 30,000 features, about 40,000 features, or about 50,000 features. In some embodiments, the array comprises at least 4,000 features. In some embodiments, the array includes approximately 5,000 features.

[0653] In some embodiments, features within an array have an irregular arrangement or relationship to one another, such that no discernable pattern or regularity is evident in the geometrical spacing relationships among the features. For example, features within an array may be positioned randomly with respect to one another. Alternatively, features within an array may be positioned irregularly, but the spacings may be selected deterministically to ensure that the resulting arrangement of features is irregular.

[0654] In some embodiments, features within an array are positioned regularly with respect to one another to form a pattern. A wide variety of different patterns of features can be implemented in arrays. Examples of such patterns include, but are not limited to, square arrays of features, rectangular arrays of features, hexagonal arrays of features (including hexagonal close-packed arrays), radial arrays of features, spiral arrays of features, triangular arrays of features, and more generally, any array in which adjacent features in the array are reached from one another by regular increments in linear and/or angular coordinate dimensions.

[0655] In some embodiments, features within an array are positioned with a degree of regularity with respect to one another such that the array of features is neither perfectly regular nor perfectly irregular (i.e., the array is “partially regular”). For example, in some embodiments, adjacent features in an array can be separated by a displacement in one or more linear and/or angular coordinate dimensions that is 10% or more (e.g., 20% or more, 30% or more, 40% or more, 50% or more, 60% or more, 70% or more, 80% or more, 90% or more, 100% or more, 110% or more, 120% or more, 130% or more, 140% or more, 150% or more, 160% or more, 170% or more, 180% or more, 190% or more, 200% or more) of an average displacement or a nominal displacement between adjacent features in the array. In certain embodiments, the distribution of displacements (linear and/or angular) between adjacent features in an array has a full-width at half-maximum of between 0% and 200% (e.g., between 0% and 100%, between 0% and 75%, between 0% and 50%, between 0% and 25%, between 0% and 15%, between 0% and 10%) of an average displacement or nominal displacement between adjacent features in the array.

[0656] In some embodiments, arrays of features can have a variable geometry. For example, a first subset of features in an array can be arranged according to a first geometrical pattern, and a second subset of features in the array can be arranged according to a second geometrical pattern that is different from the first pattern. Any of the patterns described above can correspond to the first and/or second geometrical patterns, for example.

[0657] In general, arrays of different feature densities can be prepared by adjusting the spacing between adjacent features in the array. In some embodiments, the geometric center-to-center (e.g., pitch) spacing between adjacent features in an array is between 100 nm to 10 μ m, 500 nm to 2 μ m, 1 μ m to 5 μ m, and 20 μ m to 200 μ m. For example, the center-to-center spacing can be between 100 nm to 10 μ m, 500 nm to 2 μ m, 1 μ m, to 5 μ m, 20 μ m to 40 μ m, 20 μ m to 60 μ m, 20 μ m to 80 μ m, 80 μ m to 100 μ m, 100 μ m to 120 μ m, 120 μ m to 140 μ m, 140 μ m to 160 μ m, 160 μ m to 180 μ m, 180 μ m to 200 μ m, 60 μ m to 100 μ m, or 40 μ m to 100 μ m, 50 μ m to 150 μ m, 80 μ m to 120 μ m, or 90 μ m to 110 μ m. In some embodiments, the pitch between adjacent array features is between 30 μ m and 100 μ m, 40 μ m and 90 μ m, 50 μ m and 80 μ m, 60 μ m and 70 μ m, 80 μ m and 120 μ m, or any range within the disclosed sub-ranges. In some embodiments, the pitch between adjacent array features of an array is approximately 65 μ m, approximately 60 μ m, approximately 55 μ m, approximately 50 μ m, approximately 45 μ m, approximately 40 μ m, approximately 35 μ m, approximately 30 μ m, approximately 25 μ m, approximately 20 μ m, approximately 15 μ m, approximately 10 μ m, approximately 5 μ m, approximately 4 μ m, approximately 3 μ m, approximately 2 μ m, or approximately 1 μ m. In some embodiments, the pitch between adjacent array features of an array is less than 100 μ m.

[0658] An array of features can have any appropriate resolution. In some embodiments, an array of features can have a spatially constant (e.g., within a margin of error) resolution. In general, an array with a spatially consistent resolution is an array in which the pitch between adjacent features in the array is constant (e.g., within a margin of error). Such arrays can be useful in a variety of applications. In some embodiments, an array of features can have a spatially varying resolution. In general, an array with a spatially varying resolution is an array in which the center-to-center spacing (e.g., pitch) (along linear, angular, or both linear and angular coordinate dimensions) between adjacent features in the array varies. Such arrays can be useful in a variety of applications. For example, in some embodiments, depending upon the spatial resolution at which the sample is

to be investigated, the sample can be selectively associated with the portion of the array that corresponds approximately to the desired spatial resolution of the measurement.

[0659] In some embodiments, it may be useful to describe the resolution of an array of features by functional aspects, for example, the number of reads that can be carried out per feature (which can be a proxy for sequencing saturation), the number of transcripts that can be detected per feature, or the number of genes that can be detected per feature. For example, in some embodiments, the number of reads that can be performed per feature is between 50,000 and 1,000,000. For example, the number of reads that can be performed per feature can be between 50,000 and 100,000, 50,000 and 150,000, 50,000 and 200,000, 50,000 and 250,000, 50,000 and 300,000, 50,000 and 350,000, 50,000 and 400,000, 50,000 and 500,000, 50,000 and 550,000, 50,000 and 600,000, 50,000 and 650,000, 50,000 and 700,000, 50,000 and 750,000, 50,000 and 800,000, 50,000 and 850,000, 50,000 and 900,000, 50,000 and 950,000, 50,000 and 1,000,000, 100,000 to 500,000, 150,000 to 500,000, 200,000 to 500,000, 250,000 to 500,000, 300,000 and 500,000, 350,000 and 500,000, 400,000 and 500,000, 450,000 and 500,000, 150,000 to 250,000, or 300,000 to 400,000. In some embodiments, the number of reads that can be performed per feature is about 70,000. In some embodiments, the number of reads that can be performed per feature is about 170,000. In some embodiments, the number reads that can be performed per feature is about 330,000. In some embodiments, the number reads that can be performed per feature is about 500,000. In some embodiments, the number reads that can be performed per feature is about 800,000.

[0660] In some embodiments, the number of transcripts that can be detected per feature is between 20,000 and 200,000. For example, in some embodiments, the number of transcripts that can be detected per feature can be between 20,000 and 30,000, 20,000 and 40,000, 20,000 and 50,000, 30,000 and 60,000, 40,000 and 60,000, 50,000 and 60,000, 20,000 and 100,000, 30,000 and 100,000, 40,000 and 200,000, 50,000 and 200,000, or 30,000 and 200,000. In some embodiments, the number of transcripts that can be detected per feature is about 40,000. In some embodiments, the number of transcripts that can be detected per feature is about 60,000. In some embodiments, the number of transcripts that can be detected per feature is about 80,000. In some embodiments, the number of transcripts that can be detected per feature is about 100,000.

[0661] In some embodiments, the number of genes that can be detected per feature is between 1,000 and 5,000. For example, the number of genes that can be detected per feature can be between 1,000 and 1,500, 1,000 and 2,000, 1,000 and 2,500, 1,000 and 3,000, 1,000 and 3,500, 1,000 and 4,000, 1,000 and 4,500, 1,500 and 5,000, 2,000 and 5,000, 2,500 and 5,000, 3,000 and 5,000, 3,500 and 5,000, 4,000 and 5,000, 4,500 and 5,000, 1,500 and 2,500, 2,500 and 3,500, or 3,500 and 4,000. In some embodiments, the number of genes that can be detected per feature is about 2,000. In some embodiments, the number of genes that can be detected per feature is about 3,000. In some embodiments, the number of genes that can be detected per feature is about 4,000.

[0662] In some embodiments, it may be useful to describe the resolution of an array of features by functional aspects, for example, the number of UMI counts per feature. For example, in some embodiments, the number of UMI counts that can be performed per feature is between 1,000 and 50,000. In some embodiments, the number of UMI counts can be averaged to obtain a mean UMI per feature. In some embodiments, the number of UMI counts can be averaged to obtain a median UMI count per feature. For example, the median UMI count per feature can be between 1,000 and 50,000, 1,000 and 40,000, 1,000 and 30,000, 1,000 and 20,000, 1,000 and 10,000, 1,000 and 5,000. In some embodiments, the median UMI count per feature is about 5,000. In some embodiments, the median UMI count per feature is about 10,000.

[0663] These components can be used to determine the sequencing saturation of the array. The sequencing saturation can be a measure of the library complexity and sequencing depth. For example, different cell types will have different amounts of RNA, thus different number of transcripts, influencing library complexity. Additionally, sequencing depth is related to the number of sequencing reads. In some embodiments, the inverse of sequencing saturation is the number of

additional reads it would take to detect a new transcript. One way of measuring the sequencing saturation of an array is to determine the number of reads to detect a new UMI. For example, if a new UMI is detected every 2 reads of the feature, the sequencing saturation would be 50%. As another example, if a new UMI is detected every 10 reads of a feature, the sequencing saturation would be 90%.

[0664] Arrays of spatially varying resolution can be implemented in a variety of ways. In some embodiments, for example, the pitch between adjacent features in the array varies continuously along one or more linear and/or angular coordinate directions. Thus, for a rectangular array, the spacing between successive rows of features, between successive columns of features, or between both successive rows and successive columns of features, can vary continuously.

[0665] In certain embodiments, arrays of spatially varying resolution can include discrete domains with populations of features. Within each domain, adjacent features can have a regular pitch. Thus, for example, an array can include a first domain within which adjacent features are spaced from one another along linear and/or angular coordinate dimensions by a first set of uniform coordinate displacements, and a second domain within which adjacent features are spaced from one another along linear and/or angular coordinate dimensions by a second set of uniform coordinate displacements. The first and second sets of displacements differ in at least one coordinate displacement, such that adjacent features in the two domains are spaced differently, and the resolution of the array in the first domain is therefore different from the resolution of the array in the second domain.

[0666] In some embodiments, the pitch of array features can be sufficiently small such that array features are effectively positioned continuously or nearly continuously along one or more array dimensions, with little or no displacement between array features along those dimensions. For example, in a feature array where the features correspond to regions of a substrate (i.e., oligonucleotides are directly bound to the substrate), the displacement between adjacent oligonucleotides can be very small-effectively, the molecular width of a single oligonucleotide. In such embodiments, each oligonucleotide can include a distinct spatial barcode such that the spatial location of each oligonucleotide in the array can be determined during sample analysis. Arrays of this type can have very high spatial resolution, but may only include a single oligonucleotide corresponding to each distinct spatial location in a sample.

[0667] In general, the size of the array (which corresponds to the maximum dimension of the smallest boundary that encloses all features in the array along one coordinate direction) can be selected as desired, based on criteria such as the size of the sample, the feature diameter, and the density of capture probes within each feature. For example, in some embodiments, the array can be a rectangular or square array for which the maximum array dimension along each coordinate direction is 10 mm or less (e.g., 9 mm or less, 8 mm or less, 7 mm or less, 6 mm or less, 5 mm or less, 4 mm or less, 3 mm or less). Thus, for example, a square array of features can have dimensions of 8 mm by 8 mm, 7 mm by 7 mm, 5 mm by 5 mm, or be smaller than 5 mm by 5 mm.

(v) Bead Arrays

[0668] As used herein, the term “bead array” refers to an array that includes a plurality of beads as the features in the array. In some embodiments, two or more beads are dispersed onto a substrate to create an array, where each bead is a feature on the array. In some embodiments, the beads are attached to a substrate. For example, the beads can optionally attach to a substrate such as a microscope slide and in proximity to a biological sample (e.g., a tissue section that includes cells). The beads can also be suspended in a solution and deposited on a surface (e.g., a membrane, a tissue section, or a substrate (e.g., a microscope slide)). Beads can optionally be dispersed into wells on a substrate, e.g., such that only a single bead is accommodated per well.

[0669] Examples of arrays of beads on or within a substrate include beads located in wells such as the BeadChip array for microarray genotyping (available from Illumina Inc., San Diego, CA), and array used in sequencing platforms from Ion Torrent (a subsidiary of Life Technologies, Carlsbad,

CA). Examples of bead arrays are described in, e.g., U.S. Pat. Nos. 6,266,459; 6,355,431; 6,770,441; 6,859,570; 6,210,891; 6,258,568; and 6,274,320; U.S. Pat. Application Publication Nos. 2009/0026082; 2009/0127589; 2010/0137143; 2019/0177777; and 2010/0282617; and PCT Patent Application Publication Nos. WO 00/063437 and WO 2016/162309, the entire contents of each of which is incorporated herein by reference.

[0670] In some embodiments, the bead array includes a plurality of beads. For example, the bead array can include at least 10,000 beads (e.g., at least 100,000 beads, at least 1,000,000 beads, at least 5,000,000 beads, at least 10,000,000 beads). In some embodiments, the plurality of beads includes a single type of bead (e.g., substantially uniform in volume, shape, and other physical properties, such as translucence). In some embodiments, the plurality of beads includes two or more types of different beads.

[0671] Bead arrays can be generated by attaching beads (e.g., barcoded beads) to a substrate in a regular pattern, or an irregular arrangement. In some embodiments, the barcode sequences are known before attaching them to the substrate. In some embodiments, the barcode sequences are not known before attaching them to the substrate. Beads can be attached to selective regions on a substrate by, e.g., selectively activating regions on the substrate to allow for attachment of the beads. Activating selective regions on the substrate can include activating or degrading a coating (e.g., a conditionally removable coating as described herein) at the selective regions where the coating has been applied on the substrate, rendering the selective regions more permissive to bead attachment as compared to regions outside of the selected regions. The regions that are rendered more permissive for bead attachment can be configured to fit only one bead or multiple beads (e.g., limited by well size or surface patterning, such as fabrication techniques). Beads bound to the selected regions can form a two-dimensional array on the substrate. The substrate can be uniformly or non-uniformly coated with the coating. The beads can be any suitable beads described herein, including beads that are attached to one or more spatial barcodes. Beads can be attached to the selected regions according to any of the methods suitable for attaching beads to substrates described herein, such as through covalent bonds, non-covalent bonds, or chemical linkers.

[0672] Any variety of suitable patterning techniques can be used to attach beads to a substrate surface. In some embodiments, in a non-limiting way, physical techniques such as inkjet printing, optical and optoelectronic cell trapping, laser-based patterning, acoustic patterning, dielectrophoresis, or magnetic techniques can be used to pattern the substrate. Alternatively, chemical and/or physiochemical techniques can be used such as, in a non-limiting way, surface chemistry methods, microcontact printing, microwells and filtration, DUV patterning, or patterning in microfluidic devices combined with microcontact printing (See, e.g., Martinez-Rivas, A., *Methods of micropatterning and manipulation of cells for biomedical applications*, Micromachines (Basel) 8, (2017), which is incorporated herein by reference).

[0673] The coating can be photoreactive, and selectively activating or degrading the coating involves exposing selected regions of the coating to light or radiation. Selectivity can be achieved through the application of photomasks. Regions of the coating that are exposed to light can be rendered more permissive for bead attachment (e.g., more adhesive), as compared to regions not exposed to light (e.g., regions protected from the light by a photomask). When applied to the substrate, the beads thus preferentially attach to the more permissive regions on the substrate, and un-attached beads can optionally be removed from the substrate. The light source and/or the photomask can be adjusted to allow further sites on the substrate to become more permissive for bead attachment, allowing additional beads to be attached at these sites. Alternatively, a different light source, or a different photomask can be applied. The process of photopatterning thus allows beads to be attached at pre-determined locations on the substrate, thereby generating a bead array.

[0674] Beads can be attached iteratively, e.g., a subset of the beads can be attached at one time, and the process can be repeated to attach one or more additional subsets of beads. In some embodiments, the size of the activated spot (e.g., spot on the substrate) is smaller than the size of a

bead. For example, a bead can be attached to the activated substrate (e.g., spot) such that only a single bead attaches to the activated substrate. In some embodiments, the substrate can be washed to remove unbound beads. In some embodiments, the substrate can be activated in a second location and a second bead can be attached to the activated substrate surface. This process can be done iteratively to attach beads to the entire substrate, or a portion thereof. Alternatively, beads can be attached to the substrate all in one step. Furthermore, methods of attaching beads to a substrate are known in the art. Any suitable method can be used, including, in a non-limiting way, specific chemical bonds, non-specific chemical bonds, linkers, physically trapping the beads (e.g., polymer, hydrogel), or any of the methods described herein.

[0675] An exemplary workflow for generating a bead array can include selectively rendering a first set of one or more selected regions on a coated substrate to be more permissive for bead attachment as compared to regions outside of the selected regions, applying a plurality of beads to the array and allowing the beads to attach to the first set of selected regions, optionally removing un-attached beads, rendering a second set of one or more selected regions more permissive to bead attachment as compared to regions outside the second set of selected regions, applying a plurality of beads and allowing the beads to attach to the second set of selected regions, and optionally removing the un-attached beads. This iterative process can be carried out for any number of times to generate a patterned bead array.

[0676] Another exemplary process includes activating a first region on a coated substrate and exposing the activated first region to a plurality of barcoded beads, so that a first set of one or more beads are bound to the first region; and activating a second region on the coated substrate and exposing the activated second region to a plurality of barcoded beads, so that a second set of one or more beads are bound to the second region, wherein the first set of one or more beads comprise an identical first oligonucleotide sequence unique to the first region of the surface of the substrate, and the second set of one or more beads comprise an identical second oligonucleotide sequence unique to the second region of the surface of the substrate, and wherein the first and second oligonucleotide sequences are different. Additional regions on the coated substrate may be activated and exposed to additional barcoded beads. Each set of barcoded beads can include an oligonucleotide sequence that is different from all other sets of barcoded beads and that is unique to the location of the activated region. In some instances, a set of barcoded beads includes a two-part barcode, where the first part is correlated with the location to which the set of beads are bound, and the second part is correlated with the location of a bead relative to the beads in the set. For example, the location of a bead within a set of beads bound to the substrate can be identified based on the sequence information of the first and second part barcodes. The first part of the barcode can be identical across beads within the same set, and can be attached to the beads before or after the beads are bound to the substrate. The second part of the barcode can be different between two beads within the set, and can be attached to the beads (e.g., ligated onto the first part barcode) before or after the beads are bound to the substrate. In some instances, the second part of the barcode is attached to the beads after they are bound to the substrate.

[0677] Additionally, the first set of one or more beads and the second set of one or more beads can be different. In other words, the first set of one or more beads and the second set of one or more beads can have different surface chemistries, different compositions (e.g., solid bead, gel bead, silica bead) (e.g., nanoparticles vs microparticles), and/or physical volumes. In some embodiments, a third set of one or more beads, a fourth set of one or more beads, a fifth set of one or more beads or more can have different surface chemistries, different compositions (e.g., solid bead, gel bead, silica bead) (e.g., nanoparticles vs microparticles), and/or physical volumes can be attached to the substrate surface. The methods may include removing the beads that do not bind to the first, second, and/or any of the additional regions. In some embodiments, removing the beads comprise washing the beads off the surface of the substrate. The removing may be carried out after each round of or after several rounds of activating a region (e.g., first, second or additional regions on

the surface of the substrate), and binding of beads to the activated region. In some instances, each bead is bound to the substrate at a single location. The beads bound to the first, second, and additional regions can form a two-dimensional array of beads on the substrate.

[0678] A photoreactive coating can comprise a plurality of photoreactive molecules, which can undergo a chemical reaction (e.g., hydrolysis, oxidation, photolysis) when exposed to light of certain wavelengths or range of wavelengths. A photo-reactive molecule can become reactive when exposed to light and can react with other molecules and form chemical bonds with other molecules.

[0679] The coating can comprise a polymer, and activating selected regions on the substrate include modifying the polymer at the respective regions. Modifying the polymer includes, for example, photochemically modifying the polymer by exposing the polymer to radiation or light. Alternatively or additionally, modifying the polymer can include chemically modifying the polymer by contacting the polymer with one or more chemical reagents. In some instances, the coating is a hydrogel. In some instances, the coating comprises a photoreactive polymer.

Exemplary photo-reactive polymers include poly(ethylene glycol) (PEG)-based polymers, poly(L-lysine) (PLL)-based polymer, copolymer comprising functionalized or unfunctionalized units of PEG and PLL (e.g., poly-L-lysine-grafted-polyethylene glycol (PLL-g-PEG)), and methacrylated gelatin (GelMA) polymers.

[0680] Beads can also be attached to selective regions on a substrate by selectively crosslinking beads to a coating that has been applied on the substrate. For example, a plurality of beads can be applied to a substrate having a photocrosslinkable coating, and upon crosslinking of a subset of the beads to the coating, the non-cross-linked beads can be removed, leaving only the cross-linked beads on the substrate. The process can be repeated multiple times. The coating can include a photo-crosslinkable polymer. Exemplary photo-crosslinkable polymers are described, e.g., in Shirai, *Polymer Journal* 46:859-865 (2014), Ravve, *Photocrosslinkable Polymers, Light-Associated Reactions of Synthetic Polymers*. Springer, New York, NY (2006), and Ferreira et al. *Photocrosslinkable Polymers for Biomedical Applications, Biomedical Engineering—Frontiers and Challenges*, Prof. Reza Fazel (Ed.), ISBN: 978-953-307-309-5 (2011), each of which are herein incorporated by reference in its entirety.

[0681] Suitable light sources for activating, degrading or crosslinking the coating as described herein include, but are not limited to, Ultraviolet (UV) light (e.g., 250-350 nm or 350-460 nm UV light) and visible light (e.g., broad spectrum visible light). A Digital Micromirror Device (DMD) can also be used to provide the light source.

[0682] The distance between a first pair of adjacent selected regions according to the methods described herein can be the same or different from a second pair of adjacent selected regions.

[0683] Barcoded beads, or beads comprising a plurality of barcoded probes, can be generated by first preparing a plurality of barcoded probes on a substrate, depositing a plurality of beads on the substrate, and generating probes attached to the beads using the probes on the substrate as a template.

[0684] Large scale commercial manufacturing methods allow for millions of oligonucleotides to be attached to an array. Commercially available arrays include those from Affymetrix (ThermoFisher Scientific).

[0685] In some embodiments, arrays can be prepared according to the methods set forth in WO 2012/140224, WO 2014/060483, WO 2016/162309, WO 2017/019456, WO 2018/091676, and WO 2012/140224, and U.S. Patent Application No. 2018/0245142. The entire contents of each of the foregoing documents are herein incorporated by reference.

[0686] In some embodiments, a bead array is formed when beads are embedded in a hydrogel layer where the hydrogel polymerizes and secures the relative bead positions. The bead-arrays can be pre-equilibrated and combined with reaction buffers and enzymes (e.g., reverse-transcription mix). In some embodiments, the bead arrays can be stored (e.g., frozen) long-term (e.g., days) prior to use.

(vi) Flexible Arrays

[0687] A “flexible array” includes a plurality of spatially-barcoded features attached to, or embedded in, a flexible substrate (e.g., a membrane, a hydrogel, or tape) placed onto, or proximal to, a biological sample. In some embodiments, a flexible array includes a plurality of spatially-barcoded features embedded within a hydrogel.

[0688] Flexible arrays can be highly modular. In some embodiments, spatially-barcoded features (e.g., beads) can be loaded onto a substrate (e.g., a slide) to produce a high-density self-assembled array. In some embodiments, the features (e.g., beads) can be loaded onto the substrate with a flow cell. In some embodiments, the features (e.g., beads) are embedded in a hydrogel (e.g., a hydrogel pad or layer placed on top of the self-assembled features). In some embodiments, the hydrogel can polymerize, thereby securing the features in the hydrogel. In some embodiments, the hydrogel containing the features can be removed from the substrate and used as a flexible array. In some embodiments, the flexible array can be deconvolved by optical sequencing or any other method described herein. In some embodiments, the features (e.g., beads) can be about 1 μm to about 25 μm in diameter. In some embodiments, about 25 μm diameter features in the flexible array can provide for approximately 1000 DPI and about 1 megapixel resolution. In some embodiments, the features (e.g., beads) can be about 13.2 μm in diameter. In some embodiments, the about 13.2 μm beads in the flexible array can provide for approximately 1920 $\times$ 1080 resolution.

[0689] Flexible arrays generated according to any of the methods described herein (e.g., beads embedded within a hydrogel) can contain a thermolabile polymer. In some embodiments, flexible arrays having thermolabile beads can be contacted with a biological sample. In some embodiments, a region of interest in the biological sample can be identified such that an infrared laser can be used to select a region of interest. In some embodiments, the infrared laser can cause the flexible array (e.g., thermolabile beads) to deform and become adhesive. In some embodiments, the adhesive portion of the flexible array can adhere (e.g., bind) to the region of interest (e.g., cells) directly above or underneath. The process of identifying a region of interest, applying an infrared laser to the region of interest, and adhering the underlying biological sample (e.g., cells) to the flexible array can iteratively repeated. In some embodiments, the flexible array can be removed such that only the adhered biological sample (e.g., cells) from the one or more regions of interest can also be removed with the flexible array. In some embodiments, the flexible array and the adhered biological sample can be further processed (e.g., amplified, quantitated, and/or sequenced) according to any method described herein.

[0690] Flexible arrays can be pre-equilibrated with reaction buffers and enzymes at functional concentrations (e.g., a reverse-transcription mix). In some embodiments, the flexible arrays can be stored for extended periods (e.g., days) or frozen until ready for use. In some embodiments, permeabilization of biological samples (e.g., a tissue section) can be performed with the addition of enzymes/detergents prior to contact with the flexible array. In some embodiments, the flexible array can be placed directly on the sample, or placed in indirect contact with the sample (e.g., with an intervening layer or substance between the biological sample and the flexible bead-array). In some embodiments, the flexible array can be mechanically applied (e.g., pressed downward or compressed between two surfaces) on the biological sample to enhance analyte capture. In some embodiments, a flexible array can be applied to the side of a biological sample. For example, a biological sample can be cut (e.g., sliced) in any direction and a flexible array can be applied to the exposed analytes. In some embodiments, the flexible array can be dissolvable (e.g., via heat, chemical, or enzymatic disruption). In some embodiments, once a flexible array is applied to the sample, reverse transcription and targeted capture of analytes can be performed on microspheres, or beads of a first volume and beads of a second volume, or any of the beads described herein. In some embodiments once a flexible array is applied to the biological sample and allowed to capture analytes, the flexible array can be removed (e.g., peeled) from the biological sample for further processing (e.g., amplification, quantitation, and/or sequencing) according to any method described

herein.

[0691] Flexible arrays can also be used with any of the methods (e.g., active capture methods such as electrophoresis) described herein. For example, flexible arrays can be contacted with a biological sample on a conductive substrate (e.g., an indium tin oxide coated glass slide), such that an electric field can be applied to the conductive substrate to facilitate migration of analytes through, across, within, or in the direction of the flexible array. Additionally and alternatively, flexible arrays can be contacted to a biological sample in an electrophoretic assembly (e.g., electrophoretic chamber), such that an electric field can be applied to migrate analytes in the direction of the flexible array or across, through, or within the flexible array.

[0692] In some embodiments, a flexible array can be generated with the assistance of a substrate holder (e.g., any array alignment device). For example, a spatially-barcoded bead array can be placed in one placeholder of the substrate holder and second substrate (e.g., a glass slide) can be placed in the second placeholder of the substrate holder. In some embodiments, the array is optionally optically decoded and a gel prepolymer solution is introduced between the spatially-barcoded bead array and second substrate. In some embodiments, the substrate holder is closed such that the second substrate is on top (e.g., above, parallel to) the spatially-barcoded bead array. The gel prepolymer solution can be polymerized by any method described herein and result in spatially-barcoded features cross-linked in the hydrogel, thereby generating a flexible array. In some embodiments, the substrate holder can be opened and the second substrate with the hydrogel and the spatially-barcoded cross-linked features can be removed from the substrate holder (the flexible array optionally can be removed from the second substrate) for use in spatial analysis by any of the methods described herein.

(vii) Shrinking Hydrogel Features/Arrays

[0693] As used herein “shrinking” or “reducing the size” of a hydrogel refers to any process causing the hydrogel to physically contract and/or the size of the hydrogel to decrease in volume. For example, the scaffold of the gel may shrink or “implode” upon solvent removal (see, e.g., Long and Williams. *Science*. 2018; 362 (6420): 1244-1245, and Oran et al. *Science* 2018; 362 (6420): 1281-1285; each of which is incorporated herein by reference in its entirety). As another example, the process to shrink or reduce the volume of a hydrogel may be one that removes water (i.e., a dehydrating process) from the hydrogel. There are many methods known to one of skill in the art for shrinking or reducing the volume of a hydrogel. Non-limiting examples of a method to shrink or reduce the volume of a hydrogel include exposing the hydrogel to one or more of: a dehydrating solvent, a salt, heat, a vacuum, lyophilization, desiccation, filtration, air-drying, or combinations thereof.

[0694] In some embodiments, a hydrogel bead can be decreased in volume (e.g., shrunken hydrogel bead) before being attached to or embedded in a hydrogel. In some embodiments, a hydrogel bead can be decreased in volume (e.g., shrunken hydrogel bead) after being attached to or embedded in a hydrogel. In some embodiments, one or more hydrogel beads can be attached to or embedded in a hydrogel. In some embodiments, one or more hydrogel beads can be decreased in volume (e.g., one or more shrunken hydrogel beads) before being attached to or embedded in a hydrogel. In some embodiments, one or more hydrogel beads can be decreased in volume (e.g., one or more shrunken hydrogel beads) after being attached to or embedded in a hydrogel. In some embodiments, one or more hydrogel beads attached to or embedded in a hydrogel can be decreased in volume. For example, the one or more hydrogel beads and the hydrogel that the hydrogel beads are attached to or embedded in are decreased in volume at the same time (e.g., shrunken hydrogel bead-containing hydrogel). In some embodiments, one or more hydrogel beads attached to or embedded in a hydrogel can be isometrically decreased in volume.

[0695] In some embodiments, one or more hydrogel beads attached to or embedded in a hydrogel can be decreased in volume from about 3 fold to about 4 fold. For example, one or more hydrogel beads attached to or embedded in a hydrogel can be decreased in volume by removing or

exchanging solvents, salts, or water (e.g., dehydration). In another example, one or more hydrogel beads attached to or embedded in a hydrogel can be decreased in volume by controlling temperature or pH. See e.g., Ahmed, E. M. *J. of Advanced Research*. 2015 March; 6 (2): 105-121, which is incorporated herein by reference in its entirety. In some embodiments, one or more hydrogel beads attached to or embedded in a hydrogel can be decreased in volume by removing water.

[0696] In some embodiments, decreasing the volume of one or more hydrogel beads attached to or embedded in a hydrogel can increase the spatial resolution of the subsequent analysis of the sample. The increased resolution in spatial profiling can be determined by comparison of the spatial analysis of the sample using one or more shrunken hydrogel beads attached to or embedded in a hydrogel with one or more non-shrunken hydrogel beads attached to or embedded in a hydrogel.

[0697] In some embodiments, a hydrogel bead is not decreased in volume. In some embodiments, a hydrogel bead can be decreased in volume (e.g., shrunken hydrogel bead). In some embodiments, a shrunken hydrogel gel bead is stabilized. For example, the hydrogel bead can be decreased in volume by removing solvents, salts, or water (e.g., dehydrated, desiccated, dried, exsiccated) from the hydrogel bead to form a shrunken hydrogel bead. In another example, the hydrogel bead can be decreased in volume by controlling temperature or pH. See e.g., Ahmed, E. M. *J. of Advanced Research*. 2015 March; 6 (2): 105-121, which is incorporated herein by reference in its entirety. Non-limiting examples of solvents that may be used to form a shrunken hydrogel bead or shrunken hydrogel bead array include a ketone, such as methyl ethyl ketone (MEK), isopropanol (IPA), acetone, 1-butanol, methanol (MeOH), dimethyl sulfoxide (DMSO), glycerol, propylene glycol, ethylene glycol, ethanol, (k) 1,4-dioxane, propylene carbonate, furfuryl alcohol, N,N-dimethylformamide (DMF), acetonitrile, aldehyde, such as formaldehyde or glutaraldehyde, or any combinations thereof.

[0698] In some embodiments, the hydrogel bead or hydrogel bead array is shrunken or stabilized via a cross-linking agent. For example, the cross-linking agent may comprise disuccinimidyl suberate (DSS), dimethylsuberimidate (DMS), formalin, and dimethyladipimidate (DMA), dithio-bis(-succinimidyl propionate) (DSP), disuccinimidyl tartrate (DST), and ethylene glycol bis(succinimidyl succinate) (EGS).

[0699] In some embodiments, the hydrogel bead or hydrogel bead array is processed with salts to form a shrunken hydrogel bead or shrunken hydrogel bead array. Non-limiting examples of salts that may be used to form a shrunken hydrogel bead or shrunken hydrogel bead array are inorganic salts including aluminum, ammonium, barium, beryllium, calcium, cesium, lithium, magnesium, potassium, rubidium, sodium, and strontium salts. Further non-limiting examples of inorganic salts include sodium chloride, potassium chloride, lithium chloride, cesium chloride, sodium fluoride, sodium bromide, sodium iodide, sodium nitrite, potassium sulfate, potassium nitrate, potassium carbonate, potassium bicarbonate, sodium sulfate, sodium nitrate, sodium carbonate, sodium bicarbonate, calcium sulfate, copper oxychloride, calcium chloride, calcium carbonate, calcium bicarbonate, magnesium sulfate, magnesium nitrate, magnesium chloride, magnesium carbonate, magnesium bicarbonate, ammonium sulfate, ammonium chloride, ammonium nitrate, ammonium carbonate, ammonium bicarbonate, trisodium phosphate, tripotassium phosphate, calcium phosphate, copper(II) sulfate, sodium sulfide, potassium sulfide, calcium sulfide, potassium permanganate, iron(II) chloride, iron(III) chloride, iron (2+) sulfate, iron(III) sulfate, iron(II) nitrate, iron(III) nitrate, manganese(II) chloride, manganese(III) chloride, manganese(II) sulfate, manganese(II) nitrate, zinc chloride, zinc nitrate, zinc sulfate, ammonium orthomolybdate, monopotassium phosphate, nickel(II) sulfate, nickel(II) nitrate, sodium metavanadate, sodium paravanadate, potassium dichromate, ammonium dichromate, antipyrone, ammonium nitrite, potassium fluoride, sodium fluoride, ammonium fluoride, calcium fluoride, chrome alum, potassium alum, potassium iodide, sodium hypochlorite, tin(II) sulfate, tin(II) nitrate, gold selenite,

dicesium chromate, potassium perchlorate, calcium perchlorate, aluminum sulphate, lead(II) bisulfate, barium phosphate, barium hydrogen orthophosphate, barium dihydrogen phosphate, silver dichromate, potassium bromate, sodium bromate, sodium iodate, sodium silicate, diammonium phosphate, ammonium phosphate, ammonium dihydrogen phosphate, chromium orthophosphate, copper(II) chloride, copper(I) chloride, sodium tetrametaphosphate, potassium heptafluoroniobate, zinc phosphate, sodium sulfite, copper(I) nitrate, copper(II) nitrate, potassium silicate, copper(II) carbonate basic, copper(II) carbonate salts of acrylic acid and sulfopropyl acrylate.

[0700] In some embodiments, the removal of water comprises an acid. Non-limiting examples of an acid include: HCl, HI, HBr, HClO.sub.4, HClO.sub.3, HNO.sub.3, H.sub.2SO.sub.4, phosphoric acid, phosphorous acid, acetic acid, oxalic acid, ascorbic acid, carbonic acid, sulfurous acid, tartaric acid, citric acid, malonic acid, phthalic acid, barbituric acid, cinnamic acid, glutaric acid, hexanoic acid, malic acid, folic acid, propionic acid, stearic acid, trifluoroacetic acid, acetylsalicylic acid, glutamic acid, azelaic acid, benzilic acid, fumaric acid, gluconic acid, lactic acid, oleic acid, propiolic acid, rosolic acid, tannic acid, uric acid, gallic acid, and combinations of two or more thereof. In some embodiments, the hydrogel is exposed to a different pH environment. For example, the hydrogel can be exposed to an acidic pH or a basic pH. In some embodiments, the hydrogel is exposed to a pH of less than about 6.5, e.g., a pH of about 6, about 5.5, about 5, about 4.5, about 4, about 3.5, about 3, about 2.5, about 2, about 1.5, or about 1. In some embodiments, the hydrogel is exposed to a pH of greater than about 7.5, e.g., a pH of about 8, about 8.5, about 9, about 9.5, about 10, about 10.5, about 11, about 11.5, about 12, about 12.5, about 13, about 13.5, or about 14.

[0701] In some embodiments, the removal of water comprises a dehydrating process such as heat, a vacuum, lyophilization, desiccation, filtration, and air-drying. In some embodiments, the hydrogel bead or hydrogel bead array undergoes an alteration in pH to form a shrunken hydrogel bead or shrunken hydrogel bead array (e.g., an alteration from about pH 7 to about pH 5, from about pH 7 to about pH 5.5, from about pH 7 to about pH 6, from about pH 7 to about pH 6.5, from about pH 6.5 to about pH 5, from about pH 6 to about pH 5, from about pH 6 to about pH 5.5, or any pH alteration encompassed within these ranges).

[0702] In some embodiments, the hydrogel bead or hydrogel bead array undergoes an alteration in temperature (e.g., an alteration from about 37° C., 38° C., 39° C., 40° C., 41° C., 42° C., 43° C., 44° C., 45° C., 46° C., 47° C., 48° C., 49° C. to about 50° C., 51° C., 52° C., 53° C., 54° C., 55° C., 56° C., 57° C., 58° C., 59° C. 60° C., 61° C., 62° C., 63° C., 64° C., 65° C., 66° C., 67° C., 68° C., 69° C., 70° C., or higher, or any temperature alteration encompassed within these ranges) to form a shrunken hydrogel bead or shrunken hydrogel bead array.

[0703] In some embodiments, a hydrogel bead can be decreased in size in linear dimension by about 2 fold, about 3 fold, about 4 fold, about 5 fold, about 6 fold, about 7 fold, about 8 fold, about 9 fold, or any intervals therein. In some embodiments, a hydrogel bead can be decreased in volume by about 1 fold, about 5 fold, about 10 fold, about 15 fold, about 20 fold, about 25 fold, about 30 fold, about 35 fold, about 40 fold, about 45 fold, about 50 fold, about 55 fold, about 60 fold, about 65 fold about 70 fold, about 75 fold, about 80 fold, or any intervals therein. In some embodiments, a hydrogel bead can be decreased in size such that the hydrogel bead has an average diameter of about 1 μ m to about 15 μ m.

[0704] In some embodiments, a plurality of hydrogel beads can be decreased in size in linear dimension by about 2 fold, about 3 fold, about 4 fold, about 5 fold, about 6 fold, about 7 fold, about 8 fold, about 9 fold, or any intervals therein. In some embodiments, a plurality of hydrogel beads can be decreased in size such that the average diameter of a hydrogel bead is about 1 μ m to about 15 μ m. In some embodiments, a plurality of hydrogel beads can be decreased in volume by about 1 fold, about 5 fold, about 10 fold, about 15 fold, about 20 fold, about 25 fold, about 30 fold, about 35 fold, about 40 fold, about 45 fold, about 50 fold, about 55 fold, about 60 fold, about 65 fold

about 6-13 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-13 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 11-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 10-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 9-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 8-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 7-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 6-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-12 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 10-11 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 9-11 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 8-11 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 7-11 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 6-11 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-11 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 9-10 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 8-10 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 7-10 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 6-10 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-10 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 8-9 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 7-9 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 6-9 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-9 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 7-8 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 6-8 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-8 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 6-7 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-7 μm . In some embodiments, a plurality of shrunken hydrogel beads has an average diameter of about 5-6 μm .

[0706] In some embodiments, one or more hydrogel beads can be decreased in volume at the same time. In some embodiments, one or more hydrogel beads can be decreased in volume at different times. In some embodiments, one or more hydrogel beads can be assembled into an array before decreasing the volume of the one or more hydrogel beads. In some embodiments, one or more hydrogel beads can be assembled into an array after decreasing the volume of the one or more hydrogel beads. In some embodiments, the one or more shrunken hydrogel beads can be reversibly attached to a substrate. In some embodiments, the one or more shrunken hydrogel beads can be irreversibly attached to a substrate. In some embodiments, the one or more shrunken hydrogel beads can be re-expanded. In some embodiments, the one or more shrunken hydrogel beads can be isometrically re-expanded. In some embodiments, the one or more shrunken hydrogel beads can be re-expanded primarily in the z-dimension. In some embodiments, the one or more shrunken hydrogel beads attached to a substrate (e.g., reversibly or irreversibly) can be re-expanded primarily in the z-dimension. In some embodiments, the one or more shrunken hydrogel beads attached to a substrate (e.g., reversibly or irreversibly) can be isometrically re-expanded primarily in the z-dimension.

[0707] In some embodiments, decreasing the volume of the hydrogel bead (e.g., shrunken hydrogel bead) can increase the spatial resolution of the subsequent analysis of the sample. The increased

resolution in spatial profiling can be determined by comparison of spatial analysis of the sample using a shrunken hydrogel bead with a non-shrunken hydrogel bead. For example, in some embodiments, the subsequent analysis of the sample can include any array-based spatial analysis method disclosed herein.

[0708] In some embodiments, one or more physical parameters or dimensions and/or one or more other characteristics of the hydrogel bead may be changed. For example, a cross-section of the hydrogel bead may be changed from a first cross-section to a second cross-section. The first cross-section may be smaller or larger than the second cross-section. Alternatively, or in addition, one or more other characteristics of the hydrogel bead may be changed. For example, the fluidity, density, rigidity, porosity, refractive index, polarity, and/or other characteristic of the hydrogel bead or one or more components thereof may be changed. In a non-limiting example, the hydrogel bead includes a hydrogel. In another example, the hydrogel bead hydrogel may form crosslinks within the bead. The same or different conditions may be used to change or affect different characteristics of the hydrogel bead at the same or different times. In some cases, a first condition or set of conditions may be used to change a first characteristic or set of characteristics of the hydrogel bead (e.g., a cross-section) and a second condition or set of conditions may be used to change a second characteristic or set of characteristics of the hydrogel bead. The first condition or set of conditions may be applied at the same or a different time as the second condition or set of conditions. For example, a first characteristic or set of characteristics may be changed under a first condition or set of conditions, after which a second characteristic or set of characteristics may be changed under a second condition or set of conditions.

[0709] A characteristic or set of characteristics of the hydrogel bead may be changed by one or more conditions. A condition suitable for changing a characteristic or set of characteristics of the hydrogel bead may be, for example, a temperature, a pH, an ion or salt concentration, a pressure, chemical species, any combinations thereof, or another condition. For example, hydrogel bead may be exposed to a chemical species that may bring about a change in one or more characteristics of the hydrogel bead. In some cases, a stimulus may be used to change one or more characteristics of the hydrogel bead. For example, upon application of the stimulus, one or more characteristics of the hydrogel bead may be changed. The stimulus may be, for example, a thermal stimulus, a photo stimulus, a chemical stimulus, or another stimulus. A temperature sufficient for changing one or more characteristics of the hydrogel bead may be, for example, at least about 0 degrees Celsius ($^{\circ}$ C.), 1° C., 2° C., 3° C., 4° C., 5° C., 10° C., or higher. For example, the temperature may be about 4° C. In other cases, a temperature sufficient for changing one or more characteristics of the hydrogel bead may be, for example, at least about 25° C., 30° C., 35° C., 37° C., 40° C., 45° C., 50° C., or higher. For example, the temperature may be about 37° C. A pH sufficient for changing one or more characteristics of the hydrogel bead may be, for example, between about 5 and 8, such as between about 6 and 7.

[0710] In some cases, a chemical species or a chemical stimulus may be used to change one or more characteristics of the hydrogel bead. For example, a chemical species or a chemical stimulus may be used to change a dimension of a hydrogel bead (e.g., a cross-section, diameter, or volume). In some cases, a chemical species or a chemical stimulus may be used to change a dimension of a hydrogel bead (e.g., a cross-sectional diameter) from a first dimension to a second dimension (e.g., a second cross-sectional diameter), where the second dimension is reduced compared to the first dimension. The chemical species may comprise an organic solvent, such as an alcohol, ketone, or aldehyde. For example, the chemical species may comprise acetone, methanol, ethanol, formaldehyde, or glutaraldehyde. The chemical species may comprise a cross-linking agent. For example, the cross-linking agent may comprise disuccinimidyl suberate (DSS), dimethylsuberimidate (DMS), formalin, and dimethyladipimidate (DMA), dithio-bis(-succinimidyl propionate) (DSP), disuccinimidyl tartrate (DST), and ethylene glycol bis(succinimidyl succinate) (EGS), and any combinations thereof. In some cases, a cross-linking agent may be a photo-

cleavable cross-linking agent. In some cases, a chemical stimulus may be used to change one or more characteristics of the hydrogel bead (e.g., a dimension of a hydrogel bead), where the chemical stimulus comprises one or more chemical species. For example, the chemical stimulus may comprise a first chemical species and a second chemical species, where the first chemical species is an organic solvent and the second chemical species is a cross-linking agent. In some cases, a chemical stimulus may comprise a chemical species that is a fixation agent that is capable of fixing or preserving a hydrogel bead. For example, an organic solvent such as an alcohol (e.g., ethanol or methanol), ketone (e.g., acetone), or aldehyde (e.g., formaldehyde or glutaraldehyde), or any combinations thereof may act as a fixation agent. Alternatively, or in addition, a cross-linking agent may act as a fixation agent. In some cases, a fixation agent may comprise disuccinimidyl suberate (DSS), dimethylsuberimide (DMS), formalin, and dimethyladipimide (DMA), dithio-bis(-succinimidyl propionate) (DSP), disuccinimidyl tartrate (DST), and/or ethylene glycol bis(succinimidyl succinate) (EGS), and any combinations thereof. In some cases, a first chemical species and/or fixation agent may be provided to or brought into contact with the hydrogel bead to bring about a change in a first characteristic or set of characteristics of the hydrogel bead, and a second chemical species and/or fixation agent may be provided to or brought into contact with the hydrogel bead to bring about a change in a second characteristic or set of characteristics of the hydrogel bead. For example, a first chemical species and/or fixation agent may be provided to or brought into contact with the hydrogel bead to bring about a change in a dimension of a hydrogel bead (e.g., a reduction in cross-sectional diameter), and a second chemical species and/or fixation agent may be provided to or brought into contact with the hydrogel bead to bring about a change in a second characteristic or set of characteristics of the hydrogel bead (e.g., forming crosslinks within and/or surrounding the hydrogel bead). The first and second chemical species and/or fixation agents may be provided to or brought into contact with the hydrogel bead at the same or different times.

[0711] In some embodiments, fixation may affect one or more parameters or characteristics of the hydrogel bead. For example, fixation may result in shrinkage or volumetric reduction of the hydrogel bead. Fixation may include dehydration of the hydrogel bead. Providing a fixation agent to the hydrogel bead may result in a change in a dimension of the hydrogel bead. For example, providing a fixation agent to the hydrogel bead may result in a change in the volume or diameter of the hydrogel bead. Providing a fixation agent to the hydrogel bead may result in a change in a cross-section of the hydrogel bead (e.g., a cross-sectional diameter). For example, a first cross-section of the hydrogel bead prior to fixation may be different (e.g., larger) than a second cross-section of the hydrogel bead following fixation. In an example, an approximately spherical hydrogel bead may comprise a first cross section (e.g., a cross-sectional diameter) prior to fixation that is reduced in size to a second cross-section following fixation. Providing a fixation agent to the hydrogel bead may result in a second cross-section that is reduced by at least about 5% compared to the first cross-section. In some cases, the second cross-section may be reduced by at least 6%, 8%, 10%, 15%, 25%, 30%, 35%, 40%, 45%, 50%, or more relative to the first cross-section. For example, the second cross-section may be reduced by at least about 10%, 15%, 25%, or 50% relative to the first cross-section. Fixation may also affect other features of the hydrogel bead. For example, fixation may result in a change in the porosity of a membrane or wall of a hydrogel bead, reorganization of components of the hydrogel bead, a change in hydrogel bead fluidity or rigidity, or other changes. In an example, a first fixation agent that is an organic solvent is provided to the hydrogel bead to change a first characteristic (e.g., hydrogel bead volume) and a second fixation agent that is a cross-linking agent is provided to the hydrogel bead to change a second characteristic (e.g., hydrogel bead fluidity or rigidity). The first fixation agent may be provided to the hydrogel bead before the second fixation agent.

[0712] In some instances, an approximately spherical hydrogel bead may comprise a first diameter prior to fixation (e.g., by an organic solvent) that is reduced in volume compared to a second

diameter following fixation when maintained in a non-aqueous environment. Following fixation and reduction in volume to said second diameter, when maintained in an aqueous environment, the hydrogel bead may increase in volume to have a diameter substantially similar to the first diameter. In some cases, an approximately spherical hydrogel bead may include a first diameter prior to fixation (e.g., by an organic solvent) that is reduced in volume compared to a second diameter following fixation. Following fixation and reduction in volume to said second diameter, the hydrogel bead may be cross-linked by a second fixative, wherein the second diameter is substantially maintained in an aqueous environment following cross-linking by the second fixative. [0713] A change to a characteristic or set of characteristics of the hydrogel bead may be reversible or irreversible. In some cases, a change to a characteristic or set of characteristics of the hydrogel bead may be irreversible, such that the change cannot be readily undone. For example, the volume, morphology, or other feature of the hydrogel bead may be altered in a way that cannot be readily reversed. In an example, the change from a first cross-section of the hydrogel bead to a second cross-section of the hydrogel bead is irreversible. In some cases, an irreversible change may be at least partially reversed upon the application of appropriate conditions and/or over a period of time. In other cases, a change to a characteristic or set of characteristics of the hydrogel bead may be reversible. For example, the volume of a hydrogel bead may be reduced upon being subjected to a first condition or set of conditions, and the volume of a hydrogel bead may be increased to approximately the original volume upon being subjected to a second condition or set of conditions. Thus, the change from a first cross-section of the hydrogel bead to the second cross-section may be reversible. A reversible change (e.g., a reversible volume reduction) may be useful in, for example, providing a hydrogel bead of a given volume to a given location, such as a partition. In some cases, a change to a characteristic or set of characteristics of the hydrogel bead may be only partially reversible. For example, the volume of a hydrogel bead may be reduced (e.g., by dehydration), and the reduction in hydrogel bead volume may be accompanied by reorganization of components within the hydrogel bead. Upon reversal of the volume of the hydrogel bead (e.g., by rehydration), the arrangement of one or more components may not revert to the original arrangement of the hydrogel bead prior to the volume reduction. A change to a characteristic or set of characteristics of the hydrogel bead, such as a cross-section of the hydrogel bead, may be reversible upon application of a stimulus. The stimulus may be, for example, a thermal stimulus, a photo stimulus, or a chemical stimulus. In some cases, the stimulus may comprise a change in pH and/or application of a reducing agent such as dithiothreitol. Application of the stimulus may reverse, wholly or in part, a change from, for example, a first cross-section to a second cross-section.

[0714] In some embodiments, a plurality of hydrogel beads can be shrunken hydrogel beads generated by removing water from a plurality of first hydrogel beads. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter no larger than about 15 microns. For example, the plurality of shrunken hydrogel beads has an average diameter no larger than about 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 micron. In some embodiments, each member of the plurality of shrunken hydrogel beads has a diameter no larger than about 15 microns. For example, each member of the plurality of shrunken hydrogel beads can have a diameter no larger than about 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 micron. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter no larger than 10 microns. In some embodiments, each member of the plurality of shrunken hydrogel beads has a diameter no larger than 10 microns. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter no larger than 5 microns. In some embodiments, each member of the plurality of shrunken hydrogel beads has a diameter no larger than 5 microns. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter no larger than 1 micron. In some embodiments, each member of the plurality of shrunken hydrogel beads has a diameter no larger than 1 micron. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter no larger than the diameter of a cell (e.g., a mammalian cell, a plant cell, or a fungal cell). In some

embodiments, each member of the plurality of shrunken hydrogel beads has a diameter no larger than the diameter of a cell (e.g., a mammalian cell, a plant cell, or a fungal cell).

[0715] In some embodiments, the plurality of shrunken hydrogel beads has a polydispersity index of less than about 25%. For example, the plurality of shrunken hydrogel beads can have a polydispersity index of less than about 25%, 20%, 15%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, or 1%. In some embodiments, the plurality of shrunken hydrogel beads has a polydispersity index of less than 15%. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter of about 8 to about 13 microns. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter of about 10 to about 12 microns. In some embodiments, the plurality of shrunken hydrogel beads has an average diameter of about the diameter of a cell (e.g., a mammalian cell, a plant cell, or a fungal cell). In some embodiments, the plurality of shrunken hydrogel beads has an average diameter of less than the diameter of a cell (e.g., a mammalian cell, a plant cell, or a fungal cell). In some embodiments, the plurality of capture probes on the plurality of shrunken gel beads bind cellular analytes at single-cell resolution. In some embodiments, the plurality of capture probes on the plurality of shrunken gel beads bind cellular analytes at higher than single-cell resolution (e.g., at a resolution that is at a higher density than the diameter of a cell).

[0716] In some embodiments, bead arrays having a plurality of hydrogel beads disposed on a substrate are generated by patterning or self-assembly of larger gel beads, after which the array of larger gel beads is shrunken (e.g., by any of the variety of methods provided herein). In some embodiments, the larger gel beads are not small enough for single-cell resolution, while the shrunken gel beads are small enough for single-cell resolution. In some embodiments, bead arrays having a plurality of hydrogel beads disposed on a substrate are generated by patterning or self-assembly of shrunken gel beads that have previously been generated by shrinking larger gel beads (e.g., by any of the variety of methods provided herein). Beads can be spatially confined by any of a variety of methods, including without limitation, reversible or irreversible crosslinking.

[0717] In some embodiments, bead arrays include spatially-confined gel beads with high aspect ratios (e.g., pillared arrays). For example, bead arrays having a plurality of hydrogel beads disposed on a substrate can be generated by any of the variety of methods described herein (e.g., by patterning or self-assembly of shrunken gel beads or by patterning or self-assembly of larger gel beads followed by shrinking), after which the high-density bead array is expanded (or re-expanded). When expanding, spatial constraints direct the beads to expand primarily in the Z dimension (away from the substrate), resulting in pillar arrays. In some embodiments, the gel beads of the pillar arrays have high aspect ratios. In some embodiments, aspect ratio of the expanded plurality of spatially-confined shrunken hydrogel beads is at least 2. In other embodiments, the aspect ratio of the expanded plurality of spatially-confined shrunken hydrogel beads is at least 3. In some embodiments, the plurality of spatially-confined shrunken hydrogel beads has an average aspect ratio of at least 4, 5, 6, 7, 8 or more.

[0718] In some embodiments, the method for the removal of water from a hydrogel is the same for each hydrogel (e.g., the first hydrogel, the second hydrogel, or the third hydrogel). In some embodiments, the method for the removal of water from one hydrogel (e.g., the first hydrogel) is different from the method for the removal of water for at least one other hydrogel (e.g., a second hydrogel, a third hydrogel, or a fourth hydrogel). For example, the method for the removal of water from one hydrogel can be different from the method for the removal of water for the other hydrogels (e.g., a second hydrogel, a third hydrogel, or a fourth hydrogel). In some embodiments, the method for the removal of water is different for each hydrogel (e.g., the first hydrogel, the second hydrogel, the third hydrogel, and the fourth hydrogel).

[0719] In some embodiments, the shrunken hydrogel is at least about 2-fold smaller in a linear dimension (e.g., along one axis) than the pre-shrunk hydrogel. For example, at least about 2.5, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, or more fold smaller in a

linear dimension than the pre-shrunk hydrogel.

[0720] In some embodiments, the size of the hydrogel is reduced along more than one axes, e.g., along 2 or 3 axes. In some embodiments, each axis intersects each other axis at 90 degrees. In some embodiments, the size of the hydrogel along the first axis is about 2 to about 10 or more fold smaller than the pre-shrunk hydrogel, e.g., about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10 or more fold smaller than the pre-shrunk hydrogel. In some embodiments, the size of the hydrogel along the second axis is about 2 to about 10 or more fold smaller than the pre-shrunk hydrogel, e.g., about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10 or more fold smaller than the pre-shrunk hydrogel. In some embodiments, the size of the hydrogel along the third axis is about 2 to about 10 or more fold smaller than the pre-shrunk hydrogel, e.g., about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10 or more fold smaller than the pre-shrunk hydrogel. In some embodiments, the reduction in the volume of the hydrogel is isometric.

[0721] In some embodiments, the volume of each hydrogel (e.g., a first hydrogel, a second hydrogel, a third hydrogel, or a fourth hydrogel) is the same. In some embodiments, the volume of at least one hydrogel is different. For example, in some embodiments, one hydrogel is different in volume from the other hydrogels (e.g., a second hydrogel, a third hydrogel, or a fourth hydrogel). In some embodiments, every hydrogel is different in volume from every other hydrogel.

[0722] In some embodiments, members of the plurality of features are cross-linked to a hydrogel (e.g., a first hydrogel, a second hydrogel, a third hydrogel, or a fourth hydrogel).

[0723] In one embodiment, features of an array can be copied into a hydrogel, and the volume of the hydrogel is reduced by removing water. These steps can be performed multiple times. For example, a method for preparing a high-density spatially-barcoded flexible array can include copying a plurality of spatially-barcoded features from an array into a first hydrogel, wherein the first hydrogel is in contact with the array; reducing the volume of the first hydrogel including the copied features by removing water, forming a first shrunken hydrogel including the copied features; copying the features in the first shrunken hydrogel into a second hydrogel, where the second hydrogel is in contact with the first hydrogel; and reducing the volume of the second hydrogel including the copied features by removing water, forming a second shrunken hydrogel including the copied features, thus generating a high-density spatially-barcoded array. In some instances, the array includes one or more pluralities of first oligonucleotides and the first hydrogel includes one or more pluralities of second oligonucleotides. Upon contacting the hydrogel with the array, members of the one or more pluralities of the first oligonucleotides can be attached to members of the one or more pluralities of second oligonucleotides. The array can include more species of first oligonucleotides than the number of species of the second oligonucleotides in the hydrogel such that first oligonucleotides comprising the same sequence can be coupled to second oligonucleotides comprising different sequences. The diversity of the oligonucleotides (e.g., spatial barcodes) in the first hydrogel can thereby be increased. The process of copying spatially-barcoded features from an array to a first hydrogel, removing water from the first hydrogel to form a first shrunken hydrogel, and copying spatially-barcoded features from the first shrunken hydrogel to one or more subsequent hydrogels can be performed multiple times (e.g., 2, 3, 4, 5, 6, 7, 8, 9, or 10 times). The result is a high-density flexible array including spatially-barcoded features.

[0724] In some embodiments, copying members of the plurality of features from an array includes copy by PCR. In some embodiments, the hydrogel (e.g., a first hydrogel, a second hydrogel, a third hydrogel, and/or a fourth hydrogel) comprises PCR reagents as described herein. In some embodiments, members of the plurality of features are copied using replica plating techniques (see, e.g., Mitra and Church, *Nucleic Acids Res.* 1999 Dec. 15; 27 (24): e34, which is incorporated by reference herein in its entirety). In some embodiments, after copying a plurality of features from an array into a first hydrogel, the features of the array are amplified in the first hydrogel (e.g., clonal amplification). In some embodiments, members of the plurality of features are copied into the first

hydrogel such that the pattern of the plurality of features of the first hydrogel is the same or substantially similar (e.g., at least 80%) to the pattern of the plurality of features of the array.

[0725] In some embodiments, one or more pluralities of features of the array are partitioned. For example, each partition can comprise a plurality of features different from the plurality of features of other partitions. For example, the members of the plurality of features are partitioned similar to the partitions of the plurality of features of the array. In some embodiments, the features of the array are copied into the first hydrogel, such that the volume or diameter of the pre-shrunk first hydrogel features are similar to the volume or diameter of the array features.

[0726] In some embodiments, the volume of a hydrogel comprising copied features is reduced, thus increasing the density of the copied features. In some embodiments, the copied features within a hydrogel further increases in density with each subsequent hydrogel copy and shrinking. For example, the density of the copied features of a second shrunken hydrogel is higher than the density of the copied features of a first shrunken hydrogel. Similarly, the density of the copied features of a third shrunken hydrogel is higher than the density of the copied features of a second shrunken hydrogel. Similarly, the density of the copied features of a fourth shrunken hydrogel is higher than the density of the copied features of a third shrunken hydrogel. In some embodiments, the volume of a partition of members of the plurality of features in a hydrogel is reduced when the volume of the hydrogel is reduced.

[0727] In some embodiments of the methods described herein, an array comprises shrunken gel features (e.g., beads). In some embodiments, the methods described herein generate shrunken gel bead arrays. In some embodiments, the shrunken gel beads of the array are shrunken hydrogel beads.

[0728] A “shrunken array” includes a plurality of spatially-barcoded features attached to, or embedded in, a substrate that have been reduced in volume (e.g., reduction in diameter or volume). A biological sample can be contacted with a shrunken array and further contacted with a solution capable of rehydrating the shrunken array. In some embodiments, analyte transfer and capture is driven by molecular diffusion. The process of rehydrating the shrunken array by providing a permeabilization solution or tissue stain to the sample can promote analytes (e.g., transcripts) present in the biological sample towards the spatially-barcoded features, thereby improving capture efficiency of the analytes. See, e.g., J. Vlassakis, A. E. Herr. “Effect of Polymer Hydration State on In-Gel Immunoassays.” *Anal. Chem.* 2015, 87 (21): 11030-8, herein incorporated by reference in its entirety.

[0729] A shrunken array can be generated with features (e.g., beads) containing spatial barcodes from an existing array. For example, an array (e.g., hydrogel bead array) described and prepared by any method herein can be contacted with reagents capable of dehydrating (e.g., removing water) the features (e.g., beads) to generate a shrunken array (e.g., a shrunken bead array). Methods of dehydrating features (e.g., beads) are known in the art. Any suitable method of dehydration (e.g., removing water) can be used. For example, in a non-limiting way, features (e.g., beads) can be dehydrated by a ketone, such as methyl ethyl ketone (MEK), isopropanol (IPA), acetone, 1-butanol, methanol (MeOH), dimethyl sulfoxide (DMSO), glycerol, propylene glycol, ethylene glycol, ethanol, (k) 1,4-dioxane, propylene carbonate, furfuryl alcohol, N,N-dimethylformamide (DMF), acetonitrile, aldehyde, such as formaldehyde or glutaraldehyde, or any combinations thereof. Additional dehydration agents include various salts, including inorganic salts (See, e.g., Ahmed, E. M., Hydrogel: Preparation, characterization, and applications: A review, *Journal of Advanced Research*, 6 (2) 105-121 (2015), which is incorporated herein by reference).

[0730] In some embodiments, the dehydrated features (e.g., beads) can create a shrunken array (e.g., shrunken bead array or shrunken hydrogel array) where the average diameter of the dehydrated features (e.g., beads) can be smaller than the average diameter of the features prior to dehydration. In some embodiments, the dehydrated features (e.g., beads) can have an average diameter at least two-fold smaller than the average diameter of the features prior to dehydration. In

some embodiments, the dehydrated features (e.g., beads) can have an average diameter at least three-fold smaller than the average diameter of the features prior to dehydration. In some embodiments, the dehydrated features (e.g., beads) can have an average diameter at least four-fold or smaller than the average diameter of the features (e.g., beads) prior to dehydration.

[0731] After generating a shrunken array, a biological sample (e.g., tissue sample) can be contacted with the shrunken array (e.g., shrunken bead array). A rehydrating solution can be provided to the biological sample and the shrunken array by any suitable method (e.g., by pipetting). The rehydrating solution can contain reagents to rehydrate (e.g., water or buffers) the features (e.g., beads) of the shrunken array. In some embodiments, the rehydrating solution can be applied to the entire biological sample. In some embodiments, the rehydrating solution can be selectively applied (e.g., to a region of interest). In some embodiments, absorbing water from the rehydrating solution can increase the diameter of at least one feature (e.g., bead) in the shrunken array. In some embodiments, the rehydrating solution can increase the diameter of at least one feature (e.g., bead) by at least two-fold. In some embodiments, the rehydrating solution can increase the diameter of at least one feature (e.g., bead) by at least three-fold. In some embodiments, the rehydrating solution can increase the diameter of at least one feature (e.g., bead) by at least four-fold. In some embodiments, the rehydrating solution can increase the diameter of at least one feature (e.g., bead) by at least five-fold or more.

[0732] In some embodiments, the rehydrating solution can contain permeabilization reagents. The biological sample can be permeabilized using permeabilization reagents and techniques known in the art or otherwise described herein. Biological samples from different sources (e.g., brain, liver, ovaries, kidney, breast, colon, etc.) can require different permeabilization treatments. For example, permeabilizing the biological sample (e.g., using a protease) can facilitate the migration of analytes to the substrate surface (e.g., spatially-barcoded features). In some embodiments, the permeabilization reagents can be a detergent (e.g., saponin, Triton X-100™, Tween-20™). In some embodiments, an organic solvent (e.g., methanol, acetone) can permeabilize cells of the biological sample. In some embodiments, an enzyme (e.g., trypsin) can permeabilize the biological sample. In another embodiment, an enzyme (e.g., collagenase) can permeabilize the biological sample.

[0733] In some embodiments the solution can permeabilize the biological sample and rehydrate the features (e.g., beads) of the shrunken array (e.g., shrunken hydrogel). In some embodiments, the rehydrating solution can stain the biological sample and rehydrate the features of the shrunken array (e.g., beads).

[0734] In some embodiments, the rehydrating solution (e.g., permeabilization or stain solution) can diffuse through the biological sample. In some embodiments, the rehydrating solution can reduce diffusion of analytes away from the substrate. In some embodiments, while diffusing through the biological sample, the rehydrating solution can migrate analytes toward the substrate surface and improve the efficiency of analyte capture.

(viii) Microcapillary Arrays

[0735] A “microcapillary array” is an arrayed series of features that are partitioned by microcapillaries. A “microcapillary channel” is an individual partition created by the microcapillaries. For example, microcapillary channels can be fluidically isolated from other microcapillary channels, such that fluid or other contents in one microcapillary channel in the array are separated from fluid or other contents in a neighboring microcapillary channel in the array. The density and order of the microcapillary channels can be any suitable density or order of discrete sites.

[0736] In some embodiments, microcapillary arrays are treated to generate conditions that facilitate loading. An example is the use of a corona wand (BD-20AC, Electro Technic Products) to generate a hydrophilic surface. In some embodiments, a feature (e.g., a bead with capture probes attached) is loaded onto a microcapillary array such that the exact position of the feature within the array is known. For example, a capture probe containing a spatial barcode can be placed into a

microcapillary channel so that the spatial barcode can enable identification of the location from which the barcoded nucleic acid molecule was derived. In some embodiments, features are introduced to the microcapillary array by flowing the features through microcapillary channels. In some embodiments, the microcapillary channel can reduce the cross-sectional area of a feature.

[0737] In some embodiments, when random distribution is used to distribute features, empirical testing can be performed to generate loading/distribution conditions that facilitate a single feature per microcapillary. In some embodiments, it can be desirable to achieve distribution conditions that facilitate only a single feature (e.g., bead) per microcapillary channel. In some embodiments, it can be desirable to achieve distribution conditions that facilitate more than one feature (e.g., bead) per microcapillary channel, by flowing the features through the microcapillary channel.

[0738] In some embodiments, the microcapillary array is placed in contact with a sample (e.g., on top or below) so that microcapillaries containing a feature (e.g., a bead, which can include a capture probe) are in contact with the biological sample. In some embodiments, a biological sample is placed onto an exposed side of a microcapillary array and mechanical compression is applied, moving the biological sample into the microcapillary channel to create a fluidically isolated reaction chamber containing the biological sample.

[0739] In some embodiments, a biological sample is partitioned by contacting a microcapillary array to the biological sample, thereby creating microcapillary channels including a bead and a portion of the biological sample. In some embodiments, a portion of a biological sample contained in a microcapillary channel is one or more cells. In some embodiments, a portion of a biological sample contained in a microcapillary channel is one cell. In some embodiments, a feature is introduced into a microcapillary array by flow after one or more cells are added to a microcapillary channel.

[0740] In some embodiments, reagents are added to the microcapillary array. The reagents can include enzymatic reagents or reagent mixtures for performing amplification of a nucleic acid. In some embodiments, the reagents include a reverse transcriptase, a ligase, one or more nucleotides, or any combinations thereof. One or more microcapillary channels can be sealed after reagents are added to the microcapillary channels, e.g., by using silicone oil, mineral oil, a non-porous material, or lid. In some embodiments, the microcapillary array is incubated in a humidified chamber. In some embodiments, the microcapillary array is incubated for an amount of time and at a temperature conducive to allowing amplification of a nucleic acid to occur.

[0741] In some embodiments, a reagent solution is removed from each microcapillary channel following an incubation for an amount of time and at a certain temperature or range of temperatures, e.g., following a hybridization or an amplification reaction. Reagent solutions can be processed individually for sequencing, or pooled for sequencing analysis. In some embodiments, the sequencing information from the pooled reaction solution is spatial information for one or more biological analytes.

(ix) Hydrogel/Well Arrays

[0742] In some embodiments are methods for generating patterned hydrogel arrays using wells (e.g., a nanowell or microwell array). In some embodiments, the well is a 3-dimensional structure. In some embodiments, the top view of a well is any suitable 2-dimensional shape, which when extended along the z-axis, produces a 3-dimensional structure capable of containing one or more features (e.g., beads) and/or reagents. Non-limiting examples of wells which may form an array include a triangular prism, a square or rectangular prism, a pentagonal prism, a hexagonal prism, a heptagonal prism, an octagonal prism, an n-sided prism, or a cylindrical array (e.g., “microcapillary array”). In some embodiments, a well of the well array shares at least one well wall (or a portion of the well wall, if a microcapillary array) with an adjacent well. In some embodiments, a well does not share any walls or portion of a wall in common with another well of the array. In some embodiments, the well array is attached to a substrate, such that the wells of the well array are fluidically isolated from each other. In some embodiments, one end of the well array is open (e.g.,

exposed), wherein the open end can be used to distribute features or reagents into the well. [0743] In some embodiments, the method includes providing shrunken (e.g., dehydrated) hydrogel features (e.g., beads) to a well array. The hydrogel features can be dehydrated (e.g., removing water) by any of the variety of methods described herein. The features, described elsewhere herein, can be provided such that the number of features is less than the number of wells of the array, the features can be provided such that the number of features equals the number of wells of the array, or the features can be provided in excess of the number of wells of the array. In some embodiments, the well array is manipulated such that one or more shrunken hydrogel features move from the top surface of the array down into a well. For example, a well array can be placed on a shaker for a length of time necessary for the features to distribute into the wells. Other, non-limiting examples of manipulations that can cause a shrunken hydrogel feature to enter a well are physically shaking, tilting, or rolling the well array, or a combination thereof; using forced air to blow features into a well, using a magnet to pull down hydrogel features comprising magnetic particles, using microfluidic systems to distribute features into wells, using a printer to deposit a feature into a well, or any other method to distribute features into wells. In some embodiments, once a well contains a feature, the well cannot accept or retain another feature. In other embodiments, a well can contain more than one feature.

[0744] In some embodiments, the method includes rehydrating (e.g., adding water) the shrunken hydrogel features, wherein the shrunken hydrogel features are located in the wells. Rehydrating shrunken hydrogel features can be accomplished by any method described herein. Rehydrating a shrunken hydrogel feature in the well can cause the shrunken hydrogel feature to expand. In some embodiments, the shrunken hydrogel feature expands to fill the well. In some embodiments, the shrunken hydrogel feature expands in a z direction, such that the feature expands out of the unenclosed (i.e. open) end of the well. The exposed area of the rehydrated feature can create a patterned hydrogel array (e.g., a well array). A rehydrated feature contained within a well can be stable. In some embodiments, a rehydrated feature (e.g., a rehydrated shrunken hydrogel feature) is immobilized within a well, such that typical array usage does not dissociate the rehydrated feature from the well. The patterned hydrogel array can be used for analyte capture according to the methods described herein.

(x) Bead Tethering

[0745] “Bead tethering” can refer to an arrangement of beads, wherein the arrangement may or may not form an array. The tethered beads can be contacted with a sample and processed according to methods described herein. Further, contacting a biological sample with a single bead or beads tethered together in various arrangements can allow for more precise spatial detection of analytes, e.g., a region of interest. Methods for tethering beads together are known in the art. Some suitable, non-limiting, methods of tethering beads together can be, e.g., chemical linkers, proximity ligation, or any other method described herein. In some embodiments, beads can be tethered together independent of a substrate. In some embodiments, beads can be tethered in various arrangements on an existing substrate. In some embodiments, a substrate (e.g., a hydrogel) can be formed around existing tethered beads. In some embodiments, the beads or bead arrangement can contact a portion of the biological sample. In some embodiments, the bead or bead arrangement can contact a region of interest. In some embodiments, the beads or bead arrangement can contact the entire biological sample. In some embodiments, the beads or bead arrangements are contacted to random positions on the biological sample. In some embodiments, the beads are contacted to according to a specific pattern on the biological sample.

[0746] Beads can be tethered together in various arrangements. In some embodiments, a single (e.g., one) bead can be contacted with a biological sample. In some embodiments, two or more beads can be tethered (e.g., connected to each other), in various arrangements. For example, in a non-limiting way, beads can be tethered together to form a cluster, a row, or arranged on a mesh (e.g., a net). In some embodiments, at least three beads can be tethered together in a two-

dimensional (2D) array (e.g., a cluster). In some embodiments, at least two beads can be tethered together in a one-dimensional (1D) array (e.g., a row). In such embodiments, the beads are arranged in such fashion that the beads can contact each other directly. In some embodiments, at least two beads can be tethered together in a string arrangement. In such embodiments, the beads are arranged in such fashion that the beads can contact each other indirectly (e.g., beads are connected via linker). In some embodiments, at least two beads can be tethered together in a mesh arrangement (e.g., net). In some embodiments, beads tethered together in a 2D array, a 1D array, the beads on a string arrangement, and the beads on the mesh arrangement can be used in any combination with each other on the biological sample.

[0747] In some embodiments, at least about 2 to about 10 beads can be tethered together in various arrangements. In some embodiments at least about 3, about 4, about 5, about 6, about 7, about 8, about 9, or more beads can be tethered together. In some embodiments, about 10 to about 100 beads can be tethered together in various arrangements. In some embodiments, about 10, about 20, about 30, about 40, about 50, about 60, about 70, about 80, about 90, or more beads can be tethered together in various arrangements. In some embodiments, about 100 to about 1,000 beads can be tethered together in various arrangements. In some embodiments, about 100, about 200, about 300, about 400, about 500, about 600, about 700, about 800, or about 900 or more beads can be tethered together in various arrangements. In some embodiments, about 1000 to about 10000 beads can be tethered together in various arrangements. In some embodiments, about 1000, about 2000, about 3000, about 4000, about 5000, about 6000, about 7000, about 8000, about 9000, about 10000 or more beads can be tethered together.

[0748] In some embodiments, the tethered beads can have capture probes comprising spatial barcodes, functional domains, unique molecular identifiers, cleavage domains, and capture domains, or combinations thereof. In some embodiments, each bead can be associated with a unique spatial barcode. In some embodiments, the spatial barcode is known prior to contacting the bead or bead arrangement to the biological sample. In some embodiments, the spatial barcode is not known prior to contacting the bead or bead arrangement to the biological sample. The identity of each bead (e.g., spatial barcode) in the array can be deconvolved, for example, by direct optical sequencing, as discussed herein.

(xi) Printing Arrays in Liquid

[0749] In some embodiments, an array can be printed in liquid. The resolution of conventionally-printed arrays can be limited, due to the diffusion of printed solutions. Printing the array in a highly viscous liquid can increase resolution by preventing the diffusion of the printed solution. Thus, disclosed herein are various methods and materials for attaching and/or introducing a capture probe (e.g., a nucleic acid capture probe) having a barcode (e.g., a spatial barcode) to a substrate (e.g., a slide), wherein the attaching (e.g., printing) is performed in liquid.

[0750] In some aspects, capture probes are printed on a substrate (e.g., a slide or bead). In some aspects, the substrate is a slide. In some aspects, the substrate is a 96-well microtiter plate. In some aspects, methods provided herein can also be applied to other substrates commonly used for nucleic acid analyses, including but not limited to beads, particles, membranes, filters, dipsticks, slides, plates, and microchips. In some aspects, such substrates may be composed of a number of materials known to be compatible with nucleic acid analysis, including but not limited to agarose, styrene, nylon, glass, and silicon.

(1) First Solution

[0751] In some embodiments, provided herein are methods of printing arrays on substrates using one or more liquid solution(s) (e.g., two or more solutions that include distinct capture probes). In some aspects, methods of printing arrays on substrates using one or more solution(s) can improve the resolution of the printed array. In some aspects, methods provided herein include dispensing a first solution (e.g., bulk solution) onto a substrate. In some aspects, the first solution (e.g., bulk solution) has a lower Reynolds Number relative to a second solution (e.g., a second solution that

includes capture probes to be attached to the substrate). The Reynolds Number represents an inverse relationship between the density and velocity of a fluid and its viscosity in a channel of given length. More viscous, less dense, and/or slower moving fluids will have a lower Reynolds Number, and are easier to divert, stop, start, or reverse without turbulence. In some embodiments, the first solution and the second solution are immiscible.

[0752] In some aspects, the first (e.g., bulk) solution is hydrophobic. In some aspects, after dispensing the first (e.g., bulk) solution onto the slide, the first (e.g., bulk) solution remains on the slide in discrete spatial areas on the slide. In some aspects, the first (e.g., bulk) solution is made of a solution that does not denature one or more probes and/or does not inhibit probe-to-substrate binding. In some embodiments, the bulk solution can include an aqueous solution, a high viscosity solution, or a low nucleic acid diffusivity solution. In some aspects, the first (e.g., bulk) solution is a gel. In some aspects, the first (e.g., bulk) solution is a hydrogel. In some aspects, the first (e.g., bulk) solution includes natural polymers, including for example, glycerol, collagen, gelatin, sugars such as starch, alginate, and agarose, or any combinations thereof. In some aspects, the first (e.g., bulk) solution includes a synthetic polymer. In some aspects, the synthetic polymer is prepared by any method known in the art, including for example, chemical polymerization methods. In some aspects, the gel or polymer is hydrophobic. In some aspects, the gel or polymer is hydrophilic. In some aspects, the gel or polymer is aqueous. In some aspects, the gel or polymer shrinks at room temperature. In some aspects, the gel or polymer shrinks when heated. In some aspects, the polymer is a film that shrinks when heated.

[0753] In some aspects, the first (e.g., bulk) solution includes glycerol. In some aspects, glycerol is present in the first (e.g., bulk) solution at a concentration of 5-100%. In some aspects, glycerol is present in the solution at a concentration of 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, or 70%.

[0754] In some aspects, the first (e.g., bulk) solution includes sugar. In some aspects, the sugar is a monosaccharide, a disaccharide, a polysaccharide, or combinations thereof. In some aspects, the sugar is glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, raffinose, starch cellulose, or combinations thereof. In some aspects, sugar is sourced from complex compounds such as molasses or other by-products from sugar refinement. In some aspects, sugar is present in the first (e.g., bulk) solution at a concentration of 5-100%. In some aspects, sugar is present in the solution at a concentration of 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, or 70%.

[0755] In some aspects, the first (e.g., bulk) solution has a viscosity that is about 0.1 $\times$ -fold, 0.2 $\times$ -fold, 0.3 $\times$ -fold, 0.4 $\times$ -fold, 0.5 $\times$ -fold, 0.6 $\times$ -fold, 0.7 $\times$ -fold, 0.8 $\times$ -fold, 0.9 $\times$ -fold, 1.0 $\times$ -fold, 1.1 $\times$ -fold, 1.2 $\times$ -fold, 1.3 $\times$ -fold, 1.4-fold, 1.5 $\times$ -fold, 1.6 $\times$ -fold, 1.7 $\times$ -fold, 1.8 $\times$ -fold, 1.9 $\times$ -fold, 2.0 $\times$ -fold, 2.5 $\times$ -fold, 3.0 $\times$ -fold, 4.0 $\times$ -fold, 5.0 $\times$ fold, 6.0 $\times$ -fold, 7.0 $\times$ -fold, 8.0 $\times$ fold, 9.0 $\times$ -fold, or 10 $\times$ -fold greater than the viscosity of the second solution.

(2) Second Solution

[0756] In some embodiments, printing arrays on substrates using two or more solutions includes using a second solution. In some embodiments, the second solution can include one or more capture probes. In some embodiments, the second solution is dispensed in the form of a droplet. Some embodiments include a plurality of second solutions, wherein each member of the plurality includes one or more capture probes comprising a spatial barcode unique to that particular droplet. In some embodiments, the second solution is dispensed onto a substrate covered or partially covered with the first (e.g., bulk) solution. In some embodiments, the first (e.g., bulk) solution reduces or prevents the diffusion of one or more capture probes from the second solution. In some aspects, the one or more capture probes remain entirely within the second solution when printed onto a substrate covered or partially covered with the first (e.g., bulk) solution.

[0757] In some aspects, the second solution includes one or more capture probes (e.g., any of the capture probes disclosed herein). In some aspects, the second solution includes one spatially-

barcoded capture probe. In some aspects the second solution includes at least 5, at least 10, at least 20, at least 30 at least 40, at least 50, at least 75, at least 100, at least 200, at least 300, at least 400, at least 500, at least 600, at least 700, at least 800, at least 900, at least 1,000, at least 1,500, at least 2,000, at least 3,000, at least 4,000, or at least 5,000 spatially-barcoded capture probes. In some aspects, the second solution includes compounds that facilitate binding of the one or more capture probes to the substrate. In some aspects, the second solution does not inhibit the one or more capture probes from binding the substrate and/or does not denature the one or more capture probes. In some aspects, the second solution is less viscous than the first (e.g., bulk) solution. In some aspects, the second solution and the first (e.g., bulk) solution are wholly or substantially immiscible (e.g., they do not mix). In some aspects, the second solution is an aqueous solution. In some aspects, the second solution is a hydrophilic solution.

(3) Dispensing

[0758] In some embodiments, spot printing a high-density pattern of features can include dispensing the oligonucleotides and/or features, in the form of a liquid droplet, onto the surface of the substrate in the presence of a bulk solution. In some embodiments, the second solution droplet(s) (e.g., oligonucleotides and/or feature droplet(s)) and the first (e.g., bulk) solution does not substantially mix with each other. In some aspects, the printing methods disclosed herein that include dispensing a second solution including one or more spatially-barcoded capture probes onto a substrate in the presence (e.g., through) a first (e.g., bulk) solution result in a spot size of the second solution (e.g., a cross-sectional spot size of the second solution on the plane of the substrate) that is smaller compared to the spot size that would be obtained by dispensing the same second solution onto the substrate in the absence of the first (e.g., bulk) solution. In some aspects, the spot size (e.g., the cross-sectional spot size of the second solution on the plane of the substrate) of the second solution does not increase after printing the second solution onto the substrate in the presence of the first (e.g., bulk) solution. In some aspects, the area of the spot of the second solution does not change after printing the second onto the substrate in the presence of the first (e.g., bulk) solution. In some aspects, printing of the second solution onto the substrate in the presence of the first (e.g., bulk) solution results in a desired pattern on the substrate surface. For example, a plurality of second solutions can be printed onto the substrate in the presence of the first solution such that the locations where the plurality of second solutions are printed results in a desired pattern on the substrate. In some embodiments, two or more members of a plurality of second solutions printed onto a substrate include distinct populations of capture probes, which capture probes attach to the substrate, such that an array of capture probes is generated.

(4) Density

[0759] In some embodiments, the cross-sectional area of the oligonucleotide and/or feature droplet(s) on the substrate is smaller than a corresponding cross-sectional area of the oligonucleotide and/or feature droplet(s) that would be generated by dispensing the oligonucleotides and/or features onto the surface of the substrate in the absence of the bulk solution. In some embodiments, the cross-sectional area of the oligonucleotide and/or feature droplet(s) on the substrate is about two-fold smaller than the corresponding cross-sectional area of the oligonucleotide and/or feature droplet(s) that would be generated by dispensing the oligonucleotide and/or feature droplet(s) onto the surface of the substrate in the absence of the bulk solution.

[0760] In some aspects, the cross-sectional area of the second solution on the substrate generated by dispensing the second solution onto the surface of the substrate in the presence of the first solution is smaller than a corresponding cross-sectional area of the second solution that would be generated by dispensing the second solution onto the surface of the substrate in the absence of the first solution. In some aspects, the cross-sectional area of the second solution on the substrate generated by dispensing the second solution onto the surface of the substrate in the presence of the first solution is about two-fold, about three-fold, about four-fold, or about five-fold, about 10-fold,

about 20-fold, about 30-fold, about 40-fold, about 50-fold, about 60-fold, about 70-fold, about 80-fold, about 90-fold, or about 100-fold smaller than the corresponding cross-sectional area of the second solution that would be generated by dispensing the second solution onto the surface of the substrate in the absence of the first solution.

(5) Curing

[0761] Any suitable technique or condition can be used to cure the first solution, the second solution, and/or the spot formed by the second solution after the second solution (including the capture probes) is dispensed onto the substrate. As used herein, the term “curing” and related terms can refer to treating a solution (e.g., a first solution, a second solution, or both) with an agent and/or condition that transforms a solution from a liquid state to a solid state (e.g., transformed into a matrix), wherein the solution retains a three dimensional shape after the curing process. Suitable examples of curing techniques and/or conditions include ultraviolet (UV) radiation, infrared (IR) radiation, thermal radiation, microwave radiation, visible radiation, narrow-wavelength radiation, laser light, natural light, humidity, or combinations thereof. Suitable examples of a curing source include, for example, a UV light, a heating device, a radiation device, a microwave device, a plasma device, or combinations thereof.

[0762] In some embodiments, the first solution and/or the second solution is chemically cured. In some embodiments, the oligonucleotide and/or feature droplet(s) are chemically cured. In some embodiments, the bulk solution is chemically cured. In some aspects, the spot formed by the second solution after the second solution (including the capture probes) is dispensed onto the substrate is chemically cured. In some embodiments, the oligonucleotides and/or features and the bulk solution are attached to the substrate (e.g., by curing), thus generating a feature and bulk solution-matrix. In some aspects, the matrix (e.g., the first and second solution-matrix) is chemically cured. Chemically curing a solution can be accomplished by any means known in the art. For example, a solution can include one or more hydrogel subunits that can be chemically polymerized (e.g., cross-linked) to form a three-dimensional (3D) hydrogel network. Features dispensed, in the form of a liquid droplet, onto the surface of a substrate in the presence of a bulk solution can be polymerized. In some embodiments, the features are co-polymerized with the bulk solution to create a gel feature in hydrogel flexible array, whereas in other embodiments the gel pad or feature is polymerized and the bulk solution is removed leaving a spot or bead array. Non-limiting examples of hydrogel subunits include acrylamide, bis-acrylamide, polyacrylamide and derivatives thereof, poly(ethylene glycol) and derivatives thereof (e.g., PEG-acrylate (PEG-DA), PEG-RGD), gelatin-methacryloyl (GelMA), methacrylated hyaluronic acid (MeHA), polyaliphatic polyurethanes, polyether polyurethanes, polyester polyurethanes, polyethylene copolymers, polyamides, polyvinyl alcohols, polypropylene glycol, polytetramethylene oxide, polyvinyl pyrrolidone, polyacrylamide, poly(hydroxyethyl acrylate), and poly(hydroxyethyl methacrylate), collagen, hyaluronic acid, chitosan, dextran, agarose, gelatin, alginate, protein polymers, methylcellulose, and the like, or combinations thereof. Other materials and techniques useful in forming and/or cross-linking hydrogels are described in more detail herein.

[0763] In some aspects, the first solution and/or second solution is photo-reactively cured. In some embodiments, the oligonucleotide and/or feature droplet(s) are photo-reactively cured. In some embodiments, the bulk solution is photo-reactively cured. In some aspects, the spot formed by the second solution after the second solution including the capture probe is dispensed onto the substrate is photo-reactively cured. In some aspects, the matrix (e.g., the first and second solution-matrix) is photo-reactively cured. Photo-reactive curing can be accomplished by any means known in the art.

[0764] In some aspects, methods disclosed herein include curing the first (e.g., bulk) and/or second solution. In some aspects, methods disclosed herein include curing the spot formed by the second solution after the second solution (e.g., a second solution including the capture probe) is dispensed onto the substrate. In some aspects, methods disclosed herein include curing the first solution and

the second solution after both are dispensed onto the slide. In some embodiments, curing the first and second solutions results in a first-and-second solution matrix. In some aspects, methods disclosed herein include curing the second solution followed by removal of the uncured first solution from the substrate. In some aspects, the methods disclosed herein include curing the first and second solutions on the substrate, thus generating a matrix (e.g., a first and second solution-matrix).

(6) Expanding Matrices

[0765] In some embodiments, prior to dispensing the second solution droplet(s) (e.g., oligonucleotide and/or feature droplet(s)) onto the surface of a substrate in the presence of a first (e.g., bulk) solution, the first (e.g., bulk) solution is cured to form a bulk solution-matrix, and the bulk solution matrix is reversibly expanded along one or more axes (e.g., one or more axes of a matrix or a substrate surface). In some embodiments, second solution droplet(s) can be dispensed in the presence of expanded first (e.g., bulk) solution-matrix.

[0766] In some embodiments, dispensing a plurality of second solution droplet(s) (e.g., including different capture probes having different spatial barcodes) onto a substrate covered with a first (e.g., bulk) solution, wherein the first solution is initially stretched along one or more axes. In some embodiments, the stretched first solution is cured. In some embodiments, the stretched first solution is partially cured. In some embodiments, the stretched first solution is uncured and is dispensed on a surface that itself is stretched. In some aspects, the volume (e.g., cross-sectional area) of the second solution droplet(s) is decreased after being dispensed into a first solution that is initially stretched. In some aspects, disclosed herein are methods of preparing an array including (i) providing a gel or polymer (e.g., a cured or partially cured solution) onto a substrate, (2) stretching the gel or polymer, (3) dispensing a droplet of the second solution onto the substrate while the gel or polymer is stretched, and (4) allowing the gel or polymer to relax, thereby decreasing the overall area (e.g., cross-sectional area) of the second solution droplet(s) on the substrate. In some aspects, the stretching step includes reversibly expanding the gel or polymer along one or more axes coplanar with the surface of the substrate.

(7) Removing Solution

[0767] In some embodiments, the first (e.g., bulk) solution can be removed from the substrate after the oligonucleotide and/or feature droplet(s) are attached to the substrate. In some aspects, the first (e.g., bulk) solution is removed after the second solution including the one or more capture probes (e.g., plurality of second solutions that include different capture probes) is dispensed onto the substrate. Methods of removing the first (e.g., bulk) solution are known in the art. In some aspects, removal of the first (e.g., bulk) solution results in complete removal of the first solution, leaving only the second solution including the one or more capture probes (e.g., plurality of second solutions that include different capture probes) on the substrate. In some aspects, removal of the first (e.g., bulk) solution does not change the surface area of the second solution (e.g., plurality of second solutions that include different capture probes) in contact with the substrate. In some aspects, removal of the first (e.g., bulk) solution does not change the shape of the droplet of the second solution (e.g., plurality of second solutions that include different capture probes) in contact with the substrate. In some aspects, prior to removal of the first (e.g., bulk) solution, the second solution including the one or more probes (e.g., plurality of second solutions that include different capture probes) is cured by methods disclosed herein but that the first (e.g., bulk) solution is not cured. For example, the first and second solutions can be subjected to agents and/or conditions under which the second solution (e.g., plurality of second solutions that include different capture probes) is cured, while the first (e.g., bulk) solution is not cured. In some embodiments, the second solution (e.g., plurality of second solutions that include different capture probes) includes one or more hydrogel subunits that can be polymerized (e.g., cross-linked) to form a three-dimensional (3D) hydrogel network, while the first (e.g., bulk) solution does not include such one or more hydrogel subunits. Upon subjecting the first and second solutions to curing conditions, only the

second solution(s) will be cured, allowing the first solution to be removed.

[0768] In some embodiments, the first (e.g., bulk) solution is not removed from the substrate after the oligonucleotides and/or features are attached to the substrate.

(8) Shrinking Droplet/Feature Arrays

[0769] In some embodiments, an expanded first (e.g., bulk) solution-matrix containing a second solution (e.g., oligonucleotide and/or feature droplet(s)) can be shrunk along one or more axes (e.g., of the matrix or of the substrate surface) such that the cross-sectional area of the second solution droplet(s) (e.g., oligonucleotide and/or feature droplet(s)) is smaller than a corresponding cross-sectional area of a second solution droplet(s) e.g., oligonucleotide and/or feature droplet(s)) that would be generated if the first (e.g., bulk) solution-matrix containing the second solution droplet(s) (e.g., oligonucleotide and/or feature droplet(s)) were not shrunk along the one or more axes. In some embodiments, shrinking the matrix (e.g., the first-and-second solution matrix) generates a shrunken matrix (e.g., a shrunken first-and-second solution matrix), wherein the volume of a second solution droplet (e.g., a plurality of second solution droplets) of the matrix is reduced as compared to the volume of the second solution droplet in a non-shrunken matrix. In some aspects, shrinking two or more second solutions (e.g., droplets of second solutions) results in a higher density of the two or more second solutions. In some embodiments, the second solution droplet(s) (e.g., oligonucleotide and/or feature droplet(s)) and first (e.g., bulk) solution-matrix can be shrunk by any method disclosed herein. In some embodiments, the shrinking can include removing water. In some aspects, the resolution of the array is increased after the droplet is shrunk. In some aspects, shrinking the second solution droplet results in a decrease in the cross-sectional area of the droplet (e.g., the cross-sectional area in the plane of the substrate onto which the second solution droplet is printed or dispensed). In some aspects, after shrinking the droplet, the concentration of probes on the substrate will be increased, thereby improving sensitivity.

[0770] The spatial array is contacted to the biological sample, wherein the biological sample can be any described herein (e.g., a FFPE tissue section). Once the spatial array has been placed on the biological sample, a cellular and/or nuclear permeabilization reaction can occur, such that the biological analytes (e.g., DNA, RNA, proteins, metabolites, small molecules, lipids, and the like) are released and captured onto the spatial array, preserving their spatial information. The spatial array is removed, and the molecular information therein is determined (e.g., by performing library construction for next generation sequencing). Sequencing can be followed by correlation of the expression value (e.g., gene expression of the analyte) with the feature.

[0771] In some aspects, the cross-sectional area of a second solution droplet is decreased upon shrinking by about 10%, by about 20%, by about 30%, by about 40%, by about 50%, by about 60%, by about 70%, by about 80%, or by about 90% compared to the cross-sectional area of a second solution droplet that is printed or dispensed onto the substrate. In some aspects, the cross-sectional area of a second solution droplet is decreased by about 1.0-fold, by about 1.1-fold, by about 1.2-fold, by about 1.3-fold, by about 1.4 fold, by about 1.5-fold, by about 2-fold, by about 3-fold, by about 4-fold, or by about 5-fold compared to the cross-sectional area of a second solution droplet that is printed or dispensed onto the substrate.

[0772] In some aspects, disclosed herein are methods of preparing an array including dispensing a plurality of second solution droplets of (e.g., including different capture probes having different spatial barcodes) onto the substrate covered with a first (e.g., bulk) solution, curing the first and second solutions to generate a matrix (e.g., a first and second solution-matrix), wherein the volume and shape of the plurality of second solution droplets in the matrix are substantially the same as when the plurality of second solution droplets was dispensed, and shrinking the matrix, thereby decreasing the overall volume (e.g., cross-sectional area) of the plurality of second solution droplets. In some aspects, shrinking can be accomplished using any of the variety of shrinking agents or conditions described herein. In some aspects, shrinking can be accomplished by dehydrating the matrix.

[0773] In some aspects, after the second solution is dispensed, the area of the second solution is shrunk along the one or more axes of coplanar with the surface of the substrate such that the cross-sectional area of the second solution on the substrate (e.g., the slide) is smaller than a corresponding cross-sectional area of the second solution that would be generated if the first solution matrix including the second solution were not shrunk along the one or more axes of the surface of the substrate.

[0774] With respect to orientation, shrinkage need not be equal in any two orthogonal directions on the substrate. However, in some aspects, the shrinkage of a second solution droplet is substantially uniform in shrinkage. In some aspects, a second solution droplet shrinks in substantially the same amount in each direction, regardless of position on the substrate plane.

[0775] In some aspects, after preparing the substrate with the one or more capture probes (e.g., a spatial array that includes the capture probes), disclosed herein are methods for spatially profiling an analyte (e.g., a plurality of analytes) in a biological sample, including (a) generating a spatial array including a plurality of capture probes bound to a substrate; wherein (i) at least a portion of the substrate is coated with a first solution; and (ii) a plurality of second solutions, in the form of a liquid droplets, are dispensed onto the surface of the substrate in the presence of the first solution, wherein the first solution and the plurality of second solutions do not substantially mix with each other, wherein at least two members of the plurality of the second solutions include one or more different capture probes having different spatial barcodes, and wherein at least one of the one or more capture probes of at least two members of the plurality of second solutions is bound to the substrate; (b) contacting the biological sample with the spatial array such that the analyte(s) present in the biological sample are captured by one or more of the capture probes; and (c) determining the spatial profile of the captured analyte(s) in the biological sample.

(xii) Building Arrays with Linkers

[0776] In some embodiments, an array includes a linker that is branched or dendrimeric. In some embodiments, the linker includes dendrimeric polyamines. Non-limiting examples of dendrimeric polyamines include symmetrical and unsymmetrical dendrimers, nucleic acid dendrimers, polyamidoamine dendrimers, and lysine-based dendrimers. In some embodiments, a linker is a nucleic acid dendrimer. In some embodiments, a nucleic acid dendrimer includes individual molecules of nucleic acids that share a region of at least partial complementarity located in a position of each nucleic acid molecule where upon binding at least 2 (e.g., 2, 3, 4, or more) single-strand overhangs are generated. In some embodiments, the four single-stranded overhangs are designed to interact with additional nucleic acid dendrimers or specific complementary sequence (e.g., sequences on capture probes). In some embodiments, the dendrimeric linkers (e.g., nucleic acid dendrimers) are affixed to a substrate (e.g., affixed using any of the methods described herein (e.g., covalent bonding or physical absorption)). For example, a nucleic acid dendrimer is affixed to a substrate and also interacts with capture probes via the single-stranded overhangs. In another example, a nucleic acid dendrimer is affixed to a substrate and also interacts with additional nucleic acid dendrimers. The additional nucleic acid dendrimers can also interact with capture probes via the single-strand overhangs not already interacting with other nucleic acid dendrimers. In such cases, with each additional layer of nucleic acid dendrimer, there is the potential to increase the number of sites to which capture probes can interact, thereby increasing the number of capture domains (e.g., any of the exemplary capture domains described herein) on the array.

(xiii) Array/Feature Preservation

[0777] In some embodiments, the biological sample can be preserved after completion of an assay with a feature or arrangement of features for additional rounds of spatial detection of analytes. In some embodiments, the biological sample, features, array, or any combination thereof can be preserved after the spatial profiling. In some embodiments, the biological sample, features, array, or combinations thereof can be protected from dehydration (e.g., drying, desiccation). In some embodiments, the biological sample, features, array, or combinations thereof, can be protected from

evaporation. Methods of preserving and/or protecting biological samples, features, or arrays are known in the art. For example, in a non-limiting way, the biological sample, features, array, or combinations thereof can be covered by a reversible sealing agent. Any suitable reversible sealing agent can be used. Methods of reversible sealing are known in the art (See, e.g., WO 2019/104337, which is incorporated herein by reference). In a non-limiting way, suitable reversible sealing agents can include non-porous materials, membranes, lids, or oils (e.g., silicone oil, mineral oil). In further non-limiting examples, the biological sample, features, array, or combinations thereof can be preserved in an environmental chamber (e.g., hermetically sealed) and removed for additional rounds of spatial analysis at a later time.

(e) Analyte Capture

[0778] In this section, general aspects of methods and systems for capturing analytes are described. Individual method steps and system features can be present in combination in many different embodiments; the specific combinations described herein do not in any way limit other combinations of steps or features.

(i) Conditions for Capture

[0779] Generally, analytes can be captured when contacting a biological sample with a substrate including capture probes (e.g., substrate with capture probes embedded, spotted, printed on the substrate or a substrate with features (e.g., beads, wells) comprising capture probes).

[0780] As used herein, “contact,” “contacted,” and/or “contacting,” a biological sample with a substrate refers to any contact (e.g., direct or indirect) such that capture probes can interact (e.g., capture) with analytes from the biological sample. For example, a substrate may be near or adjacent to the biological sample without direct physical contact, yet capable of capturing analytes from the biological sample. In some embodiments a biological sample is in direct physical contact with a substrate. In some embodiments, a biological sample is in indirect physical contact with a substrate. For example, a liquid layer may be between the biological sample and the substrate. In some embodiments, analytes diffuse through a liquid layer. In some embodiments capture probes diffuse through a liquid layer. In some embodiments reagents may be delivered via a liquid layer between a biological sample and a substrate. In some embodiments, indirect physical contact may include a second substrate (e.g., a hydrogel, a film, a porous membrane) between the biological sample and the first substrate comprising capture probes. In some embodiments, reagents may be delivered by a second substrate to a biological sample.

[0781] In some embodiments, a cell immobilization agent can be used to contact a biological sample with a substrate (e.g., by immobilizing non-aggregated or disaggregated sample on a spatially-barcoded array prior to analyte capture). A “cell immobilization agent” as used herein can refer to an agent (e.g., an antibody), attached to a substrate, which can bind to a cell surface marker. Non-limiting examples of a cell surface marker include CD45, CD3, CD4, CD8, CD56, CD19, CD20, CD11c, CD14, CD33, CD66b, CD34, CD41, CD61, CD235a, CD146, and epithelial cellular adhesion molecule (EpCAM). A cell immobilization agent can include any probe or component that can bind to (e.g., immobilize) a cell or tissue when on a substrate. A cell immobilization agent attached to the surface of a substrate can be used to bind a cell that has a cell surface marker. The cell surface marker can be a ubiquitous cell surface marker, wherein the purpose of the cell immobilization agent is to capture a high percentage of cells within the sample. The cell surface marker can be a specific, or more rarely expressed, cell surface marker, wherein the purpose of the cell immobilization agent is to capture a specific cell population expressing the target cell surface marker. Accordingly, a cell immobilization agent can be used to selectively capture a cell expressing the target cell surface marker from a population of cells that do not have the same cell surface marker.

[0782] Capture probes on a substrate (or on a feature on the substrate) may interact with released analytes through a capture domain, described elsewhere, to capture analytes. In some embodiments, certain steps are performed to enhance the transfer or capture of analytes to the capture probes of

the array. Examples of such modifications include, but are not limited to, adjusting conditions for contacting the substrate with a biological sample (e.g., time, temperature, orientation, pH levels, pre-treating of biological samples, etc.), using force to transport analytes (e.g., electrophoretic, centrifugal, mechanical, etc.), performing amplification reactions to increase the amount of biological analytes (e.g., PCR amplification, in situ amplification, clonal amplification), and/or using labeled probes for detecting of amplicons and barcodes.

[0783] In some embodiments, an array is adapted in order to facilitate biological analyte migration. Non-limiting examples of adapting an array to facilitate biological analyte migration include arrays with substrates containing nanopores, nanowells, and/or microfluidic channels; arrays with porous membranes; and arrays with substrates that are made of hydrogel. In some cases, the array substrate is liquid permeable. In some cases, the array is a coverslip or slide that includes nanowells or patterning, (e.g., via fabrication). In some cases where the substrate includes nanopores, nanowells, and/or microfluidic channels, these structures can facilitate exposure of the biological sample to reagents (e.g., reagents for permeabilization, biological analyte capture, and/or a nucleic acid extension reaction), thereby increasing analyte capture efficiency as compared to a substrate lacking such characteristics.

[0784] In some embodiments, analyte capture is facilitated by treating a biological sample with permeabilization reagents. If a biological sample is not permeabilized sufficiently, the amount of analyte captured on a substrate can be too low to enable adequate analysis. Conversely, if a biological sample is too permeable, an analyte can diffuse away from its origin in the biological sample, such that the relative spatial relationship of the analytes within the biological sample is lost. Hence, a balance between permeabilizing the biological sample enough to obtain good signal intensity while still maintaining the spatial resolution of the analyte distribution in the biological sample is desired. Methods of preparing biological samples to facilitate analyte capture are known in the art and can be modified depending on the biological sample and how the biological sample is prepared (e.g., fresh frozen, FFPE, etc.).

(ii) Substrate Holder

[0785] Described herein are methods in which an array with capture probes located on a substrate and a biological sample located on a different substrate, are contacted such that the array is in contact with the biological sample (e.g., the substrates are sandwiched together). In some embodiments, the array and the biological sample can be contacted (e.g., sandwiched), without the aid of a substrate holder. In some embodiments, the array and biological sample substrates can be placed in a substrate holder (e.g., an array alignment device) designed to align the biological sample and the array. For example, the substrate holder can have placeholders for two substrates. In some embodiments, an array including capture probes can be positioned on one side of the substrate holder (e.g., in a first substrate placeholder). In some embodiments, a biological sample can be placed on the adjacent side of the substrate holder in a second placeholder. In some embodiments, a hinge can be located between the two substrate placeholders that allows the substrate holder to close, e.g., make a sandwich between the two substrate placeholders. In some embodiments, when the substrate holder is closed the biological sample and the array with capture probes are contacted with one another under conditions sufficient to allow analytes present in the biological sample to interact with the capture probes of the array. For example, dried permeabilization reagents can be placed on the biological sample and rehydrated. A permeabilization solution can be flowed through the substrate holder to permeabilize the biological sample and allow analytes in the biological sample to interact with the capture probes. Additionally, the temperature of the substrates or permeabilization solution can be used to initiate or control the rate of permeabilization. For example, the substrate including the array, the substrate including the biological sample, or both substrates can be held at a low temperature to slow diffusion and permeabilization efficiency. Once sandwiched, in some embodiments, the substrates can be heated to initiate permeabilization and/or increase diffusion efficiency. Transcripts that are released from

the permeabilized tissue can diffuse to the array and be captured by the capture probes. The sandwich can be opened, and cDNA synthesis can be performed on the array.

[0786] Any of the variety of combinations described herein where a sandwich including an array with capture probes and a biological sample on two different substrates can be placed in a substrate holder designed to align the biological sample and the array. For example, the substrate holder can have placeholders for two substrates. In some embodiments, an array including capture probes can be positioned on one side of the substrate holder (e.g., in a first substrate placeholder). In some embodiments, a biological sample can be placed on the adjacent side of the substrate holder in a second placeholder. In some embodiments, in between the two substrate placeholders can be a hinge that allows the substrate holder to close, e.g., make a sandwich between the two substrate placeholders. In some embodiments, when the substrate holder is closed the biological sample and the array with capture probes can be contacted with one another under conditions sufficient to allow analytes present in the biological sample to interact with the capture probes of the array for spatial analysis by any method described herein. For example, dried permeabilization reagents can be placed on the biological sample and rehydrated. Additionally, a permeabilization solution can be flowed through the substrate holder to permeabilize the biological sample and allow analytes in the biological sample to interact with the capture probes.

[0787] In some embodiments, a flexible array described herein can be placed in the substrate holder, and sandwiched with a biological sample. In some embodiments, the flexible array can include spatially-barcoded cross-linked features. In some embodiments, the flexible array can be presoaked in permeabilization reagents before being placed into the substrate holder. In some embodiments, the flexible array can be soaked in permeabilization reagents after being placed in the substrate holder. In some embodiments, the substrate holder including a biological sample in one placeholder and a flexible array can be closed (e.g., form a sandwich) such that the permeabilization reagents allow analytes present in the biological sample to interact with capture probes of the flexible array (e.g., capture probes on the spatially-barcoded features).

[0788] In some embodiments, the substrate holder can be heated or cooled to regulate permeabilization and/or diffusion efficiency.

(iii) Passive Capture Methods

[0789] In some embodiments, analytes can be migrated from a sample to a substrate. Methods for facilitating migration can be passive (e.g., diffusion) and/or active (e.g., electrophoretic migration of nucleic acids). Non-limiting examples of passive migration can include simple diffusion and osmotic pressure created by the rehydration of dehydrated objects.

[0790] Passive migration by diffusion uses concentration gradients. Diffusion is movement of untethered objects toward equilibrium. Therefore, when there is a region of high object concentration and a region of low object concentration, the object (e.g., a capture probe, an analyte, etc.) moves to an area of lower concentration. In some embodiments, untethered analytes move down a concentration gradient.

[0791] In some embodiments, different reagents may be added to the biological sample, such that the biological sample is rehydrated while improving capture of analytes. In some embodiments, the biological sample can be contacted with a shrunken array as described herein. In some embodiments, the biological sample and/or the shrunken array can be rehydrated with permeabilization reagents. In some embodiments, the biological sample and/or the shrunken array can be rehydrated with a staining solution (e.g., hematoxylin and eosin stain).

(iv) Diffusion-Resistant Media/Lids

[0792] To increase efficiency by encouraging analyte diffusion toward the spatially-barcoded capture probes, a diffusion-resistant medium can be used. In general, molecular diffusion of biological analytes occurs in all directions, including toward the capture probes (i.e., toward the spatially-barcoded array), and away from the capture probes (i.e., into the bulk solution). Increasing diffusion toward the spatially-barcoded array reduces analyte diffusion away from the spatially-

barcoded array and increases the capturing efficiency of the capture probes.

[0793] In some embodiments, a biological sample is placed on the top of a spatially-barcoded substrate and a diffusion-resistant medium is placed on top of the biological sample. For example, the diffusion-resistant medium can be placed onto an array that has been placed in contact with a biological sample. In some embodiments, the diffusion-resistant medium and spatially-barcoded array are the same component. For example, the diffusion-resistant medium can contain spatially-barcoded capture probes within or on the diffusion-resistant medium (e.g., coverslip, slide, hydrogel, or membrane). In some embodiments, a sample is placed on a substrate and a diffusion-resistant medium is placed on top of the biological sample. Additionally, a spatially-barcoded capture probe array can be placed in close proximity over a diffusion-resistant medium. For example, a diffusion-resistant medium may be sandwiched between a spatially-barcoded array and a sample on a substrate. In some embodiments, a diffusion-resistant medium is disposed or spotted onto a sample. In other embodiments, a diffusion-resistant medium is placed in close proximity to a sample.

[0794] In general, a diffusion-resistant medium can be any material known to limit diffusivity of biological analytes. For example, a diffusion-resistant medium can be a solid lid (e.g., coverslip or glass slide). In some embodiments, a diffusion-resistant medium may be made of glass, silicon, paper, hydrogel polymer monoliths, or other material. In some embodiments, the glass slide can be an acrylated glass slide. In some embodiments, the diffusion-resistant medium is a porous membrane. In some embodiments, the material may be naturally porous. In some embodiments, the material may have pores or wells etched into solid material. In some embodiments, the pore volume can be manipulated to minimize loss of target analytes. In some embodiments, the membrane chemistry can be manipulated to minimize loss of target analytes. In some embodiments, the diffusion-resistant medium (e.g., hydrogel) is covalently attached to a substrate (e.g., glass slide). In some embodiments, a diffusion-resistant medium can be any material known to limit diffusivity of poly(A) transcripts. In some embodiments, a diffusion-resistant medium can be any material known to limit the diffusivity of proteins. In some embodiments, a diffusion-resistant medium can be any material known to limit the diffusivity of macromolecular constituents.

[0795] In some embodiments, a diffusion-resistant medium includes one or more diffusion-resistant media. For example, one or more diffusion-resistant media can be combined in a variety of ways prior to placing the media in contact with a biological sample including, without limitation, coating, layering, or spotting. As another example, a hydrogel can be placed onto a biological sample followed by placement of a lid (e.g., glass slide) on top of the hydrogel.

In some embodiments, a force (e.g., hydrodynamic pressure, ultrasonic vibration, solute contrasts, microwave radiation, vascular circulation, or other electrical, mechanical, magnetic, centrifugal, and/or thermal forces) is applied to control diffusion and enhance analyte capture. In some embodiments, one or more forces and one or more diffusion-resistant media are used to control diffusion and enhance capture. For example, a centrifugal force and a glass slide can be used contemporaneously. Any of a variety of combinations of a force and a diffusion-resistant medium can be used to control or mitigate diffusion and enhance analyte capture.

[0796] In some embodiments, a diffusion-resistant medium, along with the spatially-barcoded array and sample, is submerged in a bulk solution. In some embodiments, a bulk solution includes permeabilization reagents. In some embodiments, a diffusion-resistant medium includes at least one permeabilization reagent. In some embodiments, a diffusion-resistant medium (i.e. hydrogel) is soaked in permeabilization reagents before contacting the diffusion-resistant medium to the sample. In some embodiments, a diffusion-resistant medium can include wells (e.g., micro-, nano-, or picowells) containing a permeabilization buffer or reagents. In some embodiments, a diffusion-resistant medium can include permeabilization reagents. In some embodiments, a diffusion-resistant medium can contain dried reagents or monomers to deliver permeabilization reagents when the diffusion-resistant medium is applied to a biological sample. In some embodiments, a

diffusion-resistant medium is added to the spatially-barcoded array and sample assembly before the assembly is submerged in a bulk solution. In some embodiments, a diffusion-resistant medium is added to the spatially-barcoded array and sample assembly after the sample has been exposed to permeabilization reagents. In some embodiments, permeabilization reagents are flowed through a microfluidic chamber or channel over the diffusion-resistant medium. In some embodiments, the flow controls the sample's access to the permeabilization reagents. In some embodiments, target analytes diffuse out of the sample and toward a bulk solution and get embedded in a spatially-barcoded capture probe-embedded diffusion-resistant medium. In some embodiments, a free solution is sandwiched between the biological sample and a diffusion-resistant medium.

[0797] FIG. 13 is an illustration of an exemplary use of a diffusion-resistant medium. A diffusion-resistant medium 1302 can be contacted with a sample 1303. In FIG. 13, a glass slide 1304 is populated with spatially-barcoded capture probes 1306, and the sample 1303, 1305 is contacted with the array 1304, 1306. A diffusion-resistant medium 1302 can be applied to the sample 1303, wherein the sample 1303 is sandwiched between a diffusion-resistant medium 1302 and a capture probe coated slide 1304. When a permeabilization solution 1301 is applied to the sample, using the diffusion-resistant medium/lid 1302 directs migration of the analytes 1305 toward the capture probes 1306 by reducing diffusion of the analytes out into the medium. Alternatively, the lid may contain permeabilization reagents.

(v) Active Capture Methods

[0798] In some of the methods described herein, an analyte in a biological sample (e.g., in a cell or tissue section) can be transported (e.g., passively or actively) to a capture probe (e.g., a capture probe affixed to a substrate (e.g., a substrate or bead)).

[0799] For example, analytes can be transported to a capture probe (e.g., an immobilized capture probe) using an electric field (e.g., using electrophoresis), pressure, fluid flow, gravity, temperature, and/or a magnetic field. For example, analytes can be transported through, e.g., a gel (e.g., hydrogel), a fluid, or a permeabilized cell, to a capture probe (e.g., an immobilized capture probe) using a pressure gradient, a chemical concentration gradient, a temperature gradient, and/or a pH gradient. For example, analytes can be transported through a gel (e.g., hydrogel), a fluid, or a permeabilized cell, to a capture probe (e.g., an immobilized capture probe).

[0800] In some examples, an electrophoretic field can be applied to analytes to facilitate migration of analytes towards a capture probe. In some examples, a sample containing analytes contacts a substrate having capture probes fixed on the substrate (e.g., a slide, cover slip, or bead), and an electric current is applied to promote the directional migration of charged analytes towards capture probes on a substrate. An electrophoresis assembly (e.g., an electrophoretic chamber), where a biological sample is in contact with a cathode and capture probes (e.g., capture probes fixed on a substrate), and where the capture probes are in contact with the biological sample and an anode, can be used to apply the current.

[0801] In some embodiments, methods utilizing an active capture method can employ a conductive substrate (e.g., any of the conductive substrates described herein). In some embodiments, a conductive substrate includes paper, a hydrogel film, or a glass slide having a conductive coating. In some embodiments, a conductive substrate (e.g., any of the conductive substrates described herein) includes one or more capture probes.

[0802] FIGS. 24A and 24B show different example analytical workflows of active capture methods using an electric field (e.g., using electrophoresis). In some examples, a biological sample 2402 (e.g., a tissue sample) can be in contact with a first substrate 2404. In some embodiments, first substrate 2404 can have one or more coatings (e.g., any of the conductive substrates described herein) on its surface. Non-limiting examples of coatings include, nucleic acids (e.g., RNA) and conductive oxides (e.g., indium tin oxide). In some embodiments, first substrate 2404 can have a functionalization chemistry on its surface. In the examples shown in FIGS. 24A and 24B, first substrate 2404 is overlaid with a first coating 2406, and first coating 2406 (e.g., a conductive

coating) is further overlaid with a second coating **2408**. In some embodiments, first coating **2406** is an indium tin oxide (ITO) coating. In some embodiments, second coating **2408** is a lawn of capture probes (e.g., any of the capture probes described herein). In some embodiments, a substrate can include an ITO coating. In some embodiments, a substrate can include capture probes or capture probes attached to features on the substrate.

[0803] Biological sample **2402** and second coating **2408** (e.g., a lawn of capture probes) can be in contact with a permeabilization solution **2410**. Non-limiting examples of permeabilization solutions include, enzymes (e.g., proteinase K, pepsin, and collagenase), detergents (e.g., sodium dodecyl sulfate (SDS), polyethylene glycol tert-octylphenyl ether, polysorbate 80, and polysorbate 20), ribonuclease inhibitors, buffers optimized for electrophoresis, buffers optimized for permeabilization, buffers optimized for hybridization, or combinations thereof. Permeabilization reagents can also include but are not limited to a dried permeabilization reagent, a permeabilization buffer, a buffer without a permeabilization reagent, a permeabilization gel, and a permeabilization solution. In some examples, biological samples (e.g., tissue samples) can be permeabilized first and then be subjected to electrophoresis.

[0804] FIG. **24A** shows an example analytical workflow including a first step **2412** in which biological sample **2402** can be permeabilized prior to subjecting the sample **2402** to electrophoresis. Any of the permeabilization methods disclosed herein can be used during first step **2412**. Biological sample **2402** includes an analyte **2414**. In some embodiments, the analyte **2414** is a negatively charged analyte. First substrate **2404** can include a capture probe **2416** that is fixed or attached to the first substrate **2404** or attached to features (e.g., beads) **2418** on the substrate. In some embodiments, capture probe **2416** can include any of the capture probes disclosed herein. In some embodiments, first substrate **2402** does not include features and instead, capture probes **2416** are directly attached to the substrate surface. In some embodiments, the capture probe **2416** is positively charged.

[0805] In step **2420**, after permeabilization of biological sample **2402** concludes, the sample **2402** can be subjected to electrophoresis. During electrophoresis, the biological sample **2402** is subjected to an electric field that can be generated by sandwiching biological sample **2402** between the first substrate **2404** and a second substrate **2422**, connecting each substrate to a cathode and an anode, respectively, and running an electric current through the substrates. The application of the electric field “ $-E$ ” causes the analyte **2404** (e.g., a negatively charged analyte) to migrate towards the substrate **2404** and capture probe **2416** (e.g., a positively charged capture probe) in the direction of the arrows shown in FIG. **24A**. In some embodiments, the analyte **2414** migrates towards the capture probe **2416** for a distance “ h .” In some embodiments, the analyte **2414** migrates towards a capture probe **2416** through one or more permeabilized cells within the permeabilized biological sample (e.g., from an original location in a permeabilized cell to a final location in or close to the capture probe **2416**). Second substrate **2422** can include the first coating **2406** (e.g., a conductive coating), thereby allowing electric field “ $-E$ ” to be generated.

[0806] In some embodiments, the analyte **2414** is a protein or a nucleic acid. In some embodiments, the analyte **2414** is a negatively charged protein or a nucleic acid. In some embodiments, the analyte **2414** is a positively charged protein or a nucleic acid. In some embodiments, the capture probe **2416** is a protein or a nucleic acid. In some embodiments, the capture probe **2416** is a positively charged protein or a nucleic acid. In some embodiments, the capture probe **2416** is a negatively charged protein or a nucleic acid. In some embodiments, the analyte **2414** is a negatively charged transcript. In some embodiments, the analyte **2414** is a poly(A) transcript. In some embodiments, the capture probe **2416** is attached to a feature in a feature array. In some embodiments, permeabilization reagent **2410** can be in contact with sample **2402**, first substrate **2404** second substrate **2422**, or any combination thereof.

[0807] FIG. **24B** shows an example analytical workflow in which biological sample **2402** can be permeabilized and subjected to electrophoresis simultaneously. In some embodiments,

simultaneous permeabilization and electrophoresis of biological sample **2402** can decrease the total duration of the analytical workflow translating into a more efficient workflow.

[0808] In some embodiments, the permeabilization reaction is conducted at a chilled temperature (e.g., about 4° C.). In some embodiments, conducting the permeabilization reaction at a chilled temperature controls the enzyme activity of the permeabilization reaction. In some embodiments, the permeabilization reaction is conducted at a chilled temperature in order to prevent drift and/or diffusion of the analyte **2414** from an original location (e.g., a location in a cell of the biological sample **2402**) until a user is ready to initiate the permeabilization reaction. In some embodiments, the permeabilization reaction is conducted at a warm temperature (e.g., a temperature ranging from about 15° C. to about 37° C. or more) in order to initiate and/or increase the rate of the permeabilization reaction. In some embodiments, once electrophoresis is applied and/or the permeabilization reaction is heated, the permeabilization reaction allows for analyte migration from an original location (e.g., a location in a cell of the biological sample **2402**) to the capture probe **2416** anchored to the first substrate **2404**.

[0809] Referring generally to FIGS. **25A-C**, example substrate configurations for use in the active migration of analytes from a first location to a second location via electrophoresis are shown. FIG. **25A** shows an example substrate configuration for use in electrophoresis in which the first substrate **2502** and the second substrate **2522** are aligned at about 90 degrees with respect to each other. In this example, the first substrate **2502** including biological sample **2504** is placed beneath second substrate **2522**. Both the first substrate **2502** and the second substrate **2522** can be connected to electrical wires **2524** that direct an electric current from a power supply to the substrates, thereby generating an electric field between the substrates. FIG. **25B** shows an additional example substrate configuration for use during electrophoresis in which the first substrate **2502** and the second substrate **2522** are parallel with respect to each other. In this example, the first substrate **2502** including biological sample **2504** is also placed beneath second substrate **2522**.

[0810] FIG. **25C** shows yet an additional example substrate configuration for use in electrophoresis in which the second substrate **2522** and a third substrate **2526** are aligned at about 90 degrees with respect to the first substrate **2502**. Thus, in this example, a first biological sample **2502a** and a second biological sample **2502b** can be subjected to electrophoresis simultaneously. In some embodiments, 3, 4, 5, 6, 7, 8, 9, 10, or more biological samples can be placed on a same substrate and be subjected to electrophoresis simultaneously. In some embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more top substrates can be placed above a same bottom substrate containing one or more samples in order to simultaneously subject the one or more samples to electrophoresis. In some embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more top substrates can be perpendicularly placed (e.g., at about 90 degrees) above a same bottom substrate containing one or more samples in order to simultaneously subject the one or more samples to electrophoresis. In some embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more top substrates can be placed in a parallel orientation above a same bottom substrate containing one or more biological samples in order to simultaneously subject the one or more samples to electrophoresis. In some embodiments, a configuration of top substrates can be arranged above a same bottom substrate containing one or more biological samples in order to simultaneously subject the one or more samples to electrophoresis. In some embodiments, a first configuration of top substrates can be arranged above a second array of bottom substrates containing one or more biological samples in order to simultaneously subject the one or more biological samples to electrophoresis. In some embodiments, simultaneously subjecting two or more biological samples on a same substrate to electrophoresis can provide the advantage of a more effective workflow. In some embodiments, one or more of the top substrates can contain the biological sample.

[0811] In some embodiments, methods utilizing an active capture method can include one or more solutions between the biological sample and the substrate (e.g., a substrate including capture probes). In some embodiments, the one or more solutions between the biological sample and the

substrate including capture probes can include a permeabilization buffer (e.g., any of the permeabilization buffers described herein). In some embodiments, the one or more solutions between the biological sample and the substrate including capture probes can include an electrophoresis buffer.

[0812] In some embodiments, actively capturing analytes can include one or more porous materials between the biological sample and the substrate including capture probes. In some embodiments, the one or more porous materials between the biological sample and substrate including capture probes can include a paper or a blotting membrane. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can include a gel containing one or more solutions. For example, in a non-limiting way, the gel can be a SDS-PAGE gel. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can contain a permeabilization buffer. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can contain an electrophoresis buffer. In some embodiments, actively capturing analytes can include one or more solutions and one or more porous materials between the biological sample and the substrate including capture probes.

[0813] In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes (e.g., an array) can act as a filter to separate analytes (e.g., analytes of interest) from other molecules or analytes present in the biological sample. In some embodiments, the analytes (e.g., analytes of interest) are RNA transcripts. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can act as a filter to separate RNA transcripts from other molecules (e.g., analytes) such as proteins, lipids and/or other nucleic acids. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can act as a filter to separate the analytes and other molecules based on physicochemical properties. For example, in a non-limiting way, analytes can be separated on properties such as charge, size (e.g., length, radius of gyration, effective diameters, etc.), hydrophobicity, hydrophilicity, molecular binding (e.g., immunoaffinity), and combinations thereof. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can separate the analytes from other molecules to reduce non-specific binding near the capture probes and therefore improve binding between the analytes and the capture probes, thus improving subsequent assay performance.

[0814] In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can act as molecular sieving matrices for electrophoretic analyte separation. For example, in a non-limiting way, separation of analytes can occur based on physicochemical properties such as charge, size (e.g., length, radius of gyration, and effective diameters, etc.), electrophoretic mobility, zeta potential, isoelectric point, hydrophobicity, hydrophilicity, molecular binding (e.g., immunoaffinity), and combinations thereof. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can be of a uniform pore size. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can have discontinuities in pore sizes, as generally used in different gel electrophoresis schemes. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can have gradients in pore sizes. For example, the one or more porous materials (e.g., a hydrogel) can have a gradient of pore sizes such that the gradient separates the analytes as the analytes migrate to the substrate including capture probes (e.g., an array).

[0815] In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can separate the analytes based on length. For example, shorter analytes will have a higher electrophoretic mobility, and therefore migrate faster towards the capture probes relative to longer analytes in an electrophoretic setup. In some embodiments, the

one or more porous materials between the biological sample and the substrate including capture probes separate the analytes based on length, such that only shorter analytes can migrate through the one or more porous materials to reach the capture probes, while longer analytes cannot reach the capture probes.

[0816] In some embodiments, specific subsets of analytes (e.g., a subset of transcripts) can be captured by applying an electrophoretic field for a certain amount of time. In some embodiments, specific subsets of analytes (e.g., a subset of transcripts) can be captured by selecting different porous materials (e.g., porous materials with different compositions) between the biological sample and the substrate including capture probes. In some embodiments, specific subsets of analytes (e.g., a subset of transcripts) can be captured by applying an electrophoretic field for a certain amount of time and selecting different porous materials between the biological sample and the substrate including capture probes (e.g., an array).

[0817] In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can have discontinuities in pore sizes that can cause an increase in the concentration of the migrating analytes (e.g., “stacking”). For example, the one or more porous materials (e.g., a hydrogel) between the biological sample and the substrate including capture probes can have discontinuities in pore sizes that can cause an increase in the concentration of the analytes near the capture probes resulting in favorable binding kinetics and increased sensitivity. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can have discontinuities in pore sizes that enhance the separation between migrating analytes of different sizes and/or lengths. In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can include a first porous material and a second porous material, with the first porous material having a larger pore size than the second porous material. In some embodiments, the first porous material is located on the surface, or near the surface, of the biological sample. In some embodiments, the second porous material (e.g., second porous material with a smaller pore size than the first porous material) can be placed on the surface, or near the surface, of the first porous material. In some embodiments, as analytes migrate (e.g., migrate via electrophoresis) from the biological sample through the first porous material and the second porous material sequentially, the migrating analytes can collect (e.g., “stack”) at the interface between the first porous material and the second porous material.

[0818] In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can include a gradient in pore sizes for continuous stacking as analytes migrate through decreasing pore sizes (e.g., decreasing pore diameter). In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can include a gradient in pore sizes such that the pores decrease in diameter as the analytes migrate from the biological sample to the substrate including capture probes. In some embodiments, the pore size gradient can increase the resolution among analytes of different sizes. In some embodiments, the pore size gradient can increase the concentration of the analytes near the capture probes. In some embodiments, the pore size gradient can continuously reduce the speed at which the analytes migrate and collect (e.g., “stack”) as the analytes migrate through the gradient of decreasing pore sizes (e.g., decreasing pore diameter).

[0819] In some embodiments, the one or more porous materials between the biological sample and the substrate including capture probes can include a gradient gel for continuous stacking as analytes migrate through decreasing pore sizes (e.g., decreasing pore diameter) of the gradient gel. In some embodiments, the gradient gel can have pores with a decreasing diameter as the analytes migrate toward the capture probes. In some embodiments, the gradient gel can increase the separation resolution among analytes of different sizes. In some embodiments, the gradient gel can increase the concentration of analytes near the capture probes. In some embodiments, the gradient gel can continuously reduce the speed at which analytes migrate and collect (e.g., “stack”) as the analytes

migrate through the gradient gel of decreasing pore sizes (e.g., decreasing in diameter).

[0820] In some embodiments, a biological sample can be placed in a first substrate holder (e.g., a substrate holder described herein). In some embodiments, a spatially-barcoded capture probe array (e.g., capture probes, barcoded array) can be placed on a second substrate holder (e.g., a substrate holder described herein). In some embodiments, a biological sample can be placed in a first substrate holder that also contains capture probes. In some embodiments, the first substrate holder, the second substrate holder, or both can be conductive (e.g. any of the conductive substrates described herein). In some embodiments, the first substrate holder including the biological sample, the second substrate holder including capture probes, or both, can be contacted with permeabilization reagents (e.g., a permeabilization buffer) and analytes can be migrated from the biological sample toward the barcoded array using an electric field.

[0821] In some embodiments, electrophoresis can be applied to a biological sample on a barcoded array while in contact with a permeabilization buffer. In some embodiments, electrophoresis can be applied to a biological sample on a barcoded array while in contact with an electrophoresis buffer (e.g. a buffer that lacks permeabilization reagents). In some embodiments, the permeabilization buffer can be replaced with an electrophoresis buffer after a desired amount of time. In some embodiments, electrophoresis can be applied simultaneously with the permeabilization buffer or electrophoresis buffer. In some embodiments, electrophoresis can be applied after a desired amount of time of contact between the biological sample and the permeabilization buffer or electrophoresis buffer.

[0822] In some embodiments, the biological sample can be placed on a substrate (e.g., a porous membrane, a hydrogel, paper, etc.). In some embodiments, the biological sample placed on the substrate can have a gap (e.g., a space) between the substrate and the substrate holder (e.g., conductive substrate holder). In some embodiments, the barcoded array can be placed on a substrate (e.g., a porous membrane, a hydrogel, paper, etc.). In some embodiments, the barcoded array can have a gap between the substrate and substrate holder (e.g., conductive substrate holder). In some embodiments, the barcoded array can be placed in direct proximity to the biological sample or at a desired distance from the biological sample. In some embodiments, a buffer reservoir can be used between the substrate holder (e.g., conductive substrate holder) and the barcoded array, the biological sample, or both. This setup allows the analytes to be migrated to a barcoded array while not in proximity with the electrodes (e.g. conductive substrate holder), thus resulting in more stable electrophoresis.

[0823] In some embodiments, a combination of at least two buffers with different ionic compositions can be used to differentially migrate analytes based on their ionic mobility (e.g., isotachophoresis (ITP)). For example, using two or more buffers with different ionic compositions can increase the concentration of analytes prior to contact with a barcoded array. Isotachophoresis includes at least two buffers that contain a common counter-ion (e.g., ions that have different charge sign than the analytes) and different co-ions (e.g., ions that have the same charge sign as the analytes) (Smejkal P., et al., Microfluidic isotachophoresis: A review, *Electrophoresis*, 34.11 1493-1509, (2013) which is incorporated herein by reference in its entirety). In some embodiments, one buffer can contain a co-ion with a higher ionic mobility (e.g. speed at which they travel through solution in an electric field) than the analytes (e.g., the “leading” buffer). In some embodiments, a second buffer can contain a co-ion with a lower ionic mobility than the analytes (e.g., the “trailing” buffer). In some embodiments, a third buffer can contain a co-ion with an ionic mobility that is between the electric mobility of the analytes. In some embodiments, a biological sample can be placed on a first substrate holder (e.g., a conductive substrate holder) and the barcoded array can be placed on a second substrate holder (e.g., a second conductive substrate holder) and contacted with a permeabilization buffer and the analytes can be migrated away from the biological sample and toward the barcoded array using an electric field. As the electric field is applied to the biological sample the analytes can be concentrated in the buffer as they are migrated toward the capture

probes. In some embodiments, isotachopheresis can be used with gel-based separations (e.g., any of the gel-based separations described herein).

[0824] In some embodiments, a permeabilization buffer can be applied to a region of interest (e.g., region of interest as described herein) in a biological sample. In some embodiments, permeabilization reagents (e.g. a hydrogel containing permeabilization reagents) can be applied to a region of interest in a biological sample. For example, a region of interest can be a region that is smaller in area relative to the overall area of the biological sample. In some embodiments, the permeabilization buffer or permeabilization reagents can be contacted with the biological sample and a substrate including capture probes (e.g., an array). In some embodiments, the biological sample can have more than one region of interest (e.g. two, three). In some embodiments, the biological sample, the substrate including capture probes, or both, can be placed in a conductive substrate holder. In some embodiments, analytes can be released from the region(s) of interest and migrated from the biological sample toward the capture probes with an electric field.

[0825] In some embodiments, electrophoretic transfer of analytes can be performed while retaining the relative spatial locations of analytes in a biological sample while minimizing passive diffusion of an analyte away from its location in a biological sample. In some embodiments, an analyte captured by a capture probe (e.g., capture probes on a substrate) retains the spatial location of the analyte present in the biological sample from which it was obtained (e.g., the spatial location of the analyte that is captured by a capture probe on a substrate when the analyte is actively migrated to the capture probe by electrophoretic transfer can be more precise or representative of the spatial location of the analyte in the biological sample than when the analyte is not actively migrated to the capture probe). In some embodiments, electrophoretic transport and binding process is described by the Damköhler number (Da), which is a ratio of reaction and mass transport rates. The fraction of analytes bound and the shape of the biological sample will depend on the parameters in the Da . There parameters include electromigration velocity $U_{sub.e}$ (depending on analyte electrophoretic mobility $\mu_{sub.e}$ and electric field strength E), density of capture probes (e.g., barcoded oligonucleotides) $p_{sub.0}$, the binding rate between probes (e.g., barcoded oligonucleotides) and analytes $k_{sub.on}$, and capture area thickness L .

$$[00001] Da \sim \frac{k_{on} p_0 L}{E}$$

Fast migration (e.g., electromigration) can reduce assay time and can minimize molecular diffusion of analytes.

[0826] In some embodiments, electrophoretic transfer of analytes can be performed while retaining the relative spatial alignment of the analytes in the sample. As such, an analyte captured by the capture probes (e.g., capture probes on a substrate) retains the spatial information of the cell or the biological sample from which it was obtained. Applying an electrophoretic field to analytes can also result in an increase in temperature (e.g., heat). In some embodiments, the increased temperature (e.g., heat) can facilitate the migration of the analytes towards a capture probe.

[0827] In some examples, a spatially-addressable microelectrode array is used for spatially-constrained capture of at least one charged analyte of interest by a capture probe. For example, a spatially-addressable microelectrode array can allow for discrete (e.g., localized) application of an electric field rather than a uniform electric field. The spatially-addressable microelectrode array can be independently addressable. In some embodiments, the electric field can be applied to one or more regions of interest in a biological sample. The electrodes may be adjacent to each other or distant from each other. The microelectrode array can be configured to include a high density of discrete sites having a small area for applying an electric field to promote the migration of charged analyte(s) of interest. For example, electrophoretic capture can be performed on a region of interest using a spatially-addressable microelectrode array.

[0828] A high density of discrete sites on a microelectrode array can be used. The surface can include any suitable density of discrete sites (e.g., a density suitable for processing the sample on

the conductive substrate in a given amount of time). In one embodiment, the surface has a density of discrete sites greater than or equal to about 500 sites per 1 mm.^{sup.2}. In some embodiments, the surface has a density of discrete sites of about 100, about 200, about 300, about 400, about 500, about 600, about 700, about 800, about 900, about 1,000, about 2,000, about 3,000, about 4,000, about 5,000, about 6,000, about 7,000, about 8,000, about 9,000, about 10,000, about 20,000, about 40,000, about 60,000, about 80,000, about 100,000, or about 500,000 sites per 1 mm.^{sup.2}. In some embodiments, the surface has a density of discrete sites of at least about 200, at least about 300, at least about 400, at least about 500, at least about 600, at least about 700, at least about 800, at least about 900, at least about 1,000, at least about 2,000, at least about 3,000, at least about 4,000, at least about 5,000, at least about 6,000, at least about 7,000, at least about 8,000, at least about 9,000, at least about 10,000, at least about 20,000, at least about 40,000, at least about 60,000, at least about 80,000, at least about 100,000, or at least about 500,000 sites per 1 mm.^{sup.2}.

[0829] Schematics illustrating an electrophoretic transfer system configured to direct nucleic acid analytes (e.g., mRNA transcripts) toward a spatially-barcoded capture probe array are shown in FIG. 14A and FIG. 14B. In this exemplary configuration of an electrophoretic system, a sample 1402 is sandwiched between the cathode 1401 and the spatially-barcoded capture probe array 1404, 1405, and the spatially-barcoded capture probe array 1404, 1405 is sandwiched between the sample 1402 and the anode 1403, such that the sample 1402, 1406 is in contact with the spatially-barcoded capture probes 1407. When an electric field is applied to the electrophoretic transfer system, negatively charged nucleic acid analytes 1406 will be pulled toward the positively charged anode 1403 and into the spatially-barcoded array 1404, 1405 containing the spatially-barcoded capture probes 1407. The spatially-barcoded capture probes 1407 interact with the nucleic acid analytes (e.g., mRNA transcripts hybridize to spatially-barcoded nucleic acid capture probes forming DNA/RNA hybrids) 1406, making the analyte capture more efficient. The electrophoretic system set-up may change depending on the target analyte. For example, proteins may be positive, negative, neutral, or polar depending on the protein as well as other factors (e.g., isoelectric point, solubility, etc.). The skilled practitioner has the knowledge and experience to arrange the electrophoretic transfer system to facilitate capture of a particular target analyte.

[0830] FIG. 15 is an illustration showing an exemplary workflow protocol utilizing an electrophoretic transfer system. In the example, Panel A depicts a flexible spatially-barcoded feature array being contacted with a sample. The sample can be a flexible array, wherein the array is immobilized on a hydrogel, membrane, or other flexible substrate. Panel B depicts contact of the array with the sample and imaging of the array-sample assembly. The image of the sample/array assembly can be used to verify sample placement, choose a region of interest, or any other reason for imaging a sample on an array as described herein. Panel C depicts application of an electric field using an electrophoretic transfer system to aid in efficient capture of a target analyte. Here, negatively charged mRNA target analytes migrate toward the positively charged anode. Panel D depicts application of reverse transcription reagents and first strand cDNA synthesis of the captured target analytes. Panel E depicts array removal and preparation for library construction (Panel F) and next-generation sequencing (Panel G).

(vi) Targeted Analysis

[0831] In some aspects, arrays (e.g., glass slides) include a plurality of capture probes that bind to one or more specific biological targets in a sample. The capture probes can be directly or indirectly attached to a substrate. The capture probe can be or include, for example, DNA or RNA. In some aspects, the capture probes on an array can be immobilized, e.g., attached or bound, to the array via their 5' or 3' ends, depending on the chemical matrix of the array. In some aspects, the probes are attached via a 3' linkage, thereby leaving a free 5' end. In some aspects, the probes are attached via a 5' linkage, thereby leaving a free 3' end. In some aspects, the probes are immobilized indirectly. For example, a probe can be attached to a bead, which bead can be deposited on a substrate. A capture probe as disclosed in this section can include any of the various components of a capture

probe as provided throughout this disclosure (e.g., spatial barcodes, UMIs, functional domains, cleavage domains, etc.).

[0832] In some aspects, a capture probe or plurality of capture probes interact with an analyte specific for a particular species or organism (e.g., host or pathogen). In some aspects, the probe or plurality of probes can be used to detect a viral, bacterial, or plant protein or nucleic acid. In some aspects, the capture probe or plurality of capture probes can be used to detect the presence of a pathogen (e.g., bacteria or virus) in the biological sample. In some aspects, the capture probe or plurality of capture probes can be used to detect the expression of a particular nucleic acid associated with a pathogen (e.g., presence of 16S ribosomal RNA or Human Immunodeficiency Virus (HIV) RNA in a human sample).

[0833] In some aspects, the capture domain in the capture probe can interact with one or more specific analytes (e.g., an analyte or a subset of analytes out of the pool of total analytes). The specific analyte(s) to be detected can be any of a variety of biological molecules including but not limited to proteins, nucleic acids, lipids, carbohydrates, ions, small molecules, subcellular targets, or multicomponent complexes containing any of the above. In some embodiments, the analyte(s) can be localized to subcellular location(s), including, for example, organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplasts, endocytic vesicles, exocytic vesicles, vacuoles, lysosomes, etc. In some embodiments, analyte(s) can be peptides or proteins, including without limitation antibodies and enzymes.

[0834] In some aspects, analytes from a biological sample interact with one or more capture probes (e.g., one or more capture probes immobilized directly or indirectly on a substrate), and the capture probes interact with specific analytes in the biological sample. In some aspects, the capture probes are allowed to interact with (e.g., hybridize to) specific analytes, e.g., under appropriate conditions where oligonucleotide capture probes can hybridize to the target nucleic acids. In some aspects, analytes that did not hybridize to capture probes are removed (e.g., analytes that do not interact with capture domains of the capture probes). In some embodiments, removal of analytes that did not interact with a capture probe can be accomplished by, e.g., washing the sample to remove such analytes.

[0835] In some aspects, a capture probe or plurality of capture probes includes a capture domain that interacts with an analyte or analytes present in a biological sample. In some aspects, the capture probe or plurality of capture probes includes a capture domain that detects the presence or level amount (e.g., expression level) of a particular analyte or analytes of interest. The capture domain of a capture probe (immobilized directly or indirectly on a substrate) can be capable of binding selectively to a desired subtype or subset of nucleic acid. In some aspects, for example, the capture domain binds to a subset of nucleic acids in a genome or a subset of nucleic acids in a transcriptome. In some aspects, the analyte(s) can include one or more nucleic acids. In some aspects, the capture probe or plurality of capture probes can be used to detect the expression of a particular transcript (e.g., a particular mRNA). In some aspects, a capture probe or plurality of capture probes can be specific for (e.g., binds to) an individual change in a nucleic acid or protein (e.g., a mutation or single nucleotide polymorphism (SNP)).

[0836] In some aspects, the biological sample includes an analyte that is or includes a nucleic acid. The nucleic acid can be RNA or DNA. In some aspects, the capture probe or plurality of capture probes detects DNA copy number of a particular set of nucleic acid analyte or analytes. For example, capture probe or plurality of capture probes provided herein can be used to detect DNA copy number of nucleic acids that share homology to each other.

[0837] In some aspects, the capture probe or plurality of capture probes includes a capture domain that detects the presence or level amount (e.g., expression level) of one or more RNA transcripts (e.g., specific RNA transcripts). In some aspects, the capture probe or plurality of capture probes includes a capture domain that detects the presence or amount (e.g., expression level) of one or more non-coding RNAs (e.g., microRNA, transfer RNA (tRNA), ribosomal RNA (rRNA), small

interfering RNA (siRNA) and small nucleolar RNA (snoRNA). In some aspects, the probe or plurality of probes includes a capture domain that detects the presence or level amount (e.g., expression level) of one or more proteins (e.g., proteins expressed of a nucleic acid of interest). [0838] In some aspects, the capture probe or plurality of capture probes can be specific for a particular protein. In some aspects, the capture probe or plurality of capture probes can be used to detect the presence of a particular protein of interest. In some aspects, the capture probe or plurality of capture probes can be used to detect translation of a particular protein. In some aspects, the capture probe or plurality of capture probes can specifically interact with an active region of an enzyme, a binding domain of an immunoglobulin, defined domains of proteins, whole proteins, synthetic peptides, peptides with introduced mutations, aptamer, or any combination thereof. In some aspects, the analyte(s) can include one or more proteins. In some aspects, the analyte(s) can include one or more nucleic acids and one or more proteins.

[0839] In some aspects, the capture probe or plurality of capture probes can be used to detect particular post-translational modifications of a particular protein. In such embodiments, analyte capture agents can be specific for cell surface analytes having a given state of posttranslational modification (e.g., phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation), such that a cell surface analyte profile can include posttranslational modification information of one or more analytes.

[0840] In some aspects, the capture probe or plurality of capture probes can be specific for a particular set of nucleic acids (e.g., nucleic acids that are associated with a specific cellular pathway or pathways). In some aspects, the set of nucleic acids is DNA. In some aspects, the set of nucleic acids is RNA. In some aspects, the set of nucleic acids has similar and/or homologous sequences. In some aspects, the set of nucleic acids encodes for analytes that function in a similar cellular pathway. In some aspects, the set of nucleic acids encodes for analytes that are expressed in a certain pathological state (e.g., cancer, Alzheimer's, or Parkinson's disease). In some aspects, the set of nucleic acids encodes for analytes that are over-expressed in a certain pathological state. In some aspects, the set of nucleic acids encodes for analytes that are under-expressed in a certain pathological state.

[0841] In some aspects, the capture probe or plurality of capture probes can be specific for a particular nucleic acid, or detection or expression of a particular set of proteins (e.g., in a similar cellular pathway). In some aspects, the set of proteins has similar functional domains. In some aspects, the set of proteins functions in a similar cellular pathway. In some aspects, the set of proteins is expressed in a certain pathological state (e.g., cancer, Alzheimer's or Parkinson's disease). In some aspects, the set of proteins is over-expressed in a certain pathological state. In some aspects, the set of proteins is under-expressed in a certain pathological state.

[0842] In some embodiments, a capture probe includes a capture domain that is capable of binding to more than one analyte. In some embodiments, a capture domain can bind to one or more analytes that are about 80% identical, about 85% identical, about 90% identical, about 95% identical, about 96% identical, about 97% identical, about 98% identical, about 99% identical, 100% identical to the target analyte. In some aspects, the capture probe can bind to an analyte that is about 80% identical, about 85% identical, about 90% identical, about 95% identical, about 96% identical, about 97% identical, about 98% identical, or about 99% identical to each other. In some embodiments, a capture domain can bind to a conserved region of one or more analytes, in which the conserved regions are about 80% identical, about 85% identical, about 90% identical, about 95% identical, about 96% identical, about 97% identical, about 98% identical, about 99% identical, 100% identical to the target analyte.

[0843] In some aspects, a capture probe or plurality of capture probes interacts with two or more analytes (e.g., nucleic acids or proteins) that are not similar in sequence and/or do not share a conserved domain. In some embodiments, a capture probe includes two or more capture domains, each of which interacts with a different analyte. In such embodiments, members of the two or more

capture domains can be adjacent to each other in the capture probe and/or members of the two or more capture domains can be separated from each other in the capture probe by one or more domains (e.g., nucleic acid domains). For example, in some aspects, the sets of analytes that are detected include mutational changes in the targeted nucleic acids or proteins. In some aspects, the capture probe or plurality of capture probes detects sets of nucleic acids or proteins (e.g., non-homologous nucleic acids or proteins) that are individually mutated during a pathogenic state. In some aspects, the pathogenic state is cancer.

[0844] In some aspects, a capture probe or plurality of capture probes include capture domains that can be used to detect analytes that are typically detected using diagnostic panels. In some aspects, the capture probe or plurality of capture probes are used to detect changes in one or more analytes. In some aspects, the analyte changes include one or more of increased analyte expression, decreased analyte expression, mutated nucleic acid sequences, or any combination thereof. In some aspects, the changes in the analytes are associated with and/or lead to manifestation of a pathogenic state in a subject. In some aspects, the detected changes are compared to a reference analyte or analytes.

(vii) Polypeptide Capture

[0845] Provided herein are methods and materials for identifying the location of a polypeptide in a biological sample. In some embodiments, an analyte (e.g., a polypeptide analyte) can be directly captured on a substrate. For example, polypeptide analytes can be captured by amine groups on a functionalized substrate. In other examples, an analyte (e.g., a polypeptide analyte) can be captured via an analyte binding moiety directly attached to a substrate. In some embodiments, the substrate may be populated with analyte binding moieties directly attached to the substrate as well as spatially-barcoded capture probes directly attached to the substrate. In other embodiments, an analyte (e.g., a polypeptide analyte) can be captured via an analyte binding moiety indirectly attached to a substrate. In an example, the substrate may be populated with capture probes that are bound to an analyte capture agent, wherein the analyte capture domain of the analyte capture agent binds to the capture domain of the capture probe and the analyte binding moiety binds the polypeptide analyte.

[0846] In some embodiments, an analyte (e.g., a polypeptide analyte) can be directly captured or immobilized on a substrate. Direct immobilization may be achieved by covalently coupling the polypeptide analyte to the substrate via amide bonds between the carboxylic acid of the C-terminal amino acid residue and a functionalized substrate surface. For example, a substrate (e.g., a glass coverslip or slide) can be functionalized through amino-silanization with aminopropyltriethoxysilane. The substrate surfaces are further passivated by overnight incubation with polyethylene glycol (PEG)-NHS solution, and functionalized slides can be stored in a vacuum desiccator until use. The t-butyloxycarbonyl protecting groups can be removed by incubating the substrate with 90% TFA (v/v in water) for 5 hours before use, thus exposing free amine groups for peptide immobilization. The resulting functionalized substrate is stable to multiple cycles of Edman degradation and washing steps.

[0847] In some embodiments, methods for capturing polypeptides in a biological sample include providing a substrate where an analyte binding moiety is directly immobilized on the substrate. In some embodiments, direct immobilization is achieved through chemical modification of the substrate and/or chemical modification of the analyte binding moiety. For example, a substrate can be prepared with free amines on the surface. When exposed to an analyte binding moiety with a free carboxylic acid on the C-terminal residue, the free amines can form amide bonds with the carboxylic acid thereby covalently coupling the analyte binding moiety to the substrate. Substrates and/or analyte binding moieties can be modified in any manner that facilitates covalent bonding of the analyte binding moiety to the substrate. Non-limiting examples of chemical modification that can be used to covalently bind the analyte binding moiety to the substrate include are described herein.

[0848] In some embodiments, methods for capturing analyte polypeptides include providing a substrate (e.g., an array) where the analyte binding moiety is indirectly attached to the substrate. For example, an analyte binding moiety can be indirectly attached to a substrate via an oligonucleotide (e.g., a capture agent barcode domain or capture agent barcode domain hybridized to a capture probe) or other domain capable of binding to both the substrate and the analyte binding domain. The capture agent barcode domain is described elsewhere herein. The capture agent barcode domain can be modified to include a cleavage domain, which can attach to a substrate using any of the chemistries described herein. In some embodiments, the capture agent barcode domain can include an analyte capture sequence as described herein, wherein the analyte capture sequence can hybridize to the capture domain of a capture probe. In some embodiments, a substrate (e.g., an array) containing capture probes can be modified to capture polypeptide analytes by hybridizing the analyte capture sequence of the analyte capture agent to the capture domain of a capture probe.

[0849] In some embodiments, methods for capturing analyte polypeptides include providing a substrate (e.g., an array) and providing an analyte capture agent to the biological sample. For example, after drying and fixing sectioned tissue samples, the tissue samples can be positioned on a substrate (e.g., a spatial array), rehydrated, blocked, and permeabilized (e.g., 3×SSC, 2% BSA, 0.1% Triton X, 1 U/μl RNase inhibitor for 10 min at 4° C.) before being stained with fluorescent primary antibodies (1:100) and a pool of analyte capture agents (in 3×SSC, 2% BSA, 0.1% Triton X, 1 U/μl RNase inhibitor for 30 min at 4° C.). The biological sample can be washed, coverslipped (in glycerol+1 U/μl RNase inhibitor), imaged for detected analytes (e.g., using a confocal microscope or other apparatus capable of fluorescent detection), and washed again. The analyte-bound analyte capture agents can be released from the biological sample (e.g., the biological sample can be treated with proteinase, e.g., proteinase K) and migrated to the spatial array. An analyte capture sequence of the analyte-bound analyte capture agent can be captured by a capture probe capture domain, and the capture agent barcode domain can be extended to produce a spatially-tagged analyte capture agent. The spatially-tagged analyte capture agents can be processed according to spatial workflows described herein.

[0850] In some embodiments, methods for capturing analyte polypeptides include providing blocking probes to analyte capture agents before introducing the analyte capture agents to a biological sample. In some embodiments, the blocking probes can be alternatively or additionally provided in any of the rehydrating or blocking buffers provided herein. In some embodiments, the analyte capture agent analyte capture sequence can be blocked prior to binding to the capture probe capture domain using a blocking probe sequence complementary to the analyte capture sequence. Blocking the capture agent barcode domain, particularly the free 3' end of the capture agent barcode domain (e.g., analyte capture sequence), prior to contacting the analyte capture agents with the biological sample and/or substrate, can prevent binding of the analyte capture sequence of the capture agent barcode domains, e.g., prevents the binding of a poly(A) tail to the capture probe capture domain. In some embodiments, blocking the analyte capture agent analyte capture domain reduces non-specific background staining. In some embodiments, the blocking probes are reversible, such that the blocking probes can be removed from the analyte capture sequence during or after the time that analyte capture agents are in contact with the biological sample. In some embodiments, the blocking probe can be removed with RNase treatment (e.g., RNase H treatment).

[0851] In some embodiments, methods for capturing polypeptides in a biological sample include active transfer (e.g., electrophoresis). For example, the biological sample is placed on a conductive substrate and contacted with a spatial array including one or more analyte binding moieties. An electric field can be applied to the conductive substrate to promote migration of the polypeptides towards the analyte binding moieties, as described herein.

[0852] In some embodiments, methods for identifying the spatial location of a polypeptide in a

biological sample include determining the sequence of a captured polypeptide. In some embodiments, the sequence of the captured polypeptide is determined through detection of amino acid residues labeled with a detectable label (e.g., radiolabel of a fluorophore). Non-limiting examples of detectable labels that can be used for labelling the captured polypeptide include fluorophores and radiolabels. In some embodiments, the polypeptides are labeled at specific amino acid residues only (e.g., not all amino acid residues are labeled). In some embodiments, the polypeptide is labeled prior to contacting the biological sample with the substrate. In some embodiments, a captured polypeptide is labeled with fluorophores using standard coupling schemes (see Hernandez et al., *New J. Chem.* 41:462-469 (2017)). For example, polypeptides may be labeled by reaction with Atto647N-NHS, Atto647Niodoacetamide, TMR-NHS, or JF549-NHS, as appropriate, to label lysines (via NHS) or cysteines (via iodoacetamide). In addition, serine or threonine phosphorylation sites may be selectively labeled via beta elimination followed by conjugate addition via thiols to substitute thiol-linked fluorophores in place of phosphates (see Stevens et al., *Rapid Commun. Mass Spectrom.*, 15:2157-2162 (2005)). The number of fluorophores incorporated into a polypeptide is any number that may be spectrally resolved. In some instances, four or more fluorophores are utilized.

[0853] In some embodiments, a captured polypeptide is radiolabeled. In some embodiments, specific amino acids can be labeled with an isotope. Non-limiting examples of isotopes used to label amino acids include .sup.3H, .sup.14C, .sup.15N, .sup.32P, and .sup.125I. In some embodiments, the isotope is incorporated into the selected amino acid prior to incorporation into a polypeptide. In some embodiments, the radiolabeled amino acid can be incorporated into the polypeptide after polypeptide formation.

[0854] In some embodiments, the sequence of the captured polypeptide is determined using Edman degradation (and in some embodiments successive rounds of Edman degradation). In such cases, a polypeptide is captured, and the polypeptide sequence can be resolved by imaging the substrate following repeated rounds of Edman degradation. For example, the substrate is imaged following each Edman reaction in order to capture the detectable labels that are produced due to the removal of amino acids that are a byproduct of the reaction. The information obtained by the Edman degradation can be compiled to identify a polypeptide. In some embodiments, the biological sample is visualized or imaged using light or fluorescence microscopy.

(viii) Enrichment of Captured Analytes after Capture

[0855] In some aspects, spatial analysis of targeted analytes includes an enrichment step or steps post-capture to enrich the captured analytes for the targeted analyte. For example, the capture domain can be selected or designed for the selective capture of more analytes than the practitioner desires to analyze. In some embodiments, capture probes that include random sequences (e.g., random hexamers or similar sequences) that form all or part of the capture domain can be used to capture nucleic acids from a biological sample in an unbiased way. For example, capture probes having capture domains that include random sequences can be used to generically capture DNA, RNA, or both from a biological sample. Alternatively, capture probes can include capture domains that capture mRNA generally. As is well known in the art, this can be on the basis of hybridization to the poly-A tail of mRNAs. In some embodiments, the capture domain includes a sequence that interacts with (e.g., hybridizes to) the poly-A tail of mRNAs. Non-limiting examples of such sequences include poly-T DNA sequences and poly-U RNA sequences. In some embodiments, random sequences can be used in conjunction with poly-T (or poly-T analogue etc.) sequences. Thus, where a capture domain includes a poly-T (or a "poly-T-like") oligonucleotide, it can also include a random oligonucleotide sequence.

[0856] In some aspects, after capture of more analytes than the practitioner desires to analyze, methods disclosed herein include enrichment of particular captured analytes. In some aspects, methods include enrichment of analytes that include mutations (e.g., SNPs,) of interest, nucleic acid(s) of interest, and/or proteins(s) of interest.

[0857] In some embodiments, methods of spatial analysis provided herein include selectively enriching one or more analytes of interest (e.g., target analytes) after analyte capture. For example, one or more analytes of interest can be enriched by addition of one or more oligonucleotides to the pool of captured analytes. In some embodiments, one or more analytes of interest can be enriched by addition of one or more oligonucleotides to the pool of captured analytes on the array. In some embodiments, one or more analytes of interest can be enriched by addition of one or more oligonucleotides to the pool of captured analytes where the pool of captured analytes have been released (e.g., removed) from the array. In some embodiments, when captured analytes have been released from the array the one or more nucleotides can be complementary to a portion of a TSO and R1 sequence, or portion thereof. In some embodiments, the additional oligonucleotide(s) include a sequence used for priming a reaction by a polymerase. For example, one or more primer sequences with sequence complementarity to one or more analytes of interest can be used to amplify the one or more analyte(s) of interest, thereby selectively enriching these analytes. In some embodiments, one or more primer sequences can be complementary to other domains on the capture probe (e.g., R1 sequence, or portion thereof, as above), and not complementary to the analyte. In some embodiments, enrichment by amplification (e.g., PCR) occurs by using a first primer complementary to an analyte or analytes of interest (or another domain in the capture probe and the TSO), or complement thereof, and a second primer complementary to a region of the capture probe, or complement thereof. In some embodiments, the region of the capture probe, or complement thereof, is distal to a spatial barcode from the capture domain, such that enrichment by amplification amplifies both the captured analyte or analytes and its or their associated spatial barcodes, thus permitting spatial analysis of the enriched analyte or analytes.

[0858] In some embodiments, two or more capture probes capture two or more distinct analytes, which analytes are enriched (e.g., simultaneously or sequentially) from the pool of captured analytes. In some embodiments, enrichment by PCR amplification includes multiple rounds of amplification. For example, enrichment by PCR amplification can include nested PCR reactions using different primers that are specific for the analyte or analytes of interest. In some embodiments, enrichment by amplification can be performed using an amplification method that is not PCR. A non-limiting example of a non-PCR amplification method is rolling circle amplification. Other non-PCR amplification methods are known in the art.

[0859] In some embodiments, an oligonucleotide with sequence complementarity to a captured analyte or analytes of interest, or complement thereof, can be used to enrich the captured analyte or analytes of interest from the pool of captured analytes. In some embodiments, an oligonucleotide with sequence complementarity to a captured analyte or analytes of interest (or another domain the capture probe), or complement thereof, can include one or more functional moieties that are useful in the enrichment process. For example, biotinylated oligonucleotides with sequence complementary to one or more analytes interest, or complements thereof, can bind to the analyte(s) of interest and can be selected using biotinylation-streptavidin affinity using any of a variety of methods known in the art (e.g., streptavidin beads). In some embodiments, oligonucleotides with sequence complementary to one or more analytes interest, or complements thereof, include a magnetic moiety (e.g., a magnetic bead) that can be used in the enrichment process.

[0860] Additionally or alternatively, one or more species of analyte (e.g., mitochondrial DNA or RNA) can be down-selected (e.g., removed) using any of a variety of methods. In some embodiments, such down-selection of analytes that are not of interest can result in improved capture of other types of analytes that are of interest. For example, probes can be administered to a sample that selectively hybridize to ribosomal RNA (rRNA), thereby reducing the pool and concentration of rRNA in the sample. In some embodiments, such down-selection can result in improved capture of other types of RNA due to the reduction in non-specific RNA present in the sample. Additionally or alternatively, duplex-specific nuclease (DSN) treatment can remove rRNA (see, e.g., Archer, et al, Selective and flexible depletion of problematic sequences from RNA-seq

libraries at the cDNA stage, *BMC Genomics*, 15 401, (2014), the entire contents of which is incorporated herein by reference). In some embodiments, hydroxyapatite chromatography can be used to remove abundant species (e.g., rRNA).

(ix) RNA-Templated Ligation

[0861] In some embodiments of methods provided here, RNA-templated ligation is used to interrogate spatial gene expression in a biological sample (e.g., an FFPE tissue section). RNA-templated ligation enables sensitive measurement of specific nucleic acid analytes of interest that otherwise might be analyzed less sensitively with a whole transcriptomic approach. It provides advantages of compatibility with common histochemical stains and suitability for analysis of decade-old materials (e.g., FFPE samples) and exceedingly small microdissected tissue fragments. [0862] In some aspects, the steps of RNA-templated ligation include: (1) hybridization of pairs of probes (e.g., DNA probes) to RNA (e.g., formalin fixed RNA) within a tissue section; (2) ligation of adjacently annealed probe pairs in situ; (3) RNase H treatment that (i) releases RNA-templated ligation products from the tissue (e.g., into solution) for downstream analysis and (ii) destroys unwanted DNA-templated ligation products; and optionally, (4) amplification of RNA-templated ligation products (e.g., by multiplex PCR).

[0863] In some aspects, disclosed herein are methods of direct detection of RNA target-DNA probe duplexes without first converting RNA to cDNA by reverse transcription. In some aspects, RNA-templated ligation can include a DNA ligase. In some aspects, RNA-templated ligation can include RNA ligase. In some aspects, RNA-templated ligation can include T4 RNA ligase.

[0864] In some aspects, RNA-templated ligation is used for detection of RNA, determination of RNA sequence identity, and/or expression monitoring and transcript analysis. In some aspects, RNA-templated ligation allows for detection of a particular change in a nucleic acid (e.g., a mutation or single nucleotide polymorphism (SNP)), detection or expression of a particular nucleic acid, or detection or expression of a particular set of nucleic acids (e.g., in a similar cellular pathway or expressed in a particular pathology). In some embodiments, the methods that include RNA-templated ligation are used to analyze nucleic acids, e.g., by genotyping, quantitation of DNA copy number or RNA transcripts, localization of particular transcripts within samples, and the like. In some aspects, systems and methods provided herein that include RNA-templated ligation identify single nucleotide polymorphisms (SNPs). In some aspects, such systems and methods identify mutations.

[0865] In some aspects, disclosed herein are methods of detecting RNA expression that include bringing into contact a first probe, a second probe, and ligase (e.g., T4 RNA ligase). In some embodiments, the first probe and the second probe are designed to hybridize to a target sequence such that the 5' end of the first probe and the 3' end of the second probe are adjacent and can be ligated, wherein at least the 5'-terminal nucleotide of the first probe and at least the 3'-terminal nucleotide of the second probe are deoxyribonucleotides (DNA), and wherein the target sequence includes (e.g., is composed of) ribonucleotides (RNA). After hybridization, a ligase (e.g., T4 RNA ligase) ligates the first probe and the second probe if the target sequence is present in the target sample, but does not ligate the first probe and the second probe if the target sequence is not present in the target sample. The presence or absence of the target sequence in the biological sample can be determined by determining whether or not the first and second probes were ligated in the presence of ligase. Any of a variety of methods can be used to determine whether or not the first and second probes were ligated in the presence of ligase, including but not limited to, sequencing the ligated product, hybridizing the ligated product with a detection probe that hybridizes only when the first and second probes were ligated in the presence of ligase, restriction enzyme analysis, and other methods known in the art.

[0866] In some aspects, two or more RNA analytes are analyzed using methods that include RNA-templated ligation. In some aspects, when two or more analytes are analyzed, a first and second probe that is specific for (e.g., specifically hybridizes to) each RNA analyte are used.

[0867] In some aspects, three or more probes are used in RNA-templated ligation methods provided herein. In some embodiments, the three or more probes are designed to hybridize to a target sequence such that the three or more probes hybridize adjacent to each other such that the 5' and 3' ends of adjacent probes can be ligated. In some embodiments, the presence or absence of the target sequence in the biological sample can be determined by determining whether or not the three or more probes were ligated in the presence of ligase.

[0868] In some aspects, the first probe is a DNA probe. In some aspects, the first probe is a chimeric DNA/RNA probe. In some aspects, the second probe is a DNA probe. In some aspects, the second probe is a chimeric DNA/RNA probe.

[0869] In some aspects, methods of RNA-templated ligation utilize the T4 RNA Ligase 2 to efficiently join adjacent chimeric RNA-DNA probe pairs hybridized in situ on fixed RNA target sequences. Subsequent treatment with RNase H releases RNA-templated ligation products (e.g., into solution) for downstream analysis.

(x) Region of Interest

[0870] A biological sample can have regions that show morphological feature(s) that may indicate the presence of disease or the development of a disease phenotype. For example, morphological features at a specific site within a tumor biopsy sample can indicate the aggressiveness, therapeutic resistance, metastatic potential, migration, stage, diagnosis, and/or prognosis of cancer in a subject. A change in the morphological features at a specific site within a tumor biopsy sample often correlate with a change in the level or expression of an analyte in a cell within the specific site, which can, in turn, be used to provide information regarding the aggressiveness, therapeutic resistance, metastatic potential, migration, stage, diagnosis, and/or prognosis of cancer in a subject. A region or area within a biological sample that is selected for specific analysis (e.g., a region in a biological sample that has morphological features of interest) is often described as “a region of interest.”

[0871] A region of interest in a biological sample can be used to analyze a specific area of interest within a biological sample, and thereby, focus experimentation and data gathering to a specific region of a biological sample (rather than an entire biological sample). This results in increased time efficiency of the analysis of a biological sample.

[0872] A region of interest can be identified in a biological sample using a variety of different techniques, e.g., expansion microscopy, bright field microscopy, dark field microscopy, phase contrast microscopy, electron microscopy, fluorescence microscopy, reflection microscopy, interference microscopy, confocal microscopy, and visual identification (e.g., by eye), and combinations thereof. For example, the staining and imaging of a biological sample can be performed to identify a region of interest. In some examples, the region of interest can correspond to a specific structure of cytoarchitecture. In some embodiments, a biological sample can be stained prior to visualization to provide contrast between the different regions of the biological sample. The type of stain can be chosen depending on the type of biological sample and the region of the cells to be stained. In some embodiments, more than one stain can be used to visualize different aspects of the biological sample, e.g., different regions of the sample, specific cell structures (e.g., organelles), or different cell types. In other embodiments, the biological sample can be visualized or imaged without staining the biological sample.

[0873] In some embodiments, staining and imaging a biological sample prior to contacting the biological sample with a spatial array is performed to select samples for spatial analysis. In some embodiments, the staining includes applying a fiducial marker as described herein, including fluorescent, radioactive, chemiluminescent, or colorimetric detectable markers. In some embodiments, the staining and imaging of biological samples allows the user to identify the specific sample (or region of interest) the user wishes to assess.

[0874] In some examples, an array (e.g., any of the exemplary arrays described herein) can be contacted with only a portion of a biological sample (e.g., a cell, a tissue section, or a region of

interest). In some examples, a biological sample is contacted with only a portion of an array (e.g., any of the exemplary arrays described herein). In some embodiments, capture probes on an array corresponding to regions of interest of a biological sample (e.g., proximal to the region of interest) can be selectively cleaved and analyzed. For example, capture probes on an array may be deactivated or eliminated outside of areas corresponding to regions of interest of a biological sample. In some embodiments, capture probes including a photocleavable bond and on the array in areas corresponding to regions of interest of a biological sample can be selectively cleaved by using light. A mirror, mirror array, a lens, a moving stage, and/or a photomask can be used to direct the light to regions of the array that correspond to areas outside one or more regions of interest in the biological sample. Some embodiments include deactivating or eliminating capture probes, e.g., capture probes comprising a photocleavable bond as described herein, using light. In some embodiments, a laser, e.g., a scanning laser, can be used to deactivate or eliminate capture probes. In some embodiments, the eliminated member of the plurality of capture probes can be washed away. In some embodiments, regions of interest can be labeled with different heavy metals, and a laser can sequentially ablate these regions of interest before mass spectrometry identification. A laser can, for example, deactivate or eliminate capture probes through UV light destruction of DNA, heat, inducing a chemical reaction that prevents the capture probes from moving to the next step, inducing photocleavage of a photocleavable bond, or a combination thereof. In some examples, a portion of the array can be deactivated such that it does not interact with the analytes in the biological sample (e.g., optical deactivation, chemical deactivation, heat deactivation, or blocking of the capture probes in the array (e.g., using blocking probes)). In some embodiments, the capture probes can be blocked (e.g., masked or modified) prior to contacting the biological sample with the array. For example, the free 3' end of the capture probe can be blocked or modified prior to contacting the biological sample with the array to avoid modification of the capture probes (e.g., to avoid the removal or modification of the free 3' OH group on the end of the capture probes). In some embodiments, the capture probes can be blocked prior to contacting the biological sample to the array. In some embodiments, the blocking probe is used to block or modify the free 3' end of the capture domain of the capture probe. In some embodiments, the blocking probes can be hybridized to the capture probe. In some embodiments, the free 3' end of the capture domain can be blocked by chemical modification.

[0875] In some examples, a region of interest can be removed from a biological sample and then the region of interest can be contacted to the array (e.g., any of the arrays described herein). A region of interest can be removed from a biological sample using microsurgery, laser capture microdissection, chunking, a microtome, dicing, trypsinization, labelling, and/or fluorescence-assisted cell sorting, and the like. In some embodiments, the biological sample is dissected using laser capture microdissection, retaining one or more portions of biological sample for analysis and/or discarding one or more portions of biological sample. In some embodiments, the biological sample is dissected on the array. In some embodiments, one or more regions of interest are selected using spatially addressable microelectrode arrays.

[0876] In some examples, a region of interest can be permeabilized or lysed while areas outside the region of interest are not permeabilized or lysed (e.g., Kashyap et al. *Sci Rep.* 2016; 6:29579, herein incorporated by reference in its entirety). For example, in some embodiments, a region of interest can be contacted with a hydrogel comprising a permeabilization or lysing reagent. In some embodiments, the area(s) outside the region of interest are not contacted with the hydrogel comprising the permeabilization or lysing reagent. In some embodiments, the eliminated members of the plurality of capture probes are washed away after the permeabilization of the biological sample.

(f) Partitioning

[0877] As discussed above, in some embodiments, the sample can optionally be separated into single cells, cell groups, or other fragments/pieces that are smaller than the original, unfragmented

sample. Each of these smaller portions of the sample can be analyzed to obtain spatially-resolved analyte information for the sample.

[0878] For samples that have been separated into smaller fragments- and particularly, for samples that have been disaggregated, dissociated, or otherwise separated into individual cells-one method for analyzing the fragments involves separating the fragments into individual partitions (e.g., fluid droplets), and then analyzing the contents of the partitions. In general, each partition maintains separation of its own contents from the contents of other partitions. The partition can be a droplet in an emulsion, for example.

[0879] The partitions can be flowable within fluid streams. The partitions can include, for example, micro-vesicles that have an outer barrier surrounding an inner fluid center or core. In some cases, the partitions can include a porous matrix that is capable of entraining and/or retaining materials within its matrix. The partitions can be droplets of a first phase within a second phase, wherein the first and second phases are immiscible. For example, the partitions can be droplets of aqueous fluid within a non-aqueous continuous phase (e.g., oil phase). In another example, the partitions can be droplets of a non-aqueous fluid within an aqueous phase. In some examples, the partitions can be provided in a water-in-oil emulsion or oil-in-water emulsion. A variety of different vessels are described in, for example, U.S. Patent Application Publication No. 2014/0155295, the entire contents of which are incorporated herein by reference. Emulsion systems for creating stable droplets in non-aqueous or oil continuous phases are described, for example, in U.S. Patent Application Publication No. 2010/0105112, the entire contents of which are incorporated herein by reference.

[0880] For droplets in an emulsion, allocating individual particles to discrete partitions can be accomplished, for example, by introducing a flowing stream of particles in an aqueous fluid into a flowing stream of a non-aqueous fluid, such that droplets are generated at the junction of the two streams. Fluid properties (e.g., fluid flow rates, fluid viscosities, etc.), particle properties (e.g., volume fraction, particle volume, particle concentration, etc.), microfluidic architectures (e.g., channel geometry, etc.), and other parameters can be adjusted to control the occupancy of the resulting partitions (e.g., number of analytes per partition, number of beads per partition, etc.). For example, partition occupancy can be controlled by providing the aqueous stream at a certain concentration and/or flow rate of analytes.

[0881] To generate single analyte partitions, the relative flow rates of the immiscible fluids can be selected such that, on average, the partitions can contain less than one analyte per partition to ensure that those partitions that are occupied are primarily singly occupied. In some cases, partitions among a plurality of partitions can contain at most one analyte. In some embodiments, the various parameters (e.g., fluid properties, particle properties, microfluidic architectures, etc.) can be selected or adjusted such that a majority of partitions are occupied, for example, allowing for only a small percentage of unoccupied partitions. The flows and channel architectures can be controlled as to ensure a given number of singly occupied partitions, less than a certain level of unoccupied partitions and/or less than a certain level of multiply occupied partitions.

[0882] The channel segments described herein can be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure can have a variety of geometries. For example, a microfluidic channel structure can have one or more than one channel junction. As another example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying particles that meet at a channel junction. Fluid can be directed to flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can include compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid can also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary, and/or gravity flow.

[0883] In addition to cells and/or analytes, a partition can include additional components, and in particular, one or more beads. A partition can include a single gel bead, a single cell bead, or both a single cell bead and single gel bead. A variety of different beads can be incorporated into partitions. In some embodiments, for example, non-barcoded beads can be incorporated into the partitions. For example, where the biological particle (e.g., a cell) that is incorporated into the partitions carries one or more barcodes (e.g., spatial barcode(s), UMI(s), and combinations thereof), the bead can be a non-barcoded bead.

[0884] In some embodiments, a barcode carrying bead can be incorporated into partitions. In general, an individual bead can be coupled to any number of individual nucleic acid molecules, for example, from one to tens to hundreds of thousands or even millions of individual nucleic acid molecules. The respective barcodes for the individual nucleic acid molecules can include both common sequence segments or relatively common sequence segments and variable or unique sequence segments between different individual nucleic acid molecules coupled to the same bead. For example, a nucleic acid molecule (e.g., an oligonucleotide), can be coupled to a bead by a releasable linkage (e.g., a disulfide linker), wherein the nucleic acid molecule can be or include a barcode. For example, barcodes can be injected into droplets previous to, subsequent to, or concurrently with droplet generation. The delivery of the barcodes to a particular partition allows for the later attribution of the characteristics of the individual biological particle to the particular partition. Barcodes can be delivered, for example on a nucleic acid molecule (e.g., an oligonucleotide), to a partition via any suitable mechanism. Barcoded nucleic acid molecules can be delivered to a partition via a microcapsule. A microcapsule, in some instances, can include a bead. The same bead can be coupled (e.g., via releasable linkage) to one or more other nucleic acid molecules.

[0885] In some embodiments, a microcapillary array with spatially barcoded beads can be generated. A plurality of spatially barcoded beads can be flowed into channels on a microcapillary array such that each microcapillary channel can be loaded with one spatially barcoded bead. In some embodiments, the spatially barcoded bead microcapillary array can be contacted to a biological sample for subsequent spatial analysis of biological analytes within the biological sample. In some embodiments, a microcapillary array channel can mechanically compress the biological sample and form fluidically isolated reaction chambers. In some embodiments, reagents (e.g., enzymes, nucleic acids) are introduced into the reaction chambers. The reagents can be sealed (e.g., by silicone oil, mineral oil) within the reaction chambers and incubated, allowing for a cellular and/or nuclear permeabilization reaction to occur. In some embodiments, biological analytes (e.g., DNA, RNA, proteins, metabolites, small molecules, and lipids) are released and captured onto the spatially barcoded microcapillary array, preserving their spatial information. In some embodiments, spatial analysis using a spatially barcoded feature microcapillary array can be used to obtain spatial information of the biological sample analytes at single-cell resolution.

[0886] The nucleic acid molecule can include a functional domain that can be used in subsequent processing. For example, the functional domain can include one or more of a sequencer specific flow cell attachment sequence (e.g., a P5 sequence for Illumina® sequencing systems) and a sequencing primer sequence (e.g., a R1 primer for Illumina® sequencing systems). The nucleic acid molecule can include a barcode sequence for use in barcoding the sample (e.g., DNA, RNA, protein, etc.). In some cases, the barcode sequence can be bead-specific such that the barcode sequence is common to all nucleic acid molecules coupled to the same bead. Alternatively or in addition, the barcode sequence can be partition-specific such that the barcode sequence is common to all nucleic acid molecules coupled to one or more beads that are partitioned into the same partition. The nucleic acid molecule can include a specific priming sequence, such as an mRNA specific priming sequence (e.g., poly(T) sequence), a targeted priming sequence, and/or a random priming sequence. The nucleic acid molecule can include an anchoring sequence to ensure that the specific priming sequence hybridizes at the sequence end (e.g., of the mRNA). For example, the

anchoring sequence can include a random short sequence of nucleotides, such as a 1-mer, 2-mer, 3-mer or longer sequence, which can ensure that a poly(T) segment is more likely to hybridize at the sequence end of the poly(A) tail of the mRNA.

[0887] The nucleic acid molecule can include a unique molecular identifying sequence (e.g., unique molecular identifier (UMI)). In some embodiments, the unique molecular identifying sequence can include from about 5 to about 8 nucleotides. Alternatively, the unique molecular identifying sequence can include less than about 5 or more than about 8 nucleotides. The unique molecular identifying sequence can be a unique sequence that varies across individual nucleic acid molecules coupled to a single bead. In some embodiments, the unique molecular identifying sequence can be a random sequence (e.g., such as a random N-mer sequence). For example, the UMI can provide a unique identifier of the starting mRNA molecule that was captured, in order to allow quantitation of the number of original expressed RNA.

[0888] A partition can also include one or more reagents. Unique identifiers, such as barcodes, can be injected into the droplets previous to, subsequent to, or concurrently with droplet generation, such as via a microcapsule (e.g., bead). Microfluidic channel networks (e.g., on a chip) can be utilized to generate partitions. Alternative mechanisms can also be employed in the partitioning of individual biological particles, including porous membranes through which aqueous mixtures of cells are extruded into non-aqueous fluids.

[0889] In some embodiments, barcoded nucleic acid molecules can be initially associated with a microcapsule and then released from the microcapsule. Release of the barcoded nucleic acid molecules can be passive (e.g., by diffusion out of the microcapsule). In addition or alternatively, release from the microcapsule can be upon application of a stimulus which allows the barcoded nucleic acid molecules to dissociate or to be released from the microcapsule. Such stimulus can disrupt the microcapsule, an interaction that couples the barcoded nucleic acid molecules to or within the microcapsule, or both. Such stimulus can include, for example, a thermal stimulus, photo-stimulus, chemical stimulus (e.g., change in pH or use of a reducing agent(s)), a mechanical stimulus, a radiation stimulus; a biological stimulus (e.g., enzyme), or any combination thereof.

[0890] In some embodiments, one more barcodes (e.g., spatial barcodes, UMIs, or a combination thereof) can be introduced into a partition as part of the analyte. As described previously, barcodes can be bound to the analyte directly, or can form part of a capture probe or analyte capture agent that is hybridized to, conjugated to, or otherwise associated with an analyte, such that when the analyte is introduced into the partition, the barcode(s) are introduced as well. As described above, FIG. 16 shows an example of a microfluidical channel structure for partitioning individual analytes (e.g., cells) into discrete partitions.

[0891] FIG. 16 shows an example of a microfluidic channel structure for partitioning individual analytes (e.g., cells) into discrete partitions. The channel structure can include channel segments **1601**, **1602**, **1603**, and **1604** communicating at a channel junction **1605**. In operation, a first aqueous fluid **1606** that includes suspended biological particles (or cells) **1607** may be transported along channel segment **1601** into junction **1605**, while a second fluid **1608** that is immiscible with the aqueous fluid **1606** is delivered to the junction **1605** from each of channel segments **1602** and **1603** to create discrete droplets **1609**, **1610** of the first aqueous fluid **1606** flowing into channel segment **1604**, and flowing away from junction **1605**. The channel segment **1604** may be fluidically coupled to an outlet reservoir where the discrete droplets can be stored and/or harvested. A discrete droplet generated may include an individual biological particle **1607** (such as droplets **1609**). A discrete droplet generated may include more than one individual biological particle **1607**. A discrete droplet may contain no biological particle **1607** (such as droplet **1610**). Each discrete partition may maintain separation of its own contents (e.g., individual biological particle **1607**) from the contents of other partitions.

[0892] FIG. 17A shows another example of a microfluidic channel structure **1700** for delivering

beads to droplets. The channel structure includes channel segments **1701**, **1702**, **1703**, **1704** and **1705** communicating at a channel junction **1706**. During operation, the channel segment **1701** can transport an aqueous fluid **1707** that includes a plurality of beads **1708** along the channel segment **1701** into junction **1706**. The plurality of beads **1708** can be sourced from a suspension of beads. For example, the channel segment **1701** can be connected to a reservoir that includes an aqueous suspension of beads **1708**. The channel segment **1702** can transport the aqueous fluid **1707** that includes a plurality of particles **1709** (e.g., cells) along the channel segment **1702** into junction **1706**. In some embodiments, the aqueous fluid **1707** in either the first channel segment **1701** or the second channel segment **1702**, or in both segments, can include one or more reagents, as further described below.

[0893] A second fluid **1710** that is immiscible with the aqueous fluid **1707** (e.g., oil) can be delivered to the junction **1706** from each of channel segments **1703** and **1704**. Upon meeting of the aqueous fluid **1707** from each of channel segments **1701** and **1702** and the second fluid **1710** from each of channel segments **1703** and **1704** at the channel junction **1706**, the aqueous fluid **1707** can be partitioned as discrete droplets **1711** in the second fluid **1710** and flow away from the junction **1706** along channel segment **1705**. The channel segment **1705** can deliver the discrete droplets to an outlet reservoir fluidly coupled to the channel segment **1705**, where they can be harvested.

[0894] As an alternative, the channel segments **1701** and **1702** can meet at another junction upstream of the junction **1706**. At such junction, beads and biological particles can form a mixture that is directed along another channel to the junction **1706** to yield droplets **1711**. The mixture can provide the beads and biological particles in an alternating fashion, such that, for example, a droplet includes a single bead and a single biological particle.

[0895] The second fluid **1710** can include an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **1711**.

[0896] The partitions described herein can include small volumes, for example, less than about 10 microliters (μL), 5 μL , 1 μL , 900 picoliters (pL), 800 pL, 700 pL, 600 pL, 500 pL, 400 pL, 300 pL, 200 pL, 100 pL, 50 pL, 20 pL, 10 pL, 1 pL, 500 nanoliters (nL), 100 nL, 50 nL, or less.

[0897] In the foregoing discussion, droplets with beads were formed at the junction of different fluid streams. In some embodiments, droplets can be formed by gravity-based partitioning methods.

[0898] FIG. **17B** shows a cross-section view of another example of a microfluidic channel structure **1750** with a geometric feature for controlled partitioning. A channel structure **1750** can include a channel segment **1752** communicating at a channel junction **1758** (or intersection) with a reservoir **1754**. In some instances, the channel structure **1750** and one or more of its components can correspond to the channel structure **1700** and one or more of its components.

[0899] An aqueous fluid **1760** comprising a plurality of particles **1756** may be transported along the channel segment **1752** into the junction **1758** to meet a second fluid **1762** (e.g., oil, etc.) that is immiscible with the aqueous fluid **1760** in the reservoir **1754** to create droplets **1764** of the aqueous fluid **1760** flowing into the reservoir **1754**. At the junction **1758** where the aqueous fluid **1760** and the second fluid **1762** meet, droplets can form based on factors such as the hydrodynamic forces at the junction **1758**, relative flow rates of the two fluids **1760**, **1762**, fluid properties, and certain geometric parameters (e.g., Δh , etc.) of the channel structure **1750**. A plurality of droplets can be collected in the reservoir **1754** by continuously injecting the aqueous fluid **1760** from the channel segment **1752** at the junction **1758**.

[0900] A discrete droplet generated may comprise one or more particles of the plurality of particles **1756**. As described elsewhere herein, a particle may be any particle, such as a bead, cell bead, gel bead, biological particle, macromolecular constituents of biological particle, or other particles. Alternatively, a discrete droplet generated may not include any particles.

[0901] In some instances, the aqueous fluid **1760** can have a substantially uniform concentration or

frequency of particles **1756**. As described elsewhere herein, the particles **1756** (e.g., beads) can be introduced into the channel segment **1752** from a separate channel (not shown in FIG. **17**). The frequency of particles **1756** in the channel segment **1752** may be controlled by controlling the frequency in which the particles **1756** are introduced into the channel segment **1752** and/or the relative flow rates of the fluids in the channel segment **1752** and the separate channel. In some instances, the particles **1756** can be introduced into the channel segment **1752** from a plurality of different channels, and the frequency controlled accordingly. In some instances, different particles may be introduced via separate channels. For example, a first separate channel can introduce beads and a second separate channel can introduce biological particles into the channel segment **1752**. The first separate channel introducing the beads may be upstream or downstream of the second separate channel introducing the biological particles.

[0902] In some instances, the second fluid **1762** may not be subjected to and/or directed to any flow in or out of the reservoir **1754**. For example, the second fluid **1762** may be substantially stationary in the reservoir **1754**. In some instances, the second fluid **1762** may be subjected to flow within the reservoir **1754**, but not in or out of the reservoir **1754**, such as via application of pressure to the reservoir **1754** and/or as affected by the incoming flow of the aqueous fluid **1760** at the junction **1758**. Alternatively, the second fluid **1762** may be subjected and/or directed to flow in or out of the reservoir **1754**. For example, the reservoir **1754** can be a channel directing the second fluid **1762** from upstream to downstream, transporting the generated droplets.

[0903] The channel structure **1750** at or near the junction **1758** may have certain geometric features that at least partly determine the volumes and/or shapes of the droplets formed by the channel structure **1750**. The channel segment **1752** can have a first cross-section height, h_1 , and the reservoir **1754** can have a second cross-section height, h_2 . The first cross-section height, h_1 , and the second cross-section height, h_2 , may be different, such that at the junction **1758**, there is a height difference of Δh . The second cross-section height, h_2 , may be greater than the first cross-section height, h_1 . In some instances, the reservoir may thereafter gradually increase in cross-section height, for example, the more distant it is from the junction **1758**. In some instances, the cross-section height of the reservoir may increase in accordance with expansion angle, β , at or near the junction **1758**. The height difference, Δh , and/or expansion angle, β , can allow the tongue (portion of the aqueous fluid **1760** leaving channel segment **1752** at junction **1758** and entering the reservoir **1754** before droplet formation) to increase in depth and facilitate decrease in curvature of the intermediately formed droplet. For example, droplet volume may decrease with increasing height difference and/or increasing expansion angle.

[0904] The height difference, Δh , can be at least about 1 μm . Alternatively, the height difference can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500 μm or more. Alternatively, the height difference can be at most about 500, 400, 300, 200, 100, 90, 80, 70, 60, 50, 45, 40, 35, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 μm or less. In some instances, the expansion angle, β , may be between a range of from about 0.5° to about 4° , from about 0.1° to about 10° , or from about 0° to about 90° . For example, the expansion angle can be at least about 0.01° , 0.1° , 0.2° , 0.3° , 0.4° , 0.5° , 0.6° , 0.7° , 0.8° , 0.9° , 1° , 2° , 3° , 4° , 5° , 6° , 7° , 8° , 9° , 10° , 15° , 20° , 25° , 30° , 35° , 40° , 45° , 50° , 55° , 60° , 65° , 70° , 75° , 80° , 85° , or higher. In some instances, the expansion angle can be at most about 89° , 88° , 87° , 86° , 85° , 84° , 83° , 82° , 81° , 80° , 75° , 70° , 65° , 60° , 55° , 50° , 45° , 40° , 35° , 30° , 25° , 20° , 15° , 10° , 9° , 8° , 7° , 6° , 5° , 4° , 3° , 2° , 1° , 0.1° , 0.01° , or less.

[0905] In some instances, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be between about 0.04 microliters (μL)/minute (min) and about 40 $\mu\text{L}/\text{min}$. In some instances, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be between about 0.01 microliters (μL)/minute (min) and about 100 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be less than about 0.01 $\mu\text{L}/\text{min}$. alternatively, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be greater than about 40 $\mu\text{L}/\text{min}$, such

as 45 $\mu\text{L}/\text{min}$, 50 $\mu\text{L}/\text{min}$, 55 $\mu\text{L}/\text{min}$, 60 $\mu\text{L}/\text{min}$, 65 $\mu\text{L}/\text{min}$, 70 $\mu\text{L}/\text{min}$, 75 $\mu\text{L}/\text{min}$, 80 $\mu\text{L}/\text{min}$, 85 $\mu\text{L}/\text{min}$, 90 $\mu\text{L}/\text{min}$, 95 $\mu\text{L}/\text{min}$, 100 $\mu\text{L}/\text{min}$, 110 $\mu\text{L}/\text{min}$, 120 $\mu\text{L}/\text{min}$, 130 $\mu\text{L}/\text{min}$, 140 $\mu\text{L}/\text{min}$, 150 $\mu\text{L}/\text{min}$, or greater. At lower flow rates, such as flow rates of about less than or equal to 10 microliters/minute, the droplet radius may not be dependent on the flow rate of the aqueous fluid **1760** entering the junction **1758**. The second fluid **1762** may be stationary, or substantially stationary, in the reservoir **1754**. Alternatively, the second fluid **1762** may be flowing, such as at the above flow rates described for the aqueous fluid **1760**.

[0906] While FIG. **17B** illustrates the height difference, Δh , being abrupt at the junction **1758** (e.g., a step increase), the height difference may increase gradually (e.g., from about 0 μm to a maximum height difference). Alternatively, the height difference may decrease gradually (e.g., taper) from a maximum height difference. A gradual increase or decrease in height difference, as used herein, may refer to a continuous incremental increase or decrease in height difference, wherein an angle between any one differential segment of a height profile and an immediately adjacent differential segment of the height profile is greater than 90° . For example, at the junction **1758**, a bottom wall of the channel and a bottom wall of the reservoir can meet at an angle greater than 90° .

Alternatively or in addition, a top wall (e.g., ceiling) of the channel and a top wall (e.g., ceiling) of the reservoir can meet an angle greater than 90° . A gradual increase or decrease may be linear or non-linear (e.g., exponential, sinusoidal, etc.). Alternatively or in addition, the height difference may variably increase and/or decrease linearly or non-linearly. While FIG. **17B** illustrates the expanding reservoir cross-section height as linear (e.g., constant expansion angle, β), the cross-section height may expand non-linearly. For example, the reservoir may be defined at least partially by a dome-like (e.g., hemispherical) shape having variable expansion angles. The cross-section height may expand in any shape.

[0907] FIG. **17C** depicts a workflow wherein cells are partitioned into droplets along with barcode-bearing beads **1770**. See FIG. **17A**. The droplet forms an isolated reaction chamber wherein the cells can be lysed **1771** and target analytes within the cells can then be captured **1772** and amplified **1773**, **1774** according to previously described methods. After sequence library preparation clean-up **1775**, the material is sequenced and/or quantified **1776** according to methods described herein. For example, the workflow shown in FIG. **17C** can be used with a biological sample on an array, where the features of the array have been delivered to the substrate via a droplet manipulation system. In some embodiments, capture probes on the features can specifically bind analytes present in the biological sample. In some embodiments, the features can be removed from the substrate (e.g., removed by any method described herein) and partitioned into droplets with barcode-bearing beads for further analysis according to methods described herein.

[0908] It should be noted that while the example workflow in FIG. **17C** includes steps specifically for the analysis of mRNA, analogous workflows can be implemented for a wide variety of other analytes, including any of the analytes described previously.

[0909] By way of example, in the context of analyzing sample RNA as shown in FIG. **17C**, the poly(T) segment of one of the released nucleic acid molecules (e.g., from the bead) can hybridize to the poly(A) tail of a mRNA molecule. Reverse transcription can result in a cDNA transcript of the mRNA, which transcript includes each of the sequence segments of the nucleic acid molecule. If the nucleic acid molecule includes an anchoring sequence, it will more likely hybridize to and prime reverse transcription at the sequence end of the poly(A) tail of the mRNA.

[0910] Within any given partition, all of the cDNA transcripts of the individual mRNA molecules can include a common barcode sequence segment. However, the transcripts made from the different mRNA molecules within a given partition can vary at the unique molecular identifying sequence segment (e.g., UMI segment). Beneficially, even following any subsequent amplification of the contents of a given partition, the number of different UMIs can be indicative of the quantity of mRNA originating from a given partition. As noted above, the transcripts can be amplified, cleaned up and sequenced to identify the sequence of the cDNA transcript of the mRNA, as well as

to sequence the barcode segment and the UMI segment. While a poly(T) primer sequence is described, other targeted or random priming sequences can also be used in priming the reverse transcription reaction. Likewise, although described as releasing the barcoded oligonucleotides into the partition, in some cases, the nucleic acid molecules bound to the bead can be used to hybridize and capture the mRNA on the solid phase of the bead, for example, in order to facilitate the separation of the RNA from other cell contents.

[0911] In some embodiments, partitions include precursors that include a functional group that is reactive or capable of being activated such that it becomes reactive can be polymerized with other precursors to generate gel beads that include the activated or activatable functional group. The functional group can then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the gel beads. For example, some precursors featuring a carboxylic acid (COOH) group can co-polymerize with other precursors to form a bead that also includes a COOH functional group. In some cases, acrylic acid (a species comprising free COOH groups), acrylamide, and bis(acryloyl) cystamine can be co-polymerized together to generate a bead with free COOH groups. The COOH groups of the bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species comprising an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) comprising a moiety to be linked to the bead.

[0912] In some embodiments, a bead can be formed from materials that include degradable chemical cross-linkers, such as BAC or cystamine. Degradation of such degradable cross-linkers can be accomplished through a number of mechanisms. In some examples, a bead can be contacted with a chemical degrading agent that can induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as dithiothreitol (DTT). Additional examples of reducing agents can include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris (2-carboxyethyl) phosphine (TCEP), or combinations thereof. A reducing agent can degrade the disulfide bonds formed between gel precursors forming the bead, and thus, degrade the bead.

[0913] A degradable bead can include one or more species with a labile bond such that, when the bead/species is exposed to the appropriate stimulus, the bond is broken and the bead degrades within the partition. The labile bond can be a chemical bond (e.g., covalent bond, ionic bond) or can be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.). In some embodiments, a cross-linker used to generate a bead can include a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, a polyacrylamide bead featuring cystamine and linked, via a disulfide bond, to a barcode sequence, can be combined with a reducing agent within a droplet of a water-in-oil emulsion. Within the droplet, the reducing agent can break the various disulfide bonds, resulting in bead degradation and release of the barcode sequence into the aqueous, inner environment of the droplet. In another example, heating of a droplet with a bead-bound barcode sequence in basic solution can also result in bead degradation and release of the attached barcode sequence into the aqueous, inner environment of the droplet. The free species (e.g., oligonucleotides, nucleic acid molecules) can interact with other reagents contained in the partition.

[0914] A degradable bead can be useful in more quickly releasing an attached species (e.g., a nucleic acid molecule, a barcode sequence, a primer, etc.) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species can have greater mobility and accessibility to other species in solution upon degradation of the bead. In some embodiments, a species can also be attached to a degradable bead via a

degradable linker (e.g., disulfide linker). The degradable linker can respond to the same stimuli as the degradable bead or the two degradable species can respond to different stimuli. For example, a barcode sequence can be attached, via a disulfide bond, to a polyacrylamide bead comprising cystamine. Upon exposure of the barcoded-bead to a reducing agent, the bead degrades and the barcode sequence is released upon breakage of both the disulfide linkage between the barcode sequence and the bead and the disulfide linkages of the cystamine in the bead.

[0915] Any suitable number of species (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the species (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration can be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In some cases, the pre-defined concentration of the primer can be limited by the process of producing nucleic acid molecule (e.g., oligonucleotide) bearing beads.

[0916] As will be appreciated from the above description, while referred to as degradation of a bead, in many embodiments, degradation can refer to the disassociation of a bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species can be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore volumes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore volume due to osmotic swelling of a bead can permit the release of entrained species within the bead. In some embodiments, osmotic shrinking of a bead can cause a bead to better retain an entrained species due to pore volume contraction.

[0917] Numerous chemical triggers can be used to trigger the degradation of beads within partitions. Examples of these chemical changes can include, but are not limited to pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

[0918] In certain embodiments, a change in pH of a solution, such as an increase in pH, can trigger degradation of a bead. In other embodiments, exposure to an aqueous solution, such as water, can trigger hydrolytic degradation, and thus degradation of the bead. In some cases, any combination of stimuli can trigger degradation of a bead. For example, a change in pH can enable a chemical agent (e.g., DTT) to become an effective reducing agent.

[0919] Beads can also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat can cause melting of a bead such that a portion of the bead degrades. In other cases, heat can increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat can also act upon heat-sensitive polymers used as materials to construct beads.

[0920] In addition to beads and analytes, partitions that are formed can include a variety of different reagents and species. For example, when lysis reagents are present within the partitions, the lysis reagents can facilitate the release of analytes within the partition. Examples of lysis agents include bioactive reagents, such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, etc., such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other lysis enzymes available from, e.g., Sigma-Aldrich, Inc. (St. Louis, MO), as well as other commercially available lysis enzymes. Other lysis agents can additionally or alternatively be co-partitioned to cause the release analytes into the partitions. For example, in some cases, surfactant-based lysis solutions can be used to lyse cells, although these can be less desirable for emulsion based systems where the surfactants can interfere with stable emulsions. In some embodiments, lysis solutions can include non-ionic surfactants such as, for example, TritonX-100 and Tween 20. In some embodiments, lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption can also be used in

certain embodiments, e.g., non-emulsion based partitioning such as encapsulation of analytes that can be in addition to or in place of droplet partitioning, where any pore size of the encapsulate is sufficiently small to retain nucleic acid fragments of a given volume, following cellular disruption. [0921] Examples of other species that can be co-partitioned with analytes in the partitions include, but are not limited to, DNase and RNase inactivating agents or inhibitors, such as proteinase K, chelating agents, such as EDTA, and other reagents employed in removing or otherwise reducing negative activity or impact of different cell lysate components on subsequent processing of nucleic acids. Additional reagents can also be co-partitioned, including endonucleases to fragment DNA, DNA polymerase enzymes and dNTPs used to amplify nucleic acid fragments and to attach the barcode molecular tags to the amplified fragments.

[0922] Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers and oligonucleotides, and switch oligonucleotides (also referred to herein as “switch oligos” or “template switching oligonucleotides”) which can be used for template switching. In some embodiments, template switching can be used to increase the length of a cDNA. Template switching can be used to append a predefined nucleic acid sequence to the cDNA. In an example of template switching, cDNA can be generated from reverse transcription of a template, e.g., cellular mRNA, where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., poly(C), to the cDNA in a template independent manner. Switch oligos can include sequences complementary to the additional nucleotides, e.g., poly(G). The additional nucleotides (e.g., poly(C)) on the cDNA can hybridize to the additional nucleotides (e.g., poly(G)) on the switch oligo, whereby the switch oligo can be used by the reverse transcriptase as template to further extend the cDNA.

Template switching oligonucleotides can include a hybridization region and a template region. The hybridization region can include any sequence capable of hybridizing to the target. In some cases, the hybridization region includes a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases can include 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases or more than 5 G bases. The template sequence can include any sequence to be incorporated into the cDNA. In some cases, the template region includes at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. Switch oligos can include deoxyribonucleic acids; ribonucleic acids; bridged nucleic acids, modified nucleic acids including 2-Aminopurine, 2,6-Diaminopurine (2-Amino-dA), inverted dT, 5-Methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine), locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' Fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), and combinations of the foregoing.

[0923] In some embodiments, beads that are partitioned with the analyte can include different types of oligonucleotides bound to the bead, where the different types of oligonucleotides bind to different types of analytes. For example, a bead can include one or more first oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a first type of analyte, such as mRNA for example, and one or more second oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a second type of analyte, such as gDNA for example. Partitions can also include lysis agents that aid in releasing nucleic acids from the co-partitioned cell, and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break covalent linkages between the oligonucleotides and the bead, releasing the oligonucleotides into the partition. The released barcoded oligonucleotides (which can also be barcoded) can hybridize with mRNA released from the cell and also with gDNA released from the cell.

[0924] Barcoded constructs thus formed from hybridization can include a first type of construct that includes a sequence corresponding to an original barcode sequence from the bead and a sequence corresponding to a transcript from the cell, and a second type of construct that includes a

sequence corresponding to the original barcode sequence from the bead and a sequence corresponding to genomic DNA from the cell. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs can then be sequenced, the sequencing data processed, and the results used to spatially characterize the mRNA and the gDNA from the cell. [0925] In another example, a partition includes a bead that includes a first type of oligonucleotide (e.g., a first capture probe) with a first barcode sequence, a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript, and a UMI barcode sequence that can uniquely identify a given transcript. The bead also includes a second type of oligonucleotide (e.g., a second capture probe) with a second barcode sequence, a targeted priming sequence that is capable of specifically hybridizing with a third barcoded oligonucleotide (e.g., an analyte capture agent) coupled to an antibody that is bound to the surface of the partitioned cell. The third barcoded oligonucleotide includes a UMI barcode sequence that uniquely identifies the antibody (and thus, the particular cell surface feature to which it is bound).

[0926] In this example, the first and second barcoded oligonucleotides include the same spatial barcode sequence (e.g., the first and second barcode sequences are the same), which permits downstream association of barcoded nucleic acids with the partition. In some embodiments, however, the first and second barcode sequences are different.

[0927] The partition also includes lysis agents that aid in releasing nucleic acids from the cell and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break a covalent linkage between the barcoded oligonucleotides and the bead, releasing them into the partition. The first type of released barcoded oligonucleotide can hybridize with mRNA released from the cell and the second type of released barcoded oligonucleotide can hybridize with the third type of barcoded oligonucleotide, forming barcoded constructs.

[0928] The first type of barcoded construct includes a spatial barcode sequence corresponding to the first barcode sequence from the bead and a sequence corresponding to the UMI barcode sequence from the first type of oligonucleotide, which identifies cell transcripts. The second type of barcoded construct includes a spatial barcode sequence corresponding to the second barcode sequence from the second type of oligonucleotide, and a UMI barcode sequence corresponding to the third type of oligonucleotide (e.g., the analyte capture agent) and used to identify the cell surface feature. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs are then sequenced, sequencing data processed, and the results used to characterize the mRNA and cell surface feature of the cell.

[0929] The foregoing discussion involves two specific examples of beads with oligonucleotides for analyzing two different analytes within a partition. More generally, beads that are partitioned can have any of the structures described previously, and can include any of the described combinations of oligonucleotides for analysis of two or more (e.g., three or more, four or more, five or more, six or more, eight or more, ten or more, 12 or more, 15 or more, 20 or more, 25 or more, 30 or more, 40 or more, 50 or more) different types of analytes within a partition. Examples of beads with combinations of different types of oligonucleotides (e.g., capture probes) for concurrently analyzing different combinations of analytes within partitions include, but are not limited to: (a) genomic DNA and cell surface features (e.g., using the analyte capture agents described herein); (b) mRNA and a lineage tracing construct; (c) mRNA and cell methylation status; (d) mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq); (e) mRNA and cell surface or intracellular proteins and/or metabolites; (f) a barcoded analyte capture agent (e.g., the MHC multimers described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); and (g) mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein). In some embodiments, a perturbation agent can be a small molecule, an antibody, a drug, an aptamer, a miRNA, a physical

environmental (e.g., temperature change), or any other known perturbation agents.

[0930] Additionally, in some embodiments, the unaggregated cell or disaggregated cells introduced and processed within partitions or droplets as described herein, can be removed from the partition, contacted with a spatial array, and spatially-barcoded according to methods described herein. For example, single cells of an unaggregated cell sample can be partitioned into partitions or droplets as described herein. The partitions or droplets can include reagents to permeabilize a cell, barcode targeted cellular analyte(s) with a cellular barcode, and amplify the barcoded analytes. The partitions or droplets can be contacted with any of the spatial arrays described herein. In some embodiments, the partition can be dissolved, such that the contents of the partition are placed in contact with the capture probes of the spatial array. The capture probes of the spatial array can then capture target analytes from the ruptured partitions or the droplets, and processed by the spatial workflows described herein.

(g) Analysis of Captured Analytes

(i) Sample Removal from an Array

[0931] In some embodiments, after contacting a biological sample with a substrate that includes capture probes, a removal step can optionally be performed to remove all or a portion of the biological sample from the substrate. In some embodiments, the removal step includes enzymatic and/or chemical degradation of cells of the biological sample. For example, the removal step can include treating the biological sample with an enzyme (e.g., a proteinase, e.g., proteinase K) to remove at least a portion of the biological sample from the substrate. In some embodiments, the removal step can include ablation of the tissue (e.g., laser ablation).

[0932] In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample (e.g., present in a biological sample), the method comprising: (a) optionally staining and/or imaging a biological sample on a substrate; (b) permeabilizing (e.g., providing a solution comprising a permeabilization reagent to) the biological sample on the substrate; (c) contacting the biological sample with an array comprising a plurality of capture probes, wherein a capture probe of the plurality captures the biological analyte; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte; wherein the biological sample is fully or partially removed from the substrate.

[0933] In some embodiments, a biological sample is not removed from the substrate. For example, the biological sample is not removed from the substrate prior to releasing a capture probe (e.g., a capture probe bound to an analyte) from the substrate. In some embodiments, such releasing comprises cleavage of the capture probe from the substrate (e.g., via a cleavage domain). In some embodiments, such releasing does not comprise releasing the capture probe from the substrate (e.g., a copy of the capture probe bound to an analyte can be made and the copy can be released from the substrate, e.g., via denaturation). In some embodiments, the biological sample is not removed from the substrate prior to analysis of an analyte bound to a capture probe after it is released from the substrate. In some embodiments, the biological sample remains on the substrate during removal of a capture probe from the substrate and/or analysis of an analyte bound to the capture probe after it is released from the substrate. In some embodiments, the biological sample remains on the substrate during removal (e.g., via denaturation) of a copy of the capture probe (e.g., complement). In some embodiments, analysis of an analyte bound to capture probe from the substrate can be performed without subjecting the biological sample to enzymatic and/or chemical degradation of the cells (e.g., permeabilized cells) or ablation of the tissue (e.g., laser ablation).

[0934] In some embodiments, at least a portion of the biological sample is not removed from the substrate. For example, a portion of the biological sample can remain on the substrate prior to releasing a capture probe (e.g., a capture probe bound to an analyte) from the substrate and/or analyzing an analyte bound to a capture probe released from the substrate. In some embodiments, at least a portion of the biological sample is not subjected to enzymatic and/or chemical

degradation of the cells (e.g., permeabilized cells) or ablation of the tissue (e.g., laser ablation) prior to analysis of an analyte bound to a capture probe from the substrate.

[0935] In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample (e.g., present in a biological sample) that include: (a) optionally staining and/or imaging a biological sample on a substrate; (b) permeabilizing (e.g., providing a solution comprising a permeabilization reagent to) the biological sample on the substrate; (c) contacting the biological sample with an array comprising a plurality of capture probes, wherein a capture probe of the plurality captures the biological analyte; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte; where the biological sample is not removed from the substrate.

[0936] In some embodiments, provided herein are methods for spatially detecting a biological analyte of interest from a biological sample that include: (a) staining and imaging a biological sample on a substrate; (b) providing a solution comprising a permeabilization reagent to the biological sample on the substrate; (c) contacting the biological sample with an array on a substrate, wherein the array comprises one or more capture probe pluralities thereby allowing the one or more pluralities of capture probes to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest; where the biological sample is not removed from the substrate.

[0937] In some embodiments, the method further includes selecting a region of interest in the biological sample to subject to spatial transcriptomic analysis. In some embodiments, one or more of the one or more capture probes include a capture domain. In some embodiments, one or more of the one or more capture probe pluralities comprise a unique molecular identifier (UMI). In some embodiments, one or more of the one or more capture probe pluralities comprise a cleavage domain. In some embodiments, the cleavage domain comprises a sequence recognized and cleaved by a uracil-DNA glycosylase, apurinic/apyrimidinic (AP) endonuclease (APE1), U uracil-specific excision reagent (USER), and/or an endonuclease VIII. In some embodiments, one or more capture probes do not comprise a cleavage domain and is not cleaved from the array.

(ii) Extended Capture Probes

[0938] In some embodiments, a capture probe can be extended (an “extended capture probe,” e.g., as described herein (e.g., Section II (b) (vii))). For example, extending a capture probe can include generating cDNA from a captured (hybridized) RNA. This process involves synthesis of a complementary strand of the hybridized nucleic acid, e.g., generating cDNA based on the captured RNA template (the RNA hybridized to the capture domain of the capture probe). Thus, in an initial step of extending a capture probe, e.g., the cDNA generation, the captured (hybridized) nucleic acid, e.g., RNA, acts as a template for the extension, e.g., reverse transcription, step.

[0939] In some embodiments, the capture probe is extended using reverse transcription. For example, reverse transcription includes synthesizing cDNA (complementary or copy DNA) from RNA, e.g., (messenger RNA), using a reverse transcriptase. In some embodiments, reverse transcription is performed while the tissue is still in place, generating an analyte library, where the analyte library includes the spatial barcodes from the adjacent capture probes. In some embodiments, the capture probe is extended using one or more DNA polymerases.

[0940] In some embodiments, a capture domain of a capture probe includes a primer for producing the complementary strand of a nucleic acid hybridized to the capture probe, e.g., a primer for DNA polymerase and/or reverse transcription. The nucleic acid, e.g., DNA and/or cDNA, molecules generated by the extension reaction incorporate the sequence of the capture probe. The extension of the capture probe, e.g., a DNA polymerase and/or reverse transcription reaction, can be performed using a variety of suitable enzymes and protocols.

[0941] In some embodiments, a full-length DNA (e.g., cDNA) molecule is generated. In some embodiments, a “full-length” DNA molecule refers to the whole of the captured nucleic acid molecule. However, if a nucleic acid (e.g., RNA) was partially degraded in the tissue sample, then

the captured nucleic acid molecules will not be the same length as the initial RNA in the tissue sample. In some embodiments, the 3' end of the extended probes, e.g., first strand cDNA molecules, is modified. For example, a linker or adaptor can be ligated to the 3' end of the extended probes. This can be achieved using single stranded ligation enzymes such as T4 RNA ligase or Circligase™ (available from Lucigen, Middleton, WI). In some embodiments, template switching oligonucleotides are used to extend cDNA in order to generate a full-length cDNA (or as close to a full-length cDNA as possible). In some embodiments, a second strand synthesis helper probe (a partially double stranded DNA molecule capable of hybridizing to the 3' end of the extended capture probe), can be ligated to the 3' end of the extended probe, e.g., first strand cDNA, molecule using a double stranded ligation enzyme such as T4 DNA ligase. Other enzymes appropriate for the ligation step are known in the art and include, e.g., Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9°N) DNA ligase (9°N™ DNA ligase, New England Biolabs), Ampligase™ (available from Lucigen, Middleton, WI), and SplintR (available from New England Biolabs, Ipswich, MA). In some embodiments, a polynucleotide tail, e.g., a poly(A) tail, is incorporated at the 3' end of the extended probe molecules. In some embodiments, the polynucleotide tail is incorporated using a terminal transferase active enzyme.

[0942] In some embodiments, double-stranded extended capture probes are treated to remove any unextended capture probes prior to amplification and/or analysis, e.g., sequence analysis. This can be achieved by a variety of methods, e.g., using an enzyme to degrade the unextended probes, such as an exonuclease enzyme, or purification columns.

[0943] In some embodiments, extended capture probes are amplified to yield quantities that are sufficient for analysis, e.g., via DNA sequencing. In some embodiments, the first strand of the extended capture probes (e.g., DNA and/or cDNA molecules) acts as a template for the amplification reaction (e.g., a polymerase chain reaction).

[0944] In some embodiments, the amplification reaction incorporates an affinity group onto the extended capture probe (e.g., RNA-cDNA hybrid) using a primer including the affinity group. In some embodiments, the primer includes an affinity group and the extended capture probes includes the affinity group. The affinity group can correspond to any of the affinity groups described previously.

[0945] In some embodiments, the extended capture probes including the affinity group can be coupled to a substrate specific for the affinity group. In some embodiments, the substrate can include an antibody or antibody fragment. In some embodiments, the substrate includes avidin or streptavidin and the affinity group includes biotin. In some embodiments, the substrate includes maltose and the affinity group includes maltose-binding protein. In some embodiments, the substrate includes maltose-binding protein and the affinity group includes maltose. In some embodiments, amplifying the extended capture probes can function to release the extended probes from the surface of the substrate, insofar as copies of the extended probes are not immobilized on the substrate.

[0946] In some embodiments, the extended capture probe or complement or amplicon thereof is released. The step of releasing the extended capture probe or complement or amplicon thereof from the surface of the substrate can be achieved in a number of ways. In some embodiments, an extended capture probe or a complement thereof is released from the array by nucleic acid cleavage and/or by denaturation (e.g., by heating to denature a double-stranded molecule).

[0947] In some embodiments, the extended capture probe or complement or amplicon thereof is released from the surface of the substrate (e.g., array) by physical means. For example, where the extended capture probe is indirectly immobilized on the array substrate, e.g., via hybridization to a surface probe, it can be sufficient to disrupt the interaction between the extended capture probe and the surface probe. Methods for disrupting the interaction between nucleic acid molecules include denaturing double stranded nucleic acid molecules are known in the art. A straightforward method for releasing the DNA molecules (i.e., of stripping the array of extended probes) is to use a solution

that interferes with the hydrogen bonds of the double stranded molecules. In some embodiments, the extended capture probe is released by applying a heated solution, such as water or buffer, of at least 85° C., e.g., at least 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99° C. In some embodiments, a solution including salts, surfactants, etc. that can further destabilize the interaction between the nucleic acid molecules is added to release the extended capture probe from the substrate.

[0948] In some embodiments, where the extended capture probe includes a cleavage domain, the extended capture probe is released from the surface of the substrate by cleavage. For example, the cleavage domain of the extended capture probe can be cleaved by any of the methods described herein. In some embodiments, the extended capture probe is released from the surface of the substrate, e.g., via cleavage of a cleavage domain in the extended capture probe, prior to the step of amplifying the extended capture probe.

[0949] In some embodiments, probes complementary to the extended capture probe can be contacted with the substrate. In some embodiments, the biological sample can be in contact with the substrate when the probes are contacted with the substrate. In some embodiments, the biological sample can be removed from the substrate prior to contacting the substrate with probes. In some embodiments, the probes can be labeled with a detectable label (e.g., any of the detectable labels described herein). In some embodiments, probes that do not specially bind (e.g., hybridize) to an extended capture probe can be washed away. In some embodiments, probes complementary to the extended capture probe can be detected on the substrate (e.g., imaging, any of the detection methods described herein).

[0950] In some embodiments, probes complementary to an extended capture probe can be about 4 nucleotides to about 100 nucleotides long. In some embodiments, probes (e.g., detectable probes) complementary to an extended capture probe can be about 10 nucleotides to about 90 nucleotides long. In some embodiments, probes (e.g., detectable probes) complementary to an extended capture probe can be about 20 nucleotides to about 80 nucleotides long. In some embodiments, probes (e.g., detectable probes) complementary to an extended capture probe can be about 30 nucleotides to about 60 nucleotides long. In some embodiments, probes (e.g., detectable probes) complementary to an extended capture probe can be about 40 nucleotides to about 50 nucleotides long. In some embodiments, probes (e.g., detectable probes) complementary to an extended capture probe can be about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, about 50, about 51, about 52, about 53, about 54, about 55, about 56, about 57, about 58, about 59, about 60, about 61, about 62, about 63, about 64, about 65, about 66, about 67, about 68, about 69, about 70, about 71, about 72, about 73, about 74, about 75, about 76, about 77, about 78, about 79, about 80, about 81, about 82, about 83, about 84, about 85, about 86, about 87, about 88, about 89, about 90, about 91, about 92, about 93, about 94, about 95, about 96, about 97, about 98, and about 99 nucleotides long.

[0951] In some embodiments, about 1 to about 100 probes can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe. In some embodiments, about 1 to about 10 probes can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe. In some embodiments, about 10 to about 100 probes can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe. In some embodiments, about 20 to about 90 probes can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe. In some embodiments, about 30 to about 80 probes (e.g., detectable probes) can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe. In some embodiments, about 40 to about 70 probes can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe. In some embodiments, about 50 to about 60 probes can be contacted to the substrate and specifically bind

(e.g., hybridize) to an extended capture probe. In some embodiments, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, about 50, about 51, about 52, about 53, about 54, about 55, about 56, about 57, about 58, about 59, about 60, about 61, about 62, about 63, about 64, about 65, about 66, about 67, about 68, about 69, about 70, about 71, about 72, about 73, about 74, about 75, about 76, about 77, about 78, about 79, about 80, about 81, about 82, about 83, about 84, about 85, about 86, about 87, about 88, about 89, about 90, about 91, about 92, about 93, about 94, about 95, about 96, about 97, about 98, and about 99 probes can be contacted to the substrate and specifically bind (e.g., hybridize) to an extended capture probe.

[0952] In some embodiments, the probes can be complementary to a single analyte (e.g., a single gene). In some embodiments, the probes can be complementary to one or more analytes (e.g., analytes in a family of genes). In some embodiments, the probes (e.g., detectable probes) can be for a panel of genes associated with a disease (e.g., cancer, Alzheimer's disease, Parkinson's disease).

(iii) Cleavage Domain

[0953] Capture probes can optionally include a “cleavage domain,” where one or more segments or regions of the capture probe (e.g., spatial barcodes and/or UMIs) can be releasably, cleavably, or reversibly attached to a feature, or some other substrate, so that spatial barcodes and/or UMIs can be released or be releasable through cleavage of a linkage between the capture probe and the feature, or released through degradation of the underlying substrate or chemical substrate, allowing the spatial barcode(s) and/or UMI(s) of the cleaved capture probe to be accessed or be accessible by other reagents, or both. Non-limiting aspects of cleavage domains are described herein (e.g., in Section II(b)(ii)).

[0954] In some embodiments, the capture probe is linked, (e.g., via a disulfide bond), to a feature. In some embodiments, the capture probe is linked to a feature via a propylene group (e.g., Spacer C3). A reducing agent can be added to break the various disulfide bonds, resulting in release of the capture probe including the spatial barcode sequence. In another example, heating can also result in degradation and release of the attached capture probe. In some embodiments, the heating is done by laser (e.g., laser ablation) and features at specific locations can be degraded. In addition to thermally cleavable bonds, disulfide bonds, photo-sensitive bonds, and UV sensitive bonds, other non-limiting examples of labile bonds that can be coupled to a capture probe (e.g., spatial barcode) include an ester linkage (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)).

[0955] In some embodiments, the cleavage domain includes a poly(U) sequence which can be cleaved by a mixture of Uracil DNA glycosylase (UDG) and the DNA glycosylase-lyase Endonuclease VIII, commercially known as the USER™ enzyme. In some embodiments, the cleavage domain can be a single U. In some embodiments, the cleavage domain can be an abasic site that can be cleaved with an abasic site-specific endonuclease (e.g., Endonuclease IV or Endonuclease VIII).

[0956] In some embodiments, the cleavage domain of the capture probe is a nucleotide sequence within the capture probe that is cleaved specifically, e.g., physically by light or heat, chemically or enzymatically. The location of the cleavage domain within the capture probe will depend on whether or not the capture probe is immobilized on the substrate such that it has a free 3' end capable of functioning as an extension primer (e.g., by its 5' or 3' end). For example, if the capture probe is immobilized by its 5' end, the cleavage domain will be located 5' to the spatial barcode

and/or UMI, and cleavage of said domain results in the release of part of the capture probe including the spatial barcode and/or UMI and the sequence 3' to the spatial barcode, and optionally part of the cleavage domain, from a feature. Alternatively, if the capture probe is immobilized by its 3' end, the cleavage domain will be located 3' to the capture domain (and spatial barcode) and cleavage of said domain results in the release of part of the capture probe including the spatial barcode and the sequence 3' to the spatial barcode from a feature. In some embodiments, cleavage results in partial removal of the cleavage domain. In some embodiments, cleavage results in complete removal of the cleavage domain, particularly when the capture probes are immobilized via their 3' end as the presence of a part of the cleavage domain can interfere with the hybridization of the capture domain and the target nucleic acid and/or its subsequent extension.

(iv) Sequencing

[0957] After analytes from the sample have hybridized or otherwise been associated with capture probes, analyte capture agents, or other barcoded oligonucleotide sequences according to any of the methods described above in connection with the general spatial cell-based analytical methodology, the barcoded constructs that result from hybridization/association are analyzed via sequencing to identify the analytes.

[0958] In some embodiments, where a sample is barcoded directly via hybridization with capture probes or analyte capture agents hybridized, bound, or associated with either the cell surface, or introduced into the cell, as described above, sequencing can be performed on the intact sample. Alternatively, if the barcoded sample has been separated into fragments, cell groups, or individual cells, as described above, sequencing can be performed on individual fragments, cell groups, or cells. For analytes that have been barcoded via partitioning with beads, as described above, individual analytes (e.g., cells, or cellular contents following lysis of cells) can be extracted from the partitions by breaking the partitions, and then analyzed by sequencing to identify the analytes.

[0959] A wide variety of different sequencing methods can be used to analyze barcoded analyte constructs. In general, sequenced polynucleotides can be, for example, nucleic acid molecules such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including variants or derivatives thereof (e.g., single stranded DNA or DNA/RNA hybrids, and nucleic acid molecules with a nucleotide analog).

[0960] Sequencing of polynucleotides can be performed by various commercial systems. More generally, sequencing can be performed using nucleic acid amplification, polymerase chain reaction (PCR) (e.g., digital PCR and droplet digital PCR (ddPCR), quantitative PCR, real time PCR, multiplex PCR, PCR-based singleplex methods, emulsion PCR), and/or isothermal amplification.

[0961] Other examples of methods for sequencing genetic material include, but are not limited to, DNA hybridization methods (e.g., Southern blotting), restriction enzyme digestion methods, Sanger sequencing methods, next-generation sequencing methods (e.g., single-molecule real-time sequencing, nanopore sequencing, and Polony sequencing), ligation methods, and microarray methods. Additional examples of sequencing methods that can be used include targeted sequencing, single molecule real-time sequencing, exon sequencing, electron microscopy-based sequencing, panel sequencing, transistor-mediated sequencing, direct sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, co-amplification at lower denaturation temperature-PCR (COLD-PCR), sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, MS-PET sequencing, and any combinations thereof.

[0962] Sequence analysis of the nucleic acid molecules (including barcoded nucleic acid molecules

or derivatives thereof) can be direct or indirect. Thus, the sequence analysis substrate (which can be viewed as the molecule which is subjected to the sequence analysis step or process) can be the barcoded nucleic acid molecule or it can be a molecule which is derived therefrom (e.g., a complement thereof). Thus, for example, in the sequence analysis step of a sequencing reaction, the sequencing template can be the barcoded nucleic acid molecule or it can be a molecule derived therefrom. For example, a first and/or second strand DNA molecule can be directly subjected to sequence analysis (e.g., sequencing), i.e., can directly take part in the sequence analysis reaction or process (e.g., the sequencing reaction or sequencing process, or be the molecule which is sequenced or otherwise identified). Alternatively, the barcoded nucleic acid molecule can be subjected to a step of second strand synthesis or amplification before sequence analysis (e.g., sequencing or identification by another technique). The sequence analysis substrate (e.g., template) can thus be an amplicon or a second strand of a barcoded nucleic acid molecule.

[0963] In some embodiments, both strands of a double stranded molecule can be subjected to sequence analysis (e.g., sequenced). In some embodiments, single stranded molecules (e.g., barcoded nucleic acid molecules) can be analyzed (e.g., sequenced). To perform single molecule sequencing, the nucleic acid strand can be modified at the 3' end.

[0964] Massively parallel pyrosequencing techniques can be used for sequencing nucleic acids. In pyrosequencing, the nucleic acid is amplified inside water droplets in an oil solution (emulsion PCR), with each droplet containing a single nucleic acid template attached to a single primer-coated bead that then forms a clonal colony. The sequencing system contains many picolitre-volume wells each containing a single bead and sequencing enzymes. Pyrosequencing uses luciferase to generate light for detection of the individual nucleotides added to the nascent nucleic acid and the combined data are used to generate sequence reads.

[0965] As another example application of pyrosequencing, released PPi can be detected by being immediately converted to adenosine triphosphate (ATP) by ATP sulfurylase, and the level of ATP generated can be detected via luciferase-produced photons, such as described in Ronaghi, et al., *Anal. Biochem.* 242 (1), 84-9 (1996); Ronaghi, *Genome Res.* 11 (1), 3-11 (2001); Ronaghi et al. *Science* 281 (5375), 363 (1998); and U.S. Pat. Nos. 6,210,891, 6,258,568, and 6,274,320, the entire contents of each of which are incorporated herein by reference.

[0966] In some embodiments, a massively parallel sequencing technique can be based on reversible dye-terminators. As an example, DNA molecules are first attached to primers on, e.g., a glass or silicon substrate, and amplified so that local clonal colonies are formed (bridge amplification). Four types of ddNTPs are added, and non-incorporated nucleotides are washed away. Unlike pyrosequencing, the DNA is only extended one nucleotide at a time due to a blocking group (e.g., 3' blocking group present on the sugar moiety of the ddNTP). A detector acquires images of the fluorescently labelled nucleotides, and then the dye along with the terminal 3' blocking group is chemically removed from the DNA, as a precursor to a subsequent cycle. This process can be repeated until the required sequence data is obtained.

[0967] In some embodiments, sequencing is performed by detection of hydrogen ions that are released during the polymerization of DNA. A microwell containing a template DNA strand to be sequenced can be flooded with a single type of nucleotide. If the introduced nucleotide is complementary to the leading template nucleotide, it is incorporated into the growing complementary strand. This causes the release of a hydrogen ion that triggers a hypersensitive ion sensor, which indicates that a reaction has occurred. If homopolymer repeats are present in the template sequence, multiple nucleotides will be incorporated in a single cycle. This leads to a corresponding number of released hydrogen ions and a proportionally higher electronic signal.

[0968] In some embodiments, sequencing can be performed in situ. In situ sequencing methods are particularly useful, for example, when the biological sample remains intact after analytes on the sample surface (e.g., cell surface analytes) or within the sample (e.g., intracellular analytes) have been barcoded. In situ sequencing typically involves incorporation of a labeled nucleotide (e.g.,

fluorescently labeled mononucleotides or dinucleotides) in a sequential, template-dependent manner or hybridization of a labeled primer (e.g., a labeled random hexamer) to a nucleic acid template such that the identities (i.e., nucleotide sequence) of the incorporated nucleotides or labeled primer extension products can be determined, and consequently, the nucleotide sequence of the corresponding template nucleic acid. Aspects of in situ sequencing are described, for example, in Mitra et al., (2003) *Anal. Biochem.* 320, 55-65, and Lee et al., (2014) *Science*, 343 (6177), 1360-1363, the entire contents of each of which are incorporated herein by reference.

[0969] In addition, examples of methods and systems for performing in situ sequencing are described in PCT Patent Application Publication Nos. WO2014/163886, WO2018/045181, WO2018/045186, and in U.S. Pat. Nos. 10,138,509 and 10,179,932, the entire contents of each of which are incorporated herein by reference. Exemplary techniques for in situ sequencing include, but are not limited to, STARmap (described for example in Wang et al., (2018) *Science*, 361 (6499) 5691), MERFISH (described for example in Moffitt, (2016) *Methods in Enzymology*, 572, 1-49), and FISSEQ (described for example in U.S. Patent Application Publication No. 2019/0032121). The entire contents of each of the foregoing references are incorporated herein by reference.

[0970] For analytes that have been barcoded via partitioning, barcoded nucleic acid molecules or derivatives thereof (e.g., barcoded nucleic acid molecules to which one or more functional sequences have been added, or from which one or more features have been removed) can be pooled and processed together for subsequent analysis such as sequencing on high throughput sequencers. Processing with pooling can be implemented using barcode sequences. For example, barcoded nucleic acid molecules of a given partition can have the same barcode, which is different from barcodes of other spatial partitions. Alternatively, barcoded nucleic acid molecules of different partitions can be processed separately for subsequent analysis (e.g., sequencing).

[0971] In some embodiments, where capture probes do not contain a spatial barcode, the spatial barcode can be added after the capture probe captures analytes from a biological sample and before analysis of the analytes. When a spatial barcode is added after an analyte is captured, the barcode can be added after amplification of the analyte (e.g., reverse transcription and polymerase amplification of RNA). In some embodiments, analyte analysis uses direct sequencing of one or more captured analytes, such as direct sequencing of hybridized RNA. In some embodiments, direct sequencing is performed after reverse transcription of hybridized RNA. In some embodiments direct sequencing is performed after amplification of reverse transcription of hybridized RNA.

[0972] In some embodiments, direct sequencing of captured RNA is performed by sequencing-by-synthesis (SBS). In some embodiments, a sequencing primer is complementary to a sequence in one or more of the domains of a capture probe (e.g., functional domain). In such embodiments, sequencing-by-synthesis can include reverse transcription and/or amplification in order to generate a template sequence (e.g., functional domain) from which a primer sequence can bind.

[0973] SBS can involve hybridizing an appropriate primer, sometimes referred to as a sequencing primer, with the nucleic acid template to be sequenced, extending the primer, and detecting the nucleotides used to extend the primer. Preferably, the nucleic acid used to extend the primer is detected before a further nucleotide is added to the growing nucleic acid chain, thus allowing base-by-base in situ nucleic acid sequencing. The detection of incorporated nucleotides is facilitated by including one or more labelled nucleotides in the primer extension reaction. To allow the hybridization of an appropriate sequencing primer to the nucleic acid template to be sequenced, the nucleic acid template should normally be in a single stranded form. If the nucleic acid templates making up the nucleic acid features are present in a double stranded form these can be processed to provide single stranded nucleic acid templates using methods well known in the art, for example by denaturation, cleavage, etc. The sequencing primers which are hybridized to the nucleic acid template and used for primer extension are preferably short oligonucleotides, for example, 15 to 25 nucleotides in length. The sequencing primers can be provided in solution or in an immobilized

form. Once the sequencing primer has been annealed to the nucleic acid template to be sequenced by subjecting the nucleic acid template and sequencing primer to appropriate conditions, primer extension is carried out, for example using a nucleic acid polymerase and a supply of nucleotides, at least some of which are provided in a labelled form, and conditions suitable for primer extension if a suitable nucleotide is provided.

[0974] Preferably after each primer extension step, a washing step is included in order to remove unincorporated nucleotides which can interfere with subsequent steps. Once the primer extension step has been carried out, the nucleic acid colony is monitored to determine whether a labelled nucleotide has been incorporated into an extended primer. The primer extension step can then be repeated to determine the next and subsequent nucleotides incorporated into an extended primer. If the sequence being determined is unknown, the nucleotides applied to a given colony are usually applied in a chosen order which is then repeated throughout the analysis, for example dATP, dTTP, dCTP, dGTP.

[0975] SBS techniques which can be used are described for example, but not limited to, those in U.S. Patent App. Pub. No. 2007/0166705, U.S. Patent App. Pub. No. 2006/0188901, U.S. Pat. No. 7,057,026, U.S. Patent App. Pub. No. 2006/0240439, U.S. Patent App. Pub. No. 2006/0281109, PCT Patent App. Pub. No. WO 05/065814, U.S. Patent App. Pub. No. 2005/0100900, PCT Patent App. Pub. No. WO 06/064199, PCT Patent App. Pub. No. WO07/010,251, U.S. Patent App. Pub. No. 2012/0270305, U.S. Patent App. Pub. No. 2013/0260372, and U.S. Patent App. Pub. No. 2013/0079232, the entire contents of each of which are incorporated herein by reference.

[0976] In some embodiments, direct sequencing of captured RNA is performed by sequential fluorescence hybridization (e.g., sequencing by hybridization). In some embodiments, a hybridization reaction where RNA is hybridized to a capture probe is performed in situ. In some embodiments, captured RNA is not amplified prior to hybridization with a sequencing probe. In some embodiments, RNA is amplified prior to hybridization with sequencing probes (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, amplification is performed using single-molecule hybridization chain reaction. In some embodiments, amplification is performed using rolling chain amplification.

[0977] Sequential fluorescence hybridization can involve sequential hybridization of probes including degenerate primer sequences and a detectable label. A degenerate primer sequence is a short oligonucleotide sequence which is capable of hybridizing to any nucleic acid fragment independent of the sequence of said nucleic acid fragment. For example, such a method could include the steps of: (a) providing a mixture including four probes, each of which includes either A, C, G, or T at the 5'-terminus, further including degenerate nucleotide sequence of 5 to 11 nucleotides in length, and further including a functional domain (e.g., fluorescent molecule) that is distinct for probes with A, C, G, or T at the 5'-terminus; (b) associating the probes of step (a) to the target polynucleotide sequences, whose sequence needs will be determined by this method; (c) measuring the activities of the four functional domains and recording the relative spatial location of the activities; (d) removing the reagents from steps (a)-(b) from the target polynucleotide sequences; and repeating steps (a)-(d) for n cycles, until the nucleotide sequence of the spatial domain for each bead is determined, with modification that the oligonucleotides used in step (a) are complementary to part of the target polynucleotide sequences and the positions 1 through n flanking the part of the sequences. Because the barcode sequences are different, in some embodiments, these additional flanking sequences are degenerate sequences. The fluorescent signal from each spot on the array for cycles 1 through n can be used to determine the sequence of the target polynucleotide sequences.

[0978] In some embodiments, direct sequencing of captured RNA using sequential fluorescence hybridization is performed in vitro. In some embodiments, captured RNA is amplified prior to hybridization with a sequencing probe (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, a capture probe containing captured RNA is exposed to the

sequencing probe targeting coding regions of RNA. In some embodiments, one or more sequencing probes are targeted to each coding region. In some embodiments, the sequencing probe is designed to hybridize with sequencing reagents (e.g., a dye-labeled readout oligonucleotides). A sequencing probe can then hybridize with sequencing reagents. In some embodiments, output from the sequencing reaction is imaged. In some embodiments, a specific sequence of cDNA is resolved from an image of a sequencing reaction. In some embodiments, reverse transcription of captured RNA is performed prior to hybridization to the sequencing probe. In some embodiments, the sequencing probe is designed to target complementary sequences of the coding regions of RNA (e.g., targeting cDNA).

[0979] In some embodiments, a captured RNA is directly sequenced using a nanopore-based method. In some embodiments, direct sequencing is performed using nanopore direct RNA sequencing in which captured RNA is translocated through a nanopore. A nanopore current can be recorded and converted into a base sequence. In some embodiments, captured RNA remains attached to a substrate during nanopore sequencing. In some embodiments, captured RNA is released from the substrate prior to nanopore sequencing. In some embodiments, where the analyte of interest is a protein, direct sequencing of the protein can be performed using nanopore-based methods. Examples of nanopore-based sequencing methods that can be used are described in Deamer et al., *Trends Biotechnol.* 18, 14 7-151 (2000); Deamer et al., *Acc. Chem. Res.* 35:817-825 (2002); Li et al., *Nat. Mater.* 2:611-615 (2003); Soni et al., *Clin. Chem.* 53, 1996-2001 (2007); Healy et al., *Nanomed.* 2, 459-481 (2007); Cockroft et al., *J. Am. Chem. Soc.* 130, 818-820 (2008); and in U.S. Pat. No. 7,001,792. The entire contents of each of the foregoing references are incorporated herein by reference.

[0980] In some embodiments, direct sequencing of captured RNA is performed using single molecule sequencing by ligation. Such techniques utilize DNA ligase to incorporate oligonucleotides and identify the incorporation of such oligonucleotides. The oligonucleotides typically have different labels that are correlated with the identity of a particular nucleotide in a sequence to which the oligonucleotides hybridize. Aspects and features involved in sequencing by ligation are described, for example, in Shendure et al. *Science* (2005), 309:1728-1732, and in U.S. Pat. Nos. 5,599,675; 5,750,341; 6,969,488; 6,172,218; and 6,306,597, the entire contents of each of which are incorporated herein by reference.

[0981] In some embodiments, nucleic acid hybridization can be used for sequencing. These methods utilize labeled nucleic acid decoder probes that are complementary to at least a portion of a barcode sequence. Multiplex decoding can be performed with pools of many different probes with distinguishable labels. Non-limiting examples of nucleic acid hybridization sequencing are described for example in U.S. Pat. No. 8,460,865, and in Gunderson et al., *Genome Research* 14:870-877 (2004), the entire contents of each of which are incorporated herein by reference.

[0982] In some embodiments, commercial high-throughput digital sequencing techniques can be used to analyze barcode sequences, in which DNA templates are prepared for sequencing not one at a time, but in a bulk process, and where many sequences are read out preferably in parallel, or alternatively using an ultra-high throughput serial process that itself may be parallelized. Examples of such techniques include Illumina® sequencing (e.g., flow cell-based sequencing by synthesis techniques), using modified nucleotides (such as commercialized in HiSeq™ and additional sequencing technology instruments by Illumina, Inc., San Diego, CA), HeliScope™ by Helicos Biosciences Corporation, Cambridge, MA, and PacBio RS by Pacific Biosciences of California, Inc., Menlo Park, CA), sequencing by ion detection technologies (Ion Torrent, Inc., South San Francisco, CA), and sequencing of DNA nanoballs (Complete Genomics, Inc., Mountain View, CA).

[0983] In some embodiments, detection of a proton released upon incorporation of a nucleotide into an extension product can be used in the methods described herein. For example, the sequencing methods and systems described in U.S. Patent Application Publication Nos.

2009/0026082, 2009/0127589, 2010/0137143, and 2010/0282617, can be used to directly sequence barcodes.

[0984] In some embodiments, real-time monitoring of DNA polymerase activity can be used during sequencing. For example, nucleotide incorporations can be detected through fluorescence resonance energy transfer (FRET), as described for example in Levene et al., *Science* (2003), 299, 682-686, Lundquist et al., *Opt. Lett.* (2008), 33, 1026-1028, and Korlach et al., *Proc. Natl. Acad. Sci. USA* (2008), 105, 1176-1181. The entire contents of each of the foregoing references are incorporated herein by reference herein.

(v) Temporal Analysis

[0985] In some embodiments, the methods described herein can be used to assess analyte levels and/or expression in a cell or a biological sample over time (e.g., before or after treatment with an agent or different stages of differentiation). In some examples, the methods described herein can be performed on multiple similar biological samples or cells obtained from the subject at a different time points (e.g., before or after treatment with an agent, different stages of differentiation, different stages of disease progression, different ages of the subject, before or after physical perturbation, before or after treatment with a perturbation agent as described herein, or before or after development of resistance to an agent). As described herein, a “perturbation agent” or “perturbation reagent” can be a small molecule, an antibody, a drug, an aptamer, a nucleic acid (e.g., miRNA), a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, antisense oligonucleotide, a physical environmental (e.g., temperature change), and/or any other known perturbation agents where the agent alters equilibrium or homeostasis.

[0986] In some embodiments, the methods described herein can be performed on multiple similar biological samples or cells obtained from the subject at 2, 3, 4, 5, 6, 7, 8, 9, 10, or more times. For example, the multiple similar biological samples can be repetitive samples from the same subject, the same tissue, the same organoid, the same cell suspension, or any other biological sample described herein. In some embodiments, the methods described herein can be performed on the same biological sample or cells obtained from the subject at a different time points (e.g., before or after treatment with a perturbation agent, different stages of differentiation, different stages of disease progression, different ages of the subject, or before or after development of resistance to an agent). In some embodiments, a perturbation agent can be small-molecules, antibodies, nucleic acids, peptides, and/or other external stimuli (e.g., temperature change). In some embodiments, the biological sample is contacted with a different array at each time point.

[0987] In some embodiments, a sample can be placed in a controlled environment permissive for cellular growth and/or maintenance, and/or to prevent hypoxia. In some embodiments, a controlled environment allows a sample to be analyzed at different time points. Barcoded arrays can be placed proximal to (e.g., on top of) the sample and imaged using a microscope or other suitable instrument to register the relative position of the biological sample to the barcoded array, optionally using optically encoded fiducial markers. An electric field can be applied for a period of time, such that biological analytes (e.g., DNA, RNA, proteins, metabolites, small molecules, lipids, and the like) are released from the sample and captured by capture probes on the spatially-barcoded array, preserving spatial information of the sample. The barcoded array can be removed, and the spatial and molecular information therein is determined (e.g., by performing library construction for next generation sequencing or in situ sequencing). Sequencing can be followed by computational analysis to correlate the molecular information (e.g., gene expression values with the spatial barcode). These steps can be repeated one or more times to capture the spatial information of analytes at different time-points.

[0988] In some embodiments, methods as described herein can be combined with a cell migration assay. A cell migration assay can comprise one or more microprinted lines, or suspended 3D nanofibers, on which the cells migrate. Migration using these assays can be measured by imaging cell migration and/or contacting migrated cells with a spatially-barcoded array. An array used in a

cell migration assay can comprise one or more channels on the substrate of the array, e.g., to confine cell migration to one dimension along the substrate. Additionally, the channels can direct the migration of a cell such that it does not contact another cell on the array (e.g., the channels do not overlap with each other), and in some embodiments, the channels are about the same width as or wider than a cell (e.g., for a mammalian cell, a channel can have a width of about 2 μm to about 10 μm). Cellular location on the spatially-barcoded array can be identified using any method described herein.

[0989] In some embodiments, cells can be disposed on an array as described herein and allowed to migrate. Cell migration in cell migration assays can be used to measure target phenotypes (e.g., phenotype for invasiveness). In some embodiments, the cell migration distance can be measured and correlated to a biological analyte. Reagents can be added to the array to facilitate cell migration. For example, the array can be coated with one or more extracellular matrix (ECM) components (e.g., basement membrane extract (BME), laminin I, collagen I, collagen IV, fibronectin, vitronectin, elastin), a cell culture medium, a chemoattractant, a chemorepellant, or a combination thereof. In some embodiments, a reagent such as a chemoattractant or chemorepellant can be disposed on only a portion of the array, present as a gradient along the one or more axis or channels of the array, or a combination thereof.

(vi) Spatially Resolving Analyte Information

[0990] In some embodiments, a lookup table (LUT) can be used to associate one property with another property of a feature. These properties include, e.g., locations, barcodes (e.g., nucleic acid barcode molecules), spatial barcodes, optical labels, molecular tags, and other properties.

[0991] In some embodiments, a lookup table can associate a nucleic acid barcode molecule with a feature. In some embodiments, an optical label of a feature can permit associating the feature with a biological particle (e.g., cell or nuclei). The association of a feature with a biological particle can further permit associating a nucleic acid sequence of a nucleic acid molecule of the biological particle to one or more physical properties of the biological particle (e.g., a type of a cell or a location of the cell). For example, based on the relationship between the barcode and the optical label, the optical label can be used to determine the location of a feature, thus associating the location of the feature with the barcode sequence of the feature. Subsequent analysis (e.g., sequencing) can associate the barcode sequence and the analyte from the sample. Accordingly, based on the relationship between the location and the barcode sequence, the location of the biological analyte can be determined (e.g., in a specific type of cell or in a cell at a specific location of the biological sample).

[0992] In some embodiments, a feature can have a plurality of nucleic acid barcode molecules attached thereto. The plurality of nucleic acid barcode molecules can include barcode sequences. The plurality of nucleic acid molecules attached to a given feature can have the same barcode sequences, or two or more different barcode sequences. Different barcode sequences can be used to provide improved spatial location accuracy.

[0993] As discussed above, analytes obtained from a sample, such as RNA, DNA, peptides, lipids, and proteins, can be further processed. In particular, the contents of individual cells from the sample can be provided with unique spatial barcode sequences such that, upon characterization of the analytes, the analytes can be attributed as having been derived from the same cell. More generally, spatial barcodes can be used to attribute analytes to corresponding spatial locations in the sample. For example, hierarchical spatial positioning of multiple pluralities of spatial barcodes can be used to identify and characterize analytes over a particular spatial region of the sample. In some embodiments, the spatial region corresponds to a particular spatial region of interest previously identified, e.g., a particular structure of cytoarchitecture previously identified. In some embodiments, the spatial region corresponds to a small structure or group of cells that cannot be seen with the naked eye. In some embodiments, a unique molecular identifier can be used to identify and characterize analytes at a single cell level.

[0994] The analyte can include a nucleic acid molecule, which can be barcoded with a barcode sequence of a nucleic acid barcode molecule. In some embodiments, the barcoded analyte can be sequenced to obtain a nucleic acid sequence. In some embodiments, the nucleic acid sequence can include genetic information associated with the sample. The nucleic acid sequence can include the barcode sequence, or a complement thereof. The barcode sequence, or a complement thereof, of the nucleic acid sequence can be electronically associated with the property (e.g., color and/or intensity) of the analyte using the LUT to identify the associated feature in an array.

(vii) Proximity Capture

[0995] In some embodiments, two- or three-dimensional spatial profiling of one or more analytes present in a biological sample can be performed using a proximity capture reaction, which is a reaction that detects two analytes that are spatially close to each other and/or interacting with each other. For example, a proximity capture reaction can be used to detect sequences of DNA that are close in space to each other, e.g., the DNA sequences can be within the same chromosome, but separated by about 700 bp or less. As another example, a proximity capture reaction can be used to detect protein associations, e.g., two proteins that interact with each other. A proximity capture reaction can be performed in situ to detect two analytes that are spatially close to each other and/or interacting with each other inside a cell. Non-limiting examples of proximity capture reactions include DNA nanoscopy, DNA microscopy, and chromosome conformation capture methods. Chromosome conformation capture (3C) and derivative experimental procedures can be used to estimate the spatial proximity between different genomic elements. Non-limiting examples of chromatin capture methods include chromosome conformation capture (3-C), conformation capture-on-chip (4-C), 5-C, ChIA-PET, Hi-C, targeted chromatin capture (T2C). Examples of such methods are described, for example, in Miele et al., *Methods Mol Biol.* (2009), 464, Simonis et al., *Nat. Genet.* (2006), 38 (11): 1348-54, Raab et al., *Embo. J.* (2012), 31 (2): 330-350, and Eagen et al., *Trends Biochem. Sci.* (2018) 43 (6): 469-478, the entire contents of each of which is incorporated herein by reference.

[0996] In some embodiments, the proximity capture reaction includes proximity ligation. In some embodiments, proximity ligation can include using antibodies with attached DNA strands that can participate in ligation, replication, and sequence decoding reactions. For example, a proximity ligation reaction can include oligonucleotides attached to pairs of antibodies that can be joined by ligation if the antibodies have been brought in proximity to each oligonucleotide, e.g., by binding the same target protein (complex), and the DNA ligation products that form are then used to template PCR amplification, as described for example in Söderberg et al., *Methods.* (2008), 45 (3): 227-32, the entire contents of which are incorporated herein by reference. In some embodiments, proximity ligation can include chromosome conformation capture methods.

In some embodiments, the proximity capture reaction is performed on analytes within about 400 nm distance (e.g., about 300 nm, about 200 nm, about 150 nm, about 100 nm, about 50 nm, about 25 nm, about 10 nm, or about 5 nm) from each other. In general, proximity capture reactions can be reversible or irreversible.

(viii) Feature Removal from an Array

[0997] A spatially-barcoded array can be contacted with a biological sample to spatially detect analytes present in the biological sample. In some embodiments, the features (e.g., gel pads, beads) can be removed from the substrate surface for additional analysis (e.g., imaging, sequencing, or quantification). For example, features on a substrate delivered by a droplet manipulation system as described herein can be removed from the substrate surface. In some embodiments, the features (e.g., gel pads, beads) can be removed mechanically (e.g., scraping), by an enzymatic reaction, or by a chemical reaction. In some embodiments, the features (e.g., gel pads, beads) can be aspirated. In some embodiments, after the features are removed (by any method), the features can be combined with a uniquely barcoded bead. In some embodiments, the oligonucleotides within a feature can be ligated or hybridized to the barcode sequence on the barcoded bead. For example,

the spatial barcode oligonucleotide within a feature can be ligated to the barcode sequence on the barcoded bead. Additionally, the capture probes can be ligated to the barcode sequence on the barcoded bead. In some embodiments, the features and the bead can be partitioned. In some embodiments, the features (e.g., gels pads, beads) and the uniquely barcoded bead can be partitioned into a vesicle. In some embodiments, the vesicle can have a lipid bilayer. In some embodiments, the features and the bead can be encapsulated. In some embodiments, the features and the bead can be encapsulated in an oil emulsion. In some embodiments, the features and the bead can be encapsulated in a water-in-oil emulsion. Once partitioned, the features (e.g., gel pads, beads) can be processed for further analysis (e.g., quantitation, amplification, or sequencing) according to any method described herein.

(ix) Other Applications

[0998] The spatial analysis methods described herein can be used to detect and characterize the spatial distribution of one or more haplotypes in a biological sample. As used in the present disclosure, a haplotype is used to describe one or more mutations, DNA variations, polymorphisms in a given segment of the genome, which can be used to classify the genetic segment, or a collection of alleles or genetic segments containing single nucleotide polymorphisms (SNPs). Haplotype association studies are used to inform a greater understanding of biological conditions. For example, identifying and characterizing haplotype variants at or associated with putative disease loci in humans can provide a foundation for mapping genetic causes underlying disease susceptibility. The term “locus” (plural “loci”), as used in the art, can be a fixed location on a chromosome, including the location of a gene or a genetic marker, which can contain a plurality of haplotypes, including alleles and SNPs.

[0999] Variant haplotype detection is a technique used to identify heterozygous cells in single cell studies. In combination with spatial analysis, variant haplotype detection can further provide novel information on the distribution of heterozygous cells in biological samples (e.g., tissues) affected by or exhibiting a variety of biological conditions. These data may reveal causal relationships between variant haplotypes and disease outcomes, to aid in identification of disease-associated variants, or to reveal heterogeneity within a biological sample.

[1000] In some embodiments, variant haplotype detection is a technique that can be used in combination with, in addition to, or as a part of, the spatial analysis methods described herein. Briefly, variant haplotype detection can include providing inputs for executing an algorithm on a computer system, and performing an analysis to identify and determine the spatial distribution of haplotypes. One input can be a plurality of sequence reads obtained from a two-dimensional spatial array in contact with a biological sample and subsequently aligned to a genome. The sequence reads can also contain spatial barcodes with positional information, such that the sequence reads can be mapped to a location on the biological sample. Other inputs can include electronic data files of gene sequence variations, or haplotypes, and a reference genome. For each locus, the corresponding sequence reads and variant haplotypes are aligned to determine the haplotype identity of each sequence read. The haplotype identity and the spatial barcode of the sequence reads are then categorized to determine the spatial distribution of haplotypes within the biological sample. As described above, this spatial distribution can be used to characterize a biological condition of the sample. In some embodiments, sequence reads are obtained by in situ sequencing of the two-dimensional array of positions on the substrate, while in some embodiments, sequence reads are obtained by high-throughput sequencing. In some embodiments, other methods for generating sequence reads described herein are used, such as paired end sequencings.

[1001] In some embodiments, a respective loci in the plurality of loci is bi-allelic and the corresponding set of haplotypes for the respective loci consists of a first allele and a second allele. In some such embodiments, the respective loci includes a heterozygous single nucleotide polymorphism (SNP), a heterozygous insert, or a heterozygous deletion.

[1002] In some embodiments, analytes captured by any of the spatial analysis methods described

herein can be analyzed (e.g., sequenced) via in situ sequencing methods. For example, a substrate including a plurality of capture probes (e.g., an array), attached either directly or indirectly (e.g., via a feature), that include a spatial barcode and a capture domain. In some embodiments, the capture domain can be configured to interact (e.g., hybridize) with an analyte (e.g., mRNA). In some embodiments, a biological sample can be contacted to the array such that the capture domain of the capture probe interacts with (e.g., hybridizes) the analyte. In some embodiments, the capture probe can function as a template for a hybridization or ligation reaction with the captured analyte. For example, a reverse transcription reaction can be performed to extend the 3' end of a capture probe hybridized to the analyte using any of the exemplary reverse transcriptases described herein, thereby generating an extended capture probe (e.g., an extended capture probe including the spatial barcode and a sequence that is complementary to a sequence in the analyte). After the extended capture probe is synthesized, a second strand that is complementary to the extended capture probe can be synthesized. In some embodiments, second strand synthesis can be performed using any of the methods described herein. In some embodiments, amine-modified nucleotides can be used when generating the extended capture probe or the second strand, or both. For example, the amine-modified nucleotides can be aminoallyl (aa)-dUTP, aa-dCTP, aa-dGTP, and/or aa-dATP.

[1003] In some embodiments, after generation of the extended capture probe, the second strand, or both the extended capture probe and/or the second strand can be released from the surface of the substrate. For example, the extended capture probe and/or the second strand can be released by any of the methods described herein (e.g., heat or cleavage via a cleavage domain). In some embodiments, the amine-modified nucleotides incorporated into the extended capture probe can be cross-linked to the surface of a substrate or cross-linked to the biological sample using its amine-modified nucleotides. In some embodiments, the surface of the substrate can be coated in a hydrogel. In some embodiments, the surface of the substrate can be coated in a protein matrix. In some embodiments, the cross-linking can be irreversible. In some embodiments, the cross-linked extended capture probe and/or second strand can be circularized. For example, circular template ligation can be performed by a DNA ligase (e.g., T4 DNA ligase) or circular template-free ligation can be performed by a template independent ligase (e.g., CircLigase). In some embodiments, the extended capture probe is circularized with CircLigase. In some embodiments, the circularized extended capture probe can be amplified. For example, rolling circle amplification can be performed with a suitable DNA polymerase (e.g., phi29). In some embodiments, the capture probe includes a functional domain (e.g., sequencing adapter). In some embodiments, rolling circle amplification can be performed with a primer complementary to the functional domain (e.g., sequencing adapter). In some embodiments, the rolling circle amplification can be performed to generate two or more amplicons (e.g., one or more amplicons including any of the amine-modified nucleotides described herein). In some embodiments, the two or more amplicons produced by the rolling circle amplification can be cross-linked to the surface of the substrate and/or cross-linked to the biological sample. In some embodiments, the two or more amplicons can be sequenced in situ. The in situ sequencing can be performed by any method described herein (See, Lee, J. H., Fluorescent in situ sequencing (FISSEQ) of RNA for gene expression profiling, *Nat Protoc.*, 10 (3): 442-458, doi: 10.1038/nprot.2014.191 (2015), which is incorporated herein by reference). In some embodiments, the two or more amplicons can be imaged.

[1004] In some embodiments, spatial analysis by any of the methods described herein can be performed on ribosomal RNA (rRNA), including, endogenous ribosomal RNA (e.g., native to the biological sample), and/or exogenous RNA (e.g., microbial ribosomal RNA and/or viral RNA also present in the biological sample). As used herein, "metagenomics," can refer to the study of exogenous nucleic acids (e.g., DNA, RNA, or other nucleic acids described herein) present in a biological sample. As used herein, "spatial metagenomics," can refer to the study of the spatial location of exogenous nucleic acid present in a biological sample. Spatial metagenomics can also refer to the identification of one or more species (e.g., viral or microbial) present in the biological

sample and/or the study of identifying patterns of proximity (e.g., co-localization) amongst species. [1005] In some embodiments, microbial rRNA can be spatially detected, quantified, and/or amplified from a biological sample. In some embodiments, rRNA (e.g., 16S ribosomal RNA) can be associated with a particular microbial species. For example, microbial ribosomal RNA (e.g., 16S ribosomal RNA) can be used to identify one or more species of microbe present in the biological sample (See e.g., Kolbert, C. P., and Persing, D. H., Ribosomal DNA sequencing as a tool for identification of bacterial pathogens, *Current Opinion in Microbiology*. 2 (3): 299-305. doi: 10.1016/S1369-5274 (99) 80052-6. PMID 10383862 (1999), which is incorporated herein by reference). In some embodiments, identification of microbial species in proximity to one or more other microbial species can be identified.

[1006] In some embodiments, a biological sample be covered (e.g., coated) or embedded in with a photo-crosslinkable coating (e.g., conditionally dissolvable polymer, e.g., DTT sensitive hydrogel). A biological sample can be contacted with the photo-crosslinkable coated substrate. In some embodiments, the biological sample and photo-crosslinkable substrate are assembled into a flow-cell and the photo-crosslinkable polymer can be incubated with the biological sample. The biological sample can be cross-linked into hydrogel-voxels of defined dimensions using a light source and a photomask. In some embodiments, the flow-cell can be dismantled and washed to remove unpolymerized hydrogel. The photo-crosslinkable coating can be treated with DTT to yield single-cell portions or approximately single-cell portions.

[1007] In some embodiments, the single-cell or approximately single-cell portions can be encapsulated in a vesicle. The vesicle can contain a barcoded feature (e.g., a bead), and the barcoded feature can contain a capture domain. In some embodiments, the capture domain can bind specifically to microbial rRNA (e.g., microbial 16S rRNA). In some embodiments, the captured microbial rRNA can be amplified and analyzed (e.g., sequenced) by any of the methods described herein. In some embodiments, the amplified and sequenced microbial rRNA can identify microbial species and/or patterns of proximity (e.g., co-localization) of one or more species.

[1008] Alternatively, spatial analysis can be performed on exogenous rRNA (e.g., microbial or viral) with a plurality of capture probes on a substrate (e.g., an array), wherein the capture probes include a spatial barcode and a capture domain. In some embodiments, the capture domain can be configured to interact (e.g., hybridize) with microbial rRNA present in the biological sample. The capture probe can be configured to interact with any microbial rRNA. In some embodiments, the capture probe is configured to interact with microbial 16S rRNA. The biological sample can be treated (e.g., permeabilized) such that the capture domain and the analyte (e.g., microbial rRNA) interact (e.g., hybridize). In some embodiments, the captured analyte (e.g., microbial rRNA) can be reverse transcribed generating an extended capture probe, followed generation of a second strand that is complementary to the extended capture probe as described herein. The extended capture probe and/or the second strand can include a portion or all of a capture probe sequence, or a complement thereof. The capture probe sequence, or complement thereof, can include the spatial barcode, or complement thereof. In some embodiments, the first strand cDNA, and optionally, the second strand cDNA can be amplified by any method described herein. The amplified capture probes and analytes can be analyzed (e.g., sequenced) by any method described herein. The spatial information of the spatially-barcoded features can be used to determine the spatial location of the captured analytes (e.g., microbial rRNA) in the biological sample, or a portion thereof. In some embodiments, the captured analyte can identify the microbial species present in the biological sample, or a portion thereof. In some embodiments, the spatial information and identity of microbial species present in the biological sample can be correlated with one another, thus revealing whether certain microbial species may be found in proximity (e.g., co-localize) with one another in the biological sample.

[1009] In exemplary embodiments, provided herein are methods for detecting a nucleic acid within a portion of biological sample that include: (a) immobilizing the biological sample in a gel matrix

to produce an embedded biological sample; (b) breaking up the embedded biological sample into portions; (c) lysing cell(s) present in the portions; (d) encapsulating a portion from step (c) together with a bead having an attached capture probe comprising a spatial barcode and a capture domain that binds specifically to the nucleic acid in the portion; and (e) determining (i) all or a part of the sequence of the spatial barcode, or a complement thereof, and (ii) all or part of the sequence of the nucleic acid, or a complement thereof, and using the determined sequences of (i) and (ii) to detect the nucleic acid within the portion of the biological sample. In some embodiments of these methods, the nucleic acid comprises microbial ribosomal RNA (rRNA). In some embodiments of these methods, the microbial rRNA comprises 16S rRNA. In some embodiments of these methods, the method comprises detecting 16S rRNA from at least two different microbacteria within the portion of the biological sample. In some embodiments of these methods, the nucleic acid is an mRNA.

[1010] Provided herein are methods for spatially profiling analytes within a biological sample. Profiles of biological samples (e.g., individual cells, populations of cells, tissue sections, etc.) can be compared to profiles of other cells, e.g., “normal,” or “healthy,” biological samples. In some embodiments of any the methods for spatially profiling analytes described herein, the method can provide for diagnosis of a disease (e.g., cancer, Alzheimer's disease, Parkinson's disease). In some embodiments of any the methods for spatially profiling analytes described herein, the methods can be used in drug screening. In some embodiments of any the methods for spatially profiling analytes described herein, the methods can be used to perform drug screening with an organoid. In some embodiments of any the methods for spatially profiling analytes described herein, the methods can be used to detect changes in (e.g., altered) cellular signaling. In some embodiments of any the methods for spatially profiling analytes described herein, the methods can include the introduction of a pathogen to the biological sample and evaluation of the response of the biological sample to the pathogen. In some embodiments of any the methods for spatially profiling analytes described herein, the methods include exposing the biological sample to a perturbation agent (e.g., any of the perturbation agents described herein) and evaluating the response of the biological sample to the perturbation agent. In some embodiments of any the methods for spatially profiling analytes described herein, the methods include monitoring cell differentiation in a biological sample (e.g., an organoid). In some embodiments of any the methods for spatially profiling analytes described herein, the methods include analyzing tissue morphogenesis. In some embodiments of any the methods for spatially profiling analytes described herein, the methods include identifying spatial heterogeneity in a biological sample (e.g., identifying different cell types or populations in a biological sample). In some embodiments of any the methods for spatially profiling analytes described herein, the methods include analyzing the spatiotemporal order (e.g., timing) of molecular events. For example, the methods for spatially profiling analytes can include monitoring expression levels over the course of a disease.

[1011] The methods provided herein can also be used to determine a relative level of inflammation in a subject (e.g., determine an inflammatory score) or a subject's response to treatment or the development of resistance to treatment. The methods described herein can also be used to identify candidate targets for potential therapeutic intervention and/or to identify biomarkers associated with different disease states in a subject.

(h) Quality Control

(i) Control Sample

[1012] As used herein, the term “control sample” typically refers to a substrate that is insoluble in aqueous liquid and that allows for an accurate and traceable positioning of test analytes on the substrate. The term “control sample” and “test substrate” are used interchangeably herein. The control sample can be any suitable substrate known to the person skilled in the art. Exemplary control samples comprise a semi-porous material. Non-limiting examples of a semi-porous material include a nitrocellulose membrane, a hydrogel, and a nylon filter.

[1013] A control sample or test substrate can be of any appropriate dimension or volume (e.g., size or shape). In some embodiments, a control sample is a regular shape (e.g., a square, circle, or a rectangle). In some embodiments, a surface of a control sample has any appropriate form or format. For example, the surface of a control sample can be flat or curved (e.g., convexly or concavely curved towards the area where the interaction between the substrate and the control sample takes place). In some embodiments, a control sample has rounded corners (e.g., for increased safety or robustness). In some embodiments, a control sample has one or more cut-off corners (e.g., for use with a slide clamp or cross-table).

[1014] A control sample can comprise a plurality of test analytes. In some embodiments, the members of the plurality of test analytes are disposed on the substrate in a known amount and in a known location. For example, a plurality of test analytes are disposed at a known amount on the control sample at one or more locations. In some embodiments, the plurality of test analytes are disposed on the substrate in a defined pattern (e.g., an x-y grid pattern). In some embodiments, the defined pattern includes one or more locations or spots.

[1015] In some embodiments, each location comprises a plurality of the same species of test analyte. In some embodiments, each location comprises a plurality of one or more different species of test analytes. In some embodiments, each location on the control sample represents a different region of a biological sample, e.g., a tissue sample. In some embodiments, an area on the control sample that does not comprise a plurality of test analytes represents an area where no biological sample is present.

[1016] In some embodiments, the plurality of test analytes comprises one or more test analytes, e.g., a first test analyte, a second test analyte, a third test analyte, a fourth test analyte, etc. In some embodiments, the plurality of test analytes comprises nucleic acids. In some embodiments, each location or feature comprises a population of nucleic acid sequences. In some embodiments, the nucleic acid sequence of a first test analyte differs from the nucleic acid sequence of a second test analyte by a single nucleic acid residue. In some embodiments, each location or feature comprises a population of RNA transcripts and one or more specific surface marker proteins or one or more CRISPR guide RNAs. In some embodiments, the plurality of test analytes comprises a bacterial artificial chromosomes (BAC). In some embodiments, each location on the control sample comprises a unique blend of BACs. In some embodiments, proteins are cross-linked to the BACs, for example, to mimic histone binding on DNA.

[1017] In some embodiments, the concentration of a first test analyte differs from the concentration of a second test analyte at a different location or feature on the control sample. In some embodiments, the first test analyte and the second test analyte comprise an identical nucleotide sequence.

[1018] A control sample can be used to determine process bias. Barcoded arrays can be placed on top of a control sample comprising a plurality of test analytes, where members of the plurality of test analytes can be disposed on a substrate in a known amount and in a known location. The array can then be removed, and the molecular information therein can be determined by performing library construction for next generation sequencing, followed by computational analysis to correlate the expression values of the test analytes with the barcodes (e.g., spatial barcodes) on the array. The sequencing data can be compared with the known amount and the known locations of the plurality of test analytes to determine whether the spatial analysis workflow accurately detects the presence, amount, location, or combinations thereof, of the test analyte, thereby determining process bias of the spatial analysis workflow.

ii) Spatial RNA Integrity Number (sRIN)

[1019] As used herein, the term “spatial RNA Integrity Number” or “sRIN” refers to the in situ indication of RNA quality based on an integrity score. Higher sRIN scores correlate with higher data quality in the spatial profiling assays described herein. For example, a first biological sample with a high sRIN score will have higher data quality compared to a second biological sample with

sRIN score lower than the first biological sample. In some embodiments, a sRIN is calculated for a tissue section, one or more regions of a tissue section, or a single cell.

[1020] In some embodiments, one or more sRINs for a given biological sample (e.g., tissue section, one or more regions of a tissue, or a single cell) are calculated by: (a) providing (i) a spatial array including a plurality of capture probes on a substrate, where a capture probe comprises a capture domain and (ii) a tissue stained with a histology stain (e.g., any of the stains described herein); (b) contacting the spatial array with the biological sample (e.g., tissue); (c) capturing a biological analyte (e.g., an 18S rRNA molecule) from the biological sample (e.g., tissue) with the capture domain; (d) generating a cDNA molecule from the captured biological analyte (e.g., 18S rRNA); (e) hybridizing one or more labeled oligonucleotide probes to the cDNA; (f) imaging the labeled cDNA and the histology stain (e.g., any of the stains described herein), and (g) generating a spatial RNA integrity number for a location in the spatial array, wherein the spatial RNA integrity number comprises an analysis of a labeled cDNA image and a histology stain (e.g., any of the stains described herein) image for the location.

[1021] In some embodiments, the biological sample (e.g., tissue) is stained with a histology stain. As used herein, a “histology stain” can be any stain described herein. For example, the biological sample can be stained with IF/IHC stains described herein. For example, the biological sample (e.g., tissue) can be stained with hematoxylin and eosin (“H&E”). In some embodiments, the biological sample (e.g., tissue) is stained with a histology stain (e.g., any of the stains described herein) before, contemporaneously with, or after labelling of the cDNA with labeled oligonucleotide probes. In some embodiments, the stained biological sample can be, optionally, destained (e.g., washed 1, 2, 3, 4, 5, or more times in a low pH acid (e.g., HCl)). For example, hematoxylin, from an H&E stain, can be optionally removed from the biological sample by washing in dilute HCl (0.001M to 0.1M) prior to further processing. In some embodiments, the stained biological sample can be optionally destained after imaging and prior to permeabilization.

[1022] In some embodiments, the spatial array includes a plurality of capture probes immobilized on a substrate where the capture probes include at least a capture domain. In some embodiments, the capture domain includes a poly(T) sequence. For example, a capture domain includes a poly(T) sequence that is capable of capturing an 18S rRNA transcript from a biological sample.

[1023] In some embodiments, calculating one or more spatial RNA Integrity Numbers for a biological sample includes hybridizing at least one (e.g., at least two, at least three, at least four, or at least five) labeled oligonucleotide probes to the cDNA generated from the 18s rRNA. In some embodiments, a labeled oligonucleotide probe includes a sequence that is complementary to a portion of the 18S cDNA. In some embodiments, four labeled oligonucleotide probes (P1-P4) are designed to hybridize at four different locations spanning the entire gene body of the 18S rRNA. In some embodiments, a labeled oligonucleotide probe can include any of the detectable labels as described herein. For example, an oligonucleotide labeled probe can include a fluorescent label (e.g., Cy3). In some embodiments, one or more of the labeled oligonucleotide probes designed with complementarity to different locations within the 18S cDNA sequence include the same detectable label. For example, four labeled oligonucleotide probes, (P1-P4) each designed to have complementarity to a different location within the 18S cDNA sequence can all have the same detectable label (e.g., Cy3). In some embodiments, one or more of the labeled oligonucleotide probes designed with complementarity to different locations within the 18S cDNA sequence include a different detectable label. For example, four labeled oligonucleotide probes, (P1-P4) each designed to have complementarity to a different location within the 18S cDNA sequence can include different detectable labels.

[1024] In some embodiments, determining a spatial RNA Integrity Number for a biological sample (e.g., tissue section, one or more regions of a tissue, or a single cell) includes analyzing the images taken from a spatial array and a histology stain (e.g., any of the stains described herein) for the same location. For example, for the spatial array, all images are generated by scanning with a laser

(e.g., a 532 nm wavelength) after the fluorescently labeled (e.g., Cy3) oligonucleotide probes have been hybridized to the 18S cDNA. One image is generated per probe (P1-P4) and one image is generated where no fluorescently labeled probes were hybridized (P0). Normalization of Fluorescence Units (FU) data is performed by subtraction of the auto-fluorescence recorded with P0 and division with P1. After alignment, the five images (one image from each probe, P1-P4, and one image from an area without bound probe) are loaded into a script. The script generates two different plots, one heat-map of spatial RIN values and one image alignment error plot, which combines the histology stain (e.g., any of the stains described herein) image. The image alignment error plot is used to visualize which pixels and positions should be excluded from the analysis due to alignment errors between the images from P0-P4.

III. General Spatial Cell-Based Analytical Methodology

(a) Barcoding a Biological Sample

[1025] In some embodiments, provided herein are methods and materials for attaching and/or introducing a molecule (e.g., a peptide, a lipid, or a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a biological sample (e.g., to a cell in a biological sample) for use in spatial analysis. In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are introduced to a biological sample (e.g., to a plurality of cells in a biological sample) for use in spatial analysis.

[1026] FIG. 18 is a schematic diagram depicting cell tagging using either covalent conjugation of an analyte binding moiety to a cell surface or non-covalent interactions with cell membrane elements. FIG. 18 lists non-exhaustive examples of a covalent analyte binding moiety/cell surface interactions, including protein targeting, amine conjugation using NHS chemistry, cyanuric chloride, thiol conjugation via maleimide addition, as well as targeting glycoproteins/glycolipids expressed on the cell surface via click chemistry. Non-exhaustive examples of non-covalent interactions with cell membrane elements include lipid modified oligos, biocompatible anchor for cell membrane (BAM, e.g., oleyl-PEG-NHS), lipid modified positive neutral polymer, and antibody to membrane proteins. A cell tag can be used in combination with an analyte capture agent and cleavable or non-cleavable spatially-barcoded capture probes for spatial and multiplexing applications.

[1027] In some embodiments, a plurality of molecules (e.g., a plurality of lipid or nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are introduced to a biological sample (e.g., to a plurality of cells in a biological sample) for use in spatial analysis, wherein the plurality of molecules are introduced to the biological sample in an arrayed format. In some embodiments, a plurality of molecules (e.g., a plurality of lipid or nucleic acid molecules) having a plurality of barcodes are provided on a substrate (e.g., any of the variety of substrates described herein) in any of the variety of arrayed formats described herein, and the biological sample is contacted with the molecules on the substrate such that the molecules are introduced to the biological sample. In some embodiments, the molecules that are introduced to the biological sample are cleavably attached to the substrate, and are cleaved from the substrate and released to the biological sample when contacted with the biological sample. In some embodiments, the molecules introduced to the biological sample are covalently attached to the substrate prior to cleavage. In some embodiments, the molecules that are introduced to the biological sample are non-covalently attached to the substrate (e.g., via hybridization), and are released from the substrate to the biological sample when contacted with the biological sample.

[1028] In some embodiments, a plurality of molecules (e.g., a plurality of lipid or nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are migrated or transferred from a substrate to cells of a biological sample. In some embodiments, migrating a plurality of molecules from a substrate to cells of a biological sample includes applying a force (e.g., mechanical, centrifugal, or electrophoretic) to the substrate and/or the biological sample to facilitate migration of the plurality of molecules from the substrate to the biological sample.

[1029] In some embodiments of any of the spatial analysis methods described herein, physical force is used to facilitate attachment to or introduction of a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) into a biological sample (e.g., a cell present in a biological sample). As used herein, “physical force” refers to the use of a physical force to counteract the cell membrane barrier in facilitating intracellular delivery of molecules. Examples of physical force instruments and methods that can be used in accordance with materials and methods described herein include the use of a needle, ballistic DNA, electroporation, sonoporation, photoporation, magnetofection, hydroporation, and combinations thereof.

(i) Introducing a Cell-Tagging Agent to the Surface of a Cell

[1030] In some embodiments, biological samples (e.g., cells in a biological sample) can be labelled using cell-tagging agents where the cell-tagging agents facilitate the introduction of the molecules (e.g., nucleic acid molecules) having barcodes (e.g., spatial barcodes) into the biological sample (e.g., into cells in a biological sample). As used herein, the term “cell-tagging agent” refers to a molecule having a moiety that is capable of attaching to the surface of a cell (e.g., thus attaching the barcode to the surface of the cell) and/or penetrating and passing through the cell membrane (e.g., thus introducing the barcode to the interior of the cell). In some embodiments, a cell-tagging agent includes a barcode (e.g., a spatial barcode). The barcode of a barcoded cell-tagging agent can be any of the variety of barcodes described herein. In some embodiments, the barcode of a barcoded cell-tagging agent is a spatial barcode. In some embodiments, a cell-tagging agent comprises a nucleic acid molecule that includes the barcode (e.g., the spatial barcode). In some embodiments, the barcode of a barcoded cell-tagging agent identifies the associated molecule, where each barcode is associated with a particular molecule. In some embodiments, one or more molecules are applied to a sample. In some embodiments, a nucleic acid molecule that includes the barcode is covalently attached to the cell-tagging agent. In some embodiments, a nucleic acid molecule that includes the barcode is non-covalently attached to the cell-tagging agent. A non-limiting example of non-covalent attachment includes hybridizing the nucleic acid molecule that includes the barcode to a nucleic acid molecule on the cell-tagging agent (which nucleic acid molecule on the cell-tagging agent can be bound to the cell-tagging agent covalently or non-covalently). In some embodiments, a nucleic acid molecule attached to a cell-tagging agent that includes a barcode (e.g., a spatial barcode) also includes one or more additional domains. Such additional domains include, without limitation, a PCR handle, a sequencing priming site, a domain for hybridizing to another nucleic acid molecule, and combinations thereof.

[1031] In some embodiments, a cell-tagging agent attaches to the surface of a cell. When the cell-tagging agent includes a barcode (e.g., a nucleic acid that includes a spatial barcode), the barcode is also attached to the surface of the cell. In some embodiments of any of the spatial analysis methods described herein, a cell-tagging agent attaches covalently to the cell surface to facilitate introduction of the spatial analysis reagents. In some embodiments of any of the spatial analysis methods described herein, a cell-tagging agent attaches non-covalently to the cell surface to facilitate introduction of the spatial analysis reagents.

[1032] In some embodiments, once a cell or cells in a biological sample is spatially tagged with a cell-tagging agent(s), spatial analysis of analytes present in the biological sample is performed. In some embodiments, such spatial analysis includes dissociating the spatially-tagged cells of the biological sample (or a subset of the spatially-tagged cells of the biological sample) and analyzing analytes present in those cells on a cell-by-cell basis. Any of a variety of methods for analyzing analytes present in cells on a cell-by-cell basis can be used. Non-limiting examples include any of the variety of methods described herein and methods described in PCT Application Publication No. WO 2019/113533A1, the content of which is incorporated herein by reference in its entirety. For example, the spatially-tagged cells can be encapsulated with beads comprising one or more nucleic acid molecules having a barcode (e.g., a cellular barcode) (e.g., an emulsion). The nucleic acid present on the bead can have a domain that hybridizes to a domain on a nucleic acid present on the

tagged cell (e.g., a domain on a nucleic acid that is attached to a cell-tagging agent), thus linking the spatial barcode of the cell to the cellular barcode of the bead. Once the spatial barcode of the cell and the cellular barcode of the bead are linked, analytes present in the cell can be analyzed using capture probes (e.g., capture probes present on the bead). This allows the nucleic acids produced (using these methods) from specific cells to be amplified and sequenced separately (e.g., within separate partitions or droplets).

[1033] In some embodiments, once a cell or cells in a biological sample is spatially tagged with a cell-tagging agent(s), spatial analysis of analytes present in the biological sample is performed in which the cells of the biological sample are not dissociated into single cells. In such embodiments, various methods of spatial analysis such as any of those provided herein can be employed. For example, once a cell or cells in a biological sample is spatially tagged with a cell-tagging agent(s), analytes in the cells can be captured and assayed. In some embodiments, cell-tagging agents include both a spatial barcode and a capture domain that can be used to capture analytes present in a cell. For example, cell-tagging agents that include both a spatial barcode and a capture domain can be introduced to cells of the biological sample in a way such that locations of the cell-tagging agents are known (or can be determined after introducing them to the cells). One non-limiting example of introducing cell-tagging agents to a biological sample is to provide the cell-tagging agents in an arrayed format (e.g., arrayed on a substrate such as any of the variety of substrates and arrays provided herein), where the positions of the cell-tagging agents on the array are known at the time of introduction (or can be determined after introduction). The cells can be permeabilized as necessary (e.g., using permeabilization agents and methods described herein), reagents for analyte analysis can be provided to the cells (e.g., a reverse transcriptase, a polymerase, nucleotides, etc., in the case where the analyte is a nucleic acid that binds to the capture probe), and the analytes can be assayed. In some embodiments, the assayed analytes (and/or copies thereof) can be released from the substrate and analyzed. In some embodiments, the assayed analytes (and/or copies thereof) are assayed in situ.

[1034] Non-limiting examples of cell-tagging agents and systems that attach to the surface of a cell (e.g., thus introducing the cell-tagging agent and any barcode attached thereto to the exterior of the cell) that can be used in accordance with materials and methods provided herein for spatially analyzing an analyte or analytes in a biological sample include: lipid tagged primers/lipophilic-tagged moieties, positive or neutral oligo-conjugated polymers, antibody-tagged primers, streptavidin-conjugated oligonucleotides, dye-tagged oligonucleotides, click-chemistry, receptor-ligand systems, covalent binding systems via amine or thiol functionalities, and combinations thereof.

(ii) Introducing a Cell-Tagging Agent to the Interior of a Cell

[1035] Non-limiting examples of cell-tagging agents and systems that penetrate and/or pass through the cell membrane (e.g., thus introducing the cell-tagging agent and any barcode attached thereto to the interior of the cell) that can be used in accordance with materials and methods provided herein for spatially profiling an analyte or analytes in a biological sample include: a cell-penetrating agent (e.g., a cell-penetrating peptide), a nanoparticle, a liposome, a polymersome, a peptide-based chemical vector, electroporation, sonoporation, lentiviral vectors, retroviral vectors, and combinations thereof.

[1036] In some embodiments, a cell-tagging agent comprises a cell-penetrating agent (described below). In some embodiments, a cell-penetrating agent transports the cell-tagging agent into the cells of a biological sample. When a cell-tagging agent comprises a barcode (e.g., a nucleic acid that includes a spatial barcode), the barcode also penetrates into the cell. In some embodiments, a plurality of cell-tagging agents are cleaved (e.g., photocleaved) from an array via a cleavage domain, thus freeing the cell-tagging agents from the array and allowing at least one capture probe of the plurality to penetrate a cell. The cell-tagging agent can then interact with an intracellular biological analyte via the capture domain. In some embodiments, the plurality of capture probes is

migrated from the array into cells of the biological sample via cell-penetrating agents. In some embodiments, migrating a plurality of capture probes from the array to cells of the biological sample includes applying a force (e.g., mechanical, centrifugal, or electrophoretic) to the biological sample.

[1037] In some embodiments, the biological sample is treated with one or more reagents to facilitate migration of a plurality of freed (e.g., cleaved) capture probes into cells of a biological sample. In one embodiment, an organic solvent (e.g., methanol or acetone) may be used to permeabilize cells of a biological sample. In another embodiment, a detergent (e.g., saponin, Triton X-100™, or Tween-20™) may be used to permeabilize cells of a biological sample. In yet another embodiment, an enzyme (e.g., trypsin) may be used to permeabilize cells of a biological sample. Any suitable method of cell permeabilization may be used to practice the methods disclosed herein. In some embodiments, the biological sample can be incubated with a cellular permeabilization reagent after contacting the array with the biological sample. In some embodiments, the biological sample can be fixed according to methods described herein.

[1038] In some embodiments, migrating a plurality of freed (e.g., cleaved) capture probes into cells of a biological sample includes passive migration (e.g., diffusion). In some embodiments, migrating a plurality of freed (e.g., cleaved) capture probes into cells of a biological sample includes active migration (e.g., electrophoretic migration). In some embodiments, migrating a plurality of freed (e.g., cleaved) capture probes into cells of a biological sample comprises antibodies. In some embodiments, migrating a plurality of freed (e.g., cleaved) capture probes into cells of a biological sample comprises transfection (e.g., chemical, biological, physical, viral vectors).

[1039] FIG. 19 is a schematic showing an exemplary cell tagging method. Non-exhaustive examples of oligo delivery vehicles may include a cell penetrating peptide or a nanoparticle. Non-exhaustive examples of the delivery systems can include lipid-based polymeric and metallic nanoparticles or oligos that can be conjugated or encapsulated within the delivery system. The cell tag can be used in combination with a capture agent barcode domain and a cleavable or non-cleavable spatially-barcoded capture probes for spatial and multiplexing applications.

1. Cell-Penetrating Agent

[1040] In some embodiments of any of the spatial profiling methods described herein, identification of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a cell-penetrating agent. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is coupled to a cell-penetrating agent, and the cell-penetrating agent allows the molecule to interact with an analyte inside the cell. A “cell-penetrating agent” as used herein can refer to an agent capable of facilitating the introduction of a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain into a cell of a biological sample (see, e.g., Lovatt et al. *Nat Methods*. 2014 February; 11 (2): 190-6, which is incorporated herein by reference in its entirety). In some embodiments, a cell-penetrating agent is a cell-penetrating peptide. A “cell-penetrating peptide” as used herein refers to a peptide (e.g., a short peptide, e.g., a peptide not usually exceeding 30 residues) that has the capacity to cross cellular membranes. In some embodiments, cell-penetrating agents or cell penetrating peptides may be covalently or non-covalently coupled to a molecule (e.g., a barcoded nucleic acid molecule), likely at the 5' end of the molecule. A cell-penetrating peptide may direct the barcoded nucleic acid molecule to a specific organelle.

[1041] In some embodiments of any of the spatial profiling methods described herein, a cell-penetrating peptide coupled to a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can cross a cellular membrane using an energy dependent or an energy independent mechanism. For example, a cell-penetrating peptide can cross a cellular membrane through direct translocation through physical perturbation of the plasma membrane,

endocytosis (e.g., mediated via clathrin), adaptive translocation, pore-formation, electroporation-like permeabilization, and/or entry at microdomain boundaries. Non-limiting examples of a cell-penetrating peptide include: penetratin, tat peptide, pVEC, transportan, MPG, Pep-1, a polyarginine peptide, MAP, R6W3, (D-Arg) 9, Cys (Npys)-(D-Arg) 9, Anti-BetaGamma (MPS-Phosducin-like protein C terminus), Cys (Npys) antennapedia, Cys (Npys)-(Arg) 9, Cys (Npys)-TAT (47-57), HIV-1 Tat (48-60), KALA, mastoparan, penetratin-Arg, pep-1-cysteamine, TAT (47-57) GGG-Cys (Npys), Tat-NR2Bct, transdermal peptide, SynB1, SynB3, PTD-4, PTD-5, FHV Coat-(35-49), BMV Gag-(7-25), HTLV-II Rex-(4-16), R9-tat, SBP, FBP, MPG, MPG (Δ NLS), Pep-2, MTS, plsl, and a polylysine peptide (see, e.g., Bechara et al. FEBS Lett. 2013 Jun. 19; 587 (12): 1693-702, which is incorporated by reference herein in its entirety).

[1042] In some embodiments, there could be two orientations for cell-penetrating peptide (CPP) conjugation. For example, one orientation can be (N-terminus)-CPP-Cys-(C-terminus)-linker-NH₂C6-5'-oligo-3'; 3'-oligo-5'-NH₂C6-linker-(N-terminus)-Cys-CPP-(C-terminus). The methods herein can be performed with other CPP conjugations and orientations.

[1043] In some embodiments, cell-tagging agents further comprise a cell-penetration tag. A "cell-penetration tag" as used herein refers to an agent that can be detected as inside a cell. In some embodiments, a cell penetration tag includes a fluorophore. In some embodiments, a cell penetration tag is selected from the group consisting of: Cy2, Cy3, Cy3B, Cy3.5, Cy5, Cy5.5, Cy7, fluorescein (6-FAM), DyLight, Alexa Fluor®, and tetramethylrhodamine (TAMRA) azide.

[1044] In some embodiments, a cell penetration tag is detected. In some embodiments, a cell penetration tag is detected after releasing a plurality of capture probes from the array and removing the array from the biological sample. In some embodiments, introduction of a cell-tagging agent into the cell is determined by detecting the presence of the cell penetration tag in the cell.

[1045] In some embodiments, cell-tagging agents can optionally include an intracellular cleavage domain, wherein one or more segments or regions of the capture probe (e.g., capture domains, spatial barcodes, and/or UMIs) can be releasably or cleavably bound to the cell-penetrating agent, such that the capture domain, spatial barcode, and/or UMI can be released. In some embodiments, the cleavage of the linkage between the capture domain, spatial barcode, and/or UMI and the cell-penetrating agent is induced in an intracellular environment (e.g., the intracellular cleavage domain is cleaved after the cell-tagging agent is introduced into the cell). In some embodiments, the intracellular cleavage domain comprises a disulfide bond. For example, the intracellular cleavage domain can be a disulfide bond cleaved by reducing conditions in the cell. Any other suitable linker can be used to release or cleave the intracellular cleavage domain of the capture probe.

2. Nanoparticles

[1046] In some embodiments of any of the spatial profiling methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by an inorganic particle (e.g., a nanoparticle). In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is coupled to an inorganic particle (e.g., a nanoparticle), and the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain uses the nanoparticle to get access to analytes inside the cell. Non-limiting examples of nanoparticles that can be used in embodiments herein to deliver a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain into a cell and/or cell bead include inorganic nanoparticles prepared from metals, (e.g., iron, gold, and silver), inorganic salts, and ceramics (e.g., phosphate or carbonate salts of calcium, magnesium, or silicon). The surface of a nanoparticle can be coated to facilitate binding of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain, or the surface can be chemically modified to facilitate attachment of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain. Magnetic nanoparticles (e.g., supermagnetic iron oxide), fullerenes (e.g., soluble carbon molecules), carbon nanotubes (e.g., cylindrical fullerenes),

quantum dots, and supramolecular systems can also be used.

3. Liposomes

[1047] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a liposome. Various types of lipids, including cationic lipids, can be used in liposome delivery. In some cases, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is delivered to a cell via a lipid nano-emulsion. A lipid emulsion refers to a dispersion of one immiscible liquid in another stabilized by emulsifying agent. Labeling cells can comprise use of a solid lipid nanoparticle.

4. Polymersomes

[1048] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a polymersome. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is contained in the polymersome, and the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain uses the polymersome to get access to analytes inside the cell. A “polymersome” as referred to herein is an artificial vesicle. For example, a polymersome can be a vesicle similar to a liposome, but the membrane comprises amphiphilic synthetic block copolymers (see, e.g., Rideau et al. *Chem. Soc. Rev.*, 2018, 47, 8572-8610, which is incorporated by reference herein in its entirety). In some embodiments, polymersomes comprise di-(AB) or tri-block copolymers (e.g., ABA or ABC), where A and C are a hydrophilic block and B is a hydrophobic block. In some embodiments, a polymersome comprises poly(butadiene)-b-poly(ethylene oxide), poly(ethyl ethylene)-b-poly(ethylene oxide), polystyrene-b-poly(ethylene oxide), poly(2-vinylpyridine)-b-poly(ethylene oxide), polydimethylsiloxane-b-poly(ethylene oxide), polydimethylsiloxane-g-poly(ethylene oxide), polycaprolactone-b-poly(ethylene oxide), polyisobutylene-b-poly(ethylene oxide), polystyrene-b-polyacrylic acid, polydimethylsiloxane-b-poly-2-methyl-2-oxazoline, or a combination thereof (wherein b=block and g=grafted).

5. Peptide-Based Chemical Vectors

[1049] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a peptide-based chemical vector, e.g., a cationic peptide-based chemical vector. Cationic peptides can be rich in basic residues like lysine and/or arginine. In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a polymer-based chemical vector. Cationic polymers, when mixed with a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain, can form nanosized complexes called polyplexes. Polymer based vectors can comprise natural proteins, peptides and/or polysaccharides. Polymer based vectors can comprise synthetic polymers. In some embodiments, a polymer-based vector comprises polyethylenimine (PEI). PEI can condense DNA into positively-charged particles, which bind to anionic cell surface residues and are brought into the cell via endocytosis. In some embodiments, a polymer-based chemical vector comprises poly(L)-lysine (PLL), poly(DL-lactic acid) (PLA), poly(DL-lactide-co-glycoside) (PLGA), polyornithine, polyarginine, histones, protamines, or a combination thereof. Polymer-based vectors can comprise a mixture of polymers, for example, PEG and PLL. Other non-limiting examples of polymers include dendrimers, chitosans, synthetic amino derivatives of dextran, and cationic acrylic polymers.

6. Electroporation

[1050] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by electroporation. With electroporation, a biological

analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can enter a cell through one or more pores in the cellular membrane formed by applied electricity. The pore of the membrane can be reversible based on the applied field strength and pulse duration.

7. Sonoporation

[1051] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by sonoporation. Cell membranes can be temporarily permeabilized using sound waves, allowing cellular uptake of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain.

8. Lentiviral Vectors and Retroviral Vectors

[1052] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by vectors. For example, a vector as described herein can be an expression vector where the expression vector includes a promoter sequence operably linked to the sequence encoding the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain. Non-limiting examples of vectors include plasmids, transposons, cosmids, and viral derived vectors (e.g., any adenoviral derived vectors (AV) cytomegaloviral derived (CMV) vectors, simian viral derived (SV40) vectors, adeno-associated virus (AAV) vectors, lentivirus vectors, and retroviral vectors), and any Gateway® vectors. A vector can, for example, include sufficient cis-acting elements for expression where other elements for expression can be supplied by the host mammalian cell or in an in vitro expression system. Skilled practitioners will be capable of selecting suitable vectors and mammalian cells for introducing any of spatial profiling reagents described herein.

9. Other Methods and Cell-Tagging Agents for Intracellular Introduction of a Molecule

[1053] In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by the use of a needle, for example for injection (e.g., microinjection), particle bombardment, photoporation, magnetofection, and/or hydroporation. For example, with particle bombardment, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can be coated with heavy metal particles and delivered to a cell at a high speed. In photoporation, a transient pore in a cell membrane can be generated using a laser pulse, allowing cellular uptake of a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain. In magnetofection, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can be coupled to a magnetic particle (e.g., magnetic nanoparticle, nanowires, etc.) and localized to a target cell via an applied magnetic field. In hydroporation, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can be delivered to a cell and/or cell bead via hydrodynamic pressure.

(iii) Lipid Tagged Primers/Lipophilic-Tagged Moieties

[1054] In some embodiments of any of the spatial profiling methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to a lipophilic molecule. In some embodiments, the lipophilic molecule enables delivery of the lipophilic molecule to the cell membrane or the nuclear membrane. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) coupled to a lipophilic molecule can associate with and/or insert into lipid membranes such as cell membranes and nuclear membranes. In some cases, the insertion can be reversible. In some cases, the association between the lipophilic molecule and the cell may be such that the cell retains the lipophilic molecule (e.g., and associated components, such as nucleic acid barcode molecules) during subsequent processing (e.g., partitioning, cell permeabilization, amplification, pooling, etc.). In some embodiments, a

molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) coupled to a lipophilic molecule may enter into the intracellular space and/or a cell nucleus.

[1055] Non-limiting examples of lipophilic molecules that can be used in embodiments described herein include sterol lipids such as cholesterol, tocopherol, steryl, palmitate, lignoceric acid, and derivatives thereof. In some embodiments, the lipophilic molecules are neutral lipids that are conjugated to hydrophobic moieties (e.g., cholesterol, squalene, or fatty acids) (See Raouane et al. *Bioconjugate Chem.*, 23 (6): 1091-1104 (2012) which is herein incorporated by reference in its entirety). In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) may be attached to the lipophilic moiety via a linker, using covalent or direct attachment. In some embodiments, the linker is a tetra-ethylene glycol (TEG) linker. Other exemplary linkers include, but are not limited to, Amino Linker C6, Amino Linker C12, Spacer C3, Spacer C6, Spacer C12, Spacer 9, and Spacer 18. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is indirectly coupled (e.g., via hybridization or ligand-ligand interactions, such as biotin-streptavidin) to a lipophilic molecule. In some embodiments, the lipophilic moiety can be attached to a capture probe, spatial barcode, or other DNA sequence, at either the 5' or 3' end of the specified DNA sequence. In some embodiments, the lipophilic moiety can be coupled to a capture probe, spatial barcode, or other DNA sequence in a lipid-dependent manner. Other lipophilic molecules that may be used in accordance with methods provided herein include amphiphilic molecules wherein the headgroup (e.g., charge, aliphatic content, and/or aromatic content) and/or fatty acid chain length (e.g., C12, C14, C16, or C18) can be varied. For instance, fatty acid side chains (e.g., C12, C14, C16, or C18) can be coupled to glycerol or glycerol derivatives (e.g., 3-t-butylidiphenylsilylglycerol), which can also comprise, e.g., a cationic head group. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) disclosed herein can then be coupled (either directly or indirectly) to these amphiphilic molecules. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) coupled to an amphiphilic molecule may associate with and/or insert into a membrane (e.g., a cell, cell bead, or nuclear membrane). In some cases, an amphiphilic or lipophilic moiety may cross a cell membrane and provide a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to an internal region of a cell and/or cell bead.

[1056] In some embodiments, additives can be added to supplement lipid-based modifications. In some embodiments, the additive is low density lipoprotein (LDL). In some embodiments, the additive is the cholesterol trafficking inhibitor U-18666A. In some embodiments, U-18666A inhibits cholesterol transport from late endosomes at micromolar concentrations and/or lysosomes to the endoplasmic reticulum (ER) at nanomolar concentrations. In some embodiments, U-18666A can inhibit oxidosqualene cyclase, a key enzyme in the cholesterol biosynthesis pathway, at sufficiently high concentrations (e.g., at or about >0.5 mM).

[1057] In some embodiments, wherein the molecule (e.g., with a nucleic acid sequence) has an amino group within the molecule, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an amino group can be coupled to an amine-reactive lipophilic molecule. For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an amino group can be conjugated to DSPE-PEG (2000)-cyanuric chloride (1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[cyanur(polyethylene glycol)-2000]).

[1058] In some embodiments, a cell-tagging agent can attach to a surface of a cell through a combination of lipophilic and covalent attachment. For example, a cell-tagging agent can include an oligonucleotide attached to a lipid to target the oligonucleotide to a cell membrane, and an amine group that can be covalently linked to a cell surface protein(s) via any number of chemistries described herein. In these embodiments, the lipid can increase the surface concentration of the oligonucleotide and can promote the covalent reaction.

[1059] As used herein, an “anchor oligonucleotide” and/or “co-anchor oligonucleotide” can include

a lipid-conjugated oligonucleotide, wherein the lipid is capable of becoming embedded within a cell membrane. In some embodiments, the lipid capable of becoming embedded within a cell membrane includes but is not limited to, sterol lipids such as cholesterol, tocopherol, steryl, palmitate, lignoceric acid, and derivatives thereof. In some embodiments, the sterol lipid of the anchor oligonucleotide and/or co-anchor oligonucleotide can be attached to either the 5' or 3' end of the oligonucleotide portion. In some embodiments, the anchor oligonucleotide and/or the co-anchor oligonucleotide can integrate into the cell membrane of a cell in a biological sample (e.g., the sterol lipid of the anchor oligonucleotide and/or co-anchor oligonucleotide).

[1060] In some embodiments, a sterol lipid (e.g., lignoceric acid) anchor oligonucleotide is attached to the 5' end of the oligonucleotide. In some embodiments, the anchor oligonucleotide can have a constant sequence. In some embodiments the constant sequence of the anchor oligonucleotide can be about 15 to about 30 nucleotides long. In some embodiments, the anchor oligonucleotide can have an additional domain 3' to the constant sequence. In some embodiments, the additional domain can be an adapter sequence (e.g., sequencing adapter). In some embodiments, the adapter sequence can be about 15 to about 35 nucleotides long.

[1061] In some embodiments, the lipid (e.g., sterol lipid) of the co-anchor oligonucleotide (e.g., palmitic acid), is attached to the 3' end of the oligonucleotide. In some embodiments, the co-anchor oligonucleotide can have a constant sequence. For example, the constant sequence of the co-anchor oligonucleotide can be a reverse complement of the constant sequence from the anchor oligonucleotide. In some embodiments, the constant sequence of the anchor oligonucleotide and the constant sequence of the co-anchor oligonucleotide can bind (e.g., hybridize) to each other. In some embodiments, the lipid (e.g., sterol lipid) of the anchor oligonucleotide and the co-anchor oligonucleotide can integrate into a cell membrane in the biological sample and the respective constant sequences can hybridize to each other at the same time. In some embodiments, a barcoded oligonucleotide, which can include several domains, can be introduced to the integrated anchor oligonucleotide and co-anchor oligonucleotide hybridized to each other. The barcoded oligonucleotide can include, in a 5' to 3' direction, a functional domain (e.g., a sequencing adapter domain), a unique molecular identifier, a sample barcode, a second unique molecular identifier, and the reverse complement of a constant sequence. For example, after tagging a cell with any of the cell-tagging agents described herein the cells can be partitioned (e.g., encapsulated in a vesicle) with a barcoded feature (e.g., a bead). In some embodiments, the reverse complement of the constant sequence of the barcoded oligonucleotide can interact (e.g., hybridize) with the constant sequence (e.g., a portion of the sequence) on the barcoded feature.

(iv) Intracellular Cleavage Domain

[1062] As used herein, capture probes can optionally include an “intracellular cleavage domain,” wherein one or more segments or regions of the capture probe (e.g., capture domains, spatial barcodes, and/or UMIs) can be releasably or cleavably attached to one or more other segments or regions of the capture probe, such as a cell-penetrating agent, such that the capture domain, spatial barcode, and/or UMI can be released or be releasable through cleavage of a linkage between the capture domain, spatial barcode, and/or UMI and the cell-penetrating agent and/or cell penetration tag. In some embodiments, the cleavage of the linkage between the capture domain, spatial barcode, and/or UMI and the cell-penetrating agent is induced in an intracellular environment (e.g., the intracellular cleavage domain is cleaved after the capture probes is introduced into the cell). For example, the linkage between the capture domain, spatial barcode, and/or UMI and the cell-penetrating agent can be a disulfide bond that is cleaved by the reducing conditions in the cell, for example, when the intracellular cleavage domain comprises a disulfide bond. Any other suitable linker can be used to release or cleave the intracellular cleavage domain of the capture probe.

(v) Positive or Neutral Oligo-Conjugated Polymers

[1063] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a glycol

chitosan derivative. In some embodiments, the glycol chitosan derivative can be coupled with two or more molecules (e.g., nucleic acid molecules) having a barcode (e.g., a spatial barcode). In some embodiments, the glycol chitosan derivative can be coupled with about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10 or more molecules. The glycol chitosan derivative (e.g., glycol chitosan-cholesterol) can serve as a hydrophobic anchor (see Wang et al. *J. Mater. Chem. B.*, 30:6165 (2015), which is herein incorporated by reference in its entirety). Non-limiting examples of chitosan derivatives that can be coupled to a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be found in Cheung et al., *Marine Drugs*, 13 (8): 5156-5186 (2015), which is herein incorporated by reference in its entirety.

(vi) Bifunctional NHS Linker Cell-Tagging

[1064] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a bifunctional NHS linker. In some embodiments, the coupled bifunctional NHS linker (e.g., bifunctional linker and the molecule having a barcode) can facilitate the attachment of the spatial barcode to the surface of the cell. In some embodiments, after facilitating attachment to the surface of the cell, excess NHS linker can be removed (e.g., washed away). In some embodiments, the process of coupling the molecule having a barcode can be performed under non-anhydrous conditions to maintain the activity of unreacted bifunctional NHS. In some embodiments, the non-anhydrous condition can be in the presence of DMSO. In some embodiments, the non-anhydrous condition can be in the presence of DMF.

(vii) Antibody-Tagged Primers

[1065] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody or antigen binding fragment thereof in a manner that facilitates attachment of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to the surface of a cell. In some embodiments, facilitating attachment to the cell surface facilitates introduction of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) into the cell. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody that is directed to an antigen that is present on the surface of a cell. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody that is directed to an antigen that is present on the surface of a plurality of cells (e.g., a plurality of cells in a biological sample). In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody that is directed to an antigen that is present on the surface of a cell, a plurality of cells, or substantially all the cells present in a biological sample. In some embodiments, the barcoded-antibody is directed to an intracellular antigen. Any of the exemplary methods described herein of attaching a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to another molecule (e.g., an antibody or antigen fragment thereof) can be used.

(viii) Streptavidin-Conjugated Oligonucleotides

[1066] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can attach to the surface of a cell using biotin-streptavidin. In some embodiments, primary amines in the side chain of lysine residues of cell surface polypeptides are labelled with NHS-activated biotin reagents. For example, the N-terminus of a polypeptide can react with NHS-activated biotin reagents to form stable amide bonds. In some embodiments, cell-tagging agents include molecules (e.g., a nucleic acid molecule) having barcodes (e.g., a spatial barcode) conjugated to streptavidin. In some cases, streptavidin can be conjugated to the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) using click chemistry (e.g., maleimide modification) as described herein. In some embodiments, a cell containing NHS-activated biotin incorporated into lysine side chains of a cell

surface protein forms a non-covalent bond with the streptavidin conjugated to the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) conjugated to streptavidin is itself part of a cell-tagging agent.

(ix) Dye-Tagged Oligonucleotides

[1067] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is directly linked to a detectable label. In some embodiments, the detectable label is any of the detectable labels described herein. In some embodiments the detectable label is a fluorescent tag. In some embodiments, the physical properties of the fluorescent tags (e.g., a fluorescent tag having hydrophobic properties) can overcome the hydrophilic nature of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). For example, in some embodiments, wherein the molecule is a nucleic acid molecule, a fluorescent tag (e.g., BODIPY, Cy3, Atto 647N, and Rhodamine Red C.sub.2) can be coupled to a 5' end of the nucleic acid molecule having a barcode (e.g., a spatial barcode). In some embodiments, wherein the molecule is a nucleic acid molecule, any fluorescent tag having hydrophobic properties can be coupled to the nucleic acid molecule having a barcode (e.g., a spatial barcode) in a manner that overcomes the hydrophilic nature of the nucleic acid molecule. Non-limiting examples of fluorescent tags with hydrophobic properties include BODIPY, Cy3, Atto 647N, and Rhodamine Red C2.

(x) Click-Chemistry

[1068] In some embodiments of any of the spatial analysis methods described herein, molecules (e.g., a nucleic acid molecule) having barcodes (e.g., a spatial barcode) are coupled to click-chemistry moieties. As used herein, the term “click chemistry,” generally refers to reactions that are modular, wide in scope, give high yields, generate byproducts, such as those that can be removed by nonchromatographic methods, and are stereospecific (but not necessarily enantioselective) (see, e.g., *Angew. Chem. Int. Ed.*, 2001, 40 (11): 2004-2021, which is incorporated herein by reference in its entirety). In some cases, click chemistry can describe pairs of functional groups that can selectively react with each other in mild, aqueous conditions.

[1069] An example of a click chemistry reaction is the Huisgen 1,3-dipolar cycloaddition of an azide and an alkyne, i.e., copper-catalyzed reaction of an azide with an alkyne to form the 5-membered heteroatom ring 1,2,3-triazole. The reaction is also known as a Cu(I)-Catalyzed Azide-Alkyne Cycloaddition (CuAAC), a Cu(I) click chemistry or a Cu⁺ click chemistry. Catalysts for the click chemistry include, but are not limited to, Cu(I) salts, or Cu(I) salts made in situ by reducing Cu(II) reagents to Cu(I) reagents with a reducing reagent (*Pharm Res.* 2008, 25 (10): 2216-2230, which is incorporated herein by reference in its entirety). Known Cu(II) reagents for the click chemistry can include, but are not limited to, the Cu(II)-(TBTA) complex and the Cu(II) (THPTA) complex. TBTA, which is tris-[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl]amine, also known as tris-(benzyltriazolylmethyl)amine, can be a stabilizing ligand for Cu(I) salts. THPTA, which is tris-(hydroxypropyltriazolylmethyl)amine, is another example of a stabilizing agent for Cu(I). Other conditions can also be used to construct the 1,2,3-triazole ring from an azide and an alkyne using copper-free click chemistry, such as the Strain-promoted Azide-Alkyne Click chemistry reaction (SPAAC) (see, e.g., *Chem. Commun.*, 2011, 47:6257-6259 and *Nature*, 2015, 519 (7544): 486-90, each of which is incorporated herein by reference in its entirety).

[1070] In some embodiments of any of the spatial analysis described herein, molecules (e.g., a nucleic acid molecule) having barcodes (e.g., a spatial barcode) are coupled to click-chemistry moieties without copper being present (e.g., copper-free click chemistry). One exemplary copper-free click chemistry methodology comprises a reaction between cyclooctyne and phenyl azide that results in the product 1-phenyl-4,5,6,7,8,9-hexahydro-1H-cycloocta[d][1,2,3]triazole (See, e.g., 2010, Akeroyd, N., et al., Click chemistry for the preparation of advance macromolecular architectures, PhD dissertation, incorporated herein by reference in its entirety). Additional copper-

free click chemistry methods are known to those skilled in the art (See, e.g., 2009, Click Chemistry for Biotechnology and Materials Science, Ed. Jeorg Lahann, John Wiley & Sons, Ltd. publ., p. 410).

(xi) Receptor-Ligand Systems

[1071] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a ligand, wherein the ligand is part of a receptor-ligand interaction on the surface of a cell. For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a ligand that interacts selectively with a cell surface receptor thereby targeting the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a specific cell. Non-limiting examples of receptor-ligand systems that can be used include integrin receptor-ligand interactions, GPCR receptor-ligand interactions, RTK receptor-ligand interactions, and TLR-ligand interactions (see Juliano, *Nucleic Acids Res.*, 44 (14): 6518-6548 (2016), which is incorporated herein by reference in its entirety). Any of the methods described herein for attaching a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a ligand (e.g., any of the methods described herein relating to attaching a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to an antibody) can be used.

(xii) Covalent Binding Systems Via Amine or Thiol Functionalities

[1072] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can incorporate reactive functional groups at sites within the molecule (e.g., with a nucleic acid sequence). In such cases, the reactive functional groups can facilitate conjugation to ligands and/or surfaces. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can include thiol modifiers that are designed to react with a broad array of activated accepting groups (e.g., maleimide and gold microspheres). For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) having thiol modifiers can interact with a maleimide-conjugated peptide thereby resulting in labelling of the peptide. In some embodiments, maleimide-conjugated peptides are present on the surface of a cell whereupon interaction with the thiol-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode), the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to the surface of the cell. Non-limiting examples of thiol modifiers include: 5' thiol modifier C6 S—S, 3' thiol modifier C3 S—S, dithiol, 3'thiol modifier oxa 6-S—S, and dithiol serinol.

[1073] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can include amine modifiers, e.g., amine modifiers that are designed to attach to another molecule in the presence of an acylating agent. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can include amine modifiers that are designed to attach to a broad array of linkage groups (e.g., carbonyl amide, thiourea, sulfonamide, and carboxamide). For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an amine modifier can interact with a sulfonamide-conjugated peptide thereby resulting in labelling of the peptide. In some embodiments, sulfonamide-conjugated peptides are present on the surface of a cell whereupon interaction with the amine-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode), the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to the surface of the cell. Non-limiting example of amine modifiers include: DMS (O) MT-Amino-Modifier-C6, Amino-Modifier-C3-TFA, Amino-Modifier-C12, Amino-Modifier-C6-TFA, Amino-dT, Amino-Modifier-5, Amino-Modifier-C2-dT, Amino-Modifier-C6-dT, and 3'-Amino-Modifier-C7.

[1074] As another example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can incorporate reactive functional groups at sites within the molecule (e.g., with a nucleic acid sequence) such as N-hydroxysuccinimide (NHS). In some embodiments, amines (e.g.,

amine-containing peptides) are present on the surface of a cell whereupon interaction with the NHS-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode), the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to the surface of the cell. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is reacted with a bifunctional NHS linker to form an NHS-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode).

[1075] In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a biocompatible anchor for cell membrane (BAM). For example, a BAM can include molecules that comprise an oleyl group and PEG. The oleyl group can facilitate anchoring the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a cell, and the PEG can increase water solubility. In some embodiments, oleyl-PEG-NHS can be coupled to a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) using NHS chemistry.

(xiii) Azide-Based Systems

[1076] In some embodiments, wherein the molecule (e.g., with a nucleic acid sequence) incorporates reactive functional groups at sites within the molecule, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an azide group on a cell surface. In some embodiments, the reactive functional group is an alkynyl group. In some embodiments, click chemistry as described herein can be used to attach the alkynyl-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to an azide group on the cell surface. An azide group can be attached to the cell surface through a variety of methods. For example, NHS chemistry can be used to attach an azide group to the cell surface. In some embodiments, N-azidoacetylmannosamine-tetraacylated (Ac4ManNAz), which contains an azide group, can react with sialic acid on the surface of a cell to attach azide to the cell surface. In some embodiments, azide is attached to the cell surface by bio-orthogonal expression of azide. For example, azide is incubated with the cells. In some embodiments, the alkynyl-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can attach to the surface of a cell via an azide group in the presence of copper. In some embodiments, the alkynyl-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can attach to the surface of a cell via an azide group in the absence of copper.

(xiv) Lectin-Based Systems

[1077] In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a lectin that facilitates attachment of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a cell surface. Lectin can bind to glycans, e.g., glycans on the surface of cells. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) has an incorporated reactive functional group such as an azide group. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an azide group is reacted with a modified lectin, e.g., a lectin modified using NHS chemistry to introduce an azide reactive group. In some embodiments, a live cell is labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, a fixed cell is labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, a permeabilized cell is labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, organelles in the secretory pathway can be labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode).

(b) Methods for Separating a Sample into Single Cells or Cell Groups

[1078] Some embodiments of any of the methods described herein can include separating a biological sample into single cells, cell groups, types of cells, or a region or regions of interest. For example, a biological sample can be separated into single cells, cell groups, types of cells, or a

region or regions of interest before being contacted with one or more capture probes. In other examples, a biological sample is first contacted with one or more capture probes, and then separated into single cells, cell groups, types of cells, or a region or regions of interest.

[1079] In some embodiments, a biological sample can be separated into chunks using pixelation. Pixelation can include the steps of providing a biological sample, and punching out one or more portions of the biological sample. The punched out portions of the biological sample can then be used to perform any of the methods described herein. In some embodiments, the punched-out portions of the biological sample can be in a random pattern or a designed pattern. In some embodiments, the punched-out portions of the biological sample can be focused on a region of interest or a subcellular structure in the biological sample.

[1080] FIG. 20A is a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for “pixelating” a sample, wherein the sample is cut, stamped, microdissected, or transferred by hollow-needle or microneedle, moving a small portion of the sample into an individual partition or well.

[1081] FIG. 20B is a schematic depicting multi-needle pixelation, wherein an array of needles is punched through a sample on a scaffold, into nanowells containing beads (e.g., gel beads) and reagents. Once the needle is in the nanowell, the cell(s) are ejected.

[1082] In some embodiments, a biological sample (e.g., a tissue sample or tissue section) is divided into smaller portions as compared to the original biological sample size (“chunks”) before performance of any of the spatial analysis methods described herein. In some embodiments, the methods can include spatial barcoding of FFPE “chunks” via barcodes applied in a spatially well-defined pattern (e.g., array printing). In order to associate a spatial barcode with a particular “chunk” of biological sample, the barcode (e.g., a spatial barcode) can be of a sufficient length to prevent diffusion of the barcode in subsequent steps, or the spatial barcode can be covalently applied to the FFPE sample. In some embodiments, the spatial barcode is unique to each FFPE chunk. In some embodiments, spatial barcodes can be embedded onto an FFPE slide (e.g., within a matrix, such as a wax or a hydrogel). In some embodiments, the FFPE slide is heated (e.g., wax is heated) prior to addition of the spatial barcodes. In some embodiments, after addition of the spatial barcodes, the FFPE slide can be cooled and cut or dissociated into chunks. Methods of chunking (e.g., cutting) biological samples are known in the art. For example, in a non-limiting example, chunking of biological samples can be done in various ways such as laser microdissection, mechanical means, acoustic (e.g., sonication) means, or any other method described herein. In some embodiments, fluorophores/Qdots, etc. can be embedded in the chunk to preserve spatial information about the biological sample. Barcoding at this step enables massively parallel encapsulation of chunks while retaining local spatial information (e.g., tumor versus normal/healthy cells). In some embodiments, chunking of a biological sample (e.g., a tissue section) can result in single-cell chunks of the biological sample. In other embodiments, chunking of a biological sample can be performed to obtain chunks that correspond to diseased portions of the biological sample. In another embodiment, chunking of biological samples can be performed to obtain discrete chunks of the biological sample that correspond to diseased or healthy portions of the biological sample. In some embodiments, chunking of biological samples can be performed to obtain chunks that correspond to specific cell types (e.g., chunking based on fluorescent or chemiluminescent imaging of antibodies bound to target proteins) in the biological sample.

[1083] In some embodiments, the spatially-barcoded chunks can be further processed. For example, the spatially-barcoded chunk can be individually encapsulated (e.g., a matrix, emulsion, or hydrogel). In some embodiments, the spatially-barcoded chunk can be encapsulated in a partition (e.g., a well, droplet, channel, or vesicle). In some embodiments, the spatially-barcoded chunk can be encapsulated in a vesicle. In some embodiments, the vesicle can comprise a lipid bilayer. In some embodiments, the spatially-barcoded FFPE chunk can be encapsulated with a uniquely barcoded bead. In some embodiments, the uniquely barcoded bead can have a functional

domain, a cleavage domain, a unique molecular identifier, and a capture domain, or combinations thereof. In some embodiments, the encapsulated spatially-barcoded FFPE chunk and the uniquely barcoded bead can be heated to deparaffinize the FFPE sample. In some embodiments, the encapsulated spatially-barcoded FFPE chunk and the uniquely barcoded bead can be treated with xylene to deparaffinize the FFPE sample. In some embodiments, the deparaffinized sample can be treated to de-crosslink methylene bridges in a single step. In some embodiments, additional steps can be performed when, for example, de-crosslinking chemistry is incompatible with barcoding or library preparation steps. In some embodiments, after de-crosslinking methylene bridges, the nucleic acids originating or present in the chunk can bind to the uniquely barcoded bead. In some embodiments, after the spatial barcode binds the uniquely barcoded bead, the encapsulation can be disrupted (e.g., lysed, melted, or removed) and the barcoded beads can be collected. In some embodiments, the collected barcoded beads can be washed and re-encapsulated. In some embodiments, the nucleic acids associated with the bead (e.g., spatial barcode, unique barcode, analyte transcript) can be amplified (e.g., PCR amplified) and processed (e.g., sequenced) according to any of the methods described herein.

[1084] In some embodiments, a biological sample can be divided or portioned using laser capture microdissection (e.g., highly-multiplexed laser capture microdissection).

(c) Release and Amplification of Analytes

[1085] In some embodiments, lysis reagents can be added to the sample to facilitate release of analyte(s) from a sample. Examples of lysis agents include, but are not limited to, bioactive reagents such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other commercially available lysis enzymes. Other lysis agents can additionally or alternatively be co-partitioned with the biological sample to cause the release of the sample's contents into the partitions. In some embodiments, surfactant-based lysis solutions can be used to lyse cells, although these can be less desirable for emulsion-based systems where the surfactants can interfere with stable emulsions. Lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption can also be used in certain embodiments, e.g., non-emulsion based partitioning such as encapsulation of biological materials that can be in addition to or in place of droplet partitioning, where any pore volume of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

[1086] In addition to the permeabilization agents, other reagents can also be added to interact with the biological sample, including, for example, DNase and RNase inactivating agents or inhibitors or chelating agents, such as EDTA, and other reagents to allow for subsequent processing of analytes from the sample. In other embodiments, nucleases, such as DNase or RNase, or proteases, such as pepsin or proteinase K, are added to the sample.

[1087] Further reagents that can be added to a sample, include, for example, endonucleases to fragment DNA, DNA polymerase enzymes, and dNTPs used to amplify nucleic acids. Other enzymes that can also be added to the sample include, but are not limited to, polymerase, transposase, ligase, proteinase K, and DNase, etc. Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers, and switch oligonucleotides. In some embodiments, template switching can be used to increase the length of a cDNA, e.g., by appending a predefined nucleic acid sequence to the cDNA.

[1088] If a tissue sample is not permeabilized sufficiently, the amount of analyte captured on the substrate can be too low to enable adequate analysis. Conversely, if the tissue sample is too permeable, the analyte can diffuse away from its origin in the tissue sample, such that the relative spatial relationship of the analytes within the tissue sample is lost. Hence, a balance between permeabilizing the tissue sample enough to obtain good signal intensity while still maintaining the spatial resolution of the analyte distribution in the tissue sample is desired.

[1089] In some embodiments, where the biological sample includes live cells, permeabilization conditions can be modified so that the live cells experience only brief permeabilization (e.g., through short repetitive bursts of electric field application), thereby allowing one or more analytes to migrate from the live cells to the substrate while retaining cellular viability.

[1090] In some embodiments, after contacting a biological sample with a substrate that include capture probes, a removal step is performed to remove all or a portion of the biological sample from the substrate. In some embodiments, the removal step includes enzymatic or chemical degradation of the permeabilized cells of the biological sample. For example, the removal step can include treating the biological samples with an enzyme (e.g., proteinase K) to remove at least a portion of the biological sample from the first substrates. In some embodiments, the removal step can include ablation of the tissue (e.g., laser ablation).

[1091] In some embodiments, where RNA is captured from cells in a sample, one or more RNA species of interest can be selectively enriched. For example, one or more species of RNA of interest can be selected by addition of one or more oligonucleotides. One or more species of RNA can be selectively down-selected (e.g., removed) using any of a variety of methods. For example, probes can be administered to a sample that selectively hybridize to ribosomal RNA (rRNA), thereby reducing the pool and concentration of rRNA in the sample. Subsequent application of the capture probes to the sample can result in improved RNA capture due to the reduction in non-specific RNA present in the sample. In some embodiments, the additional oligonucleotide is a sequence used for priming a reaction by a polymerase. For example, one or more primer sequences with sequence complementarity to one or more RNAs of interest, can be used to amplify the one or more RNAs of interest, thereby selectively enriching these RNAs. In some embodiments, an oligonucleotide with sequence complementarity to the complementary strand of captured RNA (e.g., cDNA) can bind to the cDNA. In one non-limiting example, biotinylated oligonucleotides with sequence complementary to one or more cDNA of interest binds to the cDNA and can be selected using biotinylation-streptavidin affinity in any number of methods known to the field (e.g., streptavidin beads).

[1092] In some embodiments, any of the spatial analysis methods described herein can include modulating the rate of interaction between biological analytes from the biological sample and the capture probes on the array. In some embodiments, modulating the rate of interaction can occur by modulating the biological sample (e.g., modulating temperature or pH). In some embodiments, modulating the rate of interaction includes using external stimuli. Non-limiting examples of external stimuli that can be used to modulate the rate of interaction include light, temperature, small molecules, enzymes, and/or an activating reagent. In one example, light can be used to activate a polymerase in a nucleic acid extension reaction. In another example, temperature can be used to modulate hybridization between two complementary nucleic acid molecules.

[1093] Nucleic acid analytes can be amplified using a polymerase chain reaction (e.g., digital PCR, quantitative PCR, or real time PCR), isothermal amplification, or any nucleic acid amplification or extension reactions described herein, or known in the art.

(d) Partitioning

[1094] As discussed above, in some embodiments, the sample can optionally be separated into single cells, cell groups (e.g., based on cell sub-type or gene expression profile), or other fragments/pieces that are smaller than the original sample. Each of these smaller portions of the sample can be analyzed to obtain spatially-resolved analyte information from the sample. Non-limiting partitioning methods are described herein.

[1095] For samples that have been separated into smaller fragments—and particularly, for samples that have been disaggregated, dissociated, or otherwise separated into individual cells—one method for analyzing the fragments involves partitioning the fragments into individual partitions (e.g., fluid droplets), and then analyzing the contents of the partitions. In general, each partition maintains separation of its own contents from the contents of other partitions. For example, the partition can

be a droplet in an emulsion.

[1096] The methods described herein provide for the compartmentalization or partitioning of a cell (e.g., a cell) from a sample into discrete compartments or voxels. As used herein, each “voxel” represents a 3-dimensional volumetric unit. In some embodiments, a voxel maintains separation of its own contents from the contents of other voxels. A voxel can be one partition of an array of partitions of volume. For example, a voxel can be one partition of an array of discrete partitions into which a 3-dimensional object is divided. As another example, members of a plurality of photo-crosslinkable polymer precursors can be cross-linked into voxels that are part of an array of the photo-crosslinked polymer covering the substrate or a portion of the substrate. Unique identifiers, e.g., barcodes, may be previously, subsequently, or concurrently delivered to the cell, in order to allow for the later attribution of the characteristics of the cell to the particular voxel. In some embodiments, a voxel has defined dimensions. In some embodiments, a voxel comprises a single cell.

[1097] For example, a substrate can be coated with a DTT-sensitive hydrogel and then contacted with a biological sample. Optionally, capture probes attached to the substrate are released from the substrate such that the released capture probes are introduced into the biological sample and at least one released capture probe interacts with at least one biological analyte present in the biological sample via the capture domain. The biological sample and substrate can be assembled into a flow-cell and a photo-crosslinkable polymer precursor added. The cells of the biological sample can be then crosslinked into hydrogel-voxels of defined dimensions using a light source. The flow-cell can be dismantled and washed to remove unpolymerized polymer precursors. The coating can be treated with DTT to yield single-cell partitions for use in downstream applications. The capture probes/biological analytes can be analyzed, and the spatial information of the spatially-barcoded features can be used to determine the spatial location of the captured biological analytes in the biological sample.

[1098] In addition to analytes, a partition can include additional components, and in particular, one or more beads. A partition can include a single gel bead, a single cell bead, or both a single cell bead and single gel bead.

[1099] A partition can also include one or more reagents. Unique identifiers, such as barcodes, can be injected into the droplets previous to, subsequent to, or concurrently with droplet generation, such as via a microcapsule (e.g., bead). Microfluidic channel networks (e.g., on a chip) can be utilized to generate partitions. Alternative mechanisms can also be employed in the partitioning of individual biological particles, including porous membranes through which aqueous mixtures of cells are extruded into non-aqueous fluids.

[1100] The partitions can be flowable within fluid streams. The partitions can include, for example, micro-vesicles that have an outer barrier surrounding an inner fluid center or core. In some cases, the partitions can include a porous matrix that is capable of entraining and/or retaining materials within its matrix. The partitions can be droplets of a first phase within a second phase, wherein the first and second phases are immiscible. For example, the partitions can be droplets of aqueous fluid within a non-aqueous continuous phase (e.g., oil phase). In another example, the partitions can be droplets of a non-aqueous fluid within an aqueous phase. In some examples, the partitions can be provided in a water-in-oil emulsion or oil-in-water emulsion. A variety of different vessels are described in, for example, U.S. Patent Application Publication No. 2014/0155295, the entire contents of which are incorporated herein by reference. Emulsion systems for creating stable droplets in non-aqueous or oil continuous phases are described, for example, in U.S. Patent Application Publication No. 2010/0105112, the entire contents of which are incorporated herein by reference.

[1101] For droplets in an emulsion, allocating individual particles to discrete partitions can be accomplished, for example, by introducing a flowing stream of particles in an aqueous fluid into a flowing stream of a non-aqueous fluid, such that droplets are generated at the junction of the two

streams. Fluid properties (e.g., fluid flow rates, fluid viscosities, etc.), particle properties (e.g., volume fraction, particle size, particle concentration, etc.), microfluidic architectures (e.g., channel geometry, etc.), and other parameters can be adjusted to control the occupancy of the resulting partitions (e.g., number of analytes per partition, number of beads per partition, etc.) For example, partition occupancy can be controlled by providing the aqueous stream at a certain concentration and/or flow rate of analytes.

[1102] To generate single analyte partitions, the relative flow rates of the immiscible fluids can be selected such that, on average, the partitions can contain less than one analyte per partition to ensure that those partitions that are occupied are primarily singly occupied. In some cases, partitions among a plurality of partitions can contain at most one analyte. In some embodiments, the various parameters (e.g., fluid properties, particle properties, microfluidic architectures, etc.) can be selected or adjusted such that a majority of partitions are occupied, for example, allowing for only a small percentage of unoccupied partitions. The flows and channel architectures can be controlled as to ensure a given number of singly occupied partitions, less than a certain level of unoccupied partitions and/or less than a certain level of multiply occupied partitions.

[1103] The channel segments described herein can be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure can have a variety of geometries. For example, a microfluidic channel structure can have one or more than one channel junction. As another example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying particles that meet at a channel junction. Fluid can be directed to flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can include compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid can also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary, and/or gravity flow.

[1104] A partition can include one or more unique identifiers, such as barcodes. Barcodes can be previously, subsequently, or concurrently delivered to the partitions that hold the compartmentalized or partitioned biological particle. For example, barcodes can be injected into droplets previous to, subsequent to, or concurrently with droplet generation. The delivery of the barcodes to a particular partition allows for the later attribution of the characteristics of the individual biological particle to the particular partition. Barcodes can be delivered, for example on a nucleic acid molecule (e.g., an oligonucleotide), to a partition via any suitable mechanism. Barcoded nucleic acid molecules can be delivered to a partition via a microcapsule. A microcapsule, in some instances, can include a bead.

[1105] In some embodiments, barcoded nucleic acid molecules can be initially associated with the microcapsule and then released from the microcapsule. Release of the barcoded nucleic acid molecules can be passive (e.g., by diffusion out of the microcapsule). In addition or alternatively, release from the microcapsule can be upon application of a stimulus which allows the barcoded nucleic acid molecules to dissociate or to be released from the microcapsule. Such stimulus can disrupt the microcapsule, an interaction that couples the barcoded nucleic acid molecules to or within the microcapsule, or both. Such stimulus can include, for example, a thermal stimulus, photo-stimulus, chemical stimulus (e.g., change in pH or use of a reducing agent(s)), a mechanical stimulus, a radiation stimulus; a biological stimulus (e.g., enzyme), or any combination thereof.

[1106] In some embodiments, one more barcodes (e.g., spatial barcodes, UMIs, or a combination thereof) can be introduced into a partition as part of the analyte. As described previously, barcodes can be bound to the analyte directly, or can form part of a capture probe or analyte capture agent that is hybridized to, conjugated to, or otherwise associated with an analyte, such that when the analyte is introduced into the partition, the barcode(s) are introduced as well.

[1107] FIG. 21 depicts an exemplary workflow, where a sample is contacted with a spatially-barcoded capture probe array and the sample is fixed, stained, and imaged **2101**, as described elsewhere herein. The capture probes can be cleaved from the array **2102** using any method as described herein. The capture probes can diffuse toward the cells by either passive or active migration as described elsewhere herein. The capture probes may then be introduced to the sample **2103** as described elsewhere herein, wherein the capture probe is able to gain entry into the cell in the absence of cell permeabilization, using one of the cell penetrating peptides or lipid delivery systems described herein. The sample can then be optionally imaged in order to confirm probe uptake, via a reporter molecule incorporated within the capture probe **2104**. The sample can then be separated from the array and undergo dissociation **2105**, wherein the sample is separated into single cells or small groups of cells. Once the sample is dissociated, the single cells can be introduced to an oil-in water droplet **2106**, wherein a single cell is combined with reagents within the droplet and processed so that the spatial barcode that penetrated the cell labels the contents of that cell within the droplet. Other cells undergo separately partitioned reactions concurrently. The contents of the droplet is then sequenced **2107** in order to associate a particular cell or cells with a particular spatial location within the sample **2108**.

[1108] As described above, FIG. 16 shows an example of a microfluidic channel structure for partitioning individual analytes (e.g., cells) into discrete partitions. FIGS. 17A and 17C also show other examples of microfluidic channel structures that can be used for delivering beads to droplets.

[1109] A variety of different beads can be incorporated into partitions as described above. In some embodiments, for example, non-barcoded beads can be incorporated into the partitions. For example, where the biological particle (e.g., a cell) that is incorporated into the partitions carries one or more barcodes (e.g., spatial barcode(s), UMI(s), and combinations thereof), the bead can be a non-barcoded bead.

[1110] In some embodiments, a barcode carrying bead can be incorporated into partitions. For example, a nucleic acid molecule, such as an oligonucleotide, can be coupled to a bead by a releasable linkage, such as, for example, a disulfide linker. The same bead can be coupled (e.g., via releasable linkage) to one or more other nucleic acid molecules. The nucleic acid molecule can be or include a barcode. As noted elsewhere herein, the structure of the barcode can include a number of sequence elements.

[1111] The nucleic acid molecule can include a functional domain that can be used in subsequent processing. For example, the functional domain can include one or more of a sequencer specific flow cell attachment sequence (e.g., a P5 sequence for Illumina® sequencing systems) and a sequencing primer sequence (e.g., a R1 primer for Illumina® sequencing systems). The nucleic acid molecule can include a barcode sequence for use in barcoding the sample (e.g., DNA, RNA, protein, etc.). In some cases, the barcode sequence can be bead-specific such that the barcode sequence is common to all nucleic acid molecules coupled to the same bead. Alternatively or in addition, the barcode sequence can be partition-specific such that the barcode sequence is common to all nucleic acid molecules coupled to one or more beads that are partitioned into the same partition. The nucleic acid molecule can include a specific priming sequence, such as an mRNA specific priming sequence (e.g., poly (T) sequence), a targeted priming sequence, and/or a random priming sequence. The nucleic acid molecule can include an anchoring sequence to ensure that the specific priming sequence hybridizes at the sequence end (e.g., of the mRNA). For example, the anchoring sequence can include a random short sequence of nucleotides, such as a 1-mer, 2-mer, 3-mer or longer sequence, which can ensure that a poly(T) segment is more likely to hybridize at the sequence end of the poly(A) tail of the mRNA.

[1112] The nucleic acid molecule can include a unique molecular identifying sequence (e.g., unique molecular identifier (UMI)). In some embodiments, the unique molecular identifying sequence can include from about 5 to about 8 nucleotides. Alternatively, the unique molecular identifying sequence can include less than about 5 or more than about 8 nucleotides. The unique molecular

identifying sequence can be a unique sequence that varies across individual nucleic acid molecules coupled to a single bead.

[1113] In some embodiments, the unique molecular identifying sequence can be a random sequence (e.g., such as a random N-mer sequence). For example, the UMI can provide a unique identifier of the starting mRNA molecule that was captured, in order to allow quantitation of the number of original expressed RNA.

[1114] In general, an individual bead can be coupled to any number of individual nucleic acid molecules, for example, from one to tens to hundreds of thousands or even millions of individual nucleic acid molecules. The respective barcodes for the individual nucleic acid molecules can include both common sequence segments or relatively common sequence segments and variable or unique sequence segments between different individual nucleic acid molecules coupled to the same bead.

[1115] Within any given partition, all of the cDNA transcripts of the individual mRNA molecules can include a common barcode sequence segment. However, the transcripts made from the different mRNA molecules within a given partition can vary at the unique molecular identifying sequence segment (e.g., UMI segment). Beneficially, even following any subsequent amplification of the contents of a given partition, the number of different UMIs can be indicative of the quantity of mRNA originating from a given partition. As noted above, the transcripts can be amplified, cleaned up, and sequenced to identify the sequence of the cDNA transcript of the mRNA, as well as to sequence the barcode segment and the UMI segment. While a poly(T) primer sequence is described, other targeted or random priming sequences can also be used in priming the reverse transcription reaction. Likewise, although described as releasing the barcoded oligonucleotides into the partition, in some cases, the nucleic acid molecules bound to the bead can be used to hybridize and capture the mRNA on the solid phase of the bead, for example, in order to facilitate the separation of the RNA from other cell contents.

[1116] In some embodiments, precursors that include a functional group that is reactive or capable of being activated such that it becomes reactive can be polymerized with other precursors to generate gel beads that include the activated or activatable functional group. The functional group can then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the gel beads. For example, some precursors featuring a carboxylic acid (COOH) group can co-polymerize with other precursors to form a bead that also includes a COOH functional group. In some cases, acrylic acid (a species comprising free COOH groups), acrylamide, and bis(acryloyl) cystamine can be co-polymerized together to generate a bead with free COOH groups. The COOH groups of the bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species comprising an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) comprising a moiety to be linked to the bead.

In some embodiments, a degradable bead can be introduced into a partition, such that the bead degrades within the partition and any associated species (e.g., oligonucleotides) are released within the droplet when the appropriate stimulus is applied. The free species (e.g., oligonucleotides, nucleic acid molecules) can interact with other reagents contained in the partition. For example, a polyacrylamide bead featuring cystamine and linked, via a disulfide bond, to a barcode sequence, can be combined with a reducing agent within a droplet of a water-in-oil emulsion. Within the droplet, the reducing agent can break the various disulfide bonds, resulting in bead degradation and release of the barcode sequence into the aqueous, inner environment of the droplet. In another example, heating of a droplet with a bead-bound barcode sequence in basic solution can also result in bead degradation and release of the attached barcode sequence into the aqueous, inner

environment of the droplet.

[1117] Any suitable number of species (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the species (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration can be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In some cases, the pre-defined concentration of the primer can be limited by the process of producing nucleic acid molecule (e.g., oligonucleotide) bearing beads.

[1118] A degradable bead can include one or more species with a labile bond such that, when the bead/species is exposed to the appropriate stimulus, the bond is broken and the bead degrades. The labile bond can be a chemical bond (e.g., covalent bond, ionic bond) or can be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.). In some embodiments, a cross-linker used to generate a bead can include a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead that includes cystamine cross-linkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

[1119] A degradable bead can be useful in more quickly releasing an attached species (e.g., a nucleic acid molecule, a barcode sequence, a primer, etc.) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species can have greater mobility and accessibility to other species in solution upon degradation of the bead. In some embodiments, a species can also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker can respond to the same stimuli as the degradable bead or the two degradable species can respond to different stimuli. For example, a barcode sequence can be attached, via a disulfide bond, to a polyacrylamide bead comprising cystamine. Upon exposure of the barcoded-bead to a reducing agent, the bead degrades and the barcode sequence is released upon breakage of both the disulfide linkage between the barcode sequence and the bead and the disulfide linkages of the cystamine in the bead.

[1120] As will be appreciated from the above description, while referred to as degradation of a bead, in many embodiments, degradation can refer to the disassociation of a bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species can be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore volumes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore volume due to osmotic swelling of a bead can permit the release of entrained species within the bead. In some embodiments, osmotic shrinking of a bead can cause a bead to better retain an entrained species due to pore volume contraction.

[1121] Numerous chemical triggers can be used to trigger the degradation of beads within partitions. Examples of these chemical changes can include, but are not limited to pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

[1122] In some embodiments, a bead can be formed from materials that include degradable chemical cross-linkers, such as BAC or cystamine. Degradation of such degradable cross-linkers can be accomplished through a number of mechanisms. In some examples, a bead can be contacted with a chemical degrading agent that can induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as dithiothreitol (DTT). Additional examples of reducing agents can include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris (2-carboxyethyl) phosphine (TCEP), or combinations thereof. A reducing agent can degrade the disulfide bonds formed between gel precursors forming the bead, and thus, degrade the bead.

[1123] In certain embodiments, a change in pH of a solution, such as an increase in pH, can trigger

degradation of a bead. In other embodiments, exposure to an aqueous solution, such as water, can trigger hydrolytic degradation, and thus degradation of the bead. In some cases, any combination of stimuli can trigger degradation of a bead. For example, a change in pH can enable a chemical agent (e.g., DTT) to become an effective reducing agent.

[1124] Beads can also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat can cause melting of a bead such that a portion of the bead degrades. In other cases, heat can increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat can also act upon heat-sensitive polymers used as materials to construct beads.

[1125] In addition to beads and analytes, partitions that are formed can include a variety of different reagents and species. For example, when lysis reagents are present within the partitions, the lysis reagents can facilitate the release of analytes within the partition. Examples of lysis agents include bioactive reagents, such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, etc., such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other lysis enzymes available from, e.g., Sigma-Aldrich, Inc. (St. Louis, MO), as well as other commercially available lysis enzymes. Other lysis agents can additionally or alternatively be co-partitioned to cause the release analytes into the partitions. For example, in some cases, surfactant-based lysis solutions can be used to lyse cells, although these can be less desirable for emulsion based systems where the surfactants can interfere with stable emulsions. In some embodiments, lysis solutions can include non-ionic surfactants such as, for example, TritonX-100 and Tween 20. In some embodiments, lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption can also be used in certain embodiments, e.g., non-emulsion based partitioning such as encapsulation of analytes that can be in addition to or in place of droplet partitioning, where any pore volume of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

[1126] Examples of other species that can be co-partitioned with analytes in the partitions include, but are not limited to, DNase and RNase inactivating agents or inhibitors or chelating agents, such as EDTA, and other reagents employed in removing or otherwise reducing negative activity or impact of different cell lysate components on subsequent processing of nucleic acids. Additional reagents can also be co-partitioned, including endonucleases to fragment DNA, DNA polymerase enzymes and dNTPs used to amplify nucleic acid fragments and to attach the barcode molecular tags to the amplified fragments.

[1127] Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers and oligonucleotides, and switch oligonucleotides (also referred to herein as “switch oligos” or “template switching oligonucleotides”) which can be used for template switching. In some embodiments, template switching can be used to increase the length of a cDNA. Template switching can be used to append a predefined nucleic acid sequence to the cDNA. In an example of template switching, cDNA can be generated from reverse transcription of a template, e.g., cellular mRNA, where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., poly(C), to the cDNA in a template independent manner. Switch oligos can include sequences complementary to the additional nucleotides, e.g., poly(G). The additional nucleotides (e.g., poly(C)) on the cDNA can hybridize to the additional nucleotides (e.g., poly(G)) on the switch oligo, whereby the switch oligo can be used by the reverse transcriptase as template to further extend the cDNA.

[1128] Template switching oligonucleotides can include a hybridization region and a template region. The hybridization region can include any sequence capable of hybridizing to the target. In some cases, the hybridization region includes a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases can include 1 G base, 2 G bases, 3 G

bases, 4 G bases, 5 G bases, or more than 5 G bases. The template sequence can include any sequence to be incorporated into the cDNA. In some cases, the template region includes at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. Switch oligos can include deoxyribonucleic acids; ribonucleic acids; modified nucleic acids including 2-Aminopurine, 2,6-Diaminopurine (2-Amino-dA), inverted dT, 5-Methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine), locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' Fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), and combinations of the foregoing.

[1129] In some embodiments, beads that are partitioned with the analyte can include different types of oligonucleotides bound to the bead, where the different types of oligonucleotides bind to different types of analytes. For example, a bead can include one or more first oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a first type of analyte, such as mRNA for example, and one or more second oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a second type of analyte, such as gDNA for example. Partitions can also include lysis agents that aid in releasing nucleic acids from the co-partitioned cell, and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break covalent linkages between the oligonucleotides and the bead, releasing the oligonucleotides into the partition. The released barcoded oligonucleotides (which can also be barcoded) can hybridize with mRNA released from the cell and also with gDNA released from the cell.

[1130] Barcoded constructs thus formed from hybridization can include a first type of construct that includes a sequence corresponding to an original barcode sequence from the bead and a sequence corresponding to a transcript from the cell, and a second type of construct that includes a sequence corresponding to the original barcode sequence from the bead and a sequence corresponding to genomic DNA from the cell. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs can then be sequenced, the sequencing data processed, and the results used to spatially characterize the mRNA and the gDNA from the cell.

[1131] In another example, a partition includes a bead that includes a first type of oligonucleotide (e.g., a first capture probe) with a first barcode sequence, a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript, and a UMI barcode sequence that can uniquely identify a given transcript. The bead also includes a second type of oligonucleotide (e.g., a second capture probe) with a second barcode sequence, a targeted priming sequence that is capable of specifically hybridizing with a third barcoded oligonucleotide (e.g., an analyte capture agent) coupled to an antibody that is bound to the surface of the partitioned cell. The third barcoded oligonucleotide includes a UMI barcode sequence that uniquely identifies the antibody (and thus, the particular cell surface feature to which it is bound).

[1132] In this example, the first and second barcoded oligonucleotides include the same spatial barcode sequence (e.g., the first and second barcode sequences are the same), which permits downstream association of barcoded nucleic acids with the partition. In some embodiments, however, the first and second barcode sequences are different.

[1133] The partition also includes lysis agents that aid in releasing nucleic acids from the cell and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break a covalent linkage between the barcoded oligonucleotides and the bead, releasing them into the partition. The first type of released barcoded oligonucleotide can hybridize with mRNA released from the cell and the second type of released barcoded oligonucleotide can hybridize with the third type of barcoded oligonucleotide, forming barcoded constructs.

[1134] The first type of barcoded construct includes a spatial barcode sequence corresponding to the first barcode sequence from the bead and a sequence corresponding to the UMI barcode

sequence from the first type of oligonucleotide, which identifies cell transcripts. The second type of barcoded construct includes a spatial barcode sequence corresponding to the second barcode sequence from the second type of oligonucleotide, and a UMI barcode sequence corresponding to the third type of oligonucleotide (e.g., the analyte capture agent) and used to identify the cell surface feature. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs are then sequenced, sequencing data processed, and the results used to characterize the mRNA and cell surface feature of the cell.

[1135] The foregoing discussion involves two specific examples of beads with oligonucleotides for analyzing two different analytes within a partition. More generally, beads that are partitioned can have any of the structures described previously, and can include any of the described combinations of oligonucleotides for analysis of two or more (e.g., three or more, four or more, five or more, six or more, eight or more, ten or more, 12 or more, 15 or more, 20 or more, 25 or more, 30 or more, 40 or more, 50 or more) different types of analytes within a partition. Examples of beads with combinations of different types of oligonucleotides (e.g., capture probes) for concurrently analyzing different combinations of analytes within partitions include, but are not limited to: (a) genomic DNA and cell surface features (e.g., using the analyte capture agents described herein); (b) mRNA and a lineage tracing construct; (c) mRNA and cell methylation status; (d) mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq); (e) mRNA and cell surface or intracellular proteins and/or metabolites; (f) a barcoded analyte capture agent (e.g., the MHC multimers described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); and (g) mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein). In some embodiments, a perturbation agent can be a small molecule, an antibody, a drug, an aptamer, a nucleic acid (e.g., miRNA), a physical environmental (e.g., temperature change), or any other known perturbation agents.

(e) Sequencing Analysis

[1136] After analytes from the sample have hybridized or otherwise been associated with capture probes, analyte capture agents, or other barcoded oligonucleotide sequences according to any of the methods described above in connection with the general spatial cell-based analytical methodology, the barcoded constructs that result from hybridization/association are analyzed via sequencing to identify the analytes.

[1137] In some embodiments, where a sample is barcoded directly via hybridization with capture probes or analyte capture agents hybridized, bound, or associated with either the cell surface, or introduced into the cell, as described above, sequencing can be performed on the intact sample. Alternatively, if the barcoded sample has been separated into fragments, cell groups, or individual cells, as described above, sequencing can be performed on individual fragments, cell groups, or cells. For analytes that have been barcoded via partitioning with beads, as described above, individual analytes (e.g., cells, or cellular contents following lysis of cells) can be extracted from the partitions by breaking the partitions, and then analyzed by sequencing to identify the analytes.

[1138] A wide variety of different sequencing methods can be used to analyze barcoded analyte constructs. In general, sequenced polynucleotides can be, for example, nucleic acid molecules such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including variants or derivatives thereof (e.g., single stranded DNA or DNA/RNA hybrids, and nucleic acid molecules with a nucleotide analog).

[1139] Sequencing of polynucleotides can be performed by various commercial systems. More generally, sequencing can be performed using nucleic acid amplification, polymerase chain reaction (PCR) (e.g., digital PCR and droplet digital PCR (ddPCR), quantitative PCR, real time PCR, multiplex PCR, PCR-based singleplex methods, emulsion PCR), and/or isothermal amplification.

[1140] Other examples of methods for sequencing genetic material include, but are not limited to, DNA hybridization methods (e.g., Southern blotting), restriction enzyme digestion methods, Sanger sequencing methods, next-generation sequencing methods (e.g., single-molecule real-time sequencing, nanopore sequencing, and Polony sequencing), ligation methods, and microarray methods. Additional examples of sequencing methods that can be used include targeted sequencing, single molecule real-time sequencing, exon sequencing, electron microscopy-based sequencing, panel sequencing, transistor-mediated sequencing, direct sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, co-amplification at lower denaturation temperature-PCR (COLD-PCR), sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, MS-PET sequencing, and any combinations thereof.

[1141] Sequence analysis of the nucleic acid molecules (including barcoded nucleic acid molecules or derivatives thereof) can be direct or indirect. Thus, the sequence analysis substrate (which can be viewed as the molecule which is subjected to the sequence analysis step or process) can directly be the barcoded nucleic acid molecule or it can be a molecule which is derived therefrom (e.g., a complement thereof). Thus, for example, in the sequence analysis step of a sequencing reaction, the sequencing template can be the barcoded nucleic acid molecule or it can be a molecule derived therefrom. For example, a first and/or second strand DNA molecule can be directly subjected to sequence analysis (e.g., sequencing), i.e., can directly take part in the sequence analysis reaction or process (e.g., the sequencing reaction or sequencing process, or be the molecule which is sequenced or otherwise identified). Alternatively, the barcoded nucleic acid molecule can be subjected to a step of second strand synthesis or amplification before sequence analysis (e.g., sequencing or identification by another technique). The sequence analysis substrate (e.g., template) can thus be an amplicon or a second strand of a barcoded nucleic acid molecule.

[1142] In some embodiments, both strands of a double stranded molecule can be subjected to sequence analysis (e.g., sequenced). In some embodiments, single stranded molecules (e.g., barcoded nucleic acid molecules) can be analyzed (e.g., sequenced). To perform single molecule sequencing, the nucleic acid strand can be modified at the 3' end.

[1143] In some embodiments, massively parallel pyrosequencing techniques can be used for sequencing nucleic acids. In pyrosequencing, the nucleic acid is amplified inside water droplets in an oil solution (emulsion PCR), with each droplet containing a single nucleic acid template attached to a single primer-coated bead that then forms a clonal colony. The sequencing system contains many picolitre-volume wells each containing a single bead and sequencing enzymes. Pyrosequencing uses luciferase to generate light for detection of the individual nucleotides added to the nascent nucleic acid and the combined data are used to generate sequence reads.

[1144] As another example application of pyrosequencing, released PPI can be detected by being immediately converted to adenosine triphosphate (ATP) by ATP sulfurylase, and the level of ATP generated can be detected via luciferase-produced photons, such as described in Ronaghi, et al., *Anal. Biochem.* 242 (1), 84-9 (1996); Ronaghi, *Genome Res.* 11 (1), 3-11 (2001); Ronaghi et al. *Science* 281 (5375), 363 (1998); and U.S. Pat. Nos. 6,210,891, 6,258,568, and 6,274,320, the entire contents of each of which are incorporated herein by reference.

[1145] Massively parallel sequencing techniques can be used for sequencing nucleic acids, as described above. In one embodiment, a massively parallel sequencing technique can be based on reversible dye-terminators. As an example, DNA molecules are first attached to primers on, e.g., a glass or silicon substrate, and amplified so that local clonal colonies are formed (e.g., by bridge amplification). Four types of ddNTPs are added, and non-incorporated nucleotides are washed

away. Unlike pyrosequencing, the DNA is only extended one nucleotide at a time due to a blocking group (e.g., 3' blocking group present on the sugar moiety of the ddNTP). A detector acquires images of the fluorescently labelled nucleotides, and then the dye along with the terminal 3' blocking group is chemically removed from the DNA, as a precursor to a subsequent cycle. This process can be repeated until the required sequence data is obtained.

[1146] In some embodiments, sequencing is performed by detection of hydrogen ions that are released during the polymerization of DNA. A microwell containing a template DNA strand to be sequenced can be flooded with a single type of nucleotide. If the introduced nucleotide is complementary to the leading template nucleotide, it is incorporated into the growing complementary strand. This causes the release of a hydrogen ion that triggers a hypersensitive ion sensor, which indicates that a reaction has occurred. If homopolymer repeats are present in the template sequence, multiple nucleotides will be incorporated in a single cycle. This leads to a corresponding number of released hydrogen ions and a proportionally higher electronic signal.

[1147] In some embodiments, sequencing can be performed in situ. In situ sequencing methods are particularly useful, for example, when the biological sample remains intact after analytes on the sample surface (e.g., cell surface analytes) or within the sample (e.g., intracellular analytes) have been barcoded. In situ sequencing typically involves incorporation of a labeled nucleotide (e.g., fluorescently labeled mononucleotides or dinucleotides) in a sequential, template-dependent manner or hybridization of a labeled primer (e.g., a labeled random hexamer) to a nucleic acid template such that the identities (i.e., nucleotide sequence) of the incorporated nucleotides or labeled primer extension products can be determined, and consequently, the nucleotide sequence of the corresponding template nucleic acid. Aspects of in situ sequencing are described, for example, in Mitra et al., (2003) *Anal. Biochem.* 320, 55-65, and Lee et al., (2014) *Science*, 343 (6177), 1360-1363, the entire contents of each of which are incorporated herein by reference.

[1148] In addition, examples of methods and systems for performing in situ sequencing are described in PCT Patent Application Publication Nos. WO2014/163886, WO2018/045181, WO2018/045186, and in U.S. Pat. Nos. 10,138,509 and 10,179,932, the entire contents of each of which are incorporated herein by reference. Example techniques for in situ sequencing include, but are not limited to, STARmap (described for example in Wang et al., (2018) *Science*, 361 (6499) 5691), MERFISH (described for example in Moffitt, (2016) *Methods in Enzymology*, 572, 1-49), and FISSEQ (described for example in U.S. Patent Application Publication No. 2019/0032121). The entire contents of each of the foregoing references are incorporated herein by reference.

[1149] For analytes that have been barcoded via partitioning, barcoded nucleic acid molecules or derivatives thereof (e.g., barcoded nucleic acid molecules to which one or more functional sequences have been added, or from which one or more features have been removed) can be pooled and processed together for subsequent analysis such as sequencing on high throughput sequencers. Processing with pooling can be implemented using barcode sequences. For example, barcoded nucleic acid molecules of a given partition can have the same barcode, which is different from barcodes of other spatial partitions. Alternatively, barcoded nucleic acid molecules of different partitions can be processed separately for subsequent analysis (e.g., sequencing).

[1150] In some embodiments, where capture probes do not contain a spatial barcode, the spatial barcode can be added after the capture probe captures analytes from a biological sample and before analysis of the analytes. When a spatial barcode is added after an analyte is captured, the barcode can be added after amplification of the analyte (e.g., reverse transcription and polymerase amplification of RNA). In some embodiments, analyte analysis uses direct sequencing of one or more captured analytes, such as direct sequencing of hybridized RNA. In some embodiments, direct sequencing is performed after reverse transcription of hybridized RNA. In some embodiments direct sequencing is performed after amplification of reverse transcription of hybridized RNA.

[1151] In some embodiments, direct sequencing of captured RNA is performed by sequencing-by-

synthesis (SBS). In some embodiments, a sequencing primer is complementary to a sequence in one or more of the domains of a capture probe (e.g., functional domain). In such embodiments, sequencing-by-synthesis can include reverse transcription and/or amplification in order to generate a template sequence (e.g., functional domain) from which a primer sequence can bind.

[1152] SBS can involve hybridizing an appropriate primer, sometimes referred to as a sequencing primer, with the nucleic acid template to be sequenced, extending the primer, and detecting the nucleotides used to extend the primer. Preferably, the nucleic acid used to extend the primer is detected before a further nucleotide is added to the growing nucleic acid chain, thus allowing base-by-base in situ nucleic acid sequencing. The detection of incorporated nucleotides is facilitated by including one or more labelled nucleotides in the primer extension reaction. To allow the hybridization of an appropriate sequencing primer to the nucleic acid template to be sequenced, the nucleic acid template should normally be in a single stranded form. If the nucleic acid templates making up the nucleic acid features are present in a double stranded form these can be processed to provide single stranded nucleic acid templates using methods well known in the art, for example by denaturation, cleavage, etc. The sequencing primers which are hybridized to the nucleic acid template and used for primer extension are preferably short oligonucleotides, for example, 15 to 25 nucleotides in length. The sequencing primers can be greater than 25 nucleotides in length as well. For example, sequencing primers can be about 20 to about 60 nucleotides in length, or more than 60 nucleotides in length. The sequencing primers can be provided in solution or in an immobilized form. Once the sequencing primer has been annealed to the nucleic acid template to be sequenced by subjecting the nucleic acid template and sequencing primer to appropriate conditions, primer extension is carried out, for example using a nucleic acid polymerase and a supply of nucleotides, at least some of which are provided in a labelled form, and conditions suitable for primer extension if a suitable nucleotide is provided.

[1153] Preferably after each primer extension step, a washing step is included in order to remove unincorporated nucleotides which can interfere with subsequent steps. Once the primer extension step has been carried out, the nucleic acid colony is monitored to determine whether a labelled nucleotide has been incorporated into an extended primer. The primer extension step can then be repeated to determine the next and subsequent nucleotides incorporated into an extended primer. If the sequence being determined is unknown, the nucleotides applied to a given colony are usually applied in a chosen order which is then repeated throughout the analysis, for example dATP, dTTP, dCTP, dGTP.

[1154] SBS techniques which can be used are described for example, but not limited to, those in U.S. Patent App. Pub. No. 2007/0166705, U.S. Patent App. Pub. No. 2006/0188901, U.S. Pat. No. 7,057,026, U.S. Patent App. Pub. No. 2006/0240439, U.S. Patent App. Pub. No. 2006/0281109, PCT Patent App. Pub. No. WO 05/065814, U.S. Patent App. Pub. No. 2005/0100900, PCT Patent App. Pub. No. WO 06/064199, PCT Patent App. Pub. No. WO07/010,251, U.S. Patent App. Pub. No. 2012/0270305, U.S. Patent App. Pub. No. 2013/0260372, and U.S. Patent App. Pub. No. 2013/0079232, the entire contents of each of which are incorporated herein by reference.

[1155] In some embodiments, direct sequencing of captured RNA is performed by sequential fluorescence hybridization (e.g., sequencing by hybridization). In some embodiments, a hybridization reaction where RNA is hybridized to a capture probe is performed in situ. In some embodiments, captured RNA is not amplified prior to hybridization with a sequencing probe. In some embodiments, RNA is amplified prior to hybridization with sequencing probes (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, amplification is performed using single-molecule hybridization chain reaction. In some embodiments, amplification is performed using rolling chain amplification.

[1156] Sequential fluorescence hybridization can involve sequential hybridization of probes including degenerate primer sequences and a detectable label. A degenerate primer sequence is a short oligonucleotide sequence which is capable of hybridizing to any nucleic acid fragment

independent of the sequence of said nucleic acid fragment. For example, such a method could include the steps of: (a) providing a mixture including four probes, each of which includes either A, C, G, or T at the 5'-terminus, further including degenerate nucleotide sequence of 5 to 11 nucleotides in length, and further including a functional domain (e.g., fluorescent molecule) that is distinct for probes with A, C, G, or T at the 5'-terminus; (b) associating the probes of step (a) to the target polynucleotide sequences, whose sequence needs will be determined by this method; (c) measuring the activities of the four functional domains and recording the relative spatial location of the activities; (d) removing the reagents from steps (a)-(b) from the target polynucleotide sequences; and repeating steps (a)-(d) for n cycles, until the nucleotide sequence of the spatial domain for each bead is determined, with modification that the oligonucleotides used in step (a) are complementary to part of the target polynucleotide sequences and the positions 1 through n flanking the part of the sequences. Because the barcode sequences are different, in some embodiments, these additional flanking sequences are degenerate sequences. The fluorescent signal from each spot on the array for cycles 1 through n can be used to determine the sequence of the target polynucleotide sequences.

[1157] In some embodiments, direct sequencing of captured RNA using sequential fluorescence hybridization is performed in vitro. In some embodiments, captured RNA is amplified prior to hybridization with a sequencing probe (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, a capture probe containing captured RNA is exposed to the sequencing probe targeting coding regions of RNA. In some embodiments, one or more sequencing probes are targeted to each coding region. In some embodiments, the sequencing probe is designed to hybridize with sequencing reagents (e.g., a dye-labeled readout oligonucleotides). A sequencing probe can then hybridize with sequencing reagents. In some embodiments, output from the sequencing reaction is imaged. In some embodiments, a specific sequence of cDNA is resolved from an image of a sequencing reaction. In some embodiments, reverse transcription of captured RNA is performed prior to hybridization to the sequencing probe. In some embodiments, the sequencing probe is designed to target complementary sequences of the coding regions of RNA (e.g., targeting cDNA).

[1158] In some embodiments, a captured RNA is directly sequenced using a nanopore-based method. In some embodiments, direct sequencing is performed using nanopore direct RNA sequencing in which captured RNA is translocated through a nanopore. A nanopore current can be recorded and converted into a base sequence. In some embodiments, captured RNA remains attached to a substrate during nanopore sequencing. In some embodiments, captured RNA is released from the substrate prior to nanopore sequencing. In some embodiments, where the analyte of interest is a protein, direct sequencing of the protein can be performed using nanopore-based methods. Examples of nanopore-based sequencing methods that can be used are described in Deamer et al., *Trends Biotechnol.* 18, 14 7-151 (2000); Deamer et al., *Acc. Chem. Res.* 35:817-825 (2002); Li et al., *Nat. Mater.* 2:611-615 (2003); Soni et al., *Clin. Chem.* 53, 1996-2001 (2007); Healy et al., *Nanomed.* 2, 459-481 (2007); Cockroft et al., *J. Am. Chem. Soc.* 130, 818-820 (2008); and in U.S. Pat. No. 7,001,792. The entire contents of each of the foregoing references are incorporated herein by reference.

[1159] In some embodiments, direct sequencing of captured RNA is performed using single molecule sequencing by ligation. Such techniques utilize DNA ligase to incorporate oligonucleotides and identify the incorporation of such oligonucleotides. The oligonucleotides typically have different labels that are correlated with the identity of a particular nucleotide in a sequence to which the oligonucleotides hybridize. Aspects and features involved in sequencing by ligation are described, for example, in Shendure et al. *Science* (2005), 309:1728-1732, and in U.S. Pat. Nos. 5,599,675; 5,750,341; 6,969,488; 6,172,218; and 6,306,597, the entire contents of each of which are incorporated herein by reference.

[1160] In some embodiments, nucleic acid hybridization can be used for sequencing. These

methods utilize labeled nucleic acid decoder probes that are complementary to at least a portion of a barcode sequence. Multiplex decoding can be performed with pools of many different probes with distinguishable labels. Non-limiting examples of nucleic acid hybridization sequencing are described for example in U.S. Pat. No. 8,460,865, and in Gunderson et al., *Genome Research* 14:870-877 (2004), the entire contents of each of which are incorporated herein by reference.

[1161] In some embodiments, commercial high-throughput digital sequencing techniques can be used to analyze barcode sequences, in which DNA templates are prepared for sequencing not one at a time, but in a bulk process, and where many sequences are read out preferably in parallel, or alternatively using an ultra-high throughput serial process that itself may be parallelized. Examples of such techniques include Illumina® sequencing (e.g., flow cell-based sequencing techniques), sequencing by synthesis using modified nucleotides (such as commercialized in TruSeq™ and HiSeq™ technology by Illumina, Inc., San Diego, CA), HeliScope™ by Helicos Biosciences Corporation, Cambridge, MA, and PacBio RS by Pacific Biosciences of California, Inc., Menlo Park, CA), sequencing by ion detection technologies (Ion Torrent, Inc., South San Francisco, CA), and sequencing of DNA nanoballs (Complete Genomics, Inc., Mountain View, CA).

[1162] In some embodiments, detection of a proton released upon incorporation of a nucleotide into an extension product can be used in the methods described herein. For example, the sequencing methods and systems described in U.S. Patent Application Publication Nos. 2009/0026082, 2009/0127589, 2010/0137143, and 2010/0282617, can be used to directly sequence barcodes. The entire contents of each of the foregoing references are incorporated herein by reference.

[1163] In some embodiments, real-time monitoring of DNA polymerase activity can be used during sequencing. For example, nucleotide incorporations can be detected through fluorescence resonance energy transfer (FRET), as described for example in Levene et al., *Science* (2003), 299, 682-686, Lundquist et al., *Opt. Lett.* (2008), 33, 1026-1028, and Korlach et al., *Proc. Natl. Acad. Sci. USA* (2008), 105, 1176-1181. The entire contents of each of the foregoing references are herein incorporated by reference.

IV. Multiplexing

(a) Multiplexing Generally

[1164] In various embodiments of spatial analysis as described herein, features can include different types of capture probes for analyzing both intrinsic and extrinsic information for individual cells. For example, a feature can include one or more of the following: 1) a capture probe featuring a capture domain that binds to one or more endogenous nucleic acids in the cell; 2) a capture probe featuring a capture domain that binds to one or more exogenous nucleic acids in the cell (e.g., nucleic acids from a microorganism (e.g., a virus, a bacterium)) that infects the cell, nucleic acids introduced into the cell (e.g., such as plasmids or nucleic acid derived therefrom), nucleic acids for gene editing (e.g., CRISPR-related RNA such as crRNA, guide RNA); 3) a capture probe featuring a capture domain that binds to an analyte capture agent (e.g., an antibody coupled to a oligonucleotide that includes a capture agent barcode domain having an analyte capture sequence that binds the capture domain), and 4) a capture moiety featuring a domain that binds to a protein (e.g., an exogenous protein expressed in the cell, a protein from a microorganism (e.g., a virus, a bacterium)) that infects the cell, or a binding partner for a protein of the cell (e.g., an antigen for an immune cell receptor).

[1165] In some embodiments of any of the spatial analysis methods as described herein, spatial profiling includes concurrent analysis of two different types of analytes. A feature can be a gel bead, which is coupled (e.g., reversibly coupled) to one or more capture probes. The capture probes can include a spatial barcode sequence and a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript. The capture probe can also include a UMI sequence that can uniquely identify a given transcript. The capture probe can also include a spatial barcode sequence and a random N-mer priming sequence that is capable of randomly hybridizing with gDNA. In this configuration, capture probes can include the same spatial barcode sequence, which permits

association of downstream sequencing reads with the feature.

[1166] In some embodiments of any of the spatial analysis methods as described herein, a feature can be a gel bead, which is coupled (e.g., reversibly coupled) to capture probes. The Capture probe can include a spatial barcode sequence and a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript. The capture probe can also include a UMI sequence that can uniquely identify a given transcript. The capture probe can include a spatial barcode sequence and a capture domain that is capable of specifically hybridizing with an analyte capture agent. The analyte capture agent can include an oligonucleotide that includes an analyte capture sequence that interacts with the capture domain coupled to the feature. The oligonucleotide of the analyte capture agent can be coupled to an antibody that is bound to the surface of a cell. The oligonucleotide includes a barcode sequence (e.g., an analyte binding moiety barcode) that uniquely identifies the antibody (and thus, the particular cell surface feature to which it is bound). In this configuration, the capture probes include the same spatial barcode sequence, which permit downstream association of barcoded nucleic acids with the location on the spatial array. In some embodiments of any of the spatial profiling methods described herein, the analyte capture agents can be produced by any suitable route, including via example coupling schemes described elsewhere herein.

[1167] In some embodiments of any of the spatial analysis methods described herein, other combinations of two or more biological analytes that can be concurrently measured include, without limitation: (a) genomic DNA and cell surface features (e.g., via analyte capture agents that bind to a cell surface feature), (b) mRNA and a lineage tracing construct, (c) mRNA and cell methylation status, (d) mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), (e) mRNA and cell surface or intracellular proteins and/or metabolites, (f) mRNA and chromatin (spatial organization of chromatin in a cell), (g) an analyte capture agent (e.g., any of the MHC multimers described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), (h) mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein), (i) genomic DNA and a perturbation agent, (j) an analyte capture agent and a perturbation agents, (k) accessible chromatin and a perturbation agent, (l) chromatin (e.g., spatial organization of chromatin in a cell) and a perturbation agent, and (m) cell surface or intracellular proteins and/or metabolites and a perturbation agent (e.g., any of the perturbation agents described herein), or any combination thereof.

[1168] In some embodiments of any of the spatial analysis methods described herein, the first analyte can include a nucleic acid molecule with a nucleic acid sequence (e.g., mRNA, complementary DNA derived from reverse transcription of mRNA) encoding at least a portion of a V(D)J sequence of an immune cell receptor (e.g., a TCR or BCR). In some embodiments, the nucleic acid molecule with a nucleic acid sequence encoding at least a portion of a V(D)J sequence of an immune cell receptor is cDNA first generated from reverse transcription of the corresponding mRNA, using a poly(T) containing primer. The cDNA that is generated can then be barcoded using a primer, featuring a spatial barcode sequence (and optionally, a UMI sequence) that hybridizes with at least a portion of the cDNA that is generated. In some embodiments, a template switching oligonucleotide in conjunction a terminal transferase or a reverse transcriptase having terminal transferase activity can be employed to generate a priming region on the cDNA to which a barcoded primer can hybridize during cDNA generation. Terminal transferase activity can, for example, add a poly(C) tail to a 3' end of the cDNA such that the template switching oligonucleotide can bind via a poly(G) priming sequence and the 3' end of the cDNA can be further extended. The original mRNA template and template switching oligonucleotide can then be denatured from the cDNA and the barcoded primer comprising a sequence complementary to at least a portion of the generated priming region on the cDNA can then hybridize with the cDNA and a barcoded construct comprising the barcode sequence (and any optional UMI sequence) and a

complement of the cDNA generated. Additional methods and compositions suitable for barcoding cDNA generated from mRNA transcripts including those encoding V(D)J regions of an immune cell receptor and/or barcoding methods and composition including a template switch oligonucleotide are described, for example, in PCT Patent Application Publication No. WO 2018/075693, and in U.S. Patent Application Publication No. 2018/0105808, the entire contents of each of which are incorporated herein by reference.

[1169] In some embodiments, V(D)J analysis can be performed using methods similar to those described herein. For example, V(D)J analysis can be completed with the use of one or more analyte capture agents that bind to particular surface features of immune cells and are associated with barcode sequences (e.g., analyte binding moiety barcodes). The one or more analyte capture agents can include an MHC or MHC multimer. A barcoded oligonucleotide coupled to a bead that can be used for V(D)J analysis. The oligonucleotide is coupled to a bead by a releasable linkage, such as a disulfide linker. The oligonucleotide can include functional sequences that are useful for subsequent processing, such as functional sequence, which can include a sequencer specific flow cell attachment sequence, e.g., a P5 sequence, as well as functional sequence, which can include sequencing primer sequences, e.g., a R1 primer binding site. In some embodiments, the sequence can include a P7 sequence and a R2 primer binding site. A barcode sequence can be included within the structure for use in barcoding the template polynucleotide. The functional sequences can be selected for compatibility with a variety of different sequencing systems, e.g., Ion Torrent Proton or PGM, Illumina sequencing instruments, etc., and the requirements thereof. In some embodiments, the barcode sequence, functional sequences (e.g., flow cell attachment sequence) and additional sequences (e.g., sequencing primer sequences) can be common to all of the oligonucleotides attached to a given bead. The barcoded oligonucleotide can also include a sequence to facilitate template switching (e.g., a poly(G) sequence). In some embodiments, the additional sequence provides a unique molecular identifier (UMI) sequence segment, as described elsewhere herein.

[1170] In an exemplary method of cellular polynucleotide analysis using a barcode oligonucleotide, a cell is co-partitioned along with a bead bearing a barcoded oligonucleotide and additional reagents such as a reverse transcriptase, primers, oligonucleotides (e.g., template switching oligonucleotides), dNTPs, and a reducing agent into a partition (e.g., a droplet in an emulsion). Within the partition, the cell can be lysed to yield a plurality of template polynucleotides (e.g., DNA such as genomic DNA, RNA such as mRNA, etc.).

[1171] A reaction mixture featuring a template polynucleotide from a cell and (i) the primer having a sequence towards a 3' end that hybridizes to the template polynucleotide (e.g., poly(T)) and (ii) a template switching oligonucleotide that includes a first oligonucleotide towards a 5' end can be subjected to an amplification reaction to yield a first amplification product. In some embodiments, the template polynucleotide is an mRNA with a poly(A) tail and the primer that hybridizes to the template polynucleotide includes a poly(T) sequence towards a 3' end, which is complementary to the poly(A) segment. The first oligonucleotide can include at least one of an adaptor sequence, a barcode sequence, a unique molecular identifier (UMI) sequence, a primer binding site, and a sequencing primer binding site or any combination thereof. In some cases, a first oligonucleotide is a sequence that can be common to all partitions of a plurality of partitions. For example, the first oligonucleotide can include a flow cell attachment sequence, an amplification primer binding site, or a sequencing primer binding site and the first amplification reaction facilitates the attachment the oligonucleotide to the template polynucleotide from the cell. In some embodiments, the first oligonucleotide includes a primer binding site. In some embodiments, the first oligonucleotide includes a sequencing primer binding site.

[1172] The sequence towards a 3' end (e.g., poly(T)) of the primer hybridizes to the template polynucleotide. In a first amplification reaction, extension reaction reagents, e.g., reverse transcriptase, nucleoside triphosphates, co-factors (e.g., Mg.sup.2+ or Mn.sup.2+), that are also co-partitioned, can extend the primer sequence using the cell's nucleic acid as a template, to produce a

transcript, e.g., cDNA, having a fragment complementary to the strand of the cell's nucleic acid to which the primer annealed. In some embodiments, the reverse transcriptase has terminal transferase activity and the reverse transcriptase adds additional nucleotides, e.g., poly(C), to the cDNA in a template independent manner.

[1173] The template switching oligonucleotide, for example a template switching oligonucleotide which includes a poly(G) sequence, can hybridize to the cDNA and facilitate template switching in the first amplification reaction. The transcript, therefore, can include the sequence of the primer, a sequence complementary to the template polynucleotide from the cell, and a sequence complementary to the template switching oligonucleotide.

[1174] In some embodiments of any of the spatial analysis methods described herein, subsequent to the first amplification reaction, the first amplification product or transcript can be subjected to a second amplification reaction to generate a second amplification product. In some embodiments, additional sequences (e.g., functional sequences such as flow cell attachment sequence, sequencing primer binding sequences, barcode sequences, etc.) are attached. The first and second amplification reactions can be performed in the same volume, such as for example in a droplet. In some embodiments, the first amplification product is subjected to a second amplification reaction in the presence of a barcoded oligonucleotide to generate a second amplification product having a barcode sequence. The barcode sequence can be unique to a partition, that is, each partition can have a unique barcode sequence. The barcoded oligonucleotide can include a sequence of at least a segment of the template switching oligonucleotide and at least a second oligonucleotide. The segment of the template switching oligonucleotide on the barcoded oligonucleotide can facilitate hybridization of the barcoded oligonucleotide to the transcript, e.g., cDNA, to facilitate the generation of a second amplification product. In addition to a barcode sequence, the barcoded oligonucleotide can include a second oligonucleotide such as at least one of an adaptor sequence, a unique molecular identifier (UMI) sequence, a primer binding site, and a sequencing primer binding site, or any combination thereof.

[1175] In some embodiments of any of the spatial analysis methods described herein, the second amplification reaction uses the first amplification product as a template and the barcoded oligonucleotide as a primer. In some embodiments, the segment of the template switching oligonucleotide on the barcoded oligonucleotide can hybridize to the portion of the cDNA or complementary fragment having a sequence complementary to the template switching oligonucleotide or that which was copied from the template switching oligonucleotide. In the second amplification reaction, extension reaction reagents, e.g., polymerase, nucleoside triphosphates, co-factors (e.g., Mg^{sup.2+} or Mn^{sup.2+}), that are also co-partitioned, can extend the primer sequence using the first amplification product as template. The second amplification product can include a second oligonucleotide, a sequence of a segment of the template polynucleotide (e.g., mRNA), and a sequence complementary to the primer.

[1176] In some embodiments of any of the spatial analysis methods described herein, the second amplification product uses the barcoded oligonucleotide as a template and at least a portion of the first amplification product as a primer. The segment of the first amplification product (e.g., cDNA) having a sequence complementary to the template switching oligonucleotide can hybridize to the segment of the barcoded oligonucleotide comprising a sequence of at least a segment of the template switching oligonucleotide. In the second amplification reaction, extension reaction reagents, e.g., polymerase, nucleoside triphosphates, co-factors (e.g., Mg^{sup.2+} or Mn^{sup.2+}), that are also co-partitioned, can extend the primer sequence (e.g., first amplification product) using the barcoded oligonucleotide as template. The second amplification product can include the sequence of the primer, a sequence which is complementary to the sequence of the template polynucleotide (e.g., mRNA), and a sequence complementary to the second oligonucleotide.

[1177] In some embodiments of any of the spatial analysis methods described herein, three or more classes of biological analytes can be concurrently measured. For example, a feature can include

capture probes that can participate in an assay of at least three different types of analytes via three different capture domains. A bead can be coupled to a barcoded oligonucleotide that includes a capture domain that includes a poly(T) priming sequence for mRNA analysis; a barcoded oligonucleotide that includes a capture domain that includes a random N-mer priming sequence for gDNA analysis; and a barcoded oligonucleotide that includes a capture domain that can specifically bind an analyte capture agent (e.g., an antibody with a spatial barcode), via its analyte capture sequence.

[1178] In some embodiments of any of the spatial analysis methods described herein, other combinations of three or more biological analytes that can be concurrently measured include, without limitation: (a) mRNA, a lineage tracing construct, and cell surface and/or intracellular proteins and/or metabolites; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), and cell surface and/or intracellular proteins and/or metabolites; (c) mRNA, genomic DNA, and a perturbation reagent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (d) mRNA, accessible chromatin, and a perturbation reagent; (e) mRNA, an analyte capture agent (e.g., any of the MHC multimers described herein), and a perturbation reagent; (f) mRNA, cell surface and/or intracellular proteins and/or metabolites, and a perturbation agent; (g) mRNA, a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), and a perturbation reagent; (h) mRNA, an analyte capture agent, and a V(D)J sequence of an immune cell receptor; (i) cell surface and/or intracellular proteins and/or metabolites, an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor; (j) methylation status, mRNA, and cell surface and/or intracellular proteins and/or metabolites; (k) mRNA, chromatin (e.g., spatial organization of chromatin in a cell), and a perturbation reagent; (l) a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell); and a perturbation reagent; and (m) mRNA, a V(D)J sequence of an immune cell receptor, and chromatin (e.g., spatial organization of chromatin in a cell), or any combination thereof.

[1179] In some embodiments of any of the spatial analysis methods described herein, four or more classes biological analytes can be concurrently measured. A feature can be a bead that is coupled to barcoded primers that can each participate in an assay of a different type of analyte. The feature is coupled (e.g., reversibly coupled) to a capture probe that includes a capture domain that includes a poly(T) priming sequence for mRNA analysis and is also coupled (e.g., reversibly coupled) to a capture probe that includes a capture domain that includes a random N-mer priming sequence for gDNA analysis. Moreover, the feature is also coupled (e.g., reversibly coupled) to a capture probe that binds an analyte capture sequence of an analyte capture agent via its capture domain. The feature can also be coupled (e.g., reversibly coupled) to a capture probe that can specifically bind a nucleic acid molecule that can function as a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein), via its capture domain.

[1180] In some embodiments of any of the spatial analysis methods described herein, each of the various spatially-barcoded capture probes present at a given feature or on a given bead include the same spatial barcode sequence. In some embodiments, each barcoded capture probe can be released from the feature in a manner suitable for analysis of its respective analyte. For example, barcoded constructs A, B, C and D can be generated as described elsewhere herein and analyzed. Barcoded construct A can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a DNA sequence corresponding to a target mRNA. Barcoded construct B can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a sequence corresponding to genomic DNA. Barcoded construct C can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a sequence corresponding to barcode sequence associated with an analyte capture agent (e.g., an analyte binding moiety barcode). Barcoded construct D can include a sequence corresponding to the

barcode sequence from the bead (e.g., a spatial barcode) and a sequence corresponding to a CRISPR nucleic acid (which, in some embodiments, also includes a barcode sequence). Each construct can be analyzed (e.g., via any of a variety of sequencing methods) and the results can be associated with the given cell from which the various analytes originated. Barcoded (or even non-barcoded) constructs can be tailored for analyses of any given analyte associated with a nucleic acid and capable of binding with such a construct.

[1181] In some embodiments of any of the spatial analysis methods described herein, other combinations of four or more biological analytes that can be concurrently measured include, without limitation: (a) mRNA, a lineage tracing construct, cell surface and/or intracellular proteins and/or metabolites, and gDNA; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), cell surface and/or intracellular proteins and/or metabolites, and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (c) mRNA, cell surface and/or intracellular proteins and/or metabolites, an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); (d) mRNA, genomic DNA, a perturbation reagent, and accessible chromatin; (e) mRNA, cell surface and/or intracellular proteins and/or metabolites, an analyte capture agent (e.g., the MHC multimers described herein), and a perturbation reagent; (f) mRNA, cell surface and/or intracellular proteins and/or metabolites, a perturbation reagent, and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); (g) mRNA, a perturbation reagent, an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); (h) mRNA, chromatin (e.g., spatial organization of chromatin in a cell), and a perturbation reagent; (i) a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell); and a perturbation reagent; (j) mRNA, a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell), and genomic DNA; (k) mRNA, a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell), and a perturbation reagent, or any combination thereof.

(b) Construction of Spatial Arrays for Multi-Analyte Analysis

[1182] This disclosure also provides methods and materials for constructing a spatial array capable of multi-analyte analysis. In some embodiments, a spatial array includes a plurality of features on a substrate where one or more members of the plurality of features include a plurality of oligonucleotides having a first type functional sequence and oligonucleotides having a second, different type of functional sequence. In some embodiments, a feature can include oligonucleotides with two types of functional sequences. A feature can be coupled to oligonucleotides comprising a TruSeq functional sequence and also to oligonucleotides comprising a Nextera functional sequence. In some embodiments, a functional sequence can include a sequencer specific flow cell attachment sequence, e.g., a P5 sequence, as well as functional sequence, which can include sequencing primer sequences, e.g., a R1 primer binding site. In some embodiments, one or more members of the plurality of features comprises both types of functional sequences. In some embodiments, one or more members of the plurality features includes a first type of functional sequence. In some embodiments, one or more members of the plurality of features includes a second type of functional sequence. In some embodiments, an additional oligonucleotide can be added to the functional sequence to generate a full oligonucleotide where the full oligonucleotide includes a spatial barcode sequence, an optional UMI sequence, a priming sequence, and a capture domain. Attachment of these sequences can be via ligation (including via splint ligation as is described in U.S. Patent Application Publication No. 20140378345, the entire contents of which are incorporated herein by reference), or any other suitable route. As discussed herein, oligonucleotides can be hybridized with splint sequences that can be helpful in constructing complete full oligonucleotides (e.g., oligonucleotides that are capable of spatial analysis).

[1183] In some embodiments, the oligonucleotides that hybridize to the functional sequences (e.g.,

TruSeq and Nextera) located on the features include capture domains capable of capturing different types of analytes (e.g., mRNA, genomic DNA, cell surface proteins, or accessible chromatin). In some examples, oligonucleotides that can bind to the TruSeq functional sequences can include capture domains that include poly(T) capture sequences. In addition to the poly(T) capture sequences, the oligonucleotides that can bind the TruSeq functional groups can also include a capture domain that includes a random N-mer sequence for capturing genomic DNA (e.g., or any other sequence or domain as described herein capable of capturing any of the biological analytes described herein). In such cases, the spatial arrays can be constructed by applying ratios of TruSeq-poly(T) and TruSeq-N-mer oligonucleotides to the features comprising the functional TruSeq sequences. This can produce spatial arrays where a portion of the oligonucleotides can capture mRNA and a different portion of oligonucleotides can capture genomic DNA. In some embodiments, one or more members of a plurality of features include both TruSeq and Nextera functional sequences. In such cases, a feature including both types of functional sequences is capable of binding oligonucleotides specific to each functional sequence. For example, an oligonucleotide capable of binding to a TruSeq functional sequence could be used to deliver an oligonucleotide including a poly(T) capture domain and an oligonucleotide capable of binding to a Nextera functional sequence could be used to deliver an oligonucleotide including an N-mer capture domain for capturing genomic DNA. It will be appreciated by a person of ordinary skill in the art that any combination of capture domains (e.g., capture domains having any of the variety of capture sequences described herein capable of binding to any of the different types of analytes as described herein) could be combined with oligonucleotides capable of binding to TruSeq and Nextera functional sequences to construct a spatial array.

[1184] In some embodiments, an oligonucleotide that includes a capture domain (e.g., an oligonucleotide capable of coupling to an analyte) or an analyte capture agent can include an oligonucleotide sequence that is capable of binding or ligating to an assay primer. The adapter can allow the capture probe or the analyte capture agent to be attached to any suitable assay primers and used in any suitable assays. The assay primer can include a priming region and a sequence that is capable of binding or ligating to the adapter. In some embodiments, the adapter can be a non-specific primer (e.g., a 5' overhang) and the assay primer can include a 3' overhang that can be ligated to the 5' overhang. The priming region on the assay primer can be any primer described herein, e.g., a poly(T) primer, a random N-mer primer, a target-specific primer, or an analyte capture agent capture sequence.

[1185] In some examples, an oligonucleotide can include an adapter, e.g., a 5' overhang with 10 nucleotides. The adapter can be ligated to assay primers, each of which includes a 3' overhang with 10 nucleotides that complementary to the 5' overhang of the adapter. The capture probe can be used in any assay by attaching to the assay primer designed for that assay.

[1186] Adapters and assay primers can be used to allow the capture probe or the analyte capture agent to be attached to any suitable assay primers and used in any suitable assays. A capture probe that includes a spatial barcode can be attached to a bead that includes a poly(dT) sequence. A capture probe including a spatial barcode and a poly(T) sequence can be used to assay multiple biological analytes as generally described herein (e.g., the biological analyte includes a poly(A) sequence or is coupled to or otherwise is associated with an analyte capture agent comprising a poly(A) sequence as the analyte capture sequence).

[1187] A splint oligonucleotide with a poly(A) sequence can be used to facilitate coupling to a capture probe that includes a spatial barcode and a second sequence that facilitates coupling with an assay primer. Assay primers include a sequence complementary to the splint oligo second sequence and an assay-specific sequence that determines assay primer functionality (e.g., a poly(T) primer, a random N-mer primer, a target-specific primer, or an analyte capture agent capture sequence as described herein).

[1188] In some embodiments of any of the spatial profiling methods described herein, a feature can

include a capture probe that includes a spatial barcode comprising a switch oligonucleotide, e.g., with a 3' end 3rG. For example, a feature (e.g., a gel bead) with a spatial barcode functionalized with a 3rG sequence can be used that enables template switching (e.g., reverse transcriptase template switching), but is not specific for any particular assay. In some embodiments, the assay primers added to the reaction can determine which type of analytes are analyzed. For example, the assay primers can include binding domains capable of binding to target biological analytes (e.g., poly(T) for mRNA, N-mer for genomic DNA, etc.). A capture probe (e.g., an oligonucleotide capable of spatial profiling) can be generated by using a reverse transcriptase enzyme/polymerase to extend, which is followed by template switching onto the barcoded adapter oligonucleotide to incorporate the barcode and other functional sequences. In some embodiments, the assay primers include capture domains capable of binding to a poly(T) sequence for mRNA analysis, random primers for genomic DNA analysis, or a capture sequence that can bind a nucleic acid molecule coupled to an analyte binding moiety (e.g., an analyte capture sequence of an analyte capture agent) or a nucleic acid molecule that can function in as a perturbation reagent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein).

V. Systems for Sample Analysis

[1189] The methods described above for analyzing biological samples can be implemented using a variety of hardware components. In this section, examples of such components are described. However, it should be understood that in general, the various steps and techniques discussed herein can be performed using a variety of different devices and system components, not all of which are expressly set forth.

[1190] FIG. 22A is a schematic diagram showing an example sample handling apparatus **2200**. Sample handling apparatus **2200** includes a sample chamber **2202** that, when closed or sealed, is fluid-tight. Within chamber **2202**, a first holder **2204** holds a first substrate **2206** on which a sample **2208** is positioned. Sample chamber **2202** also includes a second holder **2210** that holds a second substrate **2212** with an array of features **2214**, as described above.

[1191] A fluid reservoir **2216** is connected to the interior volume of sample chamber **2202** via a fluid inlet **2218**. Fluid outlet **2220** is also connected to the interior volume of sample chamber **2202**, and to valve **2222**. In turn, valve **2222** is connected to waste reservoir **2224** and, optionally, to analysis apparatus **2226**. A control unit **2228** is electrically connected to second holder **2210**, to valve **2222**, to waste reservoir **2224**, and to fluid reservoir **2216**.

[1192] During operation of apparatus **2200**, any of the reagents, solutions, and other biochemical components described above can be delivered into sample chamber **2202** from fluid reservoir **2216** via fluid inlet **2218**. Control unit **2228**, connected to fluid reservoir **2216**, can control the delivery of reagents, solutions, and components, and adjust the volumes and flow rates according to programmed analytical protocols for various sample types and analysis procedures. In some embodiments, fluid reservoir **2216** includes a pump, which can be controlled by control unit **2228**, to facilitate delivery of substances into sample chamber **2202**.

[1193] In certain embodiments, fluid reservoir **2216** includes a plurality of chambers, each of which is connected to fluid inlet **2218** via a manifold (not shown). Control unit **2228** can selectively deliver substances from any one or more of the multiple chambers into sample chamber **2202** by adjusting the manifold to ensure that the selected chambers are fluidically connected to fluid inlet **2218**.

[1194] In general, control unit **2228** can be configured to introduce substances from fluid reservoir **2216** into sample chamber **2202** before, after, or both before and after, sample **2208** on first substrate **2206** has interacted with the array of features **2214** on first substrate **2212**. Many examples of such substances have been described previously. Examples of such substances include, but are not limited to, permeabilizing agents, buffers, fixatives, staining solutions, washing solutions, and solutions of various biological reagents (e.g., enzymes, peptides, oligonucleotides,

primers).

[1195] To initiate interaction between sample **2208** and feature array **2214**, the sample and array are brought into spatial proximity. To facilitate this step, second holder **2210**—under the control of control unit **2228**—can translate second substrate **2212** in any of the x-, y-, and z-coordinate directions. In particular, control unit **2228** can direct second holder **2210** to translate second substrate **2212** in the z-direction so that sample **2208** contacts, or nearly contacts, feature array **2214**.

[1196] In some embodiments, apparatus **2200** can optionally include an alignment sub-system **2230**, which can be electrically connected to control unit **2228**. Alignment sub-system **2230** functions to ensure that sample **2208** and feature array **2214** are aligned in the x-y plane prior to translating second substrate **2212** in the z-direction so that sample **2208** contacts, or nearly contacts, feature array **2214**.

[1197] Alignment sub-system **2230** can be implemented in a variety of ways. In some embodiments, for example, alignment sub-system **2230** includes an imaging unit that obtains one or more images showing fiducial markings on first substrate **2206** and/or second substrate **2212**. Control unit **2218** analyzes the image(s) to determine appropriate translations of second substrate **2212** in the x- and/or y-coordinate directions to ensure that sample **2208** and feature array **2214** are aligned prior to translation in the z-coordinate direction.

[1198] In certain embodiments, control unit **2228** can optionally regulate the removal of substances from sample chamber **2202**. For example, control unit **2228** can selectively adjust valve **2222** so that substances introduced into sample chamber **2202** from fluid reservoir **2216** are directed into waste reservoir **2224**. In some embodiments, waste reservoir **2224** can include a reduced-pressure source (not shown) electrically connected to control unit **2228**. Control unit **2228** can adjust the fluid pressure in fluid outlet **2220** to control the rate at which fluids are removed from sample chamber **2202** into waste reservoir **2224**.

[1199] In some embodiments, analytes from sample **2208** or from feature array **2214** can be selectively delivered to analysis apparatus **2226** via suitable adjustment of valve **2222** by control unit **2228**. As described above, in some embodiments, analysis apparatus **2226** includes a reduced-pressure source (not shown) electrically connected to control unit **2228**, so that control unit **2228** can adjust the rate at which analytes are delivered to analysis apparatus **2226**. As such, fluid outlet **2220** effectively functions as an analyte collector, while analysis of the analytes is performed by analysis apparatus **2226**. It should be noted that not all of the workflows and methods described herein are implemented via analysis apparatus **2226**. For example, in some embodiments, analytes that are captured by feature array **2214** remain bound to the array (i.e., are not cleaved from the array), and feature array **2214** is directly analyzed to identify specifically-bound sample components.

[1200] In addition to the components described above, apparatus **2200** can optionally include other features as well. In some embodiments, for example, sample chamber **2202** includes a heating sub-system **2232** electrically connected to control unit **2228**. Control unit **2228** can activate heating sub-system **2232** to heat sample **2208** and/or feature array **2214**, which can help to facilitate certain steps of the methods described herein.

[1201] In certain embodiments, sample chamber **2202** includes an electrode **2234** electrically connected to control unit **2228**. Control unit **2228** can optionally activate electrode **2234**, thereby establishing an electric field between the first and second substrates. Such fields can be used, for example, to facilitate migration of analytes from sample **2208** toward feature array **2214**.

[1202] In some of the methods described herein, one or more images of a sample and/or a feature array are acquired. Imaging apparatus that is used to obtain such images can generally be implemented in a variety of ways. FIG. **22B** shows one example of an imaging apparatus **2250**. Imaging apparatus **2250** includes a light source **2252**, light conditioning optics **2254**, light delivery optics **2256**, light collection optics **2260**, light adjusting optics **2262**, and a detection sub-system

2264. Each of the foregoing components can optionally be connected to control unit **2228**, or alternatively, to another control unit. For purposes of explanation below, it will be assumed that control unit **2228** is connected to the components of imaging apparatus **2250**.

[1203] During operation of imaging apparatus **2250**, light source **2252** generates light. In general, the light generated by source **2252** can include light in any one or more of the ultraviolet, visible, and/or infrared regions of the electromagnetic spectrum. A variety of different light source elements can be used to generate the light, including (but not limited to) light emitting diodes, laser diodes, laser sources, fluorescent sources, incandescent sources, and glow-discharge sources.

[1204] The light generated by light source **2252** is received by light conditioning optics **2254**. In general, light conditioning optics **2254** modify the light generated by light source **2252** for specific imaging applications. For example, in some embodiments, light conditioning optics **2254** modify the spectral properties of the light, e.g., by filtering out certain wavelengths of the light. For this purpose, light conditioning optics **2254** can include a variety of spectral optical elements, such as optical filters, gratings, prisms, and chromatic beam splitters.

[1205] In certain embodiments, light conditioning optics **2254** modify the spatial properties of the light generated by light source **2252**. Examples of components that can be used for this purpose include (but are not limited to) apertures, phase masks, apodizing elements, and diffusers.

[1206] After modification by light conditioning optics **2254**, the light is received by light delivery optics **2256** and directed onto sample **2208** or feature array **2214**, either of which is positioned on a mount **2258**. Light conditioning optics **2254** generally function to collect and direct light onto the surface of the sample or array. A variety of different optical elements can be used for this purpose, and examples of such elements include, but are not limited to, lenses, mirrors, beam splitters, and various other elements having non-zero optical power.

[1207] Light emerging from sample **2208** or feature array **2214** is collected by light collection optics **2260**. In general, light collection optics **2260** can include elements similar to any of those described above in connection with light delivery optics **2256**. The collected light can then optionally be modified by light adjusting optics **2262**, which can generally include any of the elements described above in connection with light conditioning optics **2254**.

[1208] The light is then detected by detection sub-system **2264**. Generally, detection sub-system **2264** functions to generate one or more images of sample **2208** or feature array **2214** by detecting light from the sample or feature array. A variety of different imaging elements can be used in detection sub-system **2264**, including CCD detectors and other image capture devices.

[1209] Each of the foregoing components can optionally be connected to control unit **2228** as shown in FIG. **22B**, so that control unit **2228** can adjust various properties of the imaging apparatus. For example, control unit **2228** can adjust the position of sample **2208** or feature array **2214** relative to the position of the incident light, and also with respect to the focal plane of the incident light (if the incident light is focused). Control unit **2228** can also selectively filter both the incident light and the light emerging from the sample.

[1210] Imaging apparatus **2250** can typically obtain images in a variety of different imaging modalities. In some embodiments, for example, the images are transmitted light images, as shown in FIG. **22B**. In certain embodiments, apparatus **2250** is configured to obtain reflection images. In some embodiments, apparatus **2250** can be configured to obtain birefringence images, fluorescence images, phosphorescence images, multiphoton absorption images, and more generally, any known image type.

[1211] In general, control unit **2228** can perform any of the method steps described herein that do not expressly require user intervention by transmitting suitable control signals to the components of sample handling apparatus **2200** and/or imaging apparatus **2250**. To perform such steps, control unit **2228** generally includes software instructions that, when executed, cause control unit **2228** to undertake specific steps. In some embodiments, control unit **2228** includes an electronic processor and software instructions that are readable by the electronic processor, and cause the processor to

carry out the steps described herein. In certain embodiments, control unit **2228** includes one or more application-specific integrated circuits having circuit configurations that effectively function as software instructions.

[1212] Control unit **2228** can be implemented in a variety of ways. FIG. **22C** is a schematic diagram showing one example of control unit **2228**, including an electronic processor **2280**, a memory unit **2282**, a storage device **2284**, and an input/output interface **2286**. Processor **2280** is capable of processing instructions stored in memory unit **2282** or in storage device **2284**, and to display information on input/output interface **2286**.

[1213] Memory unit **2282** stores information. In some embodiments, memory unit **2282** is a computer-readable medium. Memory unit **2282** can include volatile memory and/or non-volatile memory. Storage device **2284** is capable of providing mass storage, and in some embodiments, is a computer-readable medium. In certain embodiments, storage device **2284** may be a floppy disk device, a hard disk device, an optical disk device, a tape device, a solid state device, or another type of writeable medium.

[1214] The input/output interface **2286** implements input/output operations. In some embodiments, the input/output interface **2286** includes a keyboard and/or pointing device. In some embodiments, the input/output interface **2286** includes a display unit for displaying graphical user interfaces and/or display information.

[1215] Instructions that are executed and cause control unit **2228** to perform any of the steps or procedures described herein can be implemented in digital electronic circuitry, or in computer hardware, firmware, or in combinations of these. The instructions can be implemented in a computer program product tangibly embodied in an information carrier, e.g., in a machine-readable storage device, for execution by a programmable processor (e.g., processor **2280**). The computer program can be written in any form of programming language, including compiled or interpreted languages, and it can be deployed in any form, including as a stand-alone program or as a module, component, subroutine, or other unit suitable for use in a computing environment. Storage devices suitable for tangibly embodying computer program instructions and data include all forms of non-volatile memory, including by way of example semiconductor memory devices, such as EPROM, EEPROM, and flash memory devices; magnetic disks such as internal hard disks and removable disks; magneto-optical disks; and CD-ROM and DVD-ROM disks. The processor and the memory can be supplemented by, or incorporated in, ASICs (application-specific integrated circuits).

[1216] Processor **2280** can include any one or more of a variety of suitable processors. Suitable processors for the execution of a program of instructions include, by way of example, both general and special purpose microprocessors, and the sole processor or one of multiple processors of any kind of computer or computing device.

EXEMPLARY EMBODIMENTS

[1217] In some non-limiting examples of the workflows described herein, the sample can be immersed in 100% chilled methanol and incubated for 30 minutes at -20°C . After 20 minutes, the sample can be removed and rinsed in ultrapure water. After rinsing the sample, fresh eosin solution is prepared, and the sample can be covered in isopropanol. After incubating the sample in isopropanol for 1 minute, the reagent can be removed by holding the slide at an angle, where the bottom edge of the slide can be in contact with a laboratory wipe and air dried. The sample can be uniformly covered in hematoxylin solution and incubated for 7 minutes at room temperature. After incubating the sample in hematoxylin for 7 minutes, the reagent can be removed by holding the slide at an angle, where the bottom edge of the slide can be in contact with a laboratory wipe. The slide containing the sample can be immersed in water and the excess liquid can be removed. After that, the sample can be covered with bluing buffer and can be incubated for 2 minutes at room temperature. The slide containing the sample can again be immersed in water, and uniformly covered with eosin solution and incubated for 1 minute at room temperature. The slide can be air-dried for no more than 30 minutes and incubated for 5 minutes at 37°C . The sample can be imaged

using brightfield imaging setting.

[1218] Further, the biological sample can be processed by the following exemplary steps for sample permeabilization and cDNA generation. The sample can be exposed to a permeabilization enzyme and incubated at 37° C. for the pre-determined permeabilization time (which is tissue type specific). The permeabilization enzyme can be removed and the sample prepared for analyte capture by adding 0.1×SSC buffer. The sample can then subjected to a pre-equilibration thermocycling protocol (e.g., lid temperature and pre-equilibrate at 53° C., reverse transcription at 53° C. for 45 minutes, and then hold at 4° C.) and the SSC buffer can be removed. A Master Mix, containing nuclease-free water, a reverse transcriptase reagent, a template switch oligo, a reducing agent, and a reverse transcriptase enzyme can be added to the biological sample and substrate, and the sample with the Master Mix can be subjected to a thermocycling protocol (e.g., perform reverse transcription at 53° C. for 45 minutes and hold at 4° C.). Second strand synthesis can be performed on the substrate by subjecting the substrate to a thermocycling protocol (e.g., pre-equilibrate at 65° C., second strand synthesis at 65° C. for 15 minutes, then hold at 4° C.). The Master Mix reagents can be removed from the sample and 0.8M KOH can be applied and incubated for 5 minutes at room temperature. The KOH can be removed and elution buffer can be added and removed from the sample. A Second Strand Mix, including a second strand reagent, a second strand primer, and a second strand enzyme, can be added to the sample and the sample can be sealed and incubated. At the end of the incubation, the reagents can be removed and elution buffer can be added and removed from the sample, and 0.8M KOH can be added again to the sample and the sample can be incubated for 10 minutes at room temperature. Tris-HCl can be added and the reagents can be mixed. The sample can be transferred to a new tube, vortexed, and placed on ice.

[1219] Further the biological sample can be processed by the following exemplary steps for cDNA amplification and quality control. A qPCR Mix, including nuclease-free water, qPCR Master Mix, and cDNA primers, can be prepared and pipetted into wells in a qPCR plate. A small amount of sample can be added to the plated qPCR Mix, and thermocycled according to a predetermined thermocycling protocol (e.g., step 1: 98° C. for 3 minutes, step 2: 98° C. for 5 seconds, step 3: 63° C. for 30 seconds, step 4: record amplification signal, step 5: repeating 98° C. for 5 seconds, 63° C. for 30 seconds for a total of 25 cycles). After completing the thermocycling, a cDNA amplification mix, including amplification mix and cDNA primers, can be prepared and combined with the remaining sample and mixed. The sample can then be incubated and thermocycled (e.g., lid temperature at 105° C. for ~45-60 minutes; step 1: 98° C. for 3 minutes, step 2: 98° C. for 15 seconds, step 3: 63° C. for 20 seconds, step 4: 72° C. for one minute, step 5: [the number of cycles determined by qPCR Cq Values], step 6: 72° C. for 1 minute, and step 7: hold at 4° C.). The sample can then be stored at 4° C. for up to 72 hours or at -20° C. for up to 1 week, or resuspended in 0.6×SPRIselect Reagent and pipetted to ensure proper mixing. The sample can then be incubated at 5 minutes at room temperature, and cleared by placing the sample on a magnet (e.g., the magnet is in the high position). The supernatant can be removed and 80% ethanol can be added to the pellet, and incubated for 30 seconds. The ethanol can be removed and the pellet can be washed again. The sample can then be centrifuged and placed on a magnet (e.g., the magnet is on the low position). Any remaining ethanol can be removed and the sample can be air dried for up to 2 minutes. The magnet can be removed and elution buffer can be added to the sample, mixed, and incubated for 2 minutes at room temperature. The sample can then be placed on the magnet (e.g., on low position) until the solution clears. The sample can be transferred to a new tube strip and stored at 4° C. for up to 72 hours or at -20° C. for up to 4 weeks. A portion of the sample can be run on an Agilent Bioanalyzer High Sensitivity chip, where a region can be selected and the cDNA concentration can be measured to calculate the total cDNA yield. Alternatively, the quantification can be determined by Agilent Bioanalyzer or Agilent TapeStation.

[1220] Further, the biological sample can be processed by the following exemplary steps for spatial gene expression library construction. A Fragmentation Mix, including a fragmentation buffer and

fragmentation enzyme, can be prepared on ice. Elution buffer and fragmentation mix can be added to each sample, mixed, and centrifuged. The sample mix can then be placed in a thermocycler and cycled according to a predetermined protocol (e.g., lid temperature at 65° C. for ~35 minutes, pre-cool block down to 4° C. before fragmentation at 32° C. for 5 minutes, End-repair and A-tailing at 65° C. for 30 minutes, and holding at 4° C.). The 0.6× SPRIselect Reagent can be added to the sample and incubated at 5 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears, and the supernatant can be transferred to a new tube strip. 0.8× SPRIselect Reagent can be added to the sample, mixed, and incubated for 5 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears. The supernatant can be removed and 80% ethanol can be added to the pellet, the pellet can be incubated for 30 seconds, and the ethanol can be removed. The ethanol wash can be repeated and the sample placed on a magnet (e.g., in the low position) until the solution clears. The remaining ethanol can be removed and elution buffer can be added to the sample, mixed, and incubated for 2 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears, and a portion of the sample can be moved to a new tube strip. An Adaptor Ligation Mix, including ligation buffer, DNA ligase, and adaptor oligos, can be prepared and centrifuged. The Adaptor Ligation Mix can be added to the sample, pipette-mixed, and centrifuged briefly. The sample can then be thermocycled according to a predetermined protocol (e.g., lid temperature at 30° C. for ~15 minutes, step 1: 20° C. for 15 minutes, step 2: 4° C. hold). The sample can be vortexed to re-suspend SPRIselect Reagent, additional 0.8× SPRIselect Reagent can be added to the sample and incubated for 5 minutes at room temperature, and placed on a magnet (e.g., in the high position) until the solution clears. The supernatant can be removed and the pellet can be washed with 80% ethanol, incubated for 30 seconds, and the ethanol can be removed. The ethanol wash can be repeated, and the sample can be centrifuged briefly before placing the sample on a magnet (e.g., in the low position). Any remaining ethanol can be removed and the sample can be air dried for a maximum of 2 minutes. The magnet can be removed, and elution buffer can be added to the sample, and the sample can be pipette-mixed, incubated for 2 minutes at room temperature, and placed on a magnet (e.g., in the low position) until the solution clears. A portion of the sample can be transferred to a new tube strip. Amplification mix, can be prepared and combined with the sample. An individual Dual Index TT Set A can be added to the sample, pipette-mixed and subjected to a pre-determined thermocycling protocol (e.g., lid temperature at 105° C. for ~25-40 minutes, step 1: 98° C. for 45 seconds, step 2: 98° C. for 20 seconds, step 3: 54° C. for 30 seconds; step 4: 72° C. for 20 seconds, step 5: reverting to step 2 for a predetermined number of cycles, step 6: 72° C. for 1 minute, and 4° C. on hold). Vortex to re-suspend the SPRIselect Reagent, additional 0.6× SPRIselect Reagent can be added to each sample, mixed, and incubated for 5 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears, and the supernatant can be transferred to a new tube strip. The 0.8× SPRIselect Reagent can be added to each sample, pipette-mixed, and incubated for 5 minutes at room temperature. The sample can then be placed on a magnet (e.g., in the high position) until the solution clears. The supernatant can be removed, and the pellet can be washed with 80% ethanol, incubated for 30 seconds, and then the ethanol can be removed. The ethanol wash can be repeated, the sample centrifuged, and placed on a magnet (e.g., in the low position) to remove any remaining ethanol. The sample can be removed from the magnet and Elution Buffer can be added to the sample, pipette-mixed, and incubated at 2 minutes at room temperature. The sample can be placed on a magnet (e.g., in the low position) until the solution clears and a portion of the sample can be transferred to a new tube strip. The sample can be stored at 4° C. for up to 72 hours, or at -20° C. for long-term storage. The average fragment size can be determined using a Bioanalyzer trace or an Agilent TapeStation.

[1221] The library can be sequenced using available sequencing platforms, including, MiSeq, NextSeq 500/550, HiSeq 2500, HiSeq 3000/4000, NovaSeq, and iSeq.

[1222] In alternate embodiments of the above described workflows, a biological sample can be permeabilized by exposing the sample to greater than about 1.0 w/v % (e.g., greater than about 2.0 w/v %, greater than about 3.0 w/v %, greater than about 4.0 w/v %, greater than about 5.0 w/v %, greater than about 6.0 w/v %, greater than about 7.0 w/v %, greater than about 8.0 w/v %, greater than about 9.0 w/v %, greater than about 10.0 w/v %, greater than about 11.0 w/v %, greater than about 12.0 w/v %, or greater than about 13.0 w/v %) sodium dodecyl sulfate (SDS). In some embodiments, a biological sample can be permeabilized by exposing the sample (e.g., for about 5 minutes to about 1 hour, about 5 minutes to about 40 minutes, about 5 minutes to about 30 minutes, about 5 minutes to about 20 minutes, or about 5 minutes to about 10 minutes) to about 1.0 w/v % to about 14.0 w/v % (e.g., about 2.0 w/v % to about 14.0 w/v %, about 2.0 w/v % to about 12.0 w/v %, about 2.0 w/v % to about 10.0 w/v %, about 4.0 w/v % to about 14.0 w/v %, about 4.0 w/v % to about 12.0 w/v %, about 4.0 w/v % to about 10.0 w/v %, about 6.0 w/v % to about 14.0 w/v %, about 6.0 w/v % to about 12.0 w/v %, about 6.0 w/v % to about 10.0 w/v %, about 8.0 w/v % to about 14.0 w/v %, about 8.0 w/v % to about 12.0 w/v %, about 8.0 w/v % to about 10.0 w/v %, about 10.0 w/v % to about 14.0 w/v %, about 10.0 w/v % to about 12.0 w/v %, or about 12.0 w/v % to about 14.0 w/v %) SDS and/or proteinase K (e.g., at a temperature of about 35° C. to about 50° C., about 35° C. to about 45° C., about 35° C. to about 40° C., about 40° C. to about 50° C., about 40° C. to about 45° C., or about 45° C. to about 50° C.).

[1223] In alternate embodiments of the above described workflows, the substrate can include 5000 features (e.g., spots, beads, pads, etc.). For example, a substrate including 5,000 features can provide higher resolution than an array including 1,000 features (e.g., spots, beads, pads, etc.). In alternate embodiments of the above described workflows, the substrate can include fiducial markers. For example, fiducial markers can allow for proper orientation, detection, and/or rotation of the sample on the substrate.

[1224] In alternative embodiments of the above described workflows, one or more species of RNA (e.g., ribosomal, mitochondrial RNA, etc.) can be down-selected (e.g., removed or depleted). Non-limiting examples of hybridization and capture method ribosomal RNA depletion include RiboMinus™, RiboCop™, and Ribo-Zero™. Another non-limiting RNA depletion method involves hybridization of complementary DNA oligonucleotides to unwanted RNA followed by degradation of the RNA/DNA hybrids using RNase H. Non-limiting examples of a hybridization and degradation method include NEBNext® rRNA depletion, NuGEN AnyDeplete, and RiboZero Plus. Another non-limiting ribosomal RNA depletion method includes ZapR™ digestion, for example SMARTer. In the SMARTer method, random nucleic acid adapters are hybridized to RNA for first-strand synthesis and tailing by reverse transcriptase, followed by template switching and extension by reverse transcriptase.

[1225] In alternative embodiments of the above described workflows, additional reagents can be added to improve the recovery of one or more target molecules (e.g., cDNA molecules, mRNA transcripts). For example, addition of carrier RNA to a RNA sample workflow process can increase the yield of extracted RNA/DNA hybrids from the biological sample. Carrier molecules (e.g., RNA) can be useful when the concentration of input or target molecules is low as compared to remaining molecules. Generally, single target molecules cannot form a precipitate, and addition of the carrier molecules can help in forming a precipitate. Carrier RNA can be added immediately prior to a second strand synthesis step. Carrier RNA can be added immediately prior to a second strand cDNA synthesis on oligonucleotides (e.g., via denaturation) released from the substrate. Carrier RNA can be added immediately prior to a post in vitro transcription clean-up step. Carrier RNA can be added prior to amplified RNA purification and quantification. Carrier RNA can be added before RNA quantification. Carrier RNA can be added immediately prior to both a second strand cDNA synthesis step and a post in vitro transcription clean-up step.

[1226] In alternative embodiments of the above described workflows, permeabilization can be performed with acetone (e.g., with fresh frozen tissue). When acetone fixation is performed, pre-

permeabilization steps may not be performed. Alternatively, acetone fixation can be performed with permeabilization steps. In alternative embodiments of the above described workflow, PAS staining can be performed when samples are permeabilized with acetone.

[1227] In non-limiting examples of any of the workflows described herein, a nucleic acid molecule is produced that includes a contiguous nucleotide sequence comprising: (a) a first primer sequence (e.g., Read 1); (b) a spatial barcode; (c) a unique molecular sequence (UMI); (d) a capture domain; (e) a sequence complementary to a sequence present in a nucleic acid from a biological sample; (f) a second primer sequence (e.g., Read 2) that is substantially complementary to a sequence of a template switching oligonucleotide (TSO). In some embodiments of these nucleic acid molecules, the nucleic acid molecule is a single-stranded nucleic acid molecule. In some embodiments of these nucleic acid molecules, the nucleic acid molecule is a double-stranded nucleic acid molecule. In some embodiments of these nucleic acid molecules, (a) through (f) are positioned in a 5' to 3' direction in the contiguous nucleotide sequence. In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is attached to a substrate (e.g., a slide). In some embodiments of any of these nucleic acid molecules, the 5' end of the contiguous nucleic acid sequence is attached to the substrate (e.g., a slide). In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence is a chimeric RNA and DNA sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence is a DNA sequence.

[1228] In non-limiting examples of any of the workflows described herein, a nucleic acid molecule is produced that includes a contiguous nucleotide sequence comprising: (a) a sequence complementary to a first primer sequence (e.g., a sequence complementary to Read 1); (b) a sequence complementary to a spatial barcode; (c) a sequence complementary to a unique molecular sequence; (d) a sequence complementary to a capture domain; (e) a sequence present in a nucleic acid from a biological sample; and (f) a sequence of a template switching oligonucleotide (TSO). In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is single-stranded. In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is double-stranded. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence is a DNA sequence. In some embodiments of any of these nucleic acid molecules, (a) through (f) are positioned in a 3' to 5' direction in the contiguous nucleotide sequence.

[1229] In non-limiting examples of any of the workflows described herein, a nucleic acid molecule is produced that includes a contiguous nucleotide sequence comprising: (a) a first primer sequence (e.g., Read 1); (b) a spatial barcode; (c) a unique molecular sequence (UMI); (d) a capture domain; (e) a sequence complementary to a sequence present in a nucleic acid from a biological sample; and (f) a second primer sequence (Read 2). In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is a single-stranded nucleic acid molecule. In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is a double-stranded nucleic acid molecule. In some embodiments of any of these nucleic acid molecules, (a) through (f) are positioned in a 5' to 3' direction in the contiguous nucleotide sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence is a DNA sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence further comprises 3' to (f): (g) a sequence complementary to a first adaptor sequence; and (h) a sequence complementary to a third primer sequence. In some embodiments of any of the nucleic acid molecules, the first adaptor sequence is an i7 sample index sequence. In some embodiments of any of these nucleic acid molecules, the third primer sequence is a P7 primer sequence. In some embodiments of any of these nucleic acid molecules, (h) is 3' positioned relative to (g) in the contiguous nucleotide sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence further comprises 5' to (a): (i) a second adaptor sequence; and (ii) a fourth primer sequence. In some embodiments of any of these nucleic acid molecules, the second

adaptor sequence is an i5 sample index sequence. In some embodiments of any of these nucleic acid molecules, the fourth primer sequence is a P5 primer sequence. In some embodiments of any of these nucleic acid molecules, (ii) is 5' positioned relative to (i) in the contiguous nucleotide sequence.

[1230] In non-limiting examples of any of the workflows described herein, a nucleic acid molecule is produced that includes a contiguous nucleotide sequence comprising: (a) a sequence complementary to a first primer sequence; (b) a sequence complementary to a spatial barcode; (c) a sequence complementary to a unique molecular sequence; (d) a sequence complementary to a capture domain; (e) a sequence present in a nucleic acid from a biological sample; and (f) a sequence complementary to a second primer sequence. In some embodiments of these nucleic acid molecules, a sequence complementary to a first primer sequence is a sequence complementary to Read 1. In some embodiments of these nucleic acid molecules, a sequence complementary to a second primer sequence is a sequence complementary to Read 2. In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is a single-stranded nucleic acid molecule. In some embodiments of any of these nucleic acid molecules, the nucleic acid molecule is a double-stranded nucleic acid molecule. In some embodiments of any of these nucleic acid molecules, (a) through (f) are positioned in a 3' to 5' direction in the contiguous nucleotide sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence is a DNA sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence further comprises 5' to (f): (g) a first adaptor sequence; and (h) a third primer sequence. In some embodiments of any of these nucleic acid molecules, the first adaptor sequence is an i7 sample index sequence. In some embodiments of any of these nucleic acid molecules, the third primer sequence is a P7 primer sequence. In some embodiments of any of these nucleic acid molecules, (h) is 5' positioned relative to (g) in the contiguous nucleotide sequence. In some embodiments of any of these nucleic acid molecules, the contiguous nucleotide sequence further comprises 3' to (a): (i) a sequence complementary to a second adaptor sequence; and (ii) a sequence complementary to a fourth primer sequence. In some embodiments of any of these nucleic acid molecules, the second adaptor sequence is an i5 sample index sequence. In some embodiments of any of these nucleic acid molecules, the fourth primer sequence is a P5 primer sequence. In some embodiments of any of these nucleic acid molecules, (ii) is 3' positioned relative to (i) in the contiguous nucleotide sequence.

Spatial RNA Integrity Number

[1231] Provided herein is a non-limiting example of a protocol for determining the spatial RNA integrity Number (sRIN) in a tissue that can include collecting breast cancer tissue and snap freezing in liquid nitrogen. Tissue can be embedded in OCT and sectioned at 10 or 12 μM thickness at -20°C . and mounted directly on a spatial array including capture probes having an 18S rRNA capture domain. Tissue can be fixed and stained using a Hematoxylin and Eosin (H&E) staining protocol. Briefly, Mayer's Hematoxylin can be added, washed in water, incubated in Bluing buffer, washed in water, stained with Eosin, then washed in water, and finally dried. For visualizing H&E staining, sections can be mounted with 85% glycerol and covered with a coverslip. Bright field imaging can be performed using the Metafer Slide Scanning Platform (Metasystems) where raw images are stitched together with the VSlide software (Metasystems). Glycerol can be removed by holding the spatial array or glass slide in water until the coverslip falls off and then was air dry until the remaining liquid evaporates. Hematoxylin, from the H&E stain, can be optionally removed from the tissue section, for example, by washing in dilute HCl (0.01M) prior to further processing. The tissue sections are then ready for further processing.

[1232] Following staining, the 18S rRNA present in the tissue sections are captured by the 18S rRNA specific capture domains on the spatial array. The 18S rRNA is then converted to cDNA in situ. Specifically, reverse transcription is performed on the spatial array in a sealed hybridization cassette by adding 70 μl reaction mixture including 1 $\times$ First-strand buffer, 5 mM DTT, 1 M

Betaine, 6 mM MgCl₂, 1 mM dNTPs, 0.2 mg/ml BSA, 50 ng/μl Actinomycin D, 10% DMSO, 20 U/μl SuperScript III Reverse Transcriptase, 2 U/μl RNaseOUT Recombinant Ribonuclease Inhibitor. The reaction is performed overnight at 42° C. overnight. After incubation cDNA synthesis mixture is removed and the tissue was washed with 0.1×SSC buffer.

[1233] In order to prepare the spatial array for oligonucleotide probe labeling and imaging, the breast cancer tissue and rRNA is removed. Tissue removal can be performed first by incubation with β-mercaptoethanol in RLT lysis buffer at a 3:100 ratio at 56° C. for 1 hour with continuous shaking at 300 rpm. All tissues can be incubated with a 1:7 ratio of Proteinase K and PKD buffer for 1 hour at 56° C. using short intervals with gentle shaking at 300 rpm. The spatial array is then washed with continuous shaking at 300 rpm as follows: first in 2×SSC with 0.1% SDS at 50° C. for 10 min, then in 0.2×SSC at RT for 1 min and finally in 0.1×SSC at RT for 1 min. The spatial array is then spin-dried and put back into the hybridization cassette. rRNA removal can be performed using a reaction mixture containing the following final concentrations: 1× First-strand buffer, 0.4 mg/ml BSA and 16.3 mU/μl RNase H. The reaction can be performed for 1 hour at 37° C. with gentle shaking at 300 rpm using short intervals. Spatial arrays are then washed with 0.1×SSC buffer and treated with 60% DMSO at room temperature for 5 minutes and then washed three times with 0.1×SSC buffer.

[1234] In order to detect the cDNA produced from the 18S rRNA, labeled oligonucleotide probes are generated that had sequence complementarity to the 18S cDNA sequence. Oligonucleotides are identified that have a length between 18-23 nucleotides with an optimum at 20 nucleotides, a melting temperature (T_m) between 38-50° C. with the optimal temperature at 42° C., and a content of guanine and cytosine of 30-60% with an optimum at 50%. The first five bases of the 3'-ends are set to include two of either guanine or cytosine or one of each. Oligonucleotides are checked for quality using Mfold (determination of secondary structure), Oligo Calc: Oligonucleotide Properties Calculator (determination of self-dimerization and hairpin formation), and BLAST (determination of off-target binding). The sequence locations are picked for compatibility with both human (NR_003286.2) and mouse (NR_003278.3) 18S rRNA. Four of the oligonucleotide probes selected include: probe 1 (P1; SEQ ID NO: 4) GAGGAATTCCCAGTAAGT, probe 2 (P2; SEQ ID NO: 5) GAGATTGAGCAATAACAG, probe 3 (P3; SEQ ID NO: 6) GTAGTTCCGACCATAAAC, and probe 4 (P4; SEQ ID NO: 7) GGTGACTCTAGATAACCT. Control oligonucleotide probes can be designed to include complementary sequences of three detection probes at a time with a 20 bases spacer sequence between each probe. The selected oligonucleotide probes can be then designed to incorporate a Cy3 fluorophore.

[1235] Next, labeled oligonucleotide probes are hybridized to the spatial array containing cDNA produced from the 18S rRNA or containing control capture probes. This step can include at least 4 successive rounds of hybridization and imaging, with at least one round for each of the four labeled oligonucleotide probes. Following each round of hybridization and imaging, the spatial array can be washed to remove the hybridized probe before continuing with a subsequent round of hybridization and imaging.

[1236] Hybridization of labeled oligonucleotide probes includes adding to the spatial arrays a pre-heated (e.g., heated to 50° C.) hybridization mixture (10 mM Tris-HCl, 1 mM EDTA, 50 mM

[1237] NaCl, and 0.5 μM of fluorescently labelled probe) containing at least 0.5 μM of one of the fluorescently labeled oligonucleotide probes (e.g., one of P1, P2, P3 or P4). The spatial array can be then imaged using a DNA microarray scanner with the following settings: excitation wavelength 532 nm set to gain 70 and 635 nm set to 1. Following imaging, the spatial array is incubated with 60% DMSO at room temperature for 5 minutes and washed three times with 0.1×SSC buffer to remove the hybridized probe. Subsequent rounds of hybridization and imaging are performed with a different labeled oligonucleotide probe used in each round (e.g., round 2 used P2, round 3 used P3, and round 4 used P4). An initial, pre-hybridization round (P0) of imaging are performed in order to assess background fluorescence.

[1238] One image is generated per labeled oligonucleotide probe (P1-P4) and also one where no fluorescently labeled probes were hybridized (P0). Normalization of Fluorescence Units (FU) data is done by subtraction of the auto-fluorescence recorded with P0 and division with P1. After aligning the five images for a particular location in the tissue (one image from each probe, P1-P4, and one image from the location without labeling), the images are loaded into a script and run in RStudio. The Script can generate two different plots, one heat-map of spatial RIN values and one image alignment error plot.

[1239] High quality RNA is defined as full-length (or close to full-length) transcripts, whereas low quality RNA is defined as fragmented transcripts. Spatial RIN values are similar to traditional RIN values in that the RIN values range from 1 to 10, with higher numbers indicating higher quality (e.g., less degraded, less fragmented) RNA samples.

Spatial Assay for Transposase-Accessible Chromatin Compositions

[1240] Provided herein are compositions for identifying the location of an analyte in a biological sample. In some embodiments, a nucleotide molecule composition including a) a spatial barcode b) a unique molecule c) capture domain; d) a functional domain; and e) a splint oligonucleotide. In some embodiments, a partially double-stranded nucleotide molecule composition including: a) a spatial barcode; b) a unique molecular identifier; c) a capture domain; d) a functional domain; and e) a splint oligonucleotide. In some embodiments, a composition including a) a capture probe including i) a spatial barcode; ii) a unique molecular identifier; iii) a capture domain; iv) a functional domain; and v) a splint oligonucleotide and b) fragmented genomic DNA including i) a first adapter sequence comprising a transposon end sequence and a sequence complementary to the capture domain; and ii) a second adapter sequence comprising the transposon end sequence and a second adapter sequence. In some embodiments, a composition including a) a transposase enzyme monomer complexed with a first adapter including i) a transposon end sequence; and ii) a sequence complementary to the capture domain; and b) a transposase enzyme second monomer complexed with a second adapter including i) a transposon end sequence; and ii) a second adapter sequence; c) genomic DNA; and d) a capture probe, including i) a spatial barcode; ii) a unique molecular identifier; iii) a capture domain; iv) a functional domain; and v) a splint oligonucleotide. In some embodiments, a composition including a) a transposase enzyme monomer complexed with a first adapter including i) a transposon end sequence; ii) a sequence complementary to the capture domain; wherein the 5' end of the first adapter is phosphorylated and b) a transposase enzyme second monomer complexed with a second adapter including i) the transposon end sequence; ii) a second adapter sequence; wherein the 5' end of the second adapter is phosphorylated; c) genomic DNA; and d) a capture probe including i) a spatial barcode; ii) a unique molecular identifier; iii) a capture domain; iv) a functional domain; and v) a splint oligonucleotide. In some embodiments, a composition including a) a transposase enzyme dimer including i) a transposase enzyme monomer complexed with a first adapter including 1) a transposon end sequence, 2) a sequence complementary to a capture domain; ii) a transposase enzyme second monomer complexed with a second adapter including 1) the transposon end sequence; 2) a second adapter sequence; b) genomic DNA; c) a capture probe, including i) a spatial barcode; ii) a unique molecular identifier; iii) a capture domain; iv) a functional domain; and v) a splint oligonucleotide. In some embodiments, a composition including a) a transposase enzyme dimer including i) a transposase enzyme monomer complexed with a first adapter including 1) a transposon end sequence, 2) a sequence complementary to a capture domain wherein the 5' end of the first adapter is phosphorylated; ii) a transposase enzyme second monomer complexed with a second adapter including 1) the transposon end sequence; 2) a second adapter sequence wherein the 5' end of the second adapter is phosphorylated; c) genomic DNA; and d) a capture probe, including i) a spatial barcode; ii) a unique molecular identifier; iii) a capture domain; iv) a functional domain; and v) a splint oligonucleotide.

Spatial Transcriptomics

[1241] In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample (e.g., present in a biological sample such as a tissue section) that include: (a) providing a biological sample on a substrate; (b) staining the biological sample on the substrate, imaging the stained biological sample, and selecting the biological sample or subsection of the biological sample to subject to spatial analysis; (c) providing an array comprising one or more pluralities of capture probes on a substrate; (d) contacting the biological sample with the array, thereby allowing a capture probe of the one or more pluralities of capture probes to capture the biological analyte of interest; and (e) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest. Any variety of staining and imaging techniques as described herein or known in the art can be used in accordance with methods described herein. In some embodiments, the staining includes optical labels as described herein, including, but not limited to, fluorescent, radioactive, chemiluminescent, calorimetric, or colorimetric detectable labels. In some embodiments, the staining includes a fluorescent antibody directed to a target analyte (e.g., cell surface or intracellular proteins) in the biological sample. In some embodiments, the staining includes an immunohistochemistry stain directed to a target analyte (e.g., cell surface or intracellular proteins) in the biological sample. In some embodiments, the staining includes a chemical stain such as hematoxylin and eosin (H&E) or periodic acid-schiff (PAS). In some embodiments, significant time (e.g., days, months, or years) can elapse between staining and/or imaging the biological sample and performing spatial transcriptomic analysis. In some embodiments, reagents for performing spatial analysis are added to the biological sample before, contemporaneously with, or after the array is contacted to the biological sample. In some embodiments, step (d) includes placing the array onto the biological sample. In some embodiments, the array is a flexible array where the plurality of spatially-barcoded features (e.g., capture probes) are attached to a flexible substrate. In some embodiments, measures are taken to slow down a reaction (e.g., cooling the temperature of the biological sample or using enzymes that preferentially perform their primary function at lower or higher temperature as compared to their optimal functional temperature) before the array is contacted with the biological sample. In some embodiments, step (e) is performed without bringing the biological sample out of contact with the array. In some embodiments, step (e) is performed after the biological sample is no longer in contact with the array. In some embodiments, the biological sample is tagged with an analyte capture agent before, contemporaneously with, or after staining and/or imaging of the biological sample. In such cases, significant time (e.g., days, months, or years) can elapse between staining and/or imaging and performing spatial analysis. In some embodiments, the array is adapted to facilitate biological analyte migration from the stained and/or imaged biological sample onto the array (e.g., using any of the materials or methods described herein). In some embodiments, a biological sample is permeabilized before being contacted with an array. In some embodiments, the rate of permeabilization is slowed prior to contacting a biological sample with an array (e.g., to limit diffusion of analytes away from their original locations in the biological sample). In some embodiments, modulating the rate of permeabilization (e.g., modulating the activity of a permeabilization reagent) can occur by modulating a condition that the biological sample is exposed to (e.g., modulating temperature, pH, and/or light). In some embodiments, modulating the rate of permeabilization includes use of external stimuli (e.g., small molecules, enzymes, and/or activating reagents) to modulate the rate of permeabilization. For example, a permeabilization reagent can be provided to a biological sample prior to contact with an array, which permeabilization reagent is inactive until a condition (e.g., temperature, pH, and/or light) is changed or an external stimulus (e.g., a small molecule, an enzyme, and/or an activating reagent) is provided.

Spatially-Resolved Gene Expression and Clustering in Invasive Ductal Carcinoma

[1242] The spatial gene expression of invasive ductal carcinoma tissue from a female patient (ER+,

PR-, HER2+) was profiled (BioIVT: Asterand-Case ID 66320; Specimen ID 116899F). As a control, the healthy tissue sections adjacent to the tumor were obtained. 4 replicates were used for each tissue type.

[1243] Spatially-resolved gene expression and clustering in invasive ductal carcinoma reveal intra-tumor heterogeneity is shown in FIGS. 23A-H. FIG. 23A shows a histological section of an invasive ductal carcinoma annotated by a pathologist. The section contains a large proportion of invasive carcinoma (22.344 mm.sup.2 portion indicated by thick black line (outlined in black in color figure)), three separate ductal cancer in situ regions (portions indicated by medium thickness black line 1.329 mm.sup.2, 1.242 mm.sup.2, and 0.192 mm.sup.2 (outlined in green in color figure)), and fibrous tissue. FIG. 23B shows a tissue plot with spots colored by unsupervised clustering of transcripts. FIG. 23C shows a t-SNE plot of spots colored by unsupervised clustering of transcripts. FIG. 23D shows a gene expression heat map of the most variable genes between the 9 identified clusters. The region defined as fibrous tissue mostly corresponds to clusters 1, 7, and 8. Interestingly, a large region annotated as invasive carcinoma by a pathologist contained spatial spots that were assigned to DCIS (cluster 5). In addition, four subtypes of invasive carcinoma with distinct molecular properties (clusters 2, 3, 4, and 6) were identified, revealing intra-tumor heterogeneity.

[1244] The expression levels of genes corresponding to human epidermal growth factor receptor 2 (Her2 or ERBB2), estrogen receptor (ER or ESR1), and progesterone receptor (PGR) in the tissue section are shown in FIG. 23E. It is clearly visible that ERBB2 and ESR1 are highly expressed in the invasive carcinoma and DCIS regions while the expression of PR is absent, consistent with the patient's diagnosis. One of the top differentially expressed genes from each cluster in the invasive carcinoma region was selected (rectangular boxes in FIG. 23D), and its expression levels are located in the tissue as shown in FIG. 23F and overlapped in one plot as shown in FIG. 23G. With the exception of PGR, all of these genes were highly up-regulated in the carcinoma tissue compared to the adjacent normal tissue (FIG. 23H). Analysis revealed that all of these up-regulated genes have implication in cancer progression. Interestingly, in the subset of cluster 3, a long non-coding RNA, of which abnormal expression has recently been implicated in tumor development (see, e.g., Zhang T, et al. Long Non-Coding RNA and Breast Cancer. *Technol Cancer Res Treat*. 2019, 18,1533033819843889, incorporated herein by reference in its entirety), is one of the top differentially expressed genes. In glioblastoma, LINC00645 promotes epithelial-to-mesenchymal transition by inducing TGF- β (see, e.g., Li, C. et al. Long non-coding RNA linc00645 promotes TGF- β -induced epithelial-mesenchymal transition by regulating miR-205-3p-ZEB1 axis in glioma. *Cell Death & Dis*. 2019, 10, 272, incorporated herein by reference in its entirety).

[1245] During breast cancer progression, the myoepithelial cells, which continue to surround preinvasive in situ carcinoma, gradually disappear (see, e.g., Gudjonsson, T. et al. Myoepithelial Cells: Their Origin and Function in Breast Morphogenesis and Neoplasia. *J. Mammary Gland Biol. Neoplasia*. 2009, 10, 261, incorporated herein by reference in its entirety). This phenomenon is clearly visualized in FIGS. 23I and 23J where KRT14 (a gene signature of myoepithelial cells) was highly expressed around the lining of the duct in the normal tissue while it was disappearing in the DCIS region in IDC tissue (FIG. 23I). The extracellular matrix genes such as COL1A1 and FN1, key genes associated with invasion and metastasis, were highly upregulated while smooth muscles and basal keratin were down-regulated in IDC (FIG. 23J).

SEQUENCE LISTING

TABLE-US-00002 Synthetic	PURAMATRIX®	polypeptide	sequence SEQ	ID	NO:	1
RADARADARADA Synthetic	EAK16	polypeptide	sequence SEQ	ID	NO:	2
AEAEAKAKAEAEAKAK Synthetic	KLD12	polypeptide	sequence SEQ	ID	NO:	3
KLDLKLKLKL 18s	cDNA	Probe	1 (P1) SEQ	ID	NO:	4
GAGGAATTCCTCCAGTAAGT 18s	cDNA	Probe	2 (P2) SEQ	ID	NO:	5
GAGATTGAGCAATAACAG 18s	cDNA	Probe	3 (P3) SEQ	ID	NO:	6

VI. Spatial Cell-Based Analytical Methodology for Various Biological Samples

(a) Biological Samples and Analytes

[1246] It has been identified that any of the methods described herein can be used for various types of biological samples. Further, profiles of biological samples can be compared to profiles of other cells, e.g., “normal,” or “healthy,” biological samples. Thus, in some embodiments of any the methods described herein, the methods can provide for diagnosis of a disease. In some embodiments of any the methods described herein, the methods can be used in drug screening.

[1247] In some embodiments, the biological sample is an organism. In some embodiments, the biological sample is an organoid. In some embodiments, the biological sample is one or more cells from a cell culture (e.g., suspended cells or adherent cells). In some instances, the biological sample is a plurality of live cell (e.g., from a cell culture (e.g., suspended cells or adherent cells)).

[1248] Accordingly, provided herein are methods for profiling a biological analyte, e.g., any of the analytes as described herein, in a biological sample such as an organism, an organoid, or an in vitro cell culture. In some embodiments, these methods can include releasing the biological analyte from one or more cells in an organism, an organoid, or an in vitro cell culture. The released biological analyte can be bound by a capture probe as described herein at a distinct spatial position on a substrate and detected. The bound biological analyte can then be correlated with a barcode of the capture probe at a distinct spatial position of the substrate. Such methods can be useful for correlating a biological analyte to a spatial location in a cell culture, an organism, or an organoid on an array as described herein. For example, in some embodiments, provided herein are methods for identifying a location of a biological analyte present in a biological sample, wherein the biological sample is an organism, an organoid, or an in vitro cell culture, the method comprising: (a) contacting the biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain that binds specifically to a biological analyte present in the biological sample; and (b) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample. In some embodiments, provided herein are methods of identifying a location of a biological analyte in a biological sample, wherein the biological sample is an organism, an organoid, or an in vitro cell culture, the method comprising: (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain that binds specifically to the analyte capture sequence; (b) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample. In some embodiments, the biological sample is a cell culture (e.g., a suspended or adherent cell as described herein), an organism, or an organoid.

[1249] In some embodiments, the biological sample is derived from a homogenous cellular population. In some embodiments, the biological sample is derived from a heterogeneous cellular population. In some embodiments, the heterogeneous cellular population is a culture of cells, a whole organism, or an organoid).

[1250] In some embodiments, the biological sample is an organism. In some embodiments, the organism is a nematode (e.g., *Caenorhabditis elegans*), *Danio rerio* (zebrafish), *Xenopus tropicalis*,

Xenopus laevis, *Daphnia* spp., *Coelopidae*, *Drosophila*, *Dipsidae*, *Branchiostoma floridae*, *Galleria mellonella*, *Macrostomum lignano*, *Oscarella carmela*, *Pristionchus pacificus*, *Schmidtea mediterranea*, *Symsagittifera roscoffensis*, *Trichoplax adhaerens*, *Tubifex tubifex*, *Nothobranchius furzeri*, or *Poecilia reticulata*.

[1251] In some embodiments, the biological sample is obtained from a prokaryote such as a bacterium, e.g., *Escherichia coli*, Staphylococci, or *Mycoplasma pneumoniae*; or an archaea. In some embodiments, the biological sample is obtained from a virus. For example, the biological sample can be obtained from a virus such as Hepatitis C virus or human immunodeficiency virus. In some embodiments, the biological sample is obtained from a viroid.

[1252] In some embodiments, the biological sample is an organoid. As referred to herein an “organoid” is a self-organized, three-dimensional tissue culture derived from stem cells. An organoid can be a miniaturized and simplified version of an organ that shows realistic micro-anatomy. Non-limiting examples of an organoid include a cerebral organoid, an intestinal organoid, a stomach organoid, a lingual organoid, a thyroid organoid, a thymic organoid, a testicular organoid, a hepatic organoid, a pancreatic organoid, an epithelial organoid, a lung organoid, a kidney organoid, a gastruloid, a cardiac organoid, and a retinal organoid.

[1253] In some embodiments, the biological sample is a cell culture. A cell culture as described herein can comprise a plurality of cells. In some embodiments, the plurality of cells is from a cell culture of suspended cells or a cell culture of adherent cells as described herein. For example, the adherent cells can be from cell lines comprising at least one of BT549, HS 578T, MCF7, MDA-MB-231, MDAMB-468, T-47D, SF268, SF295, SF539, SNB-19, SNB-75, U251, Colo205, HCC 2998, HCT-116, HCT-15, HT29, KM12, SW620, 786-O, A498, ACHN, CAKI, RXF 393, SN12C, TK-10, UO-31, A549, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H460, NCI-H522, LOX IMVI, M14, MALME-3M, MDA-MB-435, SK-, EL-2, SK-MEL-28, SK-MEL-5, UACC-257, UACC-62, IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3, NCI-ADR-RES, DU145, PC-3, DU145, H295R, HeLa, KBM-7, LNCaP, MCF-7, MDA-MB-468, PC3, SaOS-2, SH-SY5Y, T-47D, THP-1, U87, vero, MC3T3, GH3, PC12, dog MDCK kidney epithelial, *Xenopus* A6 kidney epithelial, zebrafish AB9, or Sf9 insect epithelial cell lines. In some embodiments, the biological sample is a plurality of cells from a disassociated tissue or a tissue section. In some embodiments, the biological sample is a plant. In some embodiments, the plant is a monocotyledonous plant (i.e., a monocot). For example, a monocotyledonous plant includes but is not limited to corn, rice, sugarcane, switchgrass, meadow grass species, Chinese sorghum, barley, wheat and oat, and durum wheat. In some embodiments, the plant is a dicotyledonous plant (i.e., a dicot). For example, a dicotyledonous plant includes but is not limited to soybean, cotton, rape, flax, sunflower, legume, clover, peanut and tomato.

[1254] In some embodiments, the biological samples described herein include analytes of interest. In some embodiments, one or more biological analytes are detected and analyzed in a biological sample. In some embodiments, the analyte is any analyte as described herein, including but not limited to a nucleic acid, a lipid, a carbohydrate, or a protein. In some embodiments, the analyte is an organelle as described herein.

[1255] In some embodiments, the analyte can be a cell surface marker. In some embodiments, a plurality of analyte capture agents can be disposed on the surface of a substrate as described herein. In some embodiments, the plurality of analyte capture agents to a biological sample can be an antibody. In some embodiments, the plurality of analyte capture agents to a biological sample can bind to a cell surface marker. Non-limiting examples of a cell surface marker include CD45, CD3, CD4, CD8, CD56, CD19, CD20, CD11c, CD14, CD33, CD66b, CD34, CD41, CD61, CD235a, CD146, and epithelial cellular adhesion molecule (EpCAM). An analyte capture agent disposed on the surface of a substrate can be used to bind a cell that has the corresponding cell surface maker.

[1256] In some embodiments, the biological sample is contacted with a substrate when the biological sample is in a bulk solution. In some embodiments, the bulk solution permeabilizes one

or more membranes of a biological sample, allowing access to a biological analyte. A bulk solution can be any bulk solution as described herein.

[1257] Tissues and cells can be obtained from any source. For example, tissues and cells can be obtained from single-cell or multicellular organisms (e.g., a mammal). Tissues and cells obtained from a mammal, e.g., a human, often have varied analyte levels (e.g., gene and/or protein expression) which can result in differences in cell morphology and/or function. The position of a cell or a subset of cells (e.g., neighboring cells and/or non-neighboring cells) within a tissue can affect, e.g., the cell's fate, behavior, morphology, and signaling and cross-talk with other cells in the tissue. Information regarding the differences in analyte levels (gene and/or protein expression) within different cells in a tissue of a mammal can also help researchers identify and elucidate analyte levels in the single-cell or multicellular organisms (e.g., a mammal) based on the detected differences in analyte levels within different cells in the tissue. Differences in analyte levels within different cells in a tissue of a mammal can also provide information on how tissues (e.g., healthy and diseased tissues) function and/or develop. Differences in analyte levels within different cells in a tissue of a mammal can also provide information of different mechanisms of disease pathogenesis in a tissue and mechanism of action of a therapeutic treatment within a tissue. Differences in analyte levels within different cells in a tissue of a mammal can also provide information on drug resistance mechanisms and the development of the same in a tissue of a mammal. Differences in the presence or absence of analytes within different cells in a tissue of a multicellular organism (e.g., a mammal) can provide information on drug resistance mechanisms and the development of the same in a tissue of a multicellular organism.

(b) Substrate(s) for Cell-Based Analysis

[1258] One or more substrates can be used in the methods as described herein. In some embodiments, a biological sample can be disposed upon a substrate comprising a plurality of capture probes as described herein. In some embodiments, wherein one substrate is used, a biological sample is disposed upon a substrate comprising a plurality of capture probes as described herein (i.e., onto the same substrate). See, for example, FIG. 26A.

[1259] In some embodiments, a biological sample can be disposed upon a first substrate and contacted with a second substrate comprising a plurality of capture probes. In some embodiments, wherein two substrates are used, a biological sample is disposed upon a first substrate, and the first substrate with the biological sample is contacted with a second substrate that comprises a plurality of capture probes as described herein. See, for example, FIG. 26B. As another example, a plurality of cells, e.g., from a cell culture, can be distributed upon either a first substrate or a second substrate. In some embodiments, wherein the first substrate comprises a plurality of capture probes, a cell of the plurality of cells is distributed onto the first substrate such that the cell occupies a distinct feature on the first substrate. In some embodiments, a cell of the plurality of cells is distributed onto a first substrate and contacted with a second substrate comprising a plurality of capture probes. In other words, the biological sample (e.g., cells) are on one slide and the plurality of capture probes are on another slide.

[1260] In some embodiments, the first substrate is contacted with the biological sample. In some embodiments, the first substrate is contacted with the second substrate. In some embodiments, the first substrate is contacted with the second substrate prior to releasing the biological analyte from the biological sample. In some embodiments, the biological sample is lysed or permeabilized, allowing capture probes to associate with (e.g., hybridize to) a target analyte. In some embodiments, after association of the capture probe with an analyte, further steps, including cDNA synthesis and purification, and sequencing are performed using methods described herein.

[1261] In some embodiments, a plurality of cells (e.g., a cell culture comprising a plurality of cells) is distributed onto the substrate (e.g., the first substrate). In some embodiments, wherein the substrate (e.g., the first substrate) comprises a plurality of features, a cell of the plurality of cells is distributed onto the substrate such that the cell occupies a distinct feature on the substrate. In some

embodiments, wherein the substrate (e.g., the first substrate) comprises a plurality of features, members of the plurality of cells are distributed over the substrate (e.g., the first substrate) until about all of the features on the substrate are bound to a cell. In some embodiments, wherein the first substrate comprises a plurality of features, the features are smaller than the minimum cell diameter. In some embodiments, wherein the first substrate comprises a plurality of features, cells not bound to a feature on the substrate are washed off. In some embodiments, cells not bound to a feature on the substrate are washed off prior to releasing the biological analyte. In some embodiments, members of the plurality of cells are distributed over the substrate prior to the binding of members of the plurality of cells to members of the plurality of capture antibodies. [1262] In some embodiments, the surface of the substrate (e.g., the first substrate) is coated with a cell-permissive coating, e.g., a cell-permissive coating as described herein. In some embodiments, the cell-permissive coating includes at least one of a biological material and/or a synthetic material. In some embodiments, the cell-permissive coating includes coatings that feature one or more extracellular matrix (ECM) components (e.g., proteoglycans and fibrous proteins such as collagen, elastin, fibronectin and laminin), poly-lysine, poly(L)-ornithine, and/or a biocompatible silicone (e.g., CYTOSOFT®). For example, a cell-permissive coating that includes one or more extracellular matrix components can include collagen Type I, collagen Type II, collagen Type IV, elastin, fibronectin, laminin, and/or vitronectin. In some embodiments, the cell-permissive coating includes a solubilized basement membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma (e.g., MATRIGEL®). In some embodiments, the cell-permissive coating includes collagen.

[1263] In some embodiments, a biological sample can be immobilized on the substrate. For example, at least one cell of a plurality of cells can be immobilized on the substrate after the at least one cell is distributed onto the substrate. In some embodiments, the biological sample is immobilized on the substrate prior to releasing the biological analyte. In some embodiments, the biological sample is immobilized by adding a hydrogel. In some embodiments, the biological sample is immobilized by applying an electric field.

[1264] In some embodiments, before the biological sample is distributed onto a substrate, it is permeabilized to allow access to one or more biological analytes. In some embodiments, after the biological sample is distributed onto a substrate, it is permeabilized to allow access to one or more biological analytes. Permeabilization can be performed using any reagents and techniques described herein. In some embodiments, the releasing the biological analyte comprises permeabilizing the biological sample, e.g., using any of the permeabilization methods as described herein. In some embodiments, a method as described herein further comprises fixing the biological sample prior to permeabilization.

[1265] In some embodiments, the methods further comprise imaging the biological sample, e.g., imaging using any of the methods as described herein.

VII. Cell Migration Assays and Spatial Cell-Based Analytical Methodology

[1266] In some embodiments, methods as described herein can be combined with a cell migration assay. For example, cells can be disposed on an array as described herein and allowed to migrate (FIGS. 29A and 29B). In some embodiments, the substrate is spotted with barcoded probes and cell capture molecules (FIG. 29A, middle). In some embodiments, the substrate is spotted with barcoded probes, cell capture molecules, and one or more cells (FIG. 29A, right). In some embodiments, the substrate includes one or more capture probes and/or one or more capture antibodies. See FIG. 29B. In some embodiments, the one or more capture antibodies are specific for one or more cell surface markers. In some embodiments, the cell migration distance can be measured and correlated to a biological analyte. In some embodiments, the substrate includes a plurality of capture probes that provide information (e.g., a spatial barcode) that provide a specific position on the substrate. In some embodiments, one can use the spatial barcode information to determine the location of the migrated cell. For example, in some embodiments, a method for

profiling at least one biological analyte present in a cell comprises: (a) allowing a cell of a plurality of cells to migrate on a substrate comprising a plurality of capture probes, wherein a capture probe comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from the cell such that the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; (c) detecting the bound biological analyte; and (d) correlating the biological analyte with a molecular barcode of the capture probe at the distinct spatial position of the substrate, thus profiling at least one biological analyte as having been present in a cell at a distinct spatial position. [1267] In some embodiments, presence of analytes can be analyzed in a setting of cell migration in one or more particular cell(s). For example, in some embodiments, presence of analytes during migration of cancer cells can be detected. For example, in some embodiments, presence of analytes during migration of inflammatory cells can be detected.

[1268] A substrate used in a cell migration assay can comprise one or more channels that are disposed on the substrate (FIG. 28), e.g., to confine cell migration to one dimension along the substrate. Additionally, the channels can direct the migration of a cell such that it does not contact another cell on the substrate (e.g., the channels do not overlap with each other). In some embodiments, the one or more channels confine cell migration along one axis of the substrate. In some embodiments, the one or more channels confine cell migration along more than one axis of the substrate. In some embodiments, the one or more channels confine cell migration along two axes of the substrate. In some embodiments, a channel does not overlap with any other channel. In some embodiments, the channels are wide enough to accommodate a cell (e.g., a channel can have a width of about 2 μm to about 10 μm). For example, a channel can have a width of about 2 μm , about 3 μm , about 4 μm , about 5 μm , about 6 μm , about 7 μm , about 8 μm , about 9 μm , or about 10 μm .

[1269] In some embodiments, cells can be disposed on a substrate as described herein and allowed to migrate. Cell migration in cell migration assays can be used to measure target phenotypes (e.g., phenotype for invasiveness). In some embodiments, the cell migration distance can be measured and correlated to a biological analyte. Reagents can be added to the substrate to facilitate cell migration. For example, the substrate can be coated with one or more extracellular matrix (ECM) components (e.g., basement membrane extract (BME), laminin I, collagen I, collagen IV, fibronectin, vitronectin, elastin), a cell culture medium, a chemoattractant (i.e., a substance which attracts motile cells of a particular type), a chemorepellant, or a combination thereof. Chemoattractants and chemorepellants are inorganic or organic substances possessing chemotaxis-inducer effect in migrating cells. In some embodiments, a chemoattractant is a positive chemoattractant; that is, it promotes movement toward a higher concentration of the chemoattractant (e.g., molecule or chemical). In some embodiments, a chemoattractant is a negative chemoattractant; that is, it promotes movement against a higher concentration of the chemoattractant (e.g., molecule or chemical). Chemoattractants are discussed in Halova et al., *Front Immunol.* 2012; 3:119, which is incorporated by reference in its entirety. For example, chemoattractants include but are not limited to a stem cell factor (SCF) recognized by c-Kit, antigen, which binds to immunoglobulin E (IgE) anchored to the high affinity IgE receptor (Fc ϵ RI); highly cytokinergic (HC) IgE recognized by Fc ϵ RI; lipid mediator sphingosine-1-phosphate (S1P), which binds to G protein-coupled receptors (GPCRs); eicosanoids [prostaglandin E2 and D2, leukotriene (LT) B4, LTD4, and LTC4, and others] and chemokines (CC, CXC, C, and CX3C), which also bind to various GPCRs; isoforms of transforming growth factor (TGF) β 1-3, which are sensitively recognized by TGF- β serine/threonine type I and II β receptors; adenosine; C1q, C3a, and C5a components of the complement; 5-hydroxytryptamine; neuroendocrine peptide catestatin; and tumor necrosis factor- α .

[1270] In some embodiments, a reagent such as a chemoattractant or chemorepellant can be disposed on only a portion of the substrate, present as a gradient along the one or more axis or channels of the substrate, or a combination thereof. In some embodiments, a chemoattractant or

chemorepellant is present at only one end of the one or more channels. In some embodiments, the chemoattractant or chemorepellant is bound to one or more ECM components.

[1271] In some embodiments, the method can further comprise imaging the migration patterns of one or more cells in a biological sample (e.g., organism, organoid, cell culture) on the substrate as described herein. Imaging can be used to determine the spatial location of a cell on the substrate, e.g., the spatial location in a channel on the substrate, and/or to determine the distance of migration of a cell. For example, imaging can be performed before and/or after cell migration. In some embodiments, imaging is performed prior to releasing the biological analyte.

[1272] In some embodiments, detection of the biological analyte is correlated with the cell migration distance. For example, an analyte can be detected before, during, and/or after migration that provides information about the distance traveled. In some embodiments, detection of an analyte is correlated with the cell migration distance of at least about 1 μm , about 2 μm , about 3 μm , about 4 μm , about 5 μm , about 6 μm , about 7 μm , about 8 μm , about 9 μm , about 10 μm , about 15 μm , about 20 μm , about 25 μm , about 30 μm , about 35 μm , about 40 μm , about 45 μm , about 50 μm , about 75 μm , about 100 μm , about 150 μm , about 200 μm , about 250 μm , about 300 μm , about 350 μm , about 400 μm , about 450 μm , about 500 μm , about 550 μm , about 600 μm , about 650 μm , about 700 μm , about 750 μm , about 800 μm , about 850 μm , about 900 μm , about 950 μm , about 1 mm, or more.

VIII. Spatial Analytical Methodology and Selective Release of a Biological Analyte

(a) Methods for Profiling a Selectively-Released Biological Analyte in a Biological Sample

[1273] Provided herein are methods for profiling a biological analyte, e.g., any of the analytes as described herein, in a biological sample that include selectively releasing the biological analyte. In some embodiments, these methods can include selectively releasing the biological analyte from one or more of: a subcellular region of a cell in a biological sample, a particular type of cell in a biological sample, and a region of interest of a biological sample. The released biological analyte can be bound by a capture probe as described herein at a distinct spatial position on a substrate and detected. The bound biological analyte can then be correlated with a barcode of the capture probe at a distinct spatial position of the substrate. Such methods can be useful for correlating a subcellular region of a cell, a particular type of cell, and/or a region of interest of a biological sample to a spatial location for a biological analyte.

[1274] Biological samples can be processed to selectively release a biological analyte from a subcellular region of a cell, a particular type of cell, and/or a region of interest of a biological sample according to established methods. For example, a biological analyte can be selectively released from a region of interest using any of the methods as described herein (e.g., microsurgery, laser capture microdissection, chunking, a microtome, dicing, trypsinization, labelling, and/or fluorescence-assisted cell sorting). In some embodiments, a biological analyte can be selectively released from a subcellular region of a cell, a particular type of cell, and/or a region of interest of a biological sample by selective permeabilization.

[1275] In some embodiments, “selective permeabilization” can refer to a permeabilization method that can permeabilize a membrane of a subcellular region while leaving a different subcellular region substantially intact (e.g., biological analytes are not released from subcellular region due to the applied permeabilization method). Non-limiting examples of selective permeabilization methods used to permeabilize a membrane of a subcellular region while leaving a different subcellular region substantially intact include using electrophoresis and/or applying a permeabilization agent or a lysis agent. Several methods for selective permeabilization of a subcellular region are known to one of skill in the art including the methods described in Lu et al. *Lab Chip*. 2005 January; 5 (1): 23-9; Niklas et al. *Anal Biochem*. 2011 Sep. 15; 416 (2): 218-27; Cox and Emili. *Nat Protoc*. 2006; 1 (4): 1872-8; Chiang et al. *J Biochem. Biophys. Methods*. 2000 Nov. 20; 46 (1-2): 53-68; and Yamauchi and Herr et al. *Microsyst. Nanoeng*. 2017; 3. pii: 16079; each of which is incorporated herein by reference in its entirety. In some embodiments, “selective

permeabilization” can refer to the selective permeabilization of a specific cell type. For example, “selective permeabilization” can refer to permeabilizing one cell type while leaving a different cell type substantially intact (e.g., biological analytes are not released from the cell due to the applied permeabilization method). A cell that is a “different cell type” than another cell can refer to a cell from a different taxonomic kingdom, a prokaryotic cell versus a eukaryotic cell, a cell from a different tissue type, etc. Many methods are known to one of skill in the art for selectively permeabilizing different cell types. Non-limiting examples include applying a permeabilization reagent, electroporation, and/or sonication. See, e.g., International Application No. WO 2012/168003; Han et al. *Microsyst Nanoeng.* 2019 Jun. 17; 5:30; Kashyap et al. *Sci Rep.* 2016; 6:29579; Oren and Shai. *Biochemistry.* 1997 Feb. 18; 36 (7): 1826-35; Algayer et al. *Molecules.* 2019 May 31; 24 (11). pii: E2079; Hipp et al. *Leukemia.* 2017 October; 31 (10): 2278; International Application No. WO 2012/168003; and U.S. Pat. No. 7,785,869; all of which are incorporated by reference herein in their entireties. In some embodiments, “selective permeabilization” can refer to selectively permeabilizing a region of interest of a biological sample. For example, the region of interest can be selectively administered a permeabilization or lysis agent. See also Gould et al. *Oncotarget.* 2018 Mar. 20; 9 (21): 15606-15615. In some embodiments, the permeabilization or lysis agent is administered by contacting the biological sample with a hydrogel that comprises the permeabilization or lysis agent.

[1276] In some embodiments, a method for identifying a location of a biological analyte present in a subcellular region of a cell in a biological sample can include: (a) contacting a biological sample comprising a cell with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from a subcellular region of the cell in, wherein the biological analyte is specifically bound by the capture probe; and (c) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample. In some embodiments, a method for identifying a location of profiling a biological analyte present in a subcellular region of a cell in a biological sample can include: (a) contacting a plurality of analyte capture agents to a biological sample comprising a cell disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture proteins comprises (i) a spatial barcode, and (ii) a capture domain; (b) releasing a biological analyte from a subcellular region of the cell, wherein the analyte capture sequence is specifically bound by the capture domain; and (c) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample. In some embodiments, the biological analyte is released from the subcellular region of the cell wherein release of other biological analytes from other subcellular regions of the cell is substantially prevented. In some embodiments, “substantially prevented” refers to less than 50%, less than 45%, less than 40%, less than 35%, less than 30%, less than 25%, less than 20%, less than 18%, less than 16%, less than 14%, less than 12%, less than 10%, less than 8%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, or less than 1% of other biological analytes being released from other subcellular regions of the cell. In some embodiments, the biological analyte is released from the subcellular region of the cell wherein release of other biological analytes from other subcellular regions of the cell is completely prevented. As used herein, a “subcellular region” can refer to any subcellular region. For example, a subcellular region can refer to cytosol, a mitochondria, a nucleus, a nucleolus, an endoplasmic reticulum, a lysosome, a vesicle, a Golgi

apparatus, a plastid, a vacuole, a ribosome, and cytoskeleton. In some embodiments, the subcellular region comprises at least one of cytosol, a nucleus, a mitochondria, and a microsome. In some embodiments, the subcellular region is cytosol. In some embodiments, the subcellular region is a nucleus. In some embodiments, the subcellular region is a mitochondria. In some embodiments, the subcellular region is a microsome.

[1277] In some embodiments, the subcellular region includes the cytosol. In some embodiments, a cell membrane is permeabilized to extract one or more components in a cell. See FIG. 30. In some embodiments, a cell membrane is permeabilized to extract the cytosolic fraction and provide access to one or more analytes in the cytosolic fraction. In some embodiments, the subcellular region includes the one or more organelles. In some embodiments, the one or more organelles is permeabilized to extract the organelle fraction(s) and provide access to one or more analytes in the organelle fraction(s). In some embodiments, the subcellular region includes the nucleus. In some embodiments, a nuclear membrane is permeabilized to extract the nuclear fraction and provide access to one or more analytes in the nuclear fraction. See, for example, FIG. 30. In some embodiments, the methods disclosed herein utilize a bulk solution. See, for example, FIG. 31. In some embodiments, the biological sample is provided onto the substrate and immobilized using a bulk solution. Afterwards, downstream analysis as disclosed herein (e.g., capture transcripts, synthesize cDNA, collect barcoded cDNA for sequencing, capture nuclear fraction targets, collect barcoded targets for sequencing) can be performed. In some embodiments, the cell membrane is permeabilized to collect cytosolic fraction (e.g., for 3' transcriptomic assay). See, for example, FIG. 31, middle. In some embodiments, the nucleus is permeabilized for ATAC-seq treatment, then collect targets from nuclear fraction. See, for example, FIG. 31, bottom.

[1278] In some embodiments, a method for profiling a biological analyte present in a region of interest of a biological sample can include: (a) contacting a biological sample with a substrate comprising a plurality of capture probes, e.g., capture probes as described herein, wherein a capture probe comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from the particular cell type in the biological sample, wherein the biological analyte is bound by the capture probe at a distinct spatial position of the substrate; (c) detecting the bound biological analyte; and (d) correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, thus profiling the biological analyte as having been present in the biological sample at a distinct spatial position.

[1279] In some embodiments, a method for profiling a biological analyte present in a region of interest of a biological sample can include: (a) contacting a biological sample with a substrate comprising a plurality of capture probes, e.g., capture probes as described herein, wherein a capture probe comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from a region of interest of the biological sample, wherein the biological analyte is bound by the capture probe at a distinct spatial position of the substrate; (c) detecting the bound biological analyte; and (d) correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, thus profiling the biological analyte as having been present in the biological sample at a distinct spatial position. In some examples, a region of interest can be permeabilized or lysed while areas outside the region of interest are not permeabilized or lysed. For example, in some embodiments, a region of interest can be contacted with a hydrogel comprising a permeabilization or lysing reagent. In some embodiments, the area(s) outside the region of interest are not contacted with the hydrogel comprising the permeabilization or lysing reagent. In some embodiments, the area outside the region of interest of the biological sample is removed by microdissection.

[1280] In some embodiments, the methods further comprise imaging the biological sample, e.g., imaging using any of the methods as described herein.

(b) Methods for More than One Selective Release of a Biological Analyte in a Biological Sample

[1281] In some embodiments, biological analytes can be released from a biological sample and

bound to capture probes more than once (e.g., more than one time before and after an event (e.g., before and after treatment of a biological sample)). In some embodiments, the methods disclosed herein include more than one selective permeabilization step that releases a biological analyte from a biological sample. In some embodiments, a first biological analyte from a subcellular region of a cell, a particular type of cell, a region of interest of a biological sample, or a combination thereof, can be released from a biological sample and bound to capture probe. See e.g., FIG. 32A. In some embodiments, a first biological analyte from a biological sample is detected by a capture probe (using the target capture sequence in the capture probe). In some embodiments, the first biological analyte (e.g., an RNA sequence) is reverse transcribed into cDNA. After, copies of the first biological analyte are removed, allowing for additional biological analytes to hybridize or bind to the probe. See e.g., FIG. 32B. In some embodiments, one or more additional biological analytes (e.g., a second biological analyte, a third biological analyte, or a fourth biological analyte) can be released from a different subcellular region of a cell, particular type of cell, region of interest of a biological sample, or combination thereof, and binding of the one or more additional biological analytes to one or more capture probes.

[1282] In some embodiments, the biological sample is in a bulk solution.

[1283] In some embodiments, in between hybridization of a first biological analyte and subsequent biological analytes, one or more probes can be repaired. In some embodiments, repair of a probe includes restoring the probe to the original nucleic acid sequence and length prior to contact with the first biological analyte using any methods of extending or retracting the size of a probe (e.g., a nucleic acid) as described herein.

[1284] In some embodiments, a biological analyte from a subcellular region of a cell, a particular type of cell, a region of interest of a biological sample, or a combination thereof, that is bound by a capture probe at a distinct spatial position of the substrate can be different from a biological analyte from any other subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof, that is bound by another capture probe at a distinct spatial position of the substrate. For example, a biological analyte bound by a capture probe at a distinct spatial position of the substrate from a first subcellular region, first particular type of cell, first region of interest, or combination thereof, can be different from a biological analyte bound by a capture probe at a distinct spatial position of the substrate from a second, third, and/or fourth subcellular region, particular type of cell, region of interest, or combination thereof. In some embodiments, a biological analyte bound by a capture probe at a distinct spatial position of the substrate from a first subcellular region, first particular type of cell, first region of interest, or combination thereof, is the same as a biological analyte bound by a capture probe at a distinct spatial position of the substrate from any other subcellular region, particular type of cell, region of interest, or combination thereof, e.g., a second, third, fourth, or fifth subcellular region, particular type of cell, region of interest, or combination thereof. In some embodiments, wherein more than one biological analyte from a subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof, is bound by a capture probe at a distinct spatial position of the substrate, as least one of the bound biological analytes is different from another bound biological analyte.

[1285] In some embodiments, a biological analyte can be released from more than one subcellular region. For example, in some embodiments, the method further includes: releasing a biological analyte from a subcellular region, e.g., a second subcellular region, of a cell from the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; detecting the bound biological analyte; and correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, wherein the second subcellular region is different than a first subcellular region. In some embodiments, the detection steps are repeated one (e.g., for a third subcellular region), two (e.g., for a fourth subcellular region), three (e.g., for a fifth subcellular region), or four (e.g., for a sixth subcellular region) times.

In some embodiments, the subcellular region of a cell is cytosol and the subcellular region of a cell includes at least one of a nucleus, a mitochondria, or a microsome.

[1286] In some embodiments, a biological analyte can be released from more than one particular type of cell. For example, in some embodiments, the method further comprises: releasing a biological analyte from a particular type of cell, e.g., a second particular type of cell, of a cell from the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; detecting the bound biological analyte; and correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, wherein the particular type of second cell is different than the particular type of a first cell. In some embodiments, the detection steps are repeated one (e.g., for a third particular type of cell), two (e.g., for a fourth particular type of cell), three (e.g., for a fifth particular type of cell), or four (e.g., for a sixth particular type of cell) times.

[1287] In some embodiments, a biological analyte can be released from more than one region of interest. For example, in some embodiments, the method further comprises: releasing a biological analyte from a region of interest, e.g., a second region of interest, of a cell from the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; detecting the bound biological analyte; and correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, wherein the region of interest is different than the region of interest of a first biological analyte. In some embodiments, the detection steps are repeated one (e.g., for a third region of interest), two (e.g., for a fourth region of interest), three (e.g., for a fifth region of interest), or four (e.g., for a sixth region of interest) times.

[1288] The method steps can be performed in various orders. Additionally, the substrate can be washed before or after each step. In some embodiments, wash steps removes unbound analytes.

[1289] There are multiple methods that can be used in detecting biological analytes from a biological sample on which more than one release, e.g., selective permeabilization, step is performed. Non-limiting examples include using second strand synthesis, selective capture probe cleavage, and multiple substrates.

[1290] In some embodiments, second strand synthesis is used in the detection of a biological analyte from a biological sample on which more than one release, e.g., selective permeabilization, step is performed. For example, a biological analyte (e.g., a first biological analyte) from a first release step can be bound to a capture probe on a substrate and second strand synthesis, e.g., as described herein, can be performed to form a second strand comprising a sequence that corresponds to the sequence of at least a portion of the biological analyte (e.g., the first biological analyte). After the second strand from the first release step is collected, a second release step can be performed to bind a biological analyte (e.g., a second biological analyte) to a capture probe, and second strand synthesis can be performed again to form a second strand comprising a sequence that corresponds to at least a portion of the sequence of the biological analyte (e.g., the second biological analyte) and the second strand for the second release step can be collected. See, for example, FIGS. 32A-32B. The second strand synthesis steps may be repeated one, two, three, four, or more times. The collected second strands from each release step can be detected and used to correlate the first, second, third, or fourth biological analyte with the barcode of the capture probe to which the biological analyte was bound at a distinct spatial position of the substrate.

[1291] In some embodiments, selective capture probe cleavage can be used in the detection of a biological analyte from a biological sample on which more than one release, e.g., selective permeabilization, step is performed. For example, a biological analyte (e.g., a first biological analyte) from a first release step can be bound to a first capture probe on a substrate and a first extended capture probe can be generated (e.g., cDNA synthesis can be performed on the first capture probe bound to the biological analyte). The first extended capture probe can be selectively cleaved (e.g., leaving any second, third, or fourth capture probes intact) and collected. After the

first capture probe is collected, a second release step can be performed to bind a biological analyte (e.g., a second biological analyte) to a second capture probe on the substrate and a second extended capture probe can be generated (e.g., cDNA synthesis can be performed on the second capture probe bound to the biological analyte). The second extended capture probe can then be cleaved and collected. See, for example, FIG. 33. The selective capture probe cleavage steps may be repeated one, two, three, four, or more times. The collected extended capture probes from one or more release steps can be detected and used to correlate the first, second, third, or fourth biological analyte with the barcode of the capture probe to which the biological analyte was bound at a distinct spatial position of the substrate. In some embodiments, wherein selective capture probe cleavage is used, the substrate comprises two or more pluralities of capture probes, e.g., a plurality of first capture probes, a plurality of second capture probes, a plurality of third capture probes. In some embodiments, the capture probes of a plurality (e.g., a plurality of first capture probes) comprise a different cleavage domain from the capture probes of another plurality (e.g., a plurality of second capture probes). In some embodiments, members of a plurality of capture probes (e.g., a plurality of first capture probes) comprise a different cleavage domain than members of another plurality of capture probes (e.g., a plurality of second capture probes). In some embodiments, the capture probes of each plurality comprise a different cleavage domain from the capture probes of each other plurality. In some embodiments, the first capture probe is cleaved after step (b). In some embodiments, the first capture probe is cleaved before step (e). In some embodiments, members of the plurality of first capture probes are cleaved after step (b). In some embodiments, members of the plurality of first capture probes are cleaved before step (e).

[1292] In some embodiments, the biological sample is contacted with two or more substrates. For example, after a subcellular region of a cell, a particular type of cell, a region of interest of a biological sample, or a combination thereof, is permeabilized and a biological analyte from the subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof, is captured on a first substrate, the first substrate can be removed, and a biological analyte from a different subcellular region of a cell, particular type of cell, region of interest of a biological sample, or combination thereof, can be captured on a second substrate. See, for example, FIG. 34.

[1293] In some embodiments, after releasing at least one biological analyte from a subcellular region of a cell, a particular type of cell, a region of interest of a biological sample, or a combination thereof, the method further comprises: (e) contacting the biological sample with a second substrate comprising a plurality of capture probes, wherein a capture probe comprises a spatial barcode and a capture domain; (f) releasing a biological analyte from a subcellular region of a cell, a particular type of cell, a region of interest of a biological sample, or a combination thereof, e.g., a second subcellular region of a cell, particular type of cell, region of interest of a biological sample, or combination thereof, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; (g) detecting the bound biological analyte from step (e); and (h) correlating the biological analyte from step (e) with a barcode of the capture probe at the distinct spatial position of the substrate, wherein the subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof, in (f) is different than the subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof in (b). In some embodiments, steps (e) through (g) are repeated one (e.g., for a third subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof), two (e.g., for a fourth subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof), three (e.g., for a fifth subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof), or four (e.g., for a sixth subcellular region of the cell, particular type of cell, region of interest of the biological sample, or combination thereof) times. In some embodiments, steps (e) through (g) are repeated one, two, three, or four times. In some embodiments, the

subcellular region of one or more cells of step (b) is cytosol and the subcellular region of step (e) comprises at least one of a nucleus, a mitochondria, and a microsome. In some embodiments, the biological sample is washed prior to step (e).

IX. Methods for Temporally Profiling a Biological Analyte in a Biological Sample

[1294] In some embodiments, the methods described herein can be performed on the same biological sample or cells obtained from the subject at more than one time points (e.g., before or after treatment with an agent, different stages of differentiation, different stages of disease progression, different ages of the subject, or before or after development of resistance to an agent). In some embodiments, the biological sample is contacted with a different substrate at each time point. See, e.g., FIG. 35. In some embodiments, the amount of time that passes between two or more time points can be measured, for example, in seconds, minutes, days, weeks, or years. In some embodiments, the methods described herein are performed before and after treatment with an agent. In some embodiments, the methods described herein are performed before and after one or more stages of differentiation. In some embodiments, the methods described herein are performed before and after stages of disease progression. In some embodiments, the methods described herein are performed before and after various ages of the subject. In some embodiments, the methods described herein are performed before and after development of resistance to an agent. In some embodiments, the methods described herein are performed before and after a cell migration assay as described in Section VII above.

[1295] In some embodiments, provided herein is a method for temporal profiling the spatial location of at least one biological analyte present in a biological sample, including: (a) contacting a plurality of analyte capture agents to a biological sample comprising a cell disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture proteins comprises (i) a spatial barcode, and (ii) a capture domain; (b) releasing a biological analyte from the cell, wherein the analyte capture sequence is specifically bound by the capture domain; (c) determining (i) all or a part of the sequence of the analyte binding moiety barcode specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (a)-(c) after a period of time. In some embodiments, the same biological analyte is detected at two or more time points. In some embodiments, step (d) is repeated one, two, three, four, or more times. In some embodiments, the method further includes generating a temporal profile for the same biological analyte over time. See, e.g., FIG. 37, which shows a two-dimensional sample (e.g., a tissue sample) that is analyzed over time (t) by examining multiple identical slides. In some embodiments, the method further includes generating a temporal profile more than one biological analyte over time.

[1296] In some embodiments, provided herein is a method for temporal and spatial profiling of a biological analyte present in a biological sample, comprising: (a) contacting a biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain; (b) releasing a biological analyte from the biological sample, wherein the capture domain binds specifically to the released biological analyte; (c) determining (i) all or a part of the sequence of the biological analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (b)-(c) after a period of time.

[1297] In some embodiments, provided herein is a method for temporal and spatial profiling of a biological analyte present in a biological sample, comprising: (a) contacting a plurality of analyte

capture agents to a biological sample comprising a live cell, wherein the biological sample is disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the biological analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises (i) a spatial barcode, and (ii) a capture domain; (b) releasing a biological analyte from the live cell, wherein the analyte capture sequence is specifically bound by the capture domain; (c) determining (i) all or a part of the sequence of the analyte binding moiety barcode specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode or a complement thereof, and using the determined sequences of (i) and (ii) to identify a location of the biological analyte in the biological sample; and (d) repeating steps (b)-(c) after a period of time.

[1298] In some embodiments, the biological sample used in the method for temporal profiling includes a tissue, an organ, an organism, or a cell culture sample as described e.g., in Section VI herein. In some embodiments, the biological sample is in a controlled environment. In some embodiments, the step of releasing comprises releasing under conditions that permit cells in the biological sample to live. In some embodiments, the biological sample is a subcellular fraction as described herein (e.g., in Section VIII above).

[1299] The present disclosure also provides compositions and methods for temporal and spatial profiling of at least one species of biological analyte present in a biological sample that include:

[1300] (a) providing a substrate that includes a plurality of capture probes, wherein the capture probes include a spatial barcode and a capture domain; [1301] (b) contacting the biological sample with the solid support that includes the capture probes; [1302] (c) releasing at least one member of the species of biological analyte from the biological sample; [1303] (d) contacting the solid support that includes the capture probes with at least one released member of the species of biological analyte such that the at least one released member of the species of biological analyte interacts with at least one of the plurality of capture probes; [1304] (e) detecting the at least one released member of the species of biological analyte; [1305] (f) correlating the at least one released member of the species of biological analyte with a spatial location in the biological sample; and [1306] (g) repeating steps (a)-(e) after a period of time.

[1307] In some embodiments, the at least one released biological analyte is a nucleic acid, the capture domain is a nucleic acid, and the at least one released biological analyte hybridizes to the capture domain.

[1308] In some embodiments, the biological sample is in a controlled environment (e.g., a flow cell, a tissue culture system (e.g. a modified tissue culture system), or a tissue bath). In some embodiments, are methods for temporal and spatial profiling of at least one species of biological analyte present in a biological sample include imaging the biological sample (e.g., imaging can be performed prior to releasing at least one member of the species of biological analyte from the biological sample, after releasing at least one member of the species of biological analyte from the biological sample, or both). In some embodiments, imaging is used to determine a region of interest in the biological sample. In some embodiments, imaging comprises using fiducial markers as disclosed herein.

[1309] In some embodiments, the step of releasing comprises applying an electric field to the biological sample (e.g., electroporation and/or subjecting biological sample to brief electric pulsing conditions). In some embodiments, the electric field is applied as a short burst. In some embodiments, the electric field is applied for a period of time of about 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or 55 minutes. In some embodiments, the electric field is applied for a period of time of less than one minute. In some embodiments, the electric field is applied for a period of time of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or more hours. In some embodiments, the step of releasing comprises releasing under conditions that permit

cells in the biological sample to live (e.g., conditions that prevent hypoxia or other situations that can lead to cell death).

[1310] In some embodiments, step (g) is repeated 1, 2, 3, 4, 5, 10, 15, 20, or more times. In some embodiments, a temporal profile for the at least one member of the species of biological analyte in the biological sample is generated.

[1311] In some embodiments, the same species of biological analyte is detected at two or more time points according to any of the variety of methods described herein. In some embodiments, the two or more species of biological analytes are the same class of analyte (e.g., RNA, mRNA, DNA, proteins, antibodies, peptides, lipids, metabolites, or small molecules. In some embodiments, the two or more species of biological analytes are different classes of analytes. In some embodiments, two or more species of two or more classes of biological analytes are analyzed at multiple time points to generate a temporal and spatial profile of the analytes in the biological sample.

[1312] In some embodiments, prior to the step of repeating, the biological sample is subjected to one or more perturbants. In some embodiments provided herein is a method for determining the effect of a perturbant on a biological sample, comprising: (a) contacting the biological sample with the perturbant; and (b) performing a method for temporal profiling as described herein. In some embodiments, the perturbant comprises a small molecule, an antibody, a nucleic acid, a peptide, or a combination thereof.

[1313] A perturbant can be any physical molecule such as, without limitation, a small molecule, an antibody, a nucleic acid, a peptide, or any combination thereof. In some embodiments, a perturbant includes a DNA modification system (e.g., a CRISPR system, e.g., a system that employs a single guide RNA (sgRNA) and an enzymatic molecule or complex (e.g. Cas9 or a derivative thereof) to modify a nucleic acid (e.g., chromosomal DNA) of cells in the biological sample). In some embodiments, a perturbant is a perturbing condition that is not a physical molecule. For example, a perturbant can be, without limitation, light, heat, pressure, an electric field, or any combination thereof. In some embodiments, the biological sample is subjected to same perturbant at multiple time points (e.g., between repeating steps (a)-(f) above such that the spatial and temporal profiles of at least one analyte in the biological sample are determined after each time the biological sample is subjected to the perturbant). In some embodiments, the biological sample is subjected to a different perturbant at multiple time points (e.g., between repeating steps (a)-(f) above such that the spatial and temporal profiles of at least one analyte in the biological sample are determined after each the biological sample is subjected to multiple perturbants at different times). In some embodiments, both a physical molecule and a perturbing condition are used as perturbants. In some embodiments, a physical molecule and a perturbing condition are used simultaneously as perturbants. In some embodiments, a physical molecule and a perturbing condition are used sequentially as perturbants. In some embodiments, a perturbant can be used in accordance with compositions and methods described herein to determine the effect of an agent (e.g., the perturbant can be an agent such as, without limitation, a small molecule drug or biologic agent) on a biological sample. For example the biological sample can be a cancer sample and the agent acting as a perturbant can be a small molecule drug or biologic agent, which is tested for its effect on the cancer sample.

X. Spatially-Tagged Analyte Capture Agents for Spatial Detection

[1314] Provided herein are methods for performing immunohistochemistry and/or immunofluorescence on a capture probe substrate as described herein. Further provided are methods of spatially-tagging analyte capture agents with a capture probe spatial barcode. Further provided are methods for blocking the analyte capture sequence of an analyte capture agent. Further provided are methods for labeling multiple analytes using the same biological sample. In some embodiments, the labels for the multiple analytes are applied to the biological sample at the same time, and in some embodiments, the labels for the multiple analytes are applied to the biological sample at different times (e.g., sequentially).

(a) Spatial Array Compatible Immunohistochemistry (IHC)/Immunofluorescence (IF)

[1315] In some embodiments, IHC and/or IF protocols can be performed on any of the spatial substrates provided herein. In some embodiments, a substrate can be homogeneously covered by a plurality of capture probes. In some embodiments, IHC and/or IF protocols can be performed on a biological sample on a spatial array, wherein after the IHC and/or IF protocol is completed, the biological sample is treated to release analytes for capture by the spatial array. In some embodiments, an antibody of the IHC/IF protocol detects an extracellular protein. In some embodiments, the antibody of the IHC/IF protocol detects a cell surface protein. In some embodiments, the antibody of the IHC/IF protocol detects an intracellular protein (e.g., GFAP, NeuN). In some embodiments, the antibody of the IHC/IF protocol detects a nuclear protein. In some embodiments, the biological sample is fixed. The biological sample can be fixed by methods described herein, or according to methods known in the art. In some embodiments, the biological sample can be blocked, permeabilized, and stained by methods described herein, or according to methods known in the art. In some embodiments, IHC and/or IF protocols can be performed using a detectable label. A detectable label can include radioisotopes, fluorophores, chemiluminescent compounds, bioluminescent compounds, and dyes, including any of the non-limiting examples of detectable labels described herein. In some embodiments, a detectable label can be a fluorophore. In some embodiments, the biological sample can be imaged according to methods described herein or known in the art.

[1316] In some embodiments after the biological sample is imaged, the biological sample can be processed and analyzed according to the spatial analysis workflows described herein. In some embodiments, the IHC and/or IF antibodies are removed from the biological sample before performing the spatial analysis workflows. In other embodiments, the IHC and/or IF antibodies are not removed from the biological sample before performing the spatial analysis workflows.

(i) Immunofluorescence

[1317] Immunofluorescence or immunohistochemistry is a technique used to examine protein expression, distribution, localization, and activation in situ. In some embodiments, immunofluorescence uses the specificity of fluorescently labeled antibodies or probes that bind to specific biological analytes (e.g., within or on a cell surface), and allows for the examination of the distribution of the biological analytes through the tissue sample.

[1318] In some embodiments, disclosed herein are methods of identifying the location of a biological analyte in a biological sample comprising: (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the target analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a spatial barcode and a capture domain that binds specifically to the analyte capture sequence; (b) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample; and (c) performing immunofluorescence microscopy on the biological sample to detect the location of an additional biological analyte in the biological sample. Also disclosed are methods of identifying a location of a biological analyte in a biological sample comprising: (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the target analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence, wherein the analyte capture sequence is hybridized to a blocking probe; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; (b) releasing the blocking probe from the analyte capture sequence, thereby allowing the analyte capture sequence to bind specifically to the capture agent; and (c) determining the sequence

of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the biological analyte in the biological sample.

[1319] In some embodiments, immunofluorescence staining methods can be divided into two categories, direct and indirect methods. In some embodiments, in the direct staining method, a fluorophore is conjugated to an antibody (e.g., “the primary antibody”) which is capable of binding directly to the biological analyte of interest (e.g., cell surface or intracellular target). In some embodiments, in the indirect staining method, the primary antibody binding to the biological analyte is visualized by the binding of a fluorescently labeled second-step antibody, wherein the second-step antibody is capable of binding to the primary antibody. In some embodiments, the second-step antibody is an anti-immunoglobulin antibody.

[1320] In some embodiments, immunofluorescence is performed individually for each analyte of interest. In some embodiments, immunofluorescence can simultaneously detect one or more target proteins in the same cell or tissue (e.g., multiplexed analyte detection). In some embodiments, multiplexing (e.g., simultaneously detecting multiple analytes) allows for examination of spatial arrangement of analytes of interest (e.g., proteins, DNA, RNA) as well as analyte interaction and co-localization thereby facilitating simultaneous analysis of multiple tissue markers.

(b) Spatially-Tagged Capture Agents and Methods for Using the Same

[1321] An “analyte capture agent” refers to a molecule that interacts with a biological analyte and with a capture probe to identify the analyte. In some embodiments, an analyte capture agent includes a label (e.g., fluorescent label). In some embodiments, the analyte capture agent can include an analyte binding moiety and a capture agent barcode domain. An analyte binding moiety is a molecule capable of binding to a specific analyte. In some embodiments, the analyte binding moiety includes an antibody or antibody fragment. In some embodiments, the analyte binding moiety includes a polypeptide and/or an aptamer. In some embodiments, the analyte binding moiety includes a DNA aptamer. In some embodiments, the analyte binding moiety includes a RNA aptamer. In some embodiments, the analyte binding moiety includes an aptamer of mixed natural or unnatural occurring nucleotides (e.g., LNA, PNA). In some embodiments, the analyte is a protein (e.g., a protein on a surface of a cell or an intracellular protein).

[1322] A capture agent barcode domain can include an analyte capture sequence which can hybridize to at least a portion or an entirety of a capture domain of a capture probe. In some embodiments, the analyte capture sequence includes a poly(A) tail. In some embodiments, the analyte capture sequence includes a sequence capable of binding a poly(T) domain. In some embodiments, the analyte capture sequence can have a GC content between 1-100%, e.g., 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, etc.). In some embodiments, the analyte capture sequence has a GC content of at least 30%. In some embodiments, one or more pluralities of analyte capture agents can be provided to a biological sample, wherein one plurality of analyte capture agent differs from another plurality of analyte capture agent by the analyte capture sequence. For example, analyte capture sequence A can be correlated with analyte binding moiety A, and analyte capture sequence B can be correlated with analyte binding moiety B. In some embodiments, the two pluralities of analyte capture agents can have the same analyte binding moiety barcode sequence.

[1323] In some embodiments, the capture domain includes a poly(T) tail. In some embodiments, the capture domain includes a sequence capable of binding a poly (A) domain. In some embodiments, the capture domain can have a GC content between 1-100%, e.g., 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, etc. In some embodiments, the capture domain has a GC content of at least 30%.

[1324] In some embodiments, the capture agent barcode domain includes an analyte binding

moiety barcode. The analyte binding moiety barcode refers to a barcode that is associated with or otherwise identifies the analyte binding moiety. In some embodiments, the analyte binding moiety barcode is correlated with the type of analyte binding moiety, such that more than one plurality of analyte capture agents can be provided to a biological sample at one time. For example, analyte binding moiety barcode A can be correlated with analyte binding moiety A, and analyte binding moiety barcode B is correlated with analyte binding moiety B. The two pluralities of analyte capture agents can have the same analyte capture sequence (e.g., poly(A)). In some embodiments, one analyte binding moiety barcode plurality is correlated with one analyte capture sequence plurality. In other embodiments, an analyte binding moiety barcode plurality is not necessarily correlated with an analyte capture sequence plurality.

[1325] In some embodiments, a capture agent barcode domain includes optional sequences, such as, without limitation, a PCR handle, a sequencing priming site, a domain for hybridizing to another nucleic acid molecule, and combinations thereof. In some embodiments, the PCR handle is identical on all capture analyte barcode domains. In some embodiments, the PCR handle is included for PCR amplification. In some embodiments, an analyte capture agent includes one or more optional sequences and one or more barcode sequences (e.g., one or more analyte binding moiety barcodes and/or one or more UMIs). In some embodiments, the capture probe capture domain and/or the analyte capture agent include a cleavage domain. In some embodiments, a capture agent barcode domain can be dissociated from the analyte binding moiety by cleaving the analyte binding moiety from the capture agent barcode domain via a cleavage domain in the capture agent barcode domain. Other embodiments of an analyte capture agent useful in spatial protein detection are described herein.

[1326] Provided herein are methods for spatially profiling a biological analyte, e.g., any of the analytes as described herein, in a biological sample that use a spatially-tagged analyte capture agent. A biological analyte can be bound by an analyte capture agent at a distinct spatial position on a substrate and detected. The bound biological analyte can then be correlated with a barcode of the capture probe at a distinct spatial position of the substrate. In some embodiments, these methods can include spatially profiling the biological analyte from one or more of: an intracellular region of a cell in a biological sample, a cell surface region of a cell in a biological sample, a particular type of cell in a biological sample, and a region of interest of a biological sample.

(i) Spatially-Tagged Capture Agents

[1327] A “spatially-tagged analyte capture agent” can be a molecule that interacts with an analyte (e.g., an analyte in a sample) and with a capture probe to identify the spatial location of the analyte. In some embodiments, a spatially-tagged analyte capture agent can be an analyte capture agent with an extended capture agent barcode domain that includes a sequence complementary to a spatial barcode of a capture probe. In some embodiments, an analyte capture agent is introduced to an analyte and a capture probe at the same time. In some embodiments, an analyte capture agent is introduced to an analyte and a capture probe at different times. In some embodiments, the spatially-tagged analyte capture agent is denatured from the capture probe before the biological sample is introduced. In some embodiments, the spatially-tagged analyte capture agent binds to a biological analyte within a biological sample before the spatially-tagged analyte capture agent is denatured from the capture probe. In some embodiments, the capture probe is cleaved from the substrate while attached to the spatially-tagged analyte capture agent. In some embodiments, once the capture domain of the capture probe is bound to the analyte binding moiety barcode, the analyte capture sequence is extended towards the 3' tail to include a sequence that is complementary to the sequence of the capture probe spatial barcode (e.g., producing a spatially-tagged analyte capture agent).

[1328] For example, an analyte capture agent can be introduced to a biological sample, wherein the analyte binding moiety binds to a biological analyte, and then the biological sample can be treated to release the analyte-bound analyte capture agent from the sample. The analyte-bound analyte

capture agent can then migrate and bind to a capture probe capture domain, and the analyte-bound capture agent barcode domain can be extended to generate a spatial barcode complement at the end of the capture agent barcode domain. The analyte-bound spatially-tagged analyte capture agent can be denatured from the capture probe, and analyzed using methods described herein.

[1329] In some embodiments, the migration of the analyte capture agent is gravitational migration. In some embodiments, the migration of the analyte capture agent is rotational migration. In some embodiments, the migration of the analyte capture agent uses simple agitation of the biological sample.

[1330] In another example, an analyte capture agent can be hybridized to a capture probe capture domain on a capture probe array, wherein the capture agent barcode domain is extended to include a sequence complementary to the spatial barcode of the capture probe. A biological sample can be contacted with the analyte capture agent modified capture probe array. Analytes from the biological sample can be released from the sample, migrated to the analyte capture agent modified capture probe array, and captured by an analyte binding moiety. The capture agent barcode domain of the analyte-bound analyte capture agents can be denatured from the capture probe, and the biological sample can be dissociated and spatially processed according to methods described herein.

[1331] In some embodiments, a spatially-tagged analyte capture agent can attach to a surface of a cell through a combination of lipophilic and covalent attachment. For example, a spatially-tagged analyte capture agent can include an oligonucleotide attached to a lipid to target the oligonucleotide to a cell membrane, and an amine group that can be covalently linked to a cell surface protein(s) via any number of chemistries described herein. In these embodiments, the lipid can increase the surface concentration of the oligonucleotide and can promote the covalent reaction.

[1332] In some embodiments, a splint oligonucleotide can be used to join the capture agent barcode domain to a capture probe. A “splint oligonucleotide” is an oligonucleotide that, when hybridized to other polynucleotides, acts as a “splint” to position the polynucleotides next to one another so that they can be ligated together. In some embodiments, the splint oligonucleotide is DNA or RNA. The splint oligonucleotide can include a nucleotide sequence that is partially complimentary to nucleotide sequences from two or more different oligonucleotides. In some embodiments, a splint oligonucleotide includes a sequence useful for ligating the capture agent barcode domain onto the capture probe. For example, a capture agent barcode domain of an analyte capture agent includes a cleavage domain and an analyte binding moiety barcode, but does not include an analyte capture sequence. A splint oligonucleotide, which corresponds to a portion of the analyte binding moiety barcode and the capture probe capture domain, can hybridize to both the analyte binding moiety barcode and the capture domain, allowing the analyte binding moiety barcode to ligate to the capture probe capture domain.

[1333] In some embodiments, an analyte-bound, spatially-tagged analyte capture agent is compartmentalized into individual partitions (e.g., spatially-tagged analytes dissociated into single cells), and barcodes (e.g., the analyte binding moiety barcode and/or the spatial barcode of the spatially-tagged analyte capture agent) can be sequenced, analyzed, and correlated with the biological analyte. In some embodiments, the spatially-tagged analyte capture agent can be separated into two or more parts, e.g., the spatially-tagged capture agent barcode domain (e.g., the extended oligonucleotide conjugated to the analyte binding moiety) can be separated from the analyte-bound analyte binding moiety by disrupting a linker that joins the two entities. In some embodiments, the spatially-tagged capture agent barcode domain is analyzed in the same compartment as the analyte-bound analyte binding moiety. In some embodiments, the spatially-tagged capture agent barcode domain is separated from the analyte-bound, analyte binding moiety before compartmentalization. In some embodiments, each partition maintains separation of its own contents from the contents of other partitions.

[1334] For example, the partition can be a droplet in an emulsion or a non-emulsion based partitioning such as encapsulation. In some embodiments, a partition can include a biological

analyte bound to an analyte capture agent (e.g., a spatially-tagged analyte capture agent). In some embodiments, the 3' tail of the analyte binding moiety barcode (e.g., antibody barcode) can be captured and barcoded in an emulsion. In some embodiments, a tissue sample or tissue section is disassociated into single cells, reverse transcription (RT) reagents, and oil are combined on a microfluidic chip to form emulsion partition reaction vesicles. In some embodiments, partitions are formed in parallel within the microfluidic channels of the chip, allowing the user to process 100's to 10,000's of the single cells in a short period of time. In some embodiments, within each partition reaction vesicle, a single cell is lysed and reverse transcription of polyadenylated mRNA occurs. As a result, cDNAs from a single cell from individual partitions will retain the same barcode, allowing sequencing reads from the cDNAs to be mapped back to their original single cell of origin in the tissue section or tissue sample.

[1335] In some embodiments, a method of associating or identifying a target biological analyte with a location within a biological sample can include providing a substrate with a plurality of capture probes (e.g., a capture probe array), contacting the capture-probe array with a plurality of analyte capture agents, extending the capture agent barcode domain of the analyte capture agent to include a sequence that is complementary to the sequence of the capture probe spatial barcode, thereby producing a partially-double stranded oligonucleotide including the capture domain, spatial barcode, and other optional domains of the capture probe, as well as the analyte capture sequence, analyte binding moiety barcode and optional domains of the capture agent barcode domain, contacting the one or more spatially-tagged analyte capture agents with the biological sample, and denaturing the spatially-tagged analyte capture agent from the capture probe. In some embodiments, the method further includes dissociating the biological sample into plurality of cells of the biological sample, and using the spatially-tagged analyte capture agent to identify the location of the target biological analyte within the biological sample. See, for example, FIG. 40 and FIGS. 41A and 41B.

(ii) Releasing an Analyte Capture Agent from an Array

[1336] In some embodiments, capture probes can be initially associated with (e.g., bound to covalently or non-covalently) an array and then released from the array. In some embodiments, the capture probe can be released from an array through activation of the capture probe cleavage domain, as described herein. In some embodiments, capture probes can be coupled with analyte capture agents via the capture domain of the capture probe, and the capture probes can denature from the spatially-tagged analyte capture agents allowing the capture probes to continue to be associated with the array.

[1337] For example, a cell surface protein of a cell can be associated with an analyte capture agent, and the cell can be subjected to spatial analysis (e.g., any of the variety of spatial analysis methods described herein). In some embodiments, the analyte capture agent bound to the cell surface protein can be bound to a capture probe (e.g., a capture probe on an array) via a capture domain that interacts with an analyte capture sequence present on the capture agent barcode domain of the analyte capture agent. The capture agent barcode domain (including the analyte binding moiety barcode) can be extended with a polymerase using a 3' end of the capture domain as a priming site, thus generating an extended capture agent barcode domain that includes the complementary sequence for all or part of the capture probe (including a spatial barcode) as well as the analyte binding moiety barcode (e.g., producing a spatially-tagged analyte capture agent). In some embodiments, the spatial array with the spatially-tagged analyte capture agents can be contacted with a sample, where the spatially-tagged analyte capture agent associated with the spatial array captures the biological analyte. The spatially-tagged analyte capture agent can then be denatured from the capture probe of the spatial array, allowing the spatial array to be reused. The sample can be dissociated into non-aggregated cells (e.g. single cells) and analyzed by the single cell/droplet methods described herein. The extended capture probe can be sequenced to obtain a nucleic acid sequence, in which the spatial barcode of the capture probe is associated with the analyte binding

moiety barcode of the analyte capture agent.

[1338] In some embodiments, the capture probe can be released from the array without denaturing from the spatially-tagged analyte capture agent. In some embodiments, release of the capture probes occurs upon application of a stimulus that initiates dissociation of the capture probes from the array. The stimulus can be a photo-stimulus, e.g., through cleavage of a photo-labile linkage that releases the capture probes, an elevation of the array's temperature (e.g., resulting in linkage cleavage or another release mechanism of the capture probes from the array). In some embodiments, a chemical stimulus can be used to cleave a linkage, or otherwise release the capture probes from the array. In some embodiments, such compositions include the polyacrylamide matrices, and can be degraded for release of the attached capture probes through exposure to a reducing agent, such as DTT. Additional, non-limiting stimulus examples can include, for example, a photo-stimulus, a mechanical stimulus, a radiation stimulus, a biological stimulus (e.g., enzyme), or any combination thereof, as discussed elsewhere herein.

(c) Multiplexed Analyte Analysis

[1339] In some embodiments, the arrays and methods described herein support multiplexed spatial analysis. In some embodiments, spatial analysis of a biological analyte can be performed individually for each analyte of interest. In some embodiments, multiple biological analytes can be detected and spatially analyzed simultaneously within the same biological sample. In some embodiments, multiplexing (e.g., simultaneously detecting multiple markers) allows for examination of spatial arrangement of analytes of interest (e.g., proteins, DNA, RNA) as well as analyte interaction and co-localization thereby facilitating simultaneous analysis of multiple tissue markers. In some embodiments, multiple biological analytes can be detected and spatially analyzed at different times (e.g., detection and analysis of one category of analyte, followed by detection and analysis of another category of analyte, or detection and analysis of one analyte followed by detection and analysis of another analyte). For example, detection, examination, and analysis of one or more proteins within a biological sample can occur prior to the detection, examination, and analysis of one or more mRNA transcripts.

[1340] In some embodiments, as described above, the analyte binding moiety barcode can be correlated with, associated with, or correspond to a particular analyte binding moiety, and thus a particular analyte. For example, analyte binding moiety A can correspond to analyte binding moiety barcode A, and analyte binding moiety B can correspond to analyte binding moiety barcode B. In this way, a number (n), of analytes can be examined by introducing (n) different analyte binding moiety/analyte binding moiety barcode pluralities to the sample. In some embodiments, analyte capture agents can be used in multiplexed methods to capture 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 or more analytes. In some embodiments, all analyte capture agent pluralities possess the same analyte capture sequence, and thus can be captured by a capture probe array, wherein the capture probes possess the same capture domain.

[1341] In some embodiments, as described above, the analyte capture sequence can be correlated with, associated with, or correspond to a particular analyte binding moiety, and thus a particular analyte. For example, analyte capture sequence A can correspond to analyte binding moiety barcode A, and analyte capture sequence B can correspond to analyte binding moiety barcode B. In this way, a number (n), analytes can be examined by introducing (n) different analyte binding moiety/analyte capture sequence pluralities to the sample. In some embodiments, the analyte binding moiety barcode and the analyte capture sequence can be correlated with, associated with, or correspond to a particular analyte binding moiety.

[1342] In some embodiments, a particular type or category of biological analytes is correlated with, associated with, or corresponds to a particular capture domain on a capture probe array. For example, a capture probe array can be generated wherein a plurality of the capture probes on the array possess a homopolymer (e.g., poly(T)) capture domain which corresponds to one type of

biological analyte (e.g., mRNA). Additionally, the same capture probe array can include a plurality of capture probes that possess a N-mer priming sequence capture domain for another type of biological analyte (e.g., genomic DNA), and the same capture probe array can include a plurality of capture probes that possess a capture domain that can specifically bind the sequence of an analyte capture sequence (e.g., corresponding to a third type of analyte, e.g., proteins).

[1343] In some embodiments of any of the spatial analysis methods described herein, multiple (e.g., two or more, three or more) classes (e.g. types, categories) of biological analytes can be concurrently measured. For example, a feature can include capture probes that can participate in an assay of at least three different types of analytes via three different capture domains. For example, a feature can be coupled to a barcoded capture probe that includes a capture domain that includes a poly(T) priming sequence for mRNA analysis; a barcoded capture probe that includes a capture domain that includes a random N-mer priming sequence for gDNA analysis; and a barcoded capture probe that includes a capture domain that can specifically bind an analyte capture agent (e.g., an antibody with a spatial barcode), via its analyte capture sequence.

[1344] In some embodiments of any of the spatial analysis methods described herein, other combinations of multiple biological analytes that can be concurrently measured include, without limitation: (a) mRNA, a lineage tracing construct, and cell surface and/or intracellular proteins and/or metabolites; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), and cell surface and/or intracellular proteins and/or metabolites; (c) mRNA, genomic DNA, and a perturbation reagent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (d) mRNA, accessible chromatin, and a perturbation reagent; (e) mRNA, an analyte capture agent (e.g., any of the MHC multimers described herein), and a perturbation reagent; (f) mRNA, cell surface and/or intracellular proteins and/or metabolites, and a perturbation agent; (g) mRNA, a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), and a perturbation reagent; (h) mRNA, an analyte capture agent, and a V(D)J sequence of an immune cell receptor; (i) cell surface and/or intracellular proteins and/or metabolites, a an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor; (j) methylation status, mRNA, and cell surface and/or intracellular proteins and/or metabolites; (k) mRNA, chromatin (e.g., spatial organization of chromatin in a cell), and a perturbation reagent; (l) a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell); and a perturbation reagent; and (m) mRNA, a V(D)J sequence of an immune cell receptor, and chromatin (e.g., spatial organization of chromatin in a cell), or any combination thereof.

[1345] In some embodiments, nucleic acids from a spatial array are amplified using any of the methods disclosed herein. The resulting amplicons can be subsequently assayed as set forth below. In some embodiments, amplicon barcodes are analyzed using multiplex spatial imaging. In some embodiments, amplicon barcodes can be analyzed by RNA sequential probing of targets (RNA SPOTs), sequential fluorescent in situ hybridization (SeqFISH), single-molecule fluorescent in situ hybridization (smFISH), multiplexed error-robust fluorescence in situ hybridization (MERFISH), in situ sequencing (ISS), targeted in situ sequencing, fluorescent in situ sequencing (FISSEQ), or spatially-resolved transcript amplicon readout mapping (STARmap). In some embodiments, amplicon barcodes are analyzed by sequential hybridization and detection with a plurality of labelled probes. A variety of light-based sequencing technologies are known in the art (Landegren et al., *Genome Res.* 8:769-76 (1998); Kwok, *Pharmacogenomics* 1:95-100 (2000); and Shi, *Clin. Chem.* 47:164-172 (2001), each herein incorporated by reference in its entirety). In some embodiments, the methods provided herein involves capturing a target nucleic acid using barcoded capture probes, and optionally generating cDNA molecules that include the barcodes from the capture probes. In some embodiments, labeled probes (e.g., fluorescently labeled probes, or fluorophore-conjugated detection oligonucleotides) are used to sequence the barcodes from the capture probes, thereby visualizing the location of the target nucleic acid. In some embodiments, a

labeled probe is labeled with a detectable moiety. In some embodiments, a labeled probe includes one detectable moiety. In some embodiments, a labeled probe includes two or more detectable moieties. In some embodiments, a detectable moiety is or includes a fluorophore. In some embodiments, the barcodes from the capture probes are sequenced by introducing fluorophores in a base-specific manner. In some embodiments, an image is taken after each addition of a fluorophore. In some embodiments, provided methods include a step of removing labeled probes after each imaging step. In some embodiments, a step of removing includes degrading the labeled probes. In some embodiments, a step of removing does not significantly degrade a labeled probe. In some embodiments, labeled probes (e.g., fluorescently labeled probes, or fluorophore-conjugated detection oligonucleotides) are used to visualize the location of a target RNA directly, without the generation of cDNA molecules.

[1346] Exemplary probes labeled with fluorophores include but are not limited to probes for fluorescence in situ hybridization (FISH). Widely known and practiced by persons having ordinary skill in the art, FISH is used to, among other things, detect and localize the presence or absence of specific DNA sequences or RNA targets. Methods for designing and preparing labeled probes are widely known in the art, including but not limited to those described in US 2012/0142014.

XI. Analysis of Spatial Heterogeneity in a Biological Sample

[1347] Spatial heterogeneity in developing systems has typically been studied via RNA hybridization, immunohistochemistry, fluorescent reporters, or purification or induction of pre-defined subpopulations and subsequent genomic profiling (e.g., RNA-seq). Such approaches, however, rely on a relatively small set of pre-defined markers, therefore introducing selection bias that limits discovery. These prior approaches also rely on a priori knowledge. Spatial RNA assays traditionally relied on staining for a limited number of RNA species. In contrast, single-cell RNA-sequencing allows for deep profiling of cellular gene expression (including non-coding RNA), but the established methods separate cells from their native spatial context. Disclosed herein are methods of determining the spatial heterogeneity of a sample by detecting one or more biological analytes in a biological sample.

[1348] Subjects from which biological samples can be obtained can be healthy or asymptomatic individuals, individuals that have or are suspected of having a disease (e.g., a patient with a disease such as cancer) or a pre-disposition to a disease, and/or individuals that are in need of therapy or suspected of needing therapy. Biological samples can include one or more diseased cells. A diseased cell can have altered metabolic properties, gene expression, protein expression, and/or morphologic features. Examples of diseases include inflammatory disorders, metabolic disorders, nervous system disorders, and cancer. Cancer cells can be derived from solid tumors, hematological malignancies, cell lines, or obtained as circulating tumor cells.

[1349] A biological sample can have regions that show morphological feature(s) that may indicate the presence of disease or the development of a disease phenotype. For example, morphological features at a specific site within a tumor biopsy sample can indicate the aggressiveness, therapeutic resistance, metastatic potential, migration, stage, diagnosis, and/or prognosis of cancer in a subject. A change in the morphological features at a specific site within a tumor biopsy sample often correlate with a change in the level or expression of an analyte in a cell within the specific site, which can, in turn, be used to provide information regarding the aggressiveness, therapeutic resistance, metastatic potential, migration, stage, diagnosis, and/or prognosis of cancer in a subject. A region or area within a biological sample that is selected for specific analysis (e.g., a region in a biological sample that has morphological features of interest) is often described as “a region of interest.”

[1350] Variant haplotype detection is used to identify heterozygous cells, such that this technique provides valuable data during single cell studies. However, in combination with spatial analysis within a biological sample, variant haplotype detection can further provide novel information on the distribution of heterozygous cells in biological samples (e.g., tissues) affected by or exhibiting a

variety of biological conditions. These data are of interest due to the potential to reveal causal relationships between variant haplotypes and disease outcomes, to aid in identification of disease-associated variants, or to reveal heterogeneity within a biological sample.

[1351] In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domains released from the analyte capture agents can then be subjected to sequence analysis to identify which analyte capture agents were bound to analytes. Based upon the capture agent barcode domains that are associated with a feature (e.g., a feature at a particular location) on a spatial array and the presence of the analyte binding moiety barcode sequence, an analyte profile can be created for a biological sample. Profiles of individual cells or populations of cells can be compared to profiles from other cells, e.g., 'normal' cells, to identify variations in analytes, which can provide diagnostically relevant information. In some embodiments, these profiles can be useful in the diagnosis of a variety of disorders that are characterized by variations in cell surface receptors, such as cancer and other disorders.

(i) Blocking Probes

[1352] In some embodiments, an analyte capture sequence of a capture agent barcode domain is blocked prior to adding the analyte capture agent to a biological sample. In some embodiments, an analyte capture sequence of a capture agent barcode domain is blocked prior to adding the analyte capture agent to a capture probe array. In some embodiments, blocking probes are added to blocking buffer or other solutions applied in an IHC and/or IF protocol. In some embodiments, a blocking probe is used to block or modify the free 3' end of the capture agent barcode domain. In some embodiments, a blocking probe is used to block or modify the free 3' end of the analyte capture sequence of the capture agent barcode domain. In some embodiments, a blocking probe can be hybridized to the analyte capture sequence of a capture agent barcode domain to mask the free 3' end of the capture agent barcode domain. In some embodiments, a blocking probe can be a hairpin probe or partially double stranded probe. In some embodiments, the free 3' end of the analyte capture sequence of the capture agent barcode domain can be blocked by chemical modification, e.g., addition of an azidomethyl group as a chemically reversible capping moiety such that the capture probes do not include a free 3' end. Blocking or modifying the capture agent barcode domains, particularly at the free 3' end of the capture agent barcode domain, prior to contacting the analyte capture agents with the substrate, prevents binding of the analyte capture sequence to capture probe capture domain (e.g., prevents the binding of an analyte capture sequence poly(A) tail to a poly(T) capture domain).

[1353] In some embodiments, a blocking probe is used to block or modify the free 3' end of a capture probe. In some embodiments, a blocking probe is used to block or modify the free 3' end of a capture probe capture domain. In some embodiments, the analyte capture sequence is blocked prior to adding the analyte capture agent to a capture probe array. In some embodiments, blocking probes are added to blocking buffer or other solutions applied in an IHC and/or IF protocol. In some embodiments, a blocking probe can be hybridized to the capture domain to mask the free 3' end of the capture domain. In some embodiments, a blocking probe can be a hairpin probe or partially double stranded probe. In some embodiments, the free 3' end of the capture domain can be blocked by chemical modification, e.g., addition of an azidomethyl group as a chemically reversible capping moiety such that the capture probes do not include a free 3' end. Blocking or modifying the capture domains, particularly at the free 3' end of the capture domain, prior to contacting the analyte capture agents with the capture probe array, prevents binding of the analyte capture sequence to capture probe capture domain (e.g., prevents the binding of an analyte capture sequence poly(A) tail to a poly(T) capture domain).

[1354] In some embodiments, the blocking probes can be reversibly removed. For example, blocking probes can be applied to block the free 3' end of either or both the capture agent barcode domain and/or the capture probes. Blocking interaction between the analyte capture agent and the capture probe array can reduce non-specific background staining in IHC and/or IF applications.

After the analyte binding agents are bound to the biological analyte, the blocking probes can be removed from the 3' end of the capture agent barcode domain and/or the capture probe, and the analyte-bound analyte binding agents can migrate to and become bound by the capture probe array. In some embodiments, the removal includes denaturing the blocking probe from the analyte binding moiety barcode and/or capture probe. In some embodiments, the removal includes removing a chemically reversible capping moiety. In some embodiments, the removal includes digesting the blocking probe with an RNase (e.g., RNase H).

[1355] In some embodiments, the nucleic acid sequence of the analyte capture sequence includes a sequence that is substantially complementary to the blocking probe. "Substantially complementary to the blocking probe" includes a nucleic acid that is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% complementary to the blocking probe. In some embodiments, the nucleic acid sequence of the analyte capture sequence includes a sequence that is completely complementary to the blocking probe.

[1356] In some embodiments, the blocking probes are oligo (dT) blocking probes. In some embodiments, the oligo (dT) blocking probes can have a length of 15-30 nucleotides. In some embodiments, the oligo (dT) blocking probes can have a length of 10-50 nucleotides, e.g., 10-50, 10-45, 10-40, 10-35, 10-30, 10-25, 10-20, 10-15, 15-50, 15-45, 15-40, 15-35, 15-30, 15-25, 15-20, 20-50, 20-45, 20-40, 20-35, 20-30, 20-25, 25-50, 25-45, 25-40, 25-35, 25-30, 30-50, 30-45, 30-40, 30-35, 35-50, 35-45, 35-40, 40-50, 40-45, or 45-50 nucleotides. In some embodiments, the analyte capture agents can be blocked at different temperatures (e.g., 4° C. and 37° C.). In some embodiments, the analyte capture agents can be blocked from binding to the capture probes more effectively at lower temperatures when using shorter blocking probes.

EXEMPLARY EMBODIMENTS

Exemplary Embodiments A

[1357] Embodiment A1 is a method for profiling at least one biological analyte present in a cell, comprising: (a) providing a solid support comprising a plurality of features, wherein each feature on the solid support comprises a plurality of capture probes unique to the feature of origin, wherein each capture probe comprises a barcode and a capture domain; (b) distributing a plurality of cells onto the solid support such that at least one cell occupies a distinct feature on the solid support; (c) lysing cells such that a biological analyte from the at least one cell is released and bound by a capture probe associated with a distinct feature of the solid support; (d) detecting the bound biological analyte; and (e) correlating the bound biological analyte with the barcode of the bound capture probe associated with the distinct feature of the solid support, thus profiling the biological analyte as having been present in the at least one cell.

[1358] Embodiment A2 is the method of embodiment A1, wherein the plurality of cells are from a suspension of cells.

[1359] Embodiment A3 is the method of embodiment A1, where the plurality of cells are from disassociated tissue or tissue section.

[1360] Embodiment A4 is the method of embodiment A1, wherein the plurality of cells are from a cell culture.

[1361] Embodiment A5 is the method of any one of embodiments A1-A4, wherein the cells from the plurality of cells are immobilized on the solid support after they are flowed onto the solid support.

[1362] Embodiment A6 is the method of embodiment A5, wherein the cells from the plurality of cells are immobilized on the solid support prior to lysing.

[1363] Embodiment A7 is the method of embodiment A5 or embodiment A6, wherein the cells from the plurality of cells are immobilized by adding a hydrogel.

[1364] Embodiment A8 is the method of embodiment A7, wherein the hydrogel comprises a material comprising collagen, fibrin, alginate, polyacrylamide, polyethylene glycol, hyaluronic

acid, a polypeptide, or a combination of two or more thereof.

[1365] Embodiment A9 is the method of any one of embodiments A1-A6, wherein the cells from the plurality of cells are immobilized by applying an electric field.

[1366] Embodiment A10 is the method of any one of embodiments A1-A9, further comprising imaging one or more cells on the solid support.

[1367] Embodiment A11 is the method of embodiment A10, wherein imaging is performed prior to lysing the cells.

[1368] Embodiment A12 is the method of embodiment A10 or embodiment A11, wherein imaging is used to determine the morphology of a cell.

[1369] Embodiment A13 is the method of embodiment A12, wherein the morphology of a cell from the plurality of cells is correlated to the at least one biological analyte.

Exemplary Embodiments B

[1370] Embodiment B1 is a method for spatial profiling at least one biological analyte present in a cell culture, comprising: (a) providing a solid support comprising a plurality of capture probes, wherein each capture probe comprises a spatial barcode and a capture domain, and wherein the solid support comprises a cell-permissive coating; (b) culturing cells on the cell-permissive coating (c) permeabilizing cells of the cell culture such that at least one biological analyte is released and bound by a capture probe; (d) detecting the at least one biological analyte; and (e) correlating the at least one biological analyte with a spatial location in the cell culture.

[1371] Embodiment B2 is the method of embodiment B1, wherein the cell-permissive coating comprises at least one of a biological material and a synthetic material.

[1372] Embodiment B3 is the method of embodiment B1 or B2, wherein the cell-permissive coating comprises an extracellular matrix component.

[1373] Embodiment B4 is the method of embodiment B3, wherein the cell-permissive coating comprises one or more ECM components selected from the group consisting of: collagen Type I, collagen Type II, collagen Type IV, elastin, fibronectin, laminin, and vitronectin.

[1374] Embodiment B5 is the method of any one of embodiments B1-B4, wherein the cell-permissive coating comprises a synthetic material selected from the group consisting of: polylysine, poly-L-ornithine, and a biocompatible silicone (e.g., CYTOSOFT®).

[1375] Embodiment B6 is the method of any one of embodiments B1-B5, wherein the cell-permissive coating comprises a solubilized basement membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma.

[1376] Embodiment B7 is the method of any one of embodiments B1-B6, wherein the cell-permissive coating comprises collagen.

[1377] Embodiment B8 is the method of any one of embodiments B1-B7, wherein the cell culture comprises adherent cells.

[1378] Embodiment B9 is the method of embodiment B8, wherein the adherent cells are from cell lines selected from the group consisting of: BT549, HS 578T, MCF7, MDA-MB-231, MDAMB-468, T-47D, SF268, SF295, SF539, SNB-19, SNB-75, U251, Colo205, HCC 2998, HCT-116, HCT-15, HT29, KM12, SW620, 786-0, A498, ACHN, CAKI, RXF 393, SN12C, TK-10, UO-31, A549, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H460, NCI-H522, LOX IMVI, M14, MALME-3M, MDA-MB-435, SK-, EL-2, SK-MEL-28, SK-MEL-5, UACC-257, UACC-62, IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3, NCI-ADR-RES, DU145, PC-3, DU145, H295R, HeLa, KBM-7, LNCaP, MCF-7, MDA-MB-468, PC3, SaOS-2, SH-SY5Y, T-47D, THP-1, U87, vero, MC3T3, GH3, PC12, dog MDCK kidney epithelial, *Xenopus* A6 kidney epithelial, zebrafish AB9, and Sf9 insect epithelial cell lines.

[1379] Embodiment B10 is the method of any one of embodiments B1-B9, further comprising imaging the cell culture.

[1380] Embodiment B11 is the method of embodiment B10, wherein the imaging is performed prior to permeabilizing cells of the cell culture.

[1381] Embodiment B12 is the method of embodiment B10, wherein the imaging is performed after permeabilizing cells of the cell culture.

[1382] Embodiment B13 is the method of any one of embodiments B10-B12, wherein the imaging is used to determine a region of interest in the cell culture.

[1383] Embodiment B14 is the method of any one of embodiments B10-B13, wherein the imaging is used to determine the morphology of the cell culture or a region of the cell culture.

[1384] Embodiment B15 is the method of embodiment B14, wherein the morphology of the cell culture or a region of the cell culture is correlated to the spatial location of the at least one biological analyte.

[1385] Embodiment B16 is the method of any one of embodiments B1-B15, further comprising fixing the cells prior to permeabilization.

[1386] Embodiment B17 is the method of embodiment B16, wherein the cells in the cell culture are fixed and permeabilized prior to releasing the at least one biological analyte from the biological sample.

[1387] Embodiment B18 is the method of any one of embodiments B1-B17, wherein the at least one biological analyte is DNA.

[1388] Embodiment B19 is the method of any one of embodiments B1-B17, wherein the at least one biological analyte is RNA.

[1389] Embodiment B20 is the method of embodiment B19, wherein the capture domain comprises a poly-dT sequence.

[1390] Embodiment B21 is the method of embodiment B19 or B20, wherein the capture domain is configured to hybridize to a poly-A tail of an mRNA.

[1391] Embodiment B22 is the method of embodiment B1, wherein the capture probe further comprises a unique molecular identifier (UMI).

[1392] Embodiment B23 is the method of embodiment B1, wherein the capture probe further comprises a functional domain.

[1393] Embodiment B24 is the method of embodiment B1, wherein the capture probe further comprises a cleavage domain.

Exemplary Embodiments C

[1394] Embodiment C.sub.1 is a method for temporal and spatial profiling of at least one species of biological analyte present in a biological sample, comprising: (a) providing a solid support comprising a plurality of capture probes, wherein the capture probes comprise a spatial barcode and a capture domain; (b) contacting the biological sample with the solid support comprising the capture probes; (c) releasing at least one member of the species of biological analyte from the biological sample; (d) contacting the solid support comprising the capture probes with at least one released member of the species of biological analyte such that the at least one released member of the species of biological analyte interacts with at least one of the plurality of capture probes; (e) detecting the at least one released member of the species of biological analyte; (f) correlating the at least one released member of the species of biological analyte with a spatial location in the biological sample; and (g) repeating steps (a)-(e) after a period of time.

[1395] Embodiment C2 is the method of embodiment C1, wherein the biological sample is in a controlled environment.

[1396] Embodiment C3 is the method of embodiment C1, further comprising imaging the biological sample.

[1397] Embodiment C4 is the method of embodiment C3, wherein imaging is performed prior to releasing at least one member of the species of biological analyte from the biological sample.

[1398] Embodiment C5 is the method of embodiment C3, wherein imaging is performed after releasing the at least one member of the species of biological analyte from the biological sample.

[1399] Embodiment C6 is the method of embodiment C3, wherein imaging is used to determine a region of interest in the biological sample.

[1400] Embodiment C7 is the method of embodiment C3, wherein the imaging comprises using fiducial markers.

[1401] Embodiment C8 is the method of embodiment C1, wherein the step of releasing comprises applying an electric field to the biological sample.

[1402] Embodiment C9 is the method of embodiment C5, wherein the electric field is applied as a short burst.

[1403] Embodiment C10 is the method of embodiment C1, wherein the period of time is about 24 hours.

[1404] Embodiment C11 is the method of embodiment C1, wherein step (g) is repeated one, two, three, or four times.

[1405] Embodiment C12 is the method of embodiment C1, further comprising generating a temporal profile for the at least one member of the species of biological analyte.

[1406] Embodiment C13 is the method of embodiment C12, wherein the same species of biological analyte is detected at two or more time points.

[1407] Embodiment C14 is the method of embodiment C1, wherein prior to the step of repeating, the biological sample is subjected to one or more perturbants.

[1408] Embodiment C15 is the method of embodiment C1, wherein the step of releasing comprises releasing under conditions that permit cells in the biological sample to live.

[1409] Embodiment C16 is a method for determining the effect of one or more perturbants on a biological sample, comprising: (a) contacting the biological sample with the one or more perturbants; and (b) performing the method for temporal and spatial profiling of embodiment C1.

[1410] Embodiment C17 is the method of embodiment C16, wherein the one or more perturbants include a small molecule, an antibody, a nucleic acid, a peptide, light, heat, pressure, an electric field, and combinations thereof.

[1411] Embodiment C18 is the method of embodiment C16, wherein the perturbant comprises a DNA modification system.

[1412] Embodiment C19 is the method of embodiment C18, wherein the DNA modification system is a CRISPR system.

[1413] Embodiment C20 is the method of embodiment C18, wherein the DNA modification system employs a single guide RNA (sgRNA).

[1414] Embodiment C21 is the method of embodiment C2, wherein the controlled environment comprises a flow cell.

[1415] Embodiment C22 is the method of embodiment C2, wherein the controlled environment comprises a tissue culture system.

[1416] Embodiment C23 is the method of embodiment C2, wherein the controlled environment comprises a tissue bath.

[1417] Embodiment C24 is the method of embodiment C1, wherein the at least one released biological analyte is a nucleic acid, wherein the capture domain is a nucleic acid, and wherein the at least one released biological analyte hybridizes to the capture domain.

Exemplary Embodiments D

[1418] Embodiment D1 is a method for generating single-cell partitions from a cell-containing biological sample, comprising: (a) providing a solid support, wherein the solid support comprises a conditionally removable coating; (b) contacting the cell-containing biological sample with the solid support; (c) adding a solution comprising a plurality of photo-crosslinkable polymer precursors, wherein the solution covers the solid support and the biological sample; (d) cross-linking members of the plurality of photo-crosslinkable polymer precursors to generate a plurality of cross-linked voxels; and (e) dissociating the cross-linked voxels by treating the conditionally removable coating with a release agent, thereby generating single-cell partitions.

[1419] Embodiment D2 is the method of embodiment D1, wherein the conditionally removable coating is a DTT sensitive hydrogel.

[1420] Embodiment D3 is the method of embodiment D2, wherein the release agent is DTT.

[1421] Embodiment D4 is the method of embodiment D1 or D2, wherein the biological sample and solid support are assembled in a flow cell prior to the addition of the solution comprising a photo-crosslinkable polymer precursor.

[1422] Embodiment D5 is the method of embodiment D4, wherein the flow cell is dismantled after crosslinking the cells of the biological sample.

[1423] Embodiment D6 is the method of any one of embodiments D1-D5, wherein the cross-linked cells on the solid support are washed.

[1424] Embodiment D7 is the method of any one of embodiments D1-D6, wherein the step of cross-linking comprises subjecting members of the plurality of photo-crosslinkable polymer precursors to a light source.

[1425] Embodiment D8 is the method of embodiment D7, wherein members of the plurality of photo-crosslinkable polymer precursors are subjected to a light source having a spatial pattern.

[1426] Embodiment D9 is the method of embodiment D8, wherein subjecting members of the plurality of photo-crosslinkable polymer precursors to a light source having a spatial pattern results in the generation of at least one voxel that comprises a single cell from the biological sample.

[1427] Embodiment D10 is the method of embodiment D8, wherein the spatial pattern is generated by preventing a portion of the biological sample from being subjected to the light source.

[1428] Embodiment D11 is the method of embodiment D8, wherein the spatial pattern is generated by selectively shining the light source on one or more areas of the biological sample.

[1429] Embodiment D12 is the method of any one of embodiments D8-D11, wherein the biological sample is imaged.

[1430] Embodiment D13 is the method of embodiment D12, wherein the imaged biological sample is used to generate the spatial pattern of the light source.

[1431] Embodiment D14 is the method of embodiment D13, wherein the imaged biological sample reveals the cellular pattern of the biological sample on the solid support, and wherein the spatial pattern of the light source corresponds to the cellular pattern of the biological sample.

[1432] Embodiment D15 is the method of any one of embodiments D1-D14, wherein the solid support comprises a plurality of features on the solid support, wherein each feature comprises a plurality of capture probes comprising a spatial barcode and a capture domain, and wherein at least one spatial barcode corresponds to a particular feature associated with a distinct spatial position on the solid support.

[1433] Embodiment D16 is the method of embodiment D15, wherein the method comprises contacting the biological sample to the solid support such that at least one biological analyte from the biological sample is bound by the at least one capture probe, determining the identity of the spatial barcode of the at least one capture probe to identify the distinct spatial position of the associated feature on the solid support, and correlating the distinct spatial position of the associated feature on the solid support with the spatial location of the biological analyte in the biological sample.

[1434] Embodiment D17 is the method of embodiment D16, wherein members of the plurality of capture probes are cleaved from the plurality of features via the cleavage domain such that the cleaved capture probes are introduced into the biological sample and at least one cleaved capture probe interacts with at least one biological analyte present in the biological sample via the capture domain.

[1435] Embodiment D18 is the method of embodiment D17, wherein the members of the plurality of capture probes are cleaved from the plurality of features prior to adding a solution comprising a plurality of photo-crosslinkable polymer precursors.

[1436] Embodiment D19 is the method of any one of embodiments D16-D18, wherein the spatial location of the biological analyte in the biological sample is resolved to a single cell present in one of the members of the plurality of cross-linked voxels.

- [1437] Embodiment D20 is the method of any one of embodiments D15-D19, wherein the at least one biological analyte is DNA.
- [1438] Embodiment D21 is the method of any one of embodiments D15-D19, wherein the at least one biological analyte is RNA.
- [1439] Embodiment D22 is the method of embodiment D21, wherein the capture domain comprises a poly-dT sequence.
- [1440] Embodiment D23 is the method of embodiment D21 or D22, wherein the capture domain is configured to hybridize to a poly-A tail of an mRNA.
- [1441] Embodiment D24 is the method of any one of embodiments D15-D23, wherein the capture probe further comprises a unique molecular identifier (UMI).
- [1442] Embodiment D25 is the method of any one of embodiments D15-D24, wherein the capture probe further comprises a functional domain.
- [1443] Embodiment D26 is the method of any one of embodiments D17-D25, wherein the capture probe further comprises a cell-penetrating agent.
- [1444] Embodiment D27 is the method of embodiment D26, wherein the cell-penetrating agent comprises a cell-penetrating peptide domain.
- [1445] Embodiment D28 is the method of embodiment D26, wherein the cell-penetrating peptide domain mediates introduction of the capture probe into the cell.
- [1446] Embodiment D29 is the method of any one of embodiments D17-D28, wherein the solid support is removed from the biological sample after releasing the plurality of capture probes from the features.
- [1447] Embodiment D30 is the method of any one of embodiments D17-D29, wherein the capture probes further comprise a cell penetration tag.
- [1448] Embodiment D31 is the method of embodiment D30, wherein the cell penetration tag comprises a fluorescent dye.
- [1449] Embodiment D32 is the method of embodiment D31, wherein the cell penetration tag comprises at least one of Cy2, Cy3, Cy3B, Cy3.5, Cy5, Cy5.5, Cy7, fluorescein (6-FAM), DyLight, Alexa Fluor®, and tetramethylrhodamine (TAMRA) azide.
- [1450] Embodiment D33 is the method of any one of embodiments D30-D32, further comprising detecting the cell penetration tag.
- [1451] Embodiment D34 is the method embodiment D33, wherein the cell penetration tag is detected after cleaving the plurality of capture probes from the features and removing the solid support from the biological sample.
- [1452] Embodiment D35 is the method of embodiment D34, wherein introduction of the capture probe into the cell is determined by detecting the presence of the cell penetration tag in the cell.
- [1453] Embodiment D36 is the method of any one of embodiments D17-D35, wherein each capture probe further comprises an intracellular cleavage domain.
- [1454] Embodiment D37 is the method of embodiment D36, wherein the intracellular cleavage domain comprises a disulfide bond.
- [1455] Embodiment D38 is the method of embodiment D36 or D37, wherein the intracellular cleavage domain is cleaved after the capture probe is introduced into the cell.
- [1456] Embodiment D39 is the method of any one of embodiments D15-D38, wherein the dissociated single cells are analyzed using high-throughput instrumentation.

Exemplary Embodiments E

- [1457] Embodiment E1 is a method of determining process bias in a spatial analysis workflow comprising: (a) providing a substrate comprising a plurality of test analytes, wherein members of the plurality of test analytes are disposed on the substrate in a known amount and in a known location; (b) contacting the substrate with a solid support comprising a plurality of capture probes under conditions wherein at least one capture probe interacts with at least one test analyte, wherein the capture probes comprise a spatial barcode and a capture domain; (c) detecting the test analyte

that interacts with the capture probe; and (d) determining whether the spatial analysis workflow accurately detects the presence, amount, location, or combinations thereof, of the test analyte.

[1458] Embodiment E2 is the method of embodiment E1, wherein the substrate comprises a semi-porous material.

[1459] Embodiment E3 is the method of embodiment E2, wherein the semi-porous material is selected from the group consisting of: a nitrocellulose membrane, a hydrogel, a nylon filter, and combinations thereof.

[1460] Embodiment E4 is the method of any one of embodiments E1-E3, wherein the test analyte is selected from the group consisting of: nucleic acid, a protein, a lipid, and combinations thereof.

[1461] Embodiment E5 is the method of embodiment E4, wherein the test analyte is DNA.

[1462] Embodiment E6 is the method of embodiment E4, wherein the test analyte is RNA.

[1463] Embodiment E7 is the method of embodiment E1, wherein the capture probe comprises a capture domain.

[1464] Embodiment E8 is the method of embodiment E1, wherein the capture probe comprises a spatial barcode.

[1465] Embodiment E9 is the method of embodiment E1, wherein the capture probe comprises a unique molecular identifier (UMI).

[1466] Embodiment E10 is the method of embodiment E1, wherein the capture probe comprises a functional domain.

[1467] Embodiment E11 is the method of embodiment E1, wherein the capture probe comprises a cleavage domain.

[1468] Embodiment E12 is the method of embodiment E7, wherein the capture domain of the capture probe comprises a poly-dT.

[1469] Embodiment E13 is the method of embodiment E7, wherein the capture domain of the capture probe is configured to hybridize to a poly-A tail of an mRNA.

[1470] Embodiment E14 is the method of embodiment E1, wherein the plurality of test analytes are disposed on the substrate in a defined pattern.

[1471] Embodiment E15 is the method of embodiment E9, wherein the defined pattern comprises one or more spots.

[1472] Embodiment E16 is the method of embodiment E1, wherein the test analytes comprise nucleic acids, and wherein the nucleic acid sequence of a first test analyte differs from the nucleic acid sequence of a second test analyte by a single nucleic acid residue.

[1473] Embodiment E17 is the method of embodiment E16, further comprising determining whether the spatial analysis workflow accurately detects the presence, amount, location, or combinations thereof, of the first test analyte, the second test analyte, or both.

[1474] Embodiment E18 is the method of embodiment E1, wherein the test analytes comprise nucleic acids, and wherein the concentration of a first test analyte differs from the concentration of a second test analyte at a different location on the substrate.

[1475] Embodiment E19 is the method of embodiment E18, further comprising determining whether the spatial analysis workflow accurately detects the presence, amount, location, or combinations thereof, of the first test analyte, the second test analyte, or both.

[1476] Embodiment E20 is the method of embodiment E18, wherein the first test analyte and the second test analyte comprise an identical nucleotide sequence.

[1477] Embodiment E21 is the method of any one of embodiments E1-E19, wherein a location on the substrate comprises a single species of test analyte.

[1478] Embodiment E22 is the method of any one of embodiments E1-E21, wherein a location on the substrate comprises two or more species of test analytes.

[1479] Embodiment E23 is the method of any one of embodiments E1-E22, wherein a location on the substrate further comprises at least one surface marker protein.

[1480] Embodiment E24 is the method of embodiment E6, wherein a location on the substrate

comprises at least one CRISPR guide RNA.

[1481] Embodiment E25 is the method of embodiment E4, wherein the test analyte is a bacterial artificial chromosome cross-linked to a protein.

Exemplary Embodiments F

[1482] Embodiment F1 is a method for spatially profiling a plurality of analytes in a biological sample, comprising: (a) providing an isometrically expanded biological sample in a hydrogel matrix; (b) contacting the isometrically expanded biological sample in a hydrogel matrix with a solid substrate comprising a plurality of capture probes, wherein each capture probe comprises a spatial barcode and a capture domain, wherein one or more analytes present in the isometrically expanded biological sample are captured by one or more of the capture probes; and (c) determining the spatial profile of the one or more captured analytes in the biological sample.

[1483] Embodiment F2 is a method for spatially profiling a plurality of analytes in a biological sample, comprising: (a) immobilizing the biological sample on a solid substrate; (b) embedding the biological sample in a hydrogel matrix, thus generating a biological sample-containing hydrogel matrix; (c) isometrically expanding the biological sample-containing hydrogel matrix, thus generating an isometrically expanded biological sample; (d) contacting the isometrically expanded biological sample with a solid substrate comprising a plurality of capture probes, wherein each capture probe comprises a spatial barcode and a capture domain, wherein one or more analytes present in the isometrically expanded biological sample are captured by one or more capture probes; and (e) determining the spatial profile of the one or more captured analytes in the biological sample.

[1484] Embodiment F3 is the method of embodiment F1 or F2, wherein the analytes are immobilized in the hydrogel matrix.

[1485] Embodiment F4 is the method of embodiment F3, wherein the analytes are immobilized by cross-linking.

[1486] Embodiment F5 is the method of embodiment F1 or F2, wherein the analytes in the biological sample are migrated to the solid substrate comprising a plurality of capture probes.

[1487] Embodiment F6 is the method of embodiment F5, wherein the migrating comprises passive migration.

[1488] Embodiment F7 is the method of embodiment F5, wherein the migrating comprises active migration.

[1489] Embodiment F8 is the method of embodiment F1 or F2, further comprising providing a cell labelling agent to the biological sample, wherein the cell labelling agent comprises an analyte capture domain and a molecular tag barcode.

[1490] Embodiment F9 is the method of embodiment F8, wherein the cell labelling agent binds the biological analyte in the biological sample.

[1491] Embodiment F10 is the method of embodiment F9, further comprising cleaving the molecular tag barcode from the cell labelling agent and allowing the molecular tag barcode to migrate to the solid substrate comprising a plurality of capture probes.

[1492] Embodiment F11 is the method of embodiment F10, wherein the migration comprises passive migration.

[1493] Embodiment F12 is the method of embodiment F10, wherein the migration comprises active migration.

[1494] Embodiment F13 is the method of embodiment F1 or F2, wherein the biological sample is isometrically expanded about $1.2\times$ its original size.

[1495] Embodiment F14 is the method of embodiment F1 or F2, wherein the biological sample is isometrically expanded about $1.5\times$ its original size.

[1496] Embodiment F15 is the method of embodiment F1 or F2, wherein the biological sample is isometrically expanded about $1.7\times$ its original size.

[1497] Embodiment F16 is the method of embodiment F1 or F2, wherein the biological sample is

isometrically expanded to a size of about $2\times$ its original size.

[1498] Embodiment F17 is the method of embodiment F1 or F2, wherein the biological sample is isometrically expanded to a size of about $3\times$ its original size.

[1499] Embodiment F18 is the method of embodiment F1 or F2, wherein the biological sample is isometrically expanded by hydration.

[1500] Embodiment F19 is the method of embodiment F1 or F2, wherein the resolution of the spatial profiling is increased as compared to a corresponding method performed on a biological sample that has not been isometrically expanded.

[1501] Embodiment F20 is the method of embodiment F2, further comprising assembling the immobilized biological sample on a solid substrate in a flow cell.

[1502] Embodiment F21 is the method of embodiment F2, further comprising removing the isometrically expanded biological sample from the solid substrate prior to contacting the solid substrate comprising a plurality of capture probes.

[1503] Embodiment F22 is the method of embodiment F1 or F2, wherein the capture probe comprises a unique molecular identifier.

[1504] Embodiment F23 is the method of embodiment F1 or F2, wherein the capture probe comprises a cleavage domain.

[1505] Embodiment F24 is the method of embodiment F1 or F2, wherein the capture probe comprises a functional domain.

[1506] Embodiment F25 is the method of embodiment F1 or F2, wherein the capture probe comprises a cleavage domain, a spatial barcode, a unique molecular identifier, a functional domain, and a capture domain.

[1507] Embodiment F26 is the method of embodiment F1 or F2, wherein the biological sample is a tissue sample.

[1508] Embodiment F27 is the method of embodiment F26, wherein the tissue sample is a fresh-frozen tissue sample.

[1509] Embodiment F28 is the method of embodiment F26, wherein the tissue sample comprises a tumor cell.

[1510] Embodiment F29 is the method of embodiment F26, wherein the tissue sample comprises a tissue section.

[1511] Embodiment F30 is the method of embodiment F1 or F2, wherein the biological sample is a preserved cell or tissue.

[1512] Embodiment F31 is the method of embodiment F1 or F2, further comprising imaging the biological sample.

[1513] Embodiment F32 is the method of embodiment F31, wherein the imaging is performed prior to expansion.

[1514] Embodiment F33 is the method of embodiment F31, wherein the imaging is performed after expansion.

[1515] Embodiment F34 is the method of embodiment F31, wherein the imaging is used to determine a region of interest in the biological sample.

[1516] Embodiment F35 is the method of embodiment F31, wherein the imaging comprises using fiducial markers.

[1517] Embodiment F36 is the method of embodiment F1 or F2, wherein the analytes comprise at least one of RNA, DNA, protein, lipid, peptide, metabolite, small molecule, and a cell labeling agent.

Exemplary Embodiments G

[1518] Embodiment G1 is a method for spatial profiling at least one biological analyte present in a biological sample, comprising: (a) providing a biological sample; (b) contacting the biological sample with a solid support comprising a plurality of capture probes, wherein each capture probe comprises a cleavage domain, a spatial barcode, and a capture domain; (c) cleaving at least one

capture probe from the solid support via the cleavage domain such that the cleaved capture probe is introduced into the biological sample and at least one cleaved capture probe interacts with at least one biological analyte present in the biological sample via the capture domain; (d) dissociating the biological sample into a plurality of single cells, wherein the at least one cleaved capture probe that interacts with the at least one biological analyte is present in one of the dissociated cells; (e) detecting the at least one biological analyte present in the dissociated cell; and (f) correlating the at least one biological analyte with a spatial location in the biological sample.

[1519] Embodiment G2 is a method for spatial profiling at least one biological analyte present in a biological sample, comprising: (a) providing a biological sample; (b) contacting the biological sample with a solid support comprising a plurality of capture probes, wherein each capture probe comprise a cleavage domain, a cell-penetrating agent, a spatial barcode, and a capture domain; (c) cleaving at least one capture probe from the solid support via the cleavage domain such that the cleaved capture probe is introduced into the biological sample and at least one cleaved capture probe interacts with at least one biological analyte present in the biological sample via the capture domain; (d) dissociating the biological sample into a plurality of single cells, wherein the at least one cleaved capture probe that interacts with the at least one biological analyte is present in one of the dissociated cells; (e) detecting the at least one biological analyte present in the dissociated cell; and (f) correlating the at least one biological analyte with a spatial location in the biological sample.

[1520] Embodiment G3 is the method of embodiment G1 or G2, further comprising staining the biological sample.

[1521] Embodiment G4 is the method of embodiment G3, wherein the biological sample is stained prior to contacting the biological sample with the solid support.

[1522] Embodiment G5 is the method of any one of embodiments G1-G4, further comprising imaging the biological sample.

[1523] Embodiment G6 is the method of embodiment G5, wherein imaging is performed prior to cleaving the at least one capture probe from the solid support.

[1524] Embodiment G7 is the method of embodiment G5, wherein imaging is performed after cleaving of the at least one capture probe from the solid support.

[1525] Embodiment G8 is the method of any one of embodiments G5-G7, wherein imaging is used to determine a region of interest in the biological sample.

[1526] Embodiment G9 is the method of any one of embodiments G1-G8, wherein the step of cleaving at least one capture probe from the solid support comprises cleaving the plurality of capture probes from the solid support in bulk.

[1527] Embodiment G10 is the method of any one of embodiments G1-G8, wherein the step of cleaving at least one capture probe from the solid support comprises selectively cleaving at least one capture probe from the solid support.

[1528] Embodiment G11 is the method of any one of embodiments G1-G10, wherein the cleaving domain comprises a photocleavable domain.

[1529] Embodiment G12 is the method of embodiment G11, wherein cleaving at least one capture probe from the solid support comprises photocleaving.

[1530] Embodiment G13 is the method of embodiment G2, wherein the cell-penetrating agent comprises a cell penetrating peptide domain.

[1531] Embodiment G14 is the method of embodiment G13, wherein the cell-penetrating peptide domain mediates introduction of the capture probe into the cell.

[1532] Embodiment G15 is the method of any one of embodiments G1-G14, wherein the biological sample is permeabilized prior to contacting with the solid support.

[1533] Embodiment G16 is the method of any one of embodiments G1-G15, wherein the capture probe is introduced into the biological sample by diffusion.

[1534] Embodiment G17 is the method of any one of embodiments G1-G16, wherein the solid

support is removed from the biological sample after cleaving at least one capture probe from the solid support.

[1535] Embodiment G18 is the method of any one of embodiments G1-G17, wherein the capture probe further comprises a cell penetration tag.

[1536] Embodiment G19 is the method of embodiment G18, wherein the cell penetration tag comprises a fluorescent dye.

[1537] Embodiment G20 is the method of embodiment G19, wherein the cell penetration tag comprises at least one of Cy2, Cy3, Cy3B, Cy3.5, Cy5, Cy5.5, Cy7, fluorescein (6-FAM), DyLight, Alexa Fluor®, and tetramethylrhodamine (TAMRA) azide.

[1538] Embodiment G21 is the method of any one of embodiments G18-G20, further comprising detecting the cell penetration tag.

[1539] Embodiment G22 is the method of embodiment G21, wherein the cell penetration tag is detected after cleaving at least one capture probe from the solid support and removing the solid support from the biological sample.

[1540] Embodiment G23 is the method of embodiment G22, wherein introduction of the capture probe into the cell is determined by detecting the presence of the cell penetration tag in the cell.

[1541] Embodiment G24 is the method of any one of embodiments G1-G23, wherein the capture probe further comprises an intracellular cleavage domain.

[1542] Embodiment G25 is the method of embodiment G24, wherein the intracellular cleavage domain comprises a disulfide bond.

[1543] Embodiment G26 is the method of embodiment G24 or G25, wherein the intracellular cleavage domain is cleaved after the capture probe is introduced into the cell.

[1544] Embodiment G27 is the method of any one of embodiments G1-G26, wherein the dissociated single cells are analyzed using high-throughput instrumentation.

[1545] Embodiment G28 is the method of any one of embodiments G1-G27, wherein the at least one biological analyte is DNA.

[1546] Embodiment G29 is the method of any one of embodiments G1-G27, wherein the at least one biological analyte is RNA.

[1547] Embodiment G30 is the method of embodiment G29, wherein the capture domain comprises a poly-dT sequence.

[1548] Embodiment G31 is the method of embodiment G29 or G30, wherein the capture domain is configured to hybridize to a poly-A tail of an mRNA.

[1549] Embodiment G32 is the method of any one of embodiments G1-G31, wherein the capture probe further comprises a unique molecular identifier (UMI).

[1550] Embodiment G33 is the method of any one of embodiments G1-G32, wherein the capture probe further comprises a functional domain.

Exemplary Embodiments H

[1551] Embodiment H1 is a method for spatial profiling at least one biological analyte present in a biological sample, comprising: (a) providing a solid support comprising a plurality of capture probes, wherein the capture probes comprise a spatial barcode and a capture domain; (b) contacting one or more capture probes on the solid support with the biological sample under conditions where the at least one biological analyte present in the biological sample interacts with at least one capture probe; (c) identifying one or more regions of interest in the biological sample; (d) eliminating members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample; (e) detecting the at least one biological analyte; and (f) correlating the at least one biological analyte with a spatial location in the biological sample.

[1552] Embodiment H2 is the method of embodiment H1, wherein the conditions comprise releasing the at least one biological analyte from the biological sample.

[1553] Embodiment H3 is the method of embodiment H1 or H2, wherein the conditions comprise permeabilizing the biological sample.

[1554] Embodiment H4 is the method of embodiment H3, wherein the biological sample is fixed prior to permeabilization.

[1555] Embodiment H5 is the method of any one of embodiments H1-H4, wherein identifying one or more regions of interest in the biological sample is performed prior to contacting one or more capture probes on the solid support with the biological sample.

[1556] Embodiment H6 is the method of any one of embodiments H1-H5, wherein identifying one or more regions of interest in the biological sample is performed after contacting one or more capture probes on the solid support with the biological sample.

[1557] Embodiment H7 is the method of any one of embodiments H1-H6, further comprising imaging the biological sample.

[1558] Embodiment H8 is the method of any one of embodiments H1-H7, wherein imaging is used to determine the one or more regions of interest in the biological sample.

[1559] Embodiment H9 is the method of embodiment H1, wherein the step of eliminating comprises destroying the members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample.

[1560] Embodiment H10 is the method of embodiment H1, wherein the step of eliminating comprises removing the members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample.

[1561] Embodiment H11 is the method of embodiment H1, wherein the step of eliminating comprises blocking the members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample.

[1562] Embodiment H12 is the method of embodiment H1, wherein the step of eliminating comprises using light to eliminate members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample.

[1563] Embodiment H13 is the method of embodiment H12, wherein the light is directed to regions of the solid support that correspond to the one or more regions of interest in the biological sample by at least one of a mirror, mirror array, a lens, a moving stage, and a mask.

[1564] Embodiment H14 is the method of embodiment H12, wherein a scanning laser is used to eliminate members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample.

[1565] Embodiment H15 is the method of embodiment H14, wherein the scanning laser eliminates members of the plurality of capture probes using heat.

[1566] Embodiment H16 is the method of embodiment H14, wherein the scanning laser eliminates members of the plurality of capture probes using UV light.

[1567] Embodiment H17 is the method of embodiment H14, wherein the capture probes further comprises a photocleavable domain and wherein the scanning laser selectively cleaves members of the plurality of capture probes on the solid support that are outside the one or more regions of interest in the biological sample.

[1568] Embodiment H18 is the method of any one of embodiments H1-H17, wherein the step of correlating comprises correlating the at least one biological analyte to a spatial barcode of the capture probe.

[1569] Embodiment H19 is the method of any one of embodiments H1-H18, wherein the at least one biological analyte is DNA.

[1570] Embodiment H20 is the method of any one of embodiments H1-H18, wherein the at least one biological analyte is RNA.

[1571] Embodiment H21 is the method of embodiment H20, wherein the capture domain comprises a poly-dT sequence.

[1572] Embodiment H22 is the method of embodiment H20 or H21, wherein the capture domain is configured to hybridize to a poly-A tail of an mRNA.

[1573] Embodiment H23 is the method of any one of embodiments H1-H22, wherein the capture

probe further comprises a unique molecular identifier (UMI).

[1574] Embodiment H24 is the method of any one of embodiments H1-H24, wherein the capture probe further comprises a functional domain.

Exemplary Embodiments I

[1575] Embodiment I1 is a method of manufacturing a shrunken hydrogel bead-containing hydrogel matrix comprising: providing a hydrogel bead; embedding the hydrogel bead in a hydrogel matrix, thus generating the hydrogel bead-containing hydrogel matrix; reducing the size of the hydrogel bead-containing hydrogel matrix by removing water, thus generating the shrunken hydrogel bead-containing hydrogel matrix.

[1576] Embodiment 12 is the method of embodiment I1, wherein the water is removed from the hydrogel bead-containing hydrogel matrix by subjecting the hydrogel bead-containing hydrogel matrix to a dehydrating solvent.

[1577] Embodiment I3 is the method of embodiment 12, wherein the solvent comprises at least one of alcohol, ketone, and aldehyde.

[1578] Embodiment 14 is the method of embodiment I1, wherein the water is removed from the hydrogel bead-containing hydrogel matrix by subjecting the hydrogel bead-containing hydrogel matrix to a salt.

[1579] Embodiment 15 is the method of embodiment 14, wherein the salt comprises an inorganic salt.

[1580] Embodiment 16 is the method of embodiment I1, wherein the water is removed from the hydrogel bead-containing hydrogel matrix by subjecting the hydrogel bead-containing hydrogel matrix to a dehydrating process.

[1581] Embodiment I7 is the method of embodiment I1, wherein the shrunken hydrogel bead-containing hydrogel matrix is at least 2-fold smaller in a linear dimension than the pre-shrunken hydrogel bead-containing hydrogel matrix.

[1582] Embodiment 18 is the method of embodiment I1, wherein the shrunken hydrogel bead-containing hydrogel matrix is at least 3-fold smaller in a linear dimension than the pre-shrunken hydrogel bead-containing hydrogel matrix.

[1583] Embodiment 19 is the method of embodiment I1, wherein the shrunken hydrogel bead-containing hydrogel matrix is at least 4-fold smaller in a linear dimension than the pre-shrunken hydrogel bead-containing hydrogel matrix.

[1584] Embodiment I10 is the method of any one of the embodiments I1-I9, wherein the shrunken hydrogel bead-containing hydrogel matrix comprises a capture probe.

[1585] Embodiment I11 is the method of embodiment I10, wherein the capture probe comprises a spatial barcode.

[1586] Embodiment I12 is the method of embodiment I10, wherein the capture probe comprises a unique molecular identifier.

[1587] Embodiment I13 is the method of embodiment I10, wherein the capture probe comprises a functional domain.

[1588] Embodiment I14 is the method of any one of the embodiments I1-I9, wherein the shrunken hydrogel bead-containing hydrogel matrix comprises a capture probe comprising a cleavage domain, a spatial barcode, a unique molecular identifier, a functional domain and a capture domain.

[1589] Embodiment I15 is a method of generating a hydrogel bead array-containing hydrogel matrix comprising: providing a plurality of hydrogel beads; embedding the plurality of hydrogel beads in a hydrogel matrix, thus generating of the hydrogel bead array-containing hydrogel matrix; and reducing the size of the hydrogel bead array-containing hydrogel matrix by removing water, thus generating the plurality of shrunken hydrogel bead array-containing hydrogel matrix.

[1590] Embodiment I16 is the method of embodiment I15, wherein the water is removed from the hydrogel bead array-containing hydrogel matrix by subjecting the hydrogel bead array-containing hydrogel matrix to a dehydrating solvent.

[1591] Embodiment I17 is the method of embodiment I16, wherein the solvent comprises at least one of alcohol, ketone, and aldehyde.

[1592] Embodiment I18 is the method of embodiment I15, wherein the water is removed from the hydrogel bead array-containing hydrogel matrix by subjecting the hydrogel bead array-containing hydrogel matrix to a salt.

[1593] Embodiment I19 is the method of embodiment I18, wherein the salt comprises an inorganic salt.

[1594] Embodiment I20 is the method of embodiment I15, wherein the water is removed from the hydrogel bead array-containing hydrogel matrix by subjecting the hydrogel bead array-containing hydrogel matrix to a dehydrating process.

[1595] Embodiment I21 is the method of embodiment I15, wherein the shrunken hydrogel bead array-containing hydrogel matrix is at least 2-fold smaller in a linear dimension than the pre-shrunk hydrogel bead array-containing hydrogel matrix.

[1596] Embodiment I22 is the method of embodiment I15, wherein the shrunken hydrogel bead array-containing hydrogel matrix is at least 3-fold smaller in a linear dimension than the pre-shrunk hydrogel bead array-containing hydrogel matrix.

[1597] Embodiment I23 is the method of embodiment I15, wherein the shrunken hydrogel bead array-containing hydrogel matrix is at least 4-fold smaller in a linear dimension than the pre-shrunk hydrogel bead array-containing hydrogel matrix.

[1598] Embodiment I24 is the method of any one of the embodiments I15-I23, wherein each of the hydrogel beads of the plurality of shrunken hydrogel bead array-containing hydrogel matrix comprise a plurality of capture probes.

[1599] Embodiment I25 is the method of embodiment I24, wherein the plurality of capture probes comprise a plurality of spatial barcodes.

[1600] Embodiment I26 is the method of embodiment I24, wherein the plurality of capture probes comprise a plurality of unique molecular identifiers.

[1601] Embodiment I27 is the method of embodiment I24, wherein the plurality of capture probes comprise a plurality of functional domains.

[1602] Embodiment I28 is the method of any one of the embodiments I15-I23, wherein the plurality of hydrogel beads on the array-containing hydrogel matrix comprise a plurality of capture probes comprising a plurality of cleavage domains, spatial barcodes, unique molecular identifiers, functional domains, and capture domains.

[1603] Embodiment I29 is a bead array, comprising: a plurality of hydrogel beads dispersed on a solid substrate, wherein the plurality of hydrogel beads are spatially confined on the solid substrate, and wherein the spatially confined plurality of hydrogel beads are embedded in a hydrogel matrix.

[1604] Embodiment I30 is the bead array of embodiment I29, wherein the plurality of hydrogel beads comprises a plurality of capture probes.

[1605] Embodiment I31 is the bead array of embodiment I30, wherein the plurality of capture probes comprise a plurality of spatial barcodes.

[1606] Embodiment I32 is the bead array of embodiment I30, wherein the plurality of capture probes comprise a plurality of unique molecular identifiers.

[1607] Embodiment I33 is the bead array of embodiment I30, wherein the plurality of capture probes comprise a plurality of functional domains.

[1608] Embodiment I34 is the bead array of embodiment I29, wherein members of the plurality of hydrogel beads comprise a plurality of capture probes comprising a plurality of cleavage domains, spatial barcodes, unique molecular identifiers, functional domains, and capture domains.

[1609] Embodiment I35 is a method for spatially profiling a plurality of analytes in a biological sample, comprising: (a) providing a shrunken hydrogel bead array-containing hydrogel matrix; wherein the shrunken hydrogel bead array-containing hydrogel matrix comprises an arrayed plurality of shrunken hydrogel beads; wherein at least one member of the plurality of shrunken

hydrogel beads comprises a capture probe; (b) contacting the biological sample with the arrayed plurality of shrunken hydrogel beads under conditions such that one or more analytes present in the biological sample are captured by one or more of the capture probes; and (c) determining the spatial profile of the captured analytes.

[1610] Embodiment I36 is the method of embodiment I35, wherein the biological sample is a tissue sample.

[1611] Embodiment I37 is the method of embodiment I36, wherein the tissue sample is a fresh-frozen tissue sample.

[1612] Embodiment I38 is the method of embodiment I36, wherein the tissue sample comprises a tumor cell.

[1613] Embodiment I39 is the method of embodiment I36, wherein the tissue sample comprises a tissue section.

[1614] Embodiment I40 is the method of embodiment I35, wherein the analytes in the biological sample are migrated to the bead array.

[1615] Embodiment I41 is the method of embodiment I40, wherein the migrating comprises passive migration.

[1616] Embodiment I42 is the method of embodiment I40, wherein the migrating comprises active migration.

[1617] Embodiment I43 is the method of embodiment I35, wherein the one or more analytes comprises at least one of RNA, DNA, protein, lipid, peptide, metabolite, small molecule, and a cell labeling agent.

[1618] Embodiment I44 is a method of manufacturing a shrunken hydrogel bead-containing hydrogel matrix comprising: providing a hydrogel bead-containing hydrogel matrix comprising a hydrogel bead embedded in a hydrogel matrix; reducing the size of the hydrogel bead-containing hydrogel matrix by removing water, thus generating the shrunken hydrogel bead-containing hydrogel matrix.

[1619] Embodiment I45 is a method of manufacturing a shrunken hydrogel bead array-containing hydrogel matrix comprising: providing a hydrogel bead array-containing hydrogel matrix comprising an array of hydrogel beads embedded in a hydrogel matrix; reducing the size of the hydrogel bead array-containing hydrogel matrix by removing water, thus generating the shrunken hydrogel bead array-containing hydrogel matrix.

Exemplary Embodiments J

[1620] Embodiment J1 is a method for spatially detecting a biological analyte of interest from a biological sample, the method comprising: (a) providing a biological sample on a support; (b) staining the biological sample on the support, imaging the stained biological sample, and selecting a biological sample or subsection of biological sample to subject to spatial transcriptomic analysis; (c) providing an array comprising one or more pluralities of capture probes on a substrate; (d) contacting the biological sample with the array, thereby allowing the one or more pluralities of capture probes to capture the biological analyte of interest; and (e) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest.

[1621] Embodiment J2 is the method of embodiment J1, wherein the capture probe comprises a capture domain.

[1622] Embodiment J3 is the method of embodiment J1, wherein the capture probe further comprises a unique molecular identifier (UMI).

[1623] Embodiment J4 is the method of embodiment J1, wherein the biological analyte comprises DNA or RNA.

[1624] Embodiment J5 is the method of embodiment J1, wherein the biological analyte comprises a protein.

[1625] Embodiment J6 is the method of embodiment J1, wherein the array substrate is liquid permeable.

[1626] Embodiment J7 is the method of embodiment J1, wherein the array substrate comprises hydrogel.

[1627] Embodiment J8 is the method of embodiment J7, wherein the hydrogel comprises a permeabilization reagent.

[1628] Embodiment J9 is the method of embodiment J1, wherein the array substrate comprises one or more microfluidic channels.

[1629] Embodiment J10 is the method of embodiment J1, wherein the array substrate comprises one or more distribution channels.

[1630] Embodiment J11 is the method of embodiment J1, wherein the biological sample is exposed to a permeabilization reagent prior to contacting the biological sample with the array.

[1631] Embodiment J12 is the method of embodiment J1, wherein the array substrate comprises a permeabilization reagent.

[1632] Embodiment J13 is the method of embodiment J1, further comprising modulating the rate of interaction between biological analytes from the biological sample and the capture probes on the array.

[1633] Embodiment J14 is the method of embodiment J1, further comprising modulating the rate of chemical reactions.

[1634] Embodiment J15 is the method of embodiment J14, wherein the chemical reactions are modulated by temperature, light, a small molecule, or an enzyme.

Exemplary Embodiments K

[1635] Embodiment K1 is a method for spatially detecting a biological analyte within a biological sample, the method comprising: (a) providing a support, wherein a section of the support is homogeneously covered by a plurality of capture probes, each capture probe comprising a capture domain; (b) contacting the biological sample to the support, such that the plurality of capture probes interacts with a plurality of analytes from the biological sample to generate a plurality of tagged analyte molecules; (c) identifying an area of interest within the support; and (d) detecting the presence of one or more biological analytes within the area of interest, thereby spatially detecting the biological analyte within the biological sample.

[1636] Embodiment K2 is a method for spatially detecting a biological analyte within a biological sample, the method comprising: (a) providing a cell labelling agent to the biological sample, wherein the cell labelling agent comprises at least a molecular tag barcode; (b) allowing the cell labelling agent to bind to the biological analyte; (c) providing a support, wherein a section of the support is homogeneously covered by a plurality of capture probes, each capture probe comprising a capture domain; (d) contacting the biological sample to the support, such that the plurality of capture probes interacts with a plurality of molecular tag barcodes, creating a plurality of tagged analyte molecules; (e) identifying an area of interest within the support; and (f) detecting the presence of one or more biological analytes within the area of interest, thereby spatially detecting the biological analyte within the biological sample.

[1637] Embodiment K3 is the method of embodiment K2, wherein a molecular tag barcode further comprises a cleavage domain.

[1638] Embodiment K4 is the method of embodiment K3, wherein the cleavage domain is cleavable by at least one of light, chemical molecule, and enzyme.

[1639] Embodiment K5 is the method of embodiments K1 or K2, wherein the biological analyte comprises RNA.

[1640] Embodiment K6 is the method of embodiments K1 or K2, wherein the biological analyte comprises DNA.

[1641] Embodiment K7 is the method of embodiment K1 or K2, wherein the biological analyte comprises a protein.

[1642] Embodiment K8 is the method of embodiment K2, wherein the biological analyte comprises an antibody or an antigen-binding fragment thereof, and an oligonucleotide.

[1643] Embodiment K9 is the method of embodiments K1 or K2, wherein an electric field is applied through the biological sample and the array.

[1644] Embodiment K10 is the method of embodiments K1 or K2, or K9, wherein the biological sample is removed from the support prior to analyzing the biological analyte at the area of interest.

[1645] Embodiment K11 is the method of embodiments K1 or K2, wherein the biological sample is treated with a permeabilization reagent prior to or after contacting the support with the biological sample.

[1646] Embodiment K12 is the method of embodiments K1 or K2, wherein the plurality of probes comprise identical capture domains.

[1647] Embodiment K13 is the method of embodiments K1 or K2, wherein the plurality of probes comprise two or more non-identical capture domains.

[1648] Embodiment K14 is the method of embodiments K1 or K2, wherein the plurality of probes are directly attached to the support.

[1649] Embodiment K15 is the method of embodiments K1 or K2, wherein the plurality of probes are indirectly attached to the support.

[1650] Embodiment K16 is the method of embodiment K15, wherein the plurality of probes are attached to beads and the beads are attached to the support.

[1651] Embodiment K17 is the method of embodiment K16, wherein the beads are gel beads.

[1652] Embodiment K18 is the method of embodiments K1 or K2, wherein detecting the presence of one or more tagged analyte molecules comprises performing in-situ sequencing.

[1653] Embodiment K19 is the method of embodiments K1, K2, or K18, wherein detecting the presence of one or more tagged analyte molecules comprises one or more cycles of hybridization with labelled probes.

[1654] Embodiment K20 is the method of embodiment K19, wherein the labelled probes are labelled with at least one of a fluorescent molecule, a fluorescent protein, a radioactive molecule, or an enzyme.

[1655] Embodiment K21 is the method of embodiment K20, wherein the nucleic acid comprising the molecular tag barcode is ligated to the capture domain prior to detecting the presence of one or more tagged analyte molecules.

[1656] Embodiment K22 is the method of embodiment K19, wherein the nucleic acid comprising the molecular tag barcode is amplified prior to detecting the presence of one or more tagged analyte molecules.

[1657] Embodiment K23 is the method of embodiment K2, wherein the cell labelling agent further comprises a molecular tag.

[1658] Embodiment K24 is the method of embodiment K12, wherein the capture domain comprises an oligo (dT) sequence.

[1659] Embodiment K25 is the method of embodiments K1 or K2, wherein the support is placed into a flow cell and a set of fluorescent probes are hybridized to the molecular tag barcodes.

[1660] Embodiment K26 is the method of embodiment K25, wherein the fluorescent probes are imaged, and the support is heated for dehybridization and the fluorescent probes are washed out of the flow cell.

Exemplary Embodiments L

[1661] Embodiment L1 is a method for transferring a biological analyte from a biological sample to an array, the method comprising: (a) providing an array comprising a capture probe; (b) contacting the array with a biological sample; and (c) applying an electric field to the biological sample and the array.

[1662] Embodiment L2 is the method of embodiment L1, wherein the array comprises a conductive substrate.

[1663] Embodiment L3 is the method of embodiments L1 or L2, wherein the array comprises spots on paper.

- [1664] Embodiment L4 is the method of embodiments L1 or L2, wherein the array comprises photocured gel pads on paper.
- [1665] Embodiment L5 is the method of embodiments L1 or L2, wherein the array comprises a gel bead array on paper.
- [1666] Embodiment L6 is the method of embodiments L1 or L2, wherein the array comprises photocured gel pads on a gel film.
- [1667] Embodiment L7 is the method of embodiments L1 or L2, wherein the array comprises a gel bead array on a gel film.
- [1668] Embodiment L8 is the method of embodiments L1 or L2, wherein the array comprises spots on conductive glass.
- [1669] Embodiment L9 is the method of embodiment L1, wherein the capture probe comprises a capture domain.
- [1670] Embodiment L10 is the method of embodiment L2, wherein the capture probe is directly attached to the conductive substrate.
- [1671] Embodiment L11 is the method of embodiment L2, wherein the capture probe is indirectly attached to the conductive substrate.
- [1672] Embodiment L12 is the method of embodiment L1, wherein the capture probe is directly attached to a bead that is fixed on the array.
- [1673] Embodiment L13 is the method of embodiment L9, wherein the capture probe is indirectly attached to the conductive substrate via a solution comprising a polymer that is activated to polymerize by photocuring.
- [1674] Embodiment L14 is the method of embodiment L1, wherein the capture probe further comprises a unique molecular identifier (UMI).
- [1675] Embodiment L15 is the method of embodiment L1, wherein the capture probe further comprises a sequence for initiating a sequencing reaction.
- [1676] Embodiment L16 is the method of embodiment L1, wherein the biological analyte comprises DNA or RNA.
- [1677] Embodiment L17 is the method of embodiment L1, wherein the biological analyte comprises a protein.
- [1678] Embodiment L18 is the method of embodiments L16 or L17, wherein the electric field is applied after the tissue sample has been exposed to a solution comprising a permeabilization reagent.
- [1679] Embodiment L19 is the method of embodiment L18, wherein the electric field is applied after the tissue sample has been further exposed to a solution comprising SDS.
- [1680] Embodiment L20 is the method of embodiment L1, wherein the electric field is applied by an anode and a cathode.

Exemplary Embodiments M

- [1681] Embodiment M1 is a method for increasing the capture efficiency of an array, the method comprising: (a) providing an array comprising one or more capture probe pluralities; (b) placing a biological sample between the array and a diffusion-resistant medium, submerged in a permeabilization buffer, wherein the diffusion-resistant medium reduces diffusion of a plurality of biological analytes from the biological sample into the permeabilization buffer, thereby increasing the capture efficiency of the array.
- [1682] Embodiment M2 is a method for increasing the capture efficiency of an array, the method comprising: (a) providing an array comprising one or more capture probe pluralities; (b) placing a biological sample between the array and a diffusion-resistant medium, wherein the diffusion resistant medium comprises a support coated with dried reagents, wherein the diffusion-resistant medium focuses the diffusion of a plurality of biological analytes from the biological sample toward the array, thereby increasing the capture efficiency of the array.
- [1683] Embodiment M3 is a method for increasing the capture efficiency of an array, the method

comprising: (a) providing an array comprising one or more capture probe pluralities; (b) placing a biological sample between the array and a diffusion-resistant medium, wherein the diffusion resistant medium comprises a support containing small wells or pores containing permeabilization buffer, wherein the diffusion-resistant medium focuses the diffusion of a plurality of biological analytes from the biological sample toward the array, thereby increasing the capture efficiency of the array.

[1684] Embodiment M4 is the method of embodiments M1, M2, or M3, wherein the one or more capture probes comprise a capture domain.

[1685] Embodiment M5 is the method of embodiments M1, M2 or M3, wherein the biological sample is placed on the top of the array, and the diffusion-resistant medium is placed on the top of the biological sample.

[1686] Embodiment M6 is the method of embodiment M1, wherein the diffusion-resistant medium is applied after the biological sample has been exposed to a solution comprising a permeabilization reagent.

[1687] Embodiment M7 is the method of embodiments M2 or M3, wherein the diffusion-resistant medium comprises at least one permeabilization reagent.

[1688] Embodiment M8 is the method of embodiments M1, M2, or M3, wherein the diffusion-resistant medium is glass.

[1689] Embodiment M9 is the method of embodiments M1, M2, or M3, wherein the diffusion-resistant medium is a hydrogel.

[1690] Embodiment M10 is the method of embodiments M2 or M3, wherein the hydrogel is pre-soaked in a solution comprising a permeabilization reagent.

Exemplary Embodiments N

[1691] Embodiment N1 is a method of spatially profiling a plurality of biological analytes in a biological sample, comprising: (a) providing a microcapillary array comprising a plurality of microcapillary channels; (b) introducing to the microcapillary array a plurality of features, such that each member of the plurality of microcapillary channels receives no more than one feature on average, wherein each feature comprises a plurality of capture probes, wherein each capture probe comprises a spatial barcode and a capture domain; (c) reducing the cross-sectional areas of the features in the microcapillary channels; (d) contacting the microcapillary array to a biological sample under conditions that permit the capture domains of the capture probes to capture analytes in the biological sample; and (e) determining the spatial profile of the captured biological analytes in the biological sample.

[1692] Embodiment N2 is the method of embodiment N1, wherein the capture probe further comprises a unique molecular identifier.

[1693] Embodiment N3 is the method of embodiment N1, wherein the capture probe further comprises a cleavage domain.

[1694] Embodiment N4 is the method of embodiment N1, wherein the capture probe further comprises a functional domain.

[1695] Embodiment N5 is the method of embodiment N1, wherein each feature in the microcapillary channels comprises a plurality of capture probes, each capture probe comprising a cleavage domain, a spatial barcode, a unique molecular identifier, a functional domain, and a capture domain.

[1696] Embodiment N6 is the method of embodiment N1, wherein the conditions that permit the capture domains of the capture probes to capture analytes in the biological sample include applying pressure.

[1697] Embodiment N7 is the method of embodiment N6, wherein the pressure is mechanical pressure.

[1698] Embodiment N8 is the method of embodiment N1, wherein the conditions that permit the capture domains of the capture probes to capture analytes in the biological sample include flowing

the features through the microcapillary channels.

[1699] Embodiment N9 is the method of embodiment N1, wherein the feature comprises a bead.

[1700] Embodiment N10 is the method of embodiment N1, wherein the feature comprises a hydrogel bead.

[1701] Embodiment N11 is the method of embodiment N1, wherein the cross-sectional area of the feature is reduced by 2-fold.

[1702] Embodiment N12 is the method of embodiment N1, wherein the biological sample is immobilized on a solid substrate.

[1703] Embodiment N13 is the method of embodiment N1, wherein the biological sample is reversibly immobilized on a solid substrate.

[1704] Embodiment N14 is the method of embodiment N13, wherein the biological sample is reversibly immobilized using a dissolvable polymer.

[1705] Embodiment N15 is the method of embodiment N1, wherein determining the spatial profile of the captured biological analytes in the biological sample further comprises: (a) contacting the microcapillary channels with a solution comprising reagents; (b) sealing some or all of the microcapillary channels; (c) incubating the microcapillary array with the solution; and (d) determining the spatial profile of the captured biological analytes in the biological sample.

[1706] Embodiment N16 is the method of embodiment N15, wherein the reagents comprise enzymatic reagents.

[1707] Embodiment N17 is the method of embodiment N15, wherein the reagents comprise at least one of a reverse transcriptase, a ligase, and one or more oligonucleotides.

[1708] Embodiment N18 is the method of embodiment N15, wherein the sealing is performed using silicone oil.

[1709] Embodiment N19 is the method of embodiment N15, wherein the incubating comprises incubating for at least 1 minute.

[1710] Embodiment N20 is the method of embodiment N15, wherein the incubating comprises incubating at an appropriate temperature.

[1711] Embodiment N21 is the method of embodiment N1, wherein the biological analytes in the biological sample are migrated to the features in the microcapillary array.

[1712] Embodiment N22 is the method of embodiment N21, wherein the migrating comprises passive migration.

[1713] Embodiment N23 is the method of embodiment N21, wherein the migrating comprises active migration.

[1714] Embodiment N24 is the method of embodiment N1, wherein the biological sample is a tissue sample.

[1715] Embodiment N25 is the method of embodiment N24, wherein the tissue sample is a fresh-frozen tissue sample.

[1716] Embodiment N26 is the method of embodiment N24, wherein the tissue sample comprises a tumor cell.

[1717] Embodiment N27 is the method of embodiment N24, wherein the tissue sample comprises a tissue section.

[1718] Embodiment N28 is the method of embodiment N1, wherein the captured biological analytes comprise at least one of RNA, DNA, protein, lipid, peptide, metabolite, small molecule, and a cell labeling agent.

[1719] Embodiment N29 is the method of embodiment N1, further comprising imaging the biological sample.

[1720] Embodiment N30 is the method of embodiment N29, wherein the imaging is performed prior to contacting the biological sample with the features in the microcapillary array.

[1721] Embodiment N31 is the method of embodiment N29, wherein the imaging is performed after contacting the biological sample with the features in the microcapillary array.

[1722] Embodiment N32 is the method of embodiment N29, wherein the imaging is used to determine a region of interest in the biological sample.

[1723] Embodiment N33 is the method of embodiment N29, wherein the imaging comprises using fiducial markers.

Exemplary Embodiments O

[1724] Embodiment O1 is a method for preparing a feature array, the method comprising: (a) providing a solid substrate; (b) applying a coating comprising a plurality of photo-reactive molecules to at least one surface of the solid substrate; (c) exposing one or more regions of the coating on the solid substrate to at least one light source such that one or more feature locations are generated; and (d) providing a plurality of features, wherein a member of the plurality of features attaches to a feature location.

[1725] Embodiment O2 is the method of embodiment O1, further comprising repeating steps: (c) exposing one or more regions of the coating on the solid substrate to at least one light source such that one or more feature locations are generated; and (d) providing a plurality of features, wherein a member of the plurality of features attaches to a feature location.

[1726] Embodiment O3 is the method of embodiment O1 or O2, wherein exposing one or more regions of the coating on the solid substrate to at least one light source comprises a photomask.

[1727] Embodiment O4 is the method of embodiment O3, wherein each step of exposing one or more regions of the coating on the solid substrate to at least one light source comprises a different photomask than any previous step of exposing one or more regions of the coating on the solid substrate to at least one light source.

[1728] Embodiment O5 is the method of any one of embodiments O2-O4, wherein each step of exposing one or more regions of the coating on the solid substrate to at least one light source comprises generating one or more feature locations at different locations than any previous step of exposing one or more regions of the coating on the solid substrate to at least one light source.

[1729] Embodiment O6 is the method of any one of embodiments O1-O5, wherein the one or more exposed regions of the coating on the solid substrate correspond to the one or more feature locations.

[1730] Embodiment O7 is the method of any one of embodiments O2-O6, wherein steps (c) and (d) are repeated at least a total of two times.

[1731] Embodiment O8 is the method of any one of embodiments O1-O7, wherein members of the plurality of photoreactive molecules are degraded at the one or more feature locations when exposed to the one or more light sources.

[1732] Embodiment O9 is the method of embodiment O8, wherein the solid substrate is treated or was previously treated such that the one or more feature locations bind to at least one of the plurality of features.

[1733] Embodiment O10 is the method of embodiment O9, wherein the solid substrate further comprises an adhesive coating between the solid substrate and the coating comprising a plurality of photo-reactive molecules.

[1734] Embodiment O11 is the method of any one of embodiments O1-O10, wherein the one or more feature locations comprise exposed adhesive that binds one or more features, wherein the adhesive is exposed as a result of photo-reactive molecules being deactivated or degraded by the light source.

[1735] Embodiment O12 is the method of any one of embodiments O1-O11, wherein the plurality of photo-reactive molecules comprises a photo-degradable polymer.

[1736] Embodiment O13 is the method of embodiment O12, wherein the photo-degradable polymer comprises a photo-degradable PEG polymer, a photo degradable methacrylated gelatin (GelMA) polymer, or a combination thereof.

[1737] Embodiment O14 is the method of embodiment O13, wherein the photo-degradable PEG polymer comprises poly(L-lysine)-[g]-poly(ethylene glycol) (PLL-PEG).

[1738] Embodiment O15 is the method of any one of embodiments O1-O7, wherein the one or more feature locations comprise activated photo-reactive molecules that bind one or more features, wherein the photoreactive molecules are activated by the light source.

[1739] Embodiment O16 is the method of any one of embodiments O1-O15, wherein step (c) is performed prior to step (d).

[1740] Embodiment O17 is the method of any one of embodiments O1-O7, wherein step (d) is performed prior to step (c).

[1741] Embodiment O18 is the method of embodiment O17, wherein members of the plurality of photo-reactive molecules are activated by the light source and cross link to one or more members of the plurality of features.

[1742] Embodiment O19 is the method of embodiment O17 or O18, wherein the plurality of features are provided in excess of the feature locations.

[1743] Embodiment O20 is the method of embodiment O19, wherein the plurality of features are provided such that there are at least twice as many features as feature locations.

[1744] Embodiment O21 is the method of embodiment O20, wherein unbound features are washed away after each step of exposing one or more regions of the coating on the solid substrate to at least one light source.

[1745] Embodiment O22 is a method for preparing a feature array, the method comprising: (a) providing a solid substrate; (b) applying a coating comprising a plurality of photo-reactive molecules to at least one surface of the solid substrate; (c) providing a plurality of features to the coated solid substrate; and (d) exposing one or more regions of the coating on the solid substrate to at least one light source such that one or more members of the plurality of photo-reactive molecules crosslink to one or more members of the plurality of features at one or more feature locations on the solid substrate.

[1746] Embodiment O23 is the method of embodiment O22, wherein the plurality of features are provided in excess of the feature locations.

[1747] Embodiment O24 is the method of embodiment O23, wherein the plurality of features are provided such that there are at least twice as many features as feature locations.

[1748] Embodiment O25 is the method of any one of embodiments O22-O24, wherein unbound features are washed away.

[1749] Embodiment O26 is the method of any one of embodiments O22-O25, further comprising repeating steps: (c) providing a plurality of features to the coated solid substrate; and (d) exposing one or more regions of the coating on the solid substrate to at least one light source such that one or more members of the plurality of photo-reactive molecules crosslink to one or more members of the plurality of features at one or more feature locations on the solid substrate.

[1750] Embodiment O27 is the method of embodiment O26, wherein unbound features are washed away after each step of exposing one or more regions of the coating on the solid substrate to at least one light source.

[1751] Embodiment O28 is the method of any one of embodiments O22-O27, wherein exposing one or more regions of the coating on the solid substrate to at least one light source comprises a photomask.

[1752] Embodiment O29 is the method of embodiment O28, wherein each step of exposing one or more regions of the coating on the solid substrate to at least one light source comprises a different photomask than any previous step of exposing one or more regions of the coating on the solid substrate to at least one light source.

[1753] Embodiment O30 is the method of any one of embodiments O2-O29, wherein the plurality of features added in a particular cycle have one or more different chemical properties and/or physical properties compared to the plurality of features previously bound to the solid substrate.

[1754] Embodiment O31 is the method of any one of embodiments O1-O30, wherein the plurality of features comprise beads.

[1755] Embodiment O32 is the method of embodiment O31, wherein the beads comprise one or more capture probes.

[1756] Embodiment O33 is the method of embodiment O32, wherein each capture probe comprises a capture domain and a spatial barcode, wherein the spatial barcode is unique to a particular feature.

[1757] Embodiment O34 is the method of embodiment O31, wherein each capture probe added in the same step comprise the same barcode, wherein the barcode is unique to the plurality of features added in the same step.

[1758] Embodiment O35 is the method of embodiment O34, wherein each capture probe further comprises a unique molecular identifier.

Exemplary Embodiments P

[1759] Embodiment P1 is a method for preparing a feature array, the method comprising (a) delivering a plurality of features to one or more regions on a solid substrate, wherein each feature comprises one or more capture probes and a prepolymer solution; (b) inducing polymerization of the prepolymer solution on the solid substrate, thereby producing a plurality of features immobilized on the array.

[1760] Embodiment P2 is the method of embodiment P1, wherein the features are droplets.

[1761] Embodiment P3 is the method of embodiment P1, wherein the features are beads.

[1762] Embodiment P4 is the method of embodiment P1, wherein the delivery of the features to the solid substrate is accomplished by a system comprising droplet manipulation.

[1763] Embodiment P5 is the method of embodiment P4, wherein the system further comprises use of an electric field.

[1764] Embodiment P6 is the method of embodiment P1, wherein the polymerization step comprises exposure to one or more light sources.

[1765] Embodiment P7 is the method of embodiment P1, wherein the polymerization step comprises exposing the plurality of features to a solution comprising a chemical compound.

[1766] Embodiment P8 is the method of embodiment P1, wherein the one or more capture probes comprise a capture domain and a spatial barcode.

[1767] Embodiment P9 is the method of embodiment P8, wherein the one or more capture probes further comprise a unique molecular identifier (UMI), a functional domain, a cleavage domain, or a combination thereof.

[1768] Embodiment P10 is the method of embodiment P1, wherein the solid substrate comprises a plurality of wells.

[1769] Embodiment P11 is the method of embodiment P1, wherein the solid substrate is a glass slide.

[1770] Embodiment P12 is a method of spatially detecting a biological analyte within a biological sample, the method comprising: (a) providing a feature array made by the method of embodiment P1; (b) contacting the feature array with the biological sample, thereby allowing the capture probes to capture the biological analyte from the biological sample; (c) detecting the biological analytes captured by the feature array, thereby spatially detecting the biological analyte within the biological sample.

[1771] Embodiment P13 is the method of embodiment P12, further comprising collecting the features from the array.

[1772] Embodiment P14 is the method of embodiment P12, further comprising partitioning a feature from the plurality of features into a vesicle.

[1773] Embodiment P15 is the method of embodiment P12, wherein the vesicle further comprises a barcoded bead.

[1774] Embodiment P16 is the method of embodiment P12, further comprising ligating the biological analyte to the capture probes of the feature array prior to step (c).

[1775] Embodiment P17 is the method of embodiments P12, wherein the detecting is determined

by situ-sequencing.

[1776] Embodiment P18 is the method of embodiment P12 or P17, wherein the detecting is determined by one or more cycles of hybridization with probes comprising a detectable label.

[1777] Embodiment P19 is the method of embodiment P18, wherein the detectable label comprises at least one of a fluorescent molecule, a fluorescent protein, a radioactive molecule, and an enzyme.

[1778] Embodiment P20 is the method of embodiments P12, wherein the detecting is determined by high-throughput sequencing.

[1779] Embodiment P21 is the method of embodiment P20, wherein the method further comprises amplifying one or more nucleotide sequences.

[1780] Embodiment P22 is the method of embodiment P12, wherein the biological analyte comprises DNA or RNA.

[1781] Embodiment P23 is the method of embodiment P12, wherein the biological analyte is a non-poly(A) RNA target.

[1782] Embodiment P24 is the method of embodiment P12, wherein the biological analyte is microRNA.

[1783] Embodiment P25 is the method of embodiment P12, wherein the biological analyte comprises a protein.

[1784] Embodiment P26 is the method of embodiment P12, wherein the biological sample is exposed to a cell labeling agent prior to step (b).

Exemplary Embodiments Q

[1785] Embodiment Q1 is a method for the spatial identification of polypeptides in a tissue sample, said method comprising: (a) contacting a tissue sample with a solid support comprising a plurality of pre-defined spatial regions that correspond to positions in the tissue sample, wherein the solid support comprises a plurality of capture molecules; (b) capturing a polypeptide derived from the tissue sample via a capture molecule of the plurality of capture molecules, wherein the capture molecule is attached to the solid support in a pre-defined spatial region of the plurality of pre-defined spatial regions; (c) removing the tissue sample; (d) detecting the captured polypeptide via the capture molecule attached to the pre-defined spatial region.

[1786] Embodiment Q2 is the method of embodiment Q1, wherein the tissue sample is a cell, tissue section or cultured cells.

[1787] Embodiment Q3 is the method of embodiment Q1, wherein the polypeptides are captured by active or passive capture methods.

[1788] Embodiment Q4 is the method of embodiment Q3, wherein the active capture methods are electrophoretic directional migration, or magnet assisted directional migration.

[1789] Embodiment Q5 is the method of embodiment Q3, wherein the passive capture is diffusion.

[1790] Embodiment Q6 is the method of embodiment Q1, wherein the tissue sample is removed incubating with a Proteinase, protease (e.g., proteinase K, pepsin, etc.), collagenase, or combination thereof.

[1791] Embodiment Q7 is the method of embodiment Q6, further comprising DNase, and/or RNase.

[1792] Embodiment Q8 is the method of embodiment Q1, wherein the captured polypeptides are sequenced.

[1793] Embodiment Q9 is the method of embodiment Q1, wherein the sequencing is optical fluorosequencing.

[1794] Embodiment Q10 is the method of embodiment Q1, wherein the sequencing is radioactivity-based sequencing.

[1795] Embodiment Q11 is the method of embodiment Q1, further comprising the capture and sequencing of nucleic acids.

[1796] Embodiment Q12 is the method of embodiment Q1, wherein the plurality of pre-defined spatial regions corresponds to positions of polypeptides in the tissue sample.

[1797] Embodiment Q13 is a method for the spatial identification of polypeptides in a tissue sample, said method comprising: (a) contacting a tissue sample with a solid support, wherein the solid support comprises a plurality of capture molecules and a plurality of spatial barcodes; (b) capturing a polypeptide derived from the tissue sample via a capture molecule of the plurality of capture molecules; (c) removing the tissue sample; (d) detecting the captured polypeptide.

[1798] Embodiment Q14 is the method of embodiment Q13, wherein detecting the captured polypeptide comprises detecting via a spatial barcode of the plurality of spatial barcodes.

[1799] Embodiment Q15 is the method of embodiment Q1, wherein the plurality of capture molecules comprises (i) a first subset of capture molecules configured to capture polypeptides derived from the tissue sample and (ii) second subset of capture molecules configured to capture nucleic acid derived from the tissue sample.

[1800] Embodiment Q16 is the method of embodiment Q15, wherein the second subset of capture molecules is configured to capture ribonucleic acid (RNA).

[1801] Embodiment Q17 is the method of embodiment Q15, wherein the second subset of capture molecules is configured to capture deoxyribonucleic acid (DNA).

[1802] Embodiment Q18 is the method of embodiment Q15, wherein step b comprises capturing a polypeptide derived from the tissue sample via a capture molecule of the first subset of capture molecules.

[1803] Embodiment Q19 is the method of embodiment Q15, wherein step b further comprises capturing a nucleic acid derived from the tissue sample via a capture molecules of the second subset of capture molecules.

[1804] Embodiment Q20 is the method of embodiment Q19, further comprising, following step d, detecting the captured nucleic acid via the capture molecule attached to the pre-defined spatial region.

Exemplary Embodiments R

[1805] Embodiment R1 is a method for spatially detecting a biological analyte of interest from a biological sample, the method comprising: (a) staining and imaging a biological sample on a support; (b) providing a solution comprising a permeabilization reagent to the biological sample on the support; (c) contacting the biological sample with an array on a substrate, wherein the array comprises one or more capture probe pluralities thereby allowing the one or more pluralities of capture probes to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest; wherein the biological sample is not removed from the support.

[1806] Embodiment R2 is the method of embodiment R1, further comprising selecting a region of interest in the biological sample to subject to spatial transcriptomic analysis.

[1807] Embodiment R3 is the method of embodiment R1, wherein one or more of the one or more capture probe pluralities comprise a capture domain.

[1808] Embodiment R4 is the method of embodiment R1, wherein one or more of the one or more capture probe pluralities comprise a unique molecular identifier (UMI).

[1809] Embodiment R5 is the method of embodiment R1, wherein one or more of the one or more capture probe pluralities comprise a cleavage domain.

[1810] Embodiment R6 is the method of embodiment R5, wherein the cleavage domain comprises a sequence recognized and cleaved by a uracil-DNA glycosylase and/or an endonuclease VIII.

[1811] Embodiment R7 is the method of embodiment R1, wherein one or more capture probes do not comprise a cleavage domain and is not cleaved from the array.

[1812] Embodiment R8 is the method of embodiment R1, wherein the biological analyte comprises DNA or RNA.

[1813] Embodiment R9 is the method of embodiment R1, wherein the biological analyte comprises a protein.

[1814] Embodiment R10 is the method of embodiment R1, wherein the array substrate comprises

one or more microfluidic channels.

[1815] Embodiment R11 is the method of embodiment R1, wherein the array substrate comprises one or more distribution channels.

[1816] Embodiment R12 is the method of embodiment R1, wherein the plurality of probes are attached are directly or indirectly attached to the support.

[1817] Embodiment R13 is the method of embodiment R1, wherein analyzing the captured biological analyte comprises performing in-situ sequencing.

[1818] Embodiment R14 is the method of embodiment R12, wherein analyzing the captured biological analyte comprises one or more cycles of hybridization with labelled probes.

[1819] Embodiment R15 is the method of embodiment R13, wherein the labelled probes are labelled with at least one of a fluorescent molecule, a fluorescent protein, a radioactive molecule, or an enzyme.

[1820] Embodiment R16 is the method of embodiment R3, wherein the unique molecular identifier is amplified prior to analyzing the presence of one or more captured biological analytes.

[1821] Embodiment R17 is the method of embodiment R15, wherein analyzing the captured biological analytes comprises high-throughput sequencing.

Exemplary Embodiments S

[1822] Embodiment S1 is a method for preparing a high-density spatially barcoded array, comprising: (a) providing an array comprising a plurality of features comprising capture probes; (b) adding a first hydrogel to the array; (c) copying a capture probe of the first array such that the capture probe is copied into the first hydrogel; (d) reducing the size of the first hydrogel comprising the copied capture probe by removing water, forming a first shrunken hydrogel comprising the copied capture probe; (e) contacting the first shrunken hydrogel comprising the copied capture probe to a second hydrogel; (f) copying the capture probe in the first shrunken hydrogel such that the capture probe is copied into the second hydrogel; and (g) reducing the size of the second hydrogel comprising the copied capture probe by removing water, forming a second shrunken hydrogel comprising the copied capture probe; thus generating a high-density spatially barcoded array.

[1823] Embodiment S2 is the method of embodiment S1, wherein the reduction in the size of the hydrogel is isometric.

[1824] Embodiment S3 is the method of embodiment S1 or S2, wherein the removal of water comprises a dehydrating solvent.

[1825] Embodiment S4 is the method of embodiment S3, wherein the solvent comprises at least one of an alcohol, a ketone, an aldehyde, acetonitrile, or a combination thereof.

[1826] Embodiment S5 is the method of embodiment S3 or S4, wherein the solvent comprises a solvent selected from the group consisting of: glycerol, ethanol, acetonitrile, acetone, 2-propanol, and a combination thereof.

[1827] Embodiment S6 is the method of any one of embodiments S1-S5, wherein the removal of water comprises a salt.

[1828] Embodiment S7 is the method of embodiment S6, wherein the salt comprises an inorganic salt.

[1829] Embodiment S8 is the method of any one of embodiments S1-S7, wherein the removal of water comprises an acid.

[1830] Embodiment S9 is the method of any one of embodiments S1-S8, wherein the removal of water comprises a dehydrating process.

[1831] Embodiment S10 is the method of any one of embodiments S1-S9, wherein the first shrunken hydrogel is about 2 to about 10 fold smaller in a linear dimension than the pre-shrunk first hydrogel.

[1832] Embodiment S11 is the method of any one of embodiments S1-S10, wherein the first shrunken hydrogel is at least 2-fold smaller in a linear dimension than the pre-shrunk first

hydrogel.

[1833] Embodiment S12 is the method of any one of embodiments S1-S11, wherein the first shrunken hydrogel is at least 4-fold smaller in a linear dimension than the pre-shrunk first hydrogel.

[1834] Embodiment S13 is the method of any one of embodiments S1-S12, wherein the first shrunken hydrogel is at least 8-fold smaller in a linear dimension than the pre-shrunk first hydrogel.

[1835] Embodiment S14 is the method of any one of embodiments S1-S13, wherein the second shrunken hydrogel is about 2 to about 10 fold smaller in a linear dimension than the pre-shrunk second hydrogel.

[1836] Embodiment S15 is the method of any one of embodiments S1-S14, wherein the second shrunken hydrogel is at least 2-fold smaller in a linear dimension than the pre-shrunk second hydrogel.

[1837] Embodiment S16 is the method of any one of embodiments S1-S15, wherein the second shrunken hydrogel is at least 4-fold smaller in a linear dimension than the pre-shrunk second hydrogel.

[1838] Embodiment S17 is the method of any one of embodiments S1-S16, wherein the second shrunken hydrogel is at least 8-fold smaller in a linear dimension than the pre-shrunk second hydrogel.

[1839] Embodiment S18 is the method of any one of embodiments S1-S17, wherein the copying comprises polymerase chain reaction.

[1840] Embodiment S19 is the method of embodiment S18, wherein the first hydrogel comprises PCR reagents comprising: a DNA polymerase, a plurality of primers, and a plurality of nucleotides.

[1841] Embodiment S20 is the method of embodiment S18, wherein the second hydrogel comprises PCR reagents comprising: a DNA polymerase, a plurality of primers, and a plurality of nucleotides.

[1842] Embodiment S21 is the method of any one of embodiments S1-S20, wherein members of the plurality of features are cross-linked to the first hydrogel.

[1843] Embodiment S22 is the method of any one of embodiments S1-S21, wherein members of the plurality of features are cross-linked to the second hydrogel.

[1844] Embodiment S23 is the method of any one of embodiments S1-S22, wherein the first hydrogel is prepared from hydrogel subunits comprising at least one of: acrylamide, bis-acrylamide, a poly(ethylene glycol), a polyaliphatic polyurethane, a polyether polyurethane, a polyester polyurethane, a polyethylene copolymer, a polyamide, a polyvinyl alcohol, polypropylene glycol, polytetramethylene oxide, polyvinyl pyrrolidone, polyacrylamide, poly(hydroxyethyl acrylate), poly(hydroxyethyl methacrylate), collagen, hyaluronic acid, chitosan, dextran, agarose, gelatin, alginate, a protein polymer, methylcellulose, and a combination thereof.

[1845] Embodiment S24 is the method of any one of embodiments S1-S23, wherein the second hydrogel is prepared from hydrogel subunits comprising at least one of: acrylamide, bis-acrylamide, a poly(ethylene glycol), a polyaliphatic polyurethane, a polyether polyurethane, a polyester polyurethane, a polyethylene copolymer, a polyamide, a polyvinyl alcohol, polypropylene glycol, polytetramethylene oxide, polyvinyl pyrrolidone, polyacrylamide, poly(hydroxyethyl acrylate), poly(hydroxyethyl methacrylate), collagen, hyaluronic acid, chitosan, dextran, agarose, gelatin, alginate, a protein polymer, methylcellulose, and a combination thereof.

[1846] Embodiment S25 is the method of any one of the embodiments S1-S24, wherein members of the plurality of features comprise a plurality of capture probes.

[1847] Embodiment S26 is the method of embodiment S25, wherein each capture probe comprises a spatial barcode.

[1848] Embodiment S27 is the method of embodiment S25 or S26, wherein each capture probe from the plurality of capture probes comprises a unique molecular identifier.

[1849] Embodiment S28 is the method of any one of the embodiments S25-S27, wherein each capture probe comprises a cleavage domain, a spatial barcode, a unique molecular identifier, and a capture domain.

[1850] Embodiment S29 is a method for spatially profiling a plurality of analytes in a biological sample, comprising: (a) providing a high-density spatially barcoded array of any one of embodiments S1-S25; (b) contacting the biological sample with the high-density spatially barcoded array under conditions such that one or more analytes present in the biological sample are captured by one or more features of the array; and (c) determining the spatial profile of the captured analytes.

[1851] Embodiment S30 is the method of embodiment S29, wherein the biological sample is a tissue sample.

[1852] Embodiment S31 is the method of embodiment S30, wherein the tissue sample is a fresh-frozen tissue sample.

[1853] Embodiment S32 is the method of embodiment S30, wherein the tissue sample comprises a tumor cell.

[1854] Embodiment S33 is the method of embodiment S30, wherein the tissue sample comprises a tissue section.

[1855] Embodiment S34 is the method of any one of embodiments S29-S33, wherein the analytes in the biological sample are migrated to the bead array.

[1856] Embodiment S35 is the method of embodiment S34, wherein the migrating comprises passive migration.

[1857] Embodiment S36 is the method of embodiment S34, wherein the migrating comprises active migration.

[1858] Embodiment S37 is the method of any one of embodiments S29-S36, wherein the one or more analytes comprises at least one of RNA, DNA, protein, lipid, peptide, metabolite, small molecule, and a cell labeling agent.

Exemplary Embodiments T

[1859] Embodiment T1 is a method for spatially detecting a biological analyte of interest from a biological sample, the method comprising: (a) staining and imaging a biological sample on a support; (b) providing a solution comprising a permeabilization reagent to the biological sample on the support; (c) contacting the biological sample with an array on a substrate, wherein the array comprises one or more capture probe pluralities thereby allowing the one or more pluralities of capture probes to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest; wherein the biological sample is not removed from the support.

[1860] Embodiment T2 is the method of embodiment T1, further comprising selecting a region of interest in the biological sample to subject to spatial transcriptomic analysis.

[1861] Embodiment T3 is the method of embodiment T1, wherein one or more of the one or more capture probe pluralities comprise a capture domain.

[1862] Embodiment T4 is the method of embodiment T1, wherein one or more of the one or more capture probe pluralities comprise a unique molecular identifier (UMI).

[1863] Embodiment T5 is the method of embodiment T1, wherein one or more of the one or more capture probe pluralities comprise a cleavage domain.

[1864] Embodiment T6 is the method of embodiment T5, wherein the cleavage domain comprises a sequence recognized and cleaved by a uracil-DNA glycosylase and/or an endonuclease VIII.

[1865] Embodiment T7 is the method of embodiment T1, wherein one or more capture probes do not comprise a cleavage domain and is not cleaved from the array.

[1866] Embodiment T8 is the method of embodiment T1, wherein the biological analyte comprises DNA or RNA.

[1867] Embodiment T9 is the method of embodiment T1, wherein the biological analyte comprises

a protein.

[1868] Embodiment T10 is the method of embodiment T1, wherein the array substrate comprises one or more microfluidic channels.

[1869] Embodiment T11 is the method of embodiment T1, wherein the array substrate comprises one or more distribution channels.

[1870] Embodiment T12 is the method of embodiment T1, wherein the plurality of probes are attached are directly or indirectly attached to the support.

[1871] Embodiment T13 is the method of embodiment T1, wherein analyzing the captured biological analyte comprises performing in-situ sequencing.

[1872] Embodiment T14 is the method of embodiment T12, wherein analyzing the captured biological analyte comprises one or more cycles of hybridization with labelled probes.

[1873] Embodiment T15 is the method of embodiment T13, wherein the labelled probes are labelled with at least one of a fluorescent molecule, a fluorescent protein, a radioactive molecule, or an enzyme.

[1874] Embodiment T16 is the method of embodiment T3, wherein the unique molecular identifier is amplified prior to analyzing the presence of one or more captured biological analytes.

[1875] Embodiment T17 is the method of embodiment T15, wherein analyzing the captured biological analytes comprises high-throughput sequencing.

[1876] Embodiment T18 is the method of embodiment T1, wherein the array comprises features, wherein a feature comprises one or more capture probes.

[1877] Embodiment T19 is the method of embodiment T18, wherein the array comprises at least 4000 features.

[1878] Embodiment T20 is the method of embodiment T19, wherein the array comprises approximately 5000 features.

[1879] Embodiment T21 is the method of embodiment T18, wherein the array features are between 50 micrometers and 80 micrometers in diameter.

[1880] Embodiment T22 is the method of embodiment T21, wherein the array features are between 60 micrometers and 70 micrometers.

[1881] Embodiment T23 is the method of embodiment T18, wherein the features of the array are patterned.

[1882] Embodiment T25 is the method of embodiment T23, wherein the center of one feature of the array is between 80 micrometers and 120 micrometers from the center of another feature of the array.

[1883] Embodiment T26 is the method of embodiment T1, wherein the array is approximately 8 mm by 8 mm.

[1884] Embodiment T27 is the method of embodiment T1, wherein the array is smaller than 8 mm by 8 mm.

[1885] Embodiment T27 is the method of embodiment T1, wherein the array is surrounded by fiducial markers.

[1886] Embodiment T28 is the method of embodiment T1, wherein the fiducial markers allow for detection of rotation or mirroring.

Exemplary Embodiments U

[1887] Embodiment U1 is a method for preparing a composition comprising: providing a substrate comprising a first oligonucleotide segment attached to the substrate through its 3'-end or its 5'-end; and preparing by spatially-directed in situ synthesis on the unattached end of the first segment a plurality of second oligonucleotide segments of 8-mer to 30-mer each having a different sequence and located in a different region of the substrate.

[1888] Embodiment U2 is the method of embodiment U1, wherein the method further comprises: attaching a plurality of third oligonucleotide segments to the unattached ends of the plurality of second oligonucleotide segments.

[1889] Embodiment U3 is the method of any one of embodiments U1-U2, wherein the first oligonucleotide segment is attached to the substrate through its 5'-end and the plurality of second oligonucleotide segments are prepared by spatially-directed in situ synthesis on the 3'-end of the first segment.

[1890] Embodiment U4 is the method of any one of embodiments U1-U2, wherein the first oligonucleotide segment is attached to the substrate through its 3'-end and the plurality of second oligonucleotide segments are prepared by spatially-directed in situ synthesis on the 5'-end of the first segment.

[1891] Embodiment U5 is the method of any one of embodiments U1-U4, wherein phosphodiester and exocyclic amine moieties of the first oligonucleotide segment are protected.

[1892] Embodiment U6 is the method of any one of embodiments U1-U5, wherein the second oligonucleotide segments have a length selected from 8-mer, 9-mer, 10-mer, 11-mer, 12-mer, 13-mer, 14-mer, 15-mer, 16-mer, 17-mer, 18-mer, 19-mer, 20-mer, 21-mer, 22-mer, 23-mer, 24-mer, 25-mer, 26-mer, 27-mer, 28-mer, 29-mer, or 30-mer.

[1893] Embodiment U7 is the method of any one of embodiments U1-U6, wherein the in situ synthesis is light-directed synthesis.

[1894] Embodiment U8 is the method of any one of embodiments U1-U6, wherein the in situ synthesis is spatially-directed synthesis.

[1895] Embodiment U9 is the method of any one of embodiments U1-U8, wherein the regions of the substrate are less than about 2500 μm^2 , less than about 1600 μm^2 , less than about 400 μm^2 , less than about 100 μm^2 , less than about 50 μm^2 , less than about 25 μm^2 , less than about 15 μm^2 , less than about 10 μm^2 , or less than about 5 μm^2 .

[1896] Embodiment U10 is the method of any one of embodiments U1-U8, wherein the number of different second oligonucleotide segment sequences comprises at least about 100,000, at least about 250,000, at least about 500,000, at least about 750,000, at least about 1,000,000, at least about 1,500,000, at least about 3,000,000, at least about 5,000,000, at least about 10,000,000, at least about 100,000,000, at least about 500,000,000, or at least about 1,000,000,000.

[1897] Embodiment U11 is the method of any one of embodiments U1-U10, wherein the number of different second oligonucleotide segment sequences comprises about 100,000 to about 1,000,000,000, about 750,000 to about 1,500,000, about 1,000,000 to about 1,000,000,000, about 1,000,000 to about 10,000,000, or about 1,500,000 to about 5,000,000.

[1898] Embodiment U12 is the method of any one of embodiments U1-U11, wherein the second oligonucleotide segment in each of the different regions of the substrate has a purity of at least 75%, at least 80%, at least 85%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or more.

[1899] Embodiment U13 is the method of any one of embodiments U1-U12, wherein the first oligonucleotide segment comprises protected phosphodiester and exocyclic amine moieties.

[1900] Embodiment U14 is the method of any one of embodiments U1-U12, wherein the substrate further comprises a complementary oligonucleotide hybridized to the first oligonucleotide segment.

[1901] Embodiment U15 is the method of any one of embodiments U1-U14, wherein the first oligonucleotide segment has length of at least about 2-mer, at least about 5-mer, at least about 8-mer, at least about 10-mer, or at least about 15-mer at least about 18-mer, at least about 20-mer, at least about 22-mer, or at least about 25-mer, at least about 27-mer, at least about 30-mer, at least about 33-mer, or at least about 35-mer.

[1902] Embodiment U16 is the method of any one of embodiments U1-U15, wherein the first oligonucleotide segment has a length of about 2-mer to about 45-mer, about 5-mer to about 40-mer, about 15-mer to about 40-mer about 25-mer to about 40-mer, about 25-mer to about 35-mer, or about 27-mer to about 33-mer.

[1903] Embodiment U17 is the method of any one of embodiments U1-U16, wherein the first

oligonucleotide segment is prepared by synthesis off the substrate and subsequently attached to the substrate.

[1904] Embodiment U18 is the method of embodiment U17, wherein the phosphodiester and exocyclic amine moieties of the first oligonucleotide segment are protected prior to attachment to the substrate.

[1905] Embodiment U19 is the method of embodiment U18, wherein the phosphodiester and exocyclic amine moieties of the first oligonucleotide segment are protected after attachment to the substrate.

[1906] Embodiment U20 is the method of any one of embodiments U1-U16, wherein the first oligonucleotide segment is prepared by in situ synthesis on the substrate.

[1907] Embodiment U21 is the method of embodiment U20, wherein the in situ synthesis is not light-directed.

[1908] Embodiment U22 is the method of embodiment U20, wherein the in situ synthesis is not spatially-directed.

[1909] Embodiment U23 is the method of any one of embodiments U1-U22, wherein the first oligonucleotide segment sequence comprises a cleavage site.

[1910] Embodiment U24 is the method of any one of embodiments U1-U23, wherein the first oligonucleotide segment is attached to the substrate through a linker.

[1911] Embodiment U25 is the method of embodiment U24, wherein the linker comprises a chemical group selected from: silane, amine, peptide, polyethylene glycol, biotin, desthiobiotin, streptavidin, thiol, acrydite, azide, digoxigenin, hexynyl, or a combination thereof.

[1912] Embodiment U26 is the method of embodiment U25, wherein the linker is branched or dendrimeric.

[1913] Embodiment U27 is the method of any one of embodiments U2-U26, wherein the plurality of third oligonucleotide segments are attached by ligation.

[1914] Embodiment U28 is the method of embodiment U27, wherein any protected moieties of the first and second oligonucleotide segments are deprotected prior to ligating the plurality of third oligonucleotide segments.

[1915] Embodiment U29 is the method of any one of embodiments U2-U28 wherein the plurality of third oligonucleotide segments have a length of at least about 5-mer, at least about 7-mer, at least about 9-mer, at least about 13-mer, at least about 20-mer, at least about 25-mer, at least about 27-mer, at least about 30-mer, at least about 33-mer, or at least about 35-mer.

[1916] Embodiment U30 is the method of any one of embodiments U2-U29, wherein the plurality of third oligonucleotide segments have a length of about 5-mer to about 40-mer, 5-mer to about 20-mer, 5-mer to about 10-mer, about 10-mer to about 35-mer, about 15-mer to about 30-mer, about 25-mer to about 35-mer, or about 27-mer to about 33-mer.

[1917] Embodiment U31 is the method of any one of embodiments U2-U30, wherein each of the plurality of third oligonucleotide segments comprise a capture sequence of at least 6-mer at the 3'-end.

[1918] Embodiment U32 is the method of embodiment U31, wherein the capture sequence comprises a poly-T sequence.

[1919] Embodiment U33 is the method of any one of embodiments U2-U32, wherein each of the plurality of third oligonucleotide segments comprise a different random sequence of 6-mer to 20-mer.

[1920] Embodiment U34 is a composition comprising a plurality of oligonucleotides attached to a plurality of regions on a substrate, wherein each oligonucleotide comprises: a first oligonucleotide segment of 2-mer to 50-mer attached to the substrate through its 3'-end or its 5'-end; and a second oligonucleotide segment of 8-mer to 30-mer that is attached to the unattached end of the first segment, wherein the second segment comprises a sequence that identifies the region where it is located.

- [1921] Embodiment U35 is the composition of embodiment U34, wherein the first and second oligonucleotide segments comprise protected phosphodiester and exocyclic amine moieties.
- [1922] Embodiment U36 is the composition of any one of embodiments U34-35, wherein the first oligonucleotide segment is attached to the substrate through its 5'-end and the plurality of second oligonucleotide segments are attached to the 3'-end of the first segment.
- [1923] Embodiment U37 is the composition of any one of embodiments U34-U35, wherein the first oligonucleotide segment is attached to the substrate through its 3'-end and the plurality of second oligonucleotide segments are attached to the 5'-end of the first segment.
- [1924] Embodiment U38 is the composition any one of embodiments U34-U37, wherein each oligonucleotide further comprises: a third oligonucleotide segment of 5-mer to 40-mer that is attached to the unattached end of the second segment, wherein the third segment comprises a capture sequence.
- [1925] Embodiment U39 is the composition of any one of embodiments U34-U38, wherein the second oligonucleotide segments have a length selected from 8-mer, 9-mer, 10-mer, 11-mer, 12-mer, 13-mer, 14-mer, 15-mer, 16-mer, 17-mer, 18-mer, 19-mer, 20-mer, 21-mer, 22-mer, 23-mer, 24-mer, 25-mer, 26-mer, 27-mer, 28-mer, 29-mer, or 30-mer.
- [1926] Embodiment U40 is the composition of any one of embodiments U34-U39, wherein the regions are less than about 2500 μm^2 , less than about 1600 μm^2 , less than about 400 μm^2 , less than about 100 μm^2 , less than about 50 μm^2 , less than about 25 μm^2 , less than about 15 μm^2 , less than about 10 μm^2 , less than about 5 μm^2 , or less than about 1 μm^2 .
- [1927] Embodiment U41 is the composition of any one of embodiments U34-U40, wherein the number of different second oligonucleotide segments is at least about 100,000, at least about 250,000, at least about 500,000, at least about 750,000, at least about 1,000,000, at least about 1,500,000, at least about 3,000,000, at least about 5,000,000, at least about 10,000,000, at least about 100,000,000, at least about 500,000,000, or at least about 1,000,000,000.
- [1928] Embodiment U42 is the composition of any one of embodiments U34-U41, wherein the number of different second oligonucleotide segment sequences comprises about 100,000 to about 1,000,000,000, about 750,000 to about 1,500,000, about 1,000,000 to about 1,000,000,000, about 1,000,000 to about 10,000,000, or about 1,500,000 to about 5,000,000.
- [1929] Embodiment U43 is the composition of any one of embodiments U34-U42, wherein the second oligonucleotide segment in each of the different regions of the substrate has a purity of at least 75%, at least 80%, at least 85%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or more.
- [1930] Embodiment U44 is the composition of any one of embodiments U34-U43, wherein the length of the oligonucleotide in each region is at least 15-mer, at least 20-mer, at least 40-mer, at least 50-mer, or at least 60-mer.
- [1931] Embodiment U45 is the composition of any one of embodiments U34-U44, wherein the length of the oligonucleotide in each region is from 15-mer to 100-mer, 20-mer to 100-mer, 40-mer to 100-mer, from 50-mer to 90-mer, or from 60-mer to 80-mer.

Exemplary Embodiments V

- [1932] Embodiment V1 is a method for spatially detecting a biological analyte of interest from a biological sample, the method comprising: (a) staining and imaging a biological sample; (b) providing a solution comprising a permeabilization reagent to the biological sample; (c) contacting the biological sample with an array, wherein the array comprises one or more capture probe pluralities, wherein a capture probe plurality comprises one or more capture probes comprising a spatial barcode and a capture domain, thereby allowing the one or more capture domains to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest.

[1933] Embodiment V2 is a method for spatially detecting a biological analyte of interest from a biological sample, the method comprising: (a) fixing a biological sample; (b) staining and imaging a biological sample; (c) contacting the biological sample with an array, wherein the array comprises one or more capture probe pluralities, wherein a capture probe plurality comprises one or more capture probes comprising a spatial barcode and a capture domain, thereby allowing the one or more capture domains to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest, wherein the biological sample is not subjected to permeabilization reagents.

[1934] Embodiment V3 is the method of embodiment V1 or V2, wherein fixing the biological sample comprises acetone fixation.

[1935] Embodiment V4 is the method of embodiment V1 or V2, further comprising selecting a region of interest in the biological sample.

[1936] Embodiment V5 is the method of embodiment V1 or V2, wherein one or more capture probes comprise a unique molecular identifier (UMI).

[1937] Embodiment V6 is the method of embodiment V1 or V2, wherein one or more capture probes comprise a cleavage domain.

[1938] Embodiment V7 is the method of embodiment V1 or V2, wherein the biological analyte comprises DNA or RNA.

[1939] Embodiment V8 is the method of embodiment V1 or V2, wherein the biological analyte comprises protein.

[1940] Embodiment V9 is the method of embodiment V1 or V2, wherein the array comprises one or more microfluidic channels.

[1941] Embodiment V10 is the method of embodiment V1 or V2, wherein the plurality of probes are attached directly or indirectly to the array.

[1942] Embodiment V11 is the method of embodiment V1 or V2, wherein analyzing the captured biological analyte comprises performing in-situ sequencing.

[1943] Embodiment V12 is the method of embodiment V1 or V2, wherein staining the biological sample comprises hematoxylin and eosin staining.

[1944] Embodiment V13 is the method of embodiment V1 or V2, wherein staining the biological sample comprises Periodic Acid Schiff staining.

[1945] Embodiment V14 is the method of embodiment V13, wherein the biological sample comprises kidney tissue.

[1946] Embodiment V15 is the method of embodiment V1 or V2, wherein the one or more capture probes in step (c) captures mRNA.

[1947] Embodiment V16 is the method of embodiment V15, wherein a plurality of cDNA molecules are generated from the captured mRNA, by extending the capture probes enzymatically using the captured mRNA as an extension template.

[1948] Embodiment V17 is the method of embodiment V1 or V2, wherein the biological sample is on a solid support.

[1949] Embodiment V18 is a method for spatially detecting a biological analyte of interest from a biological sample, the method comprising: (a) staining and imaging a biological sample; (b) providing a solution comprising a permeabilization reagent to the biological sample; (c) contacting the biological sample with an array, wherein the array comprises one or more capture probe pluralities, wherein a capture probe plurality comprises one or more capture probes comprising a spatial barcode and a capture domain, thereby allowing the one or more pluralities of capture domains probes to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest.

[1950] Embodiment V19 is the method of embodiment V18, wherein the one or more capture probes in step (c) captures mRNA.

[1951] Embodiment V20 is the method of embodiment V19, further comprising performing first

strand cDNA synthesis by generating a plurality of cDNA molecules from the captured mRNA.

[1952] Embodiment V21 is the method of embodiment V20, wherein ribosomal RNA depletion probes are added when performing first strand cDNA synthesis.

[1953] Embodiment V22 is the method of embodiment V20, wherein mitochondrial RNA depletion probes are added when performing first strand cDNA synthesis.

[1954] Embodiment V23 is the method of embodiment V20, wherein ribosomal RNA depletion probes and mitochondrial RNA depletion probes are added when performing first strand cDNA synthesis.

[1955] Embodiment V24 is the method of embodiment V21, further comprising removing the biological sample from the array.

[1956] Embodiment V25 is the method of embodiment V24, further comprising cleaving the capture probes after removal of the biological sample.

[1957] Embodiment V26 is the method of embodiment V25, further comprising adding carrier RNA before performing second strand cDNA synthesis.

[1958] Embodiment V27 is the method of embodiment V25, further comprising performing second strand cDNA synthesis.

[1959] Embodiment V28 is the method of embodiment V27, further comprising performing in vitro transcription.

[1960] Embodiment V29 is the method of embodiment V28, further comprising adding carrier RNA before performing RNA purification and quantification.

[1961] Embodiment V30 is the method of embodiment V26 or V29, wherein adding carrier RNA comprises adding 100 ng of carrier RNA.

[1962] Embodiment V31 is the method of embodiment V29, wherein the RNA purification and quantification comprises performing a size selection RNA clean up.

[1963] Embodiment V32 is the method of embodiment V18, wherein the biological sample is on a solid support.

Exemplary Embodiments W

[1964] Embodiment W1 is a method for profiling at least one biological analyte present in a biological sample, wherein the biological sample is a cell culture, an organism, or an organoid, the method comprising: [1965] (a) contacting a biological sample with a first substrate comprising a plurality of capture probes, wherein a capture probe comprises a spatial barcode and a capture domain; [1966] (b) releasing a biological analyte from the biological sample such that the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; [1967] (c) detecting the bound biological analyte; and [1968] (d) correlating the biological analyte with a molecular barcode of the capture probe at the distinct spatial position of the substrate; [1969] thus profiling the biological analyte as having been present in the biological sample at a distinct spatial position.

[1970] Embodiment W2 is the method of embodiment W1, wherein the organism is a nematode or a zebrafish.

[1971] Embodiment W3 is method of embodiment W2, wherein the nematode is *Caenorhabditis elegans*.

[1972] Embodiment W4 is the method of embodiment W1, wherein the organoid is selected from the group consisting of: a cerebral organoid, an intestinal organoid, a stomach organoid, a lingual organoid, a thyroid organoid, a thymic organoid, a testicular organoid, a hepatic organoid, a pancreatic organoid, an epithelial organoid, a lung organoid, a kidney organoid, a gastruloid, a cardiac organoid, and a retinal organoid.

[1973] Embodiment W5 is the method of embodiment W1, wherein the cell culture comprises a plurality of cells.

[1974] Embodiment W6 is the method of embodiment W5, wherein the plurality of cells are from a suspension of cells.

[1975] Embodiment W7 is the method of embodiment W5 or W6, wherein the plurality of cells are non-adherent cells.

[1976] Embodiment W8 is the method of any one of embodiments W5-W7, wherein the plurality of cells are from disassociated tissue or a tissue section.

[1977] Embodiment W9 is the method of any one of embodiments W5-W7, wherein a cell of the plurality of cells is distributed onto the first substrate.

[1978] Embodiment W10 is the method of embodiment W9, wherein the first substrate comprises a plurality of features, wherein a cell of the plurality of cells is distributed onto the first substrate such that the cell occupies a distinct feature on the first substrate.

[1979] Embodiment W11 is the method of any one of embodiments W5-W10, wherein a cell of the plurality of cells is distributed onto a second substrate.

[1980] Embodiment W12 is the method of embodiment W11, wherein the second substrate is contacted with the first substrate.

[1981] Embodiment W13 is the method of embodiment W12, wherein the second substrate is contacted with the first substrate prior to releasing the biological analyte from the biological sample.

[1982] Embodiment W14 is the method of any one of embodiments W5-W13, wherein a cell of the plurality of cells is immobilized on the substrate after the cell is distributed onto the substrate.

[1983] Embodiment W15 is the method of any one of embodiments W5-W14, wherein the cell of the plurality of cells is immobilized on the substrate prior to releasing the biological analyte.

[1984] Embodiment W16 is the method of any one of embodiments W5-W15, wherein a cell of the plurality of cells is immobilized by adding a hydrogel.

[1985] Embodiment W17 is the method of any one of embodiments W5-W16, wherein a cell of the plurality of cells is immobilized by applying an electric field.

[1986] Embodiment W18 is the method of any one of embodiments W5-W17, wherein the substrate comprises a plurality of capture antibodies.

[1987] Embodiment W19 is the method of embodiment W18, wherein a cell of the plurality of cells binds to a capture antibody of the plurality of capture antibodies.

[1988] Embodiment W20 is the method of embodiment W18 or W19, wherein the substrate comprises a plurality of features, and wherein the plurality of features comprises members of the plurality of capture antibodies.

[1989] Embodiment W21 is the method of any one of embodiments W18-W20, wherein the plurality of capture antibodies comprises antibodies against a cell surface marker.

[1990] Embodiment W22 is the method of embodiment W21, wherein the cell surface marker comprises at least one of CD45, CD3, CD4, CD8, CD56, CD19, CD20, CD11c, CD14, CD33, CD66b, CD34, CD41, CD61, CD235a, CD146, epithelial cellular adhesion molecule (EpCAM), and a combination thereof.

[1991] Embodiment W23 is the method of any one of embodiments W5-W22, wherein members of the plurality of cells are distributed over the substrate until about all of the features on the substrate are bound to a cell.

[1992] Embodiment W24 is the method of any one of embodiments W5-W23, wherein the first substrate comprises a plurality of features, and wherein the features are smaller than the minimum cell diameter.

[1993] Embodiment W25 is the method of any one of embodiments W5-W24, wherein the first substrate comprises a plurality of features, and wherein cells not bound to a feature on the substrate are washed off.

[1994] Embodiment W26 is the method of embodiment W25, wherein cells not bound to a feature on the substrate are washed off prior to releasing the biological analyte.

[1995] Embodiment W27 is the method of embodiment W5, wherein the plurality of cells comprises adherent cells.

[1996] Embodiment W28 is the method of embodiment W27, wherein the adherent cells are cultured on the first substrate.

[1997] Embodiment W29 is the method of embodiment W29, wherein the adherent cells are cultured on a second substrate.

[1998] Embodiment W30 is the method of embodiment W28 or W29, wherein the substrate comprises a cell-permissive coating.

[1999] Embodiment W31 is the method of embodiment W30, wherein the cell-permissive coating comprises at least one of a biological material and a synthetic material.

[2000] Embodiment W32 is the method of embodiment W30 or W31, wherein the cell-permissive coating comprises an extracellular matrix component.

[2001] Embodiment W33 is the method of embodiment W32, wherein the cell-permissive coating comprises one or more ECM components comprising at least one of collagen type I, collagen type II, collagen type IV, elastin, fibronectin, laminin, and vitronectin.

[2002] Embodiment W34 is the method of any one of embodiments W30-W33, wherein the cell-permissive coating comprises a synthetic material comprising at least one of poly-lysine, poly-L-ornithine, and a biocompatible silicone (e.g., CYTOSOFT®).

[2003] Embodiment W35 is the method of any one of embodiments W30-W34, wherein the cell-permissive coating comprises a solubilized basement membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma.

[2004] Embodiment W36 is the method of any one of embodiments W30-W35, wherein the cell-permissive coating comprises collagen.

[2005] Embodiment W37 is the method of any one of embodiments W27-W36, wherein the adherent cells are from cell lines comprising at least one of BT549, HS 578T, MCF7, MDA-MB-231, MDAMB-468, T-47D, SF268, SF295, SF539, SNB-19, SNB-75, U251, Colo205, HCC 2998, HCT-116, HCT-15, HT29, KM12, SW620, 786-O, A498, ACHN, CAKI, RXF 393, SN12C, TK-10, UO-31, A549, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H460, NCI-H522, LOX IMVI, M14, MALME-3M, MDA-MB-435, SK-, EL-2, SK-MEL-28, SK-MEL-5, UACC-257, UACC-62, IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3, NCI-ADR-RES, DU145, PC-3, DU145, H295R, HeLa, KBM-7, LNCaP, MCF-7, MDA-MB-468, PC3, SaOS-2, SH-SY5Y, T-47D, THP-1, U87, vero, MC3T3, GH3, PC12, dog MDCK kidney epithelial, *Xenopus* A6 kidney epithelial, zebrafish AB9, and Sf9 insect epithelial cell lines.

[2006] Embodiment W38 is the method of any one of embodiments W1-W37, wherein releasing the biological analyte comprises permeabilizing one or more cells of the biological sample.

[2007] Embodiment W39 is the method of embodiment W38, further comprising fixing the cells prior to permeabilization.

[2008] Embodiment W40 is the method of embodiment W39, wherein the cells in the biological sample are fixed and permeabilized prior to releasing the biological analyte from the biological sample.

[2009] Embodiment W41 is the method of any one of embodiments W1-W40, further comprising imaging the biological sample.

[2010] Embodiment W42 is the method of embodiment W41, wherein the imaging is performed prior to releasing the biological analyte from the biological sample.

[2011] Embodiment W43 is the method of embodiment W41, wherein the imaging is performed after contacting the biological sample with the first substrate comprising a plurality of capture probes.

[2012] Embodiment W44 is the method of any one of embodiments W41-W43, wherein the imaging is used to determine a region of interest in the biological sample.

[2013] Embodiment W45 is the method of any one of embodiments W41-W44, wherein the imaging is used to determine the morphology of the biological sample or a region of the biological sample.

[2014] Embodiment W46 is the method of embodiment W45, wherein the morphology of the biological sample or a region of the biological sample is correlated to the spatial location of the biological analyte.

[2015] Embodiment W47 is the method of any one of embodiments W1-W46, further comprising imaging one or more cells of the biological sample on the substrate.

[2016] Embodiment W48 is the method of embodiment W47, wherein imaging is performed prior to releasing the biological analyte.

[2017] Embodiment W49 is the method of embodiment W47 or W48, wherein imaging is used to determine the morphology of a cell of the biological sample.

[2018] Embodiment W50 is the method of embodiment W49, wherein the morphology of a cell of the biological sample is correlated to the biological analyte.

[2019] Embodiment W51 is a method for profiling a biological analyte present in a cell, comprising: [2020] (a) binding a cell of a plurality of cells to a feature of a plurality of features on an array, wherein a feature comprises a plurality of capture antibodies and a plurality of capture probes; [2021] (b) releasing a biological analyte from the cell such that the biological analyte is bound by a capture probe at a distinct spatial position of the array; [2022] (c) detecting the bound biological analyte; and [2023] (d) correlating the biological analyte with a molecular barcode of the capture probe at the distinct spatial position of the array, [2024] thus profiling the biological analyte as having been present in the cell.

[2025] Embodiment W52 is the method of embodiment W51, wherein the plurality of capture antibodies comprises antibodies against one or more cell surface markers.

[2026] Embodiment W53 is the method of embodiment W52, wherein the one or more cell surface markers comprise at least one of CD45, CD3, CD4, CD8, CD56, CD19, CD20, CD11c, CD14, CD33, CD66b, CD34, CD41, CD61, CD235a, CD146, and epithelial cellular adhesion molecule (EpCAM).

[2027] Embodiment W54 is the method of any one of embodiments W51-W53, wherein members of the plurality of cells are distributed over the substrate prior to the binding of members of the plurality of cells to members of the plurality of capture antibodies.

[2028] Embodiment W55 is the method of embodiment W54, wherein the cells are distributed over the substrate until about all of the features on the substrate are bound to a cell.

[2029] Embodiment W56 is the method of any one of embodiments W51-W55, wherein each feature is smaller than the minimum cell diameter.

[2030] Embodiment W57 is the method of any one of embodiments W51-W56, wherein cells not bound to a feature on the substrate are washed off.

[2031] Embodiment W58 is the method of embodiment W57, wherein cells not bound to a feature on the substrate are washed off prior to releasing the biological analyte.

[2032] Embodiment W59 is the method of any one of embodiments W51-W58, further comprising imaging one or more cells on the substrate.

[2033] Embodiment W60 is the method of embodiment W59, wherein imaging is performed prior to releasing the biological analyte.

[2034] Embodiment W61 is the method of embodiment W59 or W60, wherein imaging is used to determine the morphology of a cell.

[2035] Embodiment W62 is the method of embodiment W61, wherein the morphology of a cell from the suspension of cells is correlated to the biological analyte.

[2036] Embodiment W63 is a method for profiling a biological analyte present in a cell, comprising: [2037] (a) allowing a cell of a plurality of cells to migrate on a substrate comprising a plurality of capture probes, wherein a capture probe comprises a spatial barcode and a capture domain; [2038] (b) releasing a biological analyte from the cell such that the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; [2039] (c) detecting the bound biological analyte; and [2040] (d) correlating the biological analyte with a molecular

barcode of the capture probe at the distinct spatial position of the substrate, [2041] thus profiling the biological analyte as having been present in a cell at a distinct spatial position.

[2042] Embodiment W64 is the method of embodiment W63, wherein one or more channels are applied to the substrate.

[2043] Embodiment W65 is the method of embodiment W64, wherein the one or more channels confine cell migration to one dimension along the substrate.

[2044] Embodiment W66 is the method of embodiment W64 or W65, wherein the one or more channels confine cell migration along one axis of the substrate.

[2045] Embodiment W67 is the method of any one of embodiments W64-W66, wherein each channel does not overlap with any other channel.

[2046] Embodiment W68 is the method of any one of embodiments W64-W67, wherein each channel has a width of about 2 to about 10 μm .

[2047] Embodiment W69 is the method of any one of embodiments W63-W68, wherein one or more channels are filled with cell culture medium.

[2048] Embodiment W70 is the method of any one of embodiments W63-W69, wherein the substrate and/or the one or more channels comprise one or more extracellular matrix components.

[2049] Embodiment W71 is the method of any one of embodiments W63-W70 wherein the substrate is coated with the one or more extracellular matrix (ECM) components.

[2050] Embodiment W72 is the method of any one of embodiments W64-W71, wherein the one or more channels are coated with the one or more extracellular matrix components.

[2051] Embodiment W73 is the method of any one of embodiments W71-W72, wherein the coating comprises a chemoattractant.

[2052] Embodiment W74 is the method of embodiment W73, wherein the chemoattractant is present at only one end of the one or more channels.

[2053] Embodiment W75 is the method of embodiment W73 or W74, wherein the chemoattractant is present as a gradient along the one or more channels.

[2054] Embodiment W76 is the method of any one of embodiments W73-W75, wherein the chemoattractant is bound to one or more ECM components.

[2055] Embodiment W77 is the method of any one of embodiments W71-W72, wherein the coating comprises a chemorepellant.

[2056] Embodiment W78 is the method of embodiment W77, wherein the chemorepellant is present at only one end of the one or more channels.

[2057] Embodiment W79 is the method of embodiment W77 or W78, wherein the chemorepellant is present as a gradient along the one or more channels.

[2058] Embodiment W80 is the method of any one of embodiments W77-W79, wherein the chemorepellant is bound to one or more ECM components.

[2059] Embodiment W81 is the method of any one of embodiments W70-W80, wherein the one or more extracellular matrix components comprises at least one of basement membrane extract (BME), laminin I, collagen I, collagen IV, fibronectin, vitronectin, elastin, or a combination thereof.

[2060] Embodiment W82 is the method of any one of embodiments W63-W81, further comprising imaging one or more cells on the substrate.

[2061] Embodiment W83 is the method of embodiment W82, wherein imaging is performed prior to releasing the biological analyte.

[2062] Embodiment W84 is the method of embodiment W82 or W83, wherein imaging is performed prior to allowing the cell of the plurality of cells to migrate.

[2063] Embodiment W85 is the method of any one of embodiments W82-W84, wherein imaging is performed after allowing the cell of the plurality of cells to migrate.

[2064] Embodiment W86 is the method of any one of embodiments W82-W85, wherein imaging is used to determine the spatial location of a cell on the substrate.

[2065] Embodiment W87 is the method of any one of embodiments W82-W86, wherein imaging is used to determine the spatial location of the cell in a channel.

[2066] Embodiment W88 is the method of embodiment W86, wherein imaging is used to determine the spatial location of the cell in a channel before and after allowing the cell to migrate.

[2067] Embodiment W89 is the method of embodiment W88, wherein the cell migration distance is determined.

[2068] Embodiment W90 is the method of embodiment W89, wherein the biological analyte is correlated with the cell migration distance.

[2069] Embodiment W91 is the method of any one of embodiments W82-W90, wherein imaging is used to determine the morphology of the cell.

[2070] Embodiment W92 is the method of embodiment W91, wherein the morphology of the cell from the suspension of cells is correlated to the biological analyte.

[2071] Embodiment W93 is the method of any one of embodiments W1-W92, wherein the biological analyte is DNA.

[2072] Embodiment W94 is the method of any one of embodiments W1-W92, wherein the biological analyte is RNA.

[2073] Embodiment W95 is the method of embodiment W94, wherein the capture domain comprises a poly-dT sequence.

[2074] Embodiment W96 is the method of embodiment W94 or W95, wherein the capture domain is configured to hybridize to a poly-A tail of an mRNA.

[2075] Embodiment W97 is the method of any one of embodiments W1-W96, wherein the capture probe further comprises a unique molecular identifier (UMI).

[2076] Embodiment W98 is the method of any one of Embodiments W1-W97, wherein the capture probe further comprises a functional domain.

[2077] Embodiment W99 is the method of any one of Embodiments W1-W98, wherein the capture probe further comprises a cleavage domain.

Exemplary Embodiments X

[2078] Embodiment X1 is a method for profiling a biological analyte present in a biological sample, comprising: [2079] (a) contacting a biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; [2080] (b) releasing a biological analyte from a subcellular region of a cell from the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; [2081] (c) detecting the bound biological analyte; and [2082] (d) correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, [2083] thus profiling the biological analyte as having been present in the biological sample at a distinct spatial position.

[2084] Embodiment X2 is the method of embodiment X1, wherein the subcellular region comprises at least one of cytosol, a nucleus, a mitochondria, and a microsome.

[2085] Embodiment X3 is the method of embodiment X2, wherein the subcellular region is cytosol.

[2086] Embodiment X4 is the method of embodiment X2, wherein the subcellular region is a nucleus.

[2087] Embodiment X5 is the method of embodiment X2, wherein the subcellular region is a mitochondria.

[2088] Embodiment X6 is the method of embodiment X2, wherein the subcellular region is a microsome.

[2089] Embodiment X7 is the method of any one of embodiments X1 or X3-X7, wherein the method further comprises: [2090] (e) releasing a biological analyte from a subcellular region of a cell from the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; [2091] (f) detecting the bound biological analyte from step

(e); and [2092] (g) correlating the biological analyte from step (e) with a barcode of the capture probe at the distinct spatial position of the substrate, [2093] wherein the subcellular region of step (e) is different than the subcellular region of step (b).

[2094] Embodiment X8 is the method of embodiment X7, wherein steps (e) through (g) are repeated one, two, three, or four times.

[2095] Embodiment X9 is the method of embodiment X7 or X8, wherein the subcellular region of the cell of step (b) is cytosol and the subcellular region of the cell of step (e) comprises at least one of a nucleus, a mitochondria, and a microsome.

[2096] Embodiment X10 is the method of any one of embodiments X7-X10, wherein step (e) is performed prior to step (c) detecting the bound biological analyte.

[2097] Embodiment X11 is the method of any one of embodiments X7-X10, wherein step (e) is performed after step (c) detecting the bound biological analyte.

[2098] Embodiment X12 is the method of any one of embodiments X7-X11, wherein the substrate is washed prior to step (e).

[2099] Embodiment X13 is the method of any one of embodiments X7-X12, wherein the biological sample is washed prior to step (e).

[2100] Embodiment X14 is the method of any one of embodiments X1-X6, wherein after releasing the biological analyte, the method further comprises: [2101] (e) contacting the biological sample with a second substrate comprising a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; [2102] (f) releasing a biological analyte from a subcellular region of the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; [2103] (g) detecting the bound biological analyte from step (e); and [2104] (h) correlating the biological analyte from step (e) with a barcode of the capture probe at the distinct spatial position of the substrate, [2105] wherein the subcellular region in (f) is different than the subcellular region in (b).

[2106] Embodiment X15 is the method of embodiment X14, wherein steps (e) through (g) are repeated one, two, three, or four times.

[2107] Embodiment X16 is the method of embodiment X14 or X15, wherein the subcellular region of the cell of step (b) is cytosol and the subcellular region of step (e) comprises at least one of a nucleus, a mitochondria, and a microsome.

[2108] Embodiment X17 is the method of any one of embodiments X14-X16, wherein the biological sample is washed prior to step (e).

[2109] Embodiment X18 is the method of any one of embodiments X1-X17, wherein releasing the biological analyte from the subcellular region the cell from the biological sample comprises permeabilizing the cell or subcellular region therein.

[2110] Embodiment X19 is the method of embodiment X18, wherein the permeabilization comprises electrophoresis.

[2111] Embodiment X20 is the method of embodiment X18 or X19, wherein the permeabilization comprises administering a permeabilization reagent.

[2112] Embodiment X21 is the method embodiment X20, wherein the permeabilization reagent is a selective permeabilization reagent.

[2113] Embodiment X22 is the method of embodiment X21, wherein the selective permeabilization reagent selectively permeabilizes a subcellular region comprising at least one of cytosol, a nucleus, a mitochondria, and a microsome.

[2114] Embodiment X23 is the method of any one of embodiments X20-X22, wherein administering the permeabilization reagent comprises contacting the biological sample with a hydrogel that comprises the permeabilization reagent.

[2115] Embodiment X24 is the method of any one of embodiments X1-X23, wherein the capture probe comprises a cleavage domain, and wherein the capture probe is selectively cleaved.

[2116] Embodiment X25 is the method of any one of embodiments X1-X24, wherein one or more

capture probes comprise a cleavage domain, and wherein a capture probe is selectively cleaved.
[2117] Embodiment X26 is the method of embodiment X24 or X25, wherein the capture probe is cleaved after step (b).

[2118] Embodiment X27 is the method of embodiment X24 or X25, wherein the capture probe is cleaved before step (e).

[2119] Embodiment X28 is the method of any one of embodiments X1-X27, wherein some members of the plurality of capture probes comprise a different cleavage domain than other members of the plurality of capture probes.

[2120] Embodiment X29 is the method of embodiment X28, wherein a capture probe is cleaved after step (b).

[2121] Embodiment X30 is the method of embodiment X28 or X29, wherein a capture probe is cleaved before step (e).

[2122] Embodiment X31 is a method for profiling a biological analyte present in a biological sample, comprising: [2123] (a) contacting a biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe comprises a spatial barcode and a capture domain; [2124] (b) releasing a biological analyte from a region of interest of the biological sample, wherein the biological analyte is bound by the capture probe at a distinct spatial position of the substrate; [2125] (c) detecting the bound biological analyte; and [2126] (d) correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, [2127] thus profiling the biological analyte as having been present in the biological sample at a distinct spatial position.

[2128] Embodiment X32 is the method of embodiment X31, wherein releasing the biological analyte from the region of interest of the biological sample comprises permeabilizing the one or more cells therein.

[2129] Embodiment X33 is the method of embodiment X32, wherein the permeabilization comprises electrophoresis.

[2130] Embodiment X34 is the method of embodiment X32 or X33, wherein the permeabilization comprises administering a permeabilization reagent.

[2131] Embodiment X35 is the method of any one of embodiments X31-X34, wherein administering the permeabilization reagent comprises contacting the biological sample with a hydrogel that comprises the permeabilization reagent.

[2132] Embodiment X36 is the method of embodiment X35, wherein the region of interest is contacted with the hydrogel comprising the permeabilization reagent.

[2133] Embodiment X37 is the method of embodiment X36, wherein the area outside the region of interest of the biological sample is not contacted with the hydrogel comprising the permeabilization reagent.

[2134] Embodiment X38 is the method of any one of embodiments X31-X37, wherein the area outside the region of interest of the biological sample is removed by microdissection.

[2135] Embodiment X39 is the method of any one of embodiments X31-X38, wherein the area outside the region of interest of the biological sample is removed by microdissection.

[2136] Embodiment X40 is a method for temporal profiling the spatial location of a biological analyte present in a biological sample, comprising: [2137] (a) contacting a biological sample with a substrate comprising a plurality of capture probes, wherein a capture probe comprises a spatial barcode and a capture domain, such that at least one biological analyte is bound by a capture probe at a distinct spatial position of the substrate; [2138] (b) detecting the biological analyte; [2139] (c) correlating the released biological analyte with a spatial location in the biological sample; and [2140] (d) repeating steps (a)-(c) after a period of time.

[2141] Embodiment X41 is the method of embodiment X40, wherein the biological sample comprises a tissue, an organ, an organism, or a cell culture sample.

[2142] Embodiment X42 is the method of embodiment X41, wherein the biological sample is in a

controlled environment.

[2143] Embodiment X43 is the method of any one of embodiments X1-X42, further comprising imaging the biological sample.

[2144] Embodiment X44 is the method of embodiment X43, wherein imaging is performed prior to releasing the at least one biological analyte from the biological sample.

[2145] Embodiment X45 is the method of embodiment X43, wherein imaging is performed after releasing the at least one biological analyte from the biological sample.

[2146] Embodiment X46 is the method of any one of embodiments X43-X45, wherein imaging is used to determine a region of interest of the biological sample.

[2147] Embodiment X47 is the method of embodiment X46, wherein the imaging comprises using fiducial markers.

[2148] Embodiment X48 is the method of any one of embodiments X40-X47, wherein step (d) is repeated one, two, three, or four times.

[2149] Embodiment X49 is the method of any one of embodiments X40-X48, further comprising generating a temporal profile for the biological analyte.

[2150] Embodiment X50 is the method of any one of embodiments X40-X49, wherein the same biological analyte is detected at two or more time points.

[2151] Embodiment X51 is the method of any one of embodiments X40-X50, wherein prior to the step of repeating, the biological sample is subjected to a perturbant.

[2152] Embodiment X52 is the method of any one of embodiments X40-X51, wherein the step of releasing comprises releasing under conditions that permit cells in the biological sample to live.

[2153] Embodiment X53 is a method for determining the effect of a perturbant on a biological sample, comprising: [2154] (a) contacting the biological sample with the perturbant; and [2155] (b) performing the method for temporal profiling of embodiment 50.

[2156] Embodiment X54 is the method of embodiments X51 or X53, wherein the perturbant includes a small molecule, an antibody, a nucleic acid, a peptide, and combinations thereof.

Exemplary Embodiments Y

[2157] Embodiment Y1 is a method for temporal and spatial profiling of at least one species of biological analyte present in a biological sample, comprising: [2158] (a) providing a solid support comprising a plurality of capture probes, wherein the capture probes comprise a spatial barcode and a capture domain; [2159] (b) contacting the biological sample with the solid support comprising the capture probes; [2160] (c) releasing at least one member of the species of biological analyte from the biological sample; [2161] (d) contacting the solid support comprising the capture probes with at least one released member of the species of biological analyte such that the at least one released member of the species of biological analyte interacts with at least one of the plurality of capture probes; [2162] (e) detecting the at least one released member of the species of biological analyte; [2163] (f) correlating the at least one released member of the species of biological analyte with a spatial location in the biological sample; and [2164] (g) repeating steps (a)-(e) after a period of time.

[2165] Embodiment Y2 is the method of embodiment Y1, wherein the biological sample is in a controlled environment.

[2166] Embodiment Y3 is the method of embodiment Y1, further comprising imaging the biological sample.

[2167] Embodiment Y4 is the method of embodiment Y3, wherein imaging is performed prior to releasing at least one member of the species of biological analyte from the biological sample.

[2168] Embodiment Y5 is the method of embodiment Y3, wherein imaging is performed after releasing the at least one member of the species of biological analyte from the biological sample.

[2169] Embodiment Y6 is the method of embodiment Y3, wherein imaging is used to determine a region of interest in the biological sample.

[2170] Embodiment Y7 is the method of embodiment Y3, wherein the imaging comprises using

fiducial markers.

[2171] Embodiment Y8 is the method of embodiment Y1, wherein the step of releasing comprises applying an electric field to the biological sample.

[2172] Embodiment Y9 is the method of embodiment Y5, wherein the electric field is applied as a short burst.

[2173] Embodiment Y10 is the method of embodiment Y1, wherein the period of time is about 24 hours.

[2174] Embodiment Y11 is the method of embodiment Y1, wherein step (g) is repeated one, two, three, or four times.

[2175] Embodiment Y12 is the method of embodiment Y1, further comprising generating a temporal profile for the at least one member of the species of biological analyte.

[2176] Embodiment Y13 is the method of embodiment Y12, wherein the same species of biological analyte is detected at two or more time points.

[2177] Embodiment Y14 is the method of embodiment Y1, wherein prior to the step of repeating, the biological sample is subjected to one or more perturbants.

[2178] Embodiment Y15 is the method of embodiment Y1, wherein the step of releasing comprises releasing under conditions that permit cells in the biological sample to live.

[2179] Embodiment Y16 is a method for determining the effect of one or more perturbants on a biological sample, comprising: [2180] (h) contacting the biological sample with the one or more perturbants; and [2181] (i) performing the method for temporal and spatial profiling of embodiment Y1.

[2182] Embodiment Y17 is the method of embodiment Y16, wherein the one or more perturbants include a small molecule, an antibody, a nucleic acid, a peptide, light, heat, pressure, an electric field, and combinations thereof.

[2183] Embodiment Y18 is the method of embodiment Y16, wherein the perturbant comprises a DNA modification system.

[2184] Embodiment Y19 is the method of embodiment Y18, wherein the DNA modification system is a CRISPR system.

[2185] Embodiment Y20 is the method of embodiment Y18, wherein the DNA modification system employs a single guide RNA (sgRNA).

[2186] Embodiment Y21 is the method of embodiment Y2, wherein the controlled environment comprises a flow cell.

[2187] Embodiment Y22 is the method of embodiment Y2, wherein the controlled environment comprises a tissue culture system.

[2188] Embodiment Y23 is the method of embodiment Y2, wherein the controlled environment comprises a tissue bath.

[2189] Embodiment Y24 is the method of embodiment Y1, wherein the at least one released biological analyte is a nucleic acid, wherein the capture domain is a nucleic acid, and wherein the at least one released biological analyte hybridizes to the capture domain.

Exemplary Embodiments Z

[2190] Embodiment Z1 is a method of identifying the location of a target biological analyte within a biological sample comprising: [2191] (a) contacting the biological sample with a substrate, wherein the substrate comprises a plurality of capture probes, a capture probe comprising a spatial barcode and a capture domain; [2192] (b) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent comprises an analyte binding moiety and a capture agent barcode domain; [2193] (c) labelling a cell of the biological sample with one or more analyte capture agents by contacting the one or more analyte capture agents with the biological sample, wherein at least one analyte capture agent interacts with at least one target biological analyte within the biological sample; and [2194] (d) detecting the analyte capture agent to identify the location of the target biological analyte within the biological sample.

[2195] Embodiment Z2 is the method of embodiment Z1, wherein step (d) comprises imaging the biological sample.

[2196] Embodiment Z3 is the method of embodiments Z1 or Z2, wherein the substrate is homogenously covered by the plurality of capture probes.

[2197] Embodiment Z4 is the method of any one of embodiments Z1-Z3, wherein the capture probe capture domain comprises a poly(T) sequence.

[2198] Embodiment Z5 is the method of any one of embodiments Z1-Z3, wherein the capture probe capture domain comprises a sequence composition with at least 30% GC content.

[2199] Embodiment Z6 is the method of any one of embodiments Z1-Z3, wherein the capture probe capture domain comprises a nucleic acid sequence.

[2200] Embodiment Z7 is the method of any one of embodiments Z1-Z6, wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an analyte capture sequence.

[2201] Embodiment Z8 is the method of embodiment Z7, wherein the analyte capture sequence comprises a poly(A) 3' tail.

[2202] Embodiment Z9 is the method of embodiment Z7, wherein the analyte capture sequence comprises a sequence composition with at least 30% GC content.

[2203] Embodiment Z10 is the method of embodiment Z7, wherein the analyte capture sequence comprises a nucleic acid sequence.

[2204] Embodiment Z11 is the method of any one of embodiments Z1-Z10, wherein the plurality of analyte capture agents bind a single analyte species.

[2205] Embodiment Z12 is the method of any one of embodiments Z1-Z10, wherein two or more pluralities of analyte capture agents are contacted with the biological sample, wherein an analyte capture agent plurality binds a single analyte species.

[2206] Embodiment Z13 is a method of identifying the location of a target biological analyte within a biological sample comprising: [2207] (a) contacting the biological sample with a substrate, wherein the substrate comprises a plurality of capture probes, a capture probe comprising a spatial barcode and a capture domain; [2208] (b) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent comprises an analyte binding moiety and a capture agent barcode domain; [2209] (c) labelling a cell of the biological sample with one or more analyte capture agents by contacting the one or more analyte capture agents with the biological sample, wherein at least one analyte capture agent interacts with at least one target biological analyte within the biological sample via the analyte binding moiety; [2210] (d) allowing the plurality of analyte capture agents to interact with the plurality of capture probes on the substrate, wherein the capture probe captures the analyte capture agent via the capture domain; [2211] (e) producing a spatially-tagged analyte capture agent by extending the capture agent barcode domain of the analyte capture agent to include a sequence that is complementary to the sequence of the capture probe spatial barcode; and [2212] (f) detecting the spatially-tagged analyte capture agent to identify the location of the target biological analyte within the biological sample.

[2213] Embodiment Z14 is the method of embodiment Z13, wherein the substrate is homogenously covered by the plurality of capture probes.

[2214] Embodiment Z15 is the method of embodiment Z13 or Z14, wherein the capture probe capture domain comprises a poly(T) sequence.

[2215] Embodiment Z16 is the method of embodiment Z13 or Z14, wherein the capture probe capture domain comprises a sequence composition with at least 30% GC content.

[2216] Embodiment Z17 is the method of embodiment Z13 or Z14, wherein the capture probe capture domain comprises a nucleic acid sequence.

[2217] Embodiment Z18 is the method of any one of embodiments Z13-Z17, wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an analyte capture sequence.

[2218] Embodiment Z19 is the method of embodiment Z18, wherein the analyte capture sequence

comprises a poly(A) 3' tail.

[2219] Embodiment Z20 is the method of embodiment Z18, wherein the analyte capture sequence comprises a sequence composition with at least 30% GC content.

[2220] Embodiment Z21 is the method of embodiment Z18, wherein the analyte capture sequence comprises a nucleic acid sequence.

[2221] Embodiment Z22 is the method of any one of embodiments Z18-Z21, further comprising blocking the analyte capture sequence during steps (a)-(c) using a blocking probe.

[2222] Embodiment Z23 is the method of embodiment Z22, wherein the blocking probe comprises a poly(T) sequence.

[2223] Embodiment Z24 is the method of embodiment Z22, wherein the blocking probe comprises a sequence composition with at least 30% GC content.

[2224] Embodiment Z25 is the method of embodiment Z22, wherein the blocking probe comprises a nucleic acid sequence.

[2225] Embodiment Z26 is the method of any one of embodiments Z22-Z25, further comprising prior to step (d) releasing the blocking probe from the analyte capture sequence.

[2226] Embodiment Z27 is the method of embodiment Z26, wherein the blocking probe is released by RNase digestion.

[2227] Embodiment Z28 is the method of any one of embodiments Z13-Z27, wherein the plurality of analyte capture agents bind a single analyte species.

[2228] Embodiment Z29 is the method of any one of embodiments Z13-Z27, wherein two or more pluralities of analyte capture agents are contacted with the biological sample, wherein an analyte capture agent plurality binds a single analyte species.

[2229] Embodiment Z30 is the method of any one of embodiments Z1-Z29, wherein the target biological analyte comprises at least one of RNA, DNA, a protein, a small molecule, and a metabolite.

[2230] Embodiment Z31 is the method of any one of embodiments Z1-Z30, wherein the target biological analyte is a protein.

[2231] Embodiment Z32 is the method of embodiment Z31, wherein the target biological analyte is a ubiquitous cell surface protein.

[2232] Embodiment Z33 is the method of embodiment Z31, wherein the target biological analyte is an intracellular protein.

[2233] Embodiment Z34 is the method of any one of embodiments Z1-Z33, wherein the analyte binding moiety is an antibody.

[2234] Embodiment Z35 is the method of any one of embodiments Z1-Z34, wherein the analyte capture agent comprises an analyte binding moiety linked to the capture agent barcode domain by a cleavable domain.

[2235] Embodiment Z36 is the method of any one of embodiments Z1-Z35, further comprising permeabilizing the biological sample prior to contacting the biological sample with the plurality of analyte capture agents.

[2236] Embodiment Z37 is the method of embodiment Z36, further comprising treating the biological sample with a proteinase.

[2237] Embodiment Z38 is the method of embodiment Z37, wherein the proteinase is proteinase K.

[2238] Embodiment Z39 is the method of any one of embodiments Z13-Z38, further comprising imaging the substrate.

[2239] Embodiment Z40 is the method of embodiment Z39, further comprising performing spatial transcriptomic analysis.

[2240] Embodiment Z41 is the method of embodiment Z40, wherein spatial transcriptomic analysis comprises high-throughput sequencing.

[2241] Embodiment Z42 is the method of embodiment Z39, wherein the imaging comprises using fiducial markers.

[2242] Embodiment Z43 is the method of any one of embodiments Z1-Z42, wherein the biological sample comprises a tissue, an organ, an organism, or a cell culture sample.

[2243] Embodiment Z44 is the method of any one of embodiments Z1-Z43, wherein the biological sample is fixed prior to contacting the biological sample with the plurality of analyte capture agents.

[2244] Embodiment Z45 is the method of embodiment Z44, wherein the biological sample is fixed with formaldehyde.

[2245] Embodiment Z46 is the method of embodiment Z44, wherein the biological sample is fixed with methanol.

Exemplary Embodiments AA

[2246] Embodiment AA1 is a method of identifying the location of a target analyte in a sample comprising: [2247] (a) contacting the sample with a substrate, wherein the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; [2248] (b) contacting the sample with a plurality of analyte capture agents, wherein an analyte capture agent comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety binds specifically to the target analyte, thereby labelling the target analyte in the sample; [2249] and [2250] (c) detecting the analyte capture agent to identify the location of the target analyte in the sample.

[2251] Embodiment AA2 is the method of embodiment AA1, wherein step (c) comprises imaging the sample.

[2252] Embodiment AA3 is the method of embodiment AA1 or AA2, wherein the substrate is homogeneously covered by the plurality of capture probes.

[2253] Embodiment AA4 is the method of any one of embodiments AA1-AA3, wherein the capture domain comprises a poly(T) sequence.

[2254] Embodiment AA5 is the method of any one of embodiments AA1-AA3, wherein the capture domain comprises a nucleic acid sequence.

[2255] Embodiment AA6 is the method of embodiment AA5, wherein the nucleic acid sequence has a GC content of at least 30%.

[2256] Embodiment AA7 is the method of any one of embodiments AA1-AA6, wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an analyte capture sequence.

[2257] Embodiment AA8 is the method of embodiment AA7, wherein the analyte capture sequence comprises a poly(A) 3' tail.

[2258] Embodiment AA9 is the method of embodiment AA7, wherein the analyte capture sequence comprises a nucleic acid sequence.

[2259] Embodiment AA10 is the method of embodiment AA9, wherein the nucleic acid sequence has a GC content of at least 30%.

[2260] Embodiment AA11 is the method of any one of embodiments AA1-AA10, wherein the plurality of analyte capture agents bind a single analyte species.

[2261] Embodiment AA12 is the method of any one of embodiments AA1-AA10, wherein two or more pluralities of analyte capture agents are contacted with the sample, wherein a plurality of an analyte capture agent binds a single analyte species.

[2262] Embodiment AA13 is a method of identifying a location of a target analyte in a sample comprising: [2263] (a) contacting the sample with a substrate, wherein the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; [2264] (b) contacting the sample with a plurality of analyte capture agents, wherein an analyte capture agent comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety binds specifically to the target analyte, thereby labelling the analyte in the sample; [2265] (c) allowing the plurality of analyte capture agents to interact with the plurality of capture probes on the substrate, wherein the capture probe binds

specifically to the analyte capture agent via the capture domain; [2266] (d) producing a spatially-tagged analyte capture agent; and [2267] (e) detecting the spatially-tagged analyte capture agent to identify the location of the target analyte in the sample.

[2268] Embodiment AA14 is the method of embodiment AA13, wherein the substrate is homogenously covered by the plurality of capture probes.

[2269] Embodiment AA15 is the method of embodiment AA13 or AA14, wherein the capture domain comprises a poly(T) sequence.

[2270] Embodiment AA16 is the method of embodiment AA13 or AA14, wherein the capture domain comprises a nucleic acid sequence.

[2271] Embodiment AA17 is the method of embodiment AA16, wherein the nucleic acid sequence has a GC content of at least 30%.

[2272] Embodiment AA18 is the method of any one of embodiments AA13-AA17, wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an analyte capture sequence.

[2273] Embodiment AA19 is the method of embodiment AA18, wherein the analyte capture sequence comprises a poly(A) 3' tail.

[2274] Embodiment AA20 is the method of embodiment AA18, wherein the analyte capture sequence comprises a nucleic acid sequence.

[2275] Embodiment AA21 is the method of embodiment AA20, wherein the nucleic acid sequence has a GC content of at least 30%.

[2276] Embodiment AA22 is the method of any one of embodiments AA18-AA21, further comprising blocking the analyte capture sequence during steps (a) and (b) using a blocking probe.

[2277] Embodiment AA23 is the method of embodiment AA22, wherein the blocking probe comprises a poly(T) sequence.

[2278] Embodiment AA24 is the method of embodiment AA22, wherein the blocking probe comprises a nucleic acid sequence.

[2279] Embodiment AA25 is the method of embodiment AA24, wherein the nucleic acid sequence has a GC content of at least 30%.

[2280] Embodiment AA26 is the method of any one of embodiments AA22-AA25, further comprising prior to step (c) releasing the blocking probe from the analyte capture sequence.

[2281] Embodiment AA27 is the method of embodiment AA26, wherein the blocking probe is released by nuclease digestion.

[2282] Embodiment AA28 is the method of any one of embodiments AA13-AA27, wherein the plurality of analyte capture agents bind a single analyte species.

[2283] Embodiment AA29 is the method of any one of embodiments AA13-AA27, wherein two or more pluralities of analyte capture agents are contacted with the sample, wherein a plurality of analyte capture agents binds a single analyte species.

[2284] Embodiment AA30 is the method of any one of embodiments AA1-AA29, wherein the target analyte comprises at least one of RNA, DNA, a protein, a small molecule, and a metabolite.

[2285] Embodiment AA31 is the method of any one of embodiments AA1-AA30, wherein the target analyte is a protein.

[2286] Embodiment AA32 is the method of embodiment AA31, wherein the target analyte is a cell surface protein.

[2287] Embodiment AA33 is the method of embodiment AA31, wherein the target analyte is an intracellular protein.

[2288] Embodiment AA34 is the method of any one of embodiments AA1-AA33, wherein the analyte binding moiety is an antibody or an antigen-binding fragment thereof.

[2289] Embodiment AA35 is the method of any one of embodiments AA1-AA34, wherein the analyte capture agent comprises an analyte binding moiety linked to the capture agent barcode domain by a cleavable domain.

[2290] Embodiment AA36 is the method of any one of embodiments AA1-AA35, further comprising permeabilizing the sample prior to contacting the sample with the plurality of analyte capture agents.

[2291] Embodiment AA37 is the method of embodiment AA36, further comprising treating the sample with a proteinase.

[2292] Embodiment AA38 is the method of embodiment AA37, wherein the proteinase is proteinase K.

[2293] Embodiment AA39 is the method of any one of embodiments AA13-AA38, further comprising imaging the substrate.

[2294] Embodiment AA40 is the method of embodiment AA39, further comprising performing gene expression analysis for RNA.

[2295] Embodiment AA41 is the method of embodiment AA40, wherein the gene expression analysis comprises high-throughput sequencing.

[2296] Embodiment AA42 is the method of embodiment AA39, wherein the imaging comprises using fiducial markers.

[2297] Embodiment AA43 is the method of any one of embodiments AA1-AA42, wherein the sample comprises a tissue, a tissue section, an organ, an organism, or a cell culture sample.

[2298] Embodiment AA44 is the method of any one of embodiments AA1-AA43, wherein the sample is fixed prior to contacting the sample with the plurality of analyte capture agents.

[2299] Embodiment AA45 is the method of embodiment AA44, wherein the sample is fixed with formaldehyde.

[2300] Embodiment AA46 is the method of embodiment AA44, wherein the sample is fixed with methanol.

[2301] Embodiment AA47 is the method of embodiment AA13, wherein the step of producing a spatially-tagged analyte capture agent comprises extending the capture agent barcode domain to include a sequence that is complementary to the sequence of the spatial barcode.

[2302] Embodiment AA48 is a method for generating a spatial RNA integrity number for a location on an array, the method comprising: [2303] (a) contacting a tissue sample stained with a histology stain with an array, wherein the array includes a capture probe attached to a location on the array, and the capture probe comprises a capture domain that specifically binds to a biological analyte from the tissue sample; [2304] (b) generating a cDNA molecule from the biological analyte specifically bound to the capture domain; [2305] (c) labeling the cDNA by hybridizing a labeled oligonucleotide probe to the cDNA; [2306] (e) generating an image of the labeled cDNA and an image of the histology stain, and using the image of the labeled cDNA and the image of the histology stain to generate a spatial RNA integrity number for the location on the array.

[2307] Embodiment AA49 is the method of embodiment AA48, wherein the tissue sample comprises a tissue section, a region within a tissue, or a single cell within a tissue.

[2308] Embodiment AA50 is the method of embodiment AA48, wherein the histology stain is hematoxylin and eosin.

[2309] Embodiment AA51 is the method of embodiment AA48, wherein the capture domain comprises a poly(T) sequence.

[2310] Embodiment AA52 is the method of embodiment AA48, wherein the biological analyte is 18S rRNA.

[2311] Embodiment AA53 is the method of embodiment AA48, wherein the labeled oligonucleotide probe is fluorescently labeled.

[2312] Embodiment AA54 is the method of embodiment AA48, wherein step (c) comprises hybridizing at least four different labeled oligonucleotide probes to the cDNA.

[2313] Embodiment AA55 is the method of embodiment AA54, wherein the at least four labeled oligonucleotide probes are sequentially hybridized at different sites within the cDNA.

[2314] Embodiment AA56 is the method of embodiment AA54 or AA55, wherein the labeled

oligonucleotide probes are fluorescently labeled.

[2315] Embodiment AA57 is the method of any one of embodiments AA53-AA56, wherein the step of generating an image of the labeled cDNA comprises measuring a fluorescent signal for each of the at least four labeled oligonucleotide probes.

[2316] Embodiment AA58 is the method of embodiment AA57, wherein the fluorescent signal from each of the at least four labeled oligonucleotide probes is used to generate a spatial RNA integrity number for the location on the array.

[2317] Embodiment AA59 is a method for identifying a location of an analyte in a biological sample, the method comprising: [2318] (a) contacting the biological sample with an array, wherein the array comprises a capture probe, where the capture probe comprises a spatial barcode and a capture domain that is capable of specifically binding to the analyte; [2319] (b) contacting the biological sample with a protease, wherein the protease is capable of degrading one or more histone proteins; [2320] (c) allowing the analyte to specifically bind to the capture domain; and [2321] (d) determining (i) all or a part of the sequence of the analyte specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the analyte in the biological sample.

[2322] Embodiment AA60 is the method of claim AA59, wherein step (a) is performed before step (b).

[2323] Embodiment AA61 is the method of claim AA59, wherein step (b) is performed before step (a).

[2324] Embodiment AA62 is the method of any one of embodiments AA59-AA61, wherein the protease is capable of degrading at least one linker histone protein and at least one core histone protein in the biological sample.

[2325] Embodiment AA63 is the method of any one of embodiments AA59-AA62, wherein the protease is capable of degrading at least one histone from each core histone family in the biological sample.

[2326] Embodiment AA64 is the method of any one of claims AA59 to AA63, wherein the protease is capable of degrading at least one histone protein from every histone protein family in the biological sample.

[2327] Embodiment AA65 is the method of any one of embodiments AA59 to AA64, wherein the protease is a serine protease, an aspartyl protease, a peptidase family C1 enzyme, a protease that is inhibited by the diazomethane inhibitor Z-Phe-Phe-CHN (2) or the epoxide inhibitor E-64, a lysosomal protease or an azurophilic enzyme.

[2328] Embodiment AA66 is the method of embodiment AA65, wherein the serine protease is a trypsin enzyme, trypsin-like enzyme or a functional variant or derivative thereof.

[2329] Embodiment AA67 is the method of embodiment AA66, wherein the trypsin enzyme, trypsin-like enzyme or functional variant or derivative thereof comprises an amino acid sequence with at least 80% sequence identity to a sequence as set forth in one of SEQ ID NOs: 1-3.

[2330] Embodiment AA68 is the method of embodiment AA66 or AA67, wherein the trypsin enzyme comprises at least one of P00761, P00760, Q29463, or a mixture thereof.

[2331] Embodiment AA69 is the method of any one of embodiments AA59 to AA68, further comprising treating the biological sample with an agent or a medium that protects RNA in the biological sample from nucleases.

[2332] Embodiment AA70 is the method of embodiment AA69, wherein treating the biological sample with the medium that protects RNA in the biological sample from nucleases is completed prior to step (a).

[2333] Embodiment AA71 is the method of embodiment AA69 or AA70, wherein the medium that protects RNA in the biological sample from nucleases comprises an organic solvent.

[2334] Embodiment AA72 is the method of any one of embodiments AA69 to AA71, wherein the

medium that protects RNA in the biological sample from nucleases comprises at least about 60% v/v organic solvent.

[2335] Embodiment AA73 is the method of embodiment AA71 or AA72, wherein the organic solvent comprises at least one of ethanol, methanol, propan-2-ol, acetone, trichloroacetic acid, propanol, polyethylene glycol, acetic acid, or a combination thereof.

[2336] Embodiment AA74 is the method of any one of embodiments AA69 to AA73, wherein the medium that protects RNA in the biological sample from nucleases comprises a combination of ethanol, methanol and/or propan-2-ol.

[2337] Embodiment AA75 is the method of any one of embodiments AA59 to AA74, wherein a portion of the capture domain comprises a homopolymeric nucleotide sequence.

[2338] Embodiment AA76 is the method of embodiment AA75, wherein the homopolymeric nucleotide sequence is a poly-T sequence.

[2339] Embodiment AA77 is the method of any one of embodiments AA59 to AA76, wherein the analyte is DNA.

[2340] Embodiment AA78 is the method of embodiment AA77, wherein the DNA is genomic DNA.

[2341] Embodiment AA79 is the method of any one of embodiments AA59 to AA76, wherein the analyte is RNA.

[2342] Embodiment AA80 is the method of embodiment AA79, wherein the RNA is mRNA.

[2343] Embodiment AA81 is the method of embodiment AA59, wherein: [2344] step (a) comprises contacting the biological sample with an array comprising two or more capture probes, where each of the two or more capture probes comprises a spatial barcode and a capture domain that is capable of specifically binding to a different analyte; [2345] step (c) comprises allowing the analyte(s) to specifically bind to the capture domains; and [2346] step (d) determining (i) all or a part of the sequence of the analyte(s) specifically bound to the capture domain(s), or a complement(s) thereof, and (ii) all or a part of the sequence of the spatial barcode(s), or a complement(s) thereof, and using the determined sequences of (i) and (ii) to identify the location(s) of the analyte(s) in the biological sample.

[2347] Embodiment AA82 is the method of embodiment AA81, wherein step (d) comprises hybridizing a DNA and a RNA present in the biological sample to the capture domains.

[2348] Embodiment AA83 is the method of any one of embodiments AA59-AA82, further comprising imaging the biological sample.

[2349] Embodiment AA84 is the method of embodiment AA83, further comprising correlating the location of the analyte(s) in the biological sample with an image of the biological sample.

[2350] Embodiment AA85 is the method of any one of embodiments AA59-AA84, wherein the biological sample is a tissue section.

[2351] Embodiment AA86 is a method for identifying a location of a polypeptide in a biological sample, the method comprising: [2352] (a) contacting a biological sample with an array comprising an analyte binding moiety associated with an analyte binding moiety barcode, wherein the analyte binding moiety binds to the polypeptide in the biological sample; and [2353] (b) determining (i) all or a part of the sequence of the polypeptide bound to the analyte binding moiety and (ii) all or a portion of the sequence of the analyte binding moiety barcode, and using the determined sequences of (i) and (ii) to identify the location of the polypeptide in the biological sample.

[2354] Embodiment AA87 is the method of embodiment E86, wherein the step of determining all or a part of the sequence of the polypeptide bound to the analyte binding moiety comprises the use of Edman sequencing.

[2355] Embodiment AA88 is the method of embodiment E86 or E87, wherein: [2356] step (a) comprises contacting the biological sample with an array comprising two or more analyte binding moieties, where an analyte binding moiety is associated with an analyte binding moiety barcode, wherein the analyte binding moiety/moieties bind(s) to a polypeptide(s) in the biological sample;

and [2357] (b) determining (i) all or a part of the sequence of the polypeptide(s) bound to the analyte binding moiety/moieties and (ii) all or a portion of the sequence of the analyte binding moiety barcode(s), and using the determined sequences of (i) and (ii) to identify the location(s) of the polypeptide(s) in the biological sample.

[2358] Embodiment AA89 is a method for identifying a location of a polypeptide in a biological sample, the method comprising: [2359] (a) contacting a biological sample with an array comprising an analyte binding moiety at a location on the array, wherein the analyte binding moiety binds to the polypeptide in the biological sample; and [2360] (b) determining all or a part of the sequence of the polypeptide bound to the analyte binding moiety, and using the determined sequence to identify the location of the polypeptide in the biological sample.

[2361] Embodiment AA90 is the method of embodiment AA89, wherein the step of determining all or a part of the sequence of the polypeptide bound to the analyte binding moiety comprises the use of Edman sequencing.

[2362] Embodiment AA91 is the method of embodiment AA89 or AA90, wherein: [2363] step (a) comprises contacting the biological sample with an array comprising two or more analyte binding moieties at different locations on the array, wherein the analyte binding moiety/moieties bind(s) to a polypeptide(s) in the biological sample; and [2364] (b) determining (i) all or a part of the sequence of the polypeptide(s) bound to the analyte binding moiety/moieties and using the determined sequence(s) to identify the location(s) of the polypeptide(s) in the biological sample.

[2365] Embodiment AA92 is the method of embodiment AA86, wherein the spatial array further comprises a plurality of features, wherein a feature comprises the analyte binding moiety.

[2366] Embodiment AA93 is the method of embodiment AA92, wherein the feature further comprises a capture probe, wherein the capture probe comprises a spatial barcode and a capture domain.

[2367] Embodiment AA94 is the method of embodiment AA93, wherein the capture domain comprises a poly(T) sequence.

[2368] Embodiment AA95 is the method of embodiment AA93, wherein the analyte binding moiety barcode and spatial barcode comprise identical sequences.

[2369] Embodiment AA96 is the method of embodiment AA86 or AA89, wherein step (b) comprises: [2370] labeling one or more amino acids of the polypeptide; and [2371] detecting the labeled amino acids, thereby determining all or a part of the sequence of the polypeptide.

[2372] Embodiment AA97 is the method of embodiment AA86 or AA89, wherein step (b) comprises successive rounds of imaging and Edman sequencing.

[2373] Embodiment AA98 is a nucleic acid molecule comprising a contiguous nucleotide sequence comprising: [2374] (a) a first primer sequence; [2375] (b) a spatial barcode; [2376] (c) a unique molecular sequence; [2377] (d) a capture domain; [2378] (e) a sequence complementary to a sequence present in a nucleic acid from a biological sample; [2379] (f) a second primer sequence that is substantially complementary to a sequence of a template switching oligonucleotide.

[2380] Embodiment AA99 is the nucleic acid molecule of embodiment AA98, wherein the nucleic acid molecule is a single-stranded nucleic acid molecule.

[2381] Embodiment AA100 is the nucleic acid molecule of embodiment AA98, wherein the nucleic acid molecule is a double-stranded nucleic acid molecule.

[2382] Embodiment AA101 is the nucleic acid molecule of any one of embodiments AA98-AA100, wherein (a) through (f) are positioned in a 5' to 3' direction in the contiguous nucleotide sequence.

[2383] Embodiment AA102 is the nucleic acid molecule of embodiment AA101, wherein the nucleic acid molecule is attached to a substrate.

[2384] Embodiment AA103 is the nucleic acid molecule of embodiment AA2, wherein the substrate is a slide.

[2385] Embodiment AA104 is the nucleic acid molecule of embodiment AA102, wherein the

substrate is a bead.

[2386] Embodiment AA105 is the nucleic acid molecule of embodiment AA103, wherein the 5' end of the contiguous nucleic acid sequence is attached to the slide.

[2387] Embodiment AA106 is the nucleic acid molecule of embodiment AA104, wherein the 5' end of the contiguous nucleic acid sequence is attached to the bead.

[2388] Embodiment AA107 is the nucleic acid molecule of any one of embodiments AA98-AA106, wherein the contiguous nucleotide sequence is a chimeric RNA and DNA sequence.

[2389] Embodiment AA108 is the nucleic acid molecule of any one of embodiments AA98-AA106, wherein the contiguous nucleotide sequence is a DNA sequence.

[2390] Embodiment AA109 is a nucleic acid molecule comprising a contiguous nucleotide sequence comprising: [2391] (a) a sequence complementary to a first primer sequence; [2392] (b) a sequence complementary to a spatial barcode; [2393] (c) a sequence complementary to a unique molecular sequence; [2394] (d) a sequence complementary to a capture domain; [2395] (e) a sequence present in a nucleic acid from a biological sample; and [2396] (f) a sequence of a template switching oligonucleotide.

[2397] Embodiment AA110 is the nucleic acid molecule of embodiment AA109 wherein the nucleic acid molecule is single-stranded.

[2398] Embodiment AA111 is the nucleic acid molecule of embodiment AA109, wherein the nucleic acid molecule is double-stranded.

[2399] Embodiment AA112 is the nucleic acid molecule of any one of embodiments AA109-AA111, wherein the contiguous nucleotide sequence is a DNA sequence.

[2400] Embodiment AA113 is the nucleic acid molecule of any one of embodiments AA109-AA112, wherein (a) through (f) are positioned in a 3' to 5' direction in the contiguous nucleotide sequence.

[2401] Embodiment AA114 is a nucleic acid molecule comprising a contiguous nucleotide sequence comprising: [2402] (a) a first primer sequence; [2403] (b) a spatial barcode; [2404] (c) a unique molecular sequence; [2405] (d) a capture domain; [2406] (e) a sequence complementary to a sequence present in a nucleic acid from a biological sample; and [2407] (f) a second primer sequence.

[2408] Embodiment AA115 is the nucleic acid molecule of embodiment AA114, wherein the nucleic acid molecule is a single-stranded nucleic acid molecule.

[2409] Embodiment AA116 is the nucleic acid molecule of embodiment AA114, wherein the nucleic acid molecule is a double-stranded nucleic acid molecule.

[2410] Embodiment AA117 is the nucleic acid molecule of any one of embodiments AA114-AA116, wherein (a) through (f) are positioned in a 5' to 3' direction in the contiguous nucleotide sequence.

[2411] Embodiment AA118 is the nucleic acid molecule of any one of embodiments AA114-AA117, wherein the contiguous nucleotide sequence is a DNA sequence.

[2412] Embodiment AA119 is the nucleic acid molecule of embodiment AA117 or AA118, wherein the contiguous nucleotide sequence further comprises 3' to (f): [2413] (g) a sequence complementary to a first adaptor sequence; and [2414] (h) a sequence complementary to a third primer sequence.

[2415] Embodiment AA120 is the nucleic acid molecule of embodiment AA119, wherein the first adaptor sequence is an i7 sample index sequence.

[2416] Embodiment AA121 is the nucleic acid sequence of embodiment AA119 or AA120, wherein the third primer sequence is a P7 primer sequence.

[2417] Embodiment AA122 is the nucleic acid molecule of any one of embodiments AA119-AA121, wherein (h) is 3' positioned relative to (g) in the contiguous nucleotide sequence.

[2418] Embodiment AA123 is the nucleic acid molecule of any one of embodiments AA119-AA122, wherein the contiguous nucleotide sequence further comprises 5' to (a): [2419] (i) a second

adaptor sequence; and [2420] (ii) a fourth primer sequence.

[2421] Embodiment AA124 is the nucleic acid of embodiment AA123, wherein the second adaptor sequence is an i5 sample index sequence.

[2422] Embodiment AA125 is the nucleic acid of embodiment AA123 or AA124, wherein the fourth primer sequence is a P5 primer sequence.

[2423] Embodiment AA126 is the nucleic acid molecule of any one of embodiments AA123-AA125, wherein (ii) is 5' positioned relative to (i) in the contiguous nucleotide sequence.

[2424] Embodiment AA127 is a nucleic acid molecule comprising a contiguous nucleotide sequence comprising: [2425] (a) a sequence complementary to a first primer sequence; [2426] (b) a sequence complementary to a spatial barcode; [2427] (c) a sequence complementary to a unique molecular sequence; [2428] (d) a sequence complementary to a capture domain; [2429] (e) a sequence present in a nucleic acid from a biological sample; and [2430] (f) a sequence complementary to a second primer sequence.

[2431] Embodiment AA128 is the nucleic acid molecule of embodiment AA127, wherein the nucleic acid molecule is a single-stranded nucleic acid molecule.

[2432] Embodiment AA129 is the nucleic acid molecule of embodiment AA127, wherein the nucleic acid molecule is a double-stranded nucleic acid molecule.

[2433] Embodiment AA130 is the nucleic acid molecule of any one of embodiments AA127-AA129, wherein (a) through (f) are positioned in a 3' to 5' direction in the contiguous nucleotide sequence.

[2434] Embodiment AA131 is the nucleic acid molecule of any one of embodiments AA127-AA130, wherein the contiguous nucleotide sequence is a DNA sequence.

[2435] Embodiment AA132 is the nucleic acid molecule of embodiment AA130 or AA131, wherein the contiguous nucleotide sequence further comprises 5' to (f): [2436] (g) a first adaptor sequence; and [2437] (h) a third primer sequence.

[2438] Embodiment AA133 is the nucleic acid molecule of embodiment AA132, wherein the first adaptor sequence is an i7 sample index sequence.

[2439] Embodiment AA134 is the nucleic acid sequence of embodiment AA132 or AA133, wherein the third primer sequence is a P7 primer sequence.

[2440] Embodiment AA135 is the nucleic acid molecule of any one of embodiments AA132-AA134, wherein (h) is 5' positioned relative to (g) in the contiguous nucleotide sequence.

[2441] Embodiment AA136 is the nucleic acid molecule of any one of embodiments AA132-AA135, wherein the contiguous nucleotide sequence further comprises 3' to (a): [2442] (i) a sequence complementary to a second adaptor sequence; and [2443] (ii) a sequence complementary to a fourth primer sequence.

[2444] Embodiment AA137 is the nucleic acid of embodiment AA136, wherein the second adaptor sequence is an i5 sample index sequence.

[2445] Embodiment AA138 is the nucleic acid of embodiment AA136 or AA137, wherein the fourth primer sequence is a P5 primer sequence.

[2446] Embodiment AA139 is the nucleic acid molecule of any one of embodiments AA136-AA138, wherein (ii) is 3' positioned relative to (i) in the contiguous nucleotide sequence.

[2447] Embodiment AA140 is a method of generating a capture probe bound to substrate comprising: [2448] (a) dispensing a second solution onto a portion of a surface of a substrate coated with a first solution, wherein after the dispensing, molecules within the first solution and the second solution do not substantially mix with each other, [2449] wherein the second solution comprises a capture probe, and the dispensing results in attachment of the capture probe to the surface of the substrate.

[2450] Embodiment AA141 is the method of embodiment AA140, further comprising, after step (a): [2451] curing the second solution; and [2452] removing the first solution from the substrate, wherein the cured second solution is retained on the surface of the substrate.

[2453] Embodiment AA142 is the method of embodiment AA140 or AA141, wherein the surface of the substrate to which the capture probe is attached has an area that is smaller than a corresponding area generated by dispensing the second solution onto the surface of the substrate in the absence of the first solution.

[2454] Embodiment AA143 is the method of embodiment AA140 or AA141, wherein the surface of the substrate to which the capture probe is attached has an area that is about two-fold less than the corresponding area generated by dispensing the second solution onto the surface of the substrate in the absence of the first solution.

[2455] Embodiment AA144 is the method of embodiment AA140, further comprising: [2456] curing the first and second solutions on the substrate, thus generating a matrix.

[2457] Embodiment AA145 is the method of embodiment AA144, wherein the first and second solutions are cured by chemical means.

[2458] Embodiment AA146 is the method of embodiment AA144, wherein the first and second solutions are cured by photo-reactive means.

[2459] Embodiment AA147 is the method of any one of embodiments AA144-AA146, further comprising: [2460] shrinking the matrix, thus generating a shrunken matrix having an area to which the capture probe is attached, wherein the area in the shrunken matrix is reduced as compared to a corresponding area in the matrix.

[2461] Embodiment AA148 is the method of embodiment AA140, wherein the method further comprises, prior to the step of dispensing the second solution: [2462] curing the first solution to form a first solution matrix, wherein the first solution matrix can be reversibly expanded along one or more of its axes.

[2463] Embodiment AA149 is the method of embodiment AA148, wherein the step of dispensing the second solution onto the portion of the substrate comprising dispensing the second solution onto a surface of the first solution matrix in an expanded state.

[2464] Embodiment AA150 is the method of embodiment AA149, wherein the method further comprises, after the dispensing step: [2465] shrinking the first solution matrix along the one or more axes of the surface of the substrate.

[2466] Embodiment AA151 is the method of any one of embodiments AA140-AA150, wherein the first solution has a viscosity that is about 0.1 $\times$ -fold, 0.2 $\times$ -fold, 0.3 $\times$ -fold, 0.4 $\times$ -fold, 0.5 $\times$ -fold, 0.6 $\times$ -fold, 0.7 $\times$ -fold, 0.8 $\times$ -fold, 0.9 $\times$ -fold, 1.0 $\times$ -fold, 1.1 $\times$ -fold, 1.2 $\times$ -fold, 1.3 $\times$ -fold, 1.4-fold, 1.5 $\times$ -fold, 1.6 $\times$ -fold, 1.7 $\times$ -fold, 1.8 $\times$ -fold, 1.9 $\times$ -fold, 2.0 $\times$ -fold, 2.5 $\times$ -fold, 3.0 $\times$ -fold, 4.0 $\times$ -fold, 5.0 $\times$ -fold, 6.0 $\times$ -fold, 7.0 $\times$ -fold, 8.0 $\times$ fold, 9.0 $\times$ -fold or 10 $\times$ -fold greater than the viscosity of the second solution.

[2467] Embodiment AA152 is the method of any one of embodiments AA140-AA151, wherein the first solution has a lower nucleic acid diffusivity as compared to the second solution.

[2468] Embodiment AA153 is the method of any one of embodiments AA140-AA152, wherein the capture probe comprises a spatial barcode.

[2469] Embodiment AA154 is the method of any one of embodiments AA140-AA153, wherein the capture probe comprises a unique molecular identifier.

[2470] Embodiment AA155 is the method of any one of embodiments AA140-AA154, wherein the capture probe comprises a functional domain.

[2471] Embodiment AA156 is the method of any one of embodiments AA140-AA155, wherein the capture probe comprises a cleavage domain, a spatial barcode, a unique molecular identifier, a functional domain, and a capture domain.

[2472] Embodiment AA157 is a method of generating an array of capture probes bound to a substrate comprising: [2473] (a) dispensing a droplet of a second solution onto a portion of a surface of a substrate coated with a first solution, wherein after the dispensing, molecules within the first solution and the droplet of the second solution do not substantially mix with each other, [2474] wherein the second solution comprises a capture probe, and the dispensing results in

attachment of the capture probe to the surface of the substrate.

[2475] Embodiment AA158 is the method of embodiment AA157, further comprising, after step (a): [2476] curing the second solution; and [2477] removing the first solution from the substrate, wherein the cured second solution is retained on the surface of the substrate.

[2478] Embodiment AA159 is the method of embodiment AA157 or AA158, wherein the surface of the substrate to which the capture probe is attached has an area that is smaller than a corresponding area generated by dispensing the second solution onto the surface of the substrate in the absence of the first solution.

[2479] Embodiment AA160 is the method of embodiment AA159, wherein the surface of the substrate to which the capture probe is attached has an area that is about two-fold less than the corresponding area generated by dispensing the second solution onto the surface of the substrate in the absence of the first solution.

[2480] Embodiment AA161 is the method of any one of embodiments AA157, further comprising: [2481] curing the first and second solutions on the substrate, thus generating a matrix.

[2482] Embodiment AA162 is the method of embodiment AA161, wherein the first solution and the second solutions are cured by chemical means.

[2483] Embodiment AA163 is the method of embodiment AA161, wherein the first and second solutions are cured by photo-reactive means.

[2484] Embodiment AA164 is the method of any one of embodiments AA161-AA163, further comprising: [2485] shrinking the matrix, thus generating a shrunken matrix having an area to which the capture probe is attached, wherein the area in the shrunken matrix is reduced as compared to a corresponding area in the matrix.

[2486] Embodiment AA165 is the method of any one of embodiments AA157-AA164, wherein the method further comprises, prior to the step of dispensing the second solution: curing the first solution to form a first solution matrix, wherein the first solution matrix can be reversibly expanded along one or more of its axes.

[2487] Embodiment AA166 is the method of embodiment AA165, wherein the step of dispensing the second solution onto the portion of the substrate comprising dispensing the second solution onto a surface of the first solution matrix in an expanded state.

[2488] Embodiment AA167 is the method of embodiment AA166, wherein the method further comprises, after the dispensing step: [2489] shrinking the first solution matrix along the one or more axes of the surface of the substrate.

[2490] Embodiment AA168 is the method of any one of embodiments AA157-AA167, wherein the first solution has a viscosity that is about 0.1×-fold, 0.2×-fold, 0.3×-fold, 0.4×-fold, 0.5×-fold, 0.6×-fold, 0.7×-fold, 0.8×-fold, 0.9×-fold, 1.0×-fold, 1.1×-fold, 1.2×-fold, 1.3×-fold, 1.4-fold, 1.5×-fold, 1.6×-fold, 1.7×-fold, 1.8×-fold, 1.9×-fold, 2.0×-fold, 2.5×-fold, 3.0×-fold, 4.0×-fold, 5.0× fold, 6.0×-fold, 7.0×-fold, 8.0× fold, 9.0×-fold or 10×-fold greater than the viscosity of the second solution.

[2491] Embodiment AA169 is the method of any one of embodiments AA157-AA168, wherein the first solution has a lower nucleic acid diffusivity as compared to the second solution.

[2492] Embodiment AA170 is the method of any one of embodiments AA157-AA169, wherein the capture probe comprises a spatial barcode.

[2493] Embodiment AA171 is the method of any one of embodiments AA157-AA170, wherein the at least one of the capture probe comprises a unique molecular identifier.

[2494] Embodiment AA172 is the method of any one of embodiments AA157-AA171, wherein the capture probe comprises a functional domain.

[2495] Embodiment AA173 is the method of any one of embodiments AA157-AA172, wherein the capture probe comprises a cleavage domain, a spatial barcode, a unique molecular identifier, a functional domain, and a capture domain.

[2496] Embodiment AA174 is a method for spatially profiling an analyte in a biological sample,

comprising: [2497] (a) dispensing a droplet of a second solution onto a portion of a surface of a substrate coated with a the first solution, wherein after the dispensing, molecules within the first solution and the droplet of the second solution do not substantially mix with each other, wherein the second solution comprises a capture probe, and the dispensing results in attachment of the capture probe to the surface of the substrate; [2498] (b) contacting the biological sample with the spatial array such that the analyte present in the biological sample is captured by the capture probes; [2499] thereby determining the spatial profile of the captured analyte.

[2500] Embodiment AA175 is the method of embodiment AA174, wherein the at least one of the capture probe comprises a cleavage domain, a spatial barcode, a unique molecular identifier, a functional domain, and a capture domain.

[2501] Embodiment AA176 is a method of determining process bias in a spatial analysis workflow comprising: [2502] (a) providing a substrate comprising one or more test analytes, wherein the one or more test analytes are disposed on the substrate in a known amount at a known location; [2503] (b) contacting the substrate with an array comprising one or more capture probes under conditions that allow the one or more capture probes to interact with the one or more test analytes, wherein a capture probe comprises a spatial barcode and a capture domain; [2504] (c) detecting the one or more test analytes that interact with the one or more capture probes; and [2505] (d) determining whether the spatial analysis workflow accurately detects the presence, amount, location, or combinations thereof, of the one or more test analytes based on the detecting in step (c), thereby determining process bias in the spatial analysis workflow.

[2506] Embodiment AA177 is the method of embodiment AA176, wherein the substrate comprises a semi-porous material.

[2507] Embodiment AA178 is the method of embodiment AA177, wherein the semi-porous material comprises at least one of a nitrocellulose membrane, a hydrogel, a nylon filter, or combinations thereof.

[2508] Embodiment AA179 is the method of embodiment AA176, wherein the one or more test analytes comprises at least one of nucleic acid, a protein, a lipid, or combinations thereof.

Embodiment AA180 is the method of embodiment AA179, wherein the one or more test analytes is RNA.

[2509] Embodiment AA181 is the method of embodiment AA176, wherein the one or more test analytes are disposed on the substrate in a defined pattern.

[2510] Embodiment AA182 is the method of embodiment AA181, wherein the defined pattern comprises one or more spots.

[2511] Embodiment AA183 is a method for determining a location of an analyte in a biological sample, the method comprising: [2512] (a) contacting the biological sample with an array, wherein the array comprises a capture probe, wherein the capture probe comprises a spatial barcode and a capture domain that binds specifically to the analyte; [2513] (b) generating an amplification product from the captured analyte using isothermal amplification; and [2514] (c) determining (i) the sequence of all or a portion of the analyte, or a complement thereof, using the amplification product, and (ii) the sequence of all or a portion of the spatial barcode, or a complement thereof, using the amplification product, and using the determined sequences of (i) and (ii) to identify the location of the analyte in the biological sample.

[2515] Embodiment AA184 is the method of embodiment AA183, wherein: [2516] step (a) comprises contacting the biological sample with an array that comprises two or more capture probes, wherein a capture probe comprises a spatial barcode and a capture domain that binds specifically to an analyte; [2517] step (b) comprises generating amplification product(s) from the captured analyte(s) using isothermal amplification; and [2518] step (c) comprises determining (i) the sequence of all or a portion of the analyte(s), or a complement thereof, using the amplification product(s), and (ii) the sequence of all or a portion of the spatial barcode(s), or a complement thereof, using the amplification product(s), and using the determined sequences of (i) and (ii) to

identify the location(s) of the analyte(s) in the biological sample.

[2519] Embodiment AA185 is the method of embodiment AA183 or AA184, wherein the array comprises one or more features.

[2520] Embodiment AA186 is the method of embodiment AA185, wherein the one or more features comprises a bead.

[2521] Embodiment AA187 is the method of any one of embodiments AA183-AA186, wherein the isothermal amplification comprises the use of a helicase.

[2522] Embodiment AA188 is the method of any one of embodiments AA183-AA186, wherein the isothermal amplification comprises the use of a recombinase.

[2523] Embodiment AA189 is the method of any of embodiments AA183-AA186, wherein the isothermal amplification comprises the use of a template switching oligonucleotide.

[2524] Embodiment AA190 is the method of any one of embodiments AA183-AA189, wherein the isothermal amplification is performed on the array.

[2525] Embodiment AA191 is the method of any one of embodiments AA183-AA190, further comprising blocking the 3' end of a capture probe(s) not bound to an analyte after step (a).

[2526] Embodiment AA192 is a method for barcoding a biological sample, the method comprising:

[2527] contacting a lipophilic molecule conjugated to an oligonucleotide comprising a spatial barcode, to the biological sample, wherein the lipophilic molecule inserts into a lipid membrane in the biological sample, thereby barcoding the biological sample.

[2528] Embodiment AA193 is the method of embodiment AA192, wherein the lipophilic molecule comprises cholesterol, tocopherol, steryl, palmitate, lignoceric acid, derivatives thereof, and combinations thereof.

[2529] Embodiment AA194 is the method of embodiment AA192, wherein the lipophilic molecule conjugated to the oligonucleotide inserts into the plasma membrane.

[2530] Embodiment AA195 is the method of embodiment AA192, wherein the lipophilic molecule conjugated to the oligonucleotide inserts into the nuclear membrane.

[2531] Embodiment AA196 is the method of embodiment AA192, wherein the lipophilic molecule is conjugated to the oligonucleotide via a linker.

[2532] Embodiment AA197 is the method of embodiment AA192, wherein the barcoded oligonucleotide further comprises a functional domain, a first UMI, a second UMI, a spatial barcode, or combinations thereof.

[2533] Embodiment AA198 is a method of barcoding a biological sample, the method comprising:

[2534] contacting the biological sample with an anchor oligonucleotide and a co-anchor oligonucleotide, wherein the anchor oligonucleotide and the co-anchor oligonucleotide are at least partially inserted into a lipid membrane of the biological sample insert into a lipid membrane; and

[2535] delivering a barcoded oligonucleotide to the biological sample, wherein the barcoded oligonucleotide comprises a sequence that is substantially complementary to a sequence within the anchor oligonucleotide, thereby barcoding the biological sample.

[2536] Embodiment AA199 is the method of embodiment AA198, wherein the anchor oligonucleotide comprises a lipid molecule conjugated to its 3' end.

[2537] Embodiment AA200 is the method of embodiment AA198, wherein the co-anchor oligonucleotide comprises a lipid molecule conjugated to its 5' end.

[2538] Embodiment AA201 is the method of embodiment AA198, wherein the anchor oligonucleotide further comprises a constant sequence.

[2539] Embodiment AA202 is the method of embodiment AA198, wherein the co-anchor oligonucleotide comprises a sequence that is substantially complementary to the constant sequence of the anchor oligonucleotide.

[2540] Embodiment AA203 is the method of embodiments AA201-AA202, wherein the constant sequence and the sequence substantially complementary to the constant sequence hybridize.

[2541] Embodiment AA204 is the method of embodiment AA198, wherein the barcoded

oligonucleotide further comprises a functional domain, a first UMI, a second UMI, a spatial barcode, or combinations thereof.

[2542] Embodiment AA205 is a method for barcoding a cell of a biological sample, the method comprising: [2543] contacting the biological sample with a cell tagging agent conjugated to an oligonucleotide comprising a barcode, thereby barcoding a cell of the biological sample.

[2544] Embodiment AA206 is the method of embodiment AA205, wherein the barcoded oligonucleotide further comprises a functional domain, a first UMI, a second UMI, or combinations thereof.

[2545] Embodiment AA207 is the method of embodiment AA205, wherein the cell tagging agent comprises a lipid molecule.

[2546] Embodiment AA208 is the method of embodiment AA205, wherein the cell tagging agent comprises a dye.

[2547] Embodiment AA209 is the method of embodiment AA205, wherein the cell tagging agent comprises cell penetrating peptide.

[2548] Embodiment AA210 is the method of embodiment AA205, wherein the cell tagging agent comprises an antibody.

[2549] Embodiment AA211 is the method of embodiment AA205, wherein the cell tagging agent comprises a gold nanoparticle.

[2550] Embodiment AA212 is a method for barcoding a cell of a biological sample, the method comprising: [2551] contacting the cell of the biological sample with a cell tagging agent conjugated to an oligonucleotide comprising a barcode, thereby barcoding the cell of the biological sample.

[2552] Embodiment AA213 is the method of embodiment AA212, wherein the cell tagging agent comprises a bifunctional NHS linker.

[2553] Embodiment AA214 is the method of embodiment AA212, wherein the cell tagging agent comprises an azide group.

[2554] Embodiment AA215 is the method of embodiment AA212, wherein the cell tagging agent comprises a glycol chitosan cholesterol oligonucleotide complex.

[2555] Embodiment AA216 is the method of embodiment AA212, wherein the cell tagging agent comprises amine reactive DSPE-PEG (2000) cyanuric chloride.

[2556] Embodiment AA217 is the method of embodiment AA212, wherein the cell penetrating agent is a cell penetrating peptide.

[2557] Embodiment AA218 is the method of any one of embodiments AA212-AA217, wherein the oligonucleotide further comprises a functional domain, a first UMI, a second UMI, or combinations thereof.

[2558] Embodiment AA219 is the method of any one of embodiments AA212-AA218, wherein the biological sample is a tissue section.

Exemplary Embodiments BB

[2559] Embodiment BB1 is a method of identifying the location of a target analyte in a biological sample comprising: [2560] (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: [2561] an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the target analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence; and the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain that binds specifically to the analyte capture sequence; and [2562] (b) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the target analyte in the biological sample.

[2563] Embodiment BB2 is the method of embodiment BB1, wherein the method further comprises facilitating movement or migration of the analyte capture agent to the substrate.

[2564] Embodiment BB3 is the method of embodiment BB1 or BB2, wherein the method further comprises imaging the biological sample.

[2565] Embodiment BB4 is the method of any one of embodiments BB1 to BB3, wherein the substrate is an array.

[2566] Embodiment BB5 is the method of embodiment BB4, wherein the array comprises a slide.

[2567] Embodiment BB6 is the method of embodiment BB4, wherein the array is a bead array.

[2568] Embodiment BB7 is the method of any one of embodiments BB1-BB6, wherein the capture domain comprises a nucleic acid sequence.

[2569] Embodiment BB8 is the method of embodiment BB7, wherein the nucleic acid sequence of the capture domain comprises a poly(T) sequence.

[2570] Embodiment BB9 is the method of embodiment BB7, wherein the nucleic acid sequence of the capture domain has a GC content of at least 30%.

[2571] Embodiment BB 10 is the method of embodiment BB7, wherein the analyte capture sequence comprises a nucleic acid sequence.

[2572] Embodiment BB11 is the method of embodiment BB10, wherein the nucleic acid sequence of the analyte capture sequence comprises a poly(A) sequence.

[2573] Embodiment BB 12 is the method of embodiment BB10, wherein the nucleic acid sequence of the analyte capture sequence has a GC content of at least 30%.

[2574] Embodiment BB 13 is the method of any one of embodiments BB1-BB12, wherein the analyte capture agent comprises a label.

[2575] Embodiment BB14 is the method of embodiment BB13, wherein the label is a fluorophore.

[2576] Embodiment BB 15 is the method of any one of embodiments BB1-BB14, wherein the target analyte is a protein.

[2577] Embodiment BB 16 is the method of embodiment BB15, wherein the protein is an intracellular protein.

[2578] Embodiment BB 17 is the method of embodiment BB15, wherein the protein is an extracellular protein.

[2579] Embodiment BB18 is the method of embodiment BB15, wherein the protein is a cell surface protein.

[2580] Embodiment BB 19 is the method of any one of embodiments BB15-BB18, wherein the analyte binding moiety comprises an antibody or an antigen-binding domain thereof.

[2581] Embodiment BB20 is the method of any one of embodiments BB14-BB19, wherein the method further comprises detecting the label of the analyte capture agent before the analyte capture agent specifically binds to the capture domain.

[2582] Embodiment BB21 is the method of any one of embodiments BB14-BB20, wherein the method further comprises detecting the label of the analyte capture agent after the analyte capture agent specifically binds to the capture domain.

[2583] Embodiment BB22 is the method of any one of embodiments BB1-BB21, wherein the method further comprises performing immunofluorescence microscopy on the biological sample to detect the location of an additional target analyte in the biological sample.

[2584] Embodiment BB23 is the method of embodiment BB22, wherein the additional target analyte is a protein.

[2585] Embodiment BB24 is the method of embodiment BB22, wherein the protein is an intracellular protein.

[2586] Embodiment BB25 is the method of embodiment BB22, wherein the additional target analyte is a cell surface protein.

[2587] Embodiment BB26 is the method of any one of embodiments BB22-BB25, wherein the method further comprises generating an image of the location of the target analyte in the biological sample, and generating an image of the location of the additional target analyte in the biological sample.

[2588] Embodiment BB27 is the method of embodiment BB26, wherein the method further comprises comparing or overlying the image of the location of the target analyte in the biological sample and the image of the location of the additional target analyte in the biological sample.

[2589] Embodiment BB28 is the method of any one of embodiments BB1-BB21, wherein the target analyte is a protein and the method further comprises performing immunofluorescence microscopy on the biological sample to detect the location of an additional target analyte in the biological sample, wherein the additional target analyte is a nucleic acid.

[2590] Embodiment BB29 is the method of embodiment BB28, wherein the additional target analyte is a nucleic acid encoding the target analyte.

[2591] Embodiment BB30 is the method of embodiment BB29, wherein the method further comprises generating an image of the location of the target analyte and an image of the location of the additional target analyte in the biological sample.

[2592] Embodiment BB31 is the method of embodiment BB30, wherein the method further comprises comparing or overlying the image of the location of the target analyte and the image of the location of the additional target analyte in the biological sample.

[2593] Embodiment BB32 is a method of identifying a location of a target analyte in a biological sample comprising: [2594] (a) contacting a plurality of analyte capture agents to a biological sample disposed on a substrate, wherein: [2595] an analyte capture agent of the plurality of the analyte capture agents comprises (i) an analyte binding moiety that binds specifically to the target analyte, (ii) an analyte binding moiety barcode; and (iii) an analyte capture sequence, wherein the analyte capture sequence is hybridized to a blocking probe; and [2596] the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality comprises a spatial barcode and a capture domain; [2597] (b) releasing the blocking probe from the analyte capture sequence, thereby allowing the analyte capture sequence to bind specifically to the capture agent; and [2598] (c) determining the sequence of (i) all or a portion of the spatial barcode or a complement thereof, and (ii) all or a portion of the analyte binding moiety barcode, or a complement thereof, and using the determined sequences of (i) and (ii) to identify the location of the target analyte in the biological sample.

[2599] Embodiment BB33 is the method of embodiment BB32, wherein step (b) comprises heating the analyte capture agent to release the blocking probe.

[2600] Embodiment BB34 is the method of embodiment BB32, wherein step (b) comprises contacting the analyte capture agent with RNase H.

[2601] Embodiment BB35 is the method of any one of embodiments BB32-BB34, wherein the method further comprises imaging the biological sample.

[2602] Embodiment BB36 is the method of embodiments BB32-BB35, wherein the substrate is an array.

[2603] Embodiment BB37 is the method of embodiment BB36, wherein the array comprises a slide.

[2604] Embodiment BB38 is the method of embodiment BB36, wherein the array is a bead array.

[2605] Embodiment BB39 is the method of any one of embodiments BB32-BB38, wherein the capture domain comprises a nucleic acid sequence.

[2606] Embodiment BB40 is the method of embodiment BB39, wherein the nucleic acid sequence of the capture domain comprises a poly(T) sequence.

[2607] Embodiment BB41 is the method of embodiment BB39, wherein the nucleic acid sequence of the capture domain has a GC content of at least 30%.

[2608] Embodiment BB42 is the method of embodiment BB39, wherein the analyte capture sequence comprises a nucleic acid sequence.

[2609] Embodiment BB43 is the method of embodiment BB42, wherein the nucleic acid sequence of the analyte capture sequence comprises a poly(A) sequence.

[2610] Embodiment BB44 is the method of embodiment BB42, wherein the nucleic acid sequence

of the analyte capture sequence has a GC content of at least 30%.

[2611] Embodiment BB45 is the method of any one of embodiments BB32-BB44, wherein the analyte capture agent comprises a label.

[2612] Embodiment BB46 is the method of embodiment BB45, wherein the detectable label is a fluorophore.

[2613] Embodiment BB47 is the method of any one of embodiments BB32-BB46, wherein the target analyte is a protein.

[2614] Embodiment BB48 is the method of embodiment BB45, wherein the protein is an intracellular protein.

[2615] Embodiment BB49 is the method of embodiment BB45, wherein the protein is an extracellular protein.

[2616] Embodiment BB50 is the method of embodiment BB45, wherein the protein is a cell surface protein.

[2617] Embodiment BB51 is the method of any one of embodiments BB45-BB50, wherein the analyte binding moiety comprises an antibody or an antigen-binding domain thereof.

[2618] Embodiment BB52 is the method of any one of embodiments BB46-BB51, wherein the method further comprises, between steps (a) and (b), detecting the label of the analyte capture agent.

[2619] Embodiment BB53 is the method of any one of embodiments BB46-BB52, wherein the method further comprises, after step (b), detecting the label of the analyte capture agent.

[2620] Embodiment BB54 is the method of embodiment BB52 or BB53, wherein the label is a fluorophore.

[2621] Embodiment BB55 is the method of any one of embodiments BB32-BB54, wherein the method further comprises performing immunofluorescence microscopy on the biological sample to detect the location of an additional target analyte in the biological sample.

[2622] Embodiment BB56 is the method of embodiment BB55, wherein the additional target analyte is a protein.

[2623] Embodiment BB57 is the method of embodiment BB55, wherein the protein is an intracellular protein.

[2624] Embodiment BB58 is the method of embodiment BB55, wherein the additional target analyte is a cell surface protein.

[2625] Embodiment BB59 is the method of any one of embodiments BB32-BB58, wherein the method further comprises generating an image of the location of the target analyte in the biological sample, and generating an image of the location of the additional target analyte in the biological sample.

[2626] Embodiment BB60 is the method of embodiment BB59, wherein the method further comprises comparing or overlying the image of the location of the target analyte in the biological sample and the image of the location of the additional target analyte in the biological sample.

[2627] Embodiment BB61 is the method of any one of embodiments BB32-BB54, wherein the target analyte is a protein and the method further comprises performing immunofluorescence microscopy on the biological sample to detect the location of an additional target analyte in the biological sample, wherein the additional target analyte is a nucleic acid.

[2628] Embodiment BB62 is the method of embodiment BB61, wherein the additional target analyte is a nucleic acid encoding the target analyte.

[2629] Embodiment BB63 is the method of embodiment BB62, wherein the method further comprises generating an image of the location of the target analyte and an image of the location of the additional target analyte in the biological sample.

[2630] Embodiment BB64 is the method of embodiment BB63, wherein the method further comprises comparing or overlying the image of the location of the target analyte and the image of the location of the additional target analyte in the biological sample.

[2631] Embodiment BB65 is the method of any one of embodiments BB1-BB64, wherein the analyte capture agent comprises an analyte binding moiety linked to the capture agent barcode domain by a cleavable domain.

[2632] Embodiment BB66 is the method of any one of embodiments BB1-BB65, wherein the method further comprises, before step (a), permeabilizing the biological sample.

[2633] Embodiment BB67 is the method of any one of embodiments BB1-BB66, further comprising imaging the substrate.

[2634] Embodiment BB68 is the method of any one of embodiments BB1-BB67, further comprising performing gene expression analysis for RNA.

[2635] Embodiment BB69 is the method of embodiment BB68, wherein the gene expression analysis comprises high-throughput sequencing.

[2636] Embodiment BB70 is the method of embodiment BB67, wherein the imaging comprises the use of fiducial markers.

[2637] Embodiment BB71 is the method of any one of embodiments BB1-BB70, wherein the biological sample comprises a tissue, a tissue section, an organ, an organism, an organoid, or a cell culture sample.

[2638] Embodiment BB72 is the method of any one of embodiments BB1-BB71, wherein the biological sample is fixed prior to step (a).

[2639] Embodiment BB73 is the method of embodiment BB72, wherein the biological sample is fixed with formaldehyde.

[2640] Embodiment BB74 is the method of embodiment BB72, wherein the biological sample is fixed with methanol.

EXAMPLES

Example 1. Method for Profiling a Biological Analyte in a Plurality of Cells, an Organism, or an Organoid

[2641] A biological sample (e.g., a plurality of cells, an organism, or an organoid) is disposed upon a solid support comprising a plurality of capture probes. See FIG. 25. The capture probes of the solid support individually include a molecular barcode and a capture domain. A distinct spatial position on the solid support comprises capture probes that are distinct from other capture probes of the plurality of capture probes. The cells of the biological sample are lysed such that biological analytes from the cells are released and bind to a capture probe at the distinct spatial position of the solid support. The capture probes/analytes are analyzed, and the biological analyte with a molecular barcode of the capture probe is correlated with the distinct spatial position of the solid support, thus profiling the biological analyte as having been cell.

Example 2. Method for Profiling a Biological Analyte in a Plurality of Cells Using a Substrate Comprising Capture Antibodies

[2642] A plurality of cells is distributed onto a substrate comprising features with a plurality of capture probes and a plurality of capture antibodies. The capture probes of the solid support individually comprise a molecular barcode and a capture domain. A distinct spatial position on the solid support comprises capture probes that are distinct from other capture probes of the plurality of capture probes. Cells of the plurality of cells bind to capture antibodies on the substrate. The cells are lysed such that biological analytes from the cells are released and bind to a capture probe at the distinct spatial position of the solid support. The capture probes/analytes are analyzed, and the biological analyte with a molecular barcode of the capture probe is correlated with the distinct spatial position of the solid support, thus profiling the biological analyte as having been present in the cell.

Example 3. Method for Profiling a Biological Analyte in a Migrated Cell

[2643] A plurality of cells is allowed to migrate on a substrate having a plurality of capture probes and having channels that confine cell migration to one dimension along the substrate. The capture probes of the solid support individually comprise a spatial barcode and a capture domain. A distinct

spatial position on the solid support comprises capture probes that are distinct from other capture probes of the plurality of capture probes. The cells are lysed such that biological analytes from the cells are released and bind to a capture probe at the distinct spatial position of the solid support. The capture probes/analytes are analyzed, and the biological analyte with a molecular barcode of the capture probe is correlated with the distinct spatial position of the solid support, thus profiling the biological analyte as having been present in the cell.

Example 4. Method for Profiling a Biological Analyte in Subcellular Fractions

[2644] A biological sample is placed on a slide, and the slide is contacted with an array as described herein. A lysis solution that selectively permeabilizes the plasma membrane is added to the biological sample. Target biological analytes in the cytosol are released from the biological sample and diffuse towards the array and are bound by capture probes. The array is removed, and reverse transcription/barcoding can be performed on the array. Additionally, the capture probes are collected or second strand synthesis is performed for preparation of a library. The biological sample is then contacted with a second array and a nucleus-specific lysis solution is added to the biological sample. Target biological analytes from the nuclei of cells in the biological sample are released and diffuse towards the second array and are bound by capture probes. The second array is removed, and reverse transcription/barcoding can be performed on the second array. Additionally, the capture probes from the second array are collected or second strand synthesis is performed for preparation of a library.

Example 5. Temporal Analysis of an Analyte in a Biological Sample

[2645] Live cells tissue sections are placed in a controlled environment permissive for cellular growth and/or maintenance, and/or to prevent hypoxia. Barcoded arrays are placed on top of the tissues and imaged using a microscope or other suitable instrument to register the relative position of the tissue to the barcoded array, optionally using optically encoded fiducial markers. An electric field is applied for a period of time between the live biological sample and the barcoded array, such that biological analytes (e.g., DNA, RNA, proteins, metabolites, small molecules, lipids, and the like) are released and captured onto the spatially barcoded array, preserving their spatial information. The array is removed, and the molecular information therein is determined by performing library construction for next generation sequencing. Sequencing is followed by computational analysis to correlate the expression value with spatial barcode. These steps are repeated one or more times to capture the analytes' spatial information over various time-points.

Example 6—Methods for Spatially Detecting One or More Proteins from a Biological Sample

[2646] In a non-limiting example, antibody staining is performed on a capture probe array. FIG. 36 shows an exemplary illustration of fluorescently labeled antibodies interacting with analytes of a biological sample on a capture probe array. For example, a method of antibody staining can include: (a) providing a capture probe array, as described herein; (b) contacting a tissue section (e.g., mouse spleen tissue) with the array and drying the sectioned slides for 1 minute at 37° C.; (c) fixing the tissue section with either 2% formaldehyde at room temperature or with methanol at -20° C. for 10 minutes; (d) rehydrating, blocking, and permeabilizing the tissue sample with a blocking buffer (3×SSC, 2% BSA, 0.1% Triton X, and 1U/μl RNase inhibitor) for 10 minutes at 4° C.; (e) staining the tissue section with fluorescent primary antibodies, for example, the primary antibody is diluted 1:100 in an antibody dilution buffer (3×SSC, 2% BSA, 0.1% Triton X, 1 U/μl RNase inhibitor) for 30 minutes at 4° C.; (f) removing the antibody by washing the sample 4 times in 3×SSC, 2% BSA, 0.1% Triton X, and 1U/μl RNase inhibitor; (g) mounting and cover slipping the tissue sample in glycerol and 1U/μl RNase inhibitor; (h) imaging the tissue section to detect target proteins within the tissue; (i) removing the coverslip and washing the tissue section in 3×SSC; and (j) performing any of the spatial workflows described herein. The steps of this method are depicted in FIG. 37. Performing any of the spatial workflows described herein can include determining the sequence of all or a portion of the spatial barcode or a complement thereof, and all or a portion of the analyte binding moiety barcode, or complement thereof. In some embodiments

of the example, the analyte capture agent binds to the capture domain before detecting the label associated with the analyte capture agent. In some embodiments of the example, the analyte capture agent binds to the capture domain after detecting the label associated with the analyte capture agent. [2647] The fluorescence level across the tissue can be imaged to produce an image wherein the image shows the fluorescently labeled antibodies specifically bind to the target proteins allowing for the examination of distribution of the target proteins throughout the tissue.

[2648] FIGS. 38 and 42 show exemplary staining using the method depicted above and in FIG. 37, wherein the antibody staining is performed on a capture probe array. FIG. 38 shows tissue sections of mouse spleen fixed in either 2% formaldehyde or 100% methanol. Mouse spleen contains red pulp containing stromal tissue, blood vessels, and macrophages, surrounding pockets of white pulp containing mostly immune cells, such as T cells and B cells. In both the 2% formaldehyde and 100% methanol fixation conditions, CD3, a cell surface marker for T cells, can be seen in the pockets of white pulp, and CD29, a cell surface marker for stromal cells, can be seen in the red pulp portion of the spleen (FIG. 38). FIG. 42 shows coronal sections of mouse brain fixed in either 2% formaldehyde or 100% methanol. Brain tissue is comprised mostly of neuronal and glial cells, organized into distinct compound structures, wherein the distinct compound structures may be further comprised of layers or substructures within the compound structure. Particular substructures may be composed primarily of neuronal bodies, or bundled axonal tracts. In these coronal sections, the CA1-CA3 regions of the hippocampus as well as the cerebral cortex can be clearly seen marked with NeuN, a common intracellular biomarker for neurons. The dentate gyrus and cerebral cortex can be visualized with DAPI, a common DNA stain. Additionally, GFAP (glial fibrillary acidic protein) stains glial cells intracellularly, including astrocytes which contribute to the functioning of the blood-brain barrier. FIG. 42 also demonstrates that different fixation protocols can affect the data obtained from antibody staining. For example, with 100% methanol fixation (right) it is likely the NeuN antigen epitope has been masked, denatured, or the protein washed away via the permeabilization effects of methanol. The result is no NeuN antibody staining with 100% methanol fixation.

Example 7—Methods of Using Spatial Transcriptomics Analysis to Identify Cell Type “Clusters”

[2649] In a non-limiting example, spatial analysis was performed on mouse spleen (FIGS. 39-41) and brain coronal slices (FIGS. 43-45).

[2650] A t-SNE plot was generated which identified distinct cell type clustering (FIGS. 39 and 43). The plot demonstrated the differential expression of genes captured using the spatial analysis methods described herein. The differential gene expression identified by clustering can be mapped back onto an image of the original biological sample in order to visualize the identity of the cells in the original tissue sample (FIGS. 39 and 43). FIGS. 40 and 41 show exemplary immunofluorescence staining on the left, identifying the presence and location of CD3, CD29, and DAPI. The immunofluorescence staining is overlaid with CD3 transcript counts (measured in UMI during spatial transcriptome analysis) (right), demonstrating that specific analytes can be captured and barcoded according to the spatial workflows described herein. FIGS. 44 and 45 show exemplary immunofluorescence staining of mouse brain (left) in three channels (NeuN; GFAP; DAPI). The immunofluorescence staining is overlaid with NeuN transcript counts (measured in UMI during spatial transcriptome analysis) (top) and GFAP UMI counts (bottom), demonstrating that specific analytes can be captured and barcoded according to the spatial workflows described herein.

Example 8—Methods for Blocking an Analyte Capture Sequence Binding to a Capture Probe

[2651] In a non-limiting example, non-specific antibody staining can be reduced by introducing a reversible blocking probe to the analyte capture agent(s), prior to applying the analyte capture agents to a tissue sample. FIG. 46 shows an exemplary illustration of an analyte capture sequence (a poly(A) sequence) blocked by a poly(T) sequence blocking probe, inhibiting the fluorescently labeled antibody from interacting with capture probes on a capture probe array. When the reversible

blocking probe is removed (e.g., by heating the analyte capture agent), the analyte capture sequence is free to interact with (e.g., hybridize to) the capture domain of capture probes on the array.

[2652] A method of blocking an analyte capture sequence can include DNA-barcoded antibodies including an analyte binding moiety (e.g., antibody) and a capture agent barcode domain. The capture agent barcode domain includes an analyte capture sequence, which could bind to the capture probe capture domain prior to analyte capture by the analyte binding moiety. The analyte capture sequence binding to a capture probe can be blocked by hybridizing the analyte capture sequence (e.g., the poly(A) 3' tail of the capture agent barcode domain) with a blocking probe. In this example, the blocking probes comprise an oligo (dT) sequence that can interact with the poly(A) tails of the analyte capture sequence (FIG. 46). Here, the analyte capture agent analyte capture sequence can be blocked before applying the analyte capture agents to the tissue sample. Additionally, excess blocking probes can be provided in a sample hydration solution (e.g., a permeabilization solution applied to the tissue sample).

[2653] In some embodiments, the analyte capture sequence can be reversibly blocked with a blocking probe prior to applying the analyte capture agents to the tissue sample, and when treated with a nuclease (e.g., RNase H) the blocking probe is digested, allowing the analyte capture sequence to interact with capture probes on the substrate. See, for example, FIG. 47.

[2654] FIG. 47 shows the performance of an oligo (dT) blocking probe under various conditions. The effectiveness of the oligo (dT) blocking probes varied as a function of the length of the blocking probe and the temperature of the array during the antibody staining protocol. Here, the above experiment was performed with oligo (dT) blocking probes of varying lengths at 4° C. and 37° C. FIG. 48 indicates analyte capture agents were blocked from binding to the capture probes at both temperatures, although shorter blocking probes were more effective under lower temperatures.

Example 9—Methods for Using a Spatially-Tagged Analyte Capture Agent in a Biological Sample

[2655] In a non-limiting example, DNA-barcoded antibodies are used to detect proteins in a biological sample. For example, a method of detecting proteins within a tissue sample using DNA-barcoded antibodies can include: (a) providing a capture probe array, wherein the capture probes include a spatial barcode and a capture domain; (b) contacting the substrate with a tissue sample (e.g., mouse spleen tissue) and drying the sectioned slides for 1 minute at 37° C.; (c) fixing the tissue sample with either 2% formaldehyde at room temperature or with methanol at -20° C. for 10 minutes; (d) rehydrating, blocking and permeabilizing the tissue sample with 3×SSC, 2% BSA, 0.1% Triton X, and 1U/μl RNase inhibitor for 10 minutes at 4° C.; (e) staining the tissue sample with fluorescent primary antibodies and a pool of DNA-barcoded antibodies in 3×SSC, 2% BSA, 0.1% Triton X, and 1U/μl RNase inhibitor for 30 minutes at 4° C.; (f) imaging the tissue sample to spatially detect target proteins (e.g., CD29, CD3) within the tissue using fluorescently-labelled and DNA-barcoded antibodies; (g) treating the tissue sample with a protease to permeabilize the tissue and release the antibody oligos; and (h) performing spatial transcriptomic analysis to identify the location of the target protein within the tissue sample. The steps of this method are depicted in FIG. 49.

[2656] The DNA-barcoded antibodies can include an analyte binding moiety (e.g., antibody) and a capture agent barcode domain. The antibodies interact with the target protein of the biological sample, and the capture agent barcode domains interact with the capture probes on the substrate. The fluorescence level from the primary antibodies interacting with the proteins of the biological sample is imaged in step (f), and the spatially-tagged analyte capture agents associated with the capture probes are used to identify the location of the target protein within the biological sample (FIG. 50). In some embodiments, non-specific antibody staining can be reduced by introducing a blocking probe to the analyte capture agent(s), prior to applying the analyte capture agents to a tissue sample, as described in Example 3.

[2657] In some embodiments, detecting and identifying the location of a target protein can be performed individually for each analyte of interest. In some embodiments, multiple proteins can be

detected and spatially profiled simultaneously within the same tissue sample. In some embodiments, multiplexing (e.g., simultaneously detecting multiple markers) allows for examination of the spatial arrangement of analytes of interest (e.g., proteins, DNA, RNA) as well as analyte interaction and co-localization thereby facilitating simultaneous analysis of multiple tissue markers.

[2658] FIG. 50 shows exemplary imaging results using the method depicted above and in FIG. 49, wherein the left image shows a tissue section of mouse spleen with fluorescent antibodies bound to CD29. The images on the right show the location of target proteins, CD29, CD3, F4/80, and CD169 within the tissue sample using the spatially-tagged analyte capture agents associated with the surface capture probes on the substrate. In particular, the method above was performed by fixing the tissue sample with 2% formaldehyde and permeabilizing with proteinase K.

Example 10—Methods for Using Spatially-Tagged Analyte Capture Agents to Detect Multiple Target Proteins by Introducing Antibodies Linked to Capture Agents and Molecular Identifiers in a Biological Sample

[2659] In a non-limiting example, multiple pluralities of DNA-barcoded antibodies can be used to simultaneously detect multiple target proteins (e.g., multiplexing) within a biological sample. For example, a method of detecting multiple target proteins within a biological sample can include the method described in Example 4, wherein two or more pluralities of analyte capture agents bind to two or more pluralities of analytes. Each analyte species is associated with a spatially-tagged analyte capture agent plurality, wherein each spatially-tagged analyte capture agent plurality possesses a barcode unique to the analyte. Multiple analytes can be detected and analyzed at the same time by determining the analyte binding moiety barcode that can be determined together with, or separately from the spatial transcriptome analysis using a sequencing technology, as described elsewhere herein. In other embodiments, antibody-barcodes can be determined by fluorescent in situ hybridization or in situ sequencing approaches, as described elsewhere herein or known in the art.

[2660] For example, FIG. 51 shows exemplary multiplexed DNA-barcoded and fluorescent antibody staining and sequencing results using the method depicted above, wherein the left immunofluorescent image shows tissue sections of mouse spleen with fluorescent and DNA-barcoded antibodies bound to CD29 and CD4. CD29 (Integrin beta 1) is a cell surface marker expressed in many stromal cells and can be seen in the red pulp portion of the spleen, while CD4 is a cell surface marker for T cell subsets and can be seen in the pockets of white pulp of the spleen. The images on the right show the location of the antibody barcodes recognizing target proteins, CD29, CD3, CD4, CD8, CD19, B220, F4/80, and CD169 within the tissue sample using the multiple spatially-tagged analyte capture agents. Each spatially-tagged analyte capture agent plurality possess an analyte binding moiety barcode unique to that plurality. FIG. 51 shows that CD3, CD4, and CD8, all cell surface markers for T cells, are seen to be located in the pockets of white pulp of the spleen. CD19, a cell surface marker for B cells, CD29, a cell surface marker for stromal cells, and F4/80 and CD169, both markers for macrophage cells, can be seen within the red pulp and white pulp respectively of the spleen tissue. The data of FIG. 51 indicates that protein detection using DNA-barcoded antibodies can be used for simultaneously identifying spatial locations of multiple (e.g., two or more, three or more) target proteins within a tissue sample.

OTHER EMBODIMENTS

[2661] It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

Claims

1.-84. (canceled)

85. A computer system comprising: a memory; one or more processors; and one or more programs for identifying a spatial location of a variant haplotype in a biological sample, wherein the one or more programs are stored in the memory and configured to be executed by the one or more processors, wherein the one or more programs include instructions for: (a) obtaining a plurality of sequencing reads from the biological sample; (b) aligning each locus of the plurality of sequencing reads to determine a haplotype identity of each sequence read of the plurality of sequencing reads; and (c) categorizing each haplotype of the plurality of sequencing reads to determine the spatial location in the biological sample.

86. The computer system of claim 85, further comprising using the spatial location of each haplotype to characterize a biological condition of the biological sample.

87. The computer system of claim 85, further comprising an imager configured to image the biological sample.

88. The computer system of claim 85, further comprising input data comprising the plurality of sequencing reads.

89. The computer system of claim 85, wherein the input data further comprise electronic data files of gene sequence variations, haplotypes, and/or a reference genome.

90. The computer system of claim 89, wherein the reference genome is a human reference genome.

91. The computer system of claim 85, wherein the plurality of sequencing reads is obtained from a two-dimensional array in contact with the biological sample and subsequently aligned to a reference genome.

92. The computer system of claim 85, wherein the plurality of sequencing reads comprises spatial barcodes with positional information relative to spatial location in the biological sample.

93. The computer system of claim 85, wherein the plurality of sequence reads is obtained by in situ sequencing of the two-dimensional array, high-throughput sequencing, and/or paired end sequencing.

94. The computer system of claim 85, wherein the haplotype identity is a heterozygous single nucleotide polymorphism (SNP), a heterozygous insert, or a heterozygous deletion.

95. A non-transitory computer-readable medium comprising instructions for identifying a spatial location of a variant haplotype in a biological sample, wherein the computer-readable medium stores one or more computer programs executable by a computer system for identifying the spatial location of the variant haplotype in the biological sample, wherein the computer system comprises: a memory; one or more processors; and one or more programs for identifying a spatial location of a variant haplotype in a biological sample, wherein the one or more programs are stored in the memory and configured to be executed by the one or more processors, wherein the one or more programs comprise instructions for: (a) obtaining a plurality of sequencing reads from the biological sample; (b) aligning each locus of the plurality of sequencing reads to determine a haplotype identity of each sequence read of the plurality of sequencing reads; and (c) categorizing each haplotype of the plurality of sequencing reads to determine the spatial location in the biological sample.

96. The non-transitory computer-readable medium of claim 95, wherein the one or more programs further comprise instructions for using the spatial location of each haplotype to characterize a biological condition of the biological sample.

97. The non-transitory computer-readable medium of claim 95, wherein the computer system further comprises an imager configured to image the biological sample.

98. The non-transitory computer-readable medium of claim 95, wherein the computer system further comprises input data comprising the plurality of sequencing reads.

99. The non-transitory computer-readable medium of claim 95, wherein the input data further comprise electronic data files of gene sequence variations, haplotypes, and/or a reference genome.

100. The non-transitory computer-readable medium of claim 95, wherein the plurality of sequencing reads is obtained from a two-dimensional array in contact with the biological sample and subsequently aligned to a reference genome.

101. The non-transitory computer-readable medium of claim 100, wherein the reference genome is a human reference genome.

102. The non-transitory computer-readable medium of claim 95, wherein the plurality of sequencing reads comprises spatial barcodes with positional information relative to spatial location in the biological sample.

103. The non-transitory computer-readable medium of claim 95, wherein the plurality of sequence reads is obtained by in situ sequencing of the two-dimensional array, high-throughput sequencing, and/or paired end sequencing.

104. The non-transitory computer-readable medium of claim 95, wherein the haplotype identity is a heterozygous single nucleotide polymorphism (SNP), a heterozygous insert, or a heterozygous deletion.
