



US 20250280784A1

(19) **United States**(12) **Patent Application Publication**
BARDOL et al.(10) **Pub. No.: US 2025/0280784 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **RESISTANCE TO LEVEILLULA TAURICA IN PEPPER***A01H 1/02* (2006.01)*A01H 1/04* (2006.01)*A01H 1/06* (2006.01)*C12N 5/04* (2006.01)(71) Applicant: **VILMORIN & CIE**, Paris (FR)(72) Inventors: **Nicolas BARDOL**, Loire-Authion (FR);
Julie LEDERER, Saint Rémy de
Provence (FR); **Sylvie MANIN**,
Ledenon (FR); **Benoît GORGUET**,
Chappes (FR)(52) **U.S. Cl.**CPC *A01H 6/822* (2018.05); *A01H 1/021*
(2021.01); *A01H 1/045* (2021.01); *A01H 1/06*
(2013.01); *A01H 1/1255* (2021.01); *C12N*
5/04 (2013.01)(73) Assignee: **VILMORIN & CIE**, Paris (FR)(21) Appl. No.: **18/294,738**(57) **ABSTRACT**(22) PCT Filed: **Aug. 5, 2022**(86) PCT No.: **PCT/EP2022/072057**


§ 371 (c)(1),

(2) Date: **Feb. 2, 2024**(30) **Foreign Application Priority Data**

Aug. 6, 2021 (EP) 21306101.3

Publication Classification(51) **Int. Cl.***A01H 6/82* (2018.01)*A01H 1/00* (2006.01)

A *Capsicum* plant resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*, a cell, plant part or seed of the *Capsicum* plant and to methods for identifying it, and the use of the plants for improving the yield of pepper production in an environment infected by *Leveillula taurica* and/or for limiting or controlling an infection by *Leveillula taurica*; wherein, the *Capsicum* plant includes introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein the QTL is located on chromosome 6 within the genomic interval delimited by the marker PE-0014628 (SEQ ID NO: 1) and the marker PE-0021476 (SEQ ID NO: 37).

Specification includes a Sequence Listing.




Figure 1




Figure 2




Figure 3





Figure 4

A.



B.




Figure 5

RESISTANCE TO *LEVEILLULA TAURICA* IN PEPPER

[0001] Pepper (*Capsicum* sp.), from the Solanaceae family, is one of the most important vegetable crops. Whilst its species are native to the Americas, it is now widely cultivated throughout the warm, temperate, tropical and subtropical countries, in field or greenhouse production. Pepper production is affected by a variety of diseases, such as anthracnose (*C. acutatum*), *Phytophthora* blight (*Phytophthora capsici*), bacterial spot (*Xanthomonas campestris* pv. *vesicatoria*), and bacterial wilt (*Ralstonia solanacearum*). Powdery mildew is one the most important diseases of pepper. The initial symptoms of powdery mildew on peppers are a powdery-white fungal growth on the undersides of leaves with light-green to yellow blotches on the upper leaf surfaces. These areas turn brown with time, and the affected areas coalesce, causing a general yellowing of the entire leaf. The outer edges of leaves may curl upward. The older leaves lower in the canopy are usually infected first and show symptoms before the younger leaves. Infected leaves drop from the plant prematurely. This loss of photosynthetic leaf area slows plant growth and fruit development. The defoliation also exposes fruit to direct sunlight, which can lead to sunscalding of the fruit.

[0002] *Leveillula taurica*, the fungus that causes powdery mildew on pepper, is an obligate parasite that needs living host tissue to grow and reproduce. The fungus survives during periods between pepper crops on alternate host species. *L. taurica* is somewhat unusual for a powdery mildew fungus in that it has a very wide host range. The pathogen can infect over 700 species of plants in 59 different families. Susceptible crops and horticultural plants include alfalfa, artichoke, cotton, eggplant, onion, tomato, mesquite, and monkey-flower. Susceptible weed species include sow-thistle, cocklebur, groundsel, desert tobacco, and shepherd's purse. There appears to be some host specificity among isolates of the pathogen as individual isolates may be able to infect some reported host species but not others.

[0003] The pathogen is disseminated as spores carried in the wind. Powdery mildew occurs on peppers in both humid and arid environments, as the spores can germinate over the range of relative humidity levels from zero to 100%. However, the highest rates of spore germination occur when nighttime humidity levels are between 90% and 95% and daytime levels are above 85%. The spores can germinate at temperatures from 10° C. to 35° C. with the optimum temperature range is from 15° C. to 25° C. Once germination and infection have occurred, disease development is favored by warm days and cool nights, with the most severe disease levels occurring with daytime temperatures near 30° C. and nighttime temperatures below 25° C. Disease development is suppressed at temperatures above 35° C. While infection is highest under humid conditions, defoliation is favored by drier conditions. Therefore, the impact of the disease is most severe when humidity levels change from humid to dry.

[0004] Yield reductions from powdery mildew result, in part, from a loss of photosynthetic area that inhibits plant growth and fruit development. Fewer flowers form on severely infected plants, and flowers may also drop off of infected plants. Additional losses come from sunscalding of the fruit.

[0005] With the wide host range of the pathogen and wind dispersal of the spores, crop rotation and management of nearby weed hosts may not provide an adequate level of

control without using other means of control. Low light and high humidity favor the development of powdery mildew on peppers, so cultural practices that minimize these conditions can help lower disease severity. Such practices include the selection of sites with well-drained soils and good air circulation; the adjustment of planting densities and row orientations to promote good air circulation and light penetration into the canopy; or the avoidance of excessive fertilization that would result in overly dense canopies. In greenhouse production, increasing the temperature in the greenhouse can lower humidity levels and slow disease development.

[0006] A second control method for powdery mildew on peppers is the application of fungicides. Several fungicides are available for controlling powdery mildew on peppers, although chemical control may be hampered by the semi-endoparasitic nature of the fungi. Furthermore, the use of fungicides should be accompanied by appropriate fungicide resistance management strategies to prevent the development of fungicide-resistant strains of the pathogen. In addition, the ban on certain chemical treatments, or their social perception, reduces the ability of agronomic control of the disease.

[0007] It is thus desirable to find new pepper germplasms having a moderate to high resistant to the disease. Some sources of resistance have been identified, although most of them showed a low level of resistance. Daubeze et al, 1995 identified a resistant Ethiopian line H3. Lefebvre et al 2003 analyzed doubled haploid lines derived from a cross between H3 and the susceptible line Vania. The plants were evaluated in two different infections conditions and five QTLs distributed on five distinct chromosomes, P5, P6, P9, P10 and P12 were detected, with additive effects and epistatic interactions. However, some of the already identified QTLs may be linked to undesirable traits and/or provide only a partial resistance to the pathogen. Furthermore, the effect of QTLs may be additive, such that there is an interest in finding new QTLs, the effect of which could be cumulated to improve the resistance to the pathogen. There remains thus a strong need to find QTLs and sources of resistance to *Leveillula taurica* in pepper.

SUMMARY OF THE INVENTION

[0008] The present invention first provides a *Capsicum* plant resistant to powdery mildew, comprising introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein said QTL is present on chromosome 6.

[0009] In some aspects, the invention provides a *Capsicum* plant resistant to powdery mildew, comprising introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein said QTL is located on chromosome 6 within the chromosomal region delimited by the markers PE-0014628 (SEQ ID NO: 1) and PE-0021476 (SEQ ID NO: 37).

[0010] In some aspects, the invention provides a *Capsicum* plant resistant to powdery mildew, comprising introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein said QTL is present on chromosome 6 in the genome of a seed of *Capsicum annuum* ST17-1157, deposited at NCIMB under accession number NCIMB 43602.

[0011] The present invention also provides a *Capsicum* plant resistant to powdery mildew, comprising introgressed

in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein the presence of said QTL on chromosome 6 and conferring said resistance is characterized by at least one allele selected from the group consisting of allele A of PE-0056543 (SEQ ID NO: 9), allele A of PE-0056544 (SEQ ID NO:13), allele C of PE-0056545 (SEQ ID NO: 17) and allele C of PE-0056546 (SEQ ID NO: 21).

[0012] The present invention also provides a *Capsicum* plant resistant to powdery mildew, comprising introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein said QTL encompasses the CA06g26150 gene and/or the CA06g26250 gene, in particular the CA06g26150 gene.

[0013] The present invention also provides a *Capsicum* plant resistant to powdery mildew, wherein said *Capsicum* plant comprises a mutant allele of the CA06g26150 gene and/or the CA06g26250 gene, in particular the CA06g26150 gene, wherein said mutant allele confers resistance to powdery mildew.

[0014] The present invention also provides a *Capsicum* seed, which can be grown into a *Capsicum* plant according to the invention.

[0015] Also provided is an in vitro cell or tissue culture of regenerable cells of the *Capsicum* plant according to the invention, wherein the regenerable cells are derived from an embryo, protoplast, meristematic cell, callus, pollen, leaf, anther, stem, petiole, root, root tip, seed, flower, cotyledon, and/or hypocotyl.

[0016] Further provided is a method for detecting and/or selecting *Capsicum* plants comprising in their genome a QTL on chromosome 6 as found in the genome of *Capsicum annuum* ST17-1157 deposited at the NCIMB under accession number NCIMB 43602, said QTL conferring resistance to powdery mildew, comprising the detection of at least one of the following alleles: allele A of PE-0056543 (SEQ ID NO: 9), allele A of PE-0056544 (SEQ ID NO:13), allele C of PE-0056545 (SEQ ID NO: 17) and allele C of PE-0056546 (SEQ ID NO: 21).

[0017] Also provided is the use of a plant or seed of ST17-1157, deposited at NCIMB under accession number NCIMB-43602, or progeny thereof bearing the sequences conferring the resistance to powdery mildew, as a breeding partner in a breeding program for obtaining a *Capsicum* plant resistant to powdery mildew.

[0018] The invention also relates to a method for producing a *Capsicum* plant resistant to powdery mildew, comprising the steps of

[0019] (a) crossing a plant grown from a seed of ST17-1157 deposited at NCIMB under accession number NCIMB 43602 or progeny thereof bearing the QTL conferring the resistance to powdery mildew, and a susceptible *Capsicum* plant,

[0020] (b) selecting one plant resistant to powdery mildew in the progeny thus obtained, said plant bearing the QTL conferring the resistance to powdery mildew;

[0021] (c) optionally self-pollinating one or several times the resistant plant obtained at step b) and selecting a plant resistant to powdery mildew in the progeny thus obtained;

[0022] (d) optionally backcrossing the resistant plant selected in step b) or c) with a susceptible *Capsicum* plant and

[0023] (e) selecting a plant resistant to powdery mildew,

wherein selecting plants resistant to powdery mildew in steps b), c) and/or e) comprises the detection of nucleic acid markers, in particular SNP markers.

[0024] Further provided is a method for producing *Capsicum* plants resistant to powdery mildew, comprising the steps of:

[0025] (a1) crossing a plant grown from a seed deposited under accession number NCIMB 43602 or progeny thereof bearing the sequences conferring the resistance to powdery mildew and a susceptible *Capsicum* plant, thus generating the F1 population,

[0026] (a2) advancing the F1 population to create F2 population,

[0027] (b) selecting one resistant plant in the progeny thus obtained;

[0028] c) optionally self-pollinating one or several times the resistant plant obtained at step b) and selecting a plant resistant to white rust in the progeny thus obtained;

[0029] (d) optionally backcrossing the resistant plant selected in step b) or c) with a susceptible *Capsicum* plant, and

[0030] (e) selecting a plant resistant to powdery mildew, wherein selecting plants resistant to powdery mildew in steps b), c) and/or e) comprises the detection of nucleic acid markers, in particular SNP markers.

[0031] Also provided is a method for obtaining *Capsicum* plants of agricultural interest, resistant to powdery mildew, comprising the steps of:

[0032] (a) backcrossing a plant obtained by germinating a seed deposited at the NCIMB under accession number NCIMB 43602 or progeny thereof bearing the sequences conferring the resistance to powdery mildew, with a *Capsicum* plant susceptible to powdery mildew,

[0033] (b) Selecting a plant resistant to powdery mildew, wherein the selection comprises the detection of nucleic acid markers, in particular SNP markers.

[0034] The invention also relates to a *Capsicum* plant resistant to powdery mildew obtainable by a method or use according to the invention. Said *Capsicum* plant comprises a QTL on chromosome 6 as defined herein, conferring the resistance to powdery mildew.

[0035] The invention further concerns a method for identifying a molecular marker linked with a QTL conferring the resistance to powdery mildew, said QTL being present in seeds deposited at the NCIMB under accession number NCIMB 43602, comprising:

[0036] (a) identifying a molecular marker in *Capsicum* genome, in the chromosomal region delimited on chromosome 6 by PE-0014628 (SEQ ID NO: 1) and PE-0021476 (SEQ ID NO: 37); and

[0037] (b) determining whether an allele or state of said molecular marker is associated with resistance to powdery mildew in a segregating population comprising *Capsicum* plants exhibiting said resistance.

[0038] Also provided is a method for improving the yield of pepper production in an environment infected by *Leveillula taurica*, comprising growing a *Capsicum* plant according to the invention.

[0039] Further provided is a method for protecting a field, tunnel, greenhouse or glasshouse of pepper from an infection by *Leveillula taurica*, comprising growing a *Capsicum* plant according to the invention.

[0040] The invention also relates to the use of a *Capsicum* sp. plant according to the invention for limiting or controlling an infection by *Leveillula taurica*.

[0041] Another embodiment of the invention relates to a method of producing pepper fruit comprising:

[0042] a) growing a *Capsicum* plant according to the invention;

[0043] b) allowing said plant to set fruit; and

[0044] c) harvesting fruit of said plant, preferably at maturity and/or over-maturity; and optionally

[0045] d) processing said pepper fruit into a pepper processed food.

Definitions

[0046] As used herein, the term “pepper”, “pepper plant” or “*Capsicum* plant” relates to any species, variety, cultivar, or population of the *Capsicum* genus. The *Capsicum* genus is known to comprise 20-27 species, five of which are domesticated: *Capsicum annuum*, *Capsicum baccatum*, *Capsicum frutescens*, *Capsicum chinense*, and *Capsicum pubescens*. The vast majority of commercial varieties of peppers belong to the species *Capsicum annuum*, which comprises sweet peppers (peppers with a no pungency) and hot peppers (peppers with pungency from low level to high level). Sweet peppers include bell peppers, which are fruit of plants in the Grossum cultivar group of the species *Capsicum annuum*. Sweet pepper plants can be produced in different colors, from immature green color to mature color of red, yellow, orange, white or purple. In particular, the *Capsicum* plant may be of one of the following types: Dulce Italiano, Lamuyo, Blocky Florida, Open field and blocky protected, Ancho, Anaheim, Marconi, Jalapeno, Cayenne, Charleston or Sivri.

[0047] As used herein, the term “plant part” refers to any part of a plant including but not limited to the shoot, root, stem, seeds, fruits, leaves, petals, flowers, ovules, branches, petioles, internodes, pollen, stamen, rootstock, scion and the like.

[0048] As used herein, the term “allele” refers to any of several alternative or variant forms of a genetic unit, such as a gene, which are alternative in inheritance because they are positioned at the same locus in homologous chromosomes. Such alternative or variant forms may be the result of single nucleotide polymorphisms, insertions, inversions, translocations or deletions, or the consequence of gene regulation caused, for example, by chemical or structural modification, transcription regulation or post-translational modification/regulation. In a diploid cell or organism such as pepper, the two alleles of a given gene or genetic element typically occupy corresponding loci on a pair of homologous chromosomes.

[0049] As used herein, the term “genotype” refers to the genetic makeup of an individual cell, cell culture, tissue, organism (e.g., a plant), or group of organisms.

[0050] As used herein, the term “locus” (plural: “loci”) refers to any site that has been defined genetically, this can be a si (Single Nucleotide Polymorphism), by several markers (e.g. SNPs), or by two flanking markers (e.g. SNPs).

[0051] As used herein, the terms “Quantitative Trait Locus (QTL)” (plural: “Quantitative Trait Loci”) refer to a genomic region that may comprise one or more genes or regulatory sequences. A QTL may for instance comprise one or more genes of which products confer genetic resistance or tolerance. Alternatively, a QTL may for instance comprise

regulatory genes or sequences of which products influence the expression of genes on other loci in the genome of the plant thereby conferring the resistance or tolerance. The QTL of the present invention may be defined by indicating its genetic location in the genome of the respective pathogen-resistant accession using one or more molecular genomic markers. One or more markers, in turn, indicate a specific locus. Distances between loci are usually measured by frequency of crossing-over between loci on the same chromosome. The farther apart two are, the more likely that a crossover will occur between them. Conversely, if two loci are close together, a cross over is less likely to occur between them. As a rule, one centimorgan (cM) is equal to 1% recombination between loci (marker). When a QTL can be indicated by multiple markers, the genetic distance between the end-point markers is indicative of the size of the QTL.

[0052] The term “resistance” is as defined by the ISF (International Seed Federation) Vegetable and Ornamental Crops Section for describing the reaction of plants to pests or pathogens, and abiotic stresses for the Vegetable Seed Industry. Specifically, by resistance, it is meant the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure.

[0053] By “tolerance” is meant the ability of a plant variety to endure biotic and abiotic stress without serious consequences for growth, appearance and yield.

[0054] As used herein, the term “susceptible” refers to a plant that is unable to restrict the growth and development of a specified pest or pathogen.

[0055] As used herein, the term “offspring” or “progeny” refers to any plant resulting as progeny from a vegetative or sexual reproduction from one or more parent plants or descendants thereof. For instance, an offspring plant may be obtained by cloning or selfing of a parent plant or by crossing two parent plants and include selfings as well as the F1 or F2 or still further generations. An F1 is a first-generation offspring produced from parents at least one of which is used for the first time as donor of a trait, while offspring of a second generation (F2) or subsequent generations (F3, F4, etc.) are specimens produced from selfing of F1's, F2s, etc. An F1 may thus be (and usually is) a hybrid resulting from a cross between two true breeding parents (true-breeding is homozygous for a trait), while an F2 may be (and usually is) an offspring resulting from self-pollination of said F1 hybrids.

[0056] As used herein, the terms “cross”, “crossing” refer to the process by which the pollen of one flower on one plant is applied (artificially or naturally) to the ovule (stigma) of a flower on another plant.

[0057] As used herein, the term “heterozygote” refers to a diploid or polyploidy cell or plant having different alleles (forms of a given gene or sequences) present at at least one locus.

[0058] As used herein, the term “heterozygous” refers to the presence of different alleles (forms of a given gene or sequences) at a particular locus.

[0059] As used herein, the term “homozygote” refers to an individual cell or plant having the same alleles at one or more loci on all homologous chromosomes.

[0060] As used herein, the term “homozygous” refers to the presence of identical alleles at one or more loci in homologous chromosomal segments.

[0061] As used herein, the term “hybrid” refers to any individual cell, tissue or plant resulting from a cross between parents that differ in one or more genes.

[0062] As used herein, the term “inbred” or “line” refers to a relatively true-breeding strain.

[0063] As used herein, the term “phenotype” refers to the observable characters of an individual cell, cell culture, organism (e.g. a plant), or group of organisms which results from the interaction between that individual genetic makeup (i.e. genotype) and the environment.

[0064] As used herein, the terms “introgression”, “introgressed” and “introgressing” refer to the process whereby genes of one species, variety or cultivar are moved into the genome of another species, variety or cultivar, by crossing those species. The crossing may be natural or artificial. The process may be optionally be completed by backcrossing to the recurrent parent, in which case introgression refers to infiltration of the genes of one species into the gene pool of another through repeated backcrossing of an interspecific hybrid with one of its parents. An introgression may be also described as a heterologous genetic material stably integrated in the genome of a recipient plant.

[0065] As used herein, the terms “molecular marker” refer to an indicator that is used in methods for visualizing differences in characteristics of nucleic acid sequences. Examples of such indicators are restriction fragment length polymorphism (RFLP) markers, amplification fragment length polymorphism (AFLP) markers, single nucleotide polymorphisms (SNPs), insertion mutations, microsatellite markers (SSRs), sequence-characterized amplified regions (SCARs), cleaved amplified polymorphic sequence (CAPS) markers or isozyme markers or combinations of the markers described herein which defines a specific genetic and chromosomal location. Mapping of molecular markers in the vicinity of an allele is a procedure which can be performed quite easily by the person skilled in the art using common molecular techniques.

[0066] As used herein, “marker-based selection” or “marker-assisted selection (MAS)” or “marker-assisted breeding [text missing or illegible when filed] undesirable traits, so that those plants can be used (or avoided) in a selective breeding program.

[0067] As used herein, the term “primer” refers to an oligonucleotide which is capable of annealing to the amplification target allowing a DNA polymerase to attach, thereby serving as a point of initiation of DNA synthesis when placed under conditions in which synthesis of primers extension product is induced, i.e., in the presence of nucleotides and an agent for polymerization such as DNA polymerase and at a suitable temperature and pH. The primer is preferably single stranded for maximum efficiency in amplification. Preferably, the primer is an oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact length of the primers will depend on many factors, including temperature and composition (A/T and G/C content) of primer. A pair of bi-directional primers consists of one forward and one reverse primer as commonly used in the art of DNA amplification such as in PCR amplification.

[0068] As used herein, a single nucleotide polymorphism (SNP) is a DNA sequence variation occurring when a single AAGCCTA to AAGCTTA, contain a difference in a single nucleotide. In this case there are two alleles: C and T.

[0069] In the context of the invention and unless otherwise specified, DNA strand and allele designation and orientation for the markers (i.e. SNPs) disclosed in the present application are mentioned according to the TOP/BOT method developed by Illumina (https://www.illumina.com/documents/products/technotes/technote_topbot.pdf).

[0070] By “powdery mildew”, it is meant a disease which primarily affects leaves on pepper plants. Symptoms include patchy, white, powdery growth, called sporulation, that enlarges and coalesces to cover the entire lower leaf surface. At times the powdery growth is present on the upper leaf surface as well. Powdery mildew is caused in pepper by the fungal pathogen *Leveillula taurica*.

[0071] In the context of the present application, a chromosomal region delimited by two markers X and Y (e.g. SNPs) refers to the section of the chromosome lying between the positions of these two markers and comprising said markers, therefore the nucleotide sequence of this chromosomal region begins with the nucleotide corresponding to marker X and ends with the nucleotide corresponding to marker Y, i.e. the markers are comprised within the region they delimit.

[0072] By association, or genetic association, and more specifically genetic linkage, it is to be understood that a polymorphism of a genetic marker (e.g. a specific allele of the SNP marker) and the phenotype of interest occur simultaneously, i.e. are inherited together, more often than would be expected by chance occurrence, i.e. there is a non-random association of the allele and of the genetic sequences responsible for the phenotype, as a result of their genomic proximity.

SEQUENCE LISTING

[0073] SEQ ID NO: 1 shows a sequence surrounding the marker PE-0014628.

[0074] SEQ ID NO: 2 shows the sequence of a forward primer for detecting the susceptible allele of PE-0014628.

[0075] SEQ ID NO: 3 shows the sequence of a forward primer for detecting the resistant allele of PE-0014628.

[0076] SEQ ID NO: 4 shows the sequence of a common reverse primer for PE-0014628.

[0077] SEQ ID NO: 5 shows a sequence surrounding the marker PE-0056542.

[0078] SEQ ID NO: 6 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056542.

[0079] SEQ ID NO: 7 shows the sequence of a forward primer for detecting the resistant allele of PE-0056542.

[0080] SEQ ID NO: 8 shows the sequence of a common reverse primer for PE-0056542.

[0081] SEQ ID NO: 9 shows a sequence surrounding the marker PE-0056543.

[0082] SEQ ID NO: 10 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056543.

[0083] SEQ ID NO: 11 shows the sequence of a forward primer for detecting the resistant allele of PE-0056543.

[0084] SEQ ID NO: 12 shows the sequence of a common reverse primer for PE-0056543.

[0085] SEQ ID NO: 13 shows a sequence surrounding the marker PE-0056544.

[0086] SEQ ID NO: 14 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056544.

[0087] SEQ ID NO: 15 shows the sequence of a forward primer for detecting the resistant allele of PE-0056544.

[0088] SEQ ID NO: 16 shows the sequence of a common reverse primer for PE-0056544.

[0089] SEQ ID NO: 17 shows a sequence surrounding the marker PE-0056545.

[0090] SEQ ID NO: 18 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056545.

[0091] SEQ ID NO: 19 shows the sequence of a forward primer for detecting the resistant allele of PE-0056545.

[0092] SEQ ID NO: 20 shows the sequence of a common reverse primer for PE-0056545.

[0093] SEQ ID NO: 21 shows a sequence surrounding the marker PE-0056546.

[0094] SEQ ID NO: 22 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056546.

[0095] SEQ ID NO: 23 shows the sequence of a forward primer for detecting the resistant allele of PE-0056546.

[0096] SEQ ID NO: 24 shows the sequence of a common reverse primer for PE-0056546.

[0097] SEQ ID NO: 25 shows a sequence surrounding the marker PE-0056547.

[0098] SEQ ID NO: 26 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056547.

[0099] SEQ ID NO: 27 shows the sequence of a forward primer for detecting the resistant allele of PE-0056547.

[0100] SEQ ID NO: 28 shows the sequence of a common reverse primer for PE-0056547.

[0101] SEQ ID NO: 29 shows a sequence surrounding the marker PE-0056549.

[0102] SEQ ID NO: 30 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056549.

[0103] SEQ ID NO: 31 shows the sequence of a forward primer for detecting the resistant allele of PE-0056549.

[0104] SEQ ID NO: 32 shows the sequence of a common reverse primer for PE-0056549.

[0105] SEQ ID NO: 33 shows a sequence surrounding the marker PE-0056550.

[0106] SEQ ID NO: 34 shows the sequence of a forward primer for detecting the susceptible allele of PE-0056550.

[0107] SEQ ID NO: 35 shows the sequence of a forward primer for detecting the resistant allele of PE-0056550.

[0108] SEQ ID NO: 36 shows the sequence of a common reverse primer for PE-0056550.

[0109] SEQ ID NO: 37 shows a sequence surrounding the marker PE-0021476.

[0110] SEQ ID NO: 38 shows the sequence of a forward primer for detecting the susceptible allele of PE-0021476.

[0111] SEQ ID NO: 39 shows the sequence of a forward primer for detecting the resistant allele of PE-0021476.

[0112] SEQ ID NO: 40 shows the sequence of a common reverse primer for PE-0021476.

[0113] SEQ ID NO: 41 shows a coding sequence of the CA06g26150 gene in a susceptible plant (Maor).

[0114] SEQ ID NO: 42 shows a coding sequence of the CA06g26250 gene in a susceptible plant (Maor).

[0115] SEQ ID NO: 43 shows the sequence of a forward primer for amplifying the susceptible allele of the RZ marker.

[0116] SEQ ID NO: 44 shows the sequence of a forward primer for amplifying the resistant allele of the RZ marker.

[0117] SEQ ID NO: 45 shows the sequence of a common reverse primer for amplifying the resistant or susceptible allele of the RZ marker.

[0118] SEQ ID NOs: 46 to 75 show sequences surrounding, respectively markers PE-0011447, PE-0011448, PE-0019751, PE-0021027, PE-0022402, PE-0018388, PE-0028409, PE-0006573, PE-0001206, PE-0005987, PE-0005367, PE-0006634, PE-0006718, PE-0012892, PE-0004191, PE-0005575, PE-0006320, PE-0016499, PE-0013625, PE-0022671, PE-0005787, PE-0020909, PE-0000401, PE-0003984, PE-0014672, PE-0005702, PE-0021582, PE-0007248, PE-0003669, PE-0014672.

LEGEND OF THE FIGURES

[0119] FIG. 1 shows a genetic distance tree showing, inter alia, the genetic distance of line LT17 with respect to other *Capsicum* accessions.

[0120] FIG. 2 shows the distribution of the AUDPC (Area under the disease progress curve) for the parameter "Global note" measured on a segregating F2 population from the cross between the susceptible line Maor and the LT17 Line.

[0121] FIG. 3 shows the logarithm of odds (LOD), across chromosome 6 of the pepper genome, for the phenotype of resistance to *Leveillula taurica*.

[0122] FIG. 4 shows the distribution of disease scoring function to the presence of the QTL on chromosome 6, in two test locations.

[0123] FIG. 5: Panel (A) shows the disease scoring in different plant populations: S Check 3 (sensible control), S check (sensible control), IS check (Intermediate sensible control), Parent S (sensible), QTL6 SS (homozygous sensible at QTL6), QTL6 RS (heterozygous at QTL6), QTL6 RR (homozygous at QTL6), F4-RR-deposit-2 (F4 population ST117F04-1157/BLK homozygous resistant at QTL6), R check (resistant control) and Parent R (resistant parent). Panel (B) shows the result of a Tukey's test on the disease scoring in the same different plant populations.

DETAILED DESCRIPTION OF THE INVENTION

[0124] According to a first aspect, the present invention is directed to a *Capsicum* plant resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*, comprising introgressed in its genome a quantitative trait locus (QTL) on chromosome 6, conferring resistance to powdery mildew.

[0125] In some embodiments, said QTL is homozygously present in the genome of the plant.

[0126] In some embodiments, said QTL is heterozygously present in the genome of the plant.

[0127] In some embodiments, said QTL is present on chromosome 6 in the genome of a seed of *Capsicum annuum* ST17-1157, deposited at NCIMB accession number NCIMB 43602.

[0128] In some embodiments, said QTL is located on chromosome 6 within the chromosomal region delimited by markers PE-0014628 (SEQ ID NO: 1) and PE-0021476 (SEQ ID NO: 37). In some embodiments, said QTL is located on chromosome 6 within the chromosomal region delimited by positions 232015212 and 235235179 on the public genome *Capsicum annuum* cv CM334 v.1.55 (accessible on <http://www.solgenomics.net>). In particular, said QTL can be located on chromosome 6 within the chromo-

somal region delimited by markers PE-0056542 (SEQ ID NO: 5) and PE-0056550 (SEQ ID NO: 33). More particularly, said QTL is located on chromosome 6 within the chromosomal region delimited within the chromosomal region delimited by PE-0014628 (SEQ ID NO: 1) and PE-0056550 (SEQ ID NO: 33). More specifically, said QTL is located on chromosome 6 within the chromosomal region delimited by PE-0056542 (SEQ ID NO: 5) and PE-0021476 (SEQ ID NO: 37).

[0129] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by at least one marker selected from the group consisting of PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37), preferably at least one marker selected from the group consisting of PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29) and PE-0056550 (SEQ ID NO: 33), or any other marker comprised within the chromosomal region delimited by PE-0056542 (SEQ ID NO: 5) and PE-0056550 (SEQ ID NO: 33).

[0130] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0014628 marker on chromosome 6. In particular, said detection of the PE-0014628 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 1, or a fragment thereof including the [A/G] polymorphism at position 51 of SEQ ID NO: 1. For example, the forward primer for detecting the susceptible allele of the PE-0014628 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 1 or its complementary sequence, may consist of the sequence 5'-GAAGGTCGGAGTCAACGGATT-AGAAGAATAAGCACGAATAACCGCG-3' (SEQ ID NO: 2), the forward primer for detecting the resistant allele of the PE-0014628 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 1 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCTAGAAGAATAAGCACGAATAACCGCA-3' (SEQ ID NO: 3), and the common reverse primer may consist of the sequence 5'-AATTCCTCAACCAATCTATTTTCTTGAA-3' (SEQ ID NO: 4). Using primers consisting of sequences SEQ ID NO: 2, SEQ ID NO: 3 and SEQ ID NO: 4, detection of an alanine (A) rather than a guanine (G), at position 51 of the sequence SEQ ID NO: 1, or a thymine (T) rather than a cytosine (C) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0131] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056542 marker on chromosome 6. In particular, detection of the PE-0056542 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two

forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 5, or a fragment thereof including the [T/C] polymorphism at position 151 of SEQ ID NO: 5. For example, the forward primer for detecting the susceptible allele of the PE-0056542 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 5 or its complementary sequence, may consist of the sequence 5'-GAAGGTCGGAGTCAACGGATTGTGGTTATGACTGGGCCCTT-3' (SEQ ID NO: 6), the forward primer for detecting the resistant allele of the PE-0056542 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 5 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCTGTGGTTATGACTGGGCCCTC-3' (SEQ ID NO: 7), and the common reverse primer may consist of the sequence 5'-TAAGAGCTCTGTGTCCAAAACCATCTT-3' (SEQ ID NO: 8). Using primers consisting of sequences SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 8, detection of a cytosine (C) rather than a thymine (T) at position 151 of the sequence SEQ ID NO: 5, or a guanine (G) rather than an alanine (A) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0132] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056543 marker on chromosome 6. In particular, detection of the PE-0056543 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 9, or a fragment thereof including the [G/A] polymorphism at position 151 of SEQ ID NO: 9. For example, the forward primer for detecting the susceptible allele of the PE-0056543 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 9 or its complementary sequence, may consist of the sequence 5'-GAAGGTCGGAGTCAACGGATT-CATTAGAAGATCCAAATAATACTTTGAG-3' (SEQ ID NO: 10), the forward primer for detecting the resistant allele of the PE-0056543 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 9 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCTAGATTCATTAGAAGATC-CAAATAATACTTTGAA-3' (SEQ ID NO: 11), and the common reverse primer may consist of the sequence 5'-ATGAAACCTTCCGTCAGATTATCGAAGTT-3' (SEQ ID NO: 12). Using primers consisting of sequences SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 12, detection of an alanine (A) rather than a guanine (G) at position 151 of the sequence SEQ ID NO: 9, or a thymine (T) rather than a cytosine (C) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0133] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056544 marker on chromosome 6. In particular, detection of the PE-0056544 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 13, or a fragment thereof including the [G/T] polymorphism at

position 151 of SEQ ID NO: 13. For example, the forward primer for detecting the susceptible allele of the PE-0056544 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 13 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTT-CATGCTGGTGTCTATTTTGGCCATCGG-3' (SEQ ID NO: 14), the forward primer for detecting the resistant allele of the PE-0056544 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 13 or its complementary sequence, may consist of the sequence 5'-GAAGGTGCGGAGTCAACGGATTAGGTGCTGTAT-TTTTGGCCATCGT-3' (SEQ ID NO: 15), and the common reverse primer may consist of the sequence 5'-TTCCTCTTCTGGATATCCTTCTCATATA-3' (SEQ ID NO: 16). Using primers consisting of sequences SEQ ID NO: 14, SEQ ID NO: 15 and SEQ ID NO: 16, detection of a thymine (T) rather than a guanine (G) at position 151 of the sequence SEQ ID NO: 13, or an alanine (A) rather than a cytosine (C) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0134] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056545 marker on chromosome 6. In particular, detection of the PE-0056545 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 17, or a fragment thereof including the [T/C] polymorphism at position 151 of SEQ ID NO: 17. For example, the forward primer for detecting the susceptible allele of the PE-0056545 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 17 or its complementary sequence, may consist of the sequence 5'-GAAGGTGCGGAGTCAACGGAT-TCTGTCAAGTGGCCTTCATAAGATCA-3' (SEQ ID NO: 18), the forward primer for detecting the resistant allele of the PE-0056545 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 17 or its complementary sequence, may consist of the sequence 5'-GAAGGTGAC-CAAGTTCATGCTGTCAAGTGGCCTTCATAAGATCG-3' (SEQ ID NO: 19), and the common reverse primer may consist of the sequence 5'-GTTTGAGAAGTCTACTT-CAAAAATGAGATT-3' (SEQ ID NO: 20). Using primers consisting of sequences SEQ ID NO: 18, SEQ ID NO: 19 and SEQ ID NO: 20, detection of a cytosine (C) rather than a thymine (T) at position 151 of the sequence SEQ ID NO: 17, or a guanine (G) rather than an alanine (A) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0135] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056546 marker on chromosome 6. In particular, detection of the PE-0056546 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 21, or a fragment thereof including the [A/C] polymorphism at position 151 of SEQ ID NO: 21. For example, the forward primer for detecting the susceptible allele of the PE-0056546

marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 21 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTT-CATGCTCCTTCACTCACC GAATGTAACCT-3' (SEQ ID NO: 22), the forward primer for detecting the resistant allele of the PE-0056546 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 21 or its complementary sequence, may consist of the sequence 5'-GAAGGTGCGGAGTCAACGGATTCTTCACT-CACCGAATGTAACCG-3' (SEQ ID NO: 23), and the common reverse primer may consist of the sequence 5'-TTGGTCTACAAAGACATGCCAAATGGAA-3' (SEQ ID NO: 24). Using primers consisting of sequences SEQ ID NO: 22, SEQ ID NO: 23 and SEQ ID NO: 24, detection of a cytosine (C) rather than an alanine (A) at position 151 of the sequence SEQ ID NO: 21, or a guanine (G) rather than a thymine (T) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0136] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056547 marker on chromosome 6. In particular, detection of the PE-0056547 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 25, or a fragment thereof including the [A/C] polymorphism at position 151 of SEQ ID NO: 25. For example, the forward primer for detecting the susceptible allele of the PE-0056547 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 25 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCT-CAACTCTCCGGTGAATCTAAGCTTA-3' (SEQ ID NO: 26), the forward primer for detecting the resistant allele of the PE-0056547 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 25 or its complementary sequence, may consist of the sequence 5'-GAAGGTGCGGAGTCAACGGATTAACTCTCCGGT-GAATCTAAGCTTC-3' (SEQ ID NO: 27), and the common reverse primer may consist of the sequence 5'-CATTGT-CATAACCTTTACCAGACAATGAAT-3' (SEQ ID NO: 28). Using primers consisting of sequences SEQ ID NO: 26, SEQ ID NO: 27 and SEQ ID NO: 28, detection of a cytosine (C) rather than an alanine (A) at position 151 of the sequence SEQ ID NO: 25, or a guanine (G) rather than a thymine (T) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0137] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056549 marker on chromosome 6. In particular, detection of the PE-0056549 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 29, or a fragment thereof including the [T/A] polymorphism at position 151 of SEQ ID NO: 29. For example, the forward primer for detecting the susceptible allele of the PE-0056549 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 29 or its complementary sequence, may consist

of the sequence 5'-GAAGGTCGGAGTCAACGGATTCCGATATTTTATACTTAGCTATTCTAGCATAT-3' (SEQ ID NO: 30), the forward primer for detecting the resistant allele of the PE-0056549 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 29 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCTCCGATATTTTATACTTAGCTATTCTAGCATAA-3' (SEQ ID NO: 31), and the common reverse primer may consist of the sequence 5'-GAGTTTCATGATTCATCTGTTTTGTTTCCC-3' (SEQ ID NO: 32). Using primers consisting of sequences SEQ ID NO: 30, SEQ ID NO: 31 and SEQ ID NO: 32, detection of an alanine (A) rather than a thymine (T) at position 151 of the amplification product consisting of sequence SEQ ID NO: 29, or a thymine (T) rather than alanine (A) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0138] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0056550 marker on chromosome 6. In particular, detection of the PE-0056550 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 33, or a fragment thereof including the [C/T] polymorphism at position 151 of SEQ ID NO: 33. For example, the forward primer for detecting the susceptible allele of the PE-0056550 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 33 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCTACATTAATTTTCCAGCATTATTTGACATG-3' (SEQ ID NO: 34), the forward primer for detecting the resistant allele of the PE-0056550 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 33 or its complementary sequence, may consist of the sequence 5'-GAAGGTCGGAGTCAACGGATTGAACATTAATTTTCCAGCATTATTTGACATA-3' (SEQ ID NO: 35), and the common reverse primer may consist of the sequence

5'-TAGGTAATATTTGAGTTGCACTTATGGTAT-3' (SEQ ID NO: 36). Using primers consisting of sequences SEQ ID NO: 34, SEQ ID NO: 35 and SEQ ID NO: 36, detection of a thymine (T) rather than a cytosine (C) at position 151 of the sequence SEQ ID NO: 33, or an alanine (A) rather than a guanine (G) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0139] In some embodiments, said QTL on chromosome 6 conferring resistance to powdery mildew is identified by detection of the PE-0021476 marker on chromosome 6. In particular, detection of the PE-0021476 marker on chromosome 6 is performed using two forward primers, one being specific for the resistant allele and one being specific for the susceptible allele, and one common reverse primer. Said two forward primers may be selected so as to enable amplifying a nucleic acid comprising or consisting of SEQ ID NO: 37, or a fragment thereof including the [G/C] polymorphism at position 51 of SEQ ID NO: 37. For example, the forward primer for detecting the susceptible allele of the PE-0021476 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 37 or its complementary sequence, may consist of the sequence 5'-GAAGGTCGGAGTCAACGGATTGTTGGGATTTTAAATTGACTGACTAAAC-3' (SEQ ID NO: 38), the forward primer for detecting the resistant allele of the PE-0021476 marker, by amplifying a nucleic acid consisting of sequence SEQ ID NO: 37 or its complementary sequence, may consist of the sequence 5'-GAAGGTGACCAAGTTCATGCTTGTGGGATTTTAAATTGACTGACTAAAG-3' (SEQ ID NO: 39), and the common reverse primer may consist of the sequence 5'-CTCCAAAACCACAGCTACTAATTTGTT-3' (SEQ ID NO: 40). Using primers consisting of sequences SEQ ID NO: 38, SEQ ID NO: 39 and SEQ ID NO: 40, detection of a cytosine (C) rather than a guanine (G) at position 51 of the sequence SEQ ID NO: 37, or a guanine (G) rather than a cytosine (C) in the complementary strand, indicates the presence of the QTL on chromosome 6 conferring resistance to powdery mildew (see Table 1 below).

[0140] The alleles conferring the resistances to powdery mildew amplified by the markers defined here above are as described in Table 1.

TABLE 1

Markers linked to resistance to powdery mildew on chromosome 6, location and flanking sequences. R: Resistant, S: Susceptible							
Marker	Chromo- some	Position (CM334 v.1.55)	Sequence surrounding the marker	Position in SEQ ID NO	Type	Alternative alleles (R/S)	SEQ ID NO
PE-0014628	6	232 015 212	AAATACCTGCAATGGGTCGGGCTAGAAGAAATAAGCAGCAATAACCGC [A/G]TTCCAGAAAATAGATTGGTTGAGGAATTTGATCGAACACTTGGTG GGC	51	SNP	A/G	1
PE-0056542	6	233 673 276	ATTTCTAGGTTACTGGAATTTAACTCAGGAGAAAATGGATTTGTTTGGCAA TGCAATTTTCCGGGGTGCTAGTTTATAAAGACCTAAGATGTTTCAGTCA ACTGCTTAATAGACCTTTGACTGTTGGTTATGACTGGGCCCTT[T/C] GCCAAGATGGTTTGGACACAGAGCTTTACACACTTGTCTGCTATTTTG ATCTACTTGATCTTGAGTTGCGCTGTTTGTCTGTGCTGTTCAAAGCGACATT GACTGCTGAGATCTATAAAGATTCAATAGAAGATCCAAAATAATACT	151	SNP	C/T	5
PE-0056543	6	233 673 121	AGATGGTTTGGACAAACAGAGCTCTTACCACACTTGTCTGCTATTTTGATCT ACTTGATCTTGAGTTGGCTGTTTGTCTGTGCTGTTCAAGCGACATGACT GTCTGAGATCTATAAAGATTCATTAGAAGATCCAAATATATCTTTGA[G/A] CTATAGATGGAAGTTTCGNATATCTGACGGAAGGTTTCATTTGCAAGTTTG CTGGGATTCAGTCTGGCATCCTGATGAAAAATAAGGTTCTGAGTATCAGT CTTCTCGACATGGGTCTCAAGGGCAGGTTTCCACCTTGGTCTTCGGAAATTG	151	SNP	A/G	9
PE-0056544	6	233 672 530	CTGTTCTCGAGAGACTATGCCAATAATCCCGGACTGTGTGAGAACCG TTAGATCTTGGCAGGGTACTTCTAAGAAACCCCTACCGCAATCATTTCTC GGTGCAGCTGTTGGTGGGTAAGTTTAGGTGCTGTATTTTGGCCATCG [G/T]TATGTTCTCTATAGAGAGGATATCCAGGAGAGGAAGAGGATG ATGATCCTGAAGGGAATAAATGGCGAAGAGTATTAAGGGTAAAAAGCA ATCCAGGTAAGAATTTTAACCTCTCTGAATCAATAAAAACTGTCGTTTACT G	151	SNP	T/G	13
PE-0056545	6	233 671 922	TCTTTTCATAGAAGCATTTAACTAATGAATTTTCACTGCTCGCTGCTCG TGTGTGATTTTCTTGTTCAGTTTCTTAATTTGACTTAAGTTCTCTTATT TCAGTCTTCAATGTTTGAGAGTCTACTTCAAAAATGAGATTTAGT[C/GA] TCTTATGAAGGCCACTGACAGCTTCAACAAAAACAATATTATCGGCTCAG GAAGAACCGGACTTTTACAAAGCAGTACTTGATGATGGCATTCACTTA TGGTTAAGAGGTTGCAGAAATACTCAGCACTCAGAGAAAAGATTATG	151	SNP	C/T	17
PE-0056546	6	233 671 648	ACTCAGCACTCAGAGAAAAGGTTTATGTCGAGATGGCTACATTGGGAAA TGTAAGGATCGTAATCTGGTCTCTTTTAGGTTTCTGCAATGCCAAAAA AGAAAGGCTGTGTTGCTACAAGACATGCCAAATGGAACTTCGATGAT [A/C]GGTTACATTCGGTACTGAAGCGAAAACCTCTAGAGTGGCCTATAA GATTAAAAATAGGCATTGGAGCAGCCAAAGGATTTGTCATGGCTTCACCAC AACTGCAATCCTCGTATCAATTCACAGAAAATATTAGTTCTAAATGCATCTTG	151	SNP	C/A	21

TABLE 1-continued

Markers linked to resistance to powdery mildew on chromosome 6, location and flanking sequences. R: Resistant, S: Susceptible							
Marker	Chromo- some	Position (CW334 v.1.55)	Sequence surrounding the marker	Position in SEQ ID NO	Type	Alternative alleles (R/S)	SEQ ID NO
PE-0056547	6	233 671 204	GCTACTCCAAAAGGTGATGTGATACAGTTTCGGTGTGTTACTTCTCGAGTTA	151	SNP	C/A	25
			GTTACTGGTGAGAAACCGACCTCTGTCACCAAGCTCCGAGACCTTTAA				
			GGGGAATTTGGTGGAAATGGATCACAACTCCCGGTGAATCTAAGCTT				
			[A/C]AAGATCGGATGACCAATTCATTGCTCTGGTAAAGGTTATGACAAATGAGAT				
			CTTCCAGTCTCTAAAGTTGCTTGTGATGTGTTGTTCTCCTGTTCTCTAA				
GGAGAGCCGACAAATGTTGAAGTTTACCAGCTTCTGAGAGCCATCGGA	G						
PE-0056549	6	233 493 124	TAACTGATGGTTTCAGTTGAGCTGCAAAATGGAAACATCAATATCAGTAGCT	151	SNP	A/T	29
			TCTCGTTTCAACCAATTAATCTTCTTGTCTTTAGTCACTTGCACTTTAGAG				
			AAGAATCAGCTGATTCGCGATATTTATATCTTAGCTATTCTAGCATA[T/A]G				
			GGAAACAAACACAGATGAATCATGAACTCTTCGGTAGTAAAAAATATATG				
			TGCCAAAATGTTGGATGATCTTTTTATGGACGATGATCCATACCATTGTTT				
GGATTCAAGATCCACTTTTACGATTGCATCTCCAGCTCTTCCATT							
PE-0056550	6	233 489 782	TTTGATTTTGTAACTCATTTGCAATGAAATGATATGTTTATATGTTGTCTA	151	SNP	T/C	33
			GAAATATCAAACTATTACATTTATAGGTAATATTTGAGTTGCACCTATGGT				
			ATAGTGCATTTTATTTCTATTTATGTGACATTTATTTTATAGATA[C/T]ATGTC				
			GAAATAATGCTGAAAAATTAATGTTCAACTCTTTGAAATACATGAAAGACC				
			GAAACTTCTACCTATTTTTAAAGGATGGACGCGTGAATTTAGACCTAATTA				
AGAAATATTATCTTACACTTTTTTGTGTTTATCGAGAAGCTTTT							
PE-0021476	6	235 235 179	CTCCAAAACCAACAGCTACTAATTTGTTAGATCCAAAACCTTAAAAGCTTAA	51	SNP	C/G	37
[C/G]TTTAGTCACTCAATTAATAAATCCCAACAAATTTAAAAGCTAAGAGTGAAAG							

[0141] Accordingly, in some embodiments, the presence of the QTL on chromosome 6 conferring resistance to *Leveillula taurica* is characterized by the presence of:

- [0142] allele A of PE-0014628 (SEQ ID NO: 1);
- [0143] allele C of PE-0056542 (SEQ ID NO: 5);
- [0144] allele A of PE-0056543 (SEQ ID NO: 9);
- [0145] allele T of PE-0056544 (SEQ ID NO: 13);
- [0146] allele C of PE-0056545 (SEQ ID NO: 17);
- [0147] allele C of PE-0056546 (SEQ ID NO: 21);
- [0148] allele C of PE-0056547 (SEQ ID NO: 25);
- [0149] allele A of PE-0056549 (SEQ ID NO: 29);
- [0150] allele T of PE-0056550 (SEQ ID NO: 33); and/or
- [0151] allele C of PE-0021476 (SEQ ID NO: 37).

[0152] A plant of the invention which comprises said QTL on chromosome 6 preferably comprises any combination of the alleles as defined here above.

[0153] In particular, said QTL on chromosome 6 conferring said resistance to powdery mildew is characterized by at least one of the following allele combinations:

- [0154] Allele combination “A” comprising allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), and allele C of PE-0056546 (SEQ ID NO: 21);
- [0155] Allele combination “B” comprising allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21) and allele C of PE-0056547 (SEQ ID NO: 25);
- [0156] Allele combination “C” comprising allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), and allele T of PE-0056550 (SEQ ID NO: 33);
- [0157] Allele combination “D” comprising allele A of PE-0014628 (SEQ ID NO: 1), allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), allele T of PE-0056550 (SEQ ID NO: 33), and allele C of PE-0021476 (SEQ ID NO: 37); and
- [0158] Allele combination “E” comprising allele A of PE-0056549 (SEQ ID NO: 29) and allele T of PE-0056550 (SEQ ID NO: 33).

[0159] In some embodiments, said QTL on chromosome 6 conferring said resistance to powdery mildew is characterized by one of the allele combinations “A”, “B”, “C” or “D”, in particular by allele combination “A”. Alternatively or in addition, said QTL on chromosome 6 conferring said resistance to powdery mildew is characterized by allele combination “E”.

[0160] Alternatively or in addition, said QTL is characterized by the presence of at least one of:

- [0161] allele A of PE-0006634 (SEQ ID NO: 57)
- [0162] allele A of PE-0006718 (SEQ ID NO: 58)
- [0163] allele G of PE-0012892 (SEQ ID NO: 59)
- [0164] allele A of PE-0004191 (SEQ ID NO: 60)
- [0165] allele A of PE-0005575 (SEQ ID NO: 61)

[0166] allele A of PE-0006320 (SEQ ID NO: 62)

[0167] allele T of PE-0016499 (SEQ ID NO: 63)

[0168] allele A of PE-0013625 (SEQ ID NO: 64); and/or

[0169] allele A of PE-0022671 (SEQ ID NO: 65)

[0170] Said QTL on chromosome 6 can comprise at least one of said alleles, in particular more than one, such as 2, 3, 4, 5, 6, 7, 8 or 9 of said alleles, or all of said alleles.

[0171] In a preferred embodiment, said QTL is characterized by the presence of

[0172] allele A of PE-0006718 (SEQ ID NO: 58), and/or

[0173] allele G of PE-0012892 (SEQ ID NO: 59)

[0174] In one embodiment said QTL on chromosome 6 confers a resistance to powdery mildew, in particular a moderate to high resistance to powdery mildew, in a plant which does not comprise the QTL of resistance to *Leveillula taurica* on chromosome 4 as described in International patent publication WO2013/033210 and/or which does not comprise the QTL of resistance to *Leveillula taurica* on linkage group LG1/8 as described in International Patent publication WO2014/140331.

[0175] In one embodiment, the plant of the invention does not comprise the QTL of resistance to *Leveillula taurica* on chromosome 4 as described in International patent publication WO2013/033210. Alternatively or in addition, the plant according to the invention does not comprise the QTL of resistance to *Leveillula taurica* on linkage group LG1/8 as described in International Patent publication WO2014/140331.

[0176] The inventors have identified two genes CA06g26150 and CA06g26250, located within the QTL on chromosome 6 conferring resistance to powdery mildew. Said genes have been identified as candidate genes which may confer the resistance to powdery mildew.

[0177] In some embodiments, said QTL on chromosome 6 co-segregates with the gene CA06g26150. CA06g26150 encodes a MLO-like protein. CA06g26150 is located between positions 233490196 and 233495756 of chromosome 6 of pepper genome *Capsicum annuum* cv. CM334 v.1.55.

[0178] In some embodiments, said resistance to powdery mildew is linked to the CA06g26150 gene. In particular, said resistance to powdery mildew is caused by at least one mutation within the CA06g26150 gene, in comparison with the CA06g26150 gene of a susceptible *Capsicum* plant and/or line. A coding sequence of the CA06g26150 gene of a susceptible plant (Maor) is set forth in SEQ ID NO: 41. The mutation may consist in at least one nucleotide substitution, insertion or deletion in the sequence of the CA06g26150 gene, in particular its coding sequence as set forth in SEQ ID NO: 41, including the deletion of the full gene or a fragment thereof. Preferably, the mutation is a loss-of-function mutation. The mutation may induce one or more amino acid substitutions in the sequence of the MLO-like protein encoded by the CA06g26150 gene, and impair the function of the MLO-like protein. In one embodiment, the loss-of-function mutation in the CA06g26150 gene is a null mutation. A null mutation prevents expression of an active MLO-like protein, for instance by causing a premature stop in the translation of the mRNA into a protein, resulting into the expression of a truncated form of the MLO-like protein. In one embodiment, the CA06g26150

gene of a *Capsicum* plant according to the invention comprises allele A of marker PE-0056549 (SEQ ID NO: 29).

[0179] In one embodiment, the present invention thus relates to a *Capsicum* plant resistant to powdery mildew, wherein said plant comprises a mutant allele of the CA06g26150 gene encoding a MLO-like protein, wherein said mutant allele confers said resistance.

[0180] In some embodiments, said QTL on chromosome 6 co-segregates with the gene CA06g26250. CA06g26250 encodes a leucine-rich repeat receptor-like protein kinase. CA06g26250 is located between positions 233670962 and 233673290 of chromosome 6 of pepper genome *Capsicum annuum* cv. CM334 v.1.55.

[0181] In some embodiments, said resistance to powdery mildew is linked to the CA06g26250 gene. In particular, said resistance to powdery mildew is caused by at least one mutation within the CA06g26250 gene, in comparison with the CA06g26250 gene of a susceptible *Capsicum* plant and/or line. A coding sequence of the CA06g26250 gene of a susceptible plant (Maor) is set forth in SEQ ID NO: 42. The mutation may consist in at least one nucleotide substitution, insertion or deletion in the sequence of the CA06g26250 gene, in particular its coding sequence as set forth in SEQ ID NO: 42, including the deletion of the full gene or a fragment thereof. Preferably, the mutation is a loss-of-function mutation. The mutation may induce one or more amino acid substitutions in the sequence of the leucine-rich repeat receptor-like protein kinase encoded by the CA06g26250 gene, and impair the function of the leucine-rich repeat receptor-like protein kinase. In one embodiment, the loss-of-function mutation in the CA06g26250 gene is a null mutation. A null mutation prevents expression of an active leucine-rich repeat receptor-like protein kinase, for instance by causing a premature stop in the translation of the mRNA into a protein, resulting into the expression of a truncated form of the leucine-rich repeat receptor-like protein kinase.

[0182] In one embodiment, the CA06g26250 gene of a *Capsicum* plant according to the invention comprises one or more of allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21) and allele C of PE-0056547 (SEQ ID NO: 25), in particular one or more of allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17) and allele C of PE-0056546 (SEQ ID NO: 21).

[0183] In one embodiment, the present invention thus relates to a *Capsicum* plant resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*, wherein said plant comprises a mutant allele of the CA06g26250 gene encoding a leucine-rich repeat receptor-like protein kinase, wherein said mutant allele confers said resistance.

[0184] In some embodiments, said QTL on chromosome 6 co-segregates with the gene CA06g26150 and/or the gene CA06g26250.

[0185] Genes CA06g26150 and CA06g26250 are annotated in the pepper genome *Capsicum annuum* cv. CM334 v.1.55. CA06g26150 and CA06g26250 correspond, respectively to LOC107874591 and LOC107875376, as annotated in the pepper genome *Capsicum annuum* cv. Zunla v 1.0.

[0186] In some embodiments, said *Capsicum* plant according to the invention is a plant of *C. annuum* line ST17-1157 which seeds are deposited at NCIMB accession number NCIMB 43602.

[0187] In some embodiments, a plant according to the invention may be a progeny or offspring of a plant grown from the deposited seeds of *C. annuum* line ST17-1157, deposited at the NCIMB under the accession number NCIMB 43602. Plants grown from the deposited seeds are homozygously resistant to powdery mildew. They can be used to transfer said QTL on chromosome 6 in another background by crossing and selfing and/or backcrossing. A progeny of a plant obtained from the deposited seed can be identified by one skilled in the art, for example by using one or more of the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37). Preferably, such a progeny is identified by at least 2, in particular at least 3 of said markers. Such progeny can also be identified by using one or more of the markers PE-0006634 (SEQ ID NO: 57), PE-0006718 (SEQ ID NO: 58), PE-0012892 (SEQ ID NO: 59), PE-0004191 (SEQ ID NO: 60), PE-0005575 (SEQ ID NO: 61), PE-0006320 (SEQ ID NO: 62), PE-0016499 (SEQ ID NO: 63), PE-0013625 (SEQ ID NO: 64), and PE-0022671 (SEQ ID NO: 65), in particular markers PE-0006718 (SEQ ID NO: 58) and/or PE-0012892 (SEQ ID NO: 59). Sequences and alleles for these markers are provided in the present specification.

[0188] The resistance to powdery mildew is advantageously determined by comparison to a susceptible line, preferably a susceptible commercial line, for example one or more of the Maor, CPNT, Vania or BYP lines, in particular from the Maor line. The resistance is preferably determined as detailed in example 2, on the basis of an inoculation test of the plant. The test can be a test on plantlets, or a test on an adult plant.

[0189] In one embodiment, the plant according to the invention shows a sporulation covering less than 10% of the leaves surface, preferably less than 5% of the leaves surface, still preferably no sporulation on the leaves in conditions of infection by *Leveillula taurica*, particularly in the same environmental conditions in which a susceptible *Capsicum* plant shows a sporulation covering more than 60% of the leave surfaces, preferably more than 70% of the leaves surface, still preferably more than 75% of the leaves surface. The sporulation rate can be assessed on the entire plant or on only a part of the plant, e.g. the bottom part or the medium part of the plant. The susceptible *Capsicum* plant may be a susceptible (commercial) line, for example one or more of the Maor, CPNT, Vania or BYP lines, in particular from the Maor line. In one embodiment, these properties are assessed on a population of at least 5, in particular at least 10, more particularly at least 50, even more particularly at least 100 plants.

[0190] The *Capsicum* plant according to the invention can be from any species within the *Capsicum* genus. In particular, it may be a *Capsicum annuum*, *Capsicum baccatum*, *Capsicum frutescens*, *Capsicum chinense*, *Capsicum pubescens* or *Capsicum chacoense* plant. Preferably, the plant according to the invention is a *Capsicum annuum* plant, more preferably a sweet pepper or a hot pepper, even more

preferably a bell pepper. The *Capsicum* plant can also be from any type. In particular, the *Capsicum* plant may be of one of the following types: Dulce Italiano, Lamuyo, Blocky Florida, Open field and blocky protected, Ancho, Anaheim, Marconi, Jalapeno, Cayenne, Charleston or Sivri.

[0191] In one embodiment, the QTL on chromosome 6 according to the invention provides resistance to more than one strains of *Leveillula taurica*.

[0192] The invention also relates to a population of *Capsicum* plants according to the invention, wherein said population comprises at least 5 plants, in particular at least 10 plants, more particularly at least 20 plants, even more particularly at least 50 or 100, or more particularly at least 1000 plants.

[0193] According to a second aspect, the present invention is directed to one or more parts of a plant according to the invention.

[0194] All the embodiments detailed in the preceding section in connection with the first aspect of the invention are also embodiments according to this second aspect of the invention. In particular, the different features of said QTL that have been defined in relation with the above aspects of the invention apply mutatis mutandis to this aspect of the invention.

[0195] In some embodiments, said QTL is present on chromosome 6 in the genome of a seed of *Capsicum annum* ST17-1157, deposited at NCIMB accession number NCIMB 43602.

[0196] In some embodiments, a plant part according to the invention comprises a QTL conferring resistance to powdery mildew is located on chromosome 6 within the chromosomal region delimited by PE-0014628 (SEQ ID NO: 1) and PE-0021476 (SEQ ID NO: 37).

[0197] In some embodiments, a plant part according to the invention comprises one or more alleles conferring resistance to powdery mildew as described in Table 1 and/or Table 12.

[0198] In some embodiments, the plant part according to the invention comprises any one of the allele combinations "A" to "E" as defined in the first aspect of the invention.

[0199] In some embodiments, the combination of alleles as described here above is as found in the genome of a plant corresponding to the deposited material ST17-1157 (NCIMB accession number NCIMB 43602).

[0200] According to another embodiment, the plant part is an embryo, protoplast, meristematic cell, callus, pollen, leaf, anther, stem, petiole, root, root tip, fruit, seed, flower, cotyledon, and/or hypocotyl.

[0201] In some embodiments, a part of a plant is a plant cell. The invention thus provides a cell of a *Capsicum* plant according to the invention, i.e. a plant cell that comprises comprising introgressed in its genome a quantitative trait locus (QTL) on chromosome 6, conferring resistance to powdery mildew, wherein said QTL is present on chromosome 6 and is as defined in the present specification. The different features of said QTL that have been defined in relation with the above aspects of the invention apply mutatis mutandis to this aspect of the invention.

[0202] A plant cell of the invention may have the capacity to be regenerated into a whole plant.

[0203] Alternatively, the invention is also directed to plant cells which are not regenerable, and thus are not capable of giving rise to a whole plant.

[0204] Preferably, a plant cell according to the invention is derived from an embryo, protoplast, meristematic cell, callus, pollen, leaves, anther, stem, petiole, root, root tip, fruit, seed, flower, cotyledon, or hypocotyl.

[0205] Preferably, a plant cell according to the invention comprises a QTL conferring resistance to powdery mildew is located on chromosome 6 within the chromosomal region delimited by PE-0014628 (SEQ ID NO: 1) and PE-0021476 (SEQ ID NO: 37).

[0206] The invention is more particularly concerned with a seed of a *Capsicum* plant, giving rise when grown up to a *Capsicum* plant resistant to powdery mildew as defined above. Such seed are thus 'seed of a plant of the invention', i.e. seed giving rise to a plant of the invention. The invention is also concerned with seed from a plant of the invention, i.e. obtained from such a plant after selfing or crossing, provided however that the plant obtained from said seed is resistant to powdery mildew due to the QTL on chromosome 6 as defined here above conferring said resistance.

[0207] The invention also relates to a population of *Capsicum* seeds according to the invention, wherein said population comprises at least 2 seeds, especially at least 10 seeds, particularly at least 100 seeds, even more particularly at least 1000 seeds.

[0208] The present invention is also directed to a tissue culture of regenerable cells of the plant as defined above according invention, and thus comprises in their genome the QTL on chromosome 6 conferring resistance to powdery mildew as described here above.

[0209] The tissue culture will preferably be capable of regenerating plants having the physiological and morphological charac [text missing or illegible when filed] invention.

[0210] The invention also provides a protoplast of the plant defined above, or from the tissue culture defined above, said proto [text missing or illegible when filed]

[0211] All the embodiments detailed above in connection with the first aspect of the invention are also embodiments according [text missing or illegible when filed]

[0212] According to another aspect, the present invention is also directed to the use of a *Capsicum* plant as detailed according to the first aspect of the invention, resistant to powdery mildew, as a breeding partner in a breeding program for obtaining *Capsicum* plants resistant to powdery mildew. Indeed, such a *Capsicum* plant according to the first aspect harbors in its genome the QTL as defined here above conferring said resistance. By crossing this plant with susceptible or less resistant plants, it is thus possible to transfer this QTL, conferring the desired phenotype, to the progeny. A plant according to the invention can thus be used as a breeding partner for introgressing the QTL conferring the desired phenotype into a *Capsicum* plant or germplasm. The invention is also directed to the same use with plants or seeds of ST17-1157 as deposited at NCIMB under accession number NCIMB 43602. Said plants are also suitable as introgression partners in a breeding program aiming at conferring the desired phenotype to a *Capsicum* plant or germplasm.

[0213] In such a breeding program, the selection of the progeny displaying the desired phenotype, or bearing sequences linked to the desired phenotype, can advantageously be carried out on the basis of the allele of the markers disclosed here above. The progeny is preferably selected on the presence of one or more of the following specific alleles: allele A of PE-0014628 (SEQ ID NO: 1);

allele C of PE-0056542 (SEQ ID NO: 5); allele A of PE-0056543 (SEQ ID NO: 9); allele T of PE-0056544 (SEQ ID NO: 13); allele C of PE-0056545 (SEQ ID NO: 17); allele C of PE-0056546 (SEQ ID NO: 21); allele C of PE-0056547 (SEQ ID NO: 25); allele A of PE-0056549 (SEQ ID NO: 29); allele T of PE-0056550 (SEQ ID NO: 33); and/or allele C of PE-0021476 (SEQ ID NO: 37). The progeny is preferably selected on the presence of one or more of the allele combinations “A” to “E” as defined in the first aspect of the invention. The progeny may also be selected based on the presence of one or more of allele A of PE-0006634 (SEQ ID NO: 57), allele A of PE-0006718 (SEQ ID NO: 58), allele G of PE-0012892 (SEQ ID NO: 59), allele A of PE-0004191 (SEQ ID NO: 60), allele A of PE-0005575 (SEQ ID NO: 61), allele A of PE-0006320 (SEQ ID NO: 62), allele T of PE-0016499 (SEQ ID NO: 63), allele A of PE-0013625 (SEQ ID NO: 64) and/or allele A of PE-0022671 (SEQ ID NO: 65), in particular allele A of PE-0006718 (SEQ ID NO: 58) and/or allele G of PE-0012892 (SEQ ID NO: 59).

[0214] The selection of the progeny having the desired phenotype can also be made on conditions of a *Leveillula taurica* infection, as disclosed inter alia in Example 2, or with other tests well-known to the skilled reader.

[0215] A plant according to the invention, in particular a plant as deposited under accession number NCIMB 43602, is thus particularly valuable in a marker assisted selection program for obtaining commercial *Capsicum* lines and varieties resistant to powdery mildew.

[0216] The invention is also directed to the use of said plants in a program aiming at identifying, sequencing and/or cloning the genetic sequences conferring the desired phenotype.

[0217] Any specific embodiment described for the previous aspect of the invention is also applicable to this aspect of the invention, especially with regard to the features of the QTL conferring the phenotype of interest.

[0218] According to another aspect, the invention also concerns methods for the production of *Capsicum* plants resistant to powdery mildew, especially commercial plants. A method or process for the production of a plant having these features comprises the following steps:

- [0219] (a) crossing a plant according to the first aspect of the invention (e.g. a plant corresponding to the deposited seeds (NCIMB 43602)) and a susceptible or less resistant *Capsicum* plant, in which the desired phenotype is to be imported or improved,
 - [0220] (b) selecting one plant resistant to powdery mildew in the progeny thus obtained, or one plant bearing the QTL on chromosome 6 as defined in the present invention conferring a resistance to powdery mildew,
 - [0221] (c) optionally self-pollinating one or several times the resistant plant obtained at step b) and selecting a plant resistant to powdery mildew,
 - [0222] (d) backcrossing the resistant plant selected in step b) or c) with a *Capsicum* plant susceptible to powdery mildew
 - [0223] (e) selecting a plant resistant to powdery mildew.
- [0224] Alternatively, the method or process may comprise the following steps:
- [0225] (a1) crossing a plant according to the first aspect of the invention (e.g. a plant corresponding to the deposited se

[0226] (a2) selfing the F1 population to create F2 population,

[0227] (b) selecting resistant individuals in the progeny thus obtained,

[0228] (c) optionally self-pollinating one or several times the resistant plant obtained at step b) and selecting a resistant plant in the progeny thus obtained,

[0229] (d) backcrossing the resistant plant selected in step b) or c) with a *Capsicum* plant susceptible to powdery mildew,

[0230] (e) selecting a plant resistant to powdery mildew.

[0231] In some embodiments, the plant resistant to powdery mildew caused by *Leveillula taurica* can be selected at steps b), c) and e).

[0232] The plant selected at step e) is preferably a commercial plant. In some embodiments, the selected plant is a bell pepper. In some embodiments, the selected plant is one of the following types Dulce Italiano, Lamuyo and blocky in China, Blocky Florida, Open field and blocky protected, Ancho, Anaheim, Marconi, Jalapeno, Cayenne, Charleston or Sivri.

[0233] Preferably, steps d) and e) are repeated at least twice and preferably three times, not necessarily with the same susceptible *Capsicum* plant. Said susceptible *Capsicum* plant is preferably a breeding line.

[0234] The self-pollination and backcrossing steps may be carried out in any order and can be intercalated, for example a backcross can be carried out before and after one or several self-pollinations, and self-pollinations can be envisaged before and after one or several backcrosses.

[0235] In some embodiments, such a method is advantageously carried out by using markers as described here above for one or more of the selections carried out at steps b), c) and/or e) for selecting plants resistant to powdery mildew.

[0236] In some embodiments, the markers for selecting plants resistant to powdery mildew are one or more of the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37). Further markers which can be used are one or more of PE-0006634 (SEQ ID NO: 57), PE-0006718 (SEQ ID NO: 58), PE-0012892 (SEQ ID NO: 59), PE-0004191 (SEQ ID NO: 60), PE-0005575 (SEQ ID NO: 61), PE-0006320 (SEQ ID NO: 62), PE-0016499 (SEQ ID NO: 63), PE-0013625 (SEQ ID NO: 64), and PE-0022671 (SEQ ID NO: 65), in particular markers PE-0006718 (SEQ ID NO: 58) and/or PE-0012892 (SEQ ID NO: 59).

[0237] In some embodiments, the plant selected at any one of steps b), c) and/or e) is preferably selected on the presence of one of the allele combinations “A” to “E” as defined in the first aspect of the invention.

[0238] The selection carried out at steps b), c) and/or e) can also be made using any type of genetic marker, in particular restriction fragment length polymorphisms (RFLPs), amplified fragment length polymorphisms (AFLPs), simple sequence repeats (SSRs), simple sequence length polymorphisms (SSLPs), single nucleotide polymorphisms (SNPs), insertion/deletion polymorphisms (Indels), variable number tandem repeats (VNTRs), and random

amplified polymorphic DNA (RAPD), isozymes, and other markers known to those skilled in the art.

[0239] The selection of the progeny having the desired phenotype can also be made on conditions of disease infection, as disclosed inter alia in the Examples or with other tests well-known to the skilled reader.

[0240] The method used for allele detection can be based on any technique allowing the distinction between two different alleles of a marker, on a specific chromosome. Detection of a polymorphism can be made by electrophoretic techniques including a single strand conformational polymorphism (Orita, et al. (1989) *Genomics*, 8(2), 271-278), denaturing gradient gel electrophoresis (Myers (1985) EPO 0273085), or cleavage fragment length polymorphisms (Life Technologies, Inc., Gaithersburg, Md.), but the widespread availability of DNA sequencing often makes it easier to simply sequence amplified products directly. Once the polymorphic sequence difference is known, rapid assays for the detection of a polymorphism can be designed for progeny testing, generally involving some version of PCR amplification of specific alleles (PASA; Sommer, et al. (1992) *Biotechniques* 12(1), 82-87), or PCR amplification of multiple specific alleles (PAMSA; Dutton and Sommer (1991) *Biotechniques*, 11(6), 700-7002). In particular examples, PCR detection and quantification is carried out using two labeled fluorogenic oligonucleotide forward primers and an unlabeled common reverse primer, for example, KASPar™ (KBiosciences). Detection of a polymorphism can also be made by electrophoretic techniques including a single strand conformational polymorphism (Orita, et al. (1989) *Genomics*, 8(2), 271-278), denaturing gradient gel electrophoresis (Myers (1985) EPO 0273085), or cleavage fragment length polymorphisms (Life Technologies, Inc., Gaithersburg, Md.). The widespread availability of DNA sequencing often also enables to sequence amplified products directly.

[0241] The present invention also concerns a *Capsicum* plant obtained or obtainable by the methods described herein. Such a plant is indeed a *Capsicum* plant that is resistant to powdery mildew according to the first aspect of the invention.

[0242] According to a further aspect, the present invention is also directed to hybrid *Capsicum* plants obtainable by crossing a resistant plant according to the first aspect of the invention, such as a plant ST17-1157, a representative sample of seeds which have been deposited under the NCIMB accession number NCIMB 43602, or a resistant plant obtainable by the methods disclosed above, with a second *Capsicum* plant, for example a plant susceptible to powdery mildew infection, or a plant with a different, e.g. a lower, level of resistance to powdery mildew infection. A particularly preferred hybrid *Capsicum* plant, is a plant which displays any trait or phenotype of agronomical interest.

[0243] The invention is also directed to a method for obtaining commercial *Capsicum* plants that are resistant to powdery mildew, said method comprising the steps of:

[0244] backcrossing a plant obtained by germinating the deposited seeds ST17-1157 (NCIMB accession number NCIMB 43602) or a *Capsicum* plant according to the first aspect of the invention, with a *Capsicum* plant, for example a *Capsicum* plant susceptible to powdery mildew,

[0245] selecting a plant resistant to powdery mildew in the progeny thus obtained.

[0246] The selection in the second step is preferably carried out as detailed above for the other methods of the invention. Said selection is preferably carried out on the presence of one or more of the specific alleles of the markers for the QTL on chromosome 6 as described here above, as found in line ST17-1157.

[0247] The plant selected is preferably a commercial variety, cultivar or type of pepper. In some embodiments, the selected plant is a bell pepper. In some embodiments, the selected plant is one of the following types Dulce Italiano, Lamuyo and blocky in China, Blocky Florida, Open field and blocky protected, Ancho, Anaheim, Marconi, Jalapeno, Cayenne, Charleston or Sivri.

[0248] Also provided are methods for producing *Capsicum* plants seeds. In some embodiments, the methods comprise crossing the *Capsicum* plant according to the invention with itself or with another *Capsicum* plant, and harvesting the resultant seeds.

[0249] In addition to introgression of the QTL associated to resistance to powdery mildew, as detailed in the methods of the invention, said sequences can also be introduced into *Capsicum* background by genetic engineering in order to obtain a commercial *Capsicum* plant resistant to powdery mildew. The identification and cloning of the introgressed QTL from *Capsicum* conferring the desired phenotype, inter alia from the deposit, are routine for the skilled person.

[0250] It is noted that the seeds or plants of the invention may be obtained by different processes, in particular technical processes such as mutagenesis, e.g. chemical mutagenesis or UV mutagenesis, or genetic engineering such as guided recombination, and are not exclusively obtained by means of an essentially biological process.

[0251] According to such an aspect, the invention relates to a *Capsicum* plant or seed, preferably a non-naturally occurring *Capsicum* plant or seed, which may comprise one or more mutations in its genome, which provides the mutant plant a resistance to powdery mildew, which mutation is as present, for example, in the genome of plants of which a representative sample was deposited with the NCIMB under deposit number NCIMB 43602.

[0252] The mutations can have a natural cause (spontaneous mutations) or can be induced via methods such as mutagenesis. Mutagenesis methods are known in the art and include chemical mutagenesis using ethyl methanesulfonate (EMS). Other chemical mutagenic agents include but are not limited to, diethyl sulfate (des), ethyleneimine (ei), propane sultone, N-methyl-N-nitrosourea (mnu), N-nitroso-N-methylurea (NMU), N-ethyl-N-nitrosourea (enu), and sodium azide.

[0253] Alternatively, the mutations can be induced by means of irradiation, which is for example selected from x-rays, fast neutrons, UV radiation.

[0254] Mutagenesis techniques can be followed by an identification method such as TILLING. TILLING (Targeting Induced Local Lesions IN Genomes) is a general reverse genetics technique that uses traditional chemical mutagenesis methods to create libraries of mutagenized individuals that are later subjected to high throughput screens for the discovery of mutations. TILLING combines chemical mutagenesis with mutation screens of pooled PCR products, resulting in the isolation of missense and non-sense mutant alleles of the targeted genes. Thus, TILLING uses traditional chemical mutagenesis (e.g. EMS or MNU mutagenesis) or other mutagenesis methods (e.g. radiation such as UV)

followed by high-throughput screening for mutations in specific target genes, such as *cl*, *wt* and/or *y* according to the invention. S1 nucleases, such as CEL1 or ENDO1, are used to cleave heteroduplexes of mutant and wild type target DNA and detection of cleavage products using e.g. electrophoresis such as a LI-COR gel analyzer system, see e.g. Henikoff et al. *Plant Physiology* 2004, 135: 630-636. TILLING has been applied in many plant species, including pepper (Kang, H. S., Kim, S. H., Lee, S. W. et al. *Hortic. Environ. Biotechnol.* (2018) 59: 447). Also EcoTILLING, whereby mutants in natural populations are detected, has been widely used, see Till et al. 2006 (*Nat Protoc* 1: 2465-77) and Comai et al. 2004 (*Plant J* 37: 778-86).

[0255] Preferably, the mutations are the integration of one QTL conferring resistance to powdery mildew, wherein said at least one QTL is present on chromosome 6, in replacement of the homologous sequences of a *Capsicum* plant. Even more preferably, the mutation is the substitution of the sequence delimited by marker PE-0014628 (SEQ ID NO: 1) and marker PE-0021476 (SEQ ID NO: 37) on chromosome 6 of a *Capsicum* genome, or a fragment thereof, by the homologous sequence on chromosome 6 present in the genome of a plant of which a representative sample was deposited with the NCIMB under deposit number NCIMB 43602, wherein the sequences or fragments thereof confer when combined resistance to powdery mildew.

[0256] In an embodiment, the invention relates to a method for obtaining a *Capsicum* plant or seed carrying one or more mutations in its genome, which provides the plant with a resistance to powdery mildew. Such a method is illustrated in Example 12 and may comprise:

- [0257] a) treating M0 seeds of a *Capsicum* plant to be modified with a mutagenic agent to obtain M1 seeds;
- [0258] b) growing plants from the thus obtained M1 seeds to obtain M1 plants;
- [0259] c) producing M2 seeds by self-fertilisation of M1 plants; and
- [0260] d) optionally repeating step b) and c) *n* times to obtain M2+n seeds.

[0261] The M2+n seeds are grown into plants and submitted to *Leveillula taurica* infection. The surviving plants, or those with the milder symptoms of *Leveillula taurica* infection, are multiplied one or more further generations while continuing to be selected for their resistance to powdery mildew. In this method, the M1 seeds of step a) can be obtained via chemical mutagenesis such as EMS mutagenesis. Other chemical mutagenic agents include but are not limited to, diethyl sulfate (des), ethyleneimine (ei), propane sultone, N-methyl-N-nitrosourea (mnu), N-nitroso-N-methylurea (NMU), N-ethyl-N-nitrosourea (enu), and sodium azide. Alternatively, the mutations are induced by means of irradiation, which is for example selected from x-rays, fast neutrons, UV radiation.

[0262] In another embodiment of the invention, the mutations are induced by means of genetic engineering. Such mutations also include the integration of sequences conferring the resistance to powdery mildew, as well as the substitution of residing sequences by alternative sequences conferring the resistance to powdery mildew.

[0263] The genetic engineering means which can be used include the use of all such techniques called New Breeding Techniques which are various new technologies developed and/or used to create new characteristics in plants through genetic variation, the aim being targeted mutagenesis, tar-

geted introduction of new genes or gene silencing (RdDM). Example of such new breeding techniques are targeted sequence changes facilitated thru the use of Zinc finger nuclease (ZFN) technology (ZFN-1, ZFN-2 and ZFN-3, see U.S. Pat. No. 9,145,565, incorporated by reference in its entirety), Oligonucleotide directed mutagenesis (ODM), Cisgenesis and intragenesis, RNA-dependent DNA methylation (RdDM, which does not necessarily change nucleotide sequence but can change the biological activity of the sequence), Grafting (on GM rootstock), Reverse breeding, Agro-infiltration (agro-infiltration “sensu stricto”, agro-inoculation, floral dip), Transcription Activator-Like Effector Nucleases (TALENs, see U.S. Pat. Nos. 8,586,363 and 9,181,535, incorporated by reference in their entireties), the CRISPR/Cas system (see U.S. Pat. Nos. 8,697,359; 8,771,945; 8,795,965; 8,865,406; 8,871,445; 8,889,356; 8,895,308; 8,906,616; 8,932,814; 8,945,839; 8,993,233; and 8,999,641, which are all hereby incorporated by reference), engineered meganuclease re-engineered homing endonucleases, DNA guided genome editing (Gao et al., *Nature Biotechnology* (2016), doi: 10.1038/nbt.3547, incorporated by reference in its entirety), and Synthetic 5 genomics). A major part of today’s targeted genome editing, another designation for New Breeding Techniques, is the applications to induce a DNA double strand break (DSB) at a selected location in the genome where the modification is intended. Directed repair of the DSB allows for targeted genome editing. Such applications can be utilized to generate mutations (e.g., targeted mutations or precise native gene editing) as well as precise insertion of genes (e.g., cisgenes, intragenes, or transgenes). The applications leading to mutations are often identified as site-directed nuclease (SDN) technology, such as SDN1, SDN2 and SDN3. For SDN1, the outcome is a targeted, non-specific genetic deletion mutation: the position of the DNA DSB is precisely selected, but the DNA repair by the host cell is random and results in small nucleotide deletions, additions or substitutions. For SDN2, a SDN is used to generate a targeted DSB and a DNA repair template (a short DNA sequence identical to the targeted DSB DNA sequence except for one or a few nucleotide changes) is used to repair the DSB: this results in a targeted and predetermined point mutation in the desired gene of interest. As to the SDN3, the SDN is used along with a DNA repair template that contains new DNA sequence (e.g. gene). The outcome of the technology would be the integration of that DNA sequence into the plant genome. The most likely application illustrating the use of SDN3 would be the insertion of cisgenic, intragenic, or transgenic expression cassettes at a selected genome location. A complete description of each of these techniques can be found in the report made by the Joint Research Center (JRC) Institute for Prospective Technological Studies of the European Commission in 2011 and titled “New plant breeding techniques—State-of-the-art and prospects for commercial development”, which is incorporated by reference in its entirety.

[0264] The present invention also provides methods for detecting and/or selecting a *Capsicum* plant that is resistant to powdery mildew, wherein said method comprises the step of detecting the presence of at least one QTL conferring resistance to powdery mildew, wherein said at least one QTL is present on chromosome 6.

[0265] Preferably, said QTL conferring resistance to powdery mildew that is present on chromosome 6 is located

within a chromosomal region that is delimited by marker PE-0014628 and PE-0021476. In some embodiments, said QTL that is present on chromosome 6 can be identified by amplifying any one of the following markers: PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37); or any other markers within the chromosomal region delimited by markers PE-0014628 (SEQ ID NO: 1) and PE-0021476 (SEQ ID NO: 37). Additionally, or alternatively, said QTL that is present on chromosome 6 can be identified by amplifying any one of the following markers: PE-0006634 (SEQ ID NO: 57), PE-0006718 (SEQ ID NO: 58), PE-0012892 (SEQ ID NO: 59), PE-0004191 (SEQ ID NO: 60), PE-0005575 (SEQ ID NO: 61), PE-0006320 (SEQ ID NO: 62), PE-0016499 (SEQ ID NO: 63), PE-0013625 (SEQ ID NO: 64) and PE-0022671 (SEQ ID NO: 65), in particular markers PE-0006718 (SEQ ID NO: 58) and/or PE-0012892 (SEQ ID NO: 59).

[0266] In some embodiments, a plant is selected if any one of the allele combinations “A” to “E”, as defined in the first aspect material sample of the plant to be selected.

[0267] In some embodiments, detection of the markers described in the application, in particular one or more of PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37), or one or more of PE-0006634 (SEQ ID NO: 57), PE-0006718 (SEQ ID NO: 58), PE-0012892 (SEQ ID NO: 59), PE-0004191 (SEQ ID NO: 60), PE-0005575 (SEQ ID NO: 61), PE-0006320 (SEQ ID NO: 62), PE-0016499 (SEQ ID NO: 63), PE-0013625 (SEQ ID NO: 64) and PE-0022671 (SEQ ID NO: 65), is performed by amplification, e.g. by PCR, using, for each marker, one forward primer which can be used for amplifying the resistant allele, one forward primer which can be used for amplifying the susceptible allele and one common reverse primer, for example using the KASPar (KBiosciences) technology. In particular, the primers for amplifying each of said markers may have the sequences as described in the first aspect of the invention, and detailed in table 11.

[0268] In a preferred embodiment, the amplification is as described in the examples. In a still preferred embodiment, the amplification is performed using a two-step touchdown method in which the elongation and annealing steps are incorporated into a single step. The temperature used for the annealing stage determines the specificity of the reaction and hence the ability of the primers to anneal to the DNA template. A touchdown PCR involves a first step of Taq polymerase activation, followed by a second step called the touchdown step that involves a high annealing temperature and incrementally decreasing the annealing temperature in each PCR cycle, and a third step of DNA amplification. The higher annealing temperatures in the early cycles of a touchdown ensure that only very specific base pairing will occur between the DNA and the primer, hence the first sequence to be amplified is most likely to be the sequence of interest. The annealing temperature is gradually decreased to increase the efficiency of the reaction. The regions that were originally amplified during the highly specific early touch-

down cycles will be further amplified and outcompete any non-specific amplification that may occur at the lower temperatures.

[0269] In another embodiment, the amplification of SNP markers is performed as recommended in the KASPar assay and illustrated in the examples, namely by PCR cycles, comprising a first denaturation step at 94° C. during around 15 minutes, at least 10 cycles of around 20 seconds at 94° C. followed by around 60 second at a decreasing temperature from 65° C. for the 1st cycle to 57° C. for the last cycle, and around 35 cycles of around 20 seconds at 94° C. followed by around 60 seconds at 57° C. This protocol can easily be adapted by a skilled person, depending on the type of primers used.

[0270] According to a further aspect, the present invention also provides molecular markers that are linked to the QTL on chromosome 6 as defined here above conferring the resistance to powdery mildew.

[0271] In some embodiments, said molecular markers linked to the QTL conferring the resistance to powdery mildew on chromosome 6 are any one of the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37), or all the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37) such as a combination corresponding to any one of allele combinations A to E, or any other markers within the chromosomal region delimited by marker PE-0014628 (SEQ ID NO: 1) and marker PE-0021476 (SEQ ID NO: 37). Alternatively, or in addition, said molecular markers linked to the QTL conferring the resistance to powdery mildew on chromosome 6 are any one of the markers PE-0006634 (SEQ ID NO: 57), PE-0006718 (SEQ ID NO: 58), PE-0012892 (SEQ ID NO: 59), PE-0004191 (SEQ ID NO: 60), PE-0005575 (SEQ ID NO: 61), PE-0006320 (SEQ ID NO: 62), PE-0016499 (SEQ ID NO: 63), PE-0013625 (SEQ ID NO: 64) and PE-0022671 (SEQ ID NO: 65).

[0272] The sequences of the markers as mentioned above are described in Table 1 above and Table 12 below.

[0273] Further provided is the use of one or more of the molecular markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37) or all the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37).

33) and PE-0021476 (SEQ ID NO: 37), or a combination of the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37), or any other markers within the chromosomal region delimited by marker PE-0014628 (SEQ ID NO: 1) and marker PE-0021476 (SEQ ID NO: 37), for detecting a *Capsicum* plant that is resistant to powdery mildew.

[0274] The invention is also directed to the use of at least one of the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37) or all the markers PE-0014628 (SEQ ID NO: 1), PE-0056542 (SEQ ID NO: 5), PE-0056543 (SEQ ID NO: 9), PE-0056544 (SEQ ID NO: 13), PE-0056545 (SEQ ID NO: 17), PE-0056546 (SEQ ID NO: 21), PE-0056547 (SEQ ID NO: 25), PE-0056549 (SEQ ID NO: 29), PE-0056550 (SEQ ID NO: 33) and PE-0021476 (SEQ ID NO: 37), associated with a QTL on chromosome 6 conferring the resistance to powdery mildew according to the invention, for identifying alternative molecular markers associated with said QTLs, wherein said alternative molecular markers are in the chromosomal region delimited on chromosome 6 by marker PE-0014628 (SEQ ID NO: 1) and marker PE-0021476 (SEQ ID NO: 37).

[0275] The alternative molecular markers are preferably associated with said QTL with a p-value of 0.05 or less, preferably less than 0.01. The QTL is to be found in the deposited seeds NCIMB 43602.

[0276] The invention is also directed to a method for identifying a molecular marker associated with a QTL conferring resistance to powdery mildew when present heterozygously or homozygously, comprising:

[0277] (a) identifying a molecular marker in the chromosomal region delimited on chromosome 6 by marker PE-0014628 (SEQ ID NO: 1) and marker PE-0021476 (SEQ ID NO: 37),

[0278] (b) determining whether said molecular marker is associated with or linked to the resistance to resistance to powdery mildew in a segregating population issued from a plant exhibiting said resistance.

[0279] The population is preferably derived from a plant grown from the deposited seeds NCIMB 43602 or from a progeny thereof, exhibiting the resistance to powdery mildew as described in the invention.

[0280] In particular, the QTL on chromosome 6 mentioned above, conferring the resistance to powdery mildew according to the invention, is the QTL present on chromosome 6 in ST17-1157 (NCIMB 43602).

[0281] Genetic association or linkage is as defined above. Preferably the association or linkage is with a p-value of preferably less than 0.05, and most preferably less than 0.01 or even less.

[0282] A molecular marker and the resistance phenotype are inherited together in preferably more than 90% of the meiosis, preferably more than 95%.

[0283] In a further aspect, the invention relates to method for the production of *Capsicum* plantlets or plants resistant to powdery mildew, which method comprises:

[0284] i. culturing in vitro an isolated cell or tissue of the *Capsicum* plant according to the invention to produce *Capsicum* micro-plantlets resistant to powdery mildew, and

[0285] ii. optionally further subjecting the *Capsicum* micro-plantlets to an in vivo culture phase to develop into *Capsicum* plants resistant to powdery mildew.

[0286] The isolated cell or tissue used to produce a micro-plantlet is an explant obtained under sterile conditions from a *Capsicum* parent plant of the invention to be propagated. The explant comprises or consists, for instance, of a cotyledon, hypocotyl, stem tissue, leaf, embryo, meristem, node bud, shoot apice, or protoplast. The explant can be surface sterilized before being placed on a culture medium for micropropagation.

[0287] Conditions and culture media that can be suitably used in plant micropropagation are well known to those skilled in the art of plant cultivation and are described, for example, in "Plant Propagation by Tissue Culture, Handbook and Directory of Commercial Laboratories, eds. Edwin F George and Paul D Sherrington, Exegetics Ltd, 1984".

[0288] Micropropagation typically involves:

[0289] 1. axillary shoot production: axillary shoot proliferation is induced by adding cytokinin to the shoot culture medium, to produce shoots preferably with minimum callus formation;

[0290] ii. adventitious shoot production: addition of auxin to the medium induces root formation, in order to produce plantlets that are able to be transferred into the soil. Alternatively, root formation can be induced directly into the soil.

[0291] Plantlets can be further subjected an in vivo culture phase, by culture into the soil under lab conditions, and then progressive adaptation to natural climate, to develop into *Capsicum* plant resistant to powdery mildew.

[0292] In view of the ability of the resistant plants of the invention to restrict the damages caused by powdery mildew, they are advantageously grown in an environment infested or likely to be infested or infected by *Leveillula taurica*; in these conditions, the resistant plants of the invention produce more marketable peppers than susceptible plants. The invention is thus also directed to a method for improving the yield of *Capsicum* plants and/or fruits or for increasing the number of harvestable *Capsicum* fruits, in an environment infested by powdery mildew comprising growing in said environment *Capsicum* plants resistant to powdery mildew as defined, comprising on chromosome 6 the QTLs or sequences according to the invention and conferring to said plants resistance to powdery mildew. The invention is also directed to the use of the *Capsicum* plants of the invention for improving the yield of *Capsicum* plants and/or fruits, and/or for increasing the number of harvestable *Capsicum* fruits, in an environment infested by powdery mildew.

[0293] Preferably, the method comprises a first step of choosing or selecting a *Capsicum* plant comprising said sequences of interest conferring to said plants resistance to powdery mildew. The method can also be defined as a method of increasing the productivity of a *Capsicum* field, tunnel or glasshouse, or as a method of reducing the intensity or number of chemical or fungicide applications in the production of peppers.

[0294] The invention is also directed to a method for reducing the loss on *Capsicum* production in condition of *Leveillula taurica* infection, comprising growing a *Capsicum* plant as defined above.

[0295] The resistant plants of the invention are also able to restrict the growth of the pathogens responsible for powdery mildew, thus limiting the infection of further plants and the propagation of the pathogens. Accordingly, the invention is also directed to a method for protecting a field, tunnel or glasshouse, or any other type of plantation, from powdery mildew infection, or of at least limiting the level of infection or limiting the spread of powdery mildew. Such a method preferably comprises the step of growing a resistant or tolerant plant of the invention, i.e. a plant comprising on chromosome 6 the sequences conferring resistance to powdery mildew.

[0296] The invention also concerns the use of a *Capsicum* plant resistant to powdery mildew, according to the invention, in a field, tunnel or glasshouse, or other plantation. Preferably, the invention concerns the use of a *Capsicum* plant resistant to powdery mildew according to the invention for controlling infestation in a field, tunnel, glasshouse or any other plantation, by *Leveillula taurica*.

[0297] All the preferred features of the QTL are as defined in connection with the other aspects of the invention, in particular it is preferably present in the seeds of ST17-1157 (NCIMB accession number NCIMB 43602), and it is identifiable by the markers as defined according to the present invention.

[0298] The present invention is also directed to a method for improving the yield of *Capsicum* plants in an environment infested by powdery mildew comprising:

[0299] (a) identifying *Capsicum* plants resistant to powdery mildew comprising in their genome at least one QTL conferring resistance to powdery mildew, wherein said at least one QTL is present on chromosome 6, and

[0300] (b) growing said resistant *Capsicum* plants in said infested environment.

[0301] By this method, the yield of the *Capsicum* plants is increased, inter alia more marketable peppers can be harvested, or more commercial peppers are produced, or more seeds are obtained.

[0302] In still a further aspect, the invention also relates to a method of producing pepper fruits comprising:

[0303] (a) growing a *Capsicum* plant of the invention, as defined previously;

[0304] (b) allowing said plant to set fruit; and

[0305] (c) harvesting fruit of said plant, preferably at maturity and/or before maturity.

[0306] All the preferred embodiments regarding the *Capsicum* plant are already disclosed in the context of the previous aspects of the invention.

[0307] The method may advantageously comprise a further step of processing said peppers into a processed food.

[0308] The present invention also relates to a method of producing a food product, comprising mixing a pepper fruit of the invention, or part thereof, with one or more food ingredients. Optionally, the method comprises cooking and/or processing the pepper fruit of the invention, alone or in mixture with the one or more food ingredients. Examples of food products that comprise pepper in raw, cooked or otherwise processed form include powders, soups, sauces, salsas, pastas, condiments, pastries, sweets and salads.

[0309] The present invention also relates to a food product made of a pepper fruit of the invention or parts thereof, optionally in processed form.

[0310] In another aspect, the invention relates to the use of a *Capsicum* plant according to the invention or a fruit thereof in the fresh cut market or for food processing. Techniques for using pepper in food processing are well known from the skilled person, e.g. as an ingredient in a food product such as powders, soups, sauces, salsas, pastas, condiments, pastries, sweets and salads, and described, for instance, in Handbook of Food Science, Technology and Engineering, vol. 4, Y. H. Hui, Frank Sherkat. CRC Press.

[0311] Throughout the instant application, the term "comprising" is to be interpreted as encompassing all specifically mentioned features as well optional, additional, unspecified ones. As used herein, the use of the term "comprising" also discloses the embodiment wherein no features other than the specifically mentioned features are present (i.e. "consisting of").

Seed Deposit

[0312] A representative sample of seeds from the *Capsicum* plant according to the invention (i.e. seeds from *Capsicum annuum* ST17-1157 plant) has been deposited by HM-Clause, S. A., Rue Louis Saillant, Z. I. La Motte, BP83, 26802 Portes-lès-Valence cedex, France, pursuant to, and in satisfaction of, the requirements of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure (the "Budapest Treaty") with the National Collection of Industrial, Food and Marine Bacteria (NCIMB), 23 St Machar Drive, Aberdeen, Scotland, AB21 9YA, United Kingdom, on Apr. 22, 2020, under accession number NCIMB 43602.

[0313] A deposit of the ST17-1157 seeds is maintained by HM-Clause, S. A., Rue Louis Saillant, Z. I. La Motte, BP83, 26802 Portes-lès-Valence cedex, France.

EXAMPLES

Example 1: Identification of Resistance Source to Pepper Powdery Mildew

[0314] Several *Capsicum annuum* lines were grown in a greenhouse when a natural infection of *Leveillula taurica* started to build up. At the end of the growing cycle, certain plants from one of the lines, LT17, appeared to be symptomless while the entire greenhouse had *Leveillula* symptoms. Cultivar Maor, known as a susceptible control for *Leveillula taurica* was present on either side of the resistant plants and was fully infected by *Leveillula taurica*. The inventors thus decided to further evaluate the resistance level of LT17.

Example 2: Evaluation of Resistance Level of the LT17 Line

[0315] Three seed lots of LT17 were evaluated according to an "artificial" test consisting in the artificial inoculation of the fungus *Leveillula taurica* on 4-6 leaf stage plantlets.

[0316] Artificial inoculation was conducted on several plantlets per lot. To prepare the inoculum, spores of *Leveillula taurica* were collected by rinsing leaves of sensible pepper varieties infected with the fungus, with distilled water. The spores were counted using a Nageotte counting cell (concentration of about 10^4 c/ml). 1 mL/L Triton X 100

was added (Molot, Jp Leroux, M Diop-Bruckler. Agronomie, EDP Sciences, 1990, 10 (7), pp. 551-559. fihal-00885316f.

[0317] The inoculum was sprayed on pepper plantlets at the 4-6 leaves stage. For each plantlet, the 4 oldest leaves were sprayed, on both faces. The inoculum was sprayed two times on each plantlet, one week apart. 4 weeks after the 1st inoculation, the severity of the infection on plants was scored according to the following rating scale, based on percentages of infected area:

TABLE 2

scoring scale used in the artificial test		
Score	Percentage of infected area	Descriptions
9	0%	—
8	0%	No sporulation but “small spots” present
7	<10%	—
6	10-25%	
5	25-50%	non-infected leaves are lightly affected only on the lower face
3	50-75%	presence on the upper face of the leaves
1	75-100%	strong presence on both faces

[0318] The following lines were used as control: Jaune Dupoux, Pipo, Piperade (susceptible); OCP32, OAP35, ODS200, ODS257 (intermediate) and HV12 (resistant).

[0319] The artificial test reveals that the three seed lots segregate for *Leveillula taurica* resistance. Specifically, plants with the highest level of resistance (9) are present within the three seed lots.

[0320] In order to confirm the level of resistance in tunnel and on adult plants, plants from artificial test with a resistance score of 9 were then transplanted in tunnel for a second stage of resistance evaluation.

[0321] In the same tunnel, three different lots of LT17, as well as plants from the Maor line and an F1 [Maor*LT17] were sown and planted. The sensitive plants were positioned at the entry and exit of the tunnel, as well as at regular intervals throughout the tunnel in order to ensure a homogeneous and intense *Leveillula* pressure during the assay.

[0322] 2 artificial inoculations (2 weeks interval) were carried out, by spraying the median part of the plants with inoculum with *Leveillula taurica* and evaluations were performed at the end of the test (8 weeks after the first inoculation). A specific rating scale for tunnel evaluation was used. Each plant was scored for three different notes: Note bot=percentage of infected leaves area in the bottom part of the plant (calculated as the mean of the 5 more infected leaves); Note med=percentage of infected leaves area in the medium part of the plant (calculated as the mean of the 5 more infected leaves)”; and Note Glo=percentage of infected leaves per plants (whatever the percentage of sporulation).

[0323] For each plant, an Index 1 (I1) was also calculated, corresponding to the mean of Note bot, Note med and Note glo.

[0324] Notes were attributed according to the following scale:

TABLE 3

Scoring scale for “Note bot” and “Note med” in the tunnel test Scale for Note bot and Note med		
Note	Percentage of plants with mycelium	Descriptions
9	0	nothing
8	0*	Very few spots of penetration (and not mycelium)
7	1-10%	% of the infected leaves
6	10-20%	area (Mean of the more
5	20-30%	infected leaves)
4	30-40%	
3	40-50%	
2	50-60%	
1	>60%	

TABLE 4

Scoring scale for “Note glo” in the tunnel test Scale for Note Glo		
Note	Percentage of plants with mycelium	Descriptions
9	0%	Percentage of infected
8	0.1-10%	leaves per plant
7	10-20%	(whatever the percentage
6	20-30%	of sporulation)
5	30-40%	
4	40-50%	
3	50-60%	
2	60-70%	
1	>70%	

[0325] The plants were also tested for the presence of the RZ marker. This marker is disclosed in International Patent publication WO2014/140331 as a marker linked to a QTL of resistance to *Leveillula taurica* on linkage group LG1/8 of pepper genome. In the marker linked to QTL on LG1/8, the A in susceptible plants is replaced with a G in resistant plants. For detection of the RZ marker, the Kaspar technology was used. Plant DNA was amplified using the 3 primers CATGTGAGCGACTCCTCTGACAA (primer for the susceptible allele, SEQ ID NO: 43), ATGTGAGCGACTCCTCTGACAG (primer for the resistant allele, SEQ ID NO: 44) and CGAGGAGAAT-TTCCTGCTAAAATTGAGTT (Common primer, SEQ ID NO: 45).

TABLE 5

Resistance levels and RZ marker screening of plants of the LT17 line in the tunnel test		
Plant	RZ marker	<i>L. taurica</i> Resistance level
1	A:G	9
2	G:G	9
3	A:A	7
4	A:G	9

[0326] Table 5 above shows the resistance levels for four plants of the LT17 line planted in tunnel. A high level of resistance to *L. taurica* is present in several plants of the LT17 line, which confirms the results obtained using the artificial test. The screening experiment on the RZ marker suggests that the QTL on linkage group LG1/8 is not responsible for the resistance.

[0327] Progenies of these plants obtained by selfing were tested for *Leveillula taurica* resistance in an artificial test, as described above. 2*10 plants were tested per progeny. The RZ marker was screened on individual plants.

TABLE 6

Resistance levels and RZ marker screening of progeny plants of the LT17 line in the artificial test.		
Plant number	RZ marker	<i>L. taurica</i> Resistance level in artificial test
1_17LT05-1	A:A	9
1_17LT05-3	A:A	9
1_17LT05-4	A:A	9
1_17LT05-5	A:A	9
1_17LT05-6	A:A	9
1_17LT05-7	A:A	9
2_17LT05-9	A:A	9
2_17LT05-10	A:A	9
2_17LT06-1	A:G	9
2_17LT06-2	A:G	9
3_17LT06-8	A:G	9
5_17LT09-5	G:G	9
6_17LT09-6	G:G	9
7_17LT05-17	A:A	9
7_17LT05-20	A:A	9
8_17LT06-12	A:A	9
10_17LT08-12	A:G	9
10_17LT08-15	A:G	9

[0328] The results in Table 6 shows that resistant plants comprise a diversity of genotypes at the RZ markers, i.e. homozygo [text missing or illegible when filed] is not implied in the resistance to powdery mildew as found in certain plants of the LT17 lines and their progeny.

[0329] These plants have been transplanted in tunnel for another *Leveillula taurica* resistance evaluation in autumn.

Example 3: Genetic Analysis

[0330] Genetic distance analysis of line LT17 was carried out on a high density Affymetrix SNP chip array, using 590 accessions and 5005 SNPs. The genetic distance tree in FIG. 1 shows that LT17 belongs to a *Capsicum annuum* cluster, but is clustered in a different group than line H3 (Daubeze, 1995; Lefebvre 2003). The source of the resistance, LT17 is thus genetically unrelated to the powdery mildew resistant lines H3 and its progeny HV12.

[0331] A plant from the LT17 line was also genotyped on a high density genotyping chip (Metachip), together with several plants from another source of *Leveillula taurica* resistance, the PBC167 line (International patent application WO2013/033210). About 36% of the 5360 analyzed SNPs exhibited a different allele between the LT17 line and the PBC167 line, showing that the two lines are genetically unrelated and distinct.

Example 4: Development of F2 Resistance Segregating Population

[0332] A segregating F2 population from the cross between the susceptible line Maor and the LT17 line dis-

playing resistance was developed to generate a linkage map and detect QTLs involved in the resistance (QTLs mapping).

[0333] 179 F2 plants were genotyped with 2565 SNP markers evenly spaced along the full genome and being polymorphic between both Maor and LT17 parental lines.

[0334] The same 179 F2 plants as well as the 2 parental lines and the hybrid F1 Maor×LT17 were evaluated for *Leveillula taurica* resistance in tunnel under natural and artificial pressure create through two artificial inoculations performed with a *Leveillula taurica* strain (see Example 2)

[0335] The notation was made using the same three notes (Note bot, Note med and Note Glo) and scale than in Example 2 but was performed at 3 different times with 3 weeks interval for each notation (the first notation was done 2 weeks after the first inoculation).

[0336] This kinetic allows to calculate an AUDPC (Area under the disease progress curve) which is a powerful quantitative index to perform QTLs analysis. The AUDPC was calculated for each note and for the Index 1 (I1) by using the following formula:

$$AUDPC = \sum_{i=1}^{n-1} \left(\frac{y_i + y_{i+1}}{2} \right) (t_{i+1} - t_i)$$

[0337] “n” is the number of symptom assessments, “y” the symptom intensity and “t” the time in dpi (days post inoculation).

[0338] AUDPC was calculated plant by plant, for each note.

[0339] FIG. 2 presents the distribution of the “AUDPC Glo” in the F2 population. This figure shows that the segregation of the trait follows a “normal distribution” which highlights a complex inheritance of the trait. According to this segregation, these data were used to carry out QTL mapping analyses.

Example 5: QTL Mapping Analyses in F2 Population Segregating Population

[0340] Segregation of each SNP within the F2 population was evaluated using the Chi-square test χ^2 . SNP markers exhibiting distortion of segregation tested against the Normal Mendelian expectation ratios -1:2:1 for an F2 were removed from the analyses.

[0341] Significance levels used is from P-value<0.01

[0342] A genetic linkage map was then performed with the JoinMap V4 software to assign genetic position of each SNP. Linkage groups were then assigned to chromosome using the physical position of the SNPs on a reference genome of Maor, version v3.0. QTL analyses (the association between genotypic variation and the trait variation) were performed with MapQTL V5 software.

[0343] We determined the number of genomic regions involved in the quantitative trait as well as their position and their effect, using the H0 hypothesis: no segregating QTL (locus with no genetic effect) and the alternative hypothesis H1: presence of a segregating QTL.

[0344] Putative QTLs were first detected using the Interval mapping module of the software (single-QTL model). A QTL likelihood map was determined: for each position on the genome—in our case each 2 cM—the likelihood for the presence of a segregating QTL was calculated.

[0345] Then the likelihood under H0 and H1 hypothesis were compared with a likelihood ratio statistic (LOD score): A QTL was detected when the LOD score exceeds the significance threshold, in a Linkage group. The estimated position of the QTL on the map correlates with the highest LOD value on the LG or chromosome. The significance threshold of the LOD score was determined to be 3.0, with the permutation test with a P-value of 0.05.

[0346] Then the multiple QTL mapping (MQM mapping, or composite interval mapping) module was performed: use of markers close to QTLs as cofactors to obtain the best possible final position of the QTLs.

[0347] One major QTL was identified on linkage group 6 with the favorable allele originated from the LT17 source (FIG. 3). No QTL was identified on chromosome 4, known to harbor a major QTL for *Leveillula taurica* resistance in other sources (PBC167).

[0348] The results in Example 2 have shown that the known QTL1/8 for resistance to *Leveillula taurica* does not seem to be implied in the resistance to powdery mildew in the LT17 plants. The QTL mapping experiments also show that the known major QTL4 is not present in LT17. Accordingly, the newly identified QTL on linkage group 6 seems to provide resistance to *Leveillula taurica*, even in the absence of two known major QTLs for resistance to *Leveillula taurica* on linkage group 1/8 and linkage group 4.

TABLE 7

Data for the QTL identified on chromosome 6				
Linkage group	QTL position (LOD max) (cM)	LOD Max	Confidence Interval in cM (LOD-1.5)	% of explained variance
6	157.4	5.04	147-160	9.6

[0349] Confidence interval was estimated based on the maximum LOD value minus 1.5.

Example 6: Validation of QTL Effect in F2:3 Families

[0350] Based on molecular profile at QTL position on chromosome 6, F2 plants of the cross (Maor×LT17) were selected and selfed to produce F2:3 families. A total of 25 F2 plants were selected, 12 being homozygous for favorable allele inherited from LT17 and 13 being homozygous for unfavorable allele inherited from Maor at the region of the QTL located on chromosome 6.

[0351] Three different SNP markers were used to identify molecular profile at chromosome 6 QTL position. The three SNPs were selected within the interval of the QTL on chromosome 6. One SNP was selected at the left side of the interval, one SNP corresponding to the SNP with the highest significance and one SNP at the right side of the QTL interval.

[0352] SNP information are provided in Table 8 below.

TABLE 8

SNP markers for the QTL identified on chromosome 6			
SNP Name	PE-0014628	PE-0056542	PE-0021476
Linkage Group	6	6	6
Position (cM)	150.97	157.42	158.15
LOD	3.89	5.04	4.16
Physical chromosome on CM334 v.1.55	6	6	6
Physical position on CM334 v.1.55	232 015 212	233 673 276	235 235 179
Sequence	AAGATACTCTG CATTGGGTCCG GGCTAGAAGAA TAAGCACGAAT AACCGC[A/G]TT CCAAGAAAATA GATTGGTTGA GGAATTGATC GAACACTTGGT GGGC (SEQ ID NO: 1)	ATTTTCTAGGTT ACTGGATTTAAC TCAGGAGAAAA TGGATTGTGTTT TGCAATGCATTT TTCCGGGGGTG CCTAGTTTATAA AGACCTAAGAT GTTTCAGTCAA CTGCTTAATTAG GACCTTGACT GTTGTGGTTAT GACTGGGCCCC T[T/C]GCCAAGA TGGTTTGGAC AACAGAGCTCT TACCACTTG CTGCTATTTGA TCTACTTGATCT TGAGTTGCGCT	CTCCAAAACCA CAGCTACTAATT TGTTAGATCCAA AACTTAAAGCT TAT[C/G]TTTAGT CAGTCAATTAAA AATCCCAACAAT TTAAAGCTAAG AAGTGAAG (SEQ ID NO: 37)

TABLE 8-continued

SNP markers for the QTL identified on chromosome 6			
		GTTTGCTGTGC TGTTCAAAGCG ACATTGACTGT CTGAGATCTATA AAAGATTCATTA GAAGATCCAAA TAATACT (SEQ ID NO: 5)	
Forward primer- resistant allele	GAAGGTGACCA AGTTTCATGCTA GAAGAATAAGC ACGAATAACCG CA (SEQ ID NO: 3)	GAAGGTGACCA AGTTTCATGCTG TGGTTATGACT GGGCCCTC (SEQ ID NO: 7)	GAAGGTGACCA AGTTTCATGCTT GTTGGGATTTT AATTGACTGACT AAAG (SEQ ID NO: 39)
Forward primer- susceptible allele	GAAGGTCGGAG TCAACGGATTA GAAGAATAAGC ACGAATAACCG CG (SEQ ID NO: 2)	GAAGGTCGGAG TCAACGGATTG TGGTTATGACT GGGCCCTT (SEQ ID NO: 6)	GAAGGTCGGAG TCAACGGATTG TTGGGATTTT ATTGACTGACTA AAC (SEQ ID NO: 38)
Common Reverse unlabelled	AATTCTCAACC AAATCTATTTTC TTGGAA (SEQ ID NO: 4)	TAAGAGCTCTG TTGTCCAAAC CATCTT (SEQ ID NO: 8)	CTCCAAAACCA CAGCTACTAATT TGTT (SEQ ID NO: 40)
Allele LT17 (resistant)	A/A	C/C	C/C
Allele Maor (susceptible)	G/G	T/T	G/G

[0353] The 25 F2:3 families were evaluated in tunnel under artificial inoculation in 2 different geographical locations, using two different local strains, SRY and Fondi. In both location each F2:3 family was evaluated in 2 replicated plots of 5 plants each.

[0354] In the first location, evaluation of disease was performed using a simplified scoring scale.

TABLE 9

simplified scoring scale		
Simplified scoring scale: 1 note by plants: bottom/medium leaves		
Note	Percentage of plants with mycelium	Descriptions
9	0	nothing
8	0*	Very few spots of penetration (and not mycelium)
7	1-10%	% of the infected leaves area (Mean of the more infected leaves)
6	10-20%	
5	20-30%	
4	30-40%	
3	40-50%	
2	50-60%	
1	>60%	

[0355] Each plant was scored at 29 dpi, 50 dpi and 71 dpi and AUDPC was calculated as described in Example 3.

[0356] In the second location, evaluation of disease was performed only once according to the same simplified scoring scale.

[0357] FIG. 4 represents the distribution of disease scoring at the 2 locations for the F2:3 families according to their

genotype profile at the QTL location on chromosome 6 (homozygous for favorable allele versus homozygous for the unfavorable allele).

[0358] At both location, significant differences were observed between means of disease scoring according to QTL profile. This validates the QTL effect within the genetic background in which it was initially identified. These experiments also show that the identified QTL provides resistance to different strains of *Leveillula taurica*.

Example 7: Candidate Genes Identification within QTL Interval

[0359] The confidence interval of the QTL corresponds to 3.3 Mb on the reference genome CM334 v1.55 between 232.0 Mb and 235.3 Mb on the chromosome 6. Based on physical genome gene annotation from gene prediction tools, a total of 410 genes were found within the interval.

[0360] Among all the genes found in the QTL interval, two genes have been found as good candidate genes based on their molecular functions and their physical position proximity with the peak value of the QTL (maximum LOD value).

[0361] One MLO-like protein gene (MLO-like protein, gene name=CA06g26150 in *C. annuum* cv. CM334 v1.55 genome) and one receptor-like kinase (probably inactive leucine-rich repeat receptor-like protein kinase gene, name=CA06g26250 in *C. annuum* cv. CM334 v1.55 genome).

[0362] Implication of other Mlo genes or Mlo-like sequences in the resistance of fungus and in particular for *Leveillula* in pepper has been described (Zheng Zheng et al 2013, Lim and Lee (2014), Kim and Hwang (2012)).

Example 8: Markers Development within Candidate Genes

[0363] The 2 candidates genes were sequenced by targeted sequencing by in solution fragment capture (Gnirke et al., Nat Biotechnol. 2009). Based on sequences data for a set of 6 genotypes carrying or not the full favorable segment at QTL interval, SNP mining was conducted.

[0364] Several genotypes were sequenced for the 2 candidates genes: Maor, CPNT, Vania, BYP, LT17, F3 (LT17*Maor) and HV12.

[0365] A total of 8 SNP were identified within or close to the 2 genes, with respectively 6 SNPs in CA06g26250 and 2 SNPs in CA06g26150.

TABLE 10

SNPs identified in CA06g26250 and CA06g26150.											
Gene	Chromosome	Position CM334 v1.55	Ref Allele (=Maor)	Alt Allele	Maor	CPNT	Vania	BYP	LT17	F3	HV12
CA06g26250	6	233 673 276	T	C	.	.	.	.	C/C	C/C	C/C
CA06g26250	6	233 673 121	G	A	.	.	.	.	A/A	A/A	.
CA06g26250	6	233 672 530	G	T	.	.	.	.	T/T	T/T	.
CA06g26250	6	233 671 922	T	C	.	.	.	.	C/C	C/C	.
CA06g26250	6	233 671 648	A	C	.	.	.	.	C/C	C/C	.
CA06g26250	6	233 671 204	A	C	.	.	.	.	C/C	C/C	C/C
CA06g26150	6	233 493 124	T	A	.	.	.	.	A/A	A/A	.
CA06g26150	6	233 489 782	C	T	.	.	.	.	T/T	T/T	.

“.” means that the plant homozygously comprises the Ref Allele.

[0366] For each of these SNP, Kaspar markers were developed, as shown in Table 11 below.

SNP markers in or near genes CA06g26250 and CA06g26250									
SNP marker Name	Physical position CM334 v.1.155	Sequence surrounding the marker	Forward primer (susceptible allele)	Forward primer (resistant allele)	Common reverse primer	Susceptible allele	Resistant allele		
PPB- 00056542	233 673 276	ATTTTCTAGTCTAGTGAATTAACCTCAGAGAAAATGGAT TTGTTTTCAATGCAATTTTCCGGGGTGCCCTAGTTTAT AAAGACCTAAGATGTTTCAGTCAACTGCTTAATATAGGACC TTTGACTGTTGGTTATGACTGGGCCCTT/C G GCAAG ATGGTTTTGGACAACAGAGCTTTACCACTTGCTGCTA TTTGATCTACTTGAATTTGAGTTGCGCTTTGCTGTGC TGTTCAAGCGCAATTGACTGCTGAGATCTATAAAGAT TCATTAGAAGATCCAAATAACT (SEQ ID NO: 5)	GAAGGTCGGA GTCAACGGATT GTGGTTATGAC TGGGCCCTT (SEQ ID NO: 6)	GAAGGTGA CCAAGTTC ATGCTGTG GTTATGACT GGGCCCT C (SEQ ID NO: 7)	TAAGAGCT CTGTTGTC CAAAACCA TCCTT (SEQ ID NO: 8)		C/C		
PPB- 00056543	233 673 121	AGATGGTTTTGGACAACAGAGCTCTTACCACACTTGCTG CTAATTTGACTACTTGTGATCTTGAGTTGCGCTGTTGCTG TGCTGTTCAAGCGCAATGCTGCTGAGATCTATAAAA GATTCATTAGAGATCCAAATAACTTTGA G/A CTATAC ATGGAAATGTCATACTCAGCGAAGTTTCAATTGCAAG TTTGCTGGGATTCAGTCTGGCATCTGATGAAAATAAG GTTCTGAGTATCAGTCTCTCTGACATGGTCTCAAGGCG AGTTTCCACTTGCTTCGGAATG (SEQ ID NO: 9)	GAAGGTCGGA GTCAACGGATT GTGGTTATGAC TGGGCCCTT (SEQ ID NO: 10)	GAAGGTGA CCAAGTTC ATGCTGTG GTTATGACT GGGCCCT C (SEQ ID NO: 11)	ATGABAAC TTCCGTCA GATTATCG AACTT (SEQ ID NO: 12)	G/G	A/A		
PPB- 00056544	233 672 530	CTGTTCTCTCAGAGAGCTATGCAATAATCCCGAGCTGT GTGGAAACCGTTAGATCTTGTCCAGGGTACTCTPAAGA AACCCCTACCCGAATCATTTGTCGGTGCAGCTGTTGGTG GGTAAGTTTAGTGCTGTAATTTTGGCCATCG G/T TAT GTTCTTCTATAGAGAAGTATCCAGGAAGAGAGAA GGATGAATATCTGAAGGAATAAATGGGCGAAGATAT TAAGGTAATAAGCAATCAGGTAGAATTTTAACTTCT CTGAATCAITAAAAAGTCTCTGTTACTG (SEQ ID NO: 13)	GAAGGTCGGA GTCAACGGATT GTGGTTATGAC TGGGCCCTT (SEQ ID NO: 14)	GAAGGTGA CCAAGTTC ATGCTGTG GTTATGACT GGGCCCT C (SEQ ID NO: 15)	TTCTCTTTC CTGGATAT CCTTCTCAT ATA (SEQ ID NO: 16)	G/G	T/T		
PPB- 00056545	233 671 922	TCCTTTTCAAGACATTTAACTAATGAATTTTCACTGCT CCGCTCTCGTGTGATATTTTCTCTGTTCAAGTTTCT TAATGACTTTAAGTTCCTTATTTTCAGCTTTCAATGTTTG AAGAGTCTACTTTCAAAATGAGATTTAGT/C G ATCTATG AAGGCCACTTCACAGCTTCAACAAAAACAATATTATCGGT CAGGAAGAACCGGACTTTTACAAGCAGTACTGATG ATGGCACTTCACTTATGTTTAAAGGTTGCAGAATACTCA GCACTCAGAGAAAGAGTTTATG (SEQ ID NO: 17)	GAAGGTCGGA GTCAACGGATT CTGTCAATGAC CTTCAATGAC CA (SEQ ID NO: 18)	GAAGGTGA CCAAGTTC ATGCTGTG TCATAAGAT CG (SEQ ID NO: 19)	GTTCGAGA AGTCTACTT CAAAAATG AGATT (SEQ ID NO: 20)	T/T	C/C		
PPB- 00056546	233 671 648	ACTCAGCACTCAGAGAAAGAGTTTATGTCGGAAGTGGCT ACATTCGGAAATGAAGACATCTGAATCTGGTTCCTCTT TAGGTTTCTGCATTCGCAAAAGAAAGAGCTGTGCTCTA CAAGACATGCAAAATGGAACCTTGCATAT A/C GGTTA CATTCGGTGAAGAGCGAAAAAATCTAGAGTGGCT ATAAGATTAAAAATAGGCATTGGAGCAGCCAAAGATTG ATGCGCTTACCACAACTGCAATCTCTGATCAATCCAG AAATATAGTTTAAATGATCTTGC (SEQ ID NO: 21)	GAAGGTCGGA GTCAACGGATT CTTCACTCACC GAATGAACCT (SEQ ID NO: 22)	GAAGGTGA GAGTCAAC GGATCTCT CACTCACC GAATGTA CCG (SEQ ID NO: 23)	TTGCTTAC AAAGACAT GCCAAATG GAA (SEQ ID NO: 24)	A/A	C/C		

TABLE 11-continued

SNP markers in or near genes CA06q26250 and CA06q26250									
SNP marker Name	Gene	Physical position CM334 v.1.55	Sequence surrounding the marker	Forward primer (susceptible allele)	Forward primer (resistant allele)	Common reverse primer	Susceptible allele	Resistant allele	
PE-0056547	CA06q26250	233 671 204	GCTACTCCAAAAGTGATGTGTACAGTTTCGGTCTGTGA	GAAGGTGACCA	GAAGGTCG	CATTGTCAT	A/A	C/C	
			CTTCTCAGTTAGTTACTGTGTGAGAAACACCTCTGTC	AGTTCAATGCTC	GAGTCAAC	AACCTTTAC			
			ACCAAAGTCCCAGACCTTTAAGGGGAATTTGGTGGAA	AACTCTCCGGT	GAGTCAAC	CAGACAA			
			TGGATCACACAACCTCTCCGGTGAATCTAAGCTT[A/C]AAG	GAATCTAAGCT	TCTCCGGT	GAAT (SEQ ID No: 28)			
			ATGCGATTGACCAATTCATTGCTGTGTTAAAGTTATGACAA	TA (SEQ ID No: 26)	GAATCTAA				
PE-0056549	CA06q26150	233 493 124	TGAGATCTTCCAGGTCCTTAAAGTTGCTTTCGATGTGTG		GCTTC				
			TTGTCCTCTGTTCTTAAGAGAGAGGCCGACATGTTGAA		(SEQ ID No: 27)				
			GTTTACCAGCTTCTGAGGCCATCGGAG (SEQ ID No: 25)						
			TAACTGATGGTTTCAGTTGAGCTGCAAAATGGAACATCAA	GAAGGTCGGA	GAAGGTGA	GAGTTTCAT	T/T	A/A	
			TATCAGTAGCTTCTCGTTTCAACCAATTTACTTCTTTGCTC	GTCACGGATT	CCAAGTTC	GATTCATCT			
PE-0056550	CA06q26150	233 489 782	TTTAGTCACCTTGACCTTTAGAGAAGATGACGTGATCCC	CCGATATTTTAT	ATGCTCCG	GTTTGTCT			
			GATATTTATATACTAGCTATTCTAGCATAT[A/GGGAAGA	ACTTAGCTATT	ATATTTAT	CCC (SEQ ID No: 32)			
			AAACAGATGAATCATGAACCTCTTCGGTAGTAAAAAAAT	CTAGCATAT	ACTTAGCTA				
			TATGTCCAAAATGTGGATGTATCTTTTATGGACGATG	(SEQ ID No: 30)	TTCTAGCAT				
			ATCCATACCATGTTCCGATTACGATCCACTTTTACGATT		AA (SEQ ID No: 31)				
PE-0056550	CA06q26150	233 489 782	GCATCTCCAGCTCTTCCATT (SEQ ID No: 29)						
			TTTGATTTTGTAACTCAATTTGCATGAAATGATATGTTTA	GAAGGTGACCA	GAAGGTCG	TAGGTAATA	C/C	T/T	
			TATGTTTGTCTAGAAATATCACAACTATTACATTTATAGT	AGTTCAATGCTA	GAGTCAAC	TTTGAGTTG			
			AAATTTGAGTTGCACTATGTTATAGTGCAATTTTATTTTC	CATTAAATTTTC	GAGTGA	CAGTTATG			
			TATTTATGTGACATATTTTATAGATA[C/T]ATGTCGAAATA	AGCATATTTTC	CATTAAATTT	GTAT (SEQ ID No: 36)			
PE-0056550	CA06q26150	233 489 782	ATGCTGGAATTAATGTTCACTCTTTGAATACATGAAA	GACATG (SEQ ID No: 34)	TCCAGCAT				
			GACCGAACTTCTACCCCTATTTTAAAGGATGACGCG		ATTTCGACA				
			TGAATTAGACCTAATTAAGAAATATTTATCTTACACTTTT		TA (SEQ ID No: 35)				
			GTTTTATCGAGAAGCTTTT (SEQ ID No: 33)						

Example 9: SNP Analysis on Chromosome 6

[0367] Example 8 shows that the LT17 line has distinct alleles at the identified QTL when compared with the reference Maor line, but also the CPNT, Vania and BYP lines. The LT17 line is also distinct from the resistant HV12 line at the identified QTL on chromosome 6. In order to further characterize the LT17 lines, the inventors performed a SNP metachip analysis on chromosome 6 of the CM334 v.1.55 reference genome. The inventors identified 31 SNPs for which the LT17 line has a different allelic state in comparison to the *Leveillula taurica* resistant PBC167 line. This confirms that the QTL identified on chromosome 6 is unique compared to the known resistant lines.

TABLE 12

SNP analysis on chromosome 6						
SNP marker	Sequence	Sequence Orientation	Chromosome	Position in bp (CM334 v. 1.55)	LT17	PBC167
PE-0011447	TTGAACCCCTCCTTCCCTTTTCATACATTGAAGAATAATGGTGGATGGAGACT[C/G]AG CAATCATTTGGTCACATCATACATACACCTCAAGAGCATACTAGAAGCTGA (SEQ ID NO: 46)	Top	6	219 426 890	C/C	G/G
PE-0011448	AGTTGCCTTGGTACCAATGAAATTGACTCTTTCTTCCAAAGTCCAC[A/G]GC TCCAGGACCTGCATTTCAGATCCAGACCTTGC CGCTTCGGCTTCTGTT (SEQ ID NO: 47)	Top	6	219 425 984	A/A	G/G
PE-0019751	GGTGTAGAAATCCCTCCCTCGCTCGCTGACGGCAGCTCCGTTACATCGG[A/C]G GTTTTACTAGAGTCCCTCTCCGTTGATCGGTCTATTCTTTAATCAGAGTT (SEQ ID NO: 48)	Top	6	220 448 196	C/C	A/A
PE-0021027	TGGTTCTAATGTGGTGACAAAGGTTCCATCAAGTGAATGTATTGGTGATG[G/T]TG GTGTAAGTGCTAACGTGCAGAGAGTAAAGCAAAATGAAGCAAAACGGGG (SEQ ID NO: 49)	Bottom	6	219 581 697	T/T	G/G
PE-0022402	GAGATAGAAAATGAAGTGTAACAAACTTTGGATCTCCAGCTGCACCAGT[C/T]CA GGAACGACCAATTCCAGAGATCATGGGTTCTCCCGACCTCCTCCAGT (SEQ ID NO: 50)	Bottom	6	219 545 686	T/T	C/C
PE-0018388	AAAAGAAATCCGCCCTAACTGCTTTCTGCTCTTCTTCTTCACATTGACTTC[A/C]TCG GCATCATCTAGAACCAAAATTCATGTACTCATCAAAACCAATAATACG (SEQ ID NO: 51)	Top	6	224 014 874	C/C	A/A
PE-0028409	ATAATGATGTCACCAGTCTGAACCTACCGAAGACAGTTCTTTGCTTCAATC[A/G]AT ATGCTACAGAACAAAGCTGAGAAATTACTTGAAGAGATGAAGCCTAAG (SEQ ID NO: 52)	Top	6	225 012 078	G/G	A/A
PE-0006573	GTCCCTCTTTCAAAGAGTTAAAGAGTGGTGTGGGATTTGCTTTGACTACTACC CGCTT[G/A]ATGACATAATGCTGTTGGTTGTGTCATCCTTTTGTACTATTGCT GGAAAGCTTACA (SEQ ID NO: 53)	Top	6	225 782 942	G/G	A/A
PE-0001206	AKTAAAGAGAACACCTGTAATTCACAGTACTACTACCCAGAATGGGTCAATTTAC ACC[T/G]ATCTTGACAAAATGATGACATCCATCCCTCCTATCTTATTGAAGCATG CTACATTGCTTTGGCTTTGACAGATAAGTGTTCACACATATCCAAATAAGAT GCACACCATATAACATTGATGCTGGCAAGAGTATGCACCTGCAG (SEQ ID NO: 54)	Bottom	6	227 119 514	T/T	G/G
PE-0005987	GGAGCAACAAACAAAAGTATCCCACTACCCCTCCCACTAAGATGGGGATAGTGAA ACAAGG[A/C]ATAGTGCCATTCTTGGATCAGACTCTGACAAAAGAAACAAAGGCTG CTGAGCTGTTTCA (SEQ ID NO: 55)	Top	6	227 195 529	C/C	A/A
PE-0005367	GACGCTTATGGGATTTCCAGGGTAAACATTACGTTGCTTACTTGTATATA CTTGG[C/T]CGTGAGTGAACTGGAAAGAGCTGAAGAATGTATAATAATATATGC CAGTAACCCCTGGT (SEQ ID NO: 56)	Bottom	6	232 016 457	T/T	C/C

TABLE 12-continued

SNP analysis on chromosome 6						
SNP marker	Sequence	Sequence Orientation	Chromosome	Position in bp (CM334 v. 1.55)	LT17	PBC167
PE-0014628	AAGATACTCTGCAATGGTCCGGCTAGAGAATAAGCACGAATAACCGC[A/G]T TCCAAGAAATAGATTTGGTTGAGGAATTTGATCGAACACTTGGTGGGC (SEQ ID NO: 1)	Top	6	232 015 212	A/A	G/G
PE-0006634	CGGATATTGGAGAAGATGAAGGCTCCATAAAGCCAGATGATGTGCCTGAGATC ACGCC[G/A]AGCAGCTCTTGGAAAGCTGATGGTTTCATCTTTGGTTTCCCTTCGCG TTTTGGTGTGATGG (SEQ ID NO: 57)	Top	6	232 005 084	A/A	G/G
PE-0006718	TCGGTTCTTCACAGTTATTGCAGTATTGGTGAGACCAGACAGGAATCAAAAT AAGAA[G/A]ATACTGGAAATGGTACCAATTACACAGCAGGAAGAAATTTGATATTATT GGCATCAATAAA (SEQ ID NO: 58)	Top	6	233 577 814	A/A	G/G
PE-0012892	ATGCTTTACAAACCTCCATTGACAAACAAAGATTACATTGGCGGCAAACTC[C/G]TC CACGCCATATACTCCGACTTGGCCTTTCCTCTCATACACTCCTCCTA (SEQ ID NO: 59)	Bottom	6	235 354 680	G/G	C/C
PE-0004191	TATACAAAGAAATGTTGGAAGGGGAATGGGAAATGGAATATCAGCTGGCACAAAT TGTCCTT[AJAGCAGAGTAGCTTTATTGCTTGATGGAACAATCATGTATAGATTTCAT GAAGGCAGAGGG (SEQ ID NO: 60)	Bottom	6	235 302 145	A/A	T/T
PE-0005575	TGCTAGTTGTGGCATTAAATTTGCTTCTAGCCTTTCAGAAAAGACATTACGTT GTAA[A/C]CTTGTAAGTGTATTAATTCCTTGACATGAAATGAAGTCGAGTGGAA CACAACAAAAAA (SEQ ID NO: 61)	Top	6	235 289 831	A/A	C/C
PE-0006320	CTAGGCCTAGTTGATGGGACTACACAATGGTATGCTGCAAACTAAGCAAAAGAT GGGCG[A/G]CGATGGGAAAGATCTGCATACCAGCAACTTTGTTGGGCATATA TTTTAGTCATGCTA (SEQ ID NO: 62)	Top	6	235 287 024	A/A	G/G
PE-0016499	GAATGTTTTTCGCTTTTGTGTACACACTCTCTTGATCATGCTCAAATG[C/T]TTC AAAAGCATACATGCATTTAAAGAAATCCTTCAGCAGCAATCCAGAAG (SEQ ID NO: 63)	Bottom	6	235 290 056	T/T	C/C
PE-0013625	TGAATCCAAACAGATTTGGGACACTGACTGGGACTGATGAACATCAAGT[A/C]TT GCCTTTCTTGCCCTCCTTTTGTTCGTTTTCTTCCTTTCTGTTGGTTA (SEQ ID NO: 64)	Top	6	235 238 619	A/A	C/C
PE-0022671	TGATCAATAATCAAAATCTGATGTTCAACAATACAAATATGCACAAGAGGA[A/G]AAA AACTGGTTCATAAATGGTGAATCAAGAAGGCAATACACAGCCCTTT (SEQ ID NO: 65)	Top	6	235 238 219	A/A	G/G
PE-0005787	CTCCGTCATCTGTTAAATGCTTCATTGCTTCAGACATTATTAAAGTCTAGAATT CAGG[C/T]GTTCTGTTACCTGTTTATATTGTACATGACAAATGTTGATGTCATGTGGT AGCATTTGTTG (SEQ ID NO: 66)	Bottom	6	234 323 803	C/C	T/T

TABLE 12-continued

SNP analysis on chromosome 6						
SNP marker	Sequence	Sequence Orientation	Chromosome	Position in bp (CM334 v. 1.55)	LT17	PBC167
PE-0020909	TACTCCTTCAGTGGATCTTAAGGTGAACCAAGTGGAAATTCAGGATAC[C/G]A GTCAAAGCTGCCATATTCTCTTGTGTACCTTGTAGTTGCAGCATCAT (SEQ ID NO: 67)	Top	6	234 848 256	C/C	G/G
PE-0000401	TTGAATTAGTGTGTAATGCAAGGATCTTAAAGTGAAGAGAAATCACCCCCAG GCATTGCAACTGCCATTCCGCGAGACGAGGA[A/G]CTTGACACACTTATCAA GGACCAATCTGGTGGTGTGATCCCTCACATTCACAAAGTCCCTTAATAAA AAACCTCCAAGGAGTGAATTCTCCTTTTtATCTTAGATGTATCATG (SEQ ID NO: 68)	Top	6	235 746 196	A/A	G/G
PE-0003984	TAAATGTTCAACAAGAGTCCATTCTACAGCAAAATGAGCAAGTTTGGTAAAGGG CCAA[C/T]GCGATGAGCCTGTTTCTTTATATACTTAGGGAGGAAACTCAACAT GTGATAAGTACA (SEQ ID NO: 69)	Bottom	6	235 888 699	T/T	C/C
PE-0014672	GTTCCTTAGTCTAAGAGAGACAATAGCTCTTTTAAAGCTTAATTCACCAC[C/T]ATT ATCTACTTGCACCTTGAAAAGTCCAAAGAGAAAAAGGGACAAAAC (SEQ ID NO: 70)	Bottom	6	236 289 525	C/C	T/T
PE-0005702	ACAATAGTTTTATCTCAGCACTAAATGGTTGAGCACCTGTGTTCTGTGAAGTAA TGTG[A/G]ACAATTTTGTGTAGGATTTCCATGTACCTGCAATTGAATCACATATTAC AGCGGAGTTA (SEQ ID NO: 71)	Top	6	236 546 851	G/G	A/A
PE-0021582	CTAATAATCAAGATGAGAATATTAAATATGGTGAATCAGCGGATGATGAT[A/T]TAC AAATGTTGCAGCAGCAGCCTACTGATTTTGACCCCAATGGAGGATACT (SEQ ID NO: 72)	Top	6	236 574 405	A/A	T/T
PE-0007248	GTGAAGTTATGGAATAAATCAGTAGAGGATCTGATGTCGGAGCACTAGGGTA AAGAGT[C/A]TCGCGTCTCATGTTGTAGTGTATTGTCAATCAATAATTTCGATTTTC TGCTTCTTAGTCA (SEQ ID NO: 73)	Top	6	236 762 040	A/A	C/C
PE-0003669	CTCAAATAGCCGGGAATTCATTACATTTCCCTTGTATTAACAATTATCAATACTG AGCT[C/T]CTATTCTTGACTGACTATGCACTCAGGTAGGAATTCATACATACAACT CCCCAAAAGG (SEQ ID NO: 74)	Bottom	6	236 880 605	T/T	C/C
PE-0014672	GTTCCTTAGTCTAAGAGAGACAATFAGCTCTTTTAAAGCTTAATTCACCAC[C/T]ATT ATCTACTTGCACCTTGAAAAGTCCAAAGAGAAAAAGGGACAAAAC (SEQ ID NO: 75)	Bottom	6	236 289 525	C/C	T/T

Example 10: Validation of the Effects of QTL on Chromosome 6 in Susceptible Background Though QTL Introgression and Evaluation of Dominance Effects

[0368] A three-way hybrid F2 population was created as follows: The LT17 source was crossed by an intermediate genotype (HPG117) and then the obtained hybrid was crossed again by a susceptible bell pepper line (BYP). This 3-way hybrid was then selfed to obtain a F2 population segregating for the QTL on chromosome 6 within susceptible background.

[0369] Based on the markers described in examples 6 and 8, a total of 180 F2 plants were genotyped to select different plants with different genotype profile in the QTL region. A total of 20 F2 plants were selected for each of the 3 following QTL profile: homozygous favorable allele, homozygous unfavorable allele and heterozygous.

[0370] All selected F2 plants were evaluated in artificial test following the protocol described in Example 2. Selected plants having the same QTL profile were grouped in two plots of 10 plants organized in 2 repetitions of 5 plants each, leading to a total of 6 F2 plots of 10 plants. In addition to the selected F2s, 1 F4 (Maor*LT17) genotype being homozygous for favorable alleles was included in the test, as described below in Example 11.

[0371] Results are presented in FIG. 5. Results show a clear effect of QTL6 with an additive effect. For the F4 (Maor*LT17) the seed lot F4 (ST117F04-1157/BLK) is well resistant. Seeds of the seed lot F4 (ST117F04-1157/BLK) were thus selected for deposit at the NCIMB under accession number NCIMB 43602.

Example 11: Validation of Candidate Genes within QTL6 Interval

[0372] F2 seeds were produced from a cross between Maor (used as female) and LT17 (used as male). F2 plants have been evaluated under artificial inoculation (as described in Example 6). F3 seeds from genotypically selected F2 plants harboring QTL6 were harvested then selfed to obtain F4 (Maor*LT17) seeds (ST117F04-1157/BLK) that were used as females pollinated with Pipo (Susceptible genotype) pollen. F1 [F4 (Maor*LT17)*Pipo] seeds were used to generate the population F2[F4 (Maor*LT17)*Pipo]. A total of 5940 F2 plants derived from this population were genotyped with markers developed within the 2 candidate genes as described in examples 6 and 8. Out of 5940 F2 plants genotyped, a total of 131 F2 plants were selected according to their genotypic profile for the 2 candidate genes in order to evaluate gene effects in artificial test. The artificial test was conducted according to the protocol described in Example 2. Plants selection were organized in plots of 10 plants to conduct the artificial test.

[0373] Below are molecular profiles of the plants, and the results obtained from artificial test scoring:

NAME	Gene CA06g26250	Gene CA06g26150	Number of tested F2 plants	Mean score
F2[(Maor × LT17)*Pipo]	Homozygous unfavorable	Homozygous favorable	1	8.0
F2[(Maor × LT17)*Pipo]	Homozygous favorable	Homozygous favorable	60	6.3

-continued

NAME	Gene CA06g26250	Gene CA06g26150	Number of tested F2 plants	Mean score
F2[(Maor × LT17)*Pipo]	Heterozygous	Homozygous unfavorable	6	3.6
F2[(Maor × LT17)*Pipo]	Homozygous unfavorable	Heterozygous	4	3.5
F2[(Maor × LT17)*Pipo]	Homozygous unfavorable	Homozygous unfavorable	60	2.6

[0374] These results confirm the effect of QTL and suggest a dominant mechanism involving the CA06g26150 gene encoding an MLO-like protein.

Example 12: Genetic Modification of *Capsicum* Seeds by Ethyl Methane Sulfonate (EMS)

[0375] Seeds of *Capsicum* plants are to be treated with EMS by submergence of approximately 2000 seeds into an aerated solution of either 0.5% (w/v) or 0.7% EMS for 24 hours at room temperature.

[0376] Approximately 1500 treated seeds per EMS dose are germinated and the resulting plants are grown, preferably in a greenhouse to produce seeds.

[0377] Following maturation, M2 seeds are harvested and bulked in one pool per variety per treatment. The resulting pools of M2 seeds are used as starting material to identify the individual M2 seeds and the plants resistant to powdery mildew.

BIBLIOGRAPHY

- [0378]** Daubeze, A. M. et al (1995). Resistance to *Leveillula taurica* in pepper (*Capsicum annuum*) is oligogenically controlled and stable in Mediterranean regions. *Plant Breeding*, 114(4), 327-332.
- [0379]** George, E. F. and Sherrington, P. D. (1984) Plant Propagation by Tissue Culture, Handbook and Directory of Commercial Laboratories, Exegetics Ltd
- [0380]** Gnirke, A. et al (2009). Solution hybrid selection with ultra-long oligonucleotides for massively parallel targeted sequencing. *Nature biotechnology*, 27(2), 182-189.
- [0381]** Hui (Y. H.), Sherkat F., Handbook of Food Science, Technology and Engineering, vol. 4, CRC Press
- [0382]** Kim, D. S., & Hwang, B. K. (2012). The pepper MLO gene, CaMLO2, is involved in the susceptibility cell-death response and bacterial and oomycete proliferation. *The Plant Journal*, 72(5), 843-855.
- [0383]** Lefebvre, V., Daubèze, A. M., van der Voort, J. R., Peleman, J., Bardin, M., & Palloix, A. (2003). QTLs for resistance to powdery mildew in pepper under natural and artificial infections. *Theoretical and Applied Genetics*, 107(4), 661-666.
- [0384]** Lim, C. W., & Lee, S. C. (2014). Functional roles of the pepper MLO protein gene, CaMLO2, in abscisic acid signaling and drought sensitivity. *Plant molecular biology*, 85(1-2), 1-10.
- [0385]** Molot, Jp et al. *Leveillula taurica* (Lév) Arn: cultures axéniques, biologie et spécificité parasitaire. Agronomie, EDP Sciences, 1990, 10 (7), pp. 551-559.
- [0386]** Zheng, Z. et al (2013). Loss of function in Mlo orthologs reduces susceptibility of pepper and tomato to powdery mildew disease caused by *Leveillula taurica*. *PLoS one*. 8(7), e70723.

```

Sequence total quantity: 75
SEQ ID NO: 1          moltype = DNA   length = 101
FEATURE               Location/Qualifiers
variation             51
                      note = Polymorphism: Susceptible=G;Resistant=A
source                1..101
                      mol_type = unassigned DNA
                      organism = Capsicum annuum

SEQUENCE: 1
aagatactct gcattgggtc cgggctagaa gaataagcac gaataaccgc rttccaagaa 60
aatagatttg gttgaggaat ttgatcgaac acttggtggg c 101

SEQ ID NO: 2          moltype = DNA   length = 46
FEATURE               Location/Qualifiers
misc_feature          1..46
                      note = sequence of a forward primer for detecting the
source                1..46                          susceptible alleleof PE-0014628
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 2
gaagggtcgga gtcaacggat tagaagaata agcacgaata accgcg 46

SEQ ID NO: 3          moltype = DNA   length = 46
FEATURE               Location/Qualifiers
misc_feature          1..46
                      note = sequence of a forward primer for detecting the
source                1..46                          resistant alleleof PE-0014628
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 3
gaagggtgacc aagttcatgc tagaagaata agcacgaata accgca 46

SEQ ID NO: 4          moltype = DNA   length = 30
FEATURE               Location/Qualifiers
misc_feature          1..30
                      note = sequence of a common reverse primer for PE-0014628
source                1..30
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 4
aattcctcaa ccaaatctat tttcttgga 30

SEQ ID NO: 5          moltype = DNA   length = 301
FEATURE               Location/Qualifiers
variation             151
                      note = Polymorphism: Susceptible=T;Resistant=C
source                1..301
                      mol_type = unassigned DNA
                      organism = Capsicum annuum

SEQUENCE: 5
atattctagg ttactggatt taactcagga gaaaatggat ttgtttttgc aatgcatttt 60
tcggggggtg cctagtttat aaagacctaa gatgtttcag tcaactgctt aattaggacc 120
tttgactggt gtggttatga ctgggccctt ygccaagatg gttttggaca acagagctct 180
taccacactt gctgctattt tgatctactt gatcttgagt tgcgctgttt gctgtgctgt 240
tcaaagcgac attgactgtc tgagatctat aaaagattca ttagaagatc caaataatac 300
t 301

SEQ ID NO: 6          moltype = DNA   length = 42
FEATURE               Location/Qualifiers
misc_feature          1..42
                      note = sequence of a forward primer for detecting the
source                1..42                          susceptible alleleof PE-0056542
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 6
gaagggtcgga gtcaacggat tgtggttatg actgggcccc tt 42

SEQ ID NO: 7          moltype = DNA   length = 42
FEATURE               Location/Qualifiers
misc_feature          1..42
                      note = sequence of a forward primer for detecting the
                      resistant alleleof PE-0056542

```

-continued

```

source                1..42
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 7
gaaggtgacc aagttcatgc tgtggttatg actgggcccc tc                42

SEQ ID NO: 8          moltype = DNA length = 28
FEATURE              Location/Qualifiers
misc_feature          1..28
                      note = sequence of a common reverse primer for PE-0056542
source                1..28
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 8
taagagctct gttgtccaaa accatctt                                28

SEQ ID NO: 9          moltype = DNA length = 301
FEATURE              Location/Qualifiers
variation             151
                      note = Polymorphism: Susceptible=G;Resistant=A
source                1..301
                      mol_type = unassigned DNA
                      organism = Capsicum annuum

SEQUENCE: 9
agatgggttt ggacaacaga gctcttacca cacttgctgc tattttgatc tacttgatct 60
tgagttgcgc tgtttgctgt gctgttcaaa gcgacattga ctgtctgaga tctataaaag 120
attcattaga agatccaaat aatactttga rctatacatg gaagttcgat aatctgacgg 180
aagggttcat ttgcaagttt gctgggattc agtgctggca tctgatgaa aataaggttc 240
tgagtatcag tcttctgac atgggtctca agggcaggtt tccacttggt ctctcggaatt 300
g                                                                301

SEQ ID NO: 10         moltype = DNA length = 49
FEATURE              Location/Qualifiers
misc_feature          1..49
                      note = sequence of a forward primer for detecting the
                      susceptible alleleof PE-0056543
source                1..49
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 10
gaaggtcgga gtcaacggat tcattagaag atccaaataa tactttgag        49

SEQ ID NO: 11         moltype = DNA length = 54
FEATURE              Location/Qualifiers
misc_feature          1..54
                      note = sequence of a forward primer for detecting the
                      resistant alleleof PE-0056543
source                1..54
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 11
gaaggtgacc aagttcatgc tagattcatt agaagatcca aataatactt tgaa    54

SEQ ID NO: 12         moltype = DNA length = 29
FEATURE              Location/Qualifiers
misc_feature          1..29
                      note = sequence of a common reverse primer for PE-0056543
source                1..29
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 12
atgaaacctt ccgtcagatt atcgaaactt                                29

SEQ ID NO: 13         moltype = DNA length = 301
FEATURE              Location/Qualifiers
variation             151
                      note = Polymorphism: Susceptible=G;Resistant=T
source                1..301
                      mol_type = unassigned DNA
                      organism = Capsicum annuum

SEQUENCE: 13
ctgttctctg agagagctat gccaaataac cgggactgtg tggagaaccg ttagatcctt 60
gccagggtac ttctaagaaa cccctaccg caatcattgt cgggtgcagct gttggtgggg 120
taagtttagg tgctgtatct ttggccatcg ktagttctt ctatatgaga aggatatcca 180
ggaagaggaa gaaggatgat gatcctgaag ggaataaatg ggccaagagt attaagggtta 240
aaaaggcaat ccaggtgaaga attttaactt ctctgaatca ttaaaaactg tctcgttact 300

```

-continued

g 301

SEQ ID NO: 14 moltype = DNA length = 44
 FEATURE Location/Qualifiers
 misc_feature 1..44
 note = sequence of a forward primer for detecting the
 susceptible alleleof PE-0056544
 source 1..44
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 14
 gaaggtgacc aagttcatgc tgggtgctgta tttttggcca tcgg 44

SEQ ID NO: 15 moltype = DNA length = 45
 FEATURE Location/Qualifiers
 misc_feature 1..45
 note = sequence of a forward primer for detecting the
 resistant alleleof PE-0056544
 source 1..45
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 15
 gaaggtcgga gtcaacggat taggtgctgt atttttggcc atcgt 45

SEQ ID NO: 16 moltype = DNA length = 29
 FEATURE Location/Qualifiers
 misc_feature 1..29
 note = sequence of a common reverse primer for PE-0056544
 source 1..29
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 16
 ttcctcttcc tggatatacct tctcatata 29

SEQ ID NO: 17 moltype = DNA length = 301
 FEATURE Location/Qualifiers
 variation 151
 note = Polymorphism: Susceptible=T;Resistant=C
 source 1..301
 mol_type = unassigned DNA
 organism = Capsicum annuum

SEQUENCE: 17
 tcttttcata gaagcattta actaatgaat tttcacctgc tccgctgctc gtgtgtgatt 60
 tttctcttgt tcaagttttc ttaattgact taaggttcct tattttcagc tttcaatgtt 120
 tgagaagtct acttcaaaaa tgagatttag ygatcttatg aaggccactg acagcttcaa 180
 caaaaacaat attatcgggt caggaagaac cgggactttt taaaagcag tacttgatga 240
 tggcacttca cttatgggtta agagggttga gaataactcag cactcagaga aagagtttat 300
 g 301

SEQ ID NO: 18 moltype = DNA length = 45
 FEATURE Location/Qualifiers
 misc_feature 1..45
 note = sequence of a forward primer for detecting the
 susceptible alleleof PE-0056545
 source 1..45
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 18
 gaaggtcgga gtcaacggat tctgtcagtg gccctcataa gatca 45

SEQ ID NO: 19 moltype = DNA length = 43
 FEATURE Location/Qualifiers
 misc_feature 1..43
 note = sequence of a forward primer for detecting the
 resistant alleleof PE-0056545
 source 1..43
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 19
 gaaggtgacc aagttcatgc tgtcagtggc cttcataaga tcg 43

SEQ ID NO: 20 moltype = DNA length = 30
 FEATURE Location/Qualifiers
 misc_feature 1..30
 note = sequence of a common reverse primer for PE-0056545
 source 1..30

-continued

```

mol_type = other DNA
organism = synthetic construct

SEQUENCE: 20
gtttgagaag tctacttcaa aaatgagatt 30

SEQ ID NO: 21      moltype = DNA length = 301
FEATURE           Location/Qualifiers
variation         151
                  note = Polymorphism: Susceptible=A ; Resistant=C
source            1..301
                  mol_type = unassigned DNA
                  organism = Capsicum annuum

SEQUENCE: 21
actcagcact cagagaaaga gtttatgtcc gagatggcta cattgggaaa tgtaaagcat 60
cgtaatctgg ttctctcttt aggtttctgc attgccaaaa aagaaaggct gttggtctac 120
aaagacatgc caaatggaac ctgcatgat mggttacatt cggtagtgta aggcgaaaaa 180
actctagagt ggctataaag attaaaaata ggcattggag cagccaaagg atttgcattg 240
cttcaccaca actgcaatcc tcgtatcatt cacagaaata ttagttctaa atgcatcttg 300
c 301

SEQ ID NO: 22      moltype = DNA length = 44
FEATURE           Location/Qualifiers
misc_feature      1..44
                  note = sequence of a forward primer for detecting the
                  susceptible alleleof PE-0056546
source            1..44
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 22
gaaggtgacc aagttcatgc tccttcactc accgaatgta acct 44

SEQ ID NO: 23      moltype = DNA length = 44
FEATURE           Location/Qualifiers
misc_feature      1..44
                  note = sequence of a forward primer for detecting the
                  resistant alleleof PE-0056546
source            1..44
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 23
gaaggtcgga gtcaacggat tccttcactc accgaatgta accg 44

SEQ ID NO: 24      moltype = DNA length = 28
FEATURE           Location/Qualifiers
misc_feature      1..28
                  note = sequence of a common reverse primer for PE-0056546
source            1..28
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 24
ttggtctaca aagacatgcc aaatggaa 28

SEQ ID NO: 25      moltype = DNA length = 301
FEATURE           Location/Qualifiers
variation         151
                  note = Polymorphism: Susceptible=A ; Resistant=C
source            1..301
                  mol_type = unassigned DNA
                  organism = Capsicum annuum

SEQUENCE: 25
gctactccaa aaggtgatgt gtacagtttc ggtgttgtag ttctcgagtt agttactggt 60
gagaaaccga cctctgtcac caaagctccc gagaccttta aggggaattt ggtggaatgg 120
atcacacaac tctccggtga atctaagctt maagatgcga ttgaccattc attgtctggt 180
aaaggttatg acaatgagat ctccagggtc cttaaagggt cttgtcgatg tgtgttgtct 240
cctgttccta aggagaggcc gacaatgttt gaagtttacc agcttctgag agccatcgga 300
g 301

SEQ ID NO: 26      moltype = DNA length = 46
FEATURE           Location/Qualifiers
misc_feature      1..46
                  note = sequence of a forward primer for detecting the
                  susceptible alleleof PE-0056547
source            1..46
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 26

```

-continued

```

gaaggtgacc aagttcatgc tcaactctcc ggtgaatcta agctta          46

SEQ ID NO: 27          moltype = DNA length = 45
FEATURE               Location/Qualifiers
misc_feature          1..45
                      note = sequence of a forward primer for detecting the
                      resistant alleleof PE-0056547
source                1..45
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 27
gaaggtcgga gtcaacggat taactctccg gtgaatctaa gcttc          45

SEQ ID NO: 28          moltype = DNA length = 30
FEATURE               Location/Qualifiers
misc_feature          1..30
                      note = sequence of a common reverse primer for PE-0056547
source                1..30
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 28
cattgtcata acctttacca gacaatgaat          30

SEQ ID NO: 29          moltype = DNA length = 301
FEATURE               Location/Qualifiers
variation             151
                      note = Polymorphism: Susceptible= T; Resistant=A
source                1..301
                      mol_type = unassigned DNA
                      organism = Capsicum annuum

SEQUENCE: 29
taactgatgg ttctcagtga gctgcaaaat ggaacatcaa tatcagtagc ttctcgtttc 60
aaccattttac ttcccttgct ctttagtcac ttgcacttta gagaagaatg acgtgattcc 120
cgatatattta tacttagcta ttctagcata wgggaaacaa aacagatgaa tcatgaaact 180
cttcggtagt aaaaaaaatt atgtgccaaa atgttgtagt tatcttttta tggacgatga 240
tccataccat gttcggattc aggatccact ttacgattg catcctccca gctcttccat 300
t                                                    301

SEQ ID NO: 30          moltype = DNA length = 53
FEATURE               Location/Qualifiers
misc_feature          1..53
                      note = sequence of a forward primer for detecting the
                      susceptible alleleof PE-0056549
source                1..53
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 30
gaaggtcgga gtcaacggat tccgatattt tatacttagc tattctagca tat          53

SEQ ID NO: 31          moltype = DNA length = 53
FEATURE               Location/Qualifiers
misc_feature          1..53
                      note = sequence of a forward primer for detecting the
                      resistant alleleof PE-0056549
source                1..53
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 31
gaaggtgacc aagttcatgc tccgatattt tatacttagc tattctagca taa          53

SEQ ID NO: 32          moltype = DNA length = 30
FEATURE               Location/Qualifiers
misc_feature          1..30
                      note = sequence of a common reverse primer for PE-0056549
source                1..30
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 32
gagtttcatg attcatctgt ttgtttccc          30

SEQ ID NO: 33          moltype = DNA length = 301
FEATURE               Location/Qualifiers
variation             151
                      note = Polymorphism: Susceptible=C Resistant=T
source                1..301
                      mol_type = unassigned DNA

```

-continued

```

                                organism = Capsicum annuum
SEQUENCE: 33
tttgattttg ttttaactcat ttgcatgaaa tgatatgttt atatgtttgt ctagaaatat 60
cacaactatt acatttatag gtaatatattg agttgcactt atggtatagt gcatttttat 120
ttctatttat gtgacattat tttttagata yatgtcgaaa taatgctgga aaattaatgt 180
tcaactcttt gaatacatga aagaaccgaa acttctaccc tatttttaag ggatggacgc 240
gtgaattaga cctaattaag aatatattat cttacacttt ttgttttadc gagaagcttt 300
t                                                                    301

SEQ ID NO: 34                moltype = DNA length = 51
FEATURE                      Location/Qualifiers
misc_feature                  1..51
                                note = sequence of a forward primer for detecting the
                                susceptible alleleof PE-0056550
source                        1..51
                                mol_type = other DNA
                                organism = synthetic construct
SEQUENCE: 34
gaaggtgacc aagttcatgc tacattaatt ttccagcatt atttcgacat g          51

SEQ ID NO: 35                moltype = DNA length = 53
FEATURE                      Location/Qualifiers
misc_feature                  1..53
                                note = sequence of a forward primer for detecting the
                                resistant alleleof PE-0056550
source                        1..53
                                mol_type = other DNA
                                organism = synthetic construct
SEQUENCE: 35
gaaggtcgga gtcaacggat tgaacattaa ttttcagca ttatttcgac ata      53

SEQ ID NO: 36                moltype = DNA length = 30
FEATURE                      Location/Qualifiers
misc_feature                  1..30
                                note = sequence of a common reverse primer for PE-0056550
source                        1..30
                                mol_type = other DNA
                                organism = synthetic construct
SEQUENCE: 36
taggtaatat ttgagttgca cttatggat          30

SEQ ID NO: 37                moltype = DNA length = 101
FEATURE                      Location/Qualifiers
variation                    51
                                note = Polymorphism: Susceptible=G;Resistant=C
source                        1..101
                                mol_type = unassigned DNA
                                organism = Capsicum annuum
SEQUENCE: 37
ctccaaaacc acagctacta atttggttaga tccaaaactt aaaagcttat stttagtcag 60
tcaattaaaa atcccaacaa tttaaaagct aagaagtga g          101

SEQ ID NO: 38                moltype = DNA length = 49
FEATURE                      Location/Qualifiers
misc_feature                  1..49
                                note = sequence of a forward primer for detecting the
                                susceptible alleleof PE-0021476
source                        1..49
                                mol_type = other DNA
                                organism = synthetic construct
SEQUENCE: 38
gaaggtcgga gtcaacggat tgttgggatt tttaattgac tgactaaac          49

SEQ ID NO: 39                moltype = DNA length = 50
FEATURE                      Location/Qualifiers
misc_feature                  1..50
                                note = sequence of a forward primer for detecting the
                                resistant alleleof PE-0021476
source                        1..50
                                mol_type = other DNA
                                organism = synthetic construct
SEQUENCE: 39
gaaggtgacc aagttcatgc ttgttgggat ttttaattga ctgactaaag          50

SEQ ID NO: 40                moltype = DNA length = 27
FEATURE                      Location/Qualifiers

```

-continued

misc_feature 1..27
 note = sequence of a common reverse primer for PE-0021476
 source 1..27
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 40
 ctccaaaacc acagctacta atttggt 27

SEQ ID NO: 41 moltype = DNA length = 1557
 FEATURE Location/Qualifiers
 source 1..1557
 mol_type = unassigned DNA
 organism = Capsicum annuum

SEQUENCE: 41
 atgggtggag gtggagaaga tgaagggtca ttggaattta caccaacatg ggttgttgc 60
 ggtgtagtca ctgtcattgt tgaatatct ctaactcttg aacgcctcct ccattatact 120
 ggaaagcatt tgaagaagaa gaagcaaaag catctctatg aagcactaca gaaagttaaa 180
 gaagagtga tgctcttggg gtccatttcc cttttgttaa cggttattca aagtcgaatt 240
 gtcaaaatct gtgtgcctca tctgtctgta acacacttac ttccatgctc attatcattg 300
 gaggggttctt caccattcagg ttcatcggtg ggggggttctc catctccatc tccacctgaa 360
 gtgcctcgctc ctagcaacaa ttctgttcac tatcatcctg gtcctcacca tcaacggcat 420
 ttgcttgtag cagaagagag tcctaagggt tactgcagtc ataaacacaa ggtcccggtta 480
 ctatctcttg aggcgttgca tcaccttcac atttttatct ttgtcctagc tattgtgcat 540
 gtgacttttt ctgttctgac tattgtattt ggaggagtaa agatacgtca atggaagagc 600
 tgggaggatg caatcgtaaa atgtgacatc gaatccgaac atgctcaact gaaaccatca 660
 gttactcatg tccatgaaca tgatttcac aggaatcggg ttccagggtat cggtaaacaa 720
 tcactcggtta tgggttgggc gcattctttc ttcaagcaat tttatgcttc tgtaaacgag 780
 tcagactaca gaggcgttctg ttgggggttc attatgacac attgcagggg aaatccaaga 840
 ttcaatttct ataggtagcat ggtacgtgct ctggaagacg attttacgac agttgtcggg 900
 atcaggcaag ttggcatatg tatttctgga ttgctttcgt tccttttata tacaatctct 960
 atgcaggagg aagctaactt gatttctgtg tcctttcgtt cttcttttct tttcgtgttt 1020
 tttttccagc ttctacttgc agttggaaca aagttggagc atgtgattat acagttagct 1080
 catgaaattg ctgagagaca tctagcagta gaaggtgaat tgattgtaac accatccgat 1140
 gatcaactttt ggttcgctgc cctcaaat atcctcttct tgatacaatt tattctattc 1200
 caaaatgctt ttgaaatagc atttttcttc tggattttgt tccaatatgg ctttttctcc 1260
 tgcattatgg gaaatacgt gtttgttatt cctcggtctg tcataggggt tatcattcaa 1320
 gttttatgca ccatatagcact atggcactt tatgctctag ttacacagat ggggagtcac 1380
 tataagaagt caatgttcga taaacatgtt caagcttgcc tcctgggatt ggctgataag 1440
 gcaaaaaaga agaaggagct taagattgct agccctgtcc attcaattga cggttctgtt 1500
 caaggatcac caacgtccgt aagtgggcgc aaagactacc ctcaaatag ggtttag 1557

SEQ ID NO: 42 moltype = DNA length = 1866
 FEATURE Location/Qualifiers
 source 1..1866
 mol_type = unassigned DNA
 organism = Capsicum annuum

SEQUENCE: 42
 atgactgggc cccttgccaa gatgggtttg gacaacagag ctcttaccac acttgcgtgc 60
 attttgatct acttgatctt gagttgcgtg gtttgcgtg ctgttcaaa gacattgac 120
 tgtctgagat ctataaaaga ttcatagaa gatccaaata atactttgag ctatacatgg 180
 aagttcgata atctgacgga aggtttcatt tgcagggttg ctgggattca gtgctggcat 240
 cctgatgaaa atagggttct gagtatcagt cttcctgaca tgggtctcaa gggcagggtt 300
 ccacttggtc ttccgaattg caccctctatg acatcattag atctttcgag caacaatctc 360
 catggaagca ttccatctga tatctccaaa tatataggtt tcgtggtaac cctggatctc 420
 tcactcaacc aactatcagg ggaatccca gttaactctg caaactgctc ttttttgaat 480
 aaccttaaac ttgaaaaaca ccagtttacg ggtcaaatcc ctccagaaat tggcctgctt 540
 ggtcgctca aaacttttaa ttagcaaac aatcgattga ctggaccagt tccacatttc 600
 ataaatgcaa ctgttcctgc agagagctat gccataatc cgggactgtg tggagaaccg 660
 ttagatcctt gccagggtac ttctaagaaa cccctaccg caatcattgt cgggtgcagt 720
 gttgtgggg taagtttagt tgctgtattt ttggccatcg gtatgttctt ctatatgaga 780
 aggatataca ggaagaggaa gaaggatgat gatcctgaag ggaataaatg ggcgaagagt 840
 attaagggtt aaaaaggcaat ccagctttca atgtttgaga agtctacttc aaaaatgaga 900
 tttagtgtac ttatgaaggc cactgacagc ttcaacaaaa acaatattat cgggtcagga 960
 agaaccggga ctttttacaa agcagtagct gatgatgcca cttcacttat ggttaagagg 1020
 tgcagaata ctacgactc agagaagag tttatgtccg agatggctac attgggaaat 1080
 gtaaaagcat gtaatctggt tcctctttta ggtttctgca ttgcaaaaa agaaaggctg 1140
 ttggtctaca aagacatgcc aaatggaacc ttgcatgata gggtacattc ggtgagtgaa 1200
 ggcaaaaaaa ctctagagt gctataaga ttaaaaaatg gcattggagc agccaaagga 1260
 tttgcatggc ttaccacaa ctgcaatcct cgtatcattc acagaaatat tagttctaaa 1320
 tgcattctgc tggatgtgga atttgaacct aggatatcag attttggggt tgcagggcta 1380
 atgaaatcaa ttgacactca tttagtacc tttgtcaatg gagaattcgg ggaattcgg 1440
 tatgttgcct ctgagtacac acgaactctc atggctactc caaaagggtg tgtgtacagt 1500
 ttcggtgttg tacttctcga gttagtact ggtgagaaac cgacctctgt caccaaagct 1560
 cccgagacct ttaaggggaa tttggtggaa tggatcacac aactctccgg tgaatctaag 1620
 cttaaagatg cgaattgacca ttcatgtct ggtaagggt atgacaatga gatctccag 1680
 gtccttaaa ttgcttgcgt atgtgtgttg tctcctgttc ctaaggagag gccgacaatg 1740

-continued

```

tttgaagttt accagcttct gagagccatc ggagagcgat atcatttcac aactgatgat 1800
gacattttga tgctttcaga gagtgatggc ggattccaaa tggacgagct cattgttgct 1860
caataa 1866

SEQ ID NO: 43      moltype = DNA length = 23
FEATURE           Location/Qualifiers
misc_feature       1..23
                   note = sequence of a forward primer for amplifying the
                   susceptible allele of the RZ marker
source            1..23
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 43
catgtgagcg actcctctga caa 23

SEQ ID NO: 44      moltype = DNA length = 22
FEATURE           Location/Qualifiers
misc_feature       1..22
                   note = sequence of a forward primer for amplifying the
                   resistant allele of the RZ marker
source            1..22
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 44
atgtgagcga ctctcttgac ag 22

SEQ ID NO: 45      moltype = DNA length = 29
FEATURE           Location/Qualifiers
misc_feature       1..29
                   note = sequence of a common reverse primer for amplifying
                   the resistant or susceptible allele of the RZ marker
source            1..29
                   mol_type = other DNA
                   organism = synthetic construct

SEQUENCE: 45
cgaggagaat ttcttgctaa aattgagtt 29

SEQ ID NO: 46      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=C; PBC167=G
source            1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 46
ttgaaaccct ccttccttcc ataccattgaa agaaatgggtg gatggagact sagcaaatca 60
ttggtcacat catacaacct caagagcata ctagaagctg a 101

SEQ ID NO: 47      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=A; PBC167=G
source            1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 47
agttgccttg gtaccaatga aattgactct tcttttcctt ccaagtccac rgctccagga 60
cctgcattca gatccagacc ttgccggctt cggcttctgt t 101

SEQ ID NO: 48      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=C; PBC167=A
source            1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 48
ggtggtagaa tcctccctcg tcgctctgac ggcacgctcc gttacatcgg mggttttact 60
agagtcctct cgttgatcg gtctatttct tattcagagt t 101

SEQ ID NO: 49      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=T; PBC167=G
source            1..101
                   mol_type = unassigned DNA

```

-continued

```

                                organism = Capsicum annuum
SEQUENCE: 49
tggttctaata gtggtgacaa aggttccatc aagtgaatgt tatggtgatg ktggtgtaag 60
tgctaactgtg cagaagagta aagcaaatga agcaaacggg g 101

SEQ ID NO: 50      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=T; PBC167=C
source            1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 50
gagatagaaa atgaaggtga aaaaaacttt ggatctccag ctgcaccagt ycaggaacga 60
ccaattcaga gatcatgggt tcctccccag cctcctccag t 101

SEQ ID NO: 51      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=C; PBC167=A
source            1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 51
aaaagaatcc gccctaactg ctttctgctc ttcttcttca cattgaattc mtcggcacga 60
tctagaacca aattcatgta ctcatacaaaa ccaataatac g 101

SEQ ID NO: 52      moltype = DNA length = 101
FEATURE           Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=G; PBC167=A
source            1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 52
ataatgatgt caccagtctg aactaccgaa gacagttctt tgcttcaatc ratatgctac 60
aagaacaagc tgagaaatta cttgaagaga tgaagcctaa g 101

SEQ ID NO: 53      moltype = DNA length = 121
FEATURE           Location/Qualifiers
variation          61
                   note = Polymorphism: LT17=G; PBC167=A
source            1..121
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 53
gtccctcttt caagagttaa agaagtgggtg tgtgggattt gctttgacta ctaccgcct 60
ratgacataa tgtctgttgg ttgtggtcat cctttttgta ctatttgctg gaaagcttac 120
a 121

SEQ ID NO: 54      moltype = DNA length = 208
FEATURE           Location/Qualifiers
variation          59
                   note = Polymorphism: LT17=T; PBC167=G
source            1..208
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 54
aktaaagaga acacctgtat tcaccagtac tactaccagc aatgggtcaa ttacaccka 60
tcttggaaca aatgatgaca tccatccctc ctatcttatt gaagcatgct acattcagtc 120
ttgcactttg acagataagt gtttcaccac atatccaata agatgcacac catataacat 180
tgatgctggc aagagtatgc acctgcag 208

SEQ ID NO: 55      moltype = DNA length = 121
FEATURE           Location/Qualifiers
variation          61
                   note = Polymorphism: LT17=C; PBC167=A
source            1..121
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 55
ggaggcaaca acacaaagta tcccactacc ccccactaag atgggggatag tgaacaagg 60
matagtgtcc attcttggat cagactctga caaaagaaac aaggctgctg cagctgtttc 120
a 121

SEQ ID NO: 56      moltype = DNA length = 121

```

-continued

FEATURE	Location/Qualifiers	
variation	61	
	note = Polymorphism: LT17=T; PBC167=C	
source	1..121	
	mol_type = unassigned DNA	
	organism = Capsicum annuum	
SEQUENCE: 56		
gaacgcttat gggattgttc caggggtaaa acattacgct tgcttagttg atatacttgg	60	
ycgtgcaggt gaactggaag aagctgaaga actgataaat aatatgccag taacccttgg	120	
t	121	
SEQ ID NO: 57	moltype = DNA length = 121	
FEATURE	Location/Qualifiers	
variation	61	
	note = Polymorphism: LT17=A; PBC167=G	
source	1..121	
	mol_type = unassigned DNA	
	organism = Capsicum annuum	
SEQUENCE: 57		
cggatattgg agaagatgaa ggctcctaaa aagccagatg atgtgcctga gatcacgcct	60	
ragcagctct tggaagctga tggtttcac tttggtttcc cttcgcgttt tgggtgtgatg	120	
g	121	
SEQ ID NO: 58	moltype = DNA length = 121	
FEATURE	Location/Qualifiers	
variation	61	
	note = Polymorphism: LT17=A; PBC167=G	
source	1..121	
	mol_type = unassigned DNA	
	organism = Capsicum annuum	
SEQUENCE: 58		
tcggttgctt gcaggttatt gcagatttgg tgagaccaga caaggaatca aaaataagaa	60	
ratactggaa ttggtaccat tacacagcag gaagaatttt gatattattg gcatcaataa	120	
a	121	
SEQ ID NO: 59	moltype = DNA length = 101	
FEATURE	Location/Qualifiers	
variation	51	
	note = Polymorphism: LT17=G; PBC167=C	
source	1..101	
	mol_type = unassigned DNA	
	organism = Capsicum annuum	
SEQUENCE: 59		
atgtcttaca aacctccatt gacaacaaag attacattgc gggcaaactc stccacgccc	60	
atatactcgg acttggcctt tcctctcata cactcctcct a	101	
SEQ ID NO: 60	moltype = DNA length = 121	
FEATURE	Location/Qualifiers	
variation	61	
	note = Polymorphism: LT17=A; PBC167=T	
source	1..121	
	mol_type = unassigned DNA	
	organism = Capsicum annuum	
SEQUENCE: 60		
tatacaagas aatgttggaa ggggaatggg aaaatggaat atcagctggc acaattgtcc	60	
wagcagcagt agctttattg cttgatggaa catcattgta tagattcatg aaggcagagg	120	
g	121	
SEQ ID NO: 61	moltype = DNA length = 121	
FEATURE	Location/Qualifiers	
variation	61	
	note = Polymorphism: LT17=A; PBC167=C	
source	1..121	
	mol_type = unassigned DNA	
	organism = Capsicum annuum	
SEQUENCE: 61		
tgtgtagttg ttggcattaa aatttgcttc tagcctttca gaaaagacat tacgttgtaa	60	
mettgtagtg tgtatttaat tctcttgaca tgaaatgaag tcgagtggac acaacaaaaa	120	
a	121	
SEQ ID NO: 62	moltype = DNA length = 121	
FEATURE	Location/Qualifiers	
variation	61	
	note = Polymorphism: LT17=A; PBC167=G	
source	1..121	
	mol_type = unassigned DNA	

-continued

```

                                organism = Capsicum annuum
SEQUENCE: 62
ctaggcctag ttgatgggac tacacaatgg tatgcatgca aactaagcaa aagatggggc 60
rcgatggagg aaaaagatct gcataccagc aacttggtgg gcatatattt tagtcatgct 120
a                                                    121

SEQ ID NO: 63                moltype = DNA length = 101
FEATURE                      Location/Qualifiers
variation                    51
                                note = Polymorphism: LT17=T; PBC167=C
source                      1..101
                                mol_type = unassigned DNA
                                organism = Capsicum annuum

SEQUENCE: 63
gaatgttttt gcgttttggg gtaccacact ctcttgatca tgctcaaatg yttcaaaagc 60
atacatgcat ttaaagaatc cttcagcagc aaatccagaa g 101

SEQ ID NO: 64                moltype = DNA length = 101
FEATURE                      Location/Qualifiers
variation                    51
                                note = Polymorphism: LT17=A; PBC167=C
source                      1..101
                                mol_type = unassigned DNA
                                organism = Capsicum annuum

SEQUENCE: 64
tgaatccaaa cagatttggg acactgactg ggactgatga actcataagt mttgcctttc 60
tttgctcctt tttgtttcgt tttcttcctt tctgttggtt a 101

SEQ ID NO: 65                moltype = DNA length = 101
FEATURE                      Location/Qualifiers
variation                    51
                                note = Polymorphism: LT17=A; PBC167=G
source                      1..101
                                mol_type = unassigned DNA
                                organism = Capsicum annuum

SEQUENCE: 65
tgatcataat atcaaatctg atgttcaaca atacaatatg cacaagagga raaaaactgg 60
ttcataaatg gtgaatcaag aaggcaatac gacgagcctt t 101

SEQ ID NO: 66                moltype = DNA length = 121
FEATURE                      Location/Qualifiers
variation                    61
                                note = Polymorphism: LT17=C; PBC167=T
source                      1..121
                                mol_type = unassigned DNA
                                organism = Capsicum annuum

SEQUENCE: 66
ctcccgatcat ctgttaaatg cttcattgct tcagacatta ttaagtgcta gaacttcagg 60
ygttcgttca ccttggtata ttgtacatga caatgttgat gcatgtggta gcatttggtt 120
g                                                    121

SEQ ID NO: 67                moltype = DNA length = 101
FEATURE                      Location/Qualifiers
variation                    51
                                note = Polymorphism: LT17=C; PBC167=G
source                      1..101
                                mol_type = unassigned DNA
                                organism = Capsicum annuum

SEQUENCE: 67
tactccttca ggtggatcta aggggtgaacc aagtggaaat tcaggataca sagtcaagct 60
gccatattct ctcttggtta cccttgtagt tgcagcatca t 101

SEQ ID NO: 68                moltype = DNA length = 210
FEATURE                      Location/Qualifiers
variation                    89
                                note = Polymorphism: LT17=A; PBC167=G
source                      1..210
                                mol_type = unassigned DNA
                                organism = Capsicum annuum

SEQUENCE: 68
ttgaattagc tggtaatgca agcaaggatc ttaaagtga gagaatcacc cccaggcact 60
tgcaactggc cattcgcgga gacgaggarc ttgacacact tatcaaagg accattgctg 120
gtggtggtgt catccctcac attcacaagt ccctaataaa caaacctcc aaggagtga 180
tttctccttt tttatcttag atgtatcatg 210

SEQ ID NO: 69                moltype = DNA length = 120

```

-continued

FEATURE Location/Qualifiers
variation 60
note = Polymorphism: LT17=T; PBC167=C
source 1..120
mol_type = unassigned DNA
organism = Capsicum annuum

SEQUENCE: 69
ttaaagtgtc aacaagagtc cattctacag cacaatgagc aagtttggtg aagggccaaay 60
gcatgagccc tgtttgcttt atatacttac ggaggaaaac tcaaacatgt gataagtaca 120

SEQ ID NO: 70 moltype = DNA length = 101
FEATURE Location/Qualifiers
variation 51
note = Polymorphism: LT17=C; PBC167=T
source 1..101
mol_type = unassigned DNA
organism = Capsicum annuum

SEQUENCE: 70
gttgcttagt ctaagagaga caatagctct ttttaagctt aattcaccac yattatctac 60
ttgcacttga aaagtccaaa gagaaaaaag ggaccaaacc t 101

SEQ ID NO: 71 moltype = DNA length = 121
FEATURE Location/Qualifiers
variation 61
note = Polymorphism: LT17=G; PBC167=A
source 1..121
mol_type = unassigned DNA
organism = Capsicum annuum

SEQUENCE: 71
acaatagttt tatctccagc actaaaatgg ttcagcacct gtgttctgtg aagtaaatgtg 60
racaattttg tgtaggattt ccatgtacct gcaattgaat cacatattac aggcggagtt 120
a 121

SEQ ID NO: 72 moltype = DNA length = 101
FEATURE Location/Qualifiers
variation 51
note = Polymorphism: LT17=A; PBC167=T
source 1..101
mol_type = unassigned DNA
organism = Capsicum annuum

SEQUENCE: 72
ctaataatca agatgagaat attaatatgg tgcaatcagc ggatgatgat wtacaaatgt 60
tgcagcagca gcctactgat ttgacccaa tggaggatac t 101

SEQ ID NO: 73 moltype = DNA length = 121
FEATURE Location/Qualifiers
variation 61
note = Polymorphism: LT17=A; PBC167=C
source 1..121
mol_type = unassigned DNA
organism = Capsicum annuum

SEQUENCE: 73
gtgaagttat ggaaataaat gagtagagga tctgatgtcg gagcaagtag ggtaaagagt 60
mtcgcgtctc atgttgtagt gtattgtcat cataaaattc gatttttctg cttcttagtc 120
a 121

SEQ ID NO: 74 moltype = DNA length = 121
FEATURE Location/Qualifiers
variation 61
note = Polymorphism: LT17=T; PBC167=C
source 1..121
mol_type = unassigned DNA
organism = Capsicum annuum

SEQUENCE: 74
ctcaaatagc cggaatttca ttacattttc cccttggtat aacattatca atactgagct 60
yctattcttg actgactatg cactcaggta ggaattcata catacaaaact ccccaaaaag 120
g 121

-continued

```

SEQ ID NO: 75      moltype = DNA   length = 101
FEATURE            Location/Qualifiers
variation          51
                   note = Polymorphism: LT17=C; PBC167=T
source             1..101
                   mol_type = unassigned DNA
                   organism = Capsicum annuum

SEQUENCE: 75
gttgcttagt ctaagagaga caatagctct ttttaagctt aattcaccac yattatctac 60
ttgcacttga aaagtccaaa gagaaaaaag ggaccaaacc t                      101

```

1-16. (canceled)

17. A *Capsicum* plant resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*, or a cell or plant part thereof, wherein the plant comprises introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein the QTL is located on chromosome 6 within the genomic interval delimited by the marker PE-0014628 (SEQ ID NO: 1) and the marker PE-0021476 (SEQ ID NO: 37).

18. The plant according to claim 17, wherein the QTL is present on chromosome 6 in the genome of a seed of *Capsicum annuum* ST17-1157, deposited at NCIMB accession number NCIMB 43602.

19. The plant according to claim 17, wherein the presence of the QTL comprises the presence of one or more of:

allele A of PE-0014628 (SEQ ID NO: 1);
 allele C of PE-0056542 (SEQ ID NO: 5);
 allele A of PE-0056543 (SEQ ID NO: 9);
 allele T of PE-0056544 (SEQ ID NO: 13);
 allele C of PE-0056545 (SEQ ID NO: 17);
 allele C of PE-0056546 (SEQ ID NO: 21);
 allele C of PE-0056547 (SEQ ID NO: 25);
 allele A of PE-0056549 (SEQ ID NO: 29);
 allele T of PE-0056550 (SEQ ID NO: 33); and/or
 allele C of PE-0021476 (SEQ ID NO: 37).

20. The plant according to claim 17, wherein the presence of the QTL comprises following allele combinations:

allele combination "A" comprising allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), and allele C of PE-0056546 (SEQ ID NO: 21);

allele combination "B" comprising allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21) and allele C of PE-0056547 (SEQ ID NO: 25);

allele combination "C" comprising allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), and allele T of PE-0056550 (SEQ ID NO: 33);

allele combination "D" comprising allele A of PE-0014628 (SEQ ID NO: 1), allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of

PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), allele T of PE-0056550 (SEQ ID NO: 33), and allele C of PE-0021476 (SEQ ID NO: 37); and allele combination "E" comprising allele A of PE-0056549 (SEQ ID NO: 29) and allele T of PE-0056550 (SEQ ID NO: 33).

21. The plant according to claim 17, wherein the QTL is homozygously present in the genome of the plant.

22. The plant according to claim 17, wherein the QTL is heterozygously present in the genome of the plant.

23. The plant according to claim 17, wherein the QTL co-segregates with the gene CA06g26150 and/or the gene CA06g26250.

24. The plant according to claim 23, wherein the resistance is caused by at least one loss-of-function mutation in the sequence of the gene CA06g26150 or in the sequence of the gene CA06g26250.

25. The plant according to claim 17, wherein the plant is a progeny of, or is derived from a *Capsicum annuum* plant of ST17-1157, representative seeds of which have been deposited under NCIMB accession number NCIMB 43602.

26. The plant according to claim 17, which is a *Capsicum annuum*, *Capsicum baccatum*, *Capsicum frutescens*, *Capsicum chinense*, *Capsicum pubescens* or *Capsicum chacoense* plant

27. The plant according to claim 26, which is a *Capsicum annuum* plant.

28. The plant according to claim 17, which is a bell pepper.

29. A *Capsicum* seed, which can be grown into a *Capsicum* plant according to claim 17.

30. A method for producing a *Capsicum* plant according to claim 17, or a plant part, cell or seed thereof, comprising:

(i) using a *Capsicum* plant resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*, comprising introgressed in its genome a quantitative trait locus (QTL) conferring resistance to powdery mildew, wherein the QTL is located on chromosome 6 within the genomic interval delimited by the marker PE-0014628 (SEQ ID NO: 1) and the marker PE-0021476 (SEQ ID NO: 37), as a breeding partner in a breeding program, thereby transferring the QTL to a progeny *Capsicum* plant, wherein the progeny *Capsicum* plant is resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*, or

(ii) introducing by mutagenesis or genome editing one or more mutations in the genome of *Capsicum* plants, at a locus within the genomic interval delimited by the marker PE-0014628 (SEQ ID NO: 1) and the marker PE-0021476 (SEQ ID NO: 37) and selecting a mutant

plant, wherein the plant is resistant to powdery mildew caused by the fungal pathogen *Leveillula taurica*.

- 31.** The method of claim **30**, wherein step (i) comprises:
- (i) crossing the *Capsicum* plant resistant to powdery mildew and a susceptible or less resistant *Capsicum* plant, thus generating a F1 population,
 - (ii) selfing the F1 population to create a F2 population, and
 - (iii) selecting resistant individuals in the progeny thus obtained.

32. The method of claim **31**, further comprising self-pollinating and/or backcrossing one or several times the resistant plant obtained at step c) and selecting a resistant plant in the progeny thus obtained,

33. A method for detecting and/or selecting *Capsicum* plants comprising in their genome a QTL on chromosome 6, the QTL conferring resistance to powdery mildew, comprising detecting a molecular marker linked to the QTL within the chromosomal region delimited by marker PE-0014628 (SEQ ID NO: 1) and marker PE-0021476 (SEQ ID NO: 37).

34. The method of claim **33**, comprising the detection of at least one of the following alleles: A of PE-0014628 (SEQ ID NO: 1), allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), allele T of PE-0056550 (SEQ ID NO: 33), and allele C of PE-0021476 (SEQ ID NO: 37).

35. The method of claim **34**, comprising the detection of at least one of the following alleles: allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), and allele C of PE-0056546 (SEQ ID NO: 21).

36. The method of claim **34**, comprising detecting one of the following allele combinations:

allele combination “A” comprising allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), and allele C of PE-0056546 (SEQ ID NO: 21);

allele combination “B” comprising allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21) and allele C of PE-0056547 (SEQ ID NO: 25);

allele combination “C” comprising allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), and allele T of PE-0056550 (SEQ ID NO: 33);

allele combination “D” comprising allele A of PE-0014628 (SEQ ID NO: 1), allele C of PE-0056542 (SEQ ID NO: 5), allele A of PE-0056543 (SEQ ID NO: 9), allele T of PE-0056544 (SEQ ID NO: 13), allele C of PE-0056545 (SEQ ID NO: 17), allele C of PE-0056546 (SEQ ID NO: 21), allele C of PE-0056547 (SEQ ID NO: 25), allele A of PE-0056549 (SEQ ID NO: 29), allele T of PE-0056550 (SEQ ID NO: 33), and allele C of PE-0021476 (SEQ ID NO: 37); and

allele combination “E” comprising allele A of PE-0056549 (SEQ ID NO: 29) and allele T of PE-0056550 (SEQ ID NO: 33).

* * * * *