



US 20250281642A1

(19) **United States**(12) **Patent Application Publication**
KAGEYAMA et al.(10) **Pub. No.: US 2025/0281642 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **POLYNUCLEOTIDE FOR TREATMENT OF
NEURODEGENERATIVE DISEASE, VECTOR,
CELL, PHARMACEUTICAL COMPOSITION,
AND SCREENING METHOD**(71) Applicant: **RIKEN**, Saitama (JP)(72) Inventors: **Ryoichiro KAGEYAMA**, Saitama (JP);
Takashi KAISE, Saitama (JP);
Masahiro FUKUI, Saitama (JP)(73) Assignee: **RIKEN**, Saitama (JP)(21) Appl. No.: **18/279,282**(22) PCT Filed: **Feb. 25, 2022**(86) PCT No.: **PCT/JP2022/008055**

§ 371 (c)(1),

(2) Date: **Mar. 12, 2024**(30) **Foreign Application Priority Data**

Mar. 3, 2021 (JP) 2021-033464

Publication Classification(51) **Int. Cl.****A61K 48/00** (2006.01)**A61P 25/28** (2006.01)**C12N 15/113** (2010.01)**C12N 15/86** (2006.01)(52) **U.S. Cl.**CPC **A61K 48/0058** (2013.01); **A61P 25/28**(2018.01); **C12N 15/1137** (2013.01); **C12N****15/86** (2013.01); **C12N 2740/15043** (2013.01)

(57)

ABSTRACT

The purpose of the present invention is to provide a novel method for increasing the number of newly generated neurons in the adult brain, and to provide a polynucleotide, a vector, and a pharmaceutical composition for use in the method. The present invention provides: a polynucleotide characterized by including (A) the nucleic acid sequence of the *Plagl2* gene, (B) an miR-shRNA (microRNA adapted short hairpin RNA) nucleic acid sequence for the *Dyrk1a* gene, and (C) a promoter sequence operatively connected to the nucleic acid sequences; a vector including the polynucleotide; and a pharmaceutical composition including the polynucleotide and the vector.


Specification includes a Sequence Listing.**6-month-old (1wpi)**

Fig. 1




Fig. 2




Fig. 3




Fig. 4-2




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9-1




Fig. 9-2




Fig. 10-1




Fig. 10-2




Fig. 10-3





Fig. 11





Fig. 12




Fig. 13




Fig. 14

Barnes maze test




Fig. 15

Fear conditioning test




Fig. 16





Fig. 17





Fig. 18





Fig. 19

Barnes maze test




Fig. 20

Fear conditioning test





Fig. 21



**POLYNUCLEOTIDE FOR TREATMENT OF
NEURODEGENERATIVE DISEASE, VECTOR,
CELL, PHARMACEUTICAL COMPOSITION,
AND SCREENING METHOD**

TECHNICAL FIELD

[0001] The present invention relates to a polynucleotide, a vector, cells, a pharmaceutical composition, and a screening method for treating a neurodegenerative disease.

BACKGROUND ART

[0002] It has become clear that neural stem cells and neural progenitor cells are also present in the adult brain of mammals including a human, although the absolute number thereof is small, and that these cells provide neurogenesis in some brain regions. However, most of the neural stem cells in the adult brain are quiescent and have a low proliferative ability and a neuron-producing ability. These proliferative ability and neuron producing ability further decline with age, and thus there are almost no newly born neurons (newborn neurons) in the aging brain. A remarkable decrease in newborn neurons is also thought to be a cause of brain hypofunction such as dementia, and a neurodegenerative disease such as Alzheimer's disease and Parkinson's disease.

[0003] Under such circumstances, an attempt has been made to promote functional recovery by amplifying and differentiating neural stem cells and neural progenitor cells, which are present in small amounts in the adult brain, by stimulation with a growth factor and the like (see Non Patent Documents 1). However, the effect thereof cannot be deemed to be sufficient. In addition, research on a cell transplantation therapy model using neural stem cells and neural progenitor cells derived from a fetal brain, ES cells, and iPS cells is also being actively carried out, but it is not clear even whether or not cell transplantation therapy in the brain can reconstruct a normal neural network, and the risk of a side effect thereof is also high. Further, many attempts have been made so far to increase the number of newborn neurons in the adult brain, but the effects thereof are still limited (see Non Patent Literatures 1 to 5).

PRIOR ART DOCUMENTS

Non Patent Documents

- [0004]** Non Patent Document 1: Nagahara A H et al. (2009) Neuroprotective effects of brain-derived neurotrophic factor in rodent and primate models of Alzheimer's disease, *Nat. Med.*, vol. 15 (3), pp. 331-337.
- [0005]** Non Patent Document 2: Henriette van Praag et al., Exercise enhances learning and hippocampal neurogenesis in aged mice, *J Neurosci.*, 2005 Sep. 21; 25 (38): 8680-5
- [0006]** Non Patent Document 3: Benedetta Artegiani et al., Overexpression of cdk4 and cyclinD1 triggers greater expansion of neural stem cells in the adult mouse brain. *J Exp Med.* 2011 May 9; 208 (5): 937-48
- [0007]** Non Patent Document 4: Benraiss, A. et al. (2013) Mobilization of endogenous progenitor cells regenerates functionally-integrated medium spiny striopallidal projection neurons and delays disease progression in a transgenic model of Huntington's disease. *Cell Stem Cell* 12, 787-799

- [0008]** Non Patent Document 5: Berdugo-Vega, G. et al. (2020). Increasing neurogenesis refines hippocampal activity rejuvenating navigational learning strategies and contextual memory throughout life. *Nat. Commun.* 11, 135

SUMMARY OF INVENTION

Technical Problem

[0009] An object of the present invention is to provide a novel method for increasing the number of newborn neurons in an adult brain, and also provide a polynucleotide, a vector, transformed cells, and a pharmaceutical composition used in this method, and a method for screening for a substance that increases the number of newborn neurons in an adult brain.

Solution to Problem

[0010] Under these circumstances, the present inventors have carried out intensive studies, and as a result found that by introducing a specific gene set into the aged mouse brain with a lentivirus, endogenous neural stem cells are efficiently activated and proliferated to produce a large number of neurons. Aged neural stem cells reverted to their adolescent or younger state to produce a large number of neurons, resulting in improved memory and learning abilities. In addition, even when the specific gene set was introduced into the brain of an Alzheimer's disease model mouse, endogenous neural stem cells were efficiently activated to produce a large number of neurons, resulting in improved memory and learning abilities. That is, a summary of the present invention is as follows.

[0011] [1] A polynucleotide comprising:

[0012] (A) a nucleic acid sequence of a *Plagl2* gene;

[0013] (B) a nucleic acid sequence of an miR-shRNA (microRNA adapted short hairpin RNA) against a *Dyrk1a* gene; and

[0014] (C) a promoter sequence operably linked to the above nucleic acid sequences.

[0015] [2] The polynucleotide according to [1], wherein the *Plagl2* gene and the *Dyrk1a* gene are mammalian genes.

[0016] [3] The polynucleotide according to [1] or [2], wherein the promoter is selected from the group consisting of a *Hes5* promoter, a GFAP promoter, a *Sox2* promoter, and an *Ling* promoter.

[0017] [4] A vector comprising the polynucleotide according to any one of [1] to [3].

[0018] [5] The vector according to [4], wherein the vector is a lentivirus vector, an adeno-associated virus vector, an adenovirus vector, or a plasmid vector.

[0019] [6] A pharmaceutical composition comprising the vector according to [4] or [5].

[0020] [7] A pharmaceutical composition comprising:

[0021] (a) a vector comprising a nucleic acid sequence of a *Plagl2* gene and a promoter sequence operably linked to the nucleic acid sequence of the *Plagl2* gene; and

[0022] (b) a vector comprising a nucleic acid sequence of an miR-shRNA (microRNA adapted short hairpin RNA) against a *Dyrk1a* gene and a promoter sequence operably linked to the nucleic acid sequence of the miR-shRNA, or an siRNA or an miRNA against the *Dyrk1a* gene.

[0023] [8] Cells transformed with the vector according to [4] or [5].

[0024] [9] A pharmaceutical composition comprising the cells according to [8].

[0025] [10] The pharmaceutical composition according to [6], [7], or [9], wherein the pharmaceutical composition is used for treatment of a neurodegenerative disease or an inner ear disease.

[0026] [11] The pharmaceutical composition according to [10], wherein the pharmaceutical composition is administered to a subject by intracerebroventricular injection, intrathecal bolus injection or infusion, intraganglionic injection, intraneural injection, subcutaneous injection, or intratympanic injection.

[0027] [12] A method for screening for a substance for treatment of a neurodegenerative disease or an inner ear disease, comprising:

[0028] (1) a step of causing a test substance to act on neural stem cells or inner ear supporting cells to measure expression of a *Plagl2* gene and a *Dyrk1a* gene; and

[0029] (2-1) a step of selecting a test substance having an ability to enhance *Plagl2* gene expression and an ability to suppress *Dyrk1a* gene expression from a result of the step (1) or

[0030] (2-2) a step of selecting a test substance having an ability to enhance *Plagl2* gene expression, and a test substance having an ability to suppress *Dyrk1a* gene expression from the result of the step (1).

Advantageous Effects of Invention

[0031] According to the present invention, by introducing a specific gene set into an adult brain or an aged brain, endogenous neural stem cells can be efficiently activated and proliferated to produce a large number of neurons. The specific gene set can cause neural stem cells in the aged brain to revert to their adolescent or younger state and can produce a large number of neurons, resulting in improved memory and learning abilities. In addition, a therapeutic effect thereof on a neurodegenerative disease such as Alzheimer's disease can also be expected, and endogenous neural stem cells can be efficiently activated to produce a large number of neurons, resulting in improved memory and learning abilities. Further, the specific gene set of the present invention can differentiate not only brain neural stem cells but also supporting cells in the inner ear into hair cells, and it is suggested that the specific gene set is also effective in the treatment of an inner ear disease.

BRIEF DESCRIPTION OF DRAWINGS

[0032] FIG. 1 is a diagram showing a step of screening for a gene that contributes to NSC (neural stem cell) activation.

[0033] FIG. 2 is a diagram showing results of in vitro screening for an NSC-activating gene by overexpression of genes highly expressed in an embryo.

[0034] FIG. 3 is a diagram showing results of in vivo screening for an NSC-activating gene by overexpression of genes highly expressed in an embryo.

[0035] FIG. 4-1 is a diagram showing results of in vitro screening for an NSC-activating gene by knocking down genes highly expressed in an adult.

[0036] FIG. 4-2 is a diagram showing results of in vitro screening for an NSC-activating gene by knocking down genes highly expressed in an adult.

[0037] FIG. 5 is a diagram showing results of in vivo screening for an NSC-activating gene.

[0038] FIG. 6 is a diagram showing results of in vivo screening for an NSC-activating gene.

[0039] FIG. 7 is a diagram showing long-term activation of neurogenesis in aged mouse brains by iPaD. The combination of *Plagl2* overexpression and *Dyrk1a* knockdown was termed "iPaD (inducing *Plagl2* and anti-*Dyrk1a*)."

[0040] FIG. 8 is a diagram showing long-term activation of neurogenesis in aged mouse brains by iPaD.

[0041] FIG. 9-1 is a diagram showing the decline in neurogenic ability with aging (wild-type hippocampus).

[0042] FIG. 9-2 is a diagram showing the decline in neurogenic ability with aging and the rejuvenation of aged NSCs by iPaD.

[0043] FIG. 10-1 is a diagram showing the schedule of a Barnes maze test.

[0044] FIG. 10-2 is a diagram showing improving effects of iPaD on cognitive functions of aged mice (Barnes maze test).

[0045] FIG. 10-3 is a diagram showing an improving effect of iPaD on a cognitive function of aging mice (Barnes maze test).

[0046] FIG. 11 is a diagram showing changes with age in months in the number of MCM2-positive cells in the hippocampal dentate gyrus of Alzheimer's disease model mice (5xFAD mice) and normal mice (wild-type littermate mice).

[0047] FIG. 12 is a diagram showing changes with age in months in the number of DCX-positive cells in the hippocampal dentate gyrus of Alzheimer's disease model mice (5xFAD mice) and normal mice (wild-type littermate mice).

[0048] FIG. 13 is a diagram showing an age-related increase in the number of amyloid β plaques in the hippocampal dentate gyrus of Alzheimer's disease model mice (5xFAD mice).

[0049] FIG. 14 is a diagram showing results of comparing a cognitive function between Alzheimer's disease model mice (5xFAD mice) and normal mice (wild-type littermate mice) (Barnes circular maze test).

[0050] FIG. 15 is a diagram showing results of comparing a cognitive function between Alzheimer's disease model mice (5xFAD mice) and normal mice (wild-type littermate mice) (contextual fear conditioning test).

[0051] FIG. 16 is a diagram showing an improving effect of iPaD on Alzheimer's disease (number of MCM2-positive cells).

[0052] FIG. 17 is a diagram showing an improving effect of iPaD on Alzheimer's disease (number of DCX-positive cells).

[0053] FIG. 18 is a diagram showing an improving effect of iPaD on Alzheimer's disease (number of amyloid β plaques).

[0054] FIG. 19 is a diagram showing an improving effect of iPaD on a cognitive function of Alzheimer's disease model mice (5xFAD mice) (Barnes circular maze test).

[0055] FIG. 20 is a diagram showing an improving effect of iPaD on a cognitive function of Alzheimer's disease model mice (5xFAD mice) (contextual fear conditioning test).

[0056] FIG. 21 is a diagram showing activation of supporting cells and regeneration of hair cells in the inner ear by iPAD.

DESCRIPTION OF EMBODIMENTS

[0057] Hereinafter, the present invention will be described in detail. Herein, a molecular biological technique can be carried out by a method disclosed in a general laboratory book known to those skilled in the art or a method based thereon, unless otherwise specified. In addition, the terms used herein are interpreted as having a meaning commonly used in the technical field, unless otherwise specified.

[0058] According to the present disclosure, by forcibly expressing a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and suppressing expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells, in target cells such as neural stem cells, these cells can be efficiently activated and proliferated. That is, in the present application, provided are a polynucleotide, a vector, and transformed cells that include a specific gene set that can efficiently activate and proliferate endogenous neural stem cells or the like in an adult brain, an aged brain, a diseased brain, or the like to produce a large number of neurons, a pharmaceutical composition including the polynucleotide, the vector, or the transformed cells, a method for treating a neurodegenerative disease or the like using the polynucleotide, the vector, the transformed cells, or the pharmaceutical composition, and the like.

<Polynucleotide>

[0059] A first embodiment of the polynucleotide of the present disclosure is a polynucleotide including: (A) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells; (B) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene); and (C) a promoter sequence operably linked to the nucleic acid sequences, to be introduced into target cells such as neural stem cells.

[0060] A second embodiment of the polynucleotide of the present disclosure is a combination of a polynucleotide including (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and a promoter sequence operably linked to the nucleic acid sequence of the gene, and a polynucleotide including (b1) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene), and a promoter sequence operably linked to the nucleic acid sequence that suppresses expression of the gene (for example, nucleic acid sequence of miR-shRNA), to be introduced into target cells such as neural stem cells.

[0061] A third embodiment of the polynucleotide of the present disclosure is a combination of a polynucleotide including (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and a promoter sequence operably linked to the nucleic acid sequence of the gene, and the gene and (b2) a

nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, polynucleotide of siRNA or miRNA against the gene), to be introduced into target cells such as neural stem cells.

[0062] As used herein, a “polynucleotide” is used interchangeably with a “nucleic acid”, a “gene,” or a “nucleic acid molecule” and means a polymer of a nucleotide. As used herein, the term “nucleotide sequence” is used interchangeably with a “nucleic acid sequence” or a “base sequence” and is expressed as a sequence of deoxyribonucleotides (abbreviated as A, G, C, and T). For example, “a polynucleotide including the nucleotide sequence of SEQ ID NO: 1 or a fragment thereof” means a polynucleotide including the sequence represented by each deoxynucleotide A, G, C, and/or T of SEQ ID NO: 1 or a fragment portion thereof.

[0063] Examples of target cells into which the polynucleotide of the present disclosure is introduced include tissue stem cells (somatic stem cells) such as neural stem cells, inner ear supporting cells, hematopoietic stem cells, mesenchymal stem cells, skeletal muscle stem cells, or dental pulp stem cells, retinal Muller glial cells, and tissue progenitor cells, and among these, neural stem cells and inner ear supporting cells are preferable from the viewpoint of activation and proliferation being promoted by introducing the polynucleotide of the present disclosure.

Regarding First Embodiment

[0064] The first embodiment of the polynucleotide of the present disclosure to be introduced into target cells such as neural stem cells is a polynucleotide including: (A) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells; (B) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene); and (C) a promoter sequence operably linked to the nucleic acid sequences.

[(a) Nucleic Acid Sequence of Gene that is Highly Expressed in Embryo and Low Expressed in Adult Neural Stem Cells]

[0065] The nucleic acid sequence (A) is not particularly limited as long as it is a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and above all, the nucleic acid sequence is preferably a nucleic acid sequence of a gene that promotes proliferation and activation of adult neural stem cells by overexpression thereof in the adult neural stem cells; and specific examples thereof include nucleic acid sequences of Gsx2, Dmrt3, Cdk4, Plagl2, Sox21, Asc11, Tgif2, Plagl1, Hmga2, and the like. Among these, the nucleic acid sequence of the Plagl2 gene is more preferable from the viewpoint of a superior ability to proliferate and activate adult neural stem cells. The Plagl2 gene (Pleiomorphic adenoma gene like-2) is a gene encoding a transcription factor PLAGL2 having a C2H2 zinc finger region. The Plagl2 gene is also referred to as ZNF900. PLAGL2 is known to function as a positive regulator of transcription and be localized in the nucleus. As described in detail in the Examples, in the present disclosure, PLAGL2 was found as one of the nuclear factors differentially expressed by comparing the transcriptomes (DDBJ BioProject Accession:

PRJDB9010) of G1/G0 NSCs derived from the ganglionic eminence and the dorsal cortex of E14 mouse embryos with those of NSCs mostly quiescent derived from LV-SVZ (SAM00192826) and DG-SGZ (SAM00192827) in 2- to 3-month-old mice.

[0066] In the present disclosure, a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, such as the *Plagl2* gene, are derived from mammals such as a human, a monkey, a pig, a horse, a cow, a rabbit, a sheep, a goat, a cat, a dog, or a guinea pig, and among these, the gene is preferably derived from a human, a monkey, or a mouse, and more preferably derived from a human.

[0067] In the present disclosure, the nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, such as the *Plagl2* gene, includes a coding region (CDS) within an mRNA sequence of the gene. As the nucleic acid sequence of the gene, a sequence known as a coding region (CDS) of an mRNA sequence of each of the genes derived from the above mammals can be adopted, and for example, a nucleotide sequence registered in the GenBank nucleotide sequence database or the like can be used. The nucleic acid sequence of the *Plagl2* gene in the present disclosure is a nucleic acid sequence which is the same (SEQ ID NO: 1) as the nucleotide sequence registered in the GenBank nucleotide sequence database or the like, or has a homology of 80% or more, preferably 90% or more, more preferably 95% or more, or further preferably 98% or more, and particularly preferably 99% or more therewith, and in which the peptide encoded has the transcription factor activity as *PLAGL2*. Examples of a most preferable nucleic acid sequence of the *Plagl2* gene in the present disclosure include the sequence of SEQ ID NO: 1.

[(B) Nucleic Acid Sequence that Suppresses Expression of Gene that is Low Expressed in an Embryo and Highly Expressed in Adult Neural Stem Cells]

[0068] The nucleic acid sequence (B) is not particularly limited as long as it is a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells, and such a gene that is low expressed in an embryo and highly expressed in adult neural stem cells is preferably a gene that, when knocked down, can efficiently activate quiescent neural stem cells; and examples thereof include *Cdkn1a*, *Prkcz*, *Dyrk1a*, *Zbtb7a*, *Dusp22*, *Cidea*, *Rasd1*, *Nr4a1*, *Nfe212*, *Stat6*, and *Tsc22d3*, and among these, *Cdkn1a*, *Prkcz*, *Dyrk1a*, and *Zbtb7a* are more preferable, and *Dyrk1a* is further preferable. The *Dyrk1a* gene encodes an enzyme that phosphorylates serine/threonine and tyrosine, and is a gene that is thought to be possibly involved in the development of Down's syndrome caused by trisomy of chromosome 21. As described in detail in the Examples, in the present disclosure, the *Dyrk1a* gene was found as one of the above 11 genes that, when knocked down, effectively activated quiescent NSCs in target neural stem cells.

[0069] In the present disclosure, a gene that is low expressed in an embryo and highly expressed in adult neural stem cells, such as the *Dyrk1a* gene, is derived from mammals such as a human, a monkey, a pig, a horse, a cow, a rabbit, a sheep, a goat, a cat, a dog, or a guinea pig, and among these, the gene is preferably derived from a human, a monkey, or a mouse, and more preferably derived from a human.

[0070] In the present disclosure, the nucleic acid sequence of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells, such as the *Dyrk1a* gene, includes a coding region (CDS) within an mRNA sequence of the gene. As the nucleic acid sequence of the gene, a sequence known as a coding region (CDS) of an mRNA sequence of each of the genes derived from the above mammals can be adopted, and for example, a nucleotide sequence registered in the GenBank nucleotide sequence database or the like can be used. The nucleic acid sequence of the *Dyrk1a* gene in the present disclosure is a nucleic acid sequence which is the same (SEQ ID NO: 2) as the nucleotide sequence registered in the GenBank nucleotide sequence database or the like, or has a homology of 80% or more, preferably 90% or more, more preferably 95% or more, or further preferably 98% or more, and particularly preferably 99% or more therewith, and in which the peptide encoded has the enzymatic activity as *Dyrk1a*. Examples of a most preferable nucleic acid sequence of the *Dyrk1a* gene in the present disclosure include the sequence of SEQ ID NO: 2.

[0071] In the present disclosure, the nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells is not particularly limited as long as it has a nucleic acid sequence that can suppress expression of the gene by a conventionally known method, and examples thereof include an miR-shRNA (microRNA adapted short hairpin RNA) against the gene. The miR-shRNA against the gene is an shRNA (hairpin-shaped RNA sequence used for gene silencing by RNA interference) that can suppress expression of the gene. The shRNA is not particularly limited as long as it has a nucleic acid sequence that can suppress expression of the gene, and can be appropriately designed according to the sequence of the gene. Preferable examples of a nucleic acid sequence of such an shRNA include the nucleic acid sequence of SEQ ID NO: 3, which is the nucleic acid sequence of an miR-shRNA against *Dyrk1a*, and a nucleic acid sequence that has a homology of 80% or more, preferably 90% or more, more preferably 95% or more, further preferably 98% or more, and particularly preferably 99% or more with the nucleic acid sequence of SEQ ID NO: 3 and that can suppress expression of the *Dyrk1a* gene can also be adopted as a nucleic acid sequence of an miR-shRNA (microRNA adapted short hairpin RNA) against the *Dyrk1a* gene in the present disclosure.

[(C) Promoter Sequence Operably Linked to the Nucleic Acid Sequences]

[0072] In the present disclosure, a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene) are operably linked to a promoter sequence (C). As the promoter sequence (C) in the present embodiment, a promoter sequence that functions in nervous system cells such as neural stem cells or inner ear supporting cells can be adopted. Examples of such a promoter sequence include the *Hes5* promoter (SEQ ID NO: 4) and the *GFAP* promoter,

which function in neural stem cells, and the Sox2 promoter and the Ling promoter, which function in inner ear supporting cells.

[0073] Here, “operably linked” refers to a juxtaposition in which the components described are in a relationship that enables them to function in their intended manner. In one embodiment, this term refers to a functional linkage between a nucleic acid expression control sequence (for example, promoter and/or enhancer) and a target polynucleotide sequence, and the functional linkage may be a direct linkage or an indirect linkage (in which another polynucleotide sequence intervenes therebetween). In the present disclosure, the promoter sequence directs transcription of the target polynucleotide linked.

[0074] The polynucleotide including (A) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, (B) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene), and (C) a promoter sequence operably linked to the nucleic acid sequences according to the present embodiment is preferably a polynucleotide including (A) a nucleic acid sequence of the Plagl2 gene, (B) a nucleic acid sequence that suppresses expression of the Dyrk1a gene (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the Dyrk1a gene), and (C) a promoter sequence operably linked to the nucleic acid sequences, and more preferable examples thereof include the polynucleotide of SEQ ID NO: 5, and a polynucleotide that is derived from the polynucleotide of SEQ ID NO: 5 by deletion, substitution, insertion, and/or addition of one or more (for example, 1 to 50, 1 to 40, 1 to 30, 1 to 25, 1 to 20, 1 to 15, 1 to 10, 1 to 9, 1 to 8, 1 to 7, 1 to 6, 1 to 5, 1 to 4, 1 to 3, 1 or 2, or 1) nucleotides and that, when introduced into a vector to transform target cells, can forcibly express the Plagl2 gene and knock down the Dyrk1a gene. A combinations of two or more of these deletion, substitution, insertion, and addition may be included at the same time. In general, the smaller the number of deletions, substitutions, insertions, and/or additions of the nucleotides, the more preferable. In addition, in the present invention, preferable polynucleotides include, for example, a polynucleotide that is hybridizable under a stringent hybridization condition to a sequence complementary to SEQ ID NO: 5.

[0075] Here, the “stringent condition” may be any of a low stringent condition, a moderately stringent condition, and a highly stringent condition. The “low stringent condition” is, for example, conditions of 5×SSC, 5×Denhardt’s solution, 0.5% SDS, 50% formamide, and 32° C. In addition, the “moderately stringent condition” is, for example, conditions of 5×SSC, 5×Denhardt’s solution, 0.5% SDS, 50% formamide, and 42° C. The “highly stringent condition” is, for example, conditions of 5×SSC, 5×Denhardt’s solution, 0.5% SDS, 50% formamide, and 50° C. Under these conditions, it can be expected that the higher the temperature, the more efficiently a DNA having a higher homology can be obtained. However, a plurality of factors such as temperature, probe concentration, probe length, ionic strength, time, and salt concentration can be considered as factors that affect the stringency of hybridization, and those skilled in the art can appropriately select these factors to achieve similar stringency.

[0076] The polynucleotide of the present embodiment may further include a known sequence such as an enhancer sequence, a Kozak sequence, and an appropriate polyadenylation signal sequence that assist in transcription of mRNA, translation into a protein, or the like.

[0077] The method for introducing the polynucleotide of the present embodiment into target cells is not particularly limited, and examples thereof include using a vector such as a virus, a plasmid, or an artificial chromosome, or a technique such as lipofection, a liposome, or microinjection. When the polynucleotide of the present embodiment is introduced into target neural stem cells or the like to transform the same, it is possible to forcibly express a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, such as the Plagl2 gene, and knock down a gene that is low expressed in an embryo and highly expressed in adult neural stem cells, such as the Dyrk1a gene. The target cells (neural stem cells or the like) thus transformed can be activated and proliferated to differentiate to fully exhibit cellular functions, such as producing a large number of neurons. The polynucleotide of the present embodiment can be suitably used for endogenous neural stem cells in an adult brain, an aged brain, a diseased brain, or the like, supporting cells present in the inner ear, or the like.

Regarding Second Embodiment

[0078] The second embodiment of the polynucleotide of the present disclosure to be introduced into target cells such as neural stem cells is a combination of a polynucleotide including (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and a promoter sequence operably linked to the nucleic acid sequence of the gene, and a polynucleotide including (b1) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene), and a promoter sequence operably linked to the nucleic acid sequence that suppresses expression of the gene. The combination of polynucleotides according to the present embodiment is used to, for example, introduce a gene that is highly expressed in an embryo and low expressed in adult neural stem cells and a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, miR-shRNA against the Dyrk1a gene) into separate vectors to separately transform target cells such as neural stem cells.

[0079] To (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells in the present embodiment, the description of (A) a nucleic acid sequence of a gene that is highly expressed in an embryo and highly expressed in adult neural stem cells in the first embodiment described above can be directly applied. In addition, to a promoter sequence operably linked to the nucleic acid sequence of the gene, the description of (C) a promoter sequence operably linked to the nucleic acid sequence in the first embodiment can also be applied, and specific examples thereof include the Hes5 promoter (SEQ ID NO: 4) and the GFAP promoter, which function in neural stem cells, and the Sox2 promoter and the Ling promoter, which function in inner ear supporting cells.

[0080] The polynucleotide including (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells and a promoter sequence operably linked to the nucleic acid sequence of the gene in the present embodiment is preferably a polynucleotide including (a) a nucleic acid sequence of the *Plagl2* gene and a promoter sequence operably linked to the nucleic acid sequence of the *Plagl2* gene, and more preferable examples thereof include a polynucleotide including the nucleic acid sequence of SEQ ID NO: 6. This polynucleotide may further include a known sequence such as an enhancer sequence, a Kozak sequence, and an appropriate polyadenylation signal sequence that assist in transcription of mRNA, translation into a protein, or the like.

[0081] To (b1) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene) in the present embodiment, the description of (B) a nucleic acid sequence of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells in the first embodiment described above can be directly applied. In addition, to a promoter sequence operably linked to the nucleic acid sequence, the description of (C) a promoter sequence operably linked to the nucleic acid sequences in the first embodiment can be applied, and specific examples thereof include the *Hes5* promoter (SEQ ID NO: 4) and the GFAP promoter, which function in neural stem cells, and the *Sox2* promoter and the *Ling* promoter, which function in inner ear supporting cells.

[0082] The polynucleotide including (b1) a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the *Dyrk1a* gene) and a promoter sequence operably linked to the nucleic acid sequence in the present embodiment is preferably a polynucleotide including (b1) a nucleic acid sequence of an miR-shRNA (microRNA adapted short hairpin RNA) against the *Dyrk1a* gene and a promoter sequence operably linked to the nucleic acid sequence of the miR-shRNA, and more preferable examples thereof include a polynucleotide including the nucleic acid sequence of SEQ ID NO: 7. This polynucleotide may further include a known sequence such as an enhancer sequence, a Kozak sequence, and an appropriate polyadenylation signal sequence that assist in transcription of mRNA, translation into a protein, or the like.

[0083] The method for introducing the polynucleotide of the present embodiment into target cells is not particularly limited, and examples thereof include using a vector such as a virus, a plasmid, or an artificial chromosome, or a technique such as lipofection, a liposome, or microinjection. The two types of polynucleotides of the present embodiment can be each independently introduced into target cells. For example, when the two types are incorporated into separate vectors and introduced into target neural stem cells or the like to transform the same, it is possible to forcibly express a gene that is highly expressed in an embryo and low expressed in adult neural stem cells and knock down a gene that is low expressed in an embryo and highly expressed in adult neural stem cells. The target cells (neural stem cells or the like) thus transformed can be activated and proliferated

to fully exhibit cellular functions, such as producing a large number of neurons. The polynucleotides of the present embodiment can be suitably used for endogenous neural stem cells in an adult brain, an aged brain, a diseased brain, or the like, supporting cells present in the inner ear, or the like.

Regarding Third Embodiment

[0084] The third embodiment of the polynucleotide of the present disclosure to be introduced into target cells such as neural stem cells is a combination of a polynucleotide including (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells, and a promoter sequence operably linked to the nucleic acid sequence of the gene, and (b2) a polynucleotide of an siRNA or an miRNA against a gene that is low expressed in an embryo and highly expressed in adult neural stem cells. In the present embodiment, the polynucleotide (a) is introduced into a vector, and the polynucleotide (siRNA or miRNA) (b2) is used together with a transfection reagent and the like to transform target cells.

[0085] To (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells in the present embodiment, the description of (A) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells in the first embodiment described above can be directly applied. In addition, to a promoter sequence operably linked to the nucleic acid sequence of the gene, the description of (C) a promoter sequence operably linked to the nucleic acid sequences in the first embodiment can be applied, and specific examples thereof include the *Hes5* promoter (SEQ ID NO: 4) and the GFAP promoter, which function in neural stem cells, and the *Sox2* promoter and the *Ling* promoter, which function in inner ear supporting cells.

[0086] The polynucleotide including (a) a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells and a promoter sequence operably linked to the nucleic acid sequence of the gene in the present embodiment is preferably a polynucleotide including (a) a nucleic acid sequence of the *Plagl2* gene and a promoter sequence operably linked to the nucleic acid sequence of the *Plagl2* gene, and more preferable examples thereof include a polynucleotide including the nucleic acid sequence of SEQ ID NO: 6.

[0087] The polynucleotide (b2) in the present embodiment is not particularly limited as long as it is an siRNA or an miRNA of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells and having a sequence that can suppress expression of (knock down) the gene. The siRNA or miRNA of the gene can be designed from the sequence of the gene. For example, an siRNA or an miRNA of the *Dyrk1a* gene can be designed from the sequence of the *Dyrk1a* gene, and specific examples thereof include the polynucleotide of SEQ ID NO: 8.

[0088] The method for introducing the polynucleotide of the present embodiment into target cells is not particularly limited, and examples thereof include using a vector such as a virus, a plasmid, or an artificial chromosome, or a technique such as lipofection, a liposome, or microinjection. For example, target cells (neural stem cells or the like) can be transformed with a vector into which the polynucleotide (a) of the present embodiment has been introduced to forcibly express the *Plagl2* gene or the like, and further, the target

cells can be transfected with an siRNA or an miRNA of the Dyrk1a gene or the like as the polynucleotide (b2) to suppress expression of the Dyrk1a gene or the like. A transfection method conventionally known to those skilled in the art can be used. The target cells (neural stem cells or the like) thus transformed can be activated and proliferated to fully exhibit cellular functions, such as producing a large number of neurons. The polynucleotide of the present embodiment can be suitably used for endogenous neural stem cells in an adult brain, an aged brain, a diseased brain, or the like, supporting cells present in the inner ear, or the like.

<Vector>

[0089] The disclosure also includes a vector including the polynucleotide of the disclosure described above. To the polynucleotide of the present disclosure included in the vector of the present disclosure, the description of the polynucleotide section described above can be directly applied.

[0090] In the present disclosure, the vector used to forcibly express a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like), and/or suppress the expression of (knock down) a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) in target cells is not particularly limited as long as it is a vector that can efficiently deliver the gene to target cells (neural stem cells, inner ear supporting cells, or the like). Examples of such a vector include a virus vector such as a lentivirus vector, an adeno-associated virus vector, an adenovirus vector, a herpes virus vector (for example, herpes simplex virus vector), a pox virus vector, a baculovirus vector, a papilloma virus vector, or a papovavirus vector (for example, SV40); a plasmid vector; a phagemid vector; a cosmid vector; and a bacteriophage such as lambda phage or M13 phage. Among these, a lentivirus vector, an adeno-associated virus vector, an adenovirus vector, and a plasmid vector are preferable, a lentivirus vector and an adeno-associated virus vector are more preferable, and a lentivirus vector is further preferable.

[0091] Examples of a specific expression vector include pLenti4/V5-DEST(TM), pLenti6/V5-DEST(TM), and pLenti6.2/V5-GW/lacZ (Invitrogen) for lentivirus-mediated gene transfer and expression in mammalian cells; and pCIneo Vector (Promega) for expression in mammalian cells.

[0092] The vector can include a control sequence such as a promoter, an enhancer, a ribosome binding sequence, a terminator, or a polyadenylation site so that the target gene can be expressed. In addition, if necessary, the vector can include a selection marker sequence such as a drug resistance gene (for example, kanamycin resistance gene, ampicillin resistance gene, or puromycin resistance gene), a thymidine kinase gene, or a diphtheria toxin gene, a reporter gene sequence such as mCherry (red fluorescent protein), a green fluorescent protein (GFP), β -glucuronidase (GUS), or FLAG, or the like.

<Pharmaceutical Composition>

[0093] The present disclosure also includes a pharmaceutical composition containing the vector of the present disclosure described above. In addition, the present disclosure also includes a pharmaceutical composition containing (a) a

vector including a nucleic acid sequence of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like), and a promoter sequence operably linked to the nucleic acid sequence of the gene, and (b) a vector including a nucleic acid sequence that suppresses expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) (for example, nucleic acid sequence of miR-shRNA (microRNA adapted short hairpin RNA) against the gene), and a promoter sequence operably linked to the nucleic acid sequence that suppresses expression of the gene, or an siRNA or an miRNA against the gene. Further, the present disclosure also includes a pharmaceutical composition containing the transformed cells of the present disclosure. The pharmaceutical compositions of the present disclosure may contain at least one selected from the group consisting of a pharmaceutically acceptable adjuvant, excipient, carrier, and diluent. To the vectors and transformed cells of the present disclosure, the descriptions in their respective sections can be applied. These pharmaceutical compositions of the present disclosure are used for the treatment of a neurodegenerative disease or an inner ear disease.

[0094] The present disclosure also includes a pharmaceutical composition for the treatment of a neurodegenerative disease or an inner ear disease containing a compound having the ability to enhance expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and the ability to suppress expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) or a prodrug thereof or a pharmaceutically acceptable salt thereof as an active ingredient. Further, the present disclosure also includes a pharmaceutical composition for the treatment of a neurodegenerative disease or an inner ear disease containing a compound having the ability to enhance expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) or a prodrug thereof or a pharmaceutically acceptable salt thereof, and a compound having the ability to suppress expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) or a prodrug thereof or a pharmaceutically acceptable salt thereof as active ingredients.

[0095] The expression of the gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and the expression of the gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) are preferably the expressions of the genes in neural stem cells or inner ear supporting cells. In addition, the above compounds, or prodrugs thereof, or pharmaceutically acceptable salts thereof may be any of low molecular weight compounds, middle molecular weight compounds, and high molecular weight compounds.

[0096] The pharmaceutical composition of the present disclosure is administered to a subject by intracerebroventricular injection, intrathecally bolus injection or infusion, intraganglionic injection, intraneural injection, subcutaneous injection, or intratympanic injection.

[0097] The dosage of the pharmaceutical composition of the present disclosure is not particularly limited, and an appropriate dosage can be selected according to various

conditions such as the type of the disease, the age and the symptom of the patient, the administration route, the purpose of treatment, and the presence or absence of a concomitant agent.

<Treatment Method>

[0098] The present disclosure also includes a method for treating a neurodegenerative disease including forcibly expressing a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and suppressing expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) in neural stem cells. In addition, the present disclosure also includes a method for treating an inner ear disease including forcibly expressing a gene that is highly expressed in an embryo and low expressed (Plagl2 gene or the like) and suppressing expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) in inner ear supporting cells. Examples of a method for forcibly expressing a gene that is highly expressed in an embryo and low expressed (Plagl2 gene or the like) and suppressing expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) in target cells such as inner ear supporting cells include a method using the polynucleotide, vector, the transformed cells, and the pharmaceutical composition according to the present disclosure, and to the descriptions thereof, the descriptions in the Polynucleotide, Vector, Pharmaceutical composition sections described above, respectively, can be applied.

<Transformed Cells>

[0099] The disclosure also includes target cells into which the polynucleotide of the disclosure described above has been introduced. For example, cells transformed with the vector of the disclosure described above, cells obtained by transformation with the vector of the disclosure, transfection with the polynucleotide of the disclosure, or the like are also included in the disclosure. The transformed cells of the present disclosure can be prepared in vitro and administered to a subject in need of treatment. At that time, the transformed cells of the present disclosure can be administered as a pharmaceutical composition together with a pharmaceutically acceptable adjuvant, excipient, carrier, or diluent.

[0100] Further, the present disclosure also includes cells transformed by treatment with a compound having the ability to enhance expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and the ability to suppress expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) or a prodrug thereof or a pharmaceutically acceptable salt thereof, and cells transformed by treatment with a compound having the ability to enhance expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) or a prodrug thereof or a pharmaceutically acceptable salt thereof, and a compound having the ability to suppress expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) or a prodrug thereof or a pharmaceutically acceptable salt thereof. Such cells are prefer-

ably cells obtained by the above transformation of the neural stem cells or inner ear supporting cells, and the above compounds, or prodrugs thereof, or pharmaceutically acceptable salts thereof may be any of low molecular weight compounds, middle molecular weight compounds, and high molecular weight compounds.

<Method for Screening for Substance for Treatment of Neurodegenerative Disease or Inner Ear Disease>

[0101] The present disclosure also includes a method for screening for a substance for treatment of a neurodegenerative disease or an inner ear disease, including:

[0102] (1) a step of causing a test substance to act on neural stem cells or inner ear supporting cells to measure expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like); and

[0103] (2-1) a step of selecting a test substance that has an ability to enhance expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and an ability to suppress expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) from a result of the step (1) or

[0104] (2-2) a step of selecting a test substance that has an ability to enhance expression of a gene that is highly expressed in an embryo and low expressed in adult neural stem cells (Plagl2 gene or the like) and a test substance that has an ability to suppress expression of a gene that is low expressed in an embryo and highly expressed in adult neural stem cells (Dyrk1a gene or the like) from the result of the step (1).

[0105] The substance(s) selected by the screening method of the present disclosure can efficiently activate and proliferate neural stem cells to produce a large number of neurons. Because of this, the substance(s) can cause neural stem cells in an aged brain to revert to their adolescent or younger state and can be expected to produce a large number of neurons, resulting in improved memory and learning abilities. In addition, a therapeutic effect thereof on a neurodegenerative disease such as Alzheimer's disease can also be expected, and endogenous neural stem cells can be efficiently activated to produce a large number of neurons, resulting in improved memory and learning abilities. Further, the substance(s) selected by the screening method of the present invention can differentiate not only brain neural stem cells but also supporting cells in the inner ear into hair cells, and it is expected that the screening method is also effective in the treatment of an inner ear disease.

EXAMPLES

[0106] The present disclosure will be specifically described with reference to the following Examples, but the present disclosure should not be construed as limited by these Examples.

1. Screening for Neural Stem Cell (NSC)-Activating Gene

[0107] The present inventors reasoned that it was necessary to alter expression of a plurality of genes in order to activate and proliferate aged NSCs. Therefore, most adult

NSCs are quiescent, and thus the transcriptomes (DDBJ BioProject Accession: PRJDB9010) of G1/G0 NSCs derived from the ganglionic eminence and the dorsal cortex of E14 mouse embryos was compared with those of NSCs mostly quiescent derived from LV-SVZ (SAMD00192826) and DG-SGZ (SAMD00192827) in 2- to 3-month-old mice to identify nuclear factors differentially expressed and known modulators therefor.

[0108] First, genes that were expressed at a high level in an embryo and low expressed in adult NSCs (“embryonic-high”) were identified. Next, these genes were overexpressed in vitro in NSCs maintained quiescent (EdU) by bFGF and BMP (FIG. 1). After two days, the proportions of proliferating cells that had incorporated EdU were measured to identify genes capable of proliferating and activating NSCs. In the present study, each of the top 80 genes expressed at a high level in an embryo was overexpressed in quiescent NSCs in vitro, and proliferation and activation of NSCs were monitored. Results thereof are shown in FIG. 2. Among the genes tested, *Gsx2*, *Dmrt3*, *Cdk4*, *Plagl2*, *Sox21*, *Asc11*, *Tgif2*, *Plagl1*, and *Hmga2* efficiently activated quiescent NSCs.

[0109] Further, in vivo studies were carried out on *Gsx2*, *Dmrt3*, *Cdk4*, *Plagl2*, *Sox21*, *Asc11*, *Tgif2*, *Plagl1*, and *Hmga2*. These nine genes were individually cloned into a lentivirus under the control of the *Hes5* promoter (SEQ ID NO: 9), which operates specifically in NSCs and astrocytes. Each lentivirus was injected into the hippocampal dentate gyrus of 6-month-old mice, and brain sections were investigated after one week (FIG. 3). The combination of *Cdk4* and cyclin D1 (*D1K4*) is known to activate NSCs in the adult brain, and thus these were used as positive controls. Among the genes tested, *Plagl2* (SEQ ID NO: 10, the sequence linking the *Hes5* promoter and *Plagl2* is SEQ ID NO: 11) most efficiently increased the number of MCM2⁺ mitotic cells (activated NSCs, cells likely to be intermediate progenitor cells) in the hippocampal dentate gyrus of the 6-month-old mice as compared with the control mouse.

[0110] Next, in order to identify genes that can proliferate and activate NSCs by suppressing the expression thereof, genes that are low expressed in an embryo and highly expressed in adult NSCs were used as subject genes. siRNAs of targeted subject genes were introduced into quiescent NSCs to carry out knockdown of genes expressed at a high level in adult NSCs (“adult-high”) and genes of modulators therefor. Results of in vitro knockdown screening are shown in FIG. 4-1 and FIG. 4-2. Among the 124 genes tested, 11 genes whose knockdown efficiently activated quiescent NSCs were selected.

[0111] By using a lentivirus, expression of an shRNA of each of the 11 genes selected above was induced in vivo with *Plagl2* under the control of the *Hes5* promoter (SEQ ID NO: 9). Each lentivirus was injected into the hippocampal dentate gyrus of 6-, 18-, or 20-month-old mice, and brain sections were investigated after 2 or 4 weeks. As a result, shRNAs of *Cdkn1a*, *Prkcz*, and *Dyrk1a* increased the number of MCM2⁺ active NSCs/IPCs in the 18-month-old mice (top in FIG. 5). “Control” in FIG. 5 shows the results obtained by connecting only mCherry to the *Hes5* promoter, and “Scramble” shows the results obtained by connecting mCherry, *Plagl2*, and a random sequence to the *Hes5* promoter. Determination of an increase or a decrease in the number of MCM2⁺ active NSCs/IPCs is based on comparison with “Scramble.” In addition, shRNA (SEQ ID NO: 13)

of *Dyrk1a* (SEQ ID NO: 12) more efficiently increased the number of DCX⁺ dendrite⁺ cells than others in 6-month-old and 19-month-old (18 months+4 weeks) mice (center in FIG. 5, bottom in FIG. 5, FIG. 6). The center in FIG. 5 and the bottom in FIG. 5 show the number of DCX⁺ dendrite⁺ cells in 6-month-old and 19-month-old (18 months+4 weeks) mice in which expression of shRNAs of *Prkcz*, *Dyrk1a*, and *Zbtb7a*, where the number of MCM2⁺ active NSCs/IPCs tended to increase, was induced. The above IPCs are intermediate progenitor cells.

[0112] These data show that combined use of *Plagl2* overexpression and *Dyrk1a* knockdown most efficiently activate neurogenesis in the hippocampal dentate gyrus of both 6-month-old and 19-month-old mice, and the combined use of the two gene expressions was termed iPaD (inducing *Plagl2* and anti-*Dyrk1a* activity). iPaD (SEQ ID NO: 14), which is this combination of the two gene expressions, increased the number of MCM2⁺ and/or BrdU-incorporating cells, including Sox2⁺ GFAP⁺ BrdU⁺ activated NSCs and DCX⁺ dendrite⁺ BrdU⁺ immature neurons. These results suggest the possibility that *Plagl2* overexpression and *Dyrk1a* knockdown in endogenous neural stem cells in an adult brain, an aged brain, or the like efficiently activates and proliferates these neural stem cells to produce a large number of neurons, resulting in improved memory and learning ability.

2. Long-Term Effects of iPaD on Neurogenesis

[0113] In order to investigate the long-term effects of an iPaD lentivirus on neurogenesis in an aged brain, the iPaD lentivirus was introduced into the hippocampal dentate gyrus of 18-month-old mice. Only a few activated NSCs (MCM2⁺;DCX⁻), IPCs (MCM2⁺;DCX^{+/+}), and immature neurons (MCM2⁻;DCX⁺;dendrite⁺) were present in this region in the mice at this age in months, and the control lentivirus did not affect the numbers thereof (Control/8wpi in FIG. 7). On the other hand, the iPaD lentivirus efficiently activated NSCs and induced the formation of IPCs and immature neurons 1 month after infection (19 months of age) (iPaD/4wpi in FIG. 7). A similar effect was observed even at week 8 and week 12 after infection (20 months of age and 21 months of age, respectively) (iPaD/8wpi and iPaD/12wpi, respectively, in FIG. 7), and thus it was suggested that neurogenesis is continuously activated by iPaD lentivirus-induced *Plagl2* overexpression and *Dyrk1a* knockdown in endogenous neural stem cells in the aged brain. *Plagl2* alone also activated neurogenesis, but less efficiently than the iPaD lentivirus (FIG. 8). Combining *Plagl2* overexpression with *Dyrk1a* knockdown resulted in remarkable effects on activation of neural stem cells and the formation of neurons. These results suggested that NSCs activated by the iPaD lentivirus proliferate continuously and are not exhausted even after 3 months. In wild-type hippocampi, MCM2⁺ or DCX⁺ cells decreased significantly in number with aging, and almost disappeared at 18 months of age (FIG. 9-1). However, the iPaD lentivirus increased the number of these cells in the hippocampal dentate gyrus of the aged brain (19 months of age) to a level comparable to that in 9-month-old or even younger wild-type mice (FIG. 9-2), and thus it was suggested that neurogenesis in the aged brain can be effectively reactivated by the iPaD lentivirus.

[0114] In order to check whether or not the iPaD lentivirus has a therapeutic effect on Alzheimer's disease, the iPaD lentivirus was introduced into the hippocampal dentate gyrus of Alzheimer model mice, and the same analysis as

above was carried out. As a result, the iPaD lentivirus efficiently activated NSCs in the Alzheimer model mice and induced the formation of IPCs and immature neurons. In addition, amyloid β deposition was suppressed, and memory improvement was observed in fear conditioning experiments. The details thereof are described in the following section “4. Therapeutic effects of iPaD on Alzheimer’s disease.”

[0115] In addition, in order to investigate whether or not the iPaD lentivirus can activate non-NSCs, the iPaD lentivirus was injected into a brain region in which no neural stem cells were present. The *Hes5* promoter is also active in astrocytes, but none expressed MCM2 or DCX, and thus it is suggested that the activation induced by the iPaD lentivirus is specific to NSCs.

3. Improvement of Cognitive Functions by iPaD (Barnes Maze Test)

[0116] In order to investigate whether or not activated neurogenesis improves cognitive functions in aged mice, spatial learning and memory were analyzed with a Barnes maze test.

[0117] By using this test, it was previously shown that spatial learning and memory depend on neurogenesis in the hippocampal dentate gyrus (Imayoshi, I., et al. Roles of continuous neurogenesis in the structural and functional integrity of the adult forebrain. *Nat Neurosci* 11, 1153-1161 (2008)), and it was shown that aged mice require longer distances and latencies to reach the target hole in the training session and exhibit poorer performance than young mice (Shoji, H. & Miyakawa, T. Age-related behavioral changes from young to old age in male mice of a C57BL/6J strain maintained under a genetic stability program. *Neuropsychopharmacol. Rep.* 39, 100-118 (2019)).

[0118] 19-month-old mice were divided into two groups, which were then injected with the control lentivirus and the iPaD lentivirus, respectively, before one month (FIG. 10-1). The Barnes maze test was carried out on these mice as follows. That is, 10 to 12 male mice per group were used for each behavior analysis. Automatic mouse tracking was carried out by using ANY-maze software (Stoelting Co.) and detecting a plurality of body points. One day before virus injection, 10 mg/kg of P7C3-A20 (MedKoo) was administered by intraperitoneal injection once daily until sacrifice in order to reverse surgical damage. P7C3-A20 was dissolved in a solution of 5% dextrose, 3% DMSO, and 10% Cremophor EL (Nacalai). Spatial memory was measured by using a Barnes circular maze (92 cm in diameter, 12 holes, Brain Science Idea Co., Ltd.). The behavior test space was surrounded by a black curtain, and 4 representative cues were placed at least 30 cm away from the edge of the maze. Habituation was carried out with one escape hole for 5 minutes. One day after a habituation phase, each mouse was trained twice daily to memorize the location of one escape hole, which was randomized among mice but fixed for each individual mouse. At the beginning of the training, a mouse was placed in a white cylinder, which was located in the center of the maze for 30 seconds, and the cylinder was configured to be uncovered when recording started. Each recording was automatically stopped up to 5 minutes after the mouse entered the escape hole or hovered around the escape hole for 10 seconds. When the mouse did not find the escape hole during the 5-minute training, the experimenter gently guided the mouse to the escape hole. The mouse was left undisturbed in the escape hole for 60 seconds after

entering the escape hole. After training, the first probe test was carried out for 3 minutes at intervals of one day. During the training, a camera placed directly above the maze recorded the total distance, the number of times the mouse circled around the correct hole and wrong holes, and the latency to enter the goal hole. In the probe test, the total distance, the number of visits, and the time spent around the correct and wrong holes were recorded. The intensity of light was set to 100 lux on training day 1, and increased by about 30 lux per day. The maze was thoroughly washed with a 70% ethanol solution and dried after every trial.

[0119] As a result, it was found that mice injected with the iPaD lentivirus reached the target hole at shorter distances, made fewer errors, and had shorter latencies than mice injected with the control lentivirus (FIG. 10-2). In the first probe test (1 day after training), iPaD mice stayed longer in the target hole than control mice (FIG. 10-3). These data show that injection of the iPaD lentivirus improved the cognitive functions of aged mice in the Barnes maze test.

4. Comparison Between Alzheimer’s Disease Model Mice and Normal Mice

(1) Comparison of the Number of MCM2-Positive Cells and the Number of DCX-Positive Cells in Hippocampal Dentate Gyrus and Amyloid β Deposition

[0120] 5xFAD mice were used as Alzheimer’s disease model mice. The 5xFAD mice are (Tg6799 line) mice into which a mutant human amyloid precursor protein (APP) (Swedish mutation: K670N, M671L; Florida mutation: 1716V; London mutation: V717I) gene and a mutant human presenilin 1 (PS1) (M146L; L286V) gene have been introduced under the control of the mouse *Thy1* promoter. In these mice, amyloid β is deposited in a brain region, neurogenesis is also reduced, and memory and learning abilities are impaired. The 5xFAD mice (B6/SJL genetic background) were purchased from Jackson Laboratory and crossed with C57BL/6J wild-type mice for 2 generations or more before use. Wild-type littermate mice were used as control animals. All mice were handled according to the “Regulations on Animal Experimentation at Kyoto University.” The experimental protocol was approved by the Animal Experimentation Committee of the Institute for Frontier Life and Medical Sciences, Kyoto University.

[0121] Sections of the hippocampal dentate gyrus were prepared from 4-month-old, 6-month-old, and 8-month-old Alzheimer’s disease model mice (5xFAD mice) and normal mice (wild-type littermate mice), and immunostained to determine the number of MCM2-positive cells and the number of DCX-positive cells to check changes with age in months. Results thereof are shown in FIG. 11 (MCM2-positive cells) and FIG. 12 (DCX-positive cells). Quantitative evaluation of labeled cells was carried out according to the method described in the section “Quantitative evaluation of labeled cells.” In addition, sections of the hippocampal dentate gyrus were prepared from 4-month-old, 6-month-old, and 8-month-old Alzheimer’s disease model mice (5xFAD mice), and changes with age in months in amyloid β deposition were also checked. Results thereof are shown in FIG. 13. Quantitative evaluation of amyloid β deposition was carried out according to the method described in the section “Quantitative evaluation of amyloid β deposition.”

(Quantitative Evaluation of Labeled Cells)

[0122] The hippocampal dentate gyrus of 3 or more mice in each experiment was used for analysis. For each animal, 50 μm thick serial sections of the hippocampal dentate gyrus anterior to the iPaD lentivirus injection point were prepared and evaluated every seventh section. All sections except damaged sections were used for immunostaining. After immunostaining, z-stack images were acquired by using a confocal microscope (LSM780) with a 10 $\times$ or 20 $\times$ objective lens. Stained cells were counted by using Imaris software (Bitplane) or ImageJ and quantified as the number of cells per mm^3 of the hippocampal dentate gyrus.

(Quantitative Evaluation of Amyloid β Deposition)

[0123] Every seventh brain section from each mouse was labeled with a mouse anti- β -amyloid primary antibody (Clone 6E10; 1:500; BioLegend 9320) and detected with a fluorescently labeled secondary antibody. After immunostaining, z-stack images were acquired by using a confocal microscope (LSM780) with a 10 $\times$ objective lens. The number of A β plaques was quantified by using ImageJ after adequate thresholding and denoising. Results thereof were expressed as the number of plaques per mm^3 of the hippocampal dentate gyrus.

[0124] As shown in FIG. 11, the number of cells positive for MCM2, a marker for activated NSCs, was about 1000 cells/ mm^3 in normal mice at 4 months of age, and decreased with increasing months of age, reaching 700 cells/ mm^3 at 6 months of age, and about 680 cells/ mm^3 at 8 months of age. On the other hand, in the Alzheimer's disease model mice (5 $\times$ FAD mice), the number of MCM2-positive cells was as small as about 350 cells/ mm^3 even at 4 months of age, and remained small with no great change even with increasing months of age.

[0125] As shown in FIG. 12, the number of cells positive for DCX, an undifferentiated neuron marker, was about 2700 cells/ mm^3 in normal mice at 4 months of age, and decreased with increasing months of age, reaching 1500 cells/ mm^3 at 6 months of age, and about 800 cells/ mm^3 at 8 months of age. On the other hand, in the Alzheimer's disease model mice (5 $\times$ FAD mice), the number of DCX-positive cells was as small as about 1200 cells/ mm^3 even at 4 months of age, and further decreased with increasing months of age, reaching about 800 cells/ mm^3 at 6 months of age and about 150 cells/ mm^3 at 8 months of age.

[0126] As shown in FIG. 13, in the Alzheimer's disease model mice (5 $\times$ PAD mice), amyloid β deposition (amyloid β plaques) remarkably increased with increasing months of age, and was about 1000 plaques/ mm^3 at 4 months of age, reaching about 2400 plaques/ mm^3 at 6 months of age, and about 3700 plaques/ mm^3 at 8 months of age.

(2) Behavior Tests

[0127] Behavior tests (Barnes circular maze test, contextual fear conditioning test) were carried out on Alzheimer's disease model mice (5 $\times$ FAD mice) and normal mice (wild-type littermate mice) according to the methods described below. Results thereof are shown in FIG. 14 (Barnes circular maze test) and FIG. 15 (contextual fear conditioning test).

[0128] 8 to 12 male mice per group were used for each behavior analysis. Automatic mouse tracking was carried out by using ANY-maze software (Stoelting Co.) to detect a plurality of body points. In the case of a test with virus

injection, 10 mg/kg of P7C3-A20 (manufactured by MedKoo) was administered intraperitoneally once daily for one week from 1 day before virus injection to reverse surgical damage. P7C3-A20 was dissolved in a solution of 5% dextrose, 3% DMSO, and 10% Cremophor EL (Nacalai).

[0129] Spatial memory was checked by the same Barnes maze test as in the above section "3. Improvement of cognitive functions by iPaD (Barnes maze test)."

[0130] Contextual memory was evaluated by a contextual fear conditioning test with partial modification of the protocol described in a previous report (Walgrave, H. et al., Cell Stem Cell 28, 1805-1821.e1808, doi: 10.1016/j.stem.2021.05.001 (2021)). Twenty-four stainless steel rods wired to an electric shock generator were attached to the floor of a chamber (17 $\times$ 10 $\times$ 10 cm), and the test was carried out in an acoustically attenuated cubicle having 60 dB of background white noise. On day 1, each mouse was placed in a chamber (context A, floor: stainless steel rods, walls: colorless and transparent) and allowed to explore for 3 minutes. After that, a foot shock of 80 mA for 2 seconds was applied. The mouse was allowed to remain in the chamber for an additional minute. The chamber was disinfected with 70% ethanol and dried for each animal. After 24 and 48 hours, the mouse was placed back in context A, and the same procedure as on day 1 was repeated. By using TimeFZ2 (O'HARA & CO., LTD, Japan), fear responses during the first 3 minutes were measured.

[0131] In the present Example, the data are expressed as mean $\pm$ standard error (SEM). Statistical analysis was carried out by using R software. Statistical differences were examined by using one-way analysis of variance or two-way analysis of variance, and further, a Tukey-Kramer method or a Bonferroni method was used as a post hoc test. A P-value < 0.05 was considered significant.

[0132] As shown in FIGS. 14 and 15, it can be seen that the Alzheimer's disease model mice (5 $\times$ FAD mice) are inferior to the normal mice in both spatial memory ability and contextual memory ability.

5. Therapeutic Effects of iPaD on Alzheimer's Disease

[0133] In order to investigate whether or not the iPaD lentivirus has a therapeutic effect on Alzheimer's disease, the control lentivirus or the iPaD lentivirus was introduced into the hippocampal dentate gyrus of 5-month-old 5 $\times$ FAD mice. After 4 weeks (6 months of age) and after 12 weeks (8 months of age), 50 μm thick serial sections of the hippocampal dentate gyrus anterior to the lentivirus injection point were prepared, and the number of MCM2-positive cells, the number of DCX-positive cells, and amyloid β deposition was checked and evaluated every seventh section. In addition, behavior tests (Barnes circular maze test and contextual fear conditioning test) were also carried out 4 weeks after the introduction of the iPaD lentivirus. Results thereof are shown in FIG. 16 (MCM2-positive cells), FIG. 17 (DCX-positive cells), FIG. 18 (amyloid β deposition), FIG. 19 (Barnes circular maze test), and FIG. 20 (contextual fear conditioning test). The quantitative evaluation of labeled cells was carried out according to the method described in the above section "Quantitative evaluation of labeled cells," and the quantitative evaluation of amyloid β deposition was carried out according to the method described in the above section "Quantitative evaluation of amyloid β deposition," the Barnes circular maze test was carried out according to the method described in the above

section “3. Improvement of cognitive functions by iPaD (Barnes maze test),” the contextual fear conditioning test was carried out according to the method described in “(2) Behavior tests” in the above section “4. Comparison between Alzheimer’s disease model mice and normal mice.”

[0134] As shown in FIG. 16, in the iPaD lentivirus-introduced mice, the number of cells positive for MCM2, a marker for active NSCs, remarkably increased at week 4 after administration. At week 12 after administration, the number decreased from that at week 4, but was significantly larger than in the control.

[0135] As shown in FIG. 17, in the iPaD lentivirus-introduced mice, the number of cells positive for DCX, an immature neuron marker, remarkably increased at week 4 after administration. At week 12 after administration, the number decreased from that at week 4, but was significantly larger than in the control.

[0136] As shown in FIG. 18, in amyloid β deposition, at week 4 after the introduction of the iPaD lentivirus, there was no difference from the control lentivirus-introduced control, but at week 12 after introduction, a remarkable suppression effect was observed. That is, in the control, the number of amyloid β plaques increased to twice or more that at week 4, whereas in the iPaD lentivirus-introduced mice, the increase was suppressed to about 1.2 times that at week 4.

[0137] As shown in FIG. 19 and FIG. 20, in the iPaD lentivirus-introduced mice, spatial memory and contextual memory were restored to the levels thereof in the normal mice shown in FIG. 14 and FIG. 15, respectively.

[0138] As described above, it was found that the introduction of the iPaD lentivirus remarkably improved a symptom of Alzheimer’s disease and provided excellent therapeutic effects.

6. Induction of Differentiation from Inner Ear Supporting Cells to Hair Cells by iPaD

[0139] iPaD was expressed in supporting cells (Hensen cells) of the inner ear of one-month-old young mice for two

weeks. In many cases, a population consisting of several BrdU+ cells was formed (one supporting cell is thought to have divided into several cells), but the population shown in this figure consisted of 11 BrdU+ cells (FIG. 21). It is considered that one supporting cell divided into 11 cells. Among these, 3 cells were mCherry+ (Nos. 2, 3, 7). In the inner ear, the Hes5 promoter works weakly only in supporting cells but does not work in other cells, and thus it was suggested the possibility that eight mCherry-negative cells changed from supporting cells to other cell types. One (No. 11) of these cells was Myosin6-positive and was considered to be differentiated into a hair cell. Supporting cells at this stage have already lost the cell proliferation ability thereof, and thus it was suggested the possibility that iPaD can efficiently restore the proliferation ability of the cells and differentiate at least some thereof into hair cells.

INDUSTRIAL APPLICABILITY

[0140] According to the present invention, by introducing a specific gene set into an adult brain or an aged brain, endogenous neural stem cells can be efficiently activated and proliferated to produce a large number of neurons. The introduction thereof can cause neural stem cells in the aged brain to revert to their adolescent or younger state and can produce a large number of neurons, resulting in improved memory and learning abilities. In addition, a therapeutic effect thereof on a neurodegenerative disease such as Alzheimer’s disease can also be expected, and endogenous neural stem cells can be efficiently activated to produce a large number of neurons, resulting in improved memory and learning abilities. Further, the specific gene set of the present disclosure can differentiate not only brain neural stem cells but also supporting cells in the inner ear into hair cells, and it is suggested that the specific gene set is also effective in the treatment of an inner ear disease.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 14

<210> SEQ ID NO 1

<211> LENGTH: 1491

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
atgaccacat ttttcaccag cgtccccccc tggattcaag atgcaaagca ggaggaggaa      60
gtgggctgga aactagttcc caggcctcgg ggccgggagg cggagagtca agtgaagtgc      120
caatgtgaaa tttcgggaac acctttctca aatggggaga agctgaggcc tcacagcctc      180
ccgcaaccag agcagagacc atatatgtgc cctcagctgc actgtggcaa ggcttttgct      240
tccaaataca agctgtatag gcacatggcc acccactcag ccagaaacc ccaccagtgt      300
atgtactgtg ataagatgtt tcaccgcaag gaccatctgc ggaaccatct gcagacccat      360
gatcctaaca aagaggccct ccaactgtct gagtgcggta agaattacaa tacgaagctg      420
ggctaccggc gccacctggc catgcatgct gccagcagcg gtgacctcag ctgcaaggtg      480
tgctgcaga cctttgagag taccagggcc ctgctagagc acctgaaggc ccaactcacgc      540
cgggtagcag gcggtgccaa ggagaagaag caccctgtg accactgcga cggcggttc      600
```

-continued

tatactcgta aggatgtacg gcggcaccta gtggtgcaca caggccgtaa ggacttctg	660
tgccagtact gtgcccagcg gtttgccgt aaggaccacc tgacgcgtca tgtaagaag	720
agccactcgc aggagctgct caagatcaag acagagcccg tggacatgtt aggcctactc	780
agctgcagct ccacagtcag tgtgaaggaa gagctgagcc ctgtgctgtg catggcctct	840
cgggacgtaa tggggaccaa ggccttcctt ggcattgtgc ccatgggcat gtatggtgcc	900
cacatcccta ccatgccag cagggcgctg ccacactccc tgggtgcaca cacgtgccc	960
atgggtatga gctacctctt ggaatcctca cctatctctt cccagctca gtcctctcca	1020
aaataaccagc ttggatctac ctcatcttg cccgacaaat tgcccaaagt ggaggtggat	1080
agttttctgg cggagcttcc tggaaacctg tctctctcat ccgtgaacc ccagcccgcc	1140
tcacctcagc cggcggcagc tgcggccctc ctatagtaag cactgcttgc caagagcccc	1200
gccaaacctct ctgaggccct ctgcctgctt aatgtggact tctccacct actgggcttt	1260
cttccactca acctgcccc gtgtaacca cctggggcca caggaggcct ggatcatgggc	1320
tactcccagg ctgaggcaca gcccctgctt accactttgc aagctcagcc tcaagattcc	1380
ccaggagctg ggggaccact gaactttggg cctctgcact ccttgctcc tgtcttcacg	1440
tctggcctga gtagcaccac cctgcctcgt ttccatcaag cattccagta g	1491

<210> SEQ ID NO 2

<211> LENGTH: 5518

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

gggagcgcgg cgccgggagcc cgaggctgag actcaccgga ggaagcggcg cgagcgcgcc	60
gccatcgtcc cggtgaagt gcagtggcac gatcataagt ggctactaca gtctcagatt	120
cctgggctta agcagctctc ccacctcagc ctccccagta gctgggatta cagtgttata	180
gttttgccgc tggactcttc cctcccttcc cccaccccat caggatgata tgagacttga	240
aagaagacga tgcatacagg aggagagact tcagcatgca aaccttcac tgctcggtt	300
gcaccgtcat ttctattcca tgetgctggc cttcagatgg ctggacagat gcccattca	360
catcagtaca gtgaccgtcg ccagccaaac ataagtgacc aacaggtttc tgcttatca	420
tattctgacc agattcagca acctctaact aaccagggtga tgctgatat tgctatgta	480
cagaggcgga tgcccaaac ctccgtgac ccagcaactg ctccctgag aaaactttct	540
gttgacttga tcaaacata caagcatatt aatgaggttt actatgcaa aaagaagcga	600
agacaccaac agggccaggg agacgattct agtcataaga aggaacggaa ggtttacaat	660
gatggttatg atgatgata ctatgattat attgtaaaa acggagaaaa gtggatggat	720
cgttacgaaa ttgactcctt gataggcaaa gggttccttg gacaggttgt aaaggcatat	780
gatcgtgtgg agcaagaatg ggttgccatt aaaataataa agaacaagaa ggcttttctg	840
aatcaagcac agatagaagt gcgacttctt gagctcatga acaaacatga cactgaaatg	900
aaatactaca tagtgcattt gaaacgccac ttatgtttc gaaacctct ctgtttagtt	960
tttgaatgc tgcctacaa cctctatgac ttgctgagaa acaccaattt ccgaggggtc	1020
tctttgaacc taacacgaaa gtttgcgaa cagatgtgca ctgcactgct tttccttgcg	1080
actccagaac ttagtatcat tcactgtgat ctaaaacctg aaaatctct tctttgtaac	1140

-continued

cccaaacgca	gtgcaatcaa	gatagttgac	tttggcagtt	cttgtcagtt	ggggcagagg	1200
atataccagt	atattcagag	tcgcttttat	cggtctccag	aggtgctact	gggaatgcct	1260
tatgaccttg	ccattgatat	gtggctccctc	gggtgtattt	tggttgaaat	gcacactgga	1320
gaacctctgt	tcagtgggtgc	caatgaggta	gatcagatga	ataaaatagt	ggaagtctctg	1380
ggtattccac	ctgctcatat	tcttgaccaa	gcacaaaaag	caagaaagtt	ctttgagaag	1440
ttgccagatg	gcacttgga	cttaaagaag	accaaagatg	gaaaacggga	gtacaaacca	1500
ccaggaaccc	gtaaacttca	taacattctt	ggagtggaaa	caggaggacc	tggtgggcga	1560
cgtgctgggg	agtcaggta	tacggtcgct	gactacttga	agttcaaaga	cctcatttta	1620
aggatgcttg	attatgaccc	caaaactcga	attcaacctt	attatgctct	gcagcacagt	1680
ttcttcaaga	aaacagctga	tgaaggtaca	aatacaagta	atagtgtatc	tacaagcccc	1740
gccatggagc	agtcctcagtc	ttcgggcacc	acctccagta	catcgtcaag	ctcaggtggc	1800
tcacggggga	caagcaacag	tgggagagcc	cggtcggatc	cgacgcacca	gcacggtcac	1860
agtgggtggc	acttcacagc	tgcctgtcag	gccatggact	gcgagacaca	cagtcctccag	1920
gtgagctcgc	acgtggttca	tttcttctgt	tcacctgcca	ttctcaggtg	gagcagcact	1980
ggatgccagg	tgcctttaga	atgaccgtat	catttaccct	agagggtcat	gacgttccctg	2040
tctaaggaaa	tgtgcgtcag	caatttccctg	ctcctcttgg	ttggtcaggc	actgaagctc	2100
ctacacaggt	cactgttgaa	actcctcctg	ttcaagaaac	aacctttcat	gtagccctc	2160
aacagaatgc	attgcatcat	caccatggta	acagttccca	tcaccatcac	caccaccacc	2220
accatcacca	ccaccatgga	caacaagcct	tgggtaaccg	gaccaggcca	aggggtctaca	2280
attctccaac	gaatagctcc	tctacccaag	attctatgga	ggttggccac	agtcaccact	2340
ccatgacatc	cctgtctctc	tcaacgactt	cttcctcgac	atcttccctc	tctactggta	2400
accaaggcaa	tcaggcctac	cagaatcgcc	cagtggctgc	taataccttg	gactttggac	2460
agaatggagc	tatggacgtt	aatttgaccg	tctactccaa	tcccgcctaa	gagactggca	2520
tagctggaca	tccaacatac	caattttctg	ctaatacagg	tcctgcacat	tacatgactg	2580
aaggacatct	gacaatgagg	caaggggctg	atagagaaga	gtcccccatg	acaggagttt	2640
gtgtgcaaca	gagtcctgta	gctagctcgt	gactacattg	aaacttgagt	ttgtttcttg	2700
tgtgttttta	tagaagtggt	gttttttttc	caaaaacaaa	gtgcaaagct	gcttgaatca	2760
ggaggagatt	aacacactga	accgctacaa	gagggcataa	ctgatttttt	ttttaacttg	2820
aaaagattgc	aaaggacat	tgaagtgttt	aaaagagcca	tgtccaaacc	catcttcatg	2880
gatagctcag	aggatctctc	tttttgctcc	cccattttta	cttgccacat	cccagtcaca	2940
gtgggggttt	tttgtctttc	tattcagcaa	aagttaatat	tcagatgttg	gtcttgggtca	3000
tttgccaact	aattttaaag	taaaaggcac	tgcacataat	ttgcataaag	ggccccatga	3060
gggtgttttt	tttttttctt	tttgtccccc	ccatccccct	ttttttttgt	tttgtctctg	3120
tttgttttgg	gtgggagggt	gggaaatttg	ggttttttaag	tcctctaaac	acacttgggc	3180
acggaaatgc	agtactgtaa	ggaagaggga	cctccagctt	ccacaaacac	catcttcagc	3240
tgtatgaaag	ggacggttgt	ggtgaagttt	gtcaggcaca	gtaagcatgc	tgagtggcgg	3300
ggatcagaac	tctcctatct	gaacctactg	aggagcaaa	gagcaattac	atgggatcct	3360
gtggctctcc	cggtgcagag	gccacaggaa	gataggatgg	aacgtgactg	gtctcctaac	3420

-continued

caaggtgcac tgagaagcaa tcaacgggtc ggctcgtggcc agtcctgggg aggtctgagt 3480
gggtgtcttt gggataacct ttggccttat ggatttggac tcgaaattag aagagcctac 3540
catttcagat gcaatcactt ttggacatgc ttttgcagac agtccttaat gctgaaaaca 3600
cagagaatgg gtaattcaag aggcctttct tttaaaatag acttttgtga cccactaatt 3660
gtaaggatt gcaaggtcac ttgctgtgtg tcataaagtt gacttcctta ttggttgaag 3720
gtcacagaag tagtggtttg ctttgatgga aatagctaca gctgtgtccc ttctgtctt 3780
ttacttttct ttttgccttt tctcggcacg tggatctccc accatttctt ctgcacaaag 3840
atgtcttctg ttcactctga acatttttaa aaaatgcaga attttatgtg actgcttttt 3900
tgccctacaa ttatgctgtg aattttacaa aaatttattt tcttttttga taattttattg 3960
taccaaagct gtttttatag cacatagatg tctgtaacca ataagttagc agttctgcac 4020
tttgacacaa ggtgtaacta gaccattttt aaatgtcagt tgaaaattat ggctgtacta 4080
ttgcttaaac aaaactggaa ctgttggtga atccatagcc aatacattta cagcaatctg 4140
tgtactgaac atagtagatt gacatctaata tcaagattac aacatctgtt acattctaag 4200
tgtgttcagg cttctgaagg taaagggaca ctggatccag aagctatgga accagcagtt 4260
gattcttcta ttctgtatta acctacttgt aaacttgaaa gcaagacctt gattgcacca 4320
acaggtccag agtatgagtg caagcaaacg agaactctca tgcgtgacct gagcagacag 4380
gctggtatct aacaggtgcc tctgtgtgag attacgctgc cttaatgtaa cacagtctgg 4440
cagttgctaa atttgtgttc ccatttttaa ttgaccaatt ttgggggtgtg acacttttga 4500
gcggttgaat tgggagaatg aagataagta atttacctgt ccaggatcaa aagaagccta 4560
gaaaagaagc agtaatctac ctctgccgat aacctgttta agatgactca gcagaacacc 4620
gcggttcatt ctattgtgca attccatgtg gctgactagg tcaatttttt ttctgaacaa 4680
aagcaggttt ttatatgtaa acagtgagaa aagaaaggct aaacactatg taaatgtgaa 4740
tggaacttg gaaatactcg tttttataaa ctacaaaaac ttttgtgtgt ttatcaggaa 4800
atccatattt attttgtaat taactgtcaa gcctgtggat gatttttttg aacttggtag 4860
ttcataaagg tttacagtga ataaaaggat atcatcttga gtatagcaat atcaaaagga 4920
attcagtagt tactgctgtt taggaatata aggttaagat atcatatggg tcaggtcatt 4980
tttttttct gtgctggttg ccacatctta gcaagcacca aaaaactaaa gcagttttta 5040
aacgatatt tacgtaaaaga aaatcataaa atccaatgct tctgcatact gtgttatgtt 5100
acagtcagtg tttgtgtgct ttactacaca gtttggttac aggacttctg tgcattgtaa 5160
acataaacag catggaaaag gttaaatacc tgtgttcaga ttgtaagatc tagtccggac 5220
ttgctgtgta tattgtaacg ttaaatgaaa aaagaacccc cctttgtatt atagtcatgc 5280
ggctctatgt atgataaaca gttgaataat ttgtcctcag actctttact atgctttttt 5340
aaaaattaat ttaagaaaaa tgtaaacata gtaaaaatct tcctatgcaa ttaaactggt 5400
ccaggctctg taggtatagt atcaaagtg agttaaatgt gtaaaaagga aactatttga 5460
gatacattga cataggcatc agcaatctct gaaagtaaaa attggagggt taacagat 5518

<210> SEQ ID NO 3

<211> LENGTH: 337

<212> TYPE: DNA

<213> ORGANISM: Artificial

-continued

<220> FEATURE:

<223> OTHER INFORMATION: shRNA

<400> SEQUENCE: 3

tagggataac agggtaattg tttgaatgag gcttcagtag tttacagaat cgttgccctgc	60
acatcttgga aacacttgct gggattactt cgacttctta acccaacaga aggtcgcaga	120
aggtatattg ctgttgacag tgagcgccct gattaaccta cttgtaaata gtgaagccac	180
agatgtatgtt acaagtaggt taatcaggat gcctactgcc tcggacttca aggggctaga	240
attcgagcaa ttatcttggt tactaaaact gaataccttg ctatctcttt gatacatgtt	300
tacaaagctg aattaaaatg gtataaatta aatcact	337

<210> SEQ ID NO 4

<211> LENGTH: 1506

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

aaggtctctc attggacct gccagctggc cagggaaccc cactcctgga ttcaaacctc	60
gcctaaccag ccctgatgcc agcccagct ctgggggcta aggcccttcc gctaggacct	120
ggcaggggtc caacctgtct ctaggagcca gccagggccc cgaatectct cctagtctct	180
gtgcagagc atttgaggaa ctgtgaccgc aggttagggg gtgggggtct cccaagggac	240
cctccctaa ggacccaggg tgggattcat tgctcaaag tggctgcggg cactcagctc	300
agttggccag gtaccggggt tattagctca tcccatggac cggtggttgg gggcgggggg	360
ctagggtctg catggcctct gatectcaag tctgccacct ggcacccact cctaccac	420
tggcaaatg ccagaggaaa gcaagcccta caagtgcctt ggctgccaga gagactaggt	480
ggagctgtgg catgcataaa cagtggccac agcaagcatg ccaagagccg aggcctcccg	540
agctgcgtgg aactgtagt cagatccctg gccagatagg ggtggggaga gggccctgag	600
ggacacctgc cagctcgtgg agcagcccca gtgctgctgt gggttacagt gctcctggga	660
ggacagggtg cagggaagcg ggagagctgc cgggtgcccc gctccgcggg gagagctgcc	720
gggtgcccgg ctcacgttg agcacgggccc ccagcttctt ggagctgccc ggagcccagc	780
caggaaggca caaggcccat ttcttaggtc tggagaagtc aggagtccca ggtgacagtc	840
agggggcttg taatataggc acgtaggact gatcctaaga gcgtgcccc ctctgtcccc	900
ctccctgga ggggcagcag gtcccagtc ggggaaggtg ccatccccc ggaagccgtg	960
gcacttggtc attctacagt ttttccctta tctgctccta cggcctctgc cagccagcag	1020
tgggaagccg ctggggagag gcaggggagg ggagcggagg ggagggcggg ggagggggcc	1080
ctccttctg cccccctccc gtttggcggc ctgggaaaag gcagcatatt gaggcgcggg	1140
gctcccgga tcaggccccg ggatgctaag gaggcgagc gcgttccac agcccggaca	1200
ttgtgccgcg cggcccaact gctccttggg gagcgcccat tgtgcccgcg ccaattcaca	1260
ggcaatttag cgtgcgctaa tgggcggcg cctttgtgct gcccgcccgg ccattggccg	1320
cggagtgtgg gaacggccgc ggcgcccga ctcaggcgc caggccgcgg ccgcgccta	1380
tatagggcgt cggcgcgagg ggcgggtgt cgcgcagcc cgggacgcgc ttggccttgc	1440
ccgcgcgcgc tcgctctgtc tcgcccggcc tccccggtc gcctcgtcgc ctgttccgcg	1500
ccaggc	1506

-continued

<210> SEQ ID NO 5

<211> LENGTH: 4227

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

aaggtcctc attggaccct gccagctggc cagggaaacc cactcctgga ttcaaactct	60
gcctaaccag ccctgatgcc agccccagct ctgggggcta aggcccttcc gctaggagccc	120
ggcagggtcc caacctgtct ctaggagcca gcccaggccc cgaatcctct cctagtctct	180
gtgtcagagc atttgaggaa ctgtgaccgc aggctagggg gtgggggtct cccaaggagc	240
cctcccctaa ggacccaggg tgggattcat tgcctcaaag tggctgcggg cactcagctc	300
agttggccag gtaccggggt tattagctca tcccatggac cgggtggttg gggcgggggg	360
ctagggtctg catggcctct gatectcaag tctgccacct ggcacccact ccctaccac	420
tggcaaaatg ccagagggaa gcaagcccta caagtgccct ggctgccaga gagactaggt	480
ggagctgtgg catgcataaa cagtggccac agcaagcatg ccaagagccg aggcctcccg	540
agctgcgtgg aactgtagt cagatccctg gccagatagg ggtggggaga ggcccctgag	600
ggacacctgc cagctcgtgg agcagcccca gtgctgctgt gggttacagt gctcctggga	660
ggacagggtg caggaaagcg ggagagctgc cgggtgcccg gctccgcggg gagagctgcc	720
gggtgccggg ctccacgtgg agcacgggcc ccagcttcct ggagctgccg ggagcccagc	780
caggaaaggc caaggcccat ttctaggtc tggagaagtc aggagtccca ggtgacagtc	840
agggggcttg taatataggc acgtaggact gatcctaaga gcgtgccccc ctctgtcccc	900
ctcccctgga ggggcagcag gtcccagtc ggaagggtgc ccatacccca ggaagccgtg	960
gcacttgctc attctacagt ttttcctta tctgtccta cggcctctgc cagccagcag	1020
tgggaagccg ctggggagag gcaggggagg ggagcggagg ggagggcggg ggagggggcc	1080
ctccttctct cccccctccc gtttgccggc ctgggaaaag gcagcatatt gaggcgcggg	1140
gtcctccgga tcaggccccc ggtatgtaac gagggcagac gcgttccac agcccgga	1200
ttgtgcccg cggcccacct gctccttggg gagcgcctat tgtgcccgcg ccaattcaca	1260
ggcaatttag cgtgcgctaa tgggcccggc cctttgtgcg gcccgcccg ccattggccg	1320
cggagtgttg gaacggccgc ggcgcgggga ctccaggcgc caggccgcgc ccgcgccta	1380
tatagggcgt cggcgccggg ggcgggtgtc cgcgccagcc cgggaacgcg ttggccttgc	1440
ccgcgcccg tcgcctcgtc tcgccgggcc tccccgcgtc gcctcgtcgc ctgttcgcg	1500
ccaggcttaa ttaagccacc atgaccacat ttttaccag cgtcccccc tggattcaag	1560
atgcaaaagc ggaggaggaa gtgggctgga aactagttcc caggcctcgg ggccgggagg	1620
cggagagtca agtgaagtgc caatgtgaaa ttccgggaac acctttctca aatggggaga	1680
agctgaggcc tcacagctc ccgaaccag agcagagacc atatagctgc cctcagctgc	1740
actgtggcaa ggcttttgct tccaaataca agctgtatag gcacatggcc acccactcag	1800
cccagaaacc ccaccagtgt atgtactgtg ataagatgtt tcaccgcaag gaccatctgc	1860
ggaaccatct gcagaccat gatcctaaca aagaggccct ccaactgctc gactgcggta	1920
agaattacaa tacgaagctg ggctaccggc gccacctggc catgcatgct gccagcagcg	1980
gtgacctcag ctgcaagggt tgcctgcaga cctttgagag taccagggcc ctgctagagc	2040

-continued

acctgaaggc ccactcacgc cgggtagcag gcggtgccaa ggagaagaag caccctgtg	2100
accactgcga cggcggttc tatactcgta aggatgtacg gcggcaccta gtggtgcaca	2160
caggccgtaa ggacttctcg tgccagtact gtgccacgag gtttgccgt aaggaccacc	2220
tgacgcgtca tgtcaagaag agccactcgc aggagctgct caagatcaag acagagcccg	2280
tggacatgtt aggcctactc agtcgcagct ccacagtcag tgtgaaggaa gagctgagcc	2340
ctgtgctgtg catggcctct cgggacgtaa tggggaccaa ggccttcctt ggcatgttg	2400
ccatgggcat gtatgggtgc cacatcccta ccatgccag cagggcggtg ccacactccc	2460
tggtgcacaa cacgctgcc atgggtatga gctaccctct ggaatcctca cctatctctt	2520
cccagctca gctccctcca aaataccagc ttggatctac ctcatacttg cccgacaaat	2580
tgccaaaagt ggagggtgat agttttcttg cggagcttcc tggaagcctg tctctctcat	2640
ccgctgaacc ccagcccgcc tcacctcagc cggcggcagc tgcggccctc ctagatgaag	2700
cactgcttgc caagagcccc gccaacctct ctgaggccct ctgcgctgct aatgtggact	2760
tctcccact actgggcttt ctccactca acctgcccc gtgtaacca cctggggcca	2820
caggaggcct ggtcatgggc tactcccagg ctgaggcaca gcccctgctt accactttgc	2880
aagctcagcc tcaagattcc ccaggagctg ggggaccact gaactttggg cctctgcact	2940
ccttgctcc tgtcttcacg tctggcctga gtagaccac cctgcctcgt tccatcaag	3000
cattccaggg aagcggagct actaacttca gcctgctgaa gcaggctgga gacgtggagg	3060
agaaccctgg acctgtgagc aagggcgagg aggataacat ggccatcac aaggagtcca	3120
tgcgcttcaa ggtgcacatg gagggctccg tgaacggcca cgagttcgag atcgaggcg	3180
agggcgaggg ccgcccctac gagggcacc agaccgcca gctgaagggtg accaagggtg	3240
gcccctgccc cttgcctgg gacatcctgt cccctcagtt catgtaaggc tccaaggcct	3300
acgtgaagca ccccgctgac atcccagact acttgaagct gtccctcccc gagggcttca	3360
agtgggagcg cgtgatgaac ttcgaggacg gcggcggtgt gaccgtgacc caggactcct	3420
ccctgcagga cggcgagtgc atctacaagg tgaagctcgc cggcaccaac tcccctccg	3480
acggccccgt aatgcagaag aagaccatgg gctgggaggg ctctccgag cggatgtacc	3540
ccgaggacgg cgcctgaag ggcgagatca agcagaggct gaagctgaag gacggcggcc	3600
actacgacgc tgaggtaag accaccta caaggccaaga gcccggtcag ctgccggcg	3660
cctacaacgt caacatcaag ttggacatca cctcccacaa cgaggactac accatcgtgg	3720
aacagtacga acgcgcgag ggcggccact ccaccggcg catggacgag ctgtacaagg	3780
atatccctcc aaagaagaag agaaagggtgt aaggcgcgcc tagggataac agggtaattg	3840
tttgaatgag gcttcagtac ttacagaat cgttgctcgc acatcttga aacacttgct	3900
gggattactt cgacttctta acccaacaga aggctcgaga aggtatattg ctgttgacag	3960
tgagcgccct gattaaccta cttgtaata gtgaagccac agatgtattt acaagtaggt	4020
taatcaggat gcctactgcc tcggacttca aggggctaga attcgagcaa ttatcttgtt	4080
tactaaaact gaataccttg ctatctcttt gatacatctt tacaagctg aattaaaatg	4140
gtataaatta aatcactcta gtaataaagg atcctttatc ttcattggat ccgtgtgttg	4200
gttttttgtg tgcggcccg ctagacc	4227

-continued

```

<210> SEQ ID NO 6
<211> LENGTH: 3011
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6
aaggctcttc attggacct gccagctggc cagggaaccc cactcctgga ttcaaacctct    60
gcctaaccag cctgatgcc agcccagct ctgggggcta aggcccttcc gctaggaccc    120
ggcagggctc caacctgtct ctaggagcca gcccaggccc cgaatcctct cctagtctct    180
gtgcagagc atttgaggaa ctgtgaccgc aggctagggg gtgggggtct cccaagggac    240
cctcccctaa ggaccaggg tgggattcat tgccctaaa tggtgcggg cactcagctc    300
agttggccag gtaccggggt tattagctca tcccatggac cggtggttgg gggcgggggg    360
ctagggtctg catggcctct gatectcaag tctgccacct ggcacccact cctaccac    420
tggaataatg ccagaggaaa gcaagcccta caagtgcctt ggctgcaga gagactaggt    480
ggagctgttg catcataaa cagtggccac agcaagcatg ccaagagccg aggcctcccg    540
agctgcgttg aactgtagt cagatccctg gccagatagg ggtggggaga ggccttgag    600
ggacacctgc cagctcgttg agcagcccca gtgctgctgt gggttacagt gctcctggga    660
ggacagggtg cagggaagcg ggagagctgc cgggtgcccc gctccgcggg gagagctgcc    720
gggtgcccgg ctcacgttg agcacgggcc ccagcttctt ggagctgccc ggagcccagc    780
caggaagcca caaggcccat ttcttaggtc tggagaagtc aggagtccca ggtgacagtc    840
agggggcttg taatataggc acgtaggact gatcctaaga gcgtgcccc ctctgtcccc    900
ctcccctgga ggggcagcag gtcccagtca gggaaggtgc ccatccccc ggaagccgtg    960
gcacttgctc attctacagt tttccctta tctgctcta cggcctctgc cagccagcag    1020
tgggaagccg ctggggagag gcaggggagg ggagcggagg ggagggcggg ggagggggcc    1080
ctccttctcg ccccccctcc gtttgccggc ctgggaaaag gcagcatatt gaggcgcggg    1140
gtcccccgca tcaggcccc ggatgctaag gagggcgagc gcgttccac agcccgga    1200
ttgtgcccg cggcccacct gctccttggg gagcgcccat tgtgcccgcg ccaattcaca    1260
ggcaatttag cgtgcgctaa tgggcggcg cctttgtgcy gcccgcccg ccattggccg    1320
cggagtgttg gaacggccgc ggcgcccgga ctccaggcgc caggcccgcc cccgcgcta    1380
tatagggcgt cggcgcgcg ggcgggtgtc cgcgccagcc cgggacgcgc ttggccttgc    1440
ccgcgcccg tcgcctcgtc tcgcccgcc tcccgcgct gcctcgctgc ctgttccgcg    1500
ccagccttaa ttaagccacc atgaccacat ttttaccag cgtcccccc tggattcaag    1560
atgcaaaaca ggaggaggaa gtgggctgga aactagttcc caggcctcgg ggcggggagg    1620
cggagagtca agtgaagtgc caatgtgaaa ttctgggaac acctttctca aatggggaga    1680
agctgaggcc tcacagcctc ccgcaaccag agcagagacc atatagctgc cctcagctgc    1740
actgtggcaa ggcttttct tccaaataca agctgtatag gcacatggcc acccactcag    1800
cccagaaacc ccaccagtgt atgtactgtg ataagatgtt tcaccgcaag gaccatctgc    1860
ggaaccatct gcagaccat gatcctaaca aagaggccct cactgctct gagtgcggta    1920
agaattacaa tacgaagtgc ggtacccgc gccacctggc catgcatgct gccagcagcg    1980
gtgacctcag ctgcaagggt tgctgcaga cctttgagag taccagggcc ctgctagagc    2040
acctgaagcg cactcacgc cgggtagcag gcggtgcaa ggagaagaag caccctgtg    2100

```

-continued

```

accactgcga cgggoggttc tatactcgta aggatgtacg gcggcaccta gtggtgcaca 2160
caggccgtaa ggacttcctg tgccagtact gtgcccagcg gtttggccgt aaggaccacc 2220
tgacgcgtca tgtcaagaag agccactcgc aggagctgct caagatcaag acagagcccc 2280
tggacatggt aggcotactc agctgcagct ccacagtcag tgtgaaggaa gagctgagcc 2340
ctgtgctgtg catggcctct cgggacgtaa tggggaccaa ggccttcctt ggcattgttg 2400
ccatgggcat gtatgggtgcc cacatcccta ccatgccag cagggcgctg ccacactccc 2460
tgggtgcacaa cagctgccc atgggtatga gctacctctt ggaatcctca cctatctctt 2520
ccccagctca gctcctcca aaataccagc ttggatctac ctcatacttg cccgacaaat 2580
tgcccaaagt ggaggtggat agttttcttg cggagcttcc tggaagcctg tctctctcat 2640
ccgtgaacc ccagcccgc tcacctcagc cggcggcagc tgcggccctc ctatagtaag 2700
cactgcttgc caagagcccc gccaacctct ctgaggccct ctgcgctgct aatgtggact 2760
tctcccacct actgggcttt ctccactca acctgcccc gtgtaaccca cctggggcca 2820
caggaggcct ggtcatgggc tactcccagc ctgaggcaca gccctgctt accactttgc 2880
aagctcagcc tcaagattcc ccaggagctg ggggaccact gaactttggg cctctgcaact 2940
ccttgccctc tgtcttcagc tctggcctga gtagcaccac cctgcctcgt ttccatcaag 3000
cattccagta g 3011

```

```

<210> SEQ ID NO 7
<211> LENGTH: 1921
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: promotor_shRNA

```

```

<400> SEQUENCE: 7

```

```

aaggtctctc attggacct gccagctggc cagggaaccc cactcctgga ttcaaactct 60
gcctaaccag ccctgatgcc agcccagct ctgggggcta aggcccttcc gctaggacct 120
ggcagggctc caacctgtct ctaggagcca gcccaggccc cgaatcctct cctagtctct 180
gtgcagagc atttgaggaa ctgtgaccgc aggctagggg gtgggggtct cccaagggac 240
cctcccctaa ggaccaggg tgggattcat tgctcaaag tggctgcggg cactcagctc 300
agttggccag gtaccggggt tattagctca tcccatggac cggtggttgg gggcgggggg 360
ctagggttgg catggcctct gatectcaag tctgccacct ggcacccact cctacccac 420
tggcaaaatg ccagaggaaa gcaagcccta caagtgcctt ggctgccaga gagactaggt 480
ggagctgtgg catgcataaa cagtggccac agcaagcatg ccaagagccg aggcctccc 540
agctgcgtgg aactgtagt cagatccctg gccagatagg ggtggggaga gggccctgag 600
ggacacctgc cagctcgtgg agcagcccca gtgctgctgt gggttacagt gctcctggga 660
ggacagggtg caggaagcgg ggagagctgc cgggtgcccc gctccgcggg gagagctgcc 720
gggtgccggg ctccacgtgg agcacgggcc ccagcttcct ggagctgcc ggagcccagc 780
caggaaggca caaggcccat ttctaggtc tggagaagtc aggagtccca ggtgacagtc 840
agggggcttg taatatagc acgtaggact gatectaaga gcgtgcgcc ctctgtcccc 900
ctcccctgga ggggcagcag gtcccagtc ggaaggtgc ccatcccca ggaagccgtg 960
gcacttgctc attctacagt tttccctta tctgctcta cggcctctgc cagccagcag 1020

```

-continued

tggaagccg ctggggagag gcaggggagg ggagcggagg ggagggcggg ggagggggcc	1080
ctccttctcg cccccctccc gtttgccggc ctgggaaaag gcagcatatt gaggcgcggg	1140
gctcccggca tcagggcccg gtagtctaag gagggcgagc gcgttccac agcccgga	1200
ttgtgccgag cggccacact gctccttggg gagcgcccat tgtgccgag ccaattcaca	1260
ggcaatttag cgtgcgctaa tgggcccggc cctttgtgag gcccgcccg ccattggcgg	1320
cggagtgtgg gaacggccgc ggcgcccga ctcaggcgc caggccgcg cccgcgccta	1380
tatagggcgt cggcgccggg ggcgggtgtc cgcgccagcc cgggacgcg ttggccttgc	1440
ccgcgccgc tcgcctcgtc tcgcccggc tccccgcgtc gcctcgtcgc ctgttcgcg	1500
ccaggcttaa ttaataggga taacagggtg attgtttgaa tgaggcttca gtactttaca	1560
gaatcgttgc ctgcacatct tggaaacact tgctgggatt acttcgactt ctaacccaa	1620
cagaaggctc gagaaggtag attgctgttg acagtgcgc cctgattaa cctacttga	1680
aatagtgaag ccacagatgt atttacaagt aggttaatca ggatgcctac tgcccgga	1740
ttcaaggggc tagaattcga gcaattatct tgtttactaa aactgaatac cttgctatct	1800
ctttgataca tttttacaaa gctgaattaa aatggtataa attaaatcac tctagtaata	1860
aaggatcctt tatcttcatt gtagccgtgt gttggttttt tgtgtgcggc ccgtctagac	1920
c	1921

<210> SEQ ID NO 8
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: siRNA

<400> SEQUENCE: 8

cctgattaac ctacttgtaa a	21
-------------------------	----

<210> SEQ ID NO 9
 <211> LENGTH: 1618
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 9

tgcaattata tcttgcaactg gacagaccat cctgtgctgc ctcaaagcca agccttcccc	60
tctgtccag atgttccat ccatggtaac ccaacaacc cttgtgctaa gagcttctgg	120
agcaagacct tcggaataga tcatttacag cagtgggtca ctaccatagc aggagggcgt	180
ttccctagtg ggcactgtga aggcccttca gttcctgagc tcagacatgg ctgacatcca	240
tccagagaac ctggtttcat tccacactgg ctattgtgct ctgatgccag cctcctggtt	300
tgggcctggc taggtcttag gttgacctg ggaggaagca cggggcctta tctcctcct	360
cagccccctg agagtattgg ggaatagta cagggtccag atggaggagc gctcctacat	420
ggccatcccc taggtgtcta ggggtgtgact agtaaggta cctgggtagg aaccacggtt	480
atcagttctt acaagaggc ccacccacac tggcacagca ttcatatggg ggcttatttt	540
taggtctccc acctcttacc tgccaacaca aaatgctgaa tgaaactgga acacacacac	600
agaggcatgc acgcgcgcgc ccgcacgagc tgagggggat gtcagagag gagttaaggg	660
cagccatggt ttcctaaaag ccacgcggga gtagtgatca cggacagcat gaaggaggca	720

-continued

cggtcctcct gtttcagaga tgcattgtgc acctggtgac acaagagcag tccagatacc	780
ctgtgggttc aggggttcct gagggcttcc taggtatggg gctttgtgct aggcattatc	840
caaagcaaat tcagctagcc tagaagtacg ctgggcaggc tccgcgcgtg gccatttcta	900
gctgctggaga agtcactcct ctcaactgac agaactggca ataaagcgcc taaattctaa	960
atacatgagt cctctcttca tctgtacct gcgaatggca tggcggcagc atctgtttga	1020
gagaatgtgg cccactttaa cccactgtgc ctgtgggcaa caggtcccag ctggggaggg	1080
tgccactcc tctggaagtc gctgtctgtt cattgtaaaa ctctcatttc tctgcgctg	1140
tggcctctgc cagccagcgg tggggagcct ctggggagtg ggaggggaaga agggagagaa	1200
ggggggggga gagcactcct tctgcccctc cccactccc cgcggcctgg gaaaaggcag	1260
catattgagg cgcggggctc tcagcatcag gcccgggat gctaattagg gcgagcgcgt	1320
tcccacagcc cggacattgt gccgcgcggc ccactgtctc ctggggagc gaccattgtg	1380
cccgcgccaa ttcacaggca atttagcgtg cgctaattgg ccggcgcctt tgtgcggccg	1440
gcgcgcgcct tggcgccga gtgtgggaac ggccgcggcg cccggacccc aggcgcgggg	1500
ccgtgcccc cgctatata gggtggcgt gctgggtcc aggtcgcgc agtccgggac	1560
gcctggctct gtcgcctcg ctccgctccg ctgcctaacc gcctccagag ctccaggc	1618

<210> SEQ ID NO 10

<211> LENGTH: 1491

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 10

atgaccacat ttttcaccag cgtccctccc tggattcaag atgcaaagca ggaggaggaa	60
gtgggctgga aactagtctc caggcctcgg ggccgggagg cagagagtca agtgaagtgc	120
caatgtgaaa tctcggggac accattttca aatggggaga agctaaggcc tcacagcctt	180
ccccatccgg agcagagacc atatagetgt cctcagctgc actgcggcaa ggcttttgcc	240
tccaagtaca agctgtatag gcacatggcc acccactcgg ccagaaaacc ccaccagtgc	300
atgtactgtg acaagatggt tcaccgaaag gaccatctgc ggaaccactt gcagaccat	360
gacccaaca aagaggccct ccactgctct gagtgtggta agaattataa tacaaagctg	420
ggataccgtc gtcacctggc catgcatgct gccagcagtg gtgatctcag ctgcaagggtg	480
tgcctgcaga cctttgagag taccaggcc ctattggagc acctgaaggc tcaactcacgc	540
cgggtagcag gaggtgctaa ggagaagaag caccctgtg accattgtga caggcggttc	600
tatactcgga aggatgtccg acggcaccta gtggcgaca caggtcgtaa ggacttctg	660
tgtcagtact gtgcccagag atttggccgc aaggatcacc tgacacgcca tgtcaagaag	720
agccactcac aggaactgct caagatcaag acagagccag tggatatggt aggettgtctc	780
agctgtagct ctacagtcag tgtaaaggaa gagctgagcc ctgtgctttg catggcctct	840
cgggacgtga tgggggctaa ggcctttcct ggcatgttc ccatgggtat gtatggtgcc	900
cacatccccta ccatgccagc tgcgggcatg ccactctccc tgggtgcacaa cacactgccc	960
atgggtatga gttacctctt ggaatcttct cctatctcct cctcagctca gctccctcca	1020
aaataccagc ttggatctac ctcatacttg cccgacaaac tgcccaaagt ggaggtggat	1080
agttttctgg cagagcttcc tggaagcctg tctctctcct cggctgaacc tcagcccgcc	1140

-continued

tcacctcagc	cgggcggcagc	tcagccctc	ctggatgaag	cactgctcgc	caagagtccc	1200
gccaaacctct	ccgaggccct	ctgcctgct	aatgtggact	tttcccactt	actgggcttt	1260
cttccgctca	acctaccccc	atgtaacca	ccaggggcca	caggaggctt	ggttatgggc	1320
tactcccaag	ctgaggcaca	gcctttgctc	accactttgc	aggctcagcc	tcaagattcc	1380
gctggagctg	ggggacccct	gaactttggg	cctctgcact	ccttacctcc	tgtcttcacc	1440
tctggcctga	gtaccaccac	cctgcctcgt	ttccaccaag	cattccagta	g	1491

<210> SEQ ID NO 11

<211> LENGTH: 3141

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 11

tgcacttata	tcttgcactg	gacagaccat	cctgtgctgc	ctcaaagcca	agccttcccc	60
tctgtcccag	atgcttccat	ccatggtaac	ccaacaaccc	cttgtgctaa	gagcttctgg	120
agcaagacct	tcggaataga	tcatttacag	cagtgggtca	ctaccatagc	aggagggcgt	180
ttccctagtg	ggcactgtga	aggcccttca	gttctctgagc	tcagacatgg	ctgacatcca	240
tccagagaa	ctggtttcat	tccacactgg	ctattgtgct	ctgatgccag	cctcctgggt	300
tgggcctggc	taggtcttag	gttgacccta	ggaggaagca	cggggcctta	tctcctccct	360
cagccctcgc	agagtattgg	ggaaatagta	caggttccag	atggagggac	gctcctacat	420
ggccatcccc	taggtgtcta	gggtgtgact	agtaaggcca	cctgggttagg	aaccacgggt	480
atcagttctt	acaaagaggc	ccatccacac	tggcacagca	ttcatatggg	ggcttatttt	540
taggtctccc	accttctacc	tgccaacaca	aaatgctgaa	tgaaactgga	acacacacac	600
agaggcatgc	acgcgcgcgc	ccgcacgagc	tgagggggat	gtcatgagag	gagttaaggg	660
cagccatggg	ttcctaaaag	ccacgcggga	gtagtgatca	cggacagcat	gaaggaggca	720
cggctcctct	gtttcagaga	tgcatgtggc	acctgggtgac	acaagagcag	tccagatacc	780
ctgtgggttc	aggggttcc	gagggcttcc	taggtatggg	gctttgtgct	aggcattatc	840
caaagcaaat	tcagctagcc	tagaagtacg	cttggcaggc	tccgcgcgtg	gccatttcta	900
gctgcgagga	agtcactcct	ctcaactgac	agaactggca	ataaagcgcc	ttaaattctaa	960
atacatgagt	ccctctctca	tctgtacct	gcgaatggca	tggcggcagc	atctgtttga	1020
gagaatgtgg	cccactttaa	cccactgtgc	ctgtgggcaa	caggteccag	ctggggaggg	1080
tgcccactcc	tctggaagtc	gctgtctgtt	cattgtaaaa	ctctcatttc	tctgcgectg	1140
tggcctctgc	cagccagcgg	tggggagcct	ctggggagtg	ggaggggaaga	agggagagaa	1200
ggggggggga	gagcactcct	tcttgccttc	cccacctccc	cgcggcctgg	gaaaaggcag	1260
catattgagg	cgcggggctc	tcagcatcag	gccccgggat	gctaattgagg	gcgagcgcg	1320
tcccacagcc	cggaattgt	gcgcgcgggc	ccacctgctc	ctcggggagc	gaccattgtg	1380
ccgcgcgcaa	ttcacaggca	atttagcgtg	cgctaattgg	ccggcgccct	tgtgcggccg	1440
gcgcgcgcat	tggcgcgcca	gtgtgggaac	ggcgcggcgc	cccggaaccc	aggcgccggg	1500
ccgtgcgccg	cgcctatata	gggctggcgt	gctggggctc	aggtcgcgcc	agtcggggac	1560
gcctggctct	gtccgctccg	ctccgctccg	ctcgtctaac	gcctccagag	ctccaggcct	1620
taagaggcag	gcaggcttaa	ttaagccacc	atgaccacat	ttttaccag	cgctccctcc	1680

-continued

tggattcaag atgcaaagca ggaggaggaa gtgggctgga aactagtcc caggcctcgg	1740
ggccgggagg cagagagtca agtgaagtgc caatgtgaaa tctcggggac accatattca	1800
aatggggaga agctaaggcc tcacagcctt ccccatccgg agcagagacc atatagtctgt	1860
cctcagctgc actgcgga caa ggcttttgcc tccaagtaca agctgtatag gcacatggcc	1920
acccactcgg ccagaaaacc ccaccagtgc atgtactgtg acaagatgtt tcaccgaaag	1980
gaccatctgc ggaaccactt gcagacccat gacccaaca aagaggccct ccactgctct	2040
gagtgtggta agaattataa tacaaagctg ggataccgtc gtcacctggc catgcatgct	2100
gccagcagtg gtgatctcag ctgcaagggtg tgctgcaga cctttgagag taccaggcc	2160
ctattggagc acctgaaggc tcaactcacgc cgggtagcag gaggtgctaa ggagaagaag	2220
caccctgtg accattgtga caggcggttc tatactcgga aggatgtccg acggcaccta	2280
gtggtgcaca caggctgtaa ggacttctg tgtcagtact gtgccagag atttgccgc	2340
aaggatcacc tgacacgcca tgtcaagaag agccactcac aggaactgct caagatcaag	2400
acagagccag tggatatgtt aggcttctc agctgtagct ctacagtcag tgtaaaggaa	2460
gagctgagcc ctgtgctttg catggcctct cgggacgtga tgggggctaa ggccttcct	2520
ggcatgttgc ccattgggtat gtatgggtgc cacatcccta ccatgccag tgcgggcatg	2580
ccacactccc tgggtgcaca cactgccc atgggtatga gttacctctt ggaatcttct	2640
cctatctcct cctcagctca gctccctcca aaataccagc ttggatctac ctcatacttg	2700
ccgcacaaac tgcccaaagt ggagggtgat agttttctgg cagagcttcc tggaagcctg	2760
tctctctcct cggtgaacc tcagcccgcc tcacctcagc cggcggcagc tgcagccctc	2820
ctggatgaag cactgctcgc caagagtccc gccaacctct ccgaggccct ctgcgctgct	2880
aatgtggact tttccactt actgggcttt ctccgctca acctacccc atgtaaccca	2940
ccaggggcca caggaggctt ggttatgggc tactcccaag ctgaggcaca gcctttgctc	3000
accactttgc aggtcagcc tcaagattcc gctggagctg ggggacccct gaactttggg	3060
cctctgcact cctacctcc tgtcttcacc tctggcctga gtaccaccac cctgcctcgt	3120
ttccaccaag cattccagta g	3141

<210> SEQ ID NO 12

<211> LENGTH: 5759

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 12

gcgagggcga gtgtgcgcgc ggtgagtgtg cgcgagtgtc tgtgcgggat cgggggcggg	60
cgggtccgggc tctcctggcg gagccgcgcg cgcgcgcgcg gccgctcggg ccccggcgct	120
gctcgtcgcg cagcttcgga gcgcgcgcga ttttgtgaga ttacaaaaa tcctcgtcgg	180
gaagaagcgg ccgcgcgcgg ccgccactcg gcgcgcgcgc tcgcgcgcgc cgcgcgcctc	240
cgcaccgggc cgggcgggc cccgcgcgcg aactcgggg gcccggggac tgcggtattt	300
gccggggagg ggctgtgcgc tgctccccg gcccggggcg ccgcgggccc cgcagcccg	360
acggaccctg agccagcgcg gctgcggcgc ggagccggga cgcgggggga ggagagaggc	420
ggccggggcg gcctgtgctg cggaggcgcg gccgggagcg cggcggggga gctcagggct	480
gagactcacc ggagggagcg gcgcgagcgc cccgccatcg tcccggttta tagttttgcc	540

-continued

gctggactct	tccctccctt	ccccacccc	atcaggatga	tatgagactt	gaaagaagac	600
gatgcataca	ggaggagaga	cttcagcatg	caaaccttca	tctgtccggc	ttgcaccgtc	660
gttctcatto	catgctgctg	gccttcagat	ggctgcacag	atgccccact	cacaccagta	720
cagtgaaccgt	cgccagccga	gcataagtga	ccagcaggtg	tctgccttac	catattctga	780
ccagattcag	caacctctaa	ctaaccaggt	gatgcctgac	attgtcatgt	tacagaggcg	840
gatgccccaa	accttccgtg	atccagcaac	tgctcctctg	agaaaactct	ctgtggactt	900
gatcaaaaaca	tacaagcata	ttaatgaggt	ttactatgca	aaaaagaage	gaagacacca	960
acagggccag	ggggacgatt	ccagtcataa	gaaggagcgg	aaggtttaca	atgatggtta	1020
cgatgatgat	aactatgatt	atattgtaaa	aaacggggaa	aagtggatgg	atcggtatga	1080
aatcgactcc	ttaataggca	aagggttcatt	tggacaggtt	gtgaaagctt	atgacagagt	1140
ggagcaagaa	tgggtcgcca	ttaaaatcat	caagaacaag	aaagcgtttc	tgaatcaagc	1200
ccagatagaa	gtgcggctgc	ttgagctcat	gaacaaacac	gacactgaaa	tgaagtacta	1260
catagtgcac	ttgaaacgcc	actttatggt	tcgaaaccat	ctctgtttag	tgtttgaaat	1320
gctgtcctat	aatctctatg	atttgttgag	aaacaccaac	ttccgagggg	tctctttgaa	1380
cctaacacga	aagtttgcgc	aacagatgtg	cacagcattg	ctttttcttg	cgactccaga	1440
acttagtata	attcactgtg	acttaaagcc	tgagaacatc	cttctgtgta	accccaaacy	1500
gagtgcatac	aagattgttg	attttggcag	ctcttgtcag	ttggggcaga	ggatatacca	1560
gtatatccag	agtcgctttt	atcggtctcc	agaggtgcta	ctgggaatgc	cttatgacct	1620
tgctatcgac	atgtggtccc	ttggatgtat	cttggttgaa	atgcacactg	gagagcctct	1680
gttcagtggt	gccaatgagg	tcgatcagat	gaataaaata	gtggaagtct	tgggcatccc	1740
acctgctcat	attcttgacc	aagcacccga	agcaagaaag	ttctttgaga	agttgcccg	1800
tggcacttgg	agcttaaaga	agaccaaaga	tggaaaacgg	gagtacaaac	caccaggaac	1860
ccgtaaaact	cataatattc	ttggagtaga	aacaggagga	cctggcgggc	ggcgtgctgg	1920
ggaatcgggt	catactgtag	ctgactactt	gaagttcaaa	gacctcattt	taaggatgct	1980
tgattatgac	cccaaaactc	ggattcaacc	ttattatgcc	ctgcagcaca	gttttttcaa	2040
gaaaacagct	gatgaaggta	ccaacacaag	taacagtgtg	tctaccagcc	ctgcgatgga	2100
gcagtcctag	tcttcaggca	ccacctccag	cacctcctcc	agctcagggtg	gatectcggg	2160
aacgagtaac	agtgaggagag	ccaggtcggg	tccgacgcac	cagcatcgac	acagcgggtg	2220
acacttcgct	gctgctgtcc	aggccatgga	ctgtgagaca	cacagtcccc	aggtgcgcca	2280
gcagtttccg	gctcctctgg	gatggtcagg	cactgaagct	cctacacaag	tcactgttga	2340
aactcatcct	gttcaagaga	caacctttca	tgtagccccc	cagcagaacg	cattgcatca	2400
tcaccatgga	aacagttccc	atcaccacca	ccaccaccat	caccaccacc	accaccatgg	2460
acagcaagcc	ttgggtaacc	ggaccaggcc	aagggtctac	aattctccaa	caaatagctc	2520
ctctacccag	gattctatgg	aggttggcca	cagtcaccac	tccatgacat	ccctgtcttc	2580
ctcaacaact	tcttctctga	catcttcttc	ctctactggt	aatcaaggca	atcaggccta	2640
tcagaaccgc	ccagtggtg	ctaacacctt	ggacttttga	cagaatggag	ctatggacgt	2700
taatttgacc	gtctactcca	atccccgcca	agagactggc	atagctggac	atccaacata	2760
ccaattttct	gctaatacag	gtcctgcaca	ttacatgact	gaaggacatc	tggcgatgag	2820

-continued

acaaggggct gatagagaag agtctcccat gacaggagtt tgtgtgcaac agagtccctgt	2880
agctagctcg tgactacatt gaaacttgag tttgtttctt gtgtgttttt atagaagtgg	2940
tgtttttttc ccaaaaaaca aaagtgcaca gctgcttgaa tcagaaggag attaacacac	3000
tgaaccgcta caagagggca aagctgactt ttttttttaa cttgaaaaga ttgcaaaggg	3060
acaatgaagt ttttaagagc catgtccacc catcttcacg gagagctcag agtgtcctct	3120
ttctgcctcc cccatttaac ttccatctcc tgggcacagt gggttttgtc tttctatcca	3180
acaagagtta atgttcagct gttggtcttg gtcatttgcc aactaatttt aaaataaaaa	3240
ggcactgcac ataatttgca taaagggccc cataagggtg ttttgtttg ttttcatttt	3300
tattttttcc ctttttcctt tgttttgttt tgtttcattt tgggctgggg gtgggggtggg	3360
aaatttgggt ttttatgtc ctctacacac ttgggcacgg aaatgcagta ctgtaaggaa	3420
gagggacctc tagcacacac aaacatcatc ttcagctgtg tgaaaggagc gattgtgttg	3480
acatttgtaa ggcacattaa gcatgctaag tggcggggat caggactctc ctatctgagc	3540
ctactgagga gcaaagcagc cattgcatgg gatcctgtgt gtctcctgct ggagaggcca	3600
caggaggata ggatggaacg tgactggctg gtctccttac caaggcgac tgagaagcaa	3660
tcagcgggtc ggtgtggcc agtcctgggg aggtctgagt ggtggctcctt gggataacct	3720
ttggccttac ggatttgac tcggagttag aagagccac catttcagat gcaatcactt	3780
gtggacatgc gtttgacagc agtccttaat gctgaaaaca cagagaatgg gtaattcaag	3840
aggcctttct tttaaaatag acttttgta ccactaatt gtaaggatt gcaaggtcac	3900
tttgctgtg tcataaagct gactcccttc ttggttgaag gccacagagg tagtggttg	3960
ctttgatgga aatagctaca gctgtgtccc ttctgcttt tcccttttct ttggcttttt	4020
ctcggcacgt ggtgtctcca ccatttcttc tgcacaaa ga cgtcttctgt tcacctgaa	4080
cattttaaaa tgcagaattt tatgtgactg cttttttgcc tcacaattat gctgtgaatt	4140
ttacaaaaat ttattttctt ttttgataat ttattgtacc aaagctgttt ttatagcaca	4200
tagatgtctg taaccaataa tgcgcaggc ctgcactttg acacagggtg aacgagaccg	4260
ttttaaacgt cagttgaaat gatggctgta ctattgctta aacaaaactg gaactgttgt	4320
tgaatccata gccaatatcat ttacagcaat cttgtactgg acagaataga ttgacatcta	4380
attcaagatt acaacacctg tcacattcta aatgtgttca ggcttctgaa gggaaaggga	4440
cactggatcc agaagctatg gagccagcag ttgatcttgt attcctgatt aacttccttg	4500
taaaacttgaa cttgagacct tgattgaacc gacaggcca cagtaggaac aggaagcagg	4560
cagagctctg agcatgacct gaagggaagca ggctggcact agcagttgcc ttgtctggag	4620
atgctgctgc cttcaagtag cacagtctgg cagtggctaa atctgtgttc ccattgaaat	4680
tgagcaattt taggggtgta tactctgaag cagttgaatt gggagaatga agatagagta	4740
atttcctgt ccaggatcaa aagaagccta gaaagaagca gtgatctacc tctgccaata	4800
accctgttgg aagcaacgca gcagaacacc acggtacttt ttattgctta attccacgtg	4860
gctgactagg tcaaggtttg tttggtgttt tttttttttt tttttgttcc tgagcaaaag	4920
cagggttttt tttgtttttt atatgtaaac aaaagaaagg cctaagctct gtgagtgcaa	4980
gtggagactc agagagcccc gtttctataa acacagacac ttcagggtgtt aatcaggaaa	5040
cccatattga ttttgtaatt aaatgtcaag cctgtggatt gttttgagct tggtagttga	5100

-continued

```

tgagcatttc cgtacagagg cctcaccttg agcacagcac tgtcaggagg ggcagcgcc 5160
tgtggctgcg gagccccagg tgccacacac cccccccag tggctcaggc cattcttttt 5220
ttctttctgt gctggttgcc acatttttagc aagcaccaaa aaagaaagca gttttttaag 5280
ctatatattac gtaaagagag ccacacaatc cagtgtcttc gcataccgtg tcacgttaca 5340
gtccagtttt gtgtgcttta ctacacagtt tgggttacagg acttctgtgc attgtaaaca 5400
taaacagcat ggagagggtta aatacctgtg ttcagattgt aagatctagt ccagacttgc 5460
tgtgtatatt gtaacgttaa atgaaaaaag accccccttt gtattatagt catgcggtct 5520
tatgtatgat aaacagttga ataatttgc ctcagagtct ttactatgct tttttaaaaa 5580
ttaacttaag gaaaaatgta aacagtaaaa aaacaatctt cgtgtgcagt gaagcgggtc 5640
cgggtgtggt aggtaaagca ggcacaatcc tgtgcctat gtaaaggga aagacttgtg 5700
agacagagac attggctgca agtgtctttg aaagtaaaaa ttggagggtt aacagataa 5759

```

```

<210> SEQ ID NO 13
<211> LENGTH: 337
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: shRNA

```

```

<400> SEQUENCE: 13
tagggataac agggtaattg tttgaatgag gcttcagtac tttacagaat cgttgccctgc 60
acatcttgga aacacttgct gggattactt cgacttctta acccaacaga aggctcgaga 120
aggtatatatg ctggtgacag tgagcgcgct gtgtatatatg taacgttata gtgaagccac 180
agatgtataa cgttacaata tacacagcat gcctactgcc tcggacttca aggggctaga 240
attcgagcaa ttatcttggt tactaaaaact gaataccttg ctatctcttt gatacatttt 300
tacaaaagctg aattaaaaatg gtataaatta aatcact 337

```

```

<210> SEQ ID NO 14
<211> LENGTH: 4357
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 14
tgcaattata tcttgcactg gacagacat cctgtgctgc ctcaaagcca agccttcccc 60
tctgtccag atgttccat ccatggtaac ccaacaaccc cttgtgctaa gagcttctgg 120
agcaagacct tcggaataga tcatttacag cagtgggtca ctaccatagc aggagggcgt 180
ttccctagtg ggcactgtga aggcccttca gttcctgagc tcagacatgg ctgacatcca 240
tccagagaac ctggtttcat tccacactgg ctattgtgct ctgatgccag cctcctggtt 300
tgggcctggc taggtcttag gttgacccta ggaggaagca cggggcctta tctcctccct 360
cagccccctg agagtattgg ggaatatagta caggttccag atggagggac gctcctacat 420
ggccatcccc taggtgtcta ggggttgact agtaaggta cctgggttagg aaccacggtt 480
atcagttctt acaaagaggc ccacccacac tggcacagca ttcatatggg ggcttatttt 540
taggtctccc accttctacc tgccaacaca aaatgctgaa tgaaactgga acacacacac 600
agaggcatgc acgcgcgcgc ccgcacgagc tgagggggat gtcattgagag gagttaaggg 660
cagccatggt ttcctaaaag ccacgcggga gtagtgatca cggacagcat gaaggaggca 720

```

-continued

cggtcctcct	gtttcagaga	tgcatgtggc	acctgggtgac	acaagagcag	tccagatacc	780
ctgtgggttc	aggggttcct	gagggcttcc	taggtatggg	gctttgtgct	aggcattatc	840
caagcaaat	tcagctagcc	tagaagtacg	cttggcagcg	tccgcgcgtg	gccatttcta	900
gctgcggaga	agtcactcct	ctcaactgac	agaactggca	ataaagcgcc	taaattctaa	960
atacatgagt	ccctctctca	tctgttacct	gcgaatggca	tggcggcagc	atctgtttga	1020
gagaatgtgg	cccactttaa	cccactgtgc	ctgtgggcaa	caggtcccag	ctggggaggg	1080
tgcccactcc	tctggaagtc	gctgtctgtt	cattgtaaaa	ctctcatttc	tctgcgcctg	1140
tggcctctgc	cagccagcgg	tggggagcct	ctggggagtg	ggagggaaga	aggagagaaa	1200
ggggggggga	gagcactcct	tctgcccctc	cccacctccc	cgcggcctgg	gaaaaggcag	1260
catattgagg	cgcggggctc	tcagcatcag	gccccgggat	gctaattgagg	gcgagcgcgt	1320
tcccacagcc	cggacattgt	gccgcgcggc	ccacctgctc	ctcggggagc	gaccattgtg	1380
ccgcgcgcaa	ttcacaggca	atttagcgtg	cgctaattgg	ccggcgcctt	tgtgcggccg	1440
gcgcgcgcat	tggccgcgga	gtgtgggaac	ggccgcggcg	ccgggacccc	aggcgcgggg	1500
ccgctgcccc	cgcctatata	gggctggcgt	gctggggctc	aggctgcgcc	agtccgggac	1560
gcctggctct	gtccgctccg	ctccgctccg	ctcgctaate	gcctccagag	ctccaggcct	1620
taagaggcag	gcaggcttaa	ttaagccacc	atgaccacat	ttttaccag	cgtccctccc	1680
tggattcaag	atgcaaagca	ggaggaggaa	gtgggctgga	aactagtccc	caggcctcgg	1740
ggccgggagg	cagagagtca	agtgaagtgc	caatgtgaaa	tctcggggac	accattttca	1800
aatggggaga	agctaaggcc	tcacagcctt	ccccatccgg	agcagagacc	atatagctgt	1860
cctcagctgc	actgcggcaa	ggcttttgcc	tccaagtaca	agctgtatag	gcacatggcc	1920
accactcgg	cccagaaaac	ccaccagtgc	atgtactgtg	acaagatgtt	tcaccgaaag	1980
gaccatctgc	ggaaccactt	gcagacccat	gaccccaaca	aagaggccct	ccactgctct	2040
gagtgtggta	agaattataa	tacaaagctg	ggataaccgt	gtcacctggc	catgcatgct	2100
gccagcagtg	gtgatctcag	ctgcaagggt	tgcctgcaga	cctttgagag	taccagggcc	2160
ctattggagc	acctgaaggc	tcaactcacg	cgggtagcag	gaggtgctaa	ggagaagaag	2220
caccctctgt	accattgtga	caggcgggtc	tatactcgga	aggatgtccg	acggcaccta	2280
gtggtgcaca	caggctgtaa	ggaactcctg	tgtcagtact	gtgccagag	atttgccgcg	2340
aaggatcacc	tgacacgcca	tgtcaagaag	agccactcac	aggaactgct	caagatcaag	2400
acagagccag	tggatatgtt	aggcttctc	agctgtagct	ctacagtcag	tgtaaaggaa	2460
gagctgagcc	ctgtgctttg	catggcctct	cgggacgtga	tgggggctaa	ggcctttcct	2520
ggcatgttgc	ccatgggtat	gtatggtgcc	cacatcccta	ccatgccag	tgcgggcatg	2580
ccacactccc	tgggtgcaca	cacactgccc	atgggtatga	gttaccctct	ggaatcttct	2640
cctatctcct	cctcagctca	gctccctcca	aaataccagc	ttgatctac	ctcatacttg	2700
cccagacaaac	tgccccaaagt	ggagggtgat	agttttctgg	cagagcttcc	tggaaagcctg	2760
tctctctcct	cggctgaacc	tcagcccggc	tcacctcagc	cggcggcagc	tgcagccctc	2820

-continued

ctggatgaag cactgctcgc caagagtccc gccaaacctct ccgaggccct ctgcgctgct	2880
aatgtggact tttcccactt actgggcttt cttccgctca acctaccccc atgtaacca	2940
ccaggggccca caggaggctt ggttatgggc tactcccaag ctgaggcaca gcctttgctc	3000
accactttgc aggtcagcc tcaagattcc gctggagctg ggggacctt gaactttggg	3060
cctctgcact ccttacctcc tgtcttcacc tctggcctga gtaccaccac cctgcctcgt	3120
ttccaccaag cattccaggg aagcggagct actaacttca gcctgctgaa gcaggctgga	3180
gacgtggagg agaaccttg accctgtgagc aaggcgaggg aggataacat ggccatcatc	3240
aaggagttca tgcgcttcaa ggtgcacatg gagggctccg tgaacggcca cgagttcgag	3300
atcgaggggc agggcgaggg ccgcccttac gagggcacc ccagccgcaa gctgaagggtg	3360
accaagggtg gcccctgcc cttgccttg gacatcctgt cccctcagtt catgtacggc	3420
tccaaggcct acgtgaagca ccccgctgac atccccgact acttgaagct gtccttcccc	3480
gagggcttca agtgggagcg cgtgatgaac ttcgaggacg gcggcgtggt gaccgtgacc	3540
caggactcct cctgcagga cggcgagttc atctacaagg tgaagctgcg cggcaccaac	3600
ttcccctccg acggccccgt aatgcagaag aagaccatgg gctgggaggg ctccctccgag	3660
cggatgtacc ccgaggacgg cgccctgaag ggcgagatca agcagaggct gaagctgaag	3720
gacggcggcc actacgagc tgaggtaag accacctaca aggccaaaga gcccgctgag	3780
ctgcccgcg cctacaacgt caacatcaag ttggacatca cctcccacaa cgaggactac	3840
accatcgtgg aacagtacga acgcgccgag ggccgccact ccaccggcgg catggacgag	3900
ctgtacaagg atatccctcc aaagaagaag agaaagggtg aaggcgcgcc tagggataac	3960
agggtaattg tttgaatgag gcttcagtag ttacagaat cgttgccctgc acatcttgga	4020
aacacttgct gggattactt cgacttctta acccaacaga aggctcgaga aggtatattg	4080
ctgttgacag tgagcgcgct gtgtatatgt taacgttata gtgaagccac agatgtataa	4140
cgttacaata tacacagcat gcctactgcc tcggacttca aggggctaga attcgagcaa	4200
ttatcttggt tactaaaact gaataccttg ctatctcttt gatacatctt tacaaagctg	4260
aattaaaatg gtataaatta aatcactcta gtaataaagg atcctttatc ttcattggat	4320
ccgtgtgttg gttttttgtg tgcggcccgt ctagacc	4357

1. A polynucleotide comprising:
 - (A) a nucleic acid sequence of a Plagl2 gene;
 - (B) a nucleic acid sequence of an miR-shRNA (microRNA adapted short hairpin RNA) against a Dyrk1a gene; and
 - (C) a promoter sequence operably linked to the above nucleic acid sequences.
2. The polynucleotide according to claim 1, wherein the Plagl2 gene and the Dyrk1a gene are mammalian genes.
3. The polynucleotide according to claim 1, wherein the promoter is selected from the group consisting of a Hes5 promoter, a GFAP promoter, a Sox2 promoter, and an Lfng promoter.
4. A vector comprising the polynucleotide according to claim 1.
5. The vector according to claim 4, wherein the vector is a lentivirus vector, an adeno-associated virus vector, an adenovirus vector, or a plasmid vector.
6. A pharmaceutical composition comprising the vector according to claim 4.
7. A pharmaceutical composition comprising:
 - (a) a vector comprising a nucleic acid sequence of a Plagl2 gene and a promoter sequence operably linked to the nucleic acid sequence of the Plagl2 gene; and
 - (b) a vector comprising a nucleic acid sequence of an miR-shRNA (microRNA adapted short hairpin RNA) against a Dyrk1a gene and a promoter sequence operably linked to the nucleic acid sequence of the miR-shRNA, or an siRNA or an miRNA against the Dyrk1a gene.
8. Cells transformed with the vector according to claim 4.
9. A pharmaceutical composition comprising the cells according to claim 8.
10. The pharmaceutical composition according to claim 6, wherein the pharmaceutical composition is used for treatment of a neurodegenerative disease or an inner ear disease.

11. The pharmaceutical composition according to claim 10, wherein the pharmaceutical composition is administered to a subject by intracerebroventricular injection, intrathecal bolus injection or infusion, intraganglionic injection, intraneural injection, subcutaneous injection, or intratympanic injection.

12. A method for screening for a substance for treatment of a neurodegenerative disease or an inner ear disease, comprising:

- (1) a step of causing a test substance to act on neural stem cells or inner ear supporting cells to measure expression of a *Plagl2* gene and a *Dyrk1a* gene; and
- (2-1) a step of selecting a test substance having an ability to enhance *Plagl2* gene expression and an ability to suppress *Dyrk1a* gene expression from a result of the step (1) or
- (2-2) a step of selecting a test substance having an ability to enhance *Plagl2* gene expression, and a test substance having an ability to suppress *Dyrk1a* gene expression from the result of the step (1).

* * * * *