



US 20250282871A1

(19) **United States**(12) **Patent Application Publication**
WANG et al.(10) **Pub. No.: US 2025/0282871 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **PHARMACEUTICAL COMBINATION AND
USE THEREOF****Publication Classification**(71) Applicant: **TIANJIN LIPOGEN TECHNOLOGY
CO., LTD**, Tianjin (CN)(72) Inventors: **Chenguang WANG**, Tianjin (CN);
Xiaojiao ZHANG, Tianjin (CN);
Xiaomeng DANG, Tianjin (CN); **Jin
ZHANG**, Tianjin (CN); **Yuting FU**,
Tianjin (CN); **Yihan WANG**, Tianjin
(CN); **Chao WANG**, Tianjin (CN)(51) **Int. Cl.****C07K 16/28** (2006.01)
A61K 31/7084 (2006.01)
A61K 38/21 (2006.01)
A61K 39/00 (2006.01)
A61K 45/06 (2006.01)
A61P 35/00 (2006.01)(52) **U.S. Cl.****CPC** **C07K 16/2818** (2013.01); **A61K 31/7084**
(2013.01); **A61K 38/212** (2013.01); **A61K**
38/215 (2013.01); **A61K 45/06** (2013.01);
A61P 35/00 (2018.01); **C07K 16/2827**
(2013.01); **A61K 2039/505** (2013.01); **A61K**
2039/545 (2013.01)(21) Appl. No.: **18/563,364**(22) PCT Filed: **May 20, 2022**(86) PCT No.: **PCT/CN2022/094047**

§ 371 (c)(1),

(2) Date: **Nov. 21, 2023**(30) **Foreign Application Priority Data**

May 21, 2021 (CN) 202110555249.9

(57)

ABSTRACT

A pharmaceutical combination and a use thereof. The pharmaceutical combination comprises a PD-1 inhibitor and/or a PD-L1 immune checkpoint inhibitor; and a STING pathway agonist. The pharmaceutical combination


Specification includes a Sequence Listing.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6





Figure 7



	TGI (%)	Q value
Vel.	0	
RB0005	43.7	
cGAMP -1	70.82	
cGAMP -3.16	79.63	
cGAMP -10	91.07	
RB0005-1 in combination with CGAMP-1	90.03	1.07>1, Synergism
RB0005-1 in combination with CGAMP-1	98.85	1.11>1, Synergism
RB0005-1 in combination with CGAMP-10	96.65	1.02>1, Synergism

Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28




Figure 29




Figure 30




Figure 31




Figure 32




Figure 33




Figure 34




Figure 35




Figure 36




Figure 37




Figure 38




Figure 39




Figure 40




Figure 41




Figure 42





Figure 43



PHARMACEUTICAL COMBINATION AND USE THEREOF

CROSS-REFERENCES TO RELATED APPLICATION

[0001] The present application is a National Stage of International Patent Application No: PCT/CN2022/094047 filed on May 20, 2022, which claims the benefit of the priority of the Chinese patent application with the application No. 202110555249.9, filed to the China National Intellectual Property Administration on May 21, 2021, the entire content of which is incorporated in this application by reference.

SEQUENCE LISTING

[0002] The present application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy is named `Sequence_Listing.txt` and is 33,420 bytes in size, and contains SEQ ID NO: 1 to SEQ ID NO: 44 which are identical to the sequence listing filed in the corresponding international application No. PCT/CN2022/094047 filed on May 20, 2022.

TECHNICAL FIELD

[0003] The disclosure relates to the field of biomedicine, specifically to development and application of a pharmaceutical combination.

BACKGROUND

[0004] Over the past decade, targeting PD-1 and PD-L1 has opened up a new era for clinical treatment of cancers. However, for many types of tumors, an immune checkpoint inhibitor monotherapy has poor efficacy. Even for immune responsive tumors, sustained clinical benefits have not been achieved in most patients. The vast majority of patients develop primary or acquired therapeutic drug resistance. In most cases, immunotherapy drug resistance can be attributed to the presence of immunosuppressive TME and insufficient activation of T cell anti-tumor effects due to a decrease in the number of immune cells in vivo. Therefore, compared to monotherapy, it is particularly important to actively seek new therapeutic strategies capable of reducing immune suppression in tumor environments or enhancing the response of cytotoxic cells to tumors.

[0005] The PD-L1/PD-1 signaling pathway is a very important co-inhibitory signaling pathway in the immune response. Researches have shown that when PD-L1 binds to PD-1, it will complement protein-tyrosine-phosphatases SHP-1 and SHP-2 with SH2 domains. These two phosphatases can reduce the extent of phosphorylation of the immunoreceptor tyrosine-based activation motif (ITAM) of CD3 ζ strand, weaken the activation of ZAP-70, and inhibit TCR downstream signaling, thereby playing a co-inhibitory role in T cell activation, and the autoimmune injury due to excessive activation of effector T cells can be prevented through such negative regulatory effect.

[0006] The stimulator of interferon genes (STING) is an immunostimulatory small molecule target mainly distributed in immune-related tissue cells, such as high expression in thymus, spleen, and peripheral blood leukocytes. When tumor cells undergo necrosis, cGAMP can bind and activate STING in the endoplasmic reticulum. Activation of STING

leads to nuclear translocation of transcription factors, induces the expressions of interferons (IFNs) and cytokines, promotes aggregation and activation of T cells, and ultimately kills tumor cells. The STING pathway can also be activated by synthesized cyclic dinucleotides (CDNs), triggering innate immune responses.

[0007] The combination of PD-L1/PD-1 inhibitors and STING pathway agonists, on the one hand, relieves inhibitory signals, enhances activation of T cells, and promotes adaptive immune system responses; on the other hand, it induces the expressions of IFNs and cytokines, promotes aggregation of T cells, activates the innate immune system responses, and enhances the responses of cytotoxic cells to tumors, etc. through a dual approach, which thus has very high clinical prospects and application values.

[0008] Moreover, researches have shown that some viral infections are also closely related to the PD-L1/PD-1 signaling pathway. For example, in chronic HIV infections, PD-1 was found to be highly expressed on the surface of CD8+ T cells that specifically recognize HIV. The virus activates the PD-L1/PD-1 signaling pathway, through which the activity of CD8+ T cells that specifically recognize HIV is inhibited, and the secretion ability of cytokines and the proliferation ability of T cells themselves are greatly weakened, leading to adaptive immune dysfunctions. It can be seen that this therapy may also have considerable application values in the treatment of such diseases.

SUMMARY

[0009] The disclosure provides a pharmaceutical combination and a use thereof in antitumor drugs. The pharmaceutical combination mainly consists of two parts:

[0010] (1) an immune checkpoint inhibitor (e.g., PD-1/PD-L1); (2) a STING pathway agonist (e.g., cyclic dinucleotide cGAMP or derivatives thereof). The pharmaceutical combination of the disclosure can be used to prepare efficient and low-toxicity anti-tumor drugs.

[0011] In one aspect, the disclosure provides a pharmaceutical combination including a programmed cell death protein 1 (PD-1) inhibitor and/or a programmed death ligand 1 (PD-L1) inhibitor, and a STING pathway agonist.

[0012] In some embodiments, where the STING pathway agonist includes a cyclic dinucleotide.

[0013] In some embodiments, where the cyclic dinucleotide is selected from the group consisting of c-di-AMP, c-di-GMP, c-di-GMP-F, 3',3'-cGAMP, 3',3'-cGAMP-F, 2',3'-cGAMP, Rp/Sp (CL656), ADU-S100, ADU-S100 disodium, and derivatives or combinations thereof.

[0014] In some embodiments, where the STING pathway agonist includes 2',3'-cGAMP, or derivatives thereof.

[0015] In some embodiments, where the STING pathway agonist includes flavonoids.

[0016] In some embodiments, where the flavonoid includes CMA, DMXAA, methoxyflavone, 6,4'-dimethoxyflavone, 4'-methoxyflavone, 3',6'-dihydroxyflavone, 7,2'-dihydroxyflavone, daidzein, Formononetin, retinene 7-methyl ether, xanthone, and/or any combination thereof.

[0017] In some embodiments, where the STING pathway agonist includes DNA.

[0018] In some embodiments, where the STING pathway agonist includes type I interferons (IFNs).

[0019] In some embodiments, where the type I interferon includes IFN- α or IFN- β .

[0020] In some embodiments, where the PD-1 inhibitor has one or more of the following characteristics:

[0021] a. inhibition or reduction of PD-1 expression, such as transcription or translation of PD-1;

[0022] b. inhibition or reduction of PD-1 activity, such as inhibition or reduction of PD-1 binding to its homologous ligands, such as PD-L1 or PD-L2; and

[0023] c. binding PD-1 or one or more of its ligands, such as PD-L1 or PD-L2.

[0024] In some embodiments, where the PD-1 inhibitor includes an anti PD-1 antibody or antigen-binding fragments thereof.

[0025] In some embodiments, where the anti PD-1 antibody is selected from the group consisting of Pembrolizumab, Nivolumab, Pidilizumab, SHR-1210, MEDI0680, BGB-A317, TSR-042, REGN2810, PF-06801591, RB0004, Tislelizumab, Camrelizumab, Toripalimab, Sintilimab, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof.

[0026] In some embodiments, where the anti PD-1 antibody includes at least one CDR in the antibody heavy chain variable region (VH), and the VH includes an amino acid sequence shown in SEQ ID NO: 8.

[0027] In some embodiments, where the anti PD-1 antibody includes a VH including HCDR3, and the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 3.

[0028] In some embodiments, where the VH further includes HCDR2, wherein the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 2.

[0029] In some embodiments, where the VH further includes HCDR1, wherein the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 1.

[0030] In some embodiments, where the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 3, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 1.

[0031] In some embodiments, where the VH includes a framework region HFR1, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, and the HFR1 includes an amino acid sequence shown in SEQ ID NO: 4 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 4.

[0032] In some embodiments, where the VH includes a framework region HFR2, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2; and the HFR2 includes an amino acid sequence shown in SEQ ID NO: 5 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 5.

[0033] In some embodiments, where the VH includes a framework region HFR3, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3; and the HFR3 includes an amino acid sequence shown in SEQ ID NO: 6 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 6.

[0034] In some embodiments, where the VH includes a framework region HFR4, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3, and the

HFR4 includes an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

[0035] In some embodiments, where the VH includes framework regions HFR1, HFR2, HFR3, and HFR4, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, the

[0036] N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3; among them, the HFR1 includes an amino acid sequence shown in SEQ ID NO: 4 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 4, the HFR2 includes an amino acid sequence shown in SEQ ID NO: 5 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 5, the HFR3 includes an amino acid sequence shown in SEQ ID NO: 6 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 6, and the HFR4 includes an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

[0037] In some embodiments, where the anti PD-1 antibody includes a VH, and the VH includes an amino acid sequence shown in SEQ ID NO: 8.

[0038] In some embodiments, where the anti PD-1 antibody includes an antibody heavy chain (HC), and the HC includes an amino acid sequence shown in SEQ ID NO: 9.

[0039] In some embodiments, where the anti PD-1 antibody includes at least one CDR in the antibody light chain variable region (VL), and the VL includes an amino acid sequence shown in SEQ ID NO: 17.

[0040] In some embodiments, where the anti PD-1 antibody includes at least one CDR in VH, wherein the VH includes an amino acid sequence shown in SEQ ID NO: 8, and the anti PD-1 antibody includes at least one CDR in VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 17.

[0041] In some embodiments, where the anti PD-1 antibody includes a VL including LCDR1, and the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 10.

[0042] In some embodiments, where the VL further includes LCDR2, wherein the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 11.

[0043] In some embodiments, where the VL further includes LCDR3, wherein the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 12.

[0044] In some embodiments, where the VL includes LCDR1, LCDR2 and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 10, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 12.

[0045] In some embodiments, where the anti PD-1 antibody includes VH and antibody VL, the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3

includes an amino acid sequence shown in SEQ ID NO: 3, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 1; and the VL includes LCDR1, LCDR2, and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 10, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 12.

[0046] In some embodiments, where the VL includes a framework region LFR1, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, and the LFR1 includes an amino acid sequence shown in SEQ ID NO: 13 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 13.

[0047] In some embodiments, where the VL includes a framework region LFR2, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2; and the LFR2 includes an amino acid sequence shown in SEQ ID NO: 14 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 14.

[0048] In some embodiments, where the VL includes a framework region LFR3, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3; and the LFR3 includes an amino acid sequence shown in SEQ ID NO: 15 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 15.

[0049] In some embodiments, where the VL includes a framework region LFR4, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 16 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 16.

[0050] In some embodiments, where the VL includes framework regions LFR1, LFR2, LFR3 and LFR4, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3; among them, the LFR1 includes an amino acid sequence shown in SEQ ID NO: 13 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 13, the LFR2 includes an amino acid sequence shown in SEQ ID NO: 14 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 14, the LFR3 includes an amino acid sequence shown in SEQ ID NO: 15 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 15, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 16 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 16.

[0051] In some embodiments, where the anti PD-1 antibody includes a VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 17.

[0052] In some embodiments, where the anti PD-1 antibody includes VH and VL, the VH includes an amino acid sequence shown in SEQ ID NO: 8, and the VL includes an amino acid sequence shown in SEQ ID NO: 17.

[0053] In some embodiments, where the anti PD-1 antibody includes an antibody light chain (LC), and the LC includes an amino acid sequence shown in SEQ ID NO: 18.

[0054] In some embodiments, where the anti PD-1 antibody includes HC and LC, the HC includes an amino acid sequence shown in SEQ ID NO: 9, and the LC includes an amino acid sequence shown in SEQ ID NO: 18.

[0055] In some embodiments, where the PD-L1 inhibitor has one or more of the following characteristics:

[0056] a. inhibition or reduction of PD-L1 expression, such as transcription or translation of PD-L1;

[0057] b. inhibition or reduction of PD-L1 activity, such as inhibition or reduction of PD-L1 binding to its associated receptors such as PD-1; and

[0058] c. binding of PD-L1 or its receptors such as PD-1.

[0059] In some embodiments, where the PD-L1 inhibitor includes an anti PD-L1 antibody or antigen-binding fragments thereof.

[0060] In some embodiments, where the anti PD-L1 antibody is selected from the group consisting of Durvalumab, Atezolizumab, Avelumab, Envipolimab, MDX-1105, YW243.55.S70, MDPL3280A, AMP-224, LY3300054, RB0005, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof.

[0061] In some embodiments, where the anti PD-L1 antibody includes at least one CDR in VH, the VH includes an amino acid sequence shown in SEQ ID NO: 25.

[0062] In some embodiments, where the anti PD-L1 antibody includes a VH, and the VH includes HCDR3, and the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21.

[0063] In some embodiments, where the VH further includes HCDR2, wherein the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20.

[0064] In some embodiments, where the VH further includes HCDR1, wherein the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19.

[0065] In some embodiments, where the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19.

[0066] In some embodiments, where the VH includes a framework region HFR1, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, and the HFR1 includes an amino acid sequence shown in SEQ ID NO: 22 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 22.

[0067] In some embodiments, where the VH includes a framework region HFR2, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2; and the HFR2 includes an amino acid sequence shown in SEQ ID NO: 23 or an amino acid

sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 23.

[0068] In some embodiments, where the VH includes a framework region HFR3, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3; and the HFR3 includes an amino acid sequence shown in SEQ ID NO: 24 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 24.

[0069] In some embodiments, where the VH includes a framework region HFR4, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3, and the HFR4 includes an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

[0070] In some embodiments, where the VH includes framework regions HFR1, HFR2, HFR3, and HFR4, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3; among them, the HFR1 includes an amino acid sequence shown in SEQ ID NO: 22 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 22, the HFR2 includes an amino acid sequence shown in SEQ ID NO: 23 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 23, the HFR3 includes an amino acid sequence shown in SEQ ID NO: 24 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 24, and the HFR4 includes an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

[0071] In some embodiments, where the anti PD-L1 antibody includes VH, and the VH includes an amino acid sequence shown in SEQ ID NO: 25.

[0072] In some embodiments, where the anti PD-L1 antibody includes HC, and the HC includes an amino acid sequence shown in SEQ ID NO: 26.

[0073] In some embodiments, where the anti PD-L1 antibody includes at least one CDR in VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 37.

[0074] In some embodiments, where the anti PD-L1 antibody includes at least one CDR in VH, the VH includes an amino acid sequence shown in SEQ ID NO: 25, and the anti PD-L1 antibody includes at least one CDR in VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 37.

[0075] In some embodiments, where the anti PD-L1 antibody includes at least one CDR in VH, the VH includes an amino acid sequence shown in SEQ ID NO: 25, and the anti PD-L1 antibody includes at least one CDR in VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 38, SEQ ID NO: 39 or SEQ ID NO: 40.

[0076] In some embodiments, where the anti PD-L1 antibody includes a VL including LCDR1, and the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 27.

[0077] In some embodiments, where the anti PD-L1 antibody includes a VL including LCDR1, and the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 28, SEQ ID NO: 29 or SEQ ID NO: 30.

[0078] In some embodiments, where the VL further includes LCDR2, wherein the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31.

[0079] In some embodiments, where the VL further includes LCDR3, wherein the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32.

[0080] In some embodiments, where the VL includes LCDR1, LCDR2 and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 27, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32.

[0081] In some embodiments, where the VL includes LCDR1, LCDR2 and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 28, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32;

[0082] the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 29, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32; or the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 30, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32.

[0083] In some embodiments, where the anti PD-L1 antibody includes VH and antibody VL, the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19; and the VL includes LCDR1, LCDR2, and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 27, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32.

[0084] In some embodiments, where the anti PD-L1 antibody includes VH and antibody VL, the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19; and the VL includes LCDR1, LCDR2, and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 28, SEQ ID NO: 29 or SEQ ID NO: 30, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32.

[0085] In some embodiments, where the VL includes a framework region LFR1, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, and the LFR1 includes an amino acid sequence shown in SEQ ID

NO: 33 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 33.

[0086] In some embodiments, where the VL includes a framework region LFR2, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2; and the LFR2 includes an amino acid sequence shown in SEQ ID NO: 34 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 34.

[0087] In some embodiments, where the VL includes a framework region LFR3, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3; and the LFR3 includes an amino acid sequence shown in SEQ ID NO: 35 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 35.

[0088] In some embodiments, where the VL includes a framework region LFR4, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 36 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 36.

[0089] In some embodiments, where the VL includes framework regions LFR1, LFR2, LFR3 and LFR4, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3; among them, the LFR1 includes an amino acid sequence shown in SEQ ID NO: 33 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 33, the LFR2 includes an amino acid sequence shown in SEQ ID NO: 34 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 34, the LFR3 includes an amino acid sequence shown in SEQ ID NO: 35 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 35, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 36 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 36.

[0090] In some embodiments, where the anti PD-L1 antibody includes a VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 37.

[0091] In some embodiments, where the anti PD-L1 antibody includes a VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 38, SEQ ID NO: 39 or SEQ ID NO: 40.

[0092] In some embodiments, where the anti PD-L1 antibody includes VH and VL, the VH includes an amino acid sequence shown in SEQ ID NO: 25, and the VL includes an amino acid sequence shown in SEQ ID NO: 37.

[0093] In some embodiments, where the anti PD-L1 antibody includes VH and VL, the VH includes an amino acid

sequence shown in SEQ ID NO: 25, and the VL includes an amino acid sequence shown in SEQ ID NO: 38, SEQ ID NO: 39 or SEQ ID NO: 40.

[0094] In some embodiments, where the anti PD-L1 antibody includes LC, and the LC includes an amino acid sequence shown in SEQ ID NO: 41.

[0095] In some embodiments, where the anti PD-L1 antibody includes LC, and the LC includes an amino acid sequence shown in SEQ ID NO: 42, SEQ ID NO: 43 or SEQ ID NO: 44.

[0096] In some embodiments, where the anti PD-L1 antibody includes HC and LC, the HC includes an amino acid sequence shown in SEQ ID NO: 26, and the LC includes an amino acid sequence shown in SEQ ID NO: 41.

[0097] In some embodiments, where the anti PD-L1 antibody includes HC and LC, the HC includes an amino acid sequence shown in SEQ ID NO: 26, and the LC includes an amino acid sequence shown in SEQ ID NO: 42, SEQ ID NO: 43 or SEQ ID NO: 44.

[0098] In some embodiments, i) the PD-1 inhibitor and/or PD-L1 inhibitor in the pharmaceutical combination are not mixed with ii) the STING pathway agonist to each other in the pharmaceutical combination.

[0099] In some embodiments, i) the PD-1 inhibitor and/or PD-L1 inhibitor, and ii) the STING pathway agonist are present in the pharmaceutical combination in a single dosage form.

[0100] In some embodiments, where the pharmaceutical combination is formulated into a pharmaceutical composition.

[0101] In some embodiments, where the pharmaceutical composition includes a PD-1 inhibitor or a PD-L1 inhibitor, and a STING pathway agonist.

[0102] In some embodiments, where the STING pathway agonist is present in an amount of about 0.0001 mg/kg to about 200 mg/kg.

[0103] In some embodiments, where the PD-1 inhibitor or PD-L1 inhibitor is present in an amount of 0.0001 mg/kg to about 200 mg/kg.

[0104] In some embodiments, where the pharmaceutical composition further includes one or more pharmaceutically acceptable carriers.

[0105] In another aspect, the disclosure also provides a use of the aforementioned pharmaceutical combination in the preparation of drugs for the treatment of neoplastic diseases.

[0106] In some embodiments, where the neoplastic diseases include tumors and/or wart diseases.

[0107] In another aspect, the disclosure also provides the aforementioned pharmaceutical combination for use in the treatment of neoplastic diseases.

[0108] In another aspect, the disclosure also provides a drug including the aforementioned pharmaceutical combination for treating neoplastic diseases.

[0109] In another aspect, the disclosure also provides a method for treating neoplastic diseases, which includes administering an effective amount of the aforementioned pharmaceutical combination to a subject in need thereof.

[0110] In some embodiments, where the subject suffers from neoplasm.

[0111] In some embodiments, where the neoplasm includes tumors and/or warts.

[0112] In some embodiments, where the administration includes local, intraneoplastic (e.g., in tumors or warts), or systemic administrations.

[0113] In some embodiments, where the administration includes intravenous injection, intravenous instillation, intramuscular injection, subcutaneous injection, and/or intraneoplastic injection.

[0114] In some embodiments, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered by use of the same or different administration routes.

[0115] In some embodiments, it includes injection of the STING pathway agonist into the neoplasm.

[0116] In some embodiments, it also includes injection or systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm.

[0117] In some embodiments, it includes injection of the STING pathway agonist and systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm.

[0118] In some embodiments, it includes injection of i) the PD-1 inhibitor or PD-L1 inhibitor STING pathway agonist, and ii) the STING pathway agonist in the pharmaceutical combination into the neoplasm.

[0119] In some embodiments, where i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered simultaneously or at different times.

[0120] In some embodiments, where the PD-1 inhibitor or PD-L1 inhibitor is administered before and/or after the administration of the STING pathway agonist.

[0121] In some embodiments, where the PD-1 inhibitor or PD-L1 inhibitor is administered after the administration of the STING pathway agonist.

[0122] In some embodiments, the method includes: i) injection of the STING pathway agonist into the neoplasm; ii) injection or systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm after administering the STING pathway agonist.

[0123] In some embodiments, where i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered simultaneously.

[0124] In some embodiments, where i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are simultaneously administered by way of intraneoplastic injection, and the PD-1 inhibitor or PD-L1 inhibitor and the STING pathway agonist are present in a same dosage form.

[0125] In another aspect, the disclosure provides a pharmaceutical kit including the aforementioned pharmaceutical combination.

[0126] Those skilled in the art can have an insight into other aspects and advantages of the disclosure from the detailed description below. The detailed description below only shows and describes exemplary embodiments of the disclosure. As those skilled in the art will recognize, the content of the disclosure enables those skilled in the art to make modifications to the disclosed specific embodiments without departing from the spirit and scope of the invention involved in the disclosure.

[0127] Correspondingly, the accompanying drawings and the descriptions in the specification of the disclosure are only illustrative, rather than being restrictive.

BRIEF DESCRIPTION OF THE DRAWINGS

[0128] The specific features of the invention involved in the disclosure are shown in the attached claims. The char-

acteristics and advantages of the invention involved in the disclosure can be better understood by referring to the exemplary embodiments and accompanying drawings described in detail hereinafter. A brief explanation of the attached drawings is as follows:

[0129] FIG. 1 shows the in vivo efficacy study of the pharmaceutical combination of PD-L1 inhibitor RB0005 and 2',3'-cGAMP according to the disclosure, as shown by changes in body weight of each group of mice during administration.

[0130] FIG. 2 shows the in vivo efficacy study of the pharmaceutical combination of PD-L1 inhibitor RB0005 and 2',3'-cGAMP according to the disclosure, as shown by changes in survival rate of each group of mice during administration.

[0131] FIG. 3A shows the in vivo efficacy study of the pharmaceutical combination of PD-L1 inhibitor RB0005 and 2',3'-cGAMP according to the disclosure, as shown by changes in tumor proliferation of each group of mice during administration.

[0132] FIG. 3B shows the in vivo efficacy study of the pharmaceutical combination of PD-L1 inhibitor RB0005 and 2',3'-cGAMP according to the disclosure, as shown by changes in tumor proliferation of each group of mice on day 17 after treatment.

[0133] FIG. 4 shows the in vivo efficacy study of the pharmaceutical combination of PD-L1 inhibitor RB0005 and 2',3'-cGAMP according to the disclosure, as shown by tumor weight inhibitory rate of mice obtained through dissection after the administration and observation have completed.

[0134] FIG. 5 shows the in vivo efficacy study of the pharmaceutical combination of PD-1 inhibitor RB0004 and 2',3'-cGAMP according to the disclosure, as shown by changes in tumor proliferation of each group of mice during administration.

[0135] FIG. 6 shows the in vivo efficacy study of the pharmaceutical combination of PD-1 inhibitor RB0004 and 2',3'-cGAMP according to the disclosure, as shown by tumor weight inhibitory rate of mice obtained through dissection after the administration and observation have completed.

[0136] FIG. 7 shows the in vivo efficacy study of the pharmaceutical combination of PD-L1 inhibitor RB0005 and different doses of 2',3'-cGAMP according to the disclosure, as shown by tumor weight inhibitory rate of mice obtained through dissection after the administration and observation have completed.

[0137] FIG. 8 shows the in vivo efficacy study of the pharmaceutical combination of PD-1 inhibitor RB0004 and different doses of 2',3'-cGAMP according to the disclosure, as shown by changes in tumor proliferation of each group of mice during administration.

[0138] FIG. 9 shows the in vivo efficacy study of the pharmaceutical combination of PD-1 inhibitor RB0004 and different doses of 2',3'-cGAMP according to the disclosure, as shown by changes in survival rate of each group of mice during administration.

[0139] FIG. 10 shows the trend of tumor proliferation in each group of mice during administration in Example 5 of the disclosure.

[0140] FIG. 11 shows the survival curve of each group of mice during administration in Example 5 of the disclosure.

[0141] FIG. 12 shows the mean tumor mass of each group of mice during administration in Example 5 of the disclosure.

[0142] FIG. 13 shows the trend of tumor proliferation in each group of mice during administration in Example 6 of the disclosure.

[0143] FIG. 14 shows the survival curve of each group of mice during administration in Example 6 of the disclosure.

[0144] FIG. 15 shows the mean tumor mass of each group of mice during administration in Example 6 of the disclosure.

[0145] FIG. 16 shows the trend of tumor proliferation in each group of mice during administration in Example 7.2.1 of the disclosure.

[0146] FIG. 17 shows the survival curve of each group of mice during administration in Example 7.2.1 of the disclosure.

[0147] FIG. 18 shows the mean tumor mass of each group of mice during administration in Example 7.2.1 of the disclosure.

[0148] FIG. 19 shows the trend of tumor proliferation in each group of mice during administration in Example 7.2.2 of the disclosure.

[0149] FIG. 20 shows the survival curve of each group of mice during administration in Example 7.2.2 of the disclosure.

[0150] FIG. 21 shows the mean tumor mass of each group of mice during administration in Example 7.2.2 of the disclosure.

[0151] FIG. 22 shows the trend of tumor proliferation in each group of mice during administration in Example 7.2.3 of the disclosure.

[0152] FIG. 23 shows the survival curve of each group of mice during administration in Example 7.2.3 of the disclosure.

[0153] FIG. 24 shows the trend of changes in tumor volume of single mouse in the composite formulation group during administration in Example 7.2.3 of the disclosure.

[0154] FIG. 25 shows the mean tumor mass of each group of mice during administration in Example 7.2.3 of the disclosure.

[0155] FIG. 26 shows the relative tumor volume inhibitory rate of each group of mice during administration in Example 8 of the disclosure.

[0156] FIG. 27 shows the trend of tumor proliferation in each group of mice during administration in Example 9 of the disclosure.

[0157] FIG. 28 shows the survival curve of each group of mice during administration in Example 9 of the disclosure.

[0158] FIG. 29 shows the mean tumor mass of each group of mice during administration in Example 9 of the disclosure.

[0159] FIG. 30 shows the trend of tumor proliferation in each group of mice during administration in Example 10 of the disclosure.

[0160] FIG. 31 shows the mean tumor volume of each group of mice at the endpoint of the experiment in Example 10 of the disclosure.

[0161] FIG. 32 shows the survival curve of each group of mice during administration in Example 10 of the disclosure.

[0162] FIG. 33 shows the trend of changes in tumor volume of single mouse in G5 group during administration in Example 10 of the disclosure.

[0163] FIG. 34 shows the trend of tumor proliferation in each group of mice during administration in Example 11 of the disclosure.

[0164] FIG. 35 shows the survival curve of each group of mice during administration in Example 11 of the disclosure.

[0165] FIG. 36 shows the mean tumor mass of each group of mice during administration in Example 11 of the disclosure.

[0166] FIG. 37 shows the trend of tumor proliferation in each group of mice during administration in Example 12 of the disclosure.

[0167] FIG. 38 shows the survival curve of each group of mice during administration in Example 12 of the disclosure.

[0168] FIG. 39 shows the mean tumor volume of each group of mice on day 7 in Example 12 of the disclosure.

[0169] FIG. 40 shows the mean tumor mass of each group of mice during administration in Example 12 of the disclosure.

[0170] FIG. 41 shows the trend of tumor proliferation in each group of mice during administration in Example 13 of the disclosure.

[0171] FIG. 42 shows the survival curve of each group of mice during administration in Example 13 of the disclosure.

[0172] FIG. 43 shows the mean tumor mass of each group of mice during administration in Example 13 of the disclosure.

DETAILED DESCRIPTION OF THE EMBODIMENTS

[0173] The embodiments of the present invention will be illustrated below by way of specific examples, and other advantages and effects of the present disclosure can be easily understood by those familiar with this technology from the content disclosed in this specification.

Definitions of Terms

[0174] In the disclosure, the term “PD-1” generally refers to programmed cell death protein 1, i.e. a type I membrane protein of 288 amino acids first described in 1992 (Ishida et al., EMBO J., 11(1992), 3887-3895). PD-1 is a member of the expanded CD28/CTLA-4T cell regulator family and has two ligands, i.e. PD-L1 (B7-H1 and CD274) and PD-L2 (B7-DC and CD273). The structure of this protein includes an extracellular IgV domain, followed by a transmembrane region and an intracellular tail. The intracellular tail contains two phosphorylation sites located in the immunoreceptor tyrosine-based inhibitory motif and the immunoreceptor tyrosine-based switch motif, indicating a negative regulation of TCR signaling by PD-1. This is consistent with the binding of SHP-1 and SHP-2 phosphatases to the cytoplasmic tail of PD-1 after ligand binding. Although PD-1 is not expressed on naive T cells, it is upregulated after the T cell receptor (TCR) mediated activation and observed on both activated and depleted T cells (Agata et al., Int. Immunology 8 (1996), 765-772). These depleted T cells exhibit a dysfunctional phenotype and are unable to respond appropriately. Although PD-1 has a relatively broad expression pattern, its most important role is likely to act as a co-inhibitory receptor on T cells (Chinai et al., Trends in Pharmacological Sciences 36 (2015), 587-595). The current treatment approaches thus focus on blocking the interaction between PD-1 and its ligand to enhance T cell response. In this disclosure, the PD-1 may include human PD-1 (hPD-1),

or variants, isotypes, and species homologs thereof, as well as analogues having at least one common epitope with hPD-1. The amino acid sequence of exemplary hPD-1 can be found under GenBank accession No.: U64863.

[0175] In this disclosure, the term “PD-L1” generally refers to programmed cell death 1 ligand 1, which can also be referred to as B7 homolog 1, B7-H1, differentiation cluster 274, (3) 274, or CD274, which, when combined with PD-1, downregulates T cell activation and cytokine secretion. “PD-L1” includes any natural PD-L1 from any vertebrate sources, including mammals such as primates (e.g., humans and *Macaca fascicularis*) and rodents (e.g., mice and rats). The term contemplates “full length”, unprocessed PD-L1, and any form of PD-L1 produced by cell processing. PD-L1 can exist as a transmembrane protein or as a soluble protein. “PD-L1” includes the full PD-L1 and fragments thereof, as well as functional variants, isoforms, species homologs, derivatives, analogues of PD-L1, and analogues having at least one common epitope with PD-L1. The basic structure of PD-L1 includes four domains, i.e. extracellular Ig like V-type domain and Ig like C2-type domain, transmembrane domain, and cytoplasmic domain. An exemplary human PD-L1 amino acid sequence can be found under the NCBI accession No.: NP_001254653 or UniProt accession No.: Q9NZQ7.

[0176] In this disclosure, the term “inhibitor” generally refers to a compound/substance or composition that can completely or partially prevent or reduce the physiological functions of one or more specific biomolecules (e.g., proteins (such as PD-1 or PD-L1), polypeptides, lipopolysaccharides, glycoproteins, ribonucleoprotein complexes, etc.). The reduce of physiological functions of one or more specific proteins can include a decrease in the activity of the protein itself (e.g., its ability to bind to other molecules, etc.) or a decrease in the existence quantity of the protein itself. Suitable inhibitor molecules can include antagonist antibodies or antibody fragments, small molecular fragments or derivatives, peptides, antisense oligonucleotides, small organic molecules, etc. In some embodiments, the inhibitor can block the activation of cellular signaling pathways. In some embodiments, the PD-1/PD-L1 inhibitor is an anti PD-1/PD-L1 antibody or antigen-binding fragments thereof.

[0177] In this disclosure, the terms “Pembrolizumab”, “Nivolumab”, “Pidilizumab”, “SHR-1210”, “MEDI0680”, “BGB-A317”, “TSR-042”, “REGN2810”, “PF-06801591”, “Durvalumab”, “Atezolizumab”, “Avelumab”, “Envafohimab”, “MDX-1105”, “YW243.55.S70”, “MDPL3280A”, “AMP-224”, “LY3300054”, “RB0004”, “RB0005”, “Tislelizumab”, “Camrelizumab”, “Toripalimab”, “Sintilimab” are used according to their general and ordinary meanings understood in the art.

[0178] In this disclosure, the term “STING (stimulator of interferon genes; also known as THEM173, MITA, ERIS, and MPYS)” generally refers to the endoplasmic reticulum (ER) transmembrane proteins expressed in the thymus, spleen, placenta, and THP1 human monocytes. Human STING is encoded by the gene TMEM173. The STING pathway can be activated by exogenous cyclic dinucleotides (CDNs) produced by bacterial infections or structurally different endogenous CDNs (e.g., cyclic GMP-AMP (cGAMP) produced by cyclic GMP-AMP Synthase (cGAS) in response to sensing the cytoplasmic double stranded DNAs (dsDNAs)) (Ablaser et al., 2013; Diner et al., 2013). The cytoplasmic domain of STING forms dimers, and

CDNs are bound at the interface of dimers (Burdette, D. L. et al., 2011). After the binding of ligands, the cytoplasmic tail of STING acts as an adapter for TBK-1 and IRF-3, causing their phosphorylation. Phosphorylated IRF-3 enters the nucleus to induce transcription of genes encoding type I IFNs and cytokines used to promote intercellular host immune defenses (Keating et al., 2011). The activation of STING can lead to the production of type I interferons (e.g., IFN- α and IFN- β) through the IRF3 (Interferon Regulatory Factor 3) pathway and the production of proinflammatory cytokines (IL-1 α , IL-1 β , IL-2, IL-6, and TNF- α , etc.) through the oncogenic transcription factor NF- κ B (activated B nuclear factor K-light chain enhancer) pathway. The STING pathway can regulate the innate immune recognition of immunogenic tumors and promote the anti-tumor effect of interferon. IFN- γ exerts the anti-tumor efficiency through TRAIL (tumor necrosis factor-related apoptosis-inducing ligand) in vivo, promoting apoptosis of tumor cells.

[0179] In this disclosure, the term “STING pathway agonist” generally refers to a molecule that binds to STING (stimulator of interferon genes, or TMEM173), activate STING, and triggers the activation of IRF3-TBK1 pathway, leading to increased transcription of type I interferon and other genes.

[0180] In this disclosure, the term “cyclic dinucleotide (CDN)” generally refers to a class of cyclic molecules having two phosphodiester bonds or two thiophosphate diester bonds, or one phosphodiester diester bond and one thiophosphate diester bond, between two nucleosides. Exemplary cyclic dinucleotides include but are not limited to 3'-5'-c-di-AMP (c-di-AMP), 3'-5'-c-di-GMP (c-di-GMP), c-di-GMP-F, 3',3'-cGAMP (also known as 3'-5',3'-5'-cGAMP, a product of *Vibrio cholerae* DncV protein), 3',3'-cGAMP-F, 2',3'-cGAMP (also known as 2'-5',3'-5'-cGAMP, a product of human cGAS protein), Rp/Sp (CL656), ADU-S100, ADU-S100 disodium and a combination thereof.

[0181] In this disclosure, the term “nucleoside” generally refers to a glycosylamine containing a nitrogen containing base and a pentose, wherein the nitrogen containing base binds to pentose through a B-glucosidic bond. The term “nucleotide” generally refers to any nucleoside linked to a phosphate group at positions 5', 3' or 2' in the sugar moiety.

[0182] In this disclosure, the term “antibody” generally refers to one being used in the broadest sense and specifically covering monoclonal antibodies, polyclonal antibodies, dimers, multimers, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments, as long as they exhibit the desired biological activities (Miller et al. (2003) Jour. of Immunology 170:4854-4861). Antibodies can be mouse, human, humanized, chimeric antibodies, or those derived from other species.

[0183] A full-length antibody typically refers to an antibody composed of two “full-length antibody heavy chains” and two “full-length antibody light chains”. The “full length antibody heavy chain” is generally such a polypeptide that is composed of the antibody heavy chain variable domain (VH), antibody constant heavy chain domain 1 (CH1), antibody hinge region (HR), antibody heavy chain constant domain 2 (CH2), and antibody heavy chain constant domain 3 (CH3) in the direction of N-terminal to C-terminal, abbreviated as VH-CH1-HR-CH2-CH3; and in the case of IgE subtype antibodies, optionally further including the antibody heavy chain constant domain 4 (CH4). In some embodiments, the “full length antibody heavy chain” is a polypep-

tide composed of VH, CH1, HR, CH2, and CH3 in the direction of N-terminal to C-terminal. The “full length antibody light chain” is generally a polypeptide composed of an antibody light chain variable domain (VL) and an antibody light chain constant domain (CL) in the direction of N-terminal to C-terminal, abbreviated as VL-CL. The antibody light chain constant domain (CL) can be κ (kappa) or λ (lambda). Two full-length antibody chains are linked together by an interpolypeptide disulfide bond between the CL domain and the CH1 domain, and an interpolypeptide disulfide bond between the hinge regions of the full-length antibody heavy chain. Typical examples of full-length antibodies are natural antibodies such as IgG (e.g., IgG1 and IgG2), IgM, IgA, IgD, and IgE).

[0184] In this disclosure, the term “antigen binding fragment” generally refers to a portion of the antibody molecule that contains amino acids responsible for the specific binding between the antibody and the antigen. The part of an antigen that is specifically recognized and bound by an antibody is called as an “epitope” as mentioned above. The antigen binding domain can typically include the antibody light chain variable region (VL) and the antibody heavy chain variable region (VH); however, it does not necessarily include both. Fd fragments for example have two VH regions, and generally retain some antigen-binding functions of the whole antigen-binding domain. Examples of antigen binding fragments of an antibody include (1) Fab fragments, monomeric fragments having VL, VH, constant light chain (CL), and CH1 domains; (2) F (ab')₂ fragments, dimeric fragments having two Fab fragments linked by a disulfide bridge of the hinge region; (3) Fd fragments having two VH and CH1 domains; (4) Fv fragments having single-armed VL and VH domains of antibodies, (5) dAb fragments (Ward et al., “Binding Activities of a Repertoire of Single Immunoglobulin Variable Domains Secreted From *Escherichia coli*,” *Nature* 341:544-546 (1989), incorporated by reference into the present disclosure in its entirety), having VH domains; (6) isolated complementary determining region (CDR); (7) single chain Fv (scFv), for example, derived from the scFV library. Although the two domains VL and VH of the Fv fragment are encoded by independent genes, they can be conjugated using a recombination method by a synthetic linker, the synthetic linker enables it to be made into a single protein chain (called as single chain Fv (scFv)) in which the VL and VH regions pair to form a monomeric molecule (see, for example, Huston et al., “Protein Engineering of Antibody Binding Sites: Recovery of Specific Activity in an Anti-Digoxin Single-Chain Fv Analogue Produced in *Escherichia coli*,” *Proc. Natl. Acad. Sci. USA* 85:5879-5883 (1988)); and (8) VHH, “VHH” involves variable antigen-binding domains of heavy chain antibodies from the Camelidae (camels, dromedaries, llamas, and alpacas, etc.) (see Nguyen V. K. et al., 2000, *The EMBO Journal*, 19, 921-930; Muyldermans S., 2001, *J Biotechnol.*, 74, 277-302; and Review Vanlandschoot P. et al., 2011, *Antiviral Research* 92, 389-407). VHH can also be referred to as nanobody (Nb) and/or single domain antibody. These antibody fragments can be obtained using conventional techniques known to those skilled in the art, and their functionalities can be evaluated in the same way as intact antibodies.

[0185] In this disclosure, the terms “variable region” or “variable domain” generally refer to regions where there may be significant differences in sequence of certain seg-

ments of the variable domains between antibodies. The “variable region” in a light chain can include the light chain variable region VL; and the “variable region” in a heavy chain can include the heavy chain variable region VH. Variable domains mediate antigen binding and determine the specificity of specific antibodies against their specific antigens. However, variability does not mean the uniform distribution throughout the entire scope of variable domains. It is generally concentrated in three segments called as hyper-variable regions (CDR or HVR) in the light chain and heavy chain variable domains. The more highly conserved part of the variable domains is called as the framework region (FR). Each of the variable domains of natural heavy and light chains contains four FR regions, most of which adopt B-sheet configuration and are connected by three CDRs, forming a circular connection, and in some cases forming a part of a B-sheet structure. The CDRs in each chain are closely held together through the FR region, and promote the formation of antigen-binding sites of antibodies together with the CDRs from another chain (see Kabat et al, *Sequences of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, Md. (1991)). The terms “VH” and “VH domain” can be interchangeably used to refer to the heavy chain variable region of an antibody or antigen-binding molecule thereof.

[0186] In this disclosure, the term “CDR” generally refers to the complementary determining region within the antibody variable sequence. There are three CDRs in each variable region of the heavy and light chains, and for each variable region, they are called as CDR1, CDR2, and CDR3, respectively. The precise boundaries of these CDRs have been defined differently based on different systems. The system described by Kabat (Kabat et al., *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987) and (1991)) not only provides a definite residue numbering system applicable to any variable region of the antibody, but also provides precise residue boundaries that define these three CDRs. These CDRs can be referred to as Kabat CDRs. Chothia and co-workers (Chothia&Lesk, *J. Mol. Biol.* 196:901-917 (1987) and Chothia et al., *Nature* 342:877-883 (1989)) found that some sub-moieties within Kabat CDR exhibit almost identical peptide skeleton conformations, despite significant differences at the amino acid sequence level. These sub-moieties are designated as L1, L2, and L3, or H1, H2, and H3, where “L” and “H” refer to the light and heavy chain regions, respectively. These regions can be called as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Padlan (FASEB J.9:133-139 (1995)) and MacCallum (JMol Biol 262 (5): 732-45 (1996)) have already described other boundaries that define CDRs overlapping with Kabat CDRs. Other CDR boundaries may not strictly conform to one of the above systems, but still overlap with Kabat CDR. Although they can be shortened or extended based on predictions or experimental findings below, specific residues or groups of residues, or even the whole CDRs, do not significantly affect the antigen binding. Unless otherwise explicitly stated in the specification, as used in this disclosure, the terms “CDR”, “HCDR1”, “HCDR2”, “HCDR3”, “LCDR1”, “LCDR2”, and “LCDR3” include CDRs as defined in any of the aforementioned methods (Kabat, Chothia, or IMGT).

[0187] In this disclosure, the term “percentage (%) sequence identity” generally refers to the number of matches

(“hits”) of consistent amino acids when two or more aligned amino acid sequences are compared with the number of amino acid residues making up the total length of the amino acid sequences. In other words, when alignment is used for two or more sequences, the percentage of identical amino acid residues (e.g., 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% sequence identity) can be determined when comparing and aligning these sequences against the maximum correspondence (as measured using sequence comparison algorithms known in the art), or when manually aligning and visually inspecting. Therefore, sequences that are compared to determine sequence identity can be distinguished by one or more amino acid substitutions, additions, or deletions. The suitable programs for alignment of protein sequences are known to those skilled in the art. The percentage sequence identity of protein sequences can be determined, for example, using programs such as CLUSTALW, Cluster Omega, FASTA, or BLAST, such as the use of NCBI BLAST algorithm (Altschul S F et al. (1997), *Nuclear Acids Res.* 25:3389-3402).

[0188] In this disclosure, the terms “combination”, “therapeutic combination”, “combination therapy”, or “pharmaceutical combination” generally refer to a combination including at least two active ingredients/therapeutic agents, where a STING pathway agonist and a PD-1/PD-L1 inhibitor can be administered independently at the same time or separately within time intervals that allow the combinations to exhibit cooperation (e.g. synergistic) effects. In some embodiments, each active ingredient/therapeutic agent can be respectively prepared into an independent preparation (solid, liquid, and gel, etc.). In some embodiments, each active ingredient/therapeutic agent can be contained in different containers, and can also be prepared into the desired preparations concurrently or separately with an appropriate carrier as needed. In some embodiments, each active ingredient/therapeutic agent can be of different sources (e.g., prepared, produced by different manufacturers, or sold by different merchants). In some embodiments, each active ingredient/therapeutic agent can form pharmaceutical compositions in a mixed form.

[0189] In this disclosure, the term “pharmaceutical composition” generally refers to a preparation being in a form that allows the biological activity of active ingredients to be effective and containing no other components that have unacceptable toxicity to the subject for who the composition is going to be administered. Such composition can be sterile and can contain pharmaceutically acceptable carriers, such as physiological saline. A suitable pharmaceutical composition may include one or more buffer solutions (e.g., acetate, phosphate, or citrate buffer solutions), surfactants (e.g., polysorbate), stabilizers (e.g., human albumin), preservatives (e.g., benzyl alcohol), absorption enhancers for enhancing bioavailability, and/or other conventional solubilizers or dispersants. The pharmaceutical compositions of the disclosure include but are not limited to liquid, frozen, and lyophilized compositions.

[0190] In this disclosure, the term “pharmaceutically acceptable carrier” generally refers to one or more non-toxic materials that do not interfere with the biological activity of the active ingredients. Such preparation can conventionally contain a salt, a buffer, a preservative, a compatible carrier, and optionally other therapeutic agents. Such pharmaceutically acceptable preparation can also contain a compatible solid or liquid filler, a diluent, or an encapsulating substance

suitable for human administration. Other contemplated carriers, excipients, and/or additives that can be used in the formulations described herein may include, for example, a flavouring agent, an antimicrobial agent, a sweetener, an antioxidant, an antistatic agent, lipid, a protein excipient (e.g., serum albumin, gelatin, casein), a salt-forming counterion (e.g., sodium), and the like. These and other known pharmaceutical carriers, excipients, and/or additives suitable for use in the formulations described herein are known in the art.

[0191] In this disclosure, the term “neoplastic cells” generally refers to cells that undergo new and abnormal proliferation, especially those in which proliferation is uncontrollable and progresses, leading to neoplastic diseases. Neoplastic cells can be malignant, i.e., invasive and metastatic, or benign.

[0192] In this disclosure, the term “neoplasm” generally refers to an abnormal lump of tissues, where the growth of the lump exceeds that of normal tissues and is not coordinated with the growth of normal tissues. The “neoplasm” can be defined to be “benign” or “malignant”, depending on the following characteristics: degree of cell differentiation, including morphology and function, rate of growth, local invasion and metastasis. “Benign neoplasm” is generally well-differentiated, has slower growth characteristics than malignant neoplasm, and maintains localization to the origin site. In addition, benign tumors do not have the ability to infiltrate, invade, or metastasize to the distal sites. “Malignant neoplasm” is generally poorly differentiated (anaplastic) and has characteristic rapid growth, accompanied by progressive infiltration, invasion, and destruction of surrounding tissues. In addition, malignant neoplasm has the ability to metastasize to the distal sites.

[0193] In this disclosure, the term “tumor” or “cancer” generally refers to any medical disorders characterized by the growth, proliferation, or metastasis of neoplastic or malignant cells, and the tumor can be a solid or non-solid tumor.

[0194] In this disclosure, the term “wart” generally refers to a type of superficial benign neoplasm of the skin caused by human papilloma virus (HPV), mainly characterized by cell proliferative responses. HPV belongs to the Papovaviridae A genus in the DNA virus family. HPV is divided into more than 80 subtypes and is associated with different types of warts. The term “wart” generally refers to all types of warts, including but not limiting to verruca plantaris, verruca vulgaris, and genital warts, unless otherwise specified.

[0195] In this disclosure, the term “administer” and similar terms are generally not limited to bodily administration, and suitable methods include in vitro, in vitro followed by in vivo, or in vivo methods. For example, any administration method known to those skilled in the art for bringing cells, organs, or tissues into contact with the composition can be used. For example, the compound can be introduced into the body of a subject in need of treatment through any introduction or delivery routes. In some embodiments, the composition of the disclosure can be administered orally, locally, intranasally, intramuscularly, subcutaneously, intradermally, intrathecally, intraperitoneally, percutaneously, or intratumorally.

[0196] In this disclosure, the term “effective amount” or “effective dose” generally refers to an amount sufficient to achieve or at least partially achieve the desired effect. The “therapeutically effective amount” or “therapeutically effective

tive dose” of a drug or therapeutic agent is generally any amount of a drug that promotes regression of a disease when used alone or in combination with another therapeutic agent (as evidenced by a decrease in the severity of disease symptoms, an increase in the frequency and duration of asymptomatic periods of a disease, or the prevention of damage or disability caused by suffering from a disease).

[0197] In this disclosure, the term “treatment” generally refers to the slowing down or improvement of the progression, severity, and/or duration of proliferative disorders, or the improvement of one or more symptoms of proliferative disorders (e.g., one or more identifiable symptoms) due to the administration of one or more therapies (e.g., one or more therapeutic agents such as the pharmaceutical composition of this disclosure). In this disclosure, the term “treatment” may also refer to the improvement of at least one measurable physical parameter of proliferative disorders, such as tumor growth, not necessarily to be distinguishable by the patient. The term “treatment” in this disclosure can also refer to the inhibition of the progression of proliferative disorders through, for example, stabilizing the identifiable symptoms by physical means (for example, stabilizing physical parameters), or by physiological means, or both. In some cases, the term “treatment” can refer to reduction or stabilization of tumor size or cancer cell counts.

[0198] In this disclosure, the term “synergy” generally refers to the efficacy of the combination of two or more active drugs being greater than the sum of efficacies obtained by using the individual drugs alone. Therefore, when the combination of two or more pharmaceutical ingredients leads to a “synergistic inhibition” of an activity or process, such as tumor growth, it means that the inhibitory effect on the activity or process is greater than the sum of the inhibitory effects of the active drugs alone. In this disclosure, the synergistic effect calculation of pharmaceutical combinations can be evaluated using the coefficient of drug interaction (CDI/CI). CDI(CI) is calculated according to the following formula: $CDI(CI) = AB/A * B$. AB is the ratio of the combination group of two drugs to the control group, as calculated based on the anatomical tumor weight. A or B is the ratio of single drug group to control group. If CDI is less than 1, it indicates that the two drugs have a synergistic effect; If CDI equals to 1, the two drugs have an additive effect; If CDI is greater than 1, the two drugs have an antagonistic effect. (Pikman, Y. et al, Synergistic Drug Combinations with a CDK4/6 Inhibitor in T-cell Acute Lymphoblastic Leukemia, (2017) Clin Cancer Res 23, 1012-1024. PMID: 28151717 PMCID: PMC5432118 DOI: 10.1158/1078-0432.CCR-15-2869.)

[0199] In this disclosure, the term “subject” generally refers to human or non-human animals, including but not limiting to cats, dogs, horses, pigs, cows, sheep, rabbits, mice, rats, or monkeys, etc

[0200] In this disclosure, the term “about” generally refers to a variation within a range of (0.5% to 10%) above or below the specified value, such as a variation within a range of about 0.5%, about 1%, about 1.5%, about 2%, about 2.5%, about 3%, about 3.5%, about 4%, about 4.5%, about 5%, about 5.5%, about 6%, about 6.5%, about 7%, about 7.5%, about 8%, about 8.5%, about 9%, about 9.5%, or about 10% above or below the specified value.

[0201] In this disclosure, the term “comprising” and its variant forms including “containing”, “including” and other

forms, generally refer to the inclusion of other components, elements, numerical values, and steps, etc.

DETAILED DESCRIPTION OF INVENTION

[0202] This disclosure provides an agonist that induces the expression of INF and cytokines as well as activates innate immune system responses in combination with an immune checkpoint inhibitor for use as a pharmaceutical combination, including a STING agonist (e.g., cyclic dinucleotide 2'3'-cGAMP or derivatives thereof) and an immune checkpoint inhibitor (PD-1/PD-L1), etc.

[0203] In one aspect, the disclosure provides a pharmaceutical combination, which can include include a programmed cell death protein 1 (PD-1) inhibitor and/or a programmed death ligand 1 (PD-L1) inhibitor, and a STING pathway agonist.

[0204] In some embodiments, where the STING pathway agonist includes a cyclic dinucleotide.

[0205] In some embodiments, where the cyclic dinucleotide is selected from the group consisting of c-di-AMP, c-di-GMP, c-di-GMP-F, 3',3'-cGAMP, 3',3'-cGAMP-F, 2',3'-cGAMP, Rp/Sp (CL656), ADU-S100, ADU-S100 disodium, and derivatives or combinations thereof.

[0206] In some embodiments, where the STING pathway agonist includes 2',3'-cGAMP, or derivatives thereof.

[0207] In some embodiments, where the STING pathway agonist includes flavonoids.

[0208] In some embodiments, where the flavonoid includes CMA, DMXAA, methoxyflavone, 6,4'-dimethoxyflavone, 4'-methoxyflavone, 3',6'-dihydroxyflavone, 7,2'-dihydroxyflavone, daidzein, Formononetin, retinene 7-methyl ether, xanthone, and/or any combination thereof.

[0209] In some embodiments, where the STING pathway agonist includes DNA.

[0210] In some embodiments, where the STING pathway agonist includes type I interferons (IFNs).

[0211] In some embodiments, where the type I interferon includes IFN- α or IFN- β .

[0212] In some embodiments, where the PD-1 inhibitor has one or more of the following characteristics:

[0213] a. inhibition or reduction of PD-1 expression, such as transcription or translation of PD-1;

[0214] b. inhibition or reduction of PD-1 activity, such as inhibition or reduction of PD-1 binding to its homologous ligands, such as PD-L1 or PD-L2; and

[0215] c. binding PD-1 or one or more of its ligands, such as PD-L1 or PD-L2.

[0216] In some embodiments, where the PD-1 inhibitor includes an anti PD-1 antibody or antigen-binding fragments thereof.

[0217] For example, the pharmaceutical combination can include: 1) an anti PD-1 antibody or antigen-binding fragments thereof; and 2) a cyclic dinucleotide.

[0218] In some embodiments, where the anti PD-1 antibody is selected from the group consisting of Pembrolizumab, Nivolumab, Pidilizumab, SHR-1210 (Incyte/Jiangsu Hengrui Pharmaceuticals Co., Ltd.), MEDI0680 (also known as AMP-514; Amplimmune Inc./Medimmune), BGB-A317 (BeiGene Ltd.), TSR-042 (also known as ANB011; AnaptysBio/Tesaro, nc.), REGN2810 (Regeneron Pharmaceuticals, Inc./Sanofi-Aventis), PF-06801591 (Pfizer), RB0004, Tislelizumab, Camrelizumab, Toripalimab, Sintilimab, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof.

[0219] For example, the pharmaceutical combination can include: 1) an anti PD-1 antibody, which can be selected from the group consisting of Pembrolizumab, Nivolumab, Pidilizumab, SHR-1210 (Incyte/Jiangsu Hengrui Pharmaceuticals Co., Ltd.), MEDI0680 (also known as AMP-514; Amplimmune Inc./Medimmune), BGB-A317 (BeiGene Ltd.), TSR-042 (also known as ANB011; AnaptysBio/Tesaro, nc.), REGN2810 (Regeneron Pharmaceuticals, Inc./Sanofi-Aventis), PF-06801591 (Pfizer), RB0004, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof; and 2) a cyclic dinucleotide, which can be selected from the group consisting of c-di-AMP, c-di-GMP, c-di-GMP-F, 3',3'-cGAMP, 3',3'-cGAMP-F, 2',3'-cGAMP, Rp/Sp (CL656), ADU-S100, ADU-S100 disodium or combinations thereof.

[0220] In some embodiments, the anti PD-1 antibody is RB0004. RB0004 and other humanized anti PD-1 monoclonal antibodies are disclosed in CN201610345750.1 and WO2017201766A1. For example, the pharmaceutical combination includes: 1) the anti PD-1 antibody, which is RB0004 or bio-enhancers, and bio-equivalents, as well as combinations thereof; and 2) a STING pathway agonist including 2',3'-cGAMP or derivatives thereof.

[0221] In another embodiments, the antibody PD-1 is Pembrolizumab (trade name Keytruda, previously Lambrizumab, also known as Merck 3745, MK-3475, or SCH-900475), a humanized IgG4 monoclonal antibody that binds to PD-1. Pembrolizumab is disclosed in, for example Hamid et al. (2013) New England Journal of Medicine 369 (2): 134-44, WV02009/114335 and U.S. Pat. No. 8,354,509. For example, the pharmaceutical combination includes: 1) the anti PD-1 antibody, which is Pembrolizumab or bio-enhancers, and bio-equivalents, or combinations thereof; and 2) a STING pathway agonist including 2',3'-cGAMP or derivatives thereof.

[0222] In another embodiments, the anti PD-1 antibody is Nivolumab (CAS Registry No.: 946414-94-4, alternative names include MDX-1106, MDX-1106-04, ONO-4538, or BMS-936558). Nivolumab is a complete human IgG4 monoclonal antibody that specifically blocks PD-1. Nivolumab (Clone 5C4) and other human monoclonal antibodies specifically binding to PD-1 have been disclosed in U.S. Pat. No. 8,008,449 and WO2006/121168. For example, the pharmaceutical combination includes: 1) the anti PD-1 antibody, which is Nivolumab or bio-enhancers, and bio-equivalents, as well as combinations thereof; and 2) a STING pathway agonist including 2',3'-cGAMP or derivatives thereof.

[0223] In another embodiments, the anti PD-1 antibody is Pidilizumab. Pidilizumab (CT-011; Cure Tech) is a humanized IgG1 monoclonal antibody that binds to PD-1. Pidilizumab and other humanized anti PD-1 monoclonal antibodies are disclosed in WO2009/101611. Other anti Rpd-1 antibodies have been disclosed in U.S. Pat. No. 8,609,089, US2010028330, and/or US20120114649. Other anti PD-1 antibodies include AMP514 (Amplimmune). For example, the pharmaceutical combination includes: 1) the anti PD-1 antibody, which is Pidilizumab or bio-enhancers, and bio-equivalents, as well as combinations thereof; and 2) a STING pathway agonist including 2',3'-cGAMP or derivatives thereof.

[0224] In some embodiments, where the anti PD-1 antibody includes at least one CDR in the antibody heavy chain

variable region (VH), and the VH includes an amino acid sequence shown in SEQ ID NO: 8.

[0225] In some embodiments, where the anti PD-1 antibody includes VH including HCDR3, the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 3 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 3.

[0226] In some embodiments, where the VH further includes HCDR2, wherein the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 2 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 2.

[0227] In some embodiments, where the VH further includes HCDR1, wherein the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 1 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 1.

[0228] In some embodiments, where the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 3 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 3, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 2 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 1 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 1.

[0229] In some embodiments, where the VH includes a framework region HFR1, wherein the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, and the HFR1 includes an amino acid sequence shown in SEQ ID NO: 4 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 4.

[0230] In some embodiments, where the VH includes a framework region HFR2, wherein the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2; and the HFR2 includes an amino acid sequence shown in SEQ ID NO: 5 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about

90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 8.

[0235] In some embodiments, where the anti PD-1 antibody includes an antibody heavy chain (HC), the HC includes an amino acid sequence shown in SEQ ID NO: 9 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 9.

[0236] In some embodiments, where the anti PD-1 antibody includes at least one CDR in the antibody light chain variable region (VL), and the VL includes an amino acid sequence shown in SEQ ID NO: 17.

[0237] In some embodiments, where the anti PD-1 antibody includes at least one CDR in VH, wherein the VH includes an amino acid sequence shown in SEQ ID NO: 8, and the anti PD-1 antibody includes at least one CDR in VL, and the VL includes an amino acid sequence shown in SEQ ID NO: 17.

[0238] In some embodiments, where the anti PD-1 antibody includes VL including LCDR1, the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 10 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 10.

[0239] In some embodiments, where the VL further includes LCDR2, wherein the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 11 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 11.

[0240] In some embodiments, where the VL further includes LCDR3, wherein the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 12 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 12.

[0241] In some embodiments, where the VL includes LCDR1, LCDR2 and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 10 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 10, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 11 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 12 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about

96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 12.

[0242] In some embodiments, where the anti PD-1 antibody includes VH and an antibody VL, the VH includes HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 3 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 3, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 2 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 1 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 1; and the VL includes LCDR1, LCDR2, and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO: 10 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 10, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 11 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 12 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 12.

[0243] For example, the anti PD-1 antibody can include VH and antibody VL, the VH can include HCDR1, HCDR2, and HCDR3, wherein the HCDR3 can include an amino acid sequence shown in SEQ ID NO: 3, the HCDR2 can include an amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 can include an amino acid sequence shown in SEQ ID NO: 1; and the VL can include LCDR1, LCDR2, and LCDR3, wherein the LCDR1 can include an amino acid sequence shown in SEQ ID NO: 10, the LCDR2 can include an amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 can include an amino acid sequence shown in SEQ ID NO: 12.

[0244] For example, the pharmaceutical combination can include: 1) anti PD-1 antibody, the anti PD-1 antibody can include VH and antibody VL, the VH can include HCDR1, HCDR2, and HCDR3, wherein the HCDR3 can include an amino acid sequence shown in SEQ ID NO: 3, the HCDR2 can include an amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 can include an amino acid sequence shown in SEQ ID NO: 1; and the VL can include LCDR1,

LCDR2, and LCDR3, wherein the LCDR1 can include an amino acid sequence shown in SEQ ID NO: 10, the LCDR2 can include an amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 can include an amino acid sequence shown in SEQ ID NO: 12; 2) a STING pathway agonist, which can be 2',3'-cGAMP, or derivatives thereof.

[0245] In some embodiments, where the VL includes a framework region LFR1, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, and the LFR1 includes an amino acid sequence shown in SEQ ID NO: 13 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 13.

[0246] In some embodiments, where the VL includes a framework region LFR2, wherein the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2; and the LFR2 includes an amino acid sequence shown in SEQ ID NO: 14 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 14.

[0247] In some embodiments, where the VL includes a framework region LFR3, wherein the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3; and the LFR3 includes an amino acid sequence shown in SEQ ID NO: 15 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 15.

[0248] In some embodiments, where the VL includes a framework region LFR4, wherein the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 16 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 16.

[0249] In some embodiments, where the VL includes framework regions LFR1, LFR2, LFR3 and LFR4, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3; among them, the LFR1 includes an amino acid sequence shown in SEQ ID NO: 13 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence

identity to the amino acid sequence shown in SEQ ID NO: 13, the LFR2 includes an amino acid sequence shown in SEQ ID NO: 14 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 14, the LFR3 includes an amino acid sequence shown in SEQ ID NO: 15 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 15, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 16 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 16.

[0250] In some embodiments, where the anti PD-1 antibody includes VL, the VL includes an amino acid sequence shown in SEQ ID NO: 17 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 17.

[0251] In some embodiments, where the anti PD-1 antibody includes VH and VL, the VH includes an amino acid sequence shown in SEQ ID NO: 8 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 8, and the VL includes an amino acid sequence shown in SEQ ID NO: 17 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 17.

[0252] For example, where the anti PD-1 antibody can include VH and VL, the VH can include an amino acid sequence shown in SEQ ID NO: 8, and the VL can include an amino acid sequence shown in SEQ ID NO: 17.

[0253] For example, the pharmaceutical combination can include: 1) anti PD-1 antibody, wherein the anti PD-1 antibody can include VH and VL, the VH can include an amino acid sequence shown in SEQ ID NO: 8, and the VL can include an amino acid sequence shown in SEQ ID NO: 17; 2) a STING pathway agonist, which can be 2',3'-cGAMP or derivatives thereof.

[0254] In some embodiments, where the anti PD-1 antibody includes an antibody light chain (LC), the LC includes an amino acid sequence shown in SEQ ID NO: 18 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 18.

[0255] In some embodiments, where the anti PD-1 antibody includes HC and LC, the HC includes an amino acid sequence shown in SEQ ID NO: 9 or an amino acid

sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 9, and the LC includes an amino acid sequence shown in SEQ ID NO: 18 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 18.

[0256] For example, where the anti PD-1 antibody can include HC and LC, the HC can include an amino acid sequence shown in SEQ ID NO: 9, and the LC can include an amino acid sequence shown in SEQ ID NO: 18.

[0257] For example, the pharmaceutical combination can include: 1) anti PD-1 antibody, wherein the anti PD-1 antibody can include HC and LC, the HC can include an amino acid sequence shown in SEQ ID NO: 9, and the LC can include an amino acid sequence shown in SEQ ID NO: 18; 2) a STING pathway agonist, which can be 2',3'-cGAMP or derivatives thereof.

[0258] In some embodiments, where the PD-L1 inhibitor has one or more of the following characteristics:

[0259] a. inhibition or reduction of PD-L1 expression, such as transcription or translation of PD-L1;

[0260] b. inhibition or reduction of PD-L1 activity, such as inhibition or reduction of PD-L1 binding to its associated receptors such as PD-1; and

[0261] c. binding of PD-L1 or its receptors such as PD-1.

[0262] In some embodiments, where the PD-L1 inhibitor includes an anti PD-L1 antibody or antigen-binding fragments thereof.

[0263] For example, the pharmaceutical combination can include: 1) an anti PD-L1 antibody or antigen-binding fragments thereof; and 2) a cyclic dinucleotide.

[0264] In some embodiments, where the anti PD-L1 antibody is selected from the group consisting of Durvalumab (MEDI4736, disclosed in US2013/0034559A1), Atezolizumab (MPDL3280A, disclosed in U.S. Pat. No. 8,217,149), Avelumab (MSB0010718C, disclosed in US2014/0341917A1), MDX-1105, YW243.55.S70, MDPL3280A, AMP-224 (Amplimune, GlaxoSmithKline), LY3300054 (Eli Lilly and Co.), RB0005, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof.

[0265] For example, the pharmaceutical combination can include: 1) an anti PD-L1 antibody, wherein the anti PD-L1 antibody is selected from the group consisting of Durvalumab (MEDI4736, disclosed in US2013/0034559A1), Atezolizumab (MPDL3280A, disclosed in U.S. Pat. No. 8,217,149), Avelumab (MSB0010718C, disclosed in US2014/0341917A1), MDX-1105, YW243.55.S70, MDPL3280A, AMP-224, LY3300054, RB0005, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof; and 2) a cyclic dinucleotide, which is selected from the group consisting of c-di-AMP, c-di-GMP, c-di-GMP-F, 3',3'-cGAMP, 3',3'-cGAMP-F, 2',3'-cGAMP, Rp/Sp (CL656), ADU-S100, ADU-S100 disodium, or combinations thereof.

[0266] In some embodiments, the anti PD-L1 antibody is RB0005. RB0005 and other humanized anti PD-L1 monoclonal antibodies have been disclosed in CN201610340678.3 and WO2017197667A1. For example,

the pharmaceutical combination includes: 1) an anti PD-L1 antibody, which is RB0005, or bio-enhancers, bio-equivalents, or combinations thereof; and 2) a STING pathway agonist including 2',3'-cGAMP or derivatives thereof.

[0267] In other embodiments, the PD-L1 inhibitor is MDX-1105. MDX-1105, also known as BMS-936559, is an anti PD-L1 antibody described in WO 2007/005874. For example, the pharmaceutical combination includes: 1) an anti PD-L1 antibody, which is MDX-1105, or bio-enhancers, bio-equivalents, or combinations thereof; and 2) a STING pathway agonist including 2',3'-cGAMP or derivatives thereof.

[0268] In other embodiments, the PD-L1 inhibitor is YW243.55.S70. The YW243.55.S70 antibody is an anti PD-L1 antibody described in WO 2010/077634.

[0269] In other embodiments, the PD-L1 inhibitor is MDPL3280A (Genentech/Roche). MDPL3280A is a human Fc optimized IgG1 monoclonal antibody that binds to PD-L1. Other human monoclonal antibodies of MDPL3280A and PD-L1 have been disclosed in U.S. Pat. No. 7,943,743 and US patent publication number 20120039906.

[0270] In some embodiments, where the anti PD-L1 antibody includes at least one CDR in VH, the VH includes an amino acid sequence shown in SEQ ID NO: 25.

[0271] In some embodiments, where the anti PD-L1 antibody includes VH including HCDR3, the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 21.

[0272] In some embodiments, where the VH further includes HCDR2, wherein the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 20.

[0273] In some embodiments, where the VH further includes HCDR1, wherein the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 19.

[0274] In some embodiments, where the VH includes HCDR1, HCDR2 and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 21, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 includes an amino acid sequence shown in SEQ ID

NO: 19 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 19.

[0275] For example, where the VH can include HCDR1, HCDR2, and HCDR3, wherein the HCDR3 includes an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 includes an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19.

[0276] In some embodiments, where the VH includes a framework region HFR1, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, and the HFR1 includes an amino acid sequence shown in SEQ ID NO: 22 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 22.

[0277] In some embodiments, where the VH includes a framework region HFR2, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2; and the HFR2 includes an amino acid sequence shown in SEQ ID NO: 23 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 23.

[0278] In some embodiments, where the VH includes a framework region HFR3, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3; and the HFR3 includes an amino acid sequence shown in SEQ ID NO: 24 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 24.

[0279] In some embodiments, where the VH includes a framework region HFR4, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3, and the HFR4 includes an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

[0280] In some embodiments, where the VH includes framework regions HFR1, HFR2, HFR3, and HFR4, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3; among

HCDR1 includes an amino acid sequence shown in SEQ ID NO: 19 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 19; and the VL includes LCDR1, LCDR2, and LCDR3, wherein the LCDR1 includes an amino acid sequence shown in SEQ ID NO:28, SEQ ID NO: 29 or SEQ ID NO:30 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO:28, SEQ ID NO:29 or SEQ ID NO:30, the LCDR2 includes an amino acid sequence shown in SEQ ID NO: 31 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 includes an amino acid sequence shown in SEQ ID NO: 32 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 32.

[0298] For example, where the anti PD-L1 antibody can include VH and an antibody VL, the VH can include HCDR1, HCDR2, and HCDR3, wherein the HCDR3 can include an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 can include an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 can include an amino acid sequence shown in SEQ ID NO: 19; and the VL include LCDR1, LCDR2, and LCDR3, wherein the LCDR1 can include an amino acid sequence shown in SEQ ID NO:28, SEQ ID NO:29 or SEQ ID NO:30, the LCDR2 can include an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 can include an amino acid sequence shown in SEQ ID NO: 32.

[0299] For example, the pharmaceutical combination can include: 1) anti PD-L1 antibody, wherein the anti PD-L1 antibody can include VH and antibody VL, the VH can include HCDR1, HCDR2, and HCDR3, wherein the HCDR3 can include an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 can include an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 can include an amino acid sequence shown in SEQ ID NO: 19; and the VL can include LCDR1, LCDR2, and LCDR3, wherein the LCDR1 can include an amino acid sequence shown in SEQ ID NO:28, SEQ ID NO: 29 or SEQ ID NO:30, the LCDR2 can include an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 can include an amino acid sequence shown in SEQ ID NO: 32; 2) a STING pathway agonist, which can be 2',3'-cGAMP, or derivatives thereof.

[0300] In some embodiments, where the VL includes a framework region LFR1, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, and the LFR1 includes an amino acid sequence shown in SEQ ID NO: 33 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about

96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 33.

[0301] In some embodiments, where the VL includes a framework region LFR2, wherein the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2; and the LFR2 includes an amino acid sequence shown in SEQ ID NO: 34 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 34.

[0302] In some embodiments, where the VL includes a framework region LFR3, wherein the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3; and the LFR3 includes an amino acid sequence shown in SEQ ID NO: 35 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 35.

[0303] In some embodiments, where the VL includes a framework region LFR4, wherein the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3, and the LFR4 includes an amino acid sequence shown in SEQ ID NO: 36 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 36.

[0304] In some embodiments, where the VL includes framework regions LFR1, LFR2, LFR3 and LFR4, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3; among them, the LFR1 includes an amino acid sequence shown in SEQ ID NO: 33 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 33, the LFR2 includes an amino acid sequence shown in SEQ ID NO: 34 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 34, the LFR3 includes an amino acid sequence shown in SEQ ID NO: 35 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO: 35.

SEQ ID NO:44 or an amino acid sequence having at least about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% sequence identity to the amino acid sequence shown in SEQ ID NO:42, SEQ ID NO:43 or SEQ ID NO:44.

[0319] For example, where the anti PD-L1 antibody can include HC and LC, the HC can include an amino acid sequence shown in SEQ ID NO: 26, and the LC can include an amino acid sequence shown in SEQ ID NO:42, SEQ ID NO:43 or SEQ ID NO:44.

[0320] For example, the pharmaceutical combination can include: 1) anti PD-L1 antibody, wherein the anti PD-L1 antibody can include HC and LC, the HC can include an amino acid sequence shown in SEQ ID NO: 26, and the LC can include an amino acid sequence shown in SEQ ID NO:42, SEQ ID NO: 43 or SEQ ID NO:44; 2) a STING pathway agonist, which can be 2',3'-cGAMP or derivatives thereof.

[0321] In some embodiments, i) the PD-1 inhibitor and/or PD-L1 inhibitor in the pharmaceutical combination are not mixed with ii) the STING pathway agonist to each other in the pharmaceutical combination, i.e., i) the PD-1 inhibitor and/or ii) the PD-L1 inhibitor and STING pathway agonist are located in separate dosage forms.

[0322] In some embodiments, i) the PD-1 inhibitor and/or PD-L1 inhibitor, and ii) the STING pathway agonist are present in the pharmaceutical combination in a single dosage form.

[0323] In some embodiments, where the pharmaceutical combination is formulated into a pharmaceutical composition (e.g., a composite formulation or compound formulation). This pharmaceutical combination can be directly injected into large tumors without affecting normal (surrounding) tissues, enabling the killing of cancer cells, preventing or delaying the growth of malignant mass (e.g., making the tumor smaller or regression), and enabling advanced cancer patients to live with the tumor in a similar manner to those living with parasites. When a pharmaceutical combination is injected into a tumor, the drug can flow along blood vessels or lymphatic vessels to the metastatic lesion, and it will kill the metastatic cells. The injection of pharmaceutical combination into the tumor results in minimal trauma to the patient and can be repeated multiple times a month, for example. Direct injection can also be applied simultaneously to primary tumors and secondary tumors to which cancer has metastasized.

[0324] In some embodiments, where the pharmaceutical composition includes a PD-1 inhibitor or a PD-L1 inhibitor, and a STING pathway agonist.

[0325] For example, the pharmaceutical composition can include a PD-1 inhibitor and a STING pathway agonist. For example, the pharmaceutical composition can include an anti PD-1 antibody or antigen-binding fragments thereof and a cyclic dinucleotide.

[0326] For example, the pharmaceutical composition can include a PD-L1 inhibitor and a STING pathway agonist. Yet for example, the pharmaceutical composition can include an anti PD-L1 antibody or antigen-binding fragments thereof and a cyclic dinucleotide.

[0327] In some embodiments, where the STING pathway agonist is present in an amount of about 0.0001 mg/kg to about 200 mg/kg. For example, the STING pathway agonist can be present in the combination described herein in an

amount relative to the subject's body weight (i.e., mg/kg). In some cases, the STING pathway agonist is present in an amount equivalent to about 0.0001 mg/kg to about 200 mg/kg, 0.001 mg/kg to about 200 mg/kg, 0.01 mg/kg to about 200 mg/kg, 0.01 mg/kg to about 150 mg/kg, 0.01 mg/kg to about 100 mg/kg, 0.01 mg/kg to about 50 mg/kg, 0.01 mg/kg to about 25 mg/kg, 0.01 mg/kg to about 10 mg/kg or 0.01 mg/kg to about 5 mg/kg, 0.05 mg/kg to about 200 mg/kg, 0.05 mg/kg to about 150 mg/kg, 0.05 mg/kg to about 100 mg/kg, 0.05 mg/kg to about 50 mg/kg, 0.05 mg/kg to about 25 mg/kg, 0.05 mg/kg to about 10 mg/kg or 0.05 mg/kg to about 5 mg/kg, 0.5 mg/kg to about 200 mg/kg, 0.5 mg/kg to about 150 mg/kg, 0.5 mg/kg to about 100 mg/kg, 0.5 mg/kg to about 50 mg/kg, 0.5 mg/kg to about 25 mg/kg, 0.5 mg/kg to about 10 mg/kg or 0.5 mg/kg to about 5 mg/kg. In other cases, the STING pathway agonist is present in an amount equivalent to about 1 mg/kg to about 200 mg/kg, 1 mg/kg to about 150 mg/kg, 1 mg/kg to about 100 mg/kg, 1 mg/kg to about 50 mg/kg, 1 mg/kg to about 25 mg/kg, 1 mg/kg to about 10 mg/kg, or 1 mg/kg to about 5 mg/kg.

[0328] Yet for example, the STING pathway agonist can be present in the combination in an amount of about 1 mg, 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 75 mg, 80 mg, 90 mg, 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 400 mg, 500 mg, 600 mg, 700 mg, 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, 1500 mg, 1600 mg, 1700 mg, 1800 mg, 1900 mg or 2000 mg. The STING pathway agonist can be present in the combination in an amount of about 1 mg to about 10 mg, 10 mg to about 20 mg, 25 mg to about 50 mg, 30 mg to about 60 mg, 40 mg to about 50 mg, 50 mg to about 100 mg, 75 mg to about 150 mg, 100 mg to about 200 mg, 200 mg to about 500 mg, 500 mg to about 1000 mg, 1000 mg to about 1200 mg, 1000 mg to about 1500 mg, 1200 mg to about 1500 mg, or 1500 mg to about 2000 mg.

[0329] Yet for example, the STING pathway agonist can be present in the combination in an amount of about 0.1 mg/mL, 0.5 mg/mL, 1 mg/mL, 2 mg/mL, 3 mg/mL, 4 mg/mL, 5 mg/mL, 6 mg/mL, 7 mg/mL, 8 mg/mL, 9 mg/mL, 10 mg/mL, 15 mg/mL, 20 mg/mL, 25 mg/mL, 30 mg/mL, 40 mg/mL, 50 mg/mL, 60 mg/mL, 70 mg/mL, 80 mg/mL, 90 mg/mL, 100 mg/mL, 150 mg/mL, 200 mg/mL, 250 mg/mL, 300 mg/mL, 400 mg/mL, or 500 mg/mL. In some embodiments, the STING pathway agonist can be present in the combination in an amount of about 1 mg/mL to about 10 mg/mL, 5 mg/mL to about 10 mg/mL, 5 mg/mL to about 15 mg/mL, 10 mg/mL to about 25 mg/mL, 20 mg/mL to about 30 mg/mL, 25 mg/mL to about 50 mg/mL or 50 mg/mL to about 100 mg/mL.

[0330] In some embodiments, where the PD-1 inhibitor or PD-L1 inhibitor (e.g., the anti PD-1/PD-L1 antibody) is present in an amount of about 0.0001 mg/kg to about 200 mg/kg. For example, the PD-1 inhibitor or PD-L1 inhibitor can be present in the combination described herein in an amount relative to the subject's body weight (i.e., mg/kg). In some cases, the PD-1 inhibitor or PD-L1 inhibitor is present in an amount equivalent to about 0.0001 mg/kg to about 200 mg/kg, 0.001 mg/kg to about 200 mg/kg, 0.01 mg/kg to about 200 mg/kg, 0.01 mg/kg to about 150 mg/kg, 0.01 mg/kg to about 100 mg/kg, 0.01 mg/kg to about 50 mg/kg, 0.01 mg/kg to about 25 mg/kg, 0.01 mg/kg to about 10 mg/kg or 0.01 mg/kg to about 5 mg/kg, 0.05 mg/kg to about 200 mg/kg, 0.05 mg/kg to about 150 mg/kg, 0.05 mg/kg to about 100 mg/kg, 0.05 mg/kg to about 50 mg/kg, 0.05 mg/kg to

to about 25 mg/kg, 0.05 mg/kg to about 10 mg/kg or 0.05 mg/kg to about 5 mg/kg, 0.5 mg/kg to about 200 mg/kg, 0.5 mg/kg to about 150 mg/kg, 0.5 mg/kg to about 100 mg/kg, 0.5 mg/kg to about 50 mg/kg, 0.5 mg/kg to about 25 mg/kg, 0.5 mg/kg to about 10 mg/kg or 0.5 mg/kg to about 5 mg/kg. In other cases, the PD-1 inhibitor or PD-L1 inhibitor is present in an amount equivalent to about 1 mg/kg to about 200 mg/kg, 1 mg/kg to about 150 mg/kg, 1 mg/kg to about 100 mg/kg, 1 mg/kg to about 50 mg/kg, 1 mg/kg to about 25 mg/kg, 1 mg/kg to about 10 mg/kg, or 1 mg/kg to about 5 mg/kg.

[0331] Yet for example, the anti PD-1/PD-L1 antibody can be present in the combination in an amount of about 1 mg, 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 75 mg, 80 mg, 90 mg, 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 400 mg, 500 mg, 600 mg, 700 mg, 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, 1500 mg, 1600 mg, 1700 mg, 1800 mg, 1900 mg, or 2000 mg. The anti PD-1/PD-L1 antibody can be present in the combination in an amount of about 1 mg to about 10 mg, 10 mg to about 20 mg, 25 mg to about 50 mg, 30 mg to about 60 mg, 40 mg to about 50 mg, 50 mg to about 100 mg, 75 mg to about 150 mg, 100 mg to about 200 mg, 200 mg to about 500 mg, 500 mg to about 1000 mg, 1000 mg to about 1200 mg, 1000 mg to about 1500 mg, 1200 mg to about 1500 mg, or 1500 mg to about 2000 mg.

[0332] Yet for example, the anti PD-1/PD-L1 antibody can be present in the combination in an amount of about 0.1 mg/mL, 0.5 mg/mL, 1 mg/mL, 2 mg/mL, 3 mg/mL, 4 mg/mL, 5 mg/mL, 6 mg/mL, 7 mg/mL, 8 mg/mL, 9 mg/mL, 10 mg/mL, 15 mg/mL, 20 mg/mL, 25 mg/mL, 30 mg/mL, 40 mg/mL, 50 mg/mL, 60 mg/mL, 70 mg/mL, 80 mg/mL, 90 mg/mL, 100 mg/mL, 150 mg/mL, 200 mg/mL, 250 mg/mL, 300 mg/mL, 400 mg/mL, or 500 mg/mL. In some embodiments, the anti PD-1/PD-L1 antibody can be present in the combination in an amount of about 1 mg/mL to about 10 mg/mL, 5 mg/mL to about 10 mg/mL, 5 mg/mL to about 15 mg/mL, 10 mg/mL to about 25 mg/mL, 20 mg/mL to about 30 mg/mL, 25 mg/mL to about 50 mg/mL or 50 mg/mL to about 100 mg/mL.

[0333] In some embodiments, the STING pathway agonist can be provided in a synergistic amount with the amount of the PD-1/PD-L1 inhibitor. The administration dosage will undoubtedly vary with known factors, such as the pharmacokinetic properties of a particular agent and mode and route of administration thereof; age, physical condition, and weight of the recipient; nature and extent of symptoms, type of concurrent treatment, frequency of treatment, and expected effect.

[0334] In some embodiments, where the pharmaceutical composition further includes one or more pharmaceutically acceptable carriers.

[0335] In another aspect, the disclosure also provides a use of the aforementioned pharmaceutical combination in the preparation of drugs for the treatment of neoplastic diseases.

[0336] In some embodiments, where the neoplastic diseases include tumors and/or wart diseases. In some embodiments, the tumors (or cancers) include but are not limited to: hepatocellular carcinoma, hepatic metastasis cancer, advanced hepatocellular carcinoma, pancreatic carcinoma, adenocarcinoma, mastocytoma or mast cell tumor, ovarian carcinoma, non-small-cell lung carcinoma, small-cell lung carcinoma, melanoma, retinoblastoma, breast neoplasms, colorectal carcinoma, histiocytosarcoma, brain neoplasms,

astrocytoma, glioblastoma, neuroma, neuroblastoma, colorectal carcinoma, cervical carcinoma, sarcoma, prostatic neoplasms, bladder neoplasms, reticuloendothelium neoplasms, nephroblastoma, ovarian carcinoma, osteocarcinoma, osteosarcoma, renal carcinoma, or head and neck cancer, oral carcinoma, laryngocarcinoma, or oropharyngeal carcinoma, breast carcinoma, genitourinary tract carcinoma, lung carcinoma, gastrointestinal carcinoma, epidermoid carcinoma, or melanoma.

[0337] In another aspect, the disclosure also provides the aforementioned pharmaceutical combination for use in the treatment of neoplastic diseases.

[0338] In another aspect, the disclosure also provides a drug including the aforementioned pharmaceutical combination for treating neoplastic diseases.

[0339] In another aspect, the disclosure also provides a method for treating neoplastic diseases, which includes administering an effective amount of the aforementioned pharmaceutical combination to a subject in need thereof.

[0340] In some embodiments, where the subject suffers from neoplasm.

[0341] In some embodiments, where the neoplasm of mammals is located at a site selected from the group consisting of:

[0342] brain, head, eyes, nasopharynx, mouth, tongue, neck, thyroid, gastrointestinal system, liver, pancreas, gall bladder, lungs, respiratory system, genitourinary system, kidney, bladder, breast, lymphatic system, cardiovascular system, nervous system, skin, thoracic cavity, pleura, musculoskeletal system, and abdomen, which has primary or secondary natures.

[0343] In some embodiments, where the neoplasm includes tumors and/or warts.

[0344] In some embodiments, where the administration includes local, intraneoplastic (e.g., in tumors or warts), or systemic administrations. For example, using intratumoral or intraverrucous injection allows for both less trauma to the patient and the killing of cancer cells instead of normal cells. The direct injection of pharmaceutical combinations into malignant tumors also greatly reduces or eliminates many common side effects.

[0345] In some embodiments, where the administration includes intravenous injection, intravenous instillation, intramuscular injection, subcutaneous injection, and/or intraneoplastic injection.

[0346] In some embodiments, the tumor includes:

[0347] (i) superficial malignant diseases of skin, eyes, tongue, mouth, thyroid, breast, cervix, uterus, anus, prostate, vagina, osteosarcoma, urethral carcinoma, penis, testicles, and epididymis, and the pharmaceutical combination is directly injected into the tumor body without dilution using a syringe; or

[0348] (ii) cancer of nasopharynx, and the pharmaceutical combination is injected into the tumor body through a nasopharyngoscopy using a syringe or a needle; or

[0349] (iii) cancers of liver, kidney, and gallbladder, and the pharmaceutical combination is injected into the tumor body through the skin with the assistance of ultrasound using a syringe, or into the tumor body through a pore formed in the patients' abdominal wall during laparoscopic surgery; or

[0350] (iv) cancers of ovarian, fallopian tube, pancreas, lymph node metastasis or direct peritoneal invasion of

the abdominal cavity, and abdominal lymphoma, and the pharmaceutical combination is injected into the tumor body through a pore formed in the patients' abdominal wall during laparoscopic surgery using a syringe; or

[0351] (v) cancers or sarcoma of esophagus, stomach, duodenum, and small intestine, and the pharmaceutical combination is injected into the tumor body through colonoscopy using a needle, or through a pore formed in the patients' abdominal wall during laparoscopic surgery through a long syringe, or through a pore formed in the patients' chest wall during thoracoscopic surgery;

[0352] (vi) cancers or sarcoma of large intestine and rectum, and the pharmaceutical combination is injected into the tumor body through colonoscopy using a needle, or through a pore formed in the patients' abdominal wall during laparoscopic surgery using a syringe; or

[0353] (vii) cancers or sarcoma of lung and trachea, and the pharmaceutical combination is injected into the tumor body through a fiberbronchoscopy using a needle; or

[0354] (viii) cancer of lung, and the pharmaceutical combination is injected with a syringe using ultrasound, X-ray, CT scan, or MR scan, or through a pore formed in the patients' chest wall during thoracoscopic surgery; or

[0355] (ix) cancer or sarcoma of bladder, and the pharmaceutical combination is injected into the tumor body through cystoscopy using a needle, or through a pore formed in the patients' abdominal wall during laparoscopic surgery; or

[0356] (x) cancer or sarcoma of uterus, and the injectable formulation of the pharmaceutical combination is injected into the tumor body using a syringe or hysteroscopic needle; or injected through a pore formed in the patients' abdominal wall during laparoscopic surgery; or

[0357] (xi) cancer or sarcoma of nasopharynx and larynx, and the pharmaceutical combination is injected into the tumor body through laryngoscopy using a needle; or

[0358] (xii) cancer of brain, and the pharmaceutical combination is injected into the tumor body with a syringe or fibroscopic needle by use of X-ray, CT scan, or MR scan after drilling a pore in the corresponding skull; or

[0359] (xiii) malignant lymphoma or lymph nodes with metastasis, and the pharmaceutical combination is injected into the tumor body through the patients' skin using a needle, or through a pore formed in the patients' abdominal wall during laparoscopic surgery, or through a pore formed in the patients' chest wall during thoracoscopic surgery.

[0360] In some embodiments, the neoplasm includes warts, and the pharmaceutical combination is injected into the wart body through the patients' skin using a needle.

[0361] In some embodiments, where i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered by use of the same or different administration routes.

[0362] In some embodiments, it includes injection of the STING pathway agonist into the neoplasm.

[0363] In some embodiments, it also includes injection or systemic infusion (e.g., intravenous injection and intravenous instillation) of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm.

[0364] In some embodiments, it includes injection of i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination into the neoplasm.

[0365] In some embodiments, where i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered simultaneously or at different times.

[0366] In some embodiments, where the PD-1 inhibitor or PD-L1 inhibitor is administered before and/or after the administration of the STING pathway agonist.

[0367] For example, the PD-1/PD-L1 inhibitor is administered at least 5 minutes, 10 minutes, 20 minutes, 40 minutes, 1 hour, 2 hours, 4 hours, 8 hours, 12 hours, 16 hours, 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days or 2 weeks before and/or after the administration of the STING pathway agonist.

[0368] In some embodiments, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are simultaneously administered by way of intraneoplastic injection.

[0369] In some embodiments, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are simultaneously administered by way of intraneoplastic injection, and i) the PD-1 inhibitor or PD-L1 inhibitor and ii) the STING pathway agonist are present in a same dosage form.

[0370] In some embodiments, where the PD-1 inhibitor or PD-L1 inhibitor in the pharmaceutical combination is administered by way of intravenous injection, the STING pathway agonist is simultaneously administered by way of intraneoplastic injection, and the PD-1 inhibitor or PD-L1 inhibitor and the STING pathway agonist are present in separate dosage forms.

[0371] In another aspect, the disclosure provides a pharmaceutical kit including the aforementioned pharmaceutical combination.

[0372] In some embodiments, each component of the pharmaceutical combination in the pharmaceutical kit can be provided in separated individual containers. Alternatively, the components of the pharmaceutical combination described herein can be provided in a single container. In such cases, the container can be a container prepared for disclosure to patients in need thereof, such as IV bags, ampoules, or syringes. In some embodiments, the STING pathway agonist in the pharmaceutical kit is formulated for administration through intratumoral or intraverrucous injection. The PD-1/PD-L1 inhibitor can be provided in the form of powder (e.g., lyophilized powder) or parenteral solution, for example. In some cases, the PD-1/PD-L1 inhibitor can be an anti PD-1/PD-L1 antibody as described herein, formulated for use, for example through intravenous administration or intratumoral or intraverrucous injection. In some embodiments, the STING pathway agonist and the PD-1/PD-L1 inhibitor are formulated as a compound formulation and used for administration through intratumoral or intraverrucous injection.

[0373] The content of the pharmaceutical kit described herein can be provided in sterile form. The pharmaceutical kit and content therein can be provided in the form of

ready-to-use to a subject in need thereof. In such cases, the components in the pharmaceutical kit are provided as a formulation and optionally provided in an administration device, so that the administration requires almost no further action by the user. In the case where the pharmaceutical kit includes an administration device, such devices include devices known and understood by those skilled in the art for the administration routes described herein, such as but not limited to syringes, pumps, bags, cups, inhalators, droppers, patches, creams, or injectors.

[0374] The pharmaceutical kit described herein may also include an instruction, which contains information about, for example, usage, dosage, administration, contraindications, and/or warnings about the use of such drugs.

[0375] Without being limited by any theory, the examples hereinafter are only intended to illustrate the pharmaceutical combinations and uses of the disclosure, rather than limiting the scope of invention of the disclosure.

EXAMPLES

Animal Species, Strain, Gender, Weight, Sources, Certificate of Conformity

[0376] C57/BL6JNifdc mouse, female, body weight of 17-22 g, aged 6-8 weeks, SPF grade, purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. [Experimental Animal Quality Certificate No.: SCXK (Beijing) 2016-0006].

Feeding Conditions

[0377] All mice were free to forage and drink water, and were raised at room temperature ($25\pm 2^{\circ}\text{C}$.) The feed and water were subjected to high-pressure sterilization treatment, and the entire experimental feeding process was SPF grade.

Tumor Cell Strains

[0378] Mouse colorectal carcinoma cell strain MC38, mouse lung carcinoma Lewis tumor strain LLC1, mouse melanoma cell strain B16, mouse prostatic carcinoma cell strain RM-1, TRAMP-C1, mouse breast carcinoma cell strains 4T1, and NAFA, etc.

Establishment of Subcutaneous Xenograft Tumor Model

[0379] Tumor cell strains were selected from the group consisting of: mouse colorectal carcinoma cell strains CT26 and MC38, mouse lung carcinoma Lewis tumor strain LLC1, mouse melanoma cell strain B16, mouse prostatic carcinoma cell strain RM-1, TRAMP-C1, mouse breast carcinoma cell strains 4T1, and NAFA, etc.

[0380] Tumor cells were cultured and passaged, cells were collected at the logarithmic phase, and formulated into a concentration of (1.0×10^7) per milliliter of cell suspension with 0.1 ml (with a cell number of 1.0×10^6) being injected into the right flank of the mice, the tumor grew to a diameter of about 5 mm within about 10 days, and the model was successfully established, they were randomly divided into 8 groups with 8 mice per group.

Drug Evaluation Indicators

[0381] Trend of changes in body weight and tumor volume of mice, survival rate of mice, and inhibitory rate of anatomical tumor weight.

Tumor weight inhibitory rate =

$$\left[1 - \frac{\text{average tumor weight of experimental group (G2/G3/G4)}}{\text{average tumor weight of negative control group G1}}\right] \times 100\%.$$

[0382] Drug interaction coefficient: $\text{CDI}(\text{CI}) = \text{AB}/\text{A} \times \text{B}$. AB is the ratio of the combination group of two drugs to the control group, as calculated based on the anatomical tumor weight. A or B is the ratio of single drug group to control group. If CDI is less than 1, it indicates that the two drugs have a synergistic effect; If CDI equals to 1, the two drugs have an additive effect; If CDI is greater than 1, the two drugs have an antagonistic effect.

Statistical Analysis

[0383] The data were represented as $\bar{x} \pm s$, and processed by SPSS10.0 software, using one-way ANOVA. Differences in data among groups (P value) were processed through statistical processing, and there was a significant difference among groups at $P < 0.05$.

Example 1

[0384] The anti-tumor effect of a pharmaceutical combination composed of PD-L1 inhibitor RB0005 and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

Main Steps of the Test

1.1 Xenograft Tumor Model in Mice

[0385] After successfully establishing a subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, it was randomly grouped with 6-8 mice per group. The mice were administered once every two days, with a total of three times followed by observation until 15 days.

Groups	Administration Method	Administration dosage	Administration frequency
G1 Negative Control	Intratumor injection	Solvent	Administered every other day with a total of 3 times
G2 single drug group	Intratumor injection	PD-L1 monoclonal antibody RB0005, 1 mg/kg	Administered every other day with a total of 3 times

-continued

Groups	Administration Method	Administration dosage	Administration frequency
G3 single drug group	Intratumoral injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administered every other day with a total of 3 times
G4 trial group (Single dosage form)	Intratumoral injection	PD-L1 monoclonal antibody RB0005, 1 mg/kg + 2',3'-cGAMP, 1 mg/kg;	Administered every other day with a total of 3 times

1.2 Results

[0386] FIGS. 1-4 show that both the novel pharmaceutical combination and each single drug can significantly inhibit the growth of tumor, and the anatomical tumor weight is significantly lower than the negative control group ($P<0.01$, $P<0.001$). The novel pharmaceutical combinations have better efficacy than 2',3'-cGAMP or PD-L1 antibody RB0005 alone, indicating that the novel pharmaceutical combinations have greater advantages.

[0387] As shown in FIG. 2, the mice in solvent control group began to experience "death" on day 9 (when the tumor volume exceeded 2000 mm³ in a single mouse, the mouse could be considered as death during the statistical analysis of survival rate in mouse (which was not excluded in the analysis of tumor volume change curve). During day 11 to day 13, the survival rate of mice in the solvent control group decreased most significantly; all mice in single drug group began to experience "death" on day 15, there was no significant difference between the two groups; one mouse in the G4 trial group had tumor regression and one mouse had almost complete regression (with a tumor volume of <100 mm³). At the endpoint of the experiment, the survival rates of mice in each group were 0%, 37.5%, 75%, and 87.5%, respectively.

[0388] As shown in FIG. 3A-3B, the tumor volume of mice in G1 solvent control group gradually increased over time; and the growth of tumor volume of mice in the G4 trial group (RB0005+2',3'-cGAMP) significantly slowed down and the overall volume decreased.

[0389] As shown in FIG. 4, the tumor weight of mice in the G4 trial group was significantly reduced, with a tumor weight inhibitory rate of 73%, which showed a statistically significant difference ($P<0.001$).

Example 2

[0390] The anti-tumor effect of a pharmaceutical combination composed of PD-1 inhibitor RB0004 and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

2.1 Xenograft Tumor Model in Mice

[0391] After successfully establishing a humanized PD-1 subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, they were randomly grouped with 8 mice per group and administered once every two days.

Groups	Administration Method	Administration dosage	Administration frequency
G1 Negative Control	Intratumoral injection	Solvent	Administering every other day
G2 single drug group	Intratumoral injection	2',3'-cGAMP monotherapy, 3 mg/kg	Administering every other day
G3 single drug group	Intraperitoneal injection (i.p.)	PD-1 monoclonal antibody RB0004, 10 mg/kg	Administering every other day
G4 trial group (separate dosage form)	Intratumoral injection	2',3'-cGAMP monotherapy, 3 mg/kg	Administering every other day
	Intraperitoneal injection	PD-1 monoclonal antibody RB0004, 10 mg/kg	

2.2 Results

[0392] FIGS. 5-6 show that both the novel pharmaceutical combination (separate dosage form) and each single drug can inhibit the growth of tumor, and the anatomical tumor weight is significantly lower than the negative control group ($P<0.05$, $P<0.001$). The novel pharmaceutical combinations have better efficacy than 2',3'-cGAMP or PD-1 antibody RB0004 alone, indicating that the novel pharmaceutical combinations have greater advantages.

[0393] As shown in FIG. 5, the tumor volume of mice in the solvent control group (G1) gradually increased over time; after treatment, the growth in tumor volume of mice in the single drug group (G2/G3) was slower than that in the solvent control group, and the tumor volume was smaller; the growth in tumor volume of mice in the trial group (G4) was significantly slowed down, and the overall volume showed a decrease.

[0394] As shown in FIG. 6, the tumor weight of mice in the trial group (G4) was significantly reduced, with a tumor weight inhibitory rate of 96%, which showed a statistically significant difference ($P<0.001$).

Example 3

[0395] The anti-tumor effect of a pharmaceutical combination composed of PD-L1 inhibitor RB0005 and different doses of the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model of colorectal carcinoma MC38 in mice, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

3.1 Xenograft Tumor Model in Mice

[0396] After successfully establishing a subcutaneous xenograft tumor model in mice, they were randomly grouped with 8 mice per group and administered once every two days.

Groups	Administration Method	Administration dosage	Administration frequency
A Negative Control	Intratumor injection	Solvent	Administering every other day
B single drug group	Intratumor injection	PD-L1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
C single drug group	Intratumor injection	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
D single drug group	Intratumor injection	2',3'-cGAMP monotherapy, 3.16 mg/kg	Administering every other day
E single drug group	Intratumor injection	2',3'-cGAMP monotherapy, 10 mg/kg	Administering every other day
F trial group (separate dosage form)	Intratumor injection	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G trial group (separate dosage form)	Intraperitoneal injection	PD-L1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
H trial group (separate dosage form)	Intratumor injection	2',3'-cGAMP monotherapy, 3.16 mg/kg	Administering every other day
	Intraperitoneal injection	PD-L1 monoclonal antibody RB0005, 1 mg/kg	
	Intratumor injection	2',3'-cGAMP monotherapy, 10 mg/kg	Administering every other day
	Intraperitoneal injection	PD-L1 monoclonal antibody RB0005, 1 mg/kg	

3.2 Results

[0397] FIG. 7 shows that all three pharmaceutical combinations and single drug alone can inhibit the growth of tumor, and the pharmaceutical combination has a more significant inhibitory effect, with tumor weight inhibitory rates of over 90% ($P < 0.001$). The novel pharmaceutical combinations have better efficacy than PD-L1 antibody RB0005 and 2',3'-cGAMP used alone, indicating that the novel pharmaceutical combinations have greater advantages.

Example 4

[0398] The anti-tumor effect of a pharmaceutical combination composed of PD-1 inhibitor RB0004 and different doses of the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

4.1 Xenograft Tumor Model in Mice

[0399] After successfully establishing a subcutaneous xenograft tumor model in mice, they were randomly grouped with 8 mice per group and administered once every two days.

Groups	Administration Method	Administration dosage	Administration frequency
G1 Negative Control	Intratumor injection	Solvent	Administering every other day
G2 single drug group	Intraperitoneal injection	PD-1 monoclonal antibody RB0004, 10 mg/kg	Administering every other day

-continued

Groups	Administration Method	Administration dosage	Administration frequency
G3 trial group (separate dosage form)	Intratumor injection	2',3'-cGAMP monotherapy, 0.03 mg/kg	Administering every other day
	Intraperitoneal injection	PD-1 monoclonal antibody RB0004, 10 mg/kg	
G4 trial group (separate dosage form)	Intratumor injection	2',3'-cGAMP monotherapy, 0.1 mg/kg	Administering every other day
	Intraperitoneal injection	PD-1 monoclonal antibody RB0004, 10 mg/kg	
G5 trial group (separate dosage form)	Intratumor injection	2',3'-cGAMP monotherapy, 0.3 mg/kg	Administering every other day
	Intraperitoneal injection	PD-1 monoclonal antibody RB0004, 10 mg/kg	

4.2 Results

[0400] FIGS. 8-9 show that all three pharmaceutical combinations can inhibit the growth of tumor, and the tumor weight inhibitory rate is above 70%. Especially in the combined 2',3'-cGAMP 0.1 mpk dose group, the inhibitory effect on tumor is significant and the survival rate is high ($P < 0.05$). The efficacy of the novel pharmaceutical combinations is better than that of PD-1 antibody RB0004 alone, indicating that the novel pharmaceutical combinations have greater advantages.

Example 5

[0401] The anti-tumor effect of a pharmaceutical combination composed of PD-1/PD-L1 inhibitor and the STING

agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

5.1 Xenograft Tumor Model in Mice

[0402] After successfully establishing a humanized CD274 cell subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

Groups	Admin- istration Method	Admin- istration dose	Admin- istration frequency
G1 control group	Intratumor injection (i.t.) Intraperitoneal injection (i.p.)	Solvent	Administering every other day
G2 single drug group	Intraperitoneal injection (i.p.)	PD-1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
G4 single drug group	Intraperitoneal injection (i.p.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G5 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G6 trial group (Single dosage form)	Intraperitoneal injection (i.p.)	LPG2008: PD-1 monoclonal antibody RB0005 1mg/kg + 2',3'-cGAMP, 1 mg/kg	Administering every other day
G7 trial group (Single dosage form)	Intratumor injection (i.t.)	LPG2008: PD-1 monoclonal antibody RB0005 1 mg/kg + 2',3'-cGAMP, 1 mg/kg	Administering every other day
G8 trial group (separate dosage form)	Intraperitoneal injection (i.p.) Intratumor injection (i.t.)	PD-1 monoclonal antibody RB0005, 1 mg/kg 2',3'-cGAMP, 1 mg/kg	Administering every other day Administering every other day

5.2 Results

TABLE 1

Average tumor mass and tumor inhibitory rate in each group			
Groups	Tumor Weight (g)	IR _{TW} %	CDI (Coefficient of drug interaction)
G1	4.43	—	
G2	3.24	26.8	
G3	3.55	19.9	
G4	3.48	21.5	
G5	1.84	58.5	
G6	2.40	45.9	0.94 (<1)
G7	0.81	81.7	0.55 (<1)
G8	0.97	78.1	0.72 (<1)

[0403] The results are shown in Table 1 and FIGS. 10 to 12:

[0404] The tumor weight inhibitory rates of single drug RB0005 via intraperitoneal and intratumoral (G2\G3) administration modes were 26.8% and 19.9%, respectively; compared with the intraperitoneal and intratumoral (G4\G5)

administration modes of cGAMP, the intratumoral administration had a better effect, with a tumor weight inhibitory rate of 58.5%. The intraperitoneal administration had no significant tumor inhibitory effect, and there was statistically significant difference between the intratumoral administration and the control groups ($P < 0.001$);

[0405] The novel composite formulation, single dosage form (G6\G7) composite formulation, was administered intraperitoneally and intratumorally. The tumor weight inhibitory rate of intratumoral administration was 81.7%, and the tumor inhibitory effect was superior to that of intraperitoneal administration. The survival rate of mice at the endpoint of the experiment was 100%.

[0406] The novel composite formulation in separate dosage form (G8) also exhibited tumor inhibitory effects, with a tumor weight inhibitory rate of 78.1% and a survival rate of mice of 100% at the endpoint of the experiment.

[0407] The novel pharmaceutical combinations in this disclosure exhibited good drug synergistic effects, whether in separate dosage forms or single dosage forms and regardless of administering intraperitoneally or intratumorally. Among them, the single dosage form of the composite formulation exhibited significant drug synergistic effects, with a CDI interaction value of 0.55.

Example 6

[0408] The anti-tumor effect of a pharmaceutical combination composed of PD-1 inhibitor RB0004 and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

6.1 Xenograft Tumor Model in Mice

[0409] After successfully establishing a humanized PD-1 subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

6.2 Grouping

Groups	Admin- istration Method	Admin- istration dose	Admin- istration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody RB0004, 10 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	LPG2009: PD-1 monoclonal antibody RB0004 10 mg/kg + 2',3'-cGAMP, 1 mg/kg	Administering every other day

6.3 Results

TABLE 2

Average tumor mass and tumor inhibitory rate in each group			
Groups	Tumor Weight (g)	IR _{TW} %	CDI (Coefficient of drug interaction)
G1	2.21	—	0.45
G2	0.87	60.8	
G3	0.84	61.9	
G4	0.15	93.2	

[0410] The results are shown in Table 2 and FIGS. 13 to 15:

[0411] The tumor weight inhibitory rate of RB0004 monotherapy was 60.8%; and the tumor weight inhibitory rate of cGAMP monotherapy was 61.9%;

[0412] The composite formulation (single dosage form) (G4) had a significant inhibitory effect on tumor proliferation. After administration, the tumor proliferation slowed down, and the tumor weight inhibitory rate was 93.2%, there was statistically significant difference compared with the control group (P<0.001). The survival rate of mice at the endpoint of the experiment was 100%, showing a significant drug synergistic effect (CDI value=0.45).

[0413] In summary, the single dosage form of 2',3'-cGAMP and PD-1 monoclonal antibody has a significant tumor inhibitory effect, high survival rate (P<0.05), and significant drug synergistic effects. The efficacy of the novel pharmaceutical combinations is better than that of PD-1 antibody RB0004 alone, indicating that the novel pharmaceutical combinations have greater advantages.

Example 7

[0414] The anti-tumor effect of a pharmaceutical combination composed of the marketed PD-1/PD-L1 inhibitor and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

7.1 Xenograft Tumor Model in Mice

[0415] After successfully establishing a subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

7.2.1 Grouping 1 (PD-1 Inhibitor)

Groups	Admin-istration Method	Admin-istration dose	Admin-istration frequency
G1 Control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody Tislelizumab, 10 mg/kg	Administering every other day

-continued

Groups	Admin-istration Method	Admin-istration dose	Admin-istration frequency
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody Tislelizumab, 10 mg/kg + 2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G5 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody Camrelizumab, 10 mg/kg	Administering every other day
G6 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody Camrelizumab, 10 mg/kg + 2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G7 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody Sintilimab, 10 mg/kg	Administering every other day
G8 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody Sintilimab, 10 mg/kg + 2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G9 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody Toripalimab, 10 mg/kg	Administering every other day
G10 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody Toripalimab, 10 mg/kg + 2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day

7.3.1 Result 1 (PD-1 Inhibitor)

TABLE 3

Average tumor mass and tumor inhibitory rate in each group			
Groups	Tumor Weight (g)	IR _{TW} %	CDI (Coefficient of drug interaction)
G1	1.53	—	0.41
G2	0.98	36	
G3	0.59	61.4	
G4	0.18	88.2	
G5	1.23	19.6	0.11
G6	0.32	79.1	
G7	1.47	3.9	
G8	0.1	93.5	
G9	0.92	49.7	1
G10	0.42	77.1	

[0416] The results are shown in Table 3 and FIGS. 16 to 18:

[0417] Compared with the control group, the composite formulation of the cGAMP and the marketed PD1 inhibitor G4/G8/G10 showed statistically significant difference (P<0.001), with tumor weight inhibitory rates of 88.2%, 79.1%, 93.5%, and 77.1%, respectively. All four composite formulations showed drug synergistic effects, wherein the com-

posite formulation in G8 group (Sintilimab) had the most significant drug synergistic effect.

[0418] At the endpoint of the experiment, the survival rate of G2 (LPG1501) \G3 (tislelizumab) \G9 (TERRE) in the single drug group was above 60%; and the survival rates of the composite formulation G4\G6\G8\G10S were all 100%, the survival rates were significantly improved.

[0419] In summary, the composite formulations of 2',3'-cGAMP and four marketed PD-1 monoclonal antibodies (single dosage form) showed significant tumor inhibitory effect, high survival rate ($P<0.05$), and exhibited drug synergistic effects.

7.2.2 Grouping 2 (PD-L1 Inhibitor Durvalumab)

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody Durvalumab, 1 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody Durvalumab, 1 mg/kg + 2',3'-cGAMP, 1 mg/kg	Administering every other day

7.3.2 Results (PD-L1 Inhibitor Durvalumab)

TABLE 4

Average tumor mass and tumor inhibitory rate in each group			
Groups	Tumor Weight (g)	IR _{TW} %	CDI (Coefficient of drug interaction)
G1	3.35	—	0.97
G2	2.42	27.7	
G3	1.74	48.0	
G4	1.22	63.6	

[0420] The results are shown in Table 4 and FIGS. 19 to 21:

[0421] The composite formulation (G4) of cGAMP and the marketed PD-L1 inhibitor (Durvalumab) had an inhibitory effect on tumor proliferation, and its efficacy was superior to that of cGAMP or PD-L1 antibody alone. There was a statistically significant difference ($P<0.001$), and the tumor weight inhibitory rate was 63.6%. At the endpoint of the experiment, the survival rate of mice in the composite formulation group was above 80%, the survival rate was significantly improved.

[0422] In summary, the composite formulation of 2',3'-cGAMP and the marketed PD-L1 monoclonal antibody (single dosage form) showed significant tumor inhibitory effect, high survival rate ($P<0.05$), and exhibited drug synergistic effect.

7.2.3 Grouping 3 (PD-L1 inhibitor Tecentriq)

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody Tecentriq, 1 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody Tecentriq, 1 mg/kg + 2',3'-cGAMP, 1 mg/kg	Administering every other day
G5 single drug group	Intratumor injection (i.t.)	PD-1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
G6 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody RB0005, 1 mg/kg + 2',3'-cGAMP, 1 mg/kg	Administering every other day

7.3.3 Results (PD-L1 Inhibitor Tecentriq)

TABLE 5

Average tumor mass and tumor inhibitory rate in each group		
Groups	Tumor Weight (g)	IR _{TW} %
G1	1.86	62.4
G2	0.7	
G3	0.9	
G4	0.7	
G5	0.8	56.9
G6	0.5	73.1

[0423] The results are shown in Table 5 and FIGS. 22 to 25:

[0424] The tumor weight inhibitory rate of the composite formulation of cGAMP and RB0005 (G6) was 73.1%, there was statistically significant difference compared with the control group ($p<0.05$); the tumor weight inhibitory rate of the composite formulation (G4) of cGAMP and the marketed PD-L1 inhibitor (Tecentriq) was 62.4%, and the efficacy of the composite formulation was superior to that of cGAMP or PD-L1 antibody alone.

[0425] During the administration, the tumor proliferation in the composite formulation group was slow or even disappeared, and at the endpoint of the experiment, the tumor of 50% mice in the composite formulation group disappeared ($T_v<80\text{ mm}^3$).

[0426] In summary, the composite formulation of 2',3'-cGAMP and RB0005 as well as the marketed PD-L1 monoclonal antibody (single dosage form) showed significant tumor inhibitory effect, and high survival rate ($P<0.05$).

Example 8

[0427] The anti-tumor effect of a pharmaceutical combination composed of PD-L1 inhibitor RB0005 and the STING agonist 2',3'-cGAMP during different tumor proliferation stages was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

8.1 Xenograft Tumor Model in Mice

[0428] After successfully establishing a humanized CD274 subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

8.2 Grouping

Groups	Administration Method	Initial tumor volume	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	—	PD-1 monoclonal antibody RB0005,	Administering every other day
G2 trial group (Single dosage form)	Intratumor injection (i.t.)	250 mm ³	1 mg/kg + 2',3'-cGAMP, 1 mg/kg	
G3 trial group (Single dosage form)	Intratumor injection (i.t.)	500 mm ³		
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	1000 mm ³		

8.3 Results

[0429] As shown in FIG. 26,
[0430] The tumor volume inhibitory rate of the composite formulation after administration during the early stage of tumor proliferation (with an average volume of 250 mm³) was 49.9%, which showed a significant statistical difference compared to the control group (P<0.01);
[0431] The tumor volume inhibitory rate of the composite formulation after administration during the rapid stage of tumor proliferation (with an average volume of 500 mm³) was 55.5%, which showed a statistical difference compared to the control group (P<0.05);
[0432] The tumor volume inhibitory rate of the composite formulation after administration during the late stage of tumor proliferation (with an average volume of 1000 mm³) was 26.4%, which showed no statistical difference compared to the control group (P>0.05);
[0433] In conclusion: the novel composite formulation (single dosage form) has tumor inhibitory effects during both the early stage of tumor proliferation and the rapid stage of tumor proliferation, especially during the early stage of tumor proliferation, where the inhibitory effect is more significant.

Example 9

[0434] The inhibitory effect on the growth of MC38 xenograft tumor model by a small molecule STING agonist 2',3'-cGAMP administered firstly, followed by PD-L1 inhibitor RB0005 administered after 48 hours, was tested with a xenograft tumor model.

9.1 Xenograft Tumor Model in Mice

[0435] After successfully establishing a humanized PD-1 subcutaneous xenograft tumor model of colorectal carcinoma MC38 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

9.2 Grouping

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Day 1, 3, 5
	Intraperitoneal injection (i.p.)		Day 3, 5, 7

-continued

Groups	Administration Method	Administration dose	Administration frequency
G2 single drug group	Intratumor injection (i.t.)	Solvent	Day 1, 3, 5
	Intraperitoneal injection (i.p.)	PD-1 monoclonal antibody RB0005, 1 mg/kg	Day 3, 5, 7
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Day 1, 3, 5
	Intraperitoneal injection (i.p.)	Solvent	Day 3, 5, 7
G4 trial group (	Intratumor injection (i.t.)	2',3'-cGAMP, 1 mg/kg	Day 1, 3, 5
	Intraperitoneal injection (i.p.)	PD-1 monoclonal antibody RB0005, 1 mg/kg	Day 3, 5, 7

9.3 Results

TABLE 6

Average tumor mass and tumor inhibitory rate in each group			
Groups	Tumor Weight (g)	IR _{TW} %	CDI (Coefficient of drug interaction)
G1	2.00	—	
G2	1.98	8.4	
G3	1.30	73.4	
G4	0.63	95.5	0.17

[0436] The results are shown in Table 6 and FIGS. 27 to 29:

[0437] The sequential administration group (G4) had a significant inhibitory effect on tumor proliferation, and during the observation period after administration, the tumor volume continued to decrease or even disappear. There was a statistically significant difference between the sequential administration group (G4) and the control group and

RB0005 single drug group (ANOVA, $P<0.001$), with a tumor weight inhibitory rate of 95.5%.

[0438] Sequential combination (firstly administering small molecule 2',3'-cGAMP, followed by macromolecule, i.e., PD-L1 monoclonal antibody) showed significant tumor inhibitory effect and exhibited drug synergistic effect, with $CDI=0.17$.

[0439] At the endpoint of the experiment, the survival rate of mice in the sequential combination administration group was 100%, the survival rate was significantly improved compared to the control group and the RB0005 single drug group.

[0440] In summary, the administration method of the combination of 2',3'-cGAMP and PD-L1 monoclonal antibody, where the small molecule was firstly administered, followed by the PD-L1 inhibitor, showed significant tumor inhibitory effect, improved survival rate of mice, and exhibited drug synergistic effects.

Example 10

[0441] The anti-tumor effect of a pharmaceutical combination composed of PD-L1 inhibitor and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

10.1 Xenograft Tumor Model in Mice

[0442] After successfully establishing a humanized CD274 cell subcutaneous xenograft tumor model of breast carcinoma CT26 in BALBc mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

10.2 Grouping

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
G2 trial group (Single dosage form)	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0005, 1 mg/kg + 2',3'-cGAMP monotherapy, 0.1 mg/kg	Administering every other day
G3 trial group (Single dosage form)	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0005, 1 mg/kg + 2',3'-cGAMP monotherapy, 0.316 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	PD-1 monoclonal antibody RB0005, 1 mg/kg + 2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day

10.3 Results

TABLE 7

Average tumor volume and tumor inhibitory rate in each group		
Groups	Tv (mm ³)	TGI (%)
G1	1499.1	—
G2	974.7	35.0
G3	868.0	42.1
G4	486.7	67.5
G5	142.5	90.5

[0443] The results are shown in Table 7 and FIGS. 30 to 33:

[0444] During the administration and observation period of the composite formulation of different doses of cGAMP and RB0005, the tumor proliferation was slow, wherein the tumor in the composite formulation group (G4) of different doses of cGAMP and LPG1501 (1 mpk) was decreased or even disappeared, the tumor volume at the endpoint of the experiment showed statistically significant difference compared to the control group ($p<0.05$);

[0445] In the solvent control group, the survival rate of mice on day 11 was 0, while the survival rates of mice in the composite formulation G3 and G4 groups at the endpoint of the experiment were above 50%.

[0446] In summary, the composite formulations of different doses of 2',3'-cGAMP and RB0005 (single dosage form) showed significant tumor inhibitory effect and improved survival rate ($P<0.05$).

Example 11

[0447] The anti-tumor effect of a pharmaceutical combination composed of PD-L1 inhibitor RB0005 and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

11.1 Xenograft Tumor Model in Mice

[0448] After successfully establishing a humanized CD274 cell subcutaneous xenograft tumor model of breast carcinoma 4T1 in BALBc mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

11.2 Grouping

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0005, 1 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0005, 1 mg/kg + 2',3'-cGAMP monotherapy, 1 mg/kg	Administering every other day

11.3 Results

TABLE 8

Average tumor mass and tumor inhibitory rate in each group			
Groups	Tumor Weight (g)	IR _{TW} %	CDI (Coefficient of drug interaction)
G1	0.96		
G2	0.99	—	
G3	0.63	34	
G4	0.60	37.5	0.92

[0449] The results are shown in Table 8 and FIGS. 34 to 36:

[0450] 4T1 model had poor responsiveness to RB0005 monotherapy, and thus RB0005 monotherapy had no significant tumor inhibitory effect;

[0451] cGAMP monotherapy had a certain inhibitory effect on tumor proliferation in 4T1 model, with a tumor weight inhibition ratio of 34%, but there was a great intra-group difference, and the survival rate at the endpoint of the experiment was 66.7%;

[0452] the tumor weight inhibitory rate of the composite formulation was 37.5%, which showed a statistically significant difference (P<0.05) compared to the control group and RB0005 single drug group, and exhibited drug synergistic effect as well, with a CDI value of less than 1 and a survival rate of 83.3% at the endpoint of the experiment;

[0453] In summary, the composite formulation of 2',3'-cGAMP and PD-L1 monoclonal antibody (single dosage form) had a certain tumor inhibitory effect, and improved survival rate in mice, and the responsiveness of RB0005 towards the 4T1 model had been improved by 2',3'-cGAMP.

Example 12

[0454] The anti-tumor effect of a pharmaceutical combination composed of PD-1 inhibitor RB0004 and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

12.1 Xenograft Tumor Model in Mice

[0455] After successfully establishing a humanized PD-1 subcutaneous xenograft tumor model of melanoma B16 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

12.2 Grouping

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0004, 10 mg/kg	Administering every other day

-continued

Groups	Administration Method	Administration dose	Administration frequency
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 0.1 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	LPG2009: D-1 monoclonal antibody RB0004, 10 mg/kg + 2',3'-cGAMP monotherapy, 0.1 mg/kg	Administering every other day

12.3 Results

TABLE 9

Average tumor volume and tumor inhibitory rate in each group			
Groups	Day 7-Tumor volume (mm ³)	TGI %	Q value (q ≥ 1, indicating that the combination of two drugs has a coefficient of drug interaction (CDI) of synergistic effect)
G1	1412.1	—	
G2	1849.9	—	
G3	578.8	59.0	
G4	213.4	84.9	1.8

[0456] The results are shown in Table 9 and FIGS. 37 to 40:

[0457] On day 7 after the administration of single drug cGAMP, the tumor inhibitory rate was 59%; the composite formulation (single dosage form) (G4) had a significant inhibitory effect on tumor proliferation. After administration, the tumor proliferation slowed down, and the tumor weight inhibitory rate on day 7 was 84.9%, there was statistically significant difference compared with the control group (P<0.001). The survival rate of mice at the endpoint of the experiment was 100%, showing a significant drug synergistic effect (King's Q value=1.8).

[0458] In summary, the single dosage form of 2',3'-cGAMP and PD-1 monoclonal antibody has a significant tumor inhibitory effect, high survival rate (P<0.05), and significant drug synergistic effects. The efficacy of the novel pharmaceutical combinations is better than that of PD-1 antibody RB0004 alone, indicating that the novel pharmaceutical combinations have greater advantages.

Example 13

[0459] The anti-tumor effect of a pharmaceutical combination composed of PD-1 inhibitor RB0004 and the STING agonist 2',3'-cGAMP was tested using a xenograft tumor model, i.e., the inhibitory effect on the growth of subcutaneous xenograft tumor in mice.

13.1 Xenograft Tumor Model in Mice

[0460] After successfully establishing a humanized PD-1 subcutaneous xenograft tumor model of lung carcinoma LLC-1 in mice, they were randomly grouped with 6-8 mice per group and administered once every two days, a total of 3 times.

13.2 Grouping

Groups	Administration Method	Administration dose	Administration frequency
G1 control group	Intratumor injection (i.t.)	Solvent	Administering every other day
G2 single drug group	Intratumor injection (i.t.)	D-1 monoclonal antibody RB0004, 10 mg/kg	Administering every other day
G3 single drug group	Intratumor injection (i.t.)	2',3'-cGAMP monotherapy, 0.1 mg/kg	Administering every other day
G4 trial group (Single dosage form)	Intratumor injection (i.t.)	LPG2009: D-1 monoclonal antibody RB0004, 10 mg/kg + 2',3'-cGAMP monotherapy, 0.1 mg/kg	Administering every other day

13.3 Results

TABLE 10

Average tumor inhibitory rate in each group		
Groups	TGI %	CDI (Coefficient of drug interaction)
G1	—	
G2	—	

TABLE 10-continued

Average tumor inhibitory rate in each group		
Groups	TGI %	CDI (Coefficient of drug interaction)
G3	47	
G4	72	0.55 (<1)

[0461] The results are shown in Table 10 and FIGS. 41 to 43:

[0462] RB0004 monotherapy had no significant tumor inhibitory effect on the model; cGAMP monotherapy had tumor inhibitory effect, with a tumor weight inhibitory rate of 47%;

[0463] The composite formulation (single dosage form) (G4) had an inhibitory effect on tumor proliferation. After administration, the tumor proliferation slowed down, and the tumor weight inhibitory rate was 72%, there was statistically significant difference compared with the control group ($P<0.001$), and showed a significant drug synergistic effect ($CDI=0.55$).

[0464] In summary, the single dosage form of 2',3'-cGAMP and PD-1 monoclonal antibody has a significant tumor inhibitory effect, high survival rate ($P<0.05$), and drug synergistic effects. The efficacy of the novel pharmaceutical combinations is better than that of PD-1 antibody RB0004 alone, indicating that the novel pharmaceutical combinations have greater advantages.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00004 HCDR1

<400> SEQUENCE: 1

Gly Phe Thr Phe Ser Ser Tyr Thr
 1 5

<210> SEQ ID NO 2
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00004 HCDR2

<400> SEQUENCE: 2

Ile Ser Asn Gly Gly Ser Phe Thr
 1 5

<210> SEQ ID NO 3
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00004 HCDR3

<400> SEQUENCE: 3

Ala Arg Asp Ser Asp Tyr Tyr Gly Ile Phe Asp Tyr

-continued

1	5	10
---	---	----

<210> SEQ ID NO 4
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 HFR1

<400> SEQUENCE: 4

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> SEQ ID NO 5
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 HFR2

<400> SEQUENCE: 5

Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala
1 5 10 15

Thr

<210> SEQ ID NO 6
<211> LENGTH: 38
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 HFR3

<400> SEQUENCE: 6

Tyr Tyr Pro Asp Ser Met Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn
1 5 10 15

Ser Lys Asn Thr Leu Tyr Leu Gln Met Ser Ser Leu Arg Ala Glu Asp
20 25 30

Thr Ala Val Tyr Tyr Cys
35

<210> SEQ ID NO 7
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 HFR4RB00005 HFR4

<400> SEQUENCE: 7

Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
1 5 10

<210> SEQ ID NO 8
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 VH

<400> SEQUENCE: 8

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly

-continued

1	5	10	15
Ser Leu Arg	Leu Ser Cys Ala Ala	Ser Gly Phe Thr	Phe Ser Ser Tyr
	20	25	30
Thr Met Ser	Trp Val Arg Gln Ala	Pro Gly Lys Gly	Leu Glu Trp Val
	35	40	45
Ala Thr Ile	Ser Asn Gly Gly	Ser Phe Thr Tyr Tyr	Pro Asp Ser Met
	50	55	60
Lys Gly Arg	Phe Thr Ile Ser Arg Asp	Asn Ser Lys Asn Thr	Leu Tyr
65	70	75	80
Leu Gln Met	Ser Ser Leu Arg Ala Glu	Asp Thr Ala Val Tyr	Tyr Cys
	85	90	95
Ala Arg Asp	Ser Asp Tyr Tyr Gly Ile	Phe Asp Tyr Trp Gly	Gln Gly
	100	105	110
Thr Thr Val	Thr Val Ser Ser		
	115		

<210> SEQ ID NO 9

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RB00004 HC

<400> SEQUENCE: 9

Glu Val Gln	Leu Val Glu Ser Gly	Gly Gly Leu Val Lys	Pro Gly Gly
1	5	10	15
Ser Leu Arg	Leu Ser Cys Ala Ala	Ser Gly Phe Thr	Phe Ser Ser Tyr
	20	25	30
Thr Met Ser	Trp Val Arg Gln Ala	Pro Gly Lys Gly	Leu Glu Trp Val
	35	40	45
Ala Thr Ile	Ser Asn Gly Gly	Ser Phe Thr Tyr Tyr	Pro Asp Ser Met
	50	55	60
Lys Gly Arg	Phe Thr Ile Ser Arg Asp	Asn Ser Lys Asn Thr	Leu Tyr
65	70	75	80
Leu Gln Met	Ser Ser Leu Arg Ala Glu	Asp Thr Ala Val Tyr	Tyr Cys
	85	90	95
Ala Arg Asp	Ser Asp Tyr Tyr Gly Ile	Phe Asp Tyr Trp Gly	Gln Gly
	100	105	110
Thr Thr Val	Thr Val Ser Ser Ala	Ser Thr Lys Gly	Pro Ser Val Phe
	115	120	125
Pro Leu Ala	Pro Cys Ser Arg Ser	Thr Ser Glu Ser	Thr Ala Ala Leu
	130	135	140
Gly Cys Leu	Val Lys Asp Tyr Phe	Pro Glu Pro Val	Thr Val Ser Trp
145	150	155	160
Asn Ser Gly	Ala Leu Thr Ser Gly	Val His Thr Phe	Pro Ala Val Leu
	165	170	175
Gln Ser Ser	Gly Leu Tyr Ser Leu	Ser Ser Val Val	Thr Val Pro Ser
	180	185	190
Ser Ser Leu	Gly Thr Lys Thr Tyr	Thr Cys Asn Val	Asp His Lys Pro
	195	200	205
Ser Asn Thr	Lys Val Asp Lys Arg	Val Glu Ser Lys	Tyr Gly Pro Pro
	210	215	220
Cys Pro Pro	Cys Pro Ala Pro Glu	Phe Leu Gly Gly	Pro Ser Val Phe

```

225              230              235              240
Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
      245              250              255

Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val
      260              265              270

Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
      275              280              285

Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val
      290              295              300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
305              310              315              320

Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser
      325              330              335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
      340              345              350

Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
      355              360              365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
      370              375              380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
385              390              395              400

Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp
      405              410              415

Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
      420              425              430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
      435              440              445

<210> SEQ ID NO 10
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LCDR1

<400> SEQUENCE: 10

Arg Ser Ser Lys Ser Leu Leu Tyr Lys Asp Gly Lys Thr Tyr Leu Asn
1      5      10      15

<210> SEQ ID NO 11
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LCDR2

<400> SEQUENCE: 11

Leu Met Ser Thr Arg Ala Ser
1      5

<210> SEQ ID NO 12
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LCDR3

<400> SEQUENCE: 12

```

-continued

Gln Gln Leu Val Glu Asp Pro Phe Thr
1 5

<210> SEQ ID NO 13
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LFR1

<400> SEQUENCE: 13

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys
20

<210> SEQ ID NO 14
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LFR2

<400> SEQUENCE: 14

Trp Tyr Leu Gln Lys Pro Gly Gln Ser Pro Gln Leu Leu Ile Tyr
1 5 10 15

<210> SEQ ID NO 15
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LFR3

<400> SEQUENCE: 15

Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15

Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys
20 25 30

<210> SEQ ID NO 16
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 LFR4

<400> SEQUENCE: 16

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val
1 5 10

<210> SEQ ID NO 17
<211> LENGTH: 115
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00004 VL

<400> SEQUENCE: 17

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu Tyr Lys

-continued

20	25	30
Asp Gly Lys Thr Tyr Leu Asn Trp Tyr Leu Gln Lys Pro Gly Gln Ser		
35	40	45
Pro Gln Leu Leu Ile Tyr Leu Met Ser Thr Arg Ala Ser Gly Val Pro		
50	55	60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile		
65	70	75
80		
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Leu		
85	90	95
Val Glu Asp Pro Phe Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
100	105	110
Arg Thr Val		
115		

<210> SEQ ID NO 18

<211> LENGTH: 230

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RB00004 LC

<400> SEQUENCE: 18

Ile Ala Leu Ser Leu Ala Leu Val Thr Asn Ser Asp Ile Val Met Thr		
1	5	10
15		
Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly Glu Pro Ala Ser Ile		
20	25	30
Ser Cys Arg Ser Ser Lys Ser Leu Leu Tyr Lys Asp Gly Lys Thr Tyr		
35	40	45
Leu Asn Trp Tyr Leu Gln Lys Pro Gly Gln Ser Pro Gln Leu Leu Ile		
50	55	60
Tyr Leu Met Ser Thr Arg Ala Ser Gly Val Pro Asp Arg Phe Ser Gly		
65	70	75
80		
Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile Ser Arg Val Glu Ala		
85	90	95
Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Leu Val Glu Asp Pro Phe		
100	105	110
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala		
115	120	125
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly		
130	135	140
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala		
145	150	155
160		
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln		
165	170	175
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser		
180	185	190
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr		
195	200	205
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser		
210	215	220
Phe Asn Arg Gly Glu Cys		
225	230	

-continued

<210> SEQ ID NO 19
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 HCDR1

<400> SEQUENCE: 19

Lys Tyr Ile Ile His
1 5

<210> SEQ ID NO 20
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 HCDR2

<400> SEQUENCE: 20

Trp Phe Tyr Pro Gly Ser Gly Asn Ile Arg Tyr Asn Glu Lys Ile Lys
1 5 10 15

Gly

<210> SEQ ID NO 21
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 HCDR3

<400> SEQUENCE: 21

His Gly Glu Leu Gly Gly Gly Tyr Phe Phe Asp Tyr
1 5 10

<210> SEQ ID NO 22
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 HFR1

<400> SEQUENCE: 22

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr
20 25 30

<210> SEQ ID NO 23
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 HFR2

<400> SEQUENCE: 23

Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met Gly
1 5 10

<210> SEQ ID NO 24
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 HFR3

-continued

<400> SEQUENCE: 24

Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr Met Glu
1 5 10 15
Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
20 25 30

<210> SEQ ID NO 25

<211> LENGTH: 121

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RB00005 VH

<400> SEQUENCE: 25

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Lys Tyr
20 25 30
Ile Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Trp Phe Tyr Pro Gly Ser Gly Asn Ile Arg Tyr Asn Glu Lys Ile
50 55 60
Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg His Gly Glu Leu Gly Gly Gly Tyr Phe Phe Asp Tyr Trp Gly
100 105 110
Gln Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 26

<211> LENGTH: 451

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RB00005 HC

<400> SEQUENCE: 26

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Lys Tyr
20 25 30
Ile Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Trp Phe Tyr Pro Gly Ser Gly Asn Ile Arg Tyr Asn Glu Lys Ile
50 55 60
Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg His Gly Glu Leu Gly Gly Gly Tyr Phe Phe Asp Tyr Trp Gly
100 105 110
Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
115 120 125

-continued

Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	
130						135					140					
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	
145					150					155					160	
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	
				165					170					175		
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	
			180					185					190			
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	
			195				200					205				
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	
	210					215					220					
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	
225					230					235					240	
Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	
				245					250					255		
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	
			260					265					270			
Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	
		275					280				285					
His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Ala	Ser	Thr	Tyr	
	290					295					300					
Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	
305					310					315					320	
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	
			325						330					335		
Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	
			340					345					350			
Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	
		355				360					365					
Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	
	370					375					380					
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	
385					390					395					400	
Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	
				405					410					415		
Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	
			420					425					430			
His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	
	435						440					445				
Pro	Gly	Lys														
	450															

<210> SEQ ID NO 27

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RB00005 LCDR1

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (5)..(5)

<223> OTHER INFORMATION: Xaa = Gly, Lys, or Ser

<220> FEATURE:

-continued

<221> NAME/KEY: misc_feature
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa = Ala, Thr, or Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa = Gly or Ser

<400> SEQUENCE: 27

Arg Ala Ser Ser Xaa Xaa Xaa Asn Ile His
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005a LCDR1

<400> SEQUENCE: 28

Arg Ala Ser Ser Ser Val Ser Asn Ile His
1 5 10

<210> SEQ ID NO 29
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005b LCDR1

<400> SEQUENCE: 29

Arg Ala Ser Ser Lys Thr Gly Asn Ile His
1 5 10

<210> SEQ ID NO 30
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005c LCDR1

<400> SEQUENCE: 30

Arg Ala Ser Ser Gly Ala Ser Asn Ile His
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 LCDR2

<400> SEQUENCE: 31

Ala Thr Ser Asn Leu Ala Ser
1 5

<210> SEQ ID NO 32
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 LCDR3

<400> SEQUENCE: 32

Gln Gln Trp Ser Ser Asn Pro Leu Thr

-continued

1 5

<210> SEQ ID NO 33
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 LFR1

<400> SEQUENCE: 33

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys
20

<210> SEQ ID NO 34
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 LFR2

<400> SEQUENCE: 34

Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr
1 5 10 15

<210> SEQ ID NO 35
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 LFR3

<400> SEQUENCE: 35

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Glu Phe Thr
1 5 10 15

Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys
20 25 30

<210> SEQ ID NO 36
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 LFR4

<400> SEQUENCE: 36

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
1 5 10

<210> SEQ ID NO 37
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005 VL
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (28)..(28)
<223> OTHER INFORMATION: Xaa = Gly, Lys, or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (29)..(29)
<223> OTHER INFORMATION: Xaa = Ala, Thr, or Val
<220> FEATURE:

-continued

<221> NAME/KEY: misc_feature
<222> LOCATION: (30)..(30)
<223> OTHER INFORMATION: Xaa = Gly or Ser

<400> SEQUENCE: 37

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Xaa Xaa Xaa Asn Ile
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr
35 40 45

Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> SEQ ID NO 38
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005a VL

<400> SEQUENCE: 38

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser Asn Ile
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr
35 40 45

Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> SEQ ID NO 39
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: RB00005b VL

<400> SEQUENCE: 39

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Lys Thr Gly Asn Ile
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr
35 40 45

-continued

Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
 50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
 65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
 85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
 100 105

<210> SEQ ID NO 40
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00005c VL

<400> SEQUENCE: 40

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Gly Ala Ser Asn Ile
 20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr
 35 40 45

Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
 50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu
 65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
 85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
 100 105

<210> SEQ ID NO 41
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00005 LC
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (28)..(28)
 <223> OTHER INFORMATION: Xaa = Gly, Lys, or Ser
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (29)..(29)
 <223> OTHER INFORMATION: Xaa = Ala, Thr, or Val
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (30)..(30)
 <223> OTHER INFORMATION: Xaa = Gly or Ser

<400> SEQUENCE: 41

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Xaa Xaa Xaa Asn Ile
 20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr
 35 40 45

Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser

-continued

50	55	60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu		
65	70	75 80
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr		
	85	90 95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro		
	100	105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr		
	115	120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys		
	130	135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu		
145	150	155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser		
	165	170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala		
	180	185 190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe		
	195	200 205
Asn Arg Gly Glu Cys		
210		

<210> SEQ ID NO 42

<211> LENGTH: 213

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: RB00005a LC

<400> SEQUENCE: 42

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly		
1	5	10 15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser Asn Ile		
	20	25 30
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr		
	35	40 45
Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser		
	50	55 60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu		
65	70	75 80
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr		
	85	90 95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro		
	100	105 110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr		
	115	120 125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys		
	130	135 140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu		
145	150	155 160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser		
	165	170 175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala		

-continued

180	185	190
Cys Glu Val Thr His Gln Gly	Leu Ser Ser Pro Val Thr Lys Ser Phe	
195	200	205

Asn Arg Gly Glu Cys
210

<210> SEQ ID NO 43
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00005b LC

<400> SEQUENCE: 43

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly	
1	15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Lys Thr Gly Asn Ile	
20	30
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro Trp Ile Tyr	
35	45
Ala Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser	
50	60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu	
65	80
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr	
85	95
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro	
100	110
Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr	
115	125
Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys	
130	140
Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu	
145	160
Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser	
165	175
Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala	
180	190
Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe	
195	205

Asn Arg Gly Glu Cys
210

<210> SEQ ID NO 44
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: RB00005c LC

<400> SEQUENCE: 44

Asp Ile Gln Leu Thr Gln Ser Pro Ser Phe Leu Ser Ala Ser Val Gly	
1	15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Gly Ala Ser Asn Ile	
20	30

-continued

His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Pro	Trp	Ile	Tyr
		35					40					45			
Ala	Thr	Ser	Asn	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
	50					55					60				
Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu
65				70					75					80	
Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp	Ser	Ser	Asn	Pro	Leu	Thr
			85						90					95	
Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro
			100					105					110		
Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr
		115					120					125			
Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys
	130					135					140				
Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu
145				150					155					160	
Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser
			165					170						175	
Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala
			180				185						190		
Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe
		195					200					205			
Asn	Arg	Gly	Glu	Cys											
		210													

What is claimed is:

1. A pharmaceutical combination comprising a programmed cell death protein 1 (PD-1) inhibitor and/or a programmed death ligand 1 (PD-L1) inhibitor, and a STING pathway agonist.

2. The pharmaceutical combination according to claim 1, wherein the STING pathway agonist comprises a cyclic dinucleotide, 2',3'-cGAMP, or derivatives thereof, flavonoids, DNA or type I interferons (IFNs);

preferably, the cyclic dinucleotide is selected from the group consisting of c-di-AMP, c-di-GMP, c-di-GMP-F, 3',3'-cGAMP, 3',3'-cGAMP-F, 2',3'-cGAMP, Rp/Sp (CL656), ADU-S100, ADU-S100 disodium, and derivatives or combinations thereof;

the flavonoid comprises CMA, DMXAA, methoxyflavone, 6,4'-dimethoxyflavone, 4'-methoxyflavone, 3',6'-dihydroxyflavone, 7,2'-dihydroxyflavone, daidzein, Formononetin, retinene 7-methyl ether, xanthone, and/or any combination thereof;

the type I interferon comprises IFN- α or IFN- β .

3. (canceled)

4. (canceled)

5. (canceled)

6. (canceled)

7. (canceled)

8. (canceled)

9. (canceled)

10. The pharmaceutical combination according to claim 1, wherein the PD-1 inhibitor has one or more of the following characteristics: a. inhibition or reduction of PD-L1 expression, such as transcription or translation of PD-L1; b. inhibition or reduction of PD-1 activity, such as inhibition or

reduction of PD-1 binding to its homologous ligands, such as PD-L1 or PD-L2; and c. binding PD-1 or one or more of its ligands, such as PD-L1 or PD-L2; or

the PD-1 inhibitor comprises an anti PD-1 antibody or antigen-binding fragments thereof;

preferably, the anti PD-1 antibody is selected from the group consisting of Pembrolizumab, Nivolumab, Pidilizumab, SHR-1210, MEDI0680, BGB-A317 TSR-042, REGN2810, PF-06801591, RB0004, Tislelizumab, Camrelizumab, Toripalimab, Sintilimab, bioanalogues, bio-enhancers, bio-equivalents, or combinations thereof.

more preferably, the anti PD-1 antibody comprises at least one CDR in the antibody heavy chain variable region (VH), and the VH comprises an amino acid sequence shown in SEQ ID NO: 8;

more preferably, the anti PD-1 antibody comprises VH comprising HCDR3, and the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 3;

more preferably, the VH further comprises HCDR2, wherein the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 2;

more preferably, the VH further comprises HCDR1, wherein the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 1;

further preferably, the VH comprises HCDR1, HCDR2, and HCDR3, wherein the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 3, the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 1.

11. (canceled)

12. (canceled)

13. (canceled)

14. (canceled)

15. (canceled)

16. (canceled)

17. (canceled)

18. The pharmaceutical combination according to claim 10, wherein the VH comprises a framework region HFR1, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, and the HFR1 comprises an amino acid sequence shown in SEQ ID NO: 4 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 4;

preferably, the VH comprises a framework region HFR2, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2; and the HFR2 comprises an amino acid sequence shown in SEQ ID NO: 5 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 5;

preferably, the VH comprises a framework region HFR3, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3; and the HFR3 comprises an amino acid sequence shown in SEQ ID NO: 6 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 6;

preferably, the VH comprises a framework region HFR4, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3, and the HFR4 comprises an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7;

more preferably, the VH comprises framework regions HFR1, HFR2, HFR3, and HFR4, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3; among them, the HFR1 comprises an amino acid sequence shown in SEQ ID NO: 4 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 4, the HFR2 comprises an amino acid sequence shown in SEQ ID NO: 5 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 5, the HFR3 comprises an amino acid sequence shown in SEQ ID NO: 6 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 6, and the HFR4 comprises an amino acid sequence shown in SEQ ID NO: 7 or an amino acid

sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

19. (canceled)

20. (canceled)

21. (canceled)

22. (canceled)

23. The pharmaceutical combination according to claim 10, wherein the anti PD-1 antibody comprises VH, and the VH comprises an amino acid sequence shown in SEQ ID NO: 8;

preferably, the anti PD-1 antibody comprises antibody heavy chain (HC), and the HC comprises an amino acid sequence shown in SEQ ID NO: 9;

preferably, the anti PD-1 antibody comprises at least one CDR in the antibody light chain variable region (VL), and the VL comprises an amino acid sequence shown in SEQ ID NO: 17;

preferably, the anti PD-1 antibody comprises at least one CDR in VH, the VH comprises an amino acid sequence shown in SEQ ID NO: 8, and the anti PD-1 antibody comprises at least one CDR in VL, and the VL comprises an amino acid sequence shown in SEQ ID NO: 17.

24. (canceled)

25. (canceled)

26. (canceled)

27. The pharmaceutical combination according to claim 10, wherein the anti PD-1 antibody comprises VL comprising LCDR1, and the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 10;

preferably, the VL further comprises LCDR2, wherein the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 11;

preferably, the VL further comprises LCDR3, wherein the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 12;

preferably, the VL comprises LCDR1, LCDR2 and LCDR3, wherein the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 10, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 12.

28. (canceled)

29. (canceled)

30. (canceled)

31. The pharmaceutical combination according to claim 10, wherein the anti PD-1 antibody comprises VH and antibody VL, the VH comprises HCDR1, HCDR2, and HCDR3, wherein the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 3, the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 2, and the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 1; and the VL comprises LCDR1, LCDR2, and LCDR3, wherein the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 10, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 11, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 12.

32. The pharmaceutical combination according to claim 27, wherein the VL comprises a framework region LFR1, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, and the LFR1 comprises an amino acid sequence shown in SEQ ID NO: 13 or an amino

acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 13;

preferably, the VL comprises a framework region LFR2, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2; and the LFR2 comprises an amino acid sequence shown in SEQ ID NO: 14 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 14;

preferably, the VL comprises a framework region LFR3, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3; and the LFR3 comprises an amino acid sequence shown in SEQ ID NO: 15 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 15;

preferably, the VL comprises a framework region LFR4, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3, and the LFR4 comprises an amino acid sequence shown in SEQ ID NO: 16 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 16.

33. (canceled)

34. (canceled)

35. (canceled)

36. The pharmaceutical combination according to claim 27, wherein the VL comprises framework regions LFR1, LFR2, LFR3 and LFR4, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3; among them, the LFR1 comprises an amino acid sequence shown in SEQ ID NO: 13 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 13, the LFR2 comprises an amino acid sequence shown in SEQ ID NO: 14 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 14, the LFR3 comprises an amino acid sequence shown in SEQ ID NO: 15 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 15, and the LFR4 comprises an amino acid sequence shown in SEQ ID NO: 16 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 16.

37. The pharmaceutical combination according to claim 10, wherein the anti PD-1 antibody comprises VL, and the VL comprises an amino acid sequence shown in SEQ ID NO: 17;

preferably, the anti PD-1 antibody comprises VH and VL, the VH comprises an amino acid sequence shown in SEQ ID NO: 8 and the VL comprises an amino acid sequence shown in SEQ ID NO: 17;

preferably, the anti PD-1 antibody comprises antibody light chain (LC), and the LC comprises an amino acid sequence shown in SEQ ID NO: 18;

preferably, the anti PD-1 antibody comprises HC and LC, the HC comprises an amino acid sequence shown in SEQ ID NO: 9 and the LC comprises an amino acid sequence shown in SEQ ID NO: 18.

38. (canceled)

39. (canceled)

40. (canceled)

41. The pharmaceutical combination according to claim 1, wherein the PD-L1 inhibitor has one or more of the following characteristics: a. inhibition or reduction of PD-L1 expression, such as transcription or translation of PD-L1; b. inhibition or reduction of PD-L1 activity, such as inhibition or reduction of PD-L1 binding to its associated receptors such as PD-1; and c. binding of PD-L1 or its receptors such as PD-1;

preferably, the PD-L1 inhibitor comprises an anti PD-L1 antibody or antigen-binding fragments thereof;

preferably, the anti PD-L1 antibody is selected from the group consisting of Durvalumab, Atezolizumab, Envafoimab, Avelumab, MDX-1105, YW243.55.S70, MDPL3280A, AMP-224, LY3300054, RB0005, bio-analogues, bio-enhancers, bio-equivalents, or combinations thereof.

42. (canceled)

43. (canceled)

44. The pharmaceutical combination according to claim 41, wherein the anti PD-L1 antibody comprises at least one CDR in VH, and the VH comprises an amino acid sequence shown in SEQ ID NO: 25;

preferably, the anti PD-L1 antibody comprises VH comprising HCDR3, and the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 21;

preferably, the VH further comprises HCDR2, wherein the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 20;

preferably, the VH further comprises HCDR1, wherein the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 19;

more preferably, the VH comprises HCDR1, HCDR2, and HCDR3, wherein the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 19.

45. (canceled)

46. (canceled)

47. (canceled)

48. (canceled)

49. The pharmaceutical combination according to claim 44, wherein the VH comprises a framework region HFR1, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, and the HFR1 comprises an amino acid sequence shown in SEQ ID NO: 22 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 22;

preferably, the VH comprises a framework region HFR2, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2; and the HFR2 comprises an amino acid sequence shown in SEQ ID NO: 23 or an

amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 23;

preferably, the VH comprises a framework region HFR3, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3; and the HFR3 comprises an amino acid sequence shown in SEQ ID NO: 24 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 24;

preferably, the VH comprises a framework region HFR4, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3, and the HFR4 comprises an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7;

more preferably, the VH comprises framework regions HFR1, HFR2, HFR3 and HFR4, the C-terminal of HFR1 is directly or indirectly connected to the N-terminal of HCDR1, the N-terminal of HFR2 is directly or indirectly connected to the C-terminal of HCDR1, and the C-terminal of HFR2 is directly or indirectly connected to the N-terminal of HCDR2, the N-terminal of HFR3 is directly or indirectly connected to the C-terminal of HCDR2, and the C-terminal of HFR3 is directly or indirectly connected to the N-terminal of HCDR3, the N-terminal of HFR4 is directly or indirectly connected to the C-terminal of HCDR3; among them, the HFR1 comprises an amino acid sequence shown in SEQ ID NO: 22 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 22, the HFR2 comprises an amino acid sequence shown in SEQ ID NO: 23 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 23, the HFR3 comprises an amino acid sequence shown in SEQ ID NO: 24 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 24, and the HFR4 comprises an amino acid sequence shown in SEQ ID NO: 7 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 7.

50. (canceled)

51. (canceled)

52. (canceled)

53. (canceled)

54. The pharmaceutical combination according to claim 41, wherein the anti PD-L1 antibody comprises VH, and the VH comprises an amino acid sequence shown in SEQ ID NO: 25;

preferably, the anti PD-L1 antibody comprises HC, and the HC comprises an amino acid sequence shown in SEQ ID NO: 26;

preferably, the anti PD-L1 antibody comprises at least one CDR in VL, and the VL comprises an amino acid sequence shown in SEQ ID NO: 37;

preferably, the anti PD-L1 antibody comprises at least one CDR in VH, the VH comprises an amino acid sequence shown in SEQ ID NO: 25, and the anti PD-L1 antibody

comprises at least one CDR in VL, and the VL comprises an amino acid sequence shown in SEQ ID NO: 37;

preferably, the anti PD-L1 antibody comprises at least one CDR in VH, the VH comprises an amino acid sequence shown in SEQ ID NO: 25, and the anti PD-L1 antibody comprises at least one CDR in VL, and the VL comprises an amino acid sequence shown in SEQ ID NO: 38, SEQ ID NO: 39 or SEQ ID NO: 40.

55. (canceled)

56. (canceled)

57. (canceled)

58. (canceled)

59. The pharmaceutical combination according to claim 41, wherein the anti PD-L1 antibody comprises VL, the VL comprises LCDR1, and the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 27-;

preferably, the anti PD-L1 antibody comprises VL, the VL comprises LCDR1, and the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 28, SEQ ID NO: 29 or SEQ ID NO: 30;

preferably, the VL further comprises LCDR2, wherein the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 31;

preferably, the VL further comprises LCDR3, wherein the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32;

preferably, the VL comprises LCDR1, LCDR2 and LCDR3, wherein the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 27, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32.

60. (canceled)

61. (canceled)

62. (canceled)

63. (canceled)

64. The pharmaceutical combination according to claim 59, wherein the VL comprises LCDR1, LCDR2 and LCDR3, wherein the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 28, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32; the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 29, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 31, the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32; or the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 30, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32;

preferably, the anti PD-L1 antibody comprises VH and antibody VL, the VH comprises HCDR1, HCDR2, and HCDR3, wherein the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 19; and the VL comprises LCDR1, LCDR2, and LCDR3, wherein the LCDR1 comprises an amino acid sequence shown in SEQ ID NO: 27, the LCDR2 comprises an amino acid

sequence shown in SEQ ID NO: 31, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32;

preferably, the anti PD-L1 antibody comprises VH and antibody VL, the VH comprises HCDR1, HCDR2, and HCDR3, wherein the HCDR3 comprises an amino acid sequence shown in SEQ ID NO: 21, the HCDR2 comprises an amino acid sequence shown in SEQ ID NO: 20, and the HCDR1 comprises an amino acid sequence shown in SEQ ID NO: 19; and the VL comprises LCDR1, LCDR2, and LCDR3, wherein the LCDR1 comprises an amino acid sequence shown in SEQ ID NO:28, SEQ ID NO:29 or SEQ ID NO:30, the LCDR2 comprises an amino acid sequence shown in SEQ ID NO: 31, and the LCDR3 comprises an amino acid sequence shown in SEQ ID NO: 32.

65. (canceled)

66. (canceled)

67. The pharmaceutical combination according to claim 59, wherein the VL comprises a framework region LFR1, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, and the LFR1 comprises an amino acid sequence shown in SEQ ID NO: 33 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 33;

preferably, the VL comprises a framework region LFR2, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2; and the LFR2 comprises an amino acid sequence shown in SEQ ID NO: 34 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 34;

preferably, the VL comprises a framework region LFR3, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3; and the LFR3 comprises an amino acid sequence shown in SEQ ID NO: 35 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 35;

preferably, the VL comprises a framework region LFR4, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3, and the LFR4 comprises an amino acid sequence shown in SEQ ID NO: 36 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 36;

more preferably, the VL comprises framework regions LFR1, LFR2, LFR3 and LFR4, the C-terminal of LFR1 is directly or indirectly connected to the N-terminal of LCDR1, the N-terminal of LFR2 is directly or indirectly connected to the C-terminal of LCDR1, and the C-terminal of LFR2 is directly or indirectly connected to the N-terminal of LCDR2, the N-terminal of LFR3 is directly or indirectly connected to the C-terminal of LCDR2, and the C-terminal of LFR3 is directly or indirectly connected to the N-terminal of LCDR3, the N-terminal of LFR4 is directly or indirectly connected to the C-terminal of LCDR3; among them, the LFR1 comprises an amino acid sequence shown in SEQ ID NO: 33 or an amino acid sequence having at least about

70% sequence identity to the amino acid sequence shown in SEQ ID NO: 33, the LFR2 comprises an amino acid sequence shown in SEQ ID NO: 34 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 34, the LFR3 comprises an amino acid sequence shown in SEQ ID NO: 35 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO: 35, and the LFR4 comprises an amino acid sequence shown in SEQ ID NO: 36 or an amino acid sequence having at least about 70% sequence identity to the amino acid sequence shown in SEQ ID NO:36.

68. (canceled)

69. (canceled)

70. (canceled)

71. (canceled)

72. The pharmaceutical combination according to claim 41, wherein the anti PD-L1 antibody comprises VL, and the VL comprises an amino acid sequence shown in SEQ ID NO: 37;

preferably, the anti PD-L1 antibody comprises VL, and the VL comprises an amino acid sequence shown in SEQ ID NO:38, SEQ ID NO:39 or SEQ ID NO:40;

preferably, the anti PD-L1 antibody comprises VH and VL, the VH comprises an amino acid sequence shown in SEQ ID NO: 25, and the VL comprises an amino acid sequence shown in SEQ ID NO: 37;

preferably, the anti PD-L1 antibody comprises VH and VL, the VH comprises an amino acid sequence shown in SEQ ID NO: 25, and the VL comprises an amino acid sequence shown in SEQ ID NO:38, SEQ ID NO:39 or SEQ ID NO:40;

preferably, the anti PD-L1 antibody comprises LC, and the LC comprises an amino acid sequence shown in SEQ ID NO: 41;

preferably, the anti PD-L1 antibody comprises LC, and the LC comprises an amino acid sequence shown in SEQ ID NO:42, SEQ ID NO:43 or SEQ ID NO:44;

preferably, the anti PD-L1 antibody comprises HC and LC, the HC comprises an amino acid sequence shown in SEQ ID NO: 26, and the LC comprises an amino acid sequence shown in SEQ ID NO: 41;

preferably, the anti PD-L1 antibody comprises HC and LC, the HC comprises an amino acid sequence shown in SEQ ID NO: 26, and the LC comprises an amino acid sequence shown in SEQ ID NO:42, SEQ ID NO:43 or SEQ ID NO:44.

73. (canceled)

74. (canceled)

75. (canceled)

76. (canceled)

77. (canceled)

78. (canceled)

79. (canceled)

80. The pharmaceutical combination according to claim 1, wherein i) the PD-1 inhibitor and/or PD-L1 inhibitor in the pharmaceutical combination are not mixed with ii) the STING pathway agonist to each other in the pharmaceutical combination;

preferably, i) the PD-1 inhibitor and/or PD-L1 inhibitor, and ii) the STING pathway agonist are present in the pharmaceutical combination in a single dosage form;

preferably, the pharmaceutical combination is formulated into a pharmaceutical composition;
 preferably, the pharmaceutical composition comprises a PD-1 inhibitor or a PD-L1 inhibitor, and a STING pathway agonist;
 preferably, the STING pathway agonist is present in an amount of about 0.0001 mg/kg to about 200 mg/kg;
 preferably, the PD-1 inhibitor or PD-L1 inhibitor is present in an amount of about 0.0001 mg/kg to about 200 mg/kg;
 preferably, the pharmaceutical composition further comprises one or more pharmaceutically acceptable carriers.

81. (canceled)
 82. (canceled)
 83. (canceled)
 84. (canceled)
 85. (canceled)
 86. (canceled)
 87. (canceled)
 88. (canceled)
 89. (canceled)
 90. (canceled)

91. A method for treating neoplastic diseases, comprising administering an effective amount of the pharmaceutical combination according to claim 1;
 preferably, the subject suffers from neoplasm;
 more preferably, the neoplasm comprises tumors and/or warts;
 preferably, the administration comprises local, intraneoplastic or systemic administrations;
 more preferably, the administration comprises intravenous injection, intravenous instillation, intramuscular injection, subcutaneous injection, and/or intraneoplastic injection;
 preferably, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered by use of the same or different administration routes;
 preferably, the method comprises injection of the STING pathway agonist into the neoplasm;
 more preferably, the method further comprises injection or systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm;
 preferably, the method comprises injection of the STING pathway agonist and systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm;
 preferably, the method comprises injecting the i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination into the neoplasm;

preferably, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered simultaneously or at different times;
 preferably, the PD-1 inhibitor or PD-L1 inhibitor is administered before and/or after the administration of the STING pathway agonist;
 preferably, the PD-1 inhibitor or PD-L1 inhibitor is administered after the administration of the STING pathway agonist;
 preferably, the method comprises: i) injection of the STING pathway agonist into the neoplasm; ii) injection or systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor into the neoplasm after administering the STING pathway agonist;
 preferably, the method comprises: i) injection of the STING pathway agonist into the neoplasm; and ii) systemic infusion of the PD-1 inhibitor or PD-L1 inhibitor after administering the STING pathway agonist;
 more preferably, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are administered simultaneously;
 more preferably, i) the PD-1 inhibitor or PD-L1 inhibitor, and ii) the STING pathway agonist in the pharmaceutical combination are simultaneously administered by way of intraneoplastic injection, and the PD-1 inhibitor or PD-L1 inhibitor and the STING pathway agonist are present in a same dosage form.

92. (canceled)
 93. (canceled)
 94. (canceled)
 95. (canceled)
 96. (canceled)
 97. (canceled)
 98. (canceled)
 99. (canceled)
 100. (canceled)
 101. (canceled)
 102. (canceled)
 103. (canceled)
 104. (canceled)
 105. (canceled)
 106. (canceled)
 107. (canceled)
 108. (canceled)

* * * * *