



US 20250283102A1

(19) **United States**(12) **Patent Application Publication**
KWON et al.(10) **Pub. No.: US 2025/0283102 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **METHOD FOR PRODUCING POTATO
PLANT WITH SUPPRESSED BROWNING
USING CRISPR/CAS9 SYSTEM***C12N 9/22* (2006.01)*C12N 15/11* (2006.01)(52) **U.S. Cl.**CPC *C12N 15/8243* (2013.01); *C12N 9/0059*
(2013.01); *C12N 9/222* (2025.05); *C12N*
15/111 (2013.01); *C12N 15/8201* (2013.01);
C12N 2310/20 (2017.05); *C12Y 110/03001*
(2013.01)(71) Applicant: **TOOLGEN INCORPORATED**, Seoul
(KR)(72) Inventors: **Soon Il KWON**, Seoul (KR); **Chee
Hark HARN**, Seoul (KR); **Ye Rim
LEE**, Seoul (KR); **Ba Reum KIM**,
Seoul (KR)(21) Appl. No.: **18/035,363**(22) PCT Filed: **Nov. 22, 2021**(86) PCT No.: **PCT/KR2021/017138**

§ 371 (c)(1),

(2) Date: **May 4, 2023**(30) **Foreign Application Priority Data**

Nov. 25, 2020 (KR) 10-2020-0160474

Publication Classification(51) **Int. Cl.***C12N 15/82* (2006.01)*C12N 9/02* (2006.01)

(57)

ABSTRACT

A method for producing a potato plant with suppressed browning may use CRISPR/Cas9 system. In a method of producing a genome-edited potato plant with suppressed browning, a potato protoplast is transfected with a ribonucleoprotein complex containing sgRNA which targets StPPO2 (*Solanum tuberosum* Polyphenol Oxidase 2) gene derived from potato. Since the method involves no insertion of a foreign gene and includes only tiny mutation that is hardly distinguishable from natural mutation, it is expected that, unlike genetically modified organism (GMO) crops which require a huge amount of cost and time for evaluating the safety and negative environmental impacts, a great deal of cost and time can be saved in the present invention.

Specification includes a Sequence Listing.**Target gene : StPPO2****Target site : TCCATGGATGAAAAGTTGAGAGG (SEQ ID NO: 2) (-)***** Result summary**

sgRNA		Total reads	Insertions	Deletions	Indel frequency(%)
StPPO2-1	Control	28,597	0	0	0
	Sample	33,616	0	2	2(0)

Green: RCEN

Red : Insertion sequence

Yellow shade : Mismatch sequence

ATC : Start codon

*** Sequence analysis data**CCACTTCTTTATCTTCCACTCCTAAGCCCTCTCAACTTTTCATCCATGGAAAACGTAACCAAATGTTCAAAG (SEQ ID No: 6) **Target sequence****Most frequent sequences****Type****Reads**CCACTTCTTTATCTTCCACTCCTAAGCCCTCTCAACTTTTCATCCATGGAAAACGTAACCAAATGTTCAAAG (SEQ ID No: 6) **wild-type** **33614**CCACTTCTTTATCTTCCACTCCTAAGCCCTCTC-AC TTTTCATCCATGGAAAACGTAACCAAATGTTCAAAG (SEQ ID No: 7) **1 del** **2****Total****33616**

FIG. 1

Target gene : StPPO2

Target site : TCCATGGATGAAAAGTTGAGAGG (SEQ ID NO: 2) (-)

* Result summary

sgRNA		Total reads	Insertions	Deletions	Indel frequency(%)
StPPO2-1	Control	28,597	0	0	0
	Sample	33,616	0	2	2(0)

Green: ROSE
Red : Insertion sequence
Yellow shade : Mismatch sequence
ATG : Start codon

* Sequence analysis data

Target sequence	SEQ ID No: 6)	Type	Reads
CCACTTCTTTATCTTCCACTCCCTAAGCCCCTCTCAACTTTTCATCCATGGAACGTAACCCAAATGTTCAAAG	6)	wild-type	33614
CCACTTCTTTATCTTCCACTCCCTAAGCCCCTCTCAACTTTTCATCCATGGAACGTAACCCAAATGTTCAAAG	7)	1 del	2
Total			33616

FIG. 2

Target gene : StPPO2
Target site : TGCATGAAACTTTGAACATTG (SEQ ID NO: 3) (~)

* Result summary

sgRNA		Total reads	Insertions	Deletions	Indel frequency(%)
StPPO2-2	Control	27,593	0	0	0(0)
	Sample	46,021	0	28	28(0)

Green: RGEN
Red : Insertion sequence
Yellow shade : Mismatch sequence
ATG : Start codon

* Sequence analysis data

Target sequence	Type	Reads
TCYCAACTTTTCATCCATGGAAACGTAAACCAATGTTCAAGTTTCATGCAAGGTTACCAATAATAACGG (SEQ ID No: 8)	Wild-type	45993
TCYCAACTTTTCATCCATGGAAACGTAAACCAATGTTCAAGTTTCATGCAAGGTTACCAATAATAACGG (SEQ ID No: 8)	1 del	28
TCYCAACTTTTCATCCATGGAAACGTAAACC-ATGTTCAAGTTTCATGCAAGGTTACCAATAATAACGG (SEQ ID No: 9)	Total	46021

FIG. 3

Target gene : StPPO2
Target site : ATACTACAAGACGAGAGATCAGG (SEQ ID NO: 4) (-)

* Result summary

sgRNA		Total reads	Insertions	Deletions	Indel frequency(%)
StPPO2-3	Control	53,787	0	0	0(0)
	Sample	34,035	0	0	0(0)

Green: BGR
Red : Insertion sequence
Yellow shade : Mismatch sequence
ATG : Start codon

* Sequence analysis data

Target sequence	Type	Reads
GCTGCAATCCGCTGCTCCAGCTCCACCTCCGATCTCTCGTCTGTAGTATAGCCAGGATTAAACGAAAAATCAG (SEQ ID No: 10)	Wild-type	34035
Total		34035

Most frequent sequences

GCTGCAATCCGCTGCTCCAGCTCCACCTCCGATCTCTCGTCTGTAGTATAGCCAGGATTAAACGAAAAATCAG

FIG. 4

Target gene : STPPO2

Target site : TTAGGCGGCAACAACGTGTACGG (SEQ ID NO: 5) (-)

* Result summary

igbba		Total reads	insertions	Deletions	Indel frequency(%)
STPPO2-4	Control	8,842	0	0	0(0)
	Sample	12,844	0	244	244(1.9%)

Green: KGEN
Red : Insertion sequence
Yellow shade : Mismatch sequence
ATG : Start codon

* Sequence analysis data

Most frequent sequences	Target sequence	Type	Reads
CCAGGATTAAACGAAATCAGGTGGTGGCCGTACAGTTGTTGGCGCCCTAAGCCTGATGATATGGAGAAAGTTTC (SEQ ID No: 11)			
CCAGGATTAAACGAAATCAGGTGGTGGCCGTACAGTTGTTGGCGCCCTAAGCCTGATGATATGGAGAAAGTTTC (SEQ ID No: 11)		Wild-type	12680
CCAGGATTAAACGAAATCAGGTGGTGGCCGTAC- - - - -GTTGGCGCCCTAAGCCTGATGATATGGAGAAAGTTTC (SEQ ID No: 12)		4 del	94
CCAGGATTAAACGAAATCAGGTGGTGGCCGTACA- - -TGTTGGCGCCCTAAGCCTGATGATATGGAGAAAGTTTC (SEQ ID No: 13)		2 del	73
CCAGGATTAAACGAAATCAGGTGGTGGCCGTACA- - - - -TGGCGCCCTAAGCCTGATGATATGGAGAAAGTTTC (SEQ ID No: 14)		5 del	40
CCAGGATTAAACGAAATCAGGTGGTGGCCGTACA- - - - -GCGCGCCCTAAGCCTGATGATATGGAGAAAGTTTC (SEQ ID No: 15)		6 del	37
Target site			Total
			12844

FIG. 5C




FIG. 5B




FIG. 5A



FIG. 6




FIG. 7A

Desiree #14

Target gene : StPPO2 (sgRNA : StPPO2-4)

Target site : TTAGGGGGGCAACAACGTACTACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
37,368	23	0	37,345	37,345(100%)

GGATTAAACGAAATCAGGTGGTGCCCGTACAGTTGTTGGCGGCTAATCCCTGATGATAATGGAGAAAGTTC (SEQ ID NO: 16) WT

GGATTAAACGAAATCAGGTGGTGCCCGTACA--TGTTCGCGCGCTAATCCCTGATGATAATGGAGAAAGTTC (SEQ ID NO: 17) 2bp del (x 37,101)

GGATTAAACGAAATCAGGTGGTGCCCGTAC-GTTGTTGGCGGCTAATCCCTGATGATAATGGAGAAAGTTC (SEQ ID NO: 18) 1bp del (x 38)

GGATTAAACGAAACTCATGTGGTGCCCGTAC----GTTGCGCGGCTAATCCCTGATGATAATGGAGAAAGTTC (SEQ ID NO: 19) 4bp del (x 24)

GGATTAAACGAAATCAGGTGGTGCCCGTACA-----GCCCTAATCCCTGATGATAATGGAGAAAGTTC (SEQ ID NO: 20) 10bp del (x 2)

Target site

FIG. 7B

Desiree #25

Target gene : StPPO2 (sgRNA : StPPO2-4)

Target site : TTAGGGCGGCAACAACGTACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
43,381	6	0	43,375	43,375(99.9%)

GGATTACGAAATCAGGTGGTGGCCGTTACAGTTGTGGCGGCTAAATGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 16) WT

GGATTACGAAATCAGGTGGTGGCCGTTACAA--TGTTCGGCGGCTAAATGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 17) 2bp del (x 38,099)

GGATTACGAAATCAGGTGGTGGCCGTTAC----GTTTCGGCGGCTAAATGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 19) 4bp del (x 5,212)

GGATTACGAAATCAGGTGGTGGCCGTTAC-GTTGTTCGGCGGCTAAATGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 18) 1bp del (x 60)

GGATTACGAAATCAGGTGGTGGCCGTTAC-----GCCTAAATGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 20) 10bp del (x 6)

Target site

FIG. 8A

Desiree #75

Target gene : StPPO2 (sgRNA : StPPO2-4)

Target site : TTAGGCGGCGCAACAAC**TGTACGG** (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
26,859	6	0	26,853	26,853(100%)

GGATTAAACGAAATCAGGTGGTGGCC**CTACAGT**TGTTGGCGCC**TATG**CCCTGATGATATGGAGAAAGTTTC (SEQ ID NO: 16) WT

GGATTAAACGAAACTCATGTGGTGGCC**CTAC**----GTTGGCGGCC**TATG**CCCTGATGATATGGAGAAAGTTTC (SEQ ID NO: 19) 4bp del (x 7,152)

GGATTAAACGAAATCAGGTGGTGGCC**CTAC**----TTGGCGGCC**TATG**CCCTGATGATATGGAGAAAGTTTC (SEQ ID NO: 21) 5bp del (x 7,011)

GGATTAAACGAAATCAGGTGGTGGCC**CTAC**----TGTTCGCGCC**TATG**CCCTGATGATATGGAGAAAGTTTC (SEQ ID NO: 22) 3bp del (x 6,174)

GGATTAAACGAAATCAGGTGGTGGCC**CTACA**--TTGGCGGCC**TATG**CCCTGATGATATGGAGAAAGTTTC (SEQ ID NO: 17) 2bp del (x 6,454)

GGATTAAACGAAATCAGGTGGTGGCC**CTAC**-GTTGTTGGCGCC**TATG**CCCTGATGATATGGAGAAAGTTTC (SEQ ID NO: 18) 1bp del (x 62)

Target site

FIG. 8B

Desiree #153

Target gene : STPPO2 (sgRNA : StPPO2-4)
Target site : TTAGGCGCGCAACAACTGTACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
74,069	4	0	74,065	74.065(100%)

GGATTACGAAATCAGGTGGTGGCCGTTACAGTTGTTGGCGGCTTAACCTGATGATATGGAGAAAGTTC (SEQ ID NO: 16) WT

GGATTACGAAACATCATGTGGTGCCCTTAC-----GTTCGCGGCTTAACCTGATGATATGGAGAAAGTTC (SEQ ID NO: 19) 4bp del (x 34,251)

GGATTACGAAATCAGGTGGTGGCCCTTAC-----TTGCGGCTTAACCTGATGATATGGAGAAAGTTC (SEQ ID NO: 21) 5bp del (x 31,985)

GGATTACGAAATCAGGTGGTGGCCCTTACA---TGTTCGCGGCTTAACCTGATGATATGGAGAAAGTTC (SEQ ID NO: 17) 2bp del (x 7,629)

GGATTACGAAATCAGGTGGTGGCCCTTAC-GTTGTTGGCGGCTTAACCTGATGATATGGAGAAAGTTC (SEQ ID NO: 18) 1bp del (x 102)

GGATTACGAAATCAGGTGGTGGCCCTTACA-----GCCCTTAACCTGATGATATGGAGAAAGTTC (SEQ ID NO: 20) 10bp del (x 8)

Target site

FIG. 9A

Desiree #205

Target gene : StPPO2 (sgRNA : StPPO2-4)

Target site : TTAGGCGCGCAACAACGTACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
23,185	0	0	23,185	23,185(100%)




FIG. 9B

Desiree #237

Target gene : StPPO2 (sgRNA : StPPO2-4)
Target site : TTAGGCGGCGCAACAACGTGACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	Insertions	deletions	In-del frequency
35,275	0	0	35,275	35,275(100%)

GGATTAAACGAAATCAGGTGGTGGCCCTTACAGTTGTTGCGGCGCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 16) WT

GGATTAAACGAAACTCATGTGGTGGCCCTTAC-----GTTGCGGCGGCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 19) 4bp del (x 9,018)

GGATTAAACGAAATCAGGTGGTGGCCCTTAC-----TTGCGGCGGCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 21) 5bp del (x 8,825)

GGATTAAACGAAATCAGGTGGTGGCCCTTACA--TGTTGCGGCGGCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 17) 2bp del (x 8,604)

GGATTAAACGAAATCAGGTGGTGGCCCTTAC-GTTGTTGCGGCGGCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 18) 1bp del (x 8,579)

GGATTAAACGAAATCAGGTGGTGGCCCTTACA-----GCCCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 20) 10bp del (x 175)

GGATTAAACGAAATCAGGTGGTGGCCCTTAC--TGTTGCGGCGGCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID NO: 22) 3bp del (x 64)

Target site

FIG. 10A

Desiree #494

Target gene : STPPO2 (sgRNA : STPPO2-4)

Target site : TTAGGGCGCGCAACAACTGTACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
37,183	0	0	37,183	37,183(100%)

GGATTACCGAAATCAGGTGGTGGCCCTACAGTTGTTGCGCGCCCTAGGCCTGATGATATGGAGAAAGTTC (SEQ ID No: 16) WT

GGATTACCGAAATCAGGTGGTGGCCCTACAGTTGTTGCGCGCCCTAGGCCTGATGATATGGAGAAAGTTC (SEQ ID No: 19) 4bp del (x 31,222)

GGATTACCGAAATCAGGTGGTGGCCCTACAGTTGTTGCGCGCCCTAGGCCTGATGATATGGAGAAAGTTC (SEQ ID No: 17) 2bp del (x 5,896)

GGATTACCGAAATCAGGTGGTGGCCCTACAGTTGTTGCGCGCCCTAGGCCTGATGATATGGAGAAAGTTC (SEQ ID No: 18) 1bp del (x 65)

Target site

FIG. 10B

Desiree #519

Target gene : StPPO2 (sgRNA :StPPO2-4)
Target site : TTAGGGGGCAACAAGTACGG (SEQ ID NO: 5)

* Result summary

Total reads	WT	insertions	deletions	In-del frequency
29,202	0	0	29,202	29,202(100%)

GGATTACGAAATCAGGTGGTGGCCCTACAGTTGTTGGGGCCCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID No: 16) WT

GGATTACGAAACTCATGTGGTGGCCCTAC-----GTTGGGGCCCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID No: 19) 4bp del (x 21,998)

GGATTACGAAATCAGGTGGTGGCCCTAC-----TTGGGGCCCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID No: 21) 5bp del (x 7,173)

GGATTACGAAATCAGGTGGTGGCCCTAC-----GCCCTAAGCCCTGATGATATGGAGAAAGTTC (SEQ ID No: 20) 10bp del (x 31)

Target site

FIG. 11




FIG. 12

PPO2 activity assay

Protein sample preparation

Protein concentration (Bradford)

Number	Samples	Absorbance (595 nm)
1	#30	0.6115
2	#16	0.5735
3	#14	0.4499
4	WT	0.5692
5	#25	0.5664

Samples	0.2 M Catechol	Absorbance (410 nm)				Unit/ul	Relative activity (% to Control)
		0 sec	30 sec	60 sec	90 sec	120 sec	
#30	+	0.3357	0.4020	0.4456	0.4697	0.4837	70.2%
#16	+	0.3274	0.3940	0.4318	0.4539	0.4653	71.4%
#14	+	0.2594	0.3033	0.3253	0.3400	0.3470	64.5%
WT	+	0.3954	0.4745	0.5269	0.5638	0.5862	-
#25	+	0.1652	0.2026	0.2252	0.2364	0.2456	42.2%

FIG. 13A





FIG. 13B



METHOD FOR PRODUCING POTATO PLANT WITH SUPPRESSED BROWNING USING CRISPR/CAS9 SYSTEM

CROSS REFERENCE TO RELATED APPLICATIONS AND CLAIM OF PRIORITY

[0001] This application claims benefit under 35 U.S.C. 119, 120, 121, or 365(c), and is a National Stage entry from International Application No. PCT/KR2021/017138 filed on Nov. 22, 2021, which claims priority to the benefit of Korean Patent Application No. 10-2020-0160474 filed in the Korean Intellectual Property Office on Nov. 25, 2020, the entire contents of which are incorporated herein by reference.

BACKGROUND

1. Technical Field

[0002] The present invention relates to a method for producing a potato plant with suppressed browning using CRISPR/Cas9 system.

2. Background Art

[0003] In recent years, the trend in consumer food product purchasing indicates a shift of preference to natural products and healthy foods, and simultaneously, due to ever increasing one-person and nuclear family households, there is increasing demand for minimally processed crops which guarantee the freshness and convenience. As the minimally processed crops are provided after being subjected to processes like washing, peeling, cutting, or the like during production stage, they are advantageous in that the consumers can, after making a purchase, use them directly without carrying out a pre-treatment process or the like.

[0004] Potato is one of the minimally processed crops. However, in accordance with the air exposure of surface and breakdown of cell wall during the processing, enzymatic browning easily occurs in potato.

[0005] Main cause of the enzymatic browning in a plant including potato is PPO (polyphenol oxidase) protein, which is an enzyme containing Cu^{2+} . The polyphenol compounds present in the cell are converted to L-DOPA (L-3,4-dihydroxyphenylalanine) according to oxidization by PPO. L-DOPA is subsequently converted to DOPA quinone and forms, after an oxidative polymerization, a melanin pigment with dark brown color to exhibit the browning phenomenon.

[0006] The browning reaction is known to cause undesirable color and smell of minimally processed crops and also to lower the nutritional value of the crops. In addition, as the product undergone browning during storage and transport of minimally processed crops has a low commercial value to yield economic loss, it is quite necessary to suppress the browning.

[0007] The genome editing technique in recent years suggests a possibility of inducing genetic mutation at target genomic DNA site by using site-specific nucleases of various types. In particular, in relation with plants and other various organisms, CRISPR/Cas9 system is known to be a potent tool for genome editing. The advantage of CRISPR/Cas9 system is that it allows accurate and efficient introduction of mutation to a target site. It is also shown that the genome edited crops are not different from the crops which have been developed by a traditional breeding technique.

[0008] Meanwhile, in Japanese Patent Application Publication No. 2019-507595, "Plant showing a reduced wound-induced surface discoloration" is disclosed, and, in Korean Patent Application Publication No. 2016-0087651, "Browning prevented colored potato and method for preventing browning of colored potato" is disclosed. However, "Method for producing a potato plant with suppressed browning using CRISPR/Cas9 system" has not been described before.

SUMMARY

[0009] The present invention is devised under the circumstances that are described in the above. To suppress the browning by editing StPPO2 (*Solanum tuberosum* Polyphenol Oxidase 2) gene using CRISPR system instead of common cross breeding or gene modification method, inventors of the present invention introduced RNP complex (StPPO2 sgRNA+Cas9) by PEG-mediated transfection after isolating protoplast from young germ-free potato plant obtained by in vitro culture, and eventually selected a plant with edited StPPO2 gene based on cell reproduction and generation of the protoplasts which have been introduced with the complex. Further, by choosing a potato line with suppressed activity of PPO2 protein and suppressed browning from the gene-edited plants obtained after the above selection, the inventors accordingly completed the present invention.

[0010] To achieve the object described in the above, the present invention provides a composition for genome editing to suppress browning in a potato plant comprising, as an active ingredient, a ribonucleoprotein complex of a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and an endonuclease protein; or a recombination vector comprising a DNA encoding a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and a nucleic acid sequence encoding an endonuclease protein.

[0011] The present invention further provides a guide nucleic acid which is an RNA sequence specific to a target region in the nucleotide sequence constituting StPPO2 gene, in which the target region is the nucleotide sequence of SEQ ID NO: 5.

[0012] The present invention further provides a method of producing a genome-edited potato plant with suppressed browning including steps of (a) introducing a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and an endonuclease protein to a potato plant cell to have genome editing; and (b) regenerating a potato plant from the potato plant cell that is genome-edited.

[0013] The present invention still further provides a genome-edited potato plant with suppressed browning which is produced by the aforementioned method, and genome-edited seed potato thereof.

[0014] The genome-edited potato plant with suppressed browning produced by the method of the present invention has delayed browning even when it is previously cut or mashed, and thus it can be conveniently used for processing or cooking and additional processes like heating or adding an artificial additive for suppressing the browning can be minimized. In addition, as the method of the present invention involves no insertion of a foreign gene and includes only tiny mutation that is hardly distinguishable from natural mutation, it is expected that, unlike GMO (Genetically Modified Organism) crops which require a huge amount of

cost and time for evaluation of the safety and negative environmental impacts, a great deal of cost and time can be saved in the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] FIG. 1 to FIG. 4 show the result of the target deep-sequencing in in vitro assay which has been carried out for determining candidate sgRNA.

[0016] FIGS. 5A to 5C show the separation of protoplast from the leaf of a young potato plant, in which FIG. 5A shows VCP treatment of the leaf of a young potato plant, FIG. 5B shows the intact protoplast (indicated with red arrow) which has been determined by separation of the VCP-treated protoplast based on sucrose concentration gradient, and FIG. 5C shows a photographic image of the protoplast under microscope which has been finally separated.

[0017] FIG. 6 shows the photographic image of a plant regenerated from a potato protoplast which has been introduced with CRISPR/Cas9 RNP complex.

[0018] FIG. 7A to FIG. 10B show the result of the deep-sequencing of several plants from the plants regenerated from a protoplast which has been introduced with CRISPR/Cas9 RNP complex.

[0019] FIG. 11 shows the result of StPPO2 editing efficiency in the potato protoplast obtained according to the method of the present invention (left column), and the InDel pattern of StPPO2 edited product derived from the potato protoplast (right column).

[0020] FIG. 12 shows the result of measuring the activity of PPO2 protein in which the measurement was carried out by using potato tuber.

[0021] FIGS. 13A and 13B show the result of observing the browning of potato tuber, in which FIG. 13A represents the result of comparing, after slicing potato tuber, the control (WT) with the gene edited product (GE #14, #25), and FIG. 13B represents the result of comparing, after removing only the skin of potato tuber, the control (WT) with the gene edited product (GE #14, #25).

DETAILED DESCRIPTION

[0022] To achieve the object of the present invention, the present invention provides a composition for genome editing to suppress browning in a potato plant comprising, as an active ingredient, a ribonucleoprotein complex of a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and an endonuclease protein; or a recombination vector comprising a DNA encoding a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and a nucleic acid sequence encoding an endonuclease protein.

[0023] In an embodiment of the composition of the present invention, the target gene for genome editing to suppress the browning in potato plant is StPPO2 gene which consists of the nucleotide sequence of SEQ ID NO: 1.

[0024] As described herein, the term “genome/gene editing” means a technique for introducing a target-oriented mutation to a genome nucleotide sequence of a plant or an animal cell including human cell. Specifically, it indicates a technique for knock-out or knock-in of a specific gene by deletion, insertion, or substitution of one more nucleic acid molecules by DNA cutting, or a technique for introducing a mutation even to a non-coding DNA sequence which does

not produce any protein. According to the purpose of the present invention, the genome editing may be an introduction of a mutation to a plant by using an endonuclease, for example, Cas9 (CRISPR associated protein 9) protein, and a guide RNA. Furthermore, the term “gene editing” may be interchangeably used with “gene engineering”.

[0025] Furthermore, the term “target gene” means part of DNA present in the genome of a plant to be edited according to the present invention. Type of the target gene is not limited, and it may include a coding region and also a non-coding region. Depending on the purpose, a person who is skilled in the pertinent art may select the target gene according to the mutation of a plant desired to be produced by genome editing.

[0026] Furthermore, the term “guide RNA” means a ribonucleic acid which includes RNA specific to a target DNA in nucleotide sequence encoding the target gene and, according to complementary binding between the whole or partial sequence of the guide RNA and the nucleotide sequence of target DNA, the guide RNA plays the role of guiding an endonuclease to the target DNA nucleotide sequence. The guide RNA indicates RNA in single guide RNA (sgRNA) form comprising the first site which includes a dual RNA having two RNAs, i.e., crRNA (CRISPR RNA) and tracrRNA (trans-activating crRNA), as a constitutional element; or a sequence which is fully or partially complementary to the nucleotide sequence in the target gene, and the second site which includes a sequence interacting with an endonuclease (in particular, RNA-guide nuclease), but those in the form in which the endonuclease is active for the target nucleotide sequence are also within the scope of the invention without any limitation. Considering the type of an endonuclease to be used together, microorganisms from which the endonuclease is derived, or the like, the guide RNA may be used after being prepared in consideration of a method well known in the pertinent art.

[0027] Furthermore, the guide RNA may be those transcribed from a plasmid template, those obtained by in vitro transcription (e.g., oligonucleotide double strand), or those obtained by synthesis, or the like, but it is not limited thereto.

[0028] In an embodiment of the composition for genome editing of the present invention, the guide RNA is devised to have specificity for the target nucleotide sequence in StPPO2 gene consisting of the nucleotide sequence of SEQ ID NO: 5, in which the guide RNA devised to have specificity for the nucleotide sequence of StPPO2 gene is characterized in that it has higher efficiency of inducing InDel mutation compared to other guide RNAs for editing StPPO2 gene that are devised to have specificity for the target nucleotide sequence of SEQ ID NO: 2 to SEQ ID NO: 4. In particular, because potato is a tetraploid crop, when CRISPR/Cas system is applied to potato by using a RNA guide having high InDel efficiency, it is likely to have higher efficiency of obtaining individual potato plant in which all four alleles are edited.

[0029] Furthermore, in an embodiment of the composition for genome editing of the present invention, the endonuclease protein may be one or more selected from a group consisting of Cas9, Cpf1 (CRISPR from *Prevotella* and *Francisella* 1), TALEN (Transcription activator-like effector nuclease), ZFN (Zinc Finger Nuclease), and a functional analog thereof. It may be preferably Cas9 protein, but it is not limited thereto.

[0030] Furthermore, the Cas9 protein may be one or more selected from a group consisting of Cas9 protein derived from *Streptococcus pyogenes*, Cas9 protein derived from *Campylobacter jejuni*, Cas9 protein derived from *S. thermophilus*, Cas9 protein derived from *S. aureus*, Cas9 protein derived from *Neisseria meningitidis*, Cas9 protein derived from *Pasteurella multocida*, Cas9 protein derived from *Francisella novicida*, and the like, but it is not limited thereto. Information of Cas9 protein or gene of Cas9 protein can be obtained from a known database like GenBank of NCBI (National Center for Biotechnology Information).

[0031] Cas9 protein is an RNA-guided DNA endonuclease enzyme which induces breakage of a double stranded DNA. In order for Cas9 protein to cause DNA breakage after precise binding to a target nucleotide sequence, a short nucleotide sequence consisting of three nucleotides, which is known as PAM (Protospacer Adjacent Motif), should be present next to the target nucleotide sequence, and Cas9 protein causes the breakage by assuming the position between the 3rd and the 4th base pairs from the PAM sequence (NGG).

[0032] In an embodiment of the composition for genome editing of the present invention, the guide RNA and endonuclease protein may function as RNA gene scissors (RNA-Guided Engineered Nuclease, RGEN) by forming a ribonucleic acid-protein (i.e., ribonucleoprotein) complex.

[0033] The present invention further provides a guide nucleic acid specific to a target region in the nucleotide sequence constituting StPPO2 gene, in which the target region is the nucleotide sequence of SEQ ID NO: 5.

[0034] The “guide nucleic acid” according to the present invention means the aforementioned “guide RNA”, and it includes an endonuclease binding site in addition to the RNA sequence specific to the target region. The endonuclease may be preferably an RNA-guide nuclease, but it is not limited thereto.

[0035] The guide nucleic acid of the present invention may be preferably in the form of a single guide RNA (sgRNA), but it is not limited thereto.

[0036] According to the guide nucleic acid of one embodiment of the present invention, the RNA sequence specific to a target region may be synthesized by replacing adenine (A), thymine (T), or cytosine (C) of the target sequence present at the 20th position from the PAM nucleotide with guanine (G), but it is not limited thereto.

[0037] The guide nucleic acid according to the present invention is characterized in that, when an insertion-deletion (InDel) mutation is induced in a target sequence of StPPO2 gene consisting of the nucleotide sequence of SEQ ID NO: 5, deletion mutation with size of several base pairs (i.e., 1 to 10 bp) is caused. Thus, in terms of the genome mutation, it has relatively higher safety than a guide RNA which may cause an insertion or a deletion mutation with size of several hundred base pairs.

[0038] The present invention further provides a method of producing a genome-edited potato plant with suppressed browning including (a) introducing a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and an endonuclease protein to a potato plant cell to have genome editing; and (b) regenerating a potato plant from the potato plant cell that is genome-edited.

[0039] According to the production method of one embodiment of the present invention, the guide RNA spe-

cific to the target nucleotide sequence in StPPO2 gene and endonuclease protein are as defined in the above.

[0040] The CRISPR/Cas9 system employed in the present invention is a method of gene editing based on NHEJ (non-homologous end joining) mechanism in which insertion-deletion (InDel) mutation resulting from incomplete repair, which is induced during a process of DNA repair, by introducing breakage of a double strand at specific site of a specific gene to be edited.

[0041] In an embodiment of the production method of the present invention, the aforementioned step (a) for introducing a guide RNA and an endonuclease protein to a potato plant cell may use a ribonucleoprotein complex of a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and an endonuclease protein; or a recombination vector comprising a DNA encoding a guide RNA specific to the target nucleotide sequence in StPPO2 gene derived from potato and a nucleic acid sequence encoding an endonuclease protein, but it is not limited thereto.

[0042] In an embodiment of the production method of the present invention, the transfection method for introducing the complex of a guide RNA and an endonuclease protein to a plant cell may be appropriately selected from a calcium/polyethylene glycol method for protoplasts (Krens, F. A. et al., 1982, Nature 296, 72-74; Negrutiu I. et al., June 1987, Plant Mol. Biol. 8, 363-373), an electroporation method for protoplasts (Shillito R. D. et al., 1985 Bio/Technol. 3, 1099-1102), a microscopic injection method for plant components (Crossway A. et al., 1986, Mol. Gen. Genet. 202, 179-185), a (DNA or RNA-coated) particle bombardment method for various plant components (Klein T. M. et al., 1987, Nature 327, 70), or a (non-complete) viral infection method in *Agrobacterium tumefaciens* mediated gene transfer by plant invasion, or the like.

[0043] Furthermore, the introduction of a recombination vector comprising a DNA encoding a guide RNA specific to the target nucleotide sequence and a nucleic acid sequence encoding an endonuclease protein means transformation. Transformation of a plant species is now generally carried out for all plant species including not only a dicot plant but also a monocot plant. In principle, any method for transformation may be used for introducing the recombination vector of the present invention to a suitable progenitor cell.

[0044] Furthermore, in an embodiment of the production method of the present invention, the “plant cell” to which a guide RNA specific to the target nucleotide sequence and an endonuclease are introduced may be any type of a plant cell. The plant cell may be a cultured cell, a cultured tissue, a cultured organ, or a whole plant. The “plant tissue” is either a generated tissue or a non-generated tissue of a plant, for example, root, stem, leaf, pollen, small spore, egg cell, seed, and cells in various form that are used for culture, i.e., single cell, protoplast, shoot, and callus tissue, although it is not limited thereto. The plant tissue may be an in planta tissue or in the form of cultured organ, cultured tissue, or cultured cell. The plant cell preferred in the present invention is a protoplast.

[0045] In an embodiment of the production method of the present invention, the potato plant cell of step (b) which is obtained after genome editing may be a cell in which deletion mutation of 1 to 10 bp in size has been induced in the target nucleotide sequence in StPPO2 gene, but it is not limited thereto.

[0046] In an embodiment of the production method of the present invention, any method well known in the pertinent art can be used as a method of regenerating a genome-edited plant from a plant cell with edited genome. A plant cell with edited genome needs to be regenerated to a whole plant. Techniques for regeneration into a mature plant from culture of callus or protoplast are well known in the pertinent art for various types of plants (Handbook of Plant Cell Culture, volume 1 to 5, 1983-1989 Momillan, N.Y.).

[0047] The present invention still further provides a genome-edited potato plant with suppressed browning which is produced by the aforementioned method, and genome-edited seed potato thereof.

[0048] In the genome-edited potato plant with suppressed browning of the present invention, StPPO2 gene involved in browning is edited by using CRISPR/Cas9 system, and it is a genome-edited potato plant which has, due to the knock-out of StPPO2 gene derived from potato, a trait with suppressed browning compared to a potato plant without any genome editing.

[0049] In particular, the genome-edited potato plant of the present invention may have a deletion mutation of 1 to 10 bp in size that is induced in the target nucleotide sequence in StPPO2 gene, and it is characterized by having, in terms of the genome mutation, relatively higher safety than a genome-edited plant in which an insertion or a deletion mutation of several hundred base pairs in size has been made.

[0050] As described herein, the term “StPPO2” may be interchangeably used with the term “StuPPO2”.

[0051] Hereinbelow, the present invention is explained in greater detail in view of the Examples. However, the following Examples are given only for exemplification of the present invention and the scope of the present invention is not limited by them.

EXAMPLES

Example 1. Selection of StPPO2 Gene sgRNA

[0052] Four different sgRNAs were selected from the website of CRISPR RGEN Tools by using the sequence of potato StPPO2 gene (Table 1). The sgRNAs described in the following Table 1 have been selected by considering various conditions such as GC contents, out-of-frame score, mismatch number in genome sequence, and the like. In the present invention, the nucleotide sequences of SEQ ID NO: 2 to 5 represent the sequence of RGEN target sequence of the following Table 1 excluding the PAM base (underlined).

TABLE 1									
sgRNA Target sequence for editing potato StPPO2 gene									
sgRNA	(SEQ ID NO)	RGEN target* (5'→3')	Direc- tion	PAM	GC content (%)	Out- of- w/o frame	Mis- matches	score	
StPPO2-1	TCCATGGATGAAAAGTTG	AGAGG	—	40	91.2	0	2	1	2
(SEQ ID NO: 2)									

TABLE 1-continued									
sgRNA Target sequence for editing potato StPPO2 gene									
sgRNA	(SEQ ID NO)	RGEN target* (5'→3')	Direc- tion	PAM	GC content (%)	Out- of- w/o frame	Mis- matches	score	
StPPO2-2	TGCATGAAACTTTGAACA	TTTGG	—	30	75.0	2	0	0	0
(SEQ ID NO: 3)									
StPPO2-3	ATACTACAAGACGAGAGA	TCAGG	—	40	79.1	0	0	0	2
(SEQ ID NO: 4)									
StPPO2-4	TTAGGCGCGCAACAACTG	TACGG	—	50	72.5	1	0	0	0
(SEQ ID NO: 5)									

*underlined; PAM base

[0053] After synthesizing each of the four selected sgRNAs against target sequence, in vitro cleavage assay and in vivo assay were carried for their determination. In vitro cleavage assay is an assay for examining the activity of sgRNA against target DNA in which a test tube containing PCR fragments of target DNA is added with a sample not containing any Cas9 and sgRNA (sample 1), a sample containing Cas9 and sgRNA at ratio of 1:1 (sample 2), or a sample containing Cas9 and sgRNA at ratio of 1:3 (sample 3) followed by reaction and the reaction product is analyzed by electrophoresis. When sgRNA is used for the analysis acts on the target RNA, the target DNA is broken to exhibit at least 2 bands. On the other hand, when sgRNA does not act on the target RNA, no breakage product is identified.

[0054] Furthermore, in vivo assay is an assay for determining InDel efficiency in which each of the selected sgRNAs is introduced, together with Cas9, by PEG-mediated transfection to protoplast cells (5×10⁵), and after 3 days or so (i.e., time for RNP [Cas9+sgRNA] complex to have a reaction with the target gene in nucleus), genomic DNA is isolated from the protoplast and target deep-sequencing is carried out by PCR.

[0055] Four sgRNAs selected by in vivo assay are found to have InDel efficiency of 0% for StPPO2-1 to StPPO2-3, and 1.9% for StPPO2-4 (FIGS. 1 to 4). Based on these assay results, the inventors of the present invention finally selected StPPO2-4 (SEQ ID NO: 5) as a target sequence of sgRNA for StPPO2 gene editing.

Example 2. Separation of Potato Protoplast and Method of Culturing Potato Protoplast

[0056] To a petri dish, VCP (Viscozyme, Celluclast, PectinEX) enzyme solution (10 ml) was added. About 5 to 7 leaves of young potato plant of Desiree variety (*Solanum tuberosum* L. cv. Desiree) which has been obtained by in vitro culture were cut and added to the petri dish such that the backside of leaf faces is on the top. Then, the reaction was allowed to occur for 5 hours or so at 25° C. and stirring speed of 45 rpm under dark conditions. After making an observation of the degree of enzyme degradation, when it is found that the leaf tissues turn to be transparent, separation of protoplast was carried out. Once the VCP enzyme reac-

tion is completed, the mixture was filtered using a disposable mesh (Falcon, 40 μm), and the resultant was transferred to a 14 ml round tube and centrifuged for 5 minutes at 800 rpm. After the centrifuge, the supernatant was discarded. W5 solution (Menczel L. et al. (1981) Theor. Appl. Genet. 59; 191-195) was added (1 ml) and the protoplast was carefully dissolved. After adding slowly 9 ml of the W5 solution thereto, the resultant was centrifuged for 5 minutes at 800 rpm to have first washing. By repeating the same procedure as above, a second washing was carried out. Thereafter, the supernatant was discarded and 5 ml of the W5 solution was slowly added, and the resultant was carefully transferred to a 15 ml round tube which has been added with 20% sucrose (5 ml) and centrifuged for 5 minutes at 800 rpm. After the centrifuge, a protoplast layer (i.e., green band) was seen near the middle part of the tube while the broken protoplast or part of the tissues was found to be settled at the bottom of the tube. Using a blunt-end pipette tip, the protoplast layer was carefully extracted, which was then transferred to a fresh 15 ml round tube added with 10 ml of the W5 solution, and centrifuged for 5 minutes at 800 rpm. The supernatant was discarded after the centrifuge and the protoplast was dissolved well by adding 1 ml of the W5 solution. Then, the state of the protoplast was observed under a microscope (FIGS. 5A to 5C).

[0057] If the separated protoplast is in good state, it was added to a 6-well plate with PIM (Protoplast Inducing Media, 3 ml), and the protoplast and cell division pattern was observed every day. Cell division is generally observed after 5 days or so. After 2 weeks or so, cell accumulation starts to appear so that microcalli (observed with a naked eye, less than 1 mm) are observed after 3 weeks or so. After 4 weeks or so, it is possible to observe with a naked eye white spots in culture using 1.2% low melting agar.

TABLE 2

Composition of solution used for protoplast separation			
[Method for preparing CPW solution]			
Composition	Stock name	1X CPW concentration (Final Conc.)	
CPW salts	$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	Stock B	1480 mg/L
	KH_2PO_4	Stock A	27.2 mg/L
	KNO_3		101 mg/L
	$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$		246 mg/L
	KI		0.16 mg/L
	$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$		0.025 mg/L
pH 5.7			

Stock A solution was prepared as 100X solution and used after dilution to appropriate concentration

Stock B solution was prepared as 10X solution and used after dilution to appropriate concentration

[Method for preparing VCP solution]	
Composition	Content (for preparing 500 ml)
CPW stock A (X 100)	5 ml
CPW stock B (X 10)	50 ml
Mannitol	45 g
MES buffer (A)	533 mg
Viscozyme (B)	5 ml
Celluclast (C)	2.5 ml
PectinEX	2.5 ml
pH 5.7	

Prepared solution was subjected to sterile filtration at 0.22 μm pore size without any autoclave, and stored in refrigerator until use

Example 3. Gene Editing of Potato Protoplast Using CRISPR/Cas9 RNP Complex

[0058] The protoplast separated by the method described in Example 2 was counted using a hemocytometer under a microscope. About 100,000 protoplasts were used for the test of CRISPR/Cas9 gene editing.

[0059] After completing the protoplast separation, CRISPR/Cas9 RNP complex was produced as follows. First, 15 μg of StPPO2-4 sgRNA (sgRNA of 5 $\mu\text{g}/\mu\text{l}$ concentration was used in an amount of 3 μl) and 30 μg of Cas9 (Cas9 protein of 5 $\mu\text{g}/\mu\text{l}$ concentration was used in an amount of 6 μl) were added to a 2 ml tube, added with 1 μl of NEB buffer (#3), and then reacted for 10 minutes at room temperature. Thereafter, CRISPR/Cas9 RNP complex was introduced to the prepared protoplast using PEG-mediated transfection. Namely, the protoplast (1×10^5 , 200 μl) was admixed with CRISPR/Cas9 RNP complex (10 μA), and after adding 40% PEG solution in the same volume as the mixture (i.e., 210 μl), the reaction was allowed to occur for 10 minutes at room temperature. Upon the completion of the reaction after 10 minutes, the W5 solution in the same volume as the mixture (i.e., 420 μl) was added for having first washing followed by centrifuge for 5 minutes at 600 rpm. The supernatant was discarded after the centrifuge. After second washing by adding again the W5 solution (1 ml) to remove residual PEG, centrifuge was carried out at the same condition as above. During the washing, PIM was aliquoted in advance in an amount of 3 ml to a 6-well plate. To the protoplast from which the supernatant has been removed after the centrifuge, 1 ml PIM solution was added and mixed well. The mixture was then aliquoted in an amount of 500 μl to each well of the 6-well plate followed by sealing. Then, culture was carried out at 25° C., dark condition and the state of protoplast was examined.

[0060] Once the protoplast shows relatively stable division after 2 weeks or so, it was added with low melting agar with concentration of 1.2% followed by solidification. Then, observation was made under culture at the same condition. After 3 weeks or so following the introduction of CRISPR/Cas9 RNP complex, microcalli starts to appear and the microcalli under culture in low melting agar/PIM were transferred to CFM (Callus Forming Media) and continuously cultured therein. Callus under culture in CFM (diameter of about 0.5 cm) was transferred to SIM (Shoot Inducing Media) and culture therein. To induce roots from a plant with shoots, transfer to RIM (Root Inducing Media) was made followed by culture therein. It was found that, after 16 weeks or so, the protoplast introduced with CRISPR/Cas9 RNP complex is fully grown to a single plant (FIG. 6). Meanwhile, with regard to the composition of the media including PIM, CFM, SIM, and RIM which have been used for protoplast culture described above, reference was made to the report by Ehsanpour and Jones (J. Sci. I. R. Iran. (2001), 12 (2): 103-110).

TABLE 3

Composition of PEG solution for transfection	
40% PEG Solution	For preparing 2.5 ml
PEG 4000	1 g
1M Mannitol	0.5 ml
1M CaCl_2	0.25 ml
ddH_2O	To be 2.5 ml (about 1 ml or so)

After addition to a 14 ml round tube and adjustment to 2.5 ml, it was placed in a microwave for 20 seconds, mixed well to have full dissolving, cooled down at room temperature, filtered, and used.

Fresh PEG solution was used by preparing it at each time of the experiment

Example 4. Selection of StPPO2 Edited Product
Derived from Potato Protoplast

[0061] For 590 plants of the total 640 plants obtained from Example 3, editing of StPPO2 gene was examined by deep-sequencing. As a result, it was finally found that the gene editing is induced in at least one allele in 110 plants while 3 plants show that the 4 alleles are edited in all different InDel patterns. FIGS. 7A to 10B illustrate the result of deep-sequencing of some of those plants. The result of deep-sequencing of the total 110 plants, which have been finally found to have edited StPPO2 gene, is the same as those described in Table 4.

TABLE 4

Result of deep-sequencing of plants which have been found to have edited StPPO2 gene						
Target gene	Sample	Total reads	WT	Inser- tions	Dele- tions	In-del frequency (%)
StPPO2-4	14	37368	23	0	37345	37345 (100%)
	16	41402	22	0	41380	41380 (99.9%)
	24	47184	91	0	47093	47093 (99.9%)
	25	43381	6	0	43375	43375 (99.9%)
	30	21108	73	0	21035	21035(99.8%)
	31	25462	19	0	25443	25443 (99.9%)
	38	16599	0	0	16599	16599 (100%)
	50	16485	0	0	16485	16485 (100%)
	51	21638	0	0	21638	21638 (100%)
	52	11835	0	0	11835	11835 (100%)
	62	32061	0	0	32059	32059 (100%)
	69	31237	15	0	31222	31222(99.9%)
	75	26859	6	0	26853	26853(100%)
	83	45162	19	0	45143	45143(99.9%)
	89	14081	0	0	14081	14081 (100%)
	93	32996	0	0	32996	32996(100%)
	102	26501	2	0	26499	26499(100%)
	112	37448	0	0	37448	37448(100%)
	127	26125	0	0	26125	26125(100%)
	128	42639	0	0	42639	42639(100%)
	130	43563	0	0	43563	43563(100%)
	132	43212	0	0	43212	43212(100%)
	136	69955	0	0	69955	69955(100%)
	150	50291	0	0	50291	50291(100%)
	153	74069	4	0	74065	74065(100%)
	159	44769	0	0	44769	44769(100%)
	161	44838	2	0	44836	44836(100%)
	165	54908	2	0	54906	54906(100%)
	166	24901	0	0	24901	24901(100%)
	174	29221	0	0	29221	29221(100%)
	175	23590	0	0	23590	23590(100%)
	176	55272	0	0	55272	55272(100%)
	180	32055	0	0	32055	32055(100%)
	183	39704	0	0	39704	39704(100%)
	186	34078	0	0	34078	34078(100%)
	195	41677	2	0	41675	41675(100%)
	197	19526	0	0	19526	19526(100%)
	205	23185	0	0	23185	23185(100%)
	209	54624	0	0	54624	54624(100%)
	212	41742	0	0	41742	41742(100%)
	229	18983	0	0	18983	18983(100%)
	231	10076	0	0	10076	10076(100%)
	237	35275	0	0	35275	35275(100%)
	239	30244	0	0	30244	30244(100%)
	255	58667	0	0	58667	58667(100%)
	256	49204	0	0	49204	49204(100%)
	258	55621	0	0	55621	55621(100%)
	284	46562	0	0	46562	46562(100%)
	288	22916	0	0	22916	22916(100%)
	289	17928	0	0	17928	17928(100%)
	290	26859	0	0	26859	26859(100%)
	295	32534	0	0	32534	32534(100%)
	301	20127	0	0	20127	20127(100%)

TABLE 4-continued

Result of deep-sequencing of plants which have been found to have edited StPPO2 gene						
Target gene	Sample	Total reads	WT	Inser- tions	Dele- tions	In-del frequency (%)
	306	15473	2	0	15471	15471(100%)
	311	16179	0	0	16179	16179(100%)
	323	18987	0	0	18987	18987(100%)
	331	15570	0	0	15570	15570(100%)
	338	38796	0	0	38796	38796(100%)
	346	28216	0	0	28216	28216(100%)
	350	36566	0	0	36566	36566(100%)
	353	18016	0	0	18016	18016(100%)
	355	36073	0	0	36073	36073(100%)
	356	10496	0	0	10496	10496(100%)
	365	28178	0	0	28178	28178(100%)
	371	26021	0	0	26021	26021(100%)
	376	23475	0	0	23475	23475(100%)
	380	28852	0	0	28852	28852(100%)
	390	16213	8	0	16205	16205(100%)
	392	13176	0	0	13176	13176(100%)
	395	19293	0	0	19293	19293(100%)
	406	21283	0	0	21283	21283(100%)
	407	24463	0	0	24463	24463(100%)
	413	31737	0	0	31737	31737(100%)
	414	12820	0	0	12820	12820(100%)
	418	12609	0	0	12609	12609(100%)
	418	12609	0	0	12609	12609(100%)
	427	38892	0	0	38892	38892(100%)
	433	12647	0	0	12647	12647(100%)
	440	25470	0	0	25470	25470(100%)
	447	26169	0	0	26169	26169(100%)
	448	31817	0	0	31817	31817(100%)
	461	19279	0	0	19279	19279(100%)
	466	13381	0	0	13381	13381(100%)
	470	18672	0	0	18672	18672(100%)
	472	10305	0	0	10305	10305(100%)
	484	22718	0	0	22718	22718(100%)
	494	37183	0	0	37183	37183(100%)
	496	33914	0	0	33914	33914(100%)
	502	29990	0	0	29990	29990(100%)
	503	28795	0	0	28795	28795(100%)
	505	31215	0	0	31215	31215(100%)
	509	25079	0	0	25079	25079(100%)
	512	38333	0	0	38333	38333(100%)
	514	29645	0	0	29645	29645(100%)
	519	29202	0	0	29202	29202(100%)
	529	23519	0	0	23519	23519(100%)
	530	34261	0	0	34261	34261(100%)
	534	31146	0	0	31146	31146(100%)
	542	28272	0	0	28272	28272(100%)
	543	31396	0	0	31396	31396(100%)
	544	39558	0	0	39558	39558(100%)
	548	18958	0	0	18958	18958(100%)
	553	10698	0	0	10698	10698(100%)
	554	14002	0	0	14002	14002(100%)
	575	14373	0	0	14373	14373(100%)
	582	17559	0	0	17559	17559(100%)
	584	23013	0	0	23013	23013(100%)
	586	25502	0	0	25502	25502(100%)
	587	18379	0	0	18379	18379(100%)
	589	22951	0	0	22951	22951(100%)
	590	19561	0	0	19561	19561(100%)

[0062] As a result of deep-sequencing of the above 110 plants, in particular, it was found that, in the plant with edited StPPO2 gene produced by the method of the present invention, the mutation in which −2 bp or −4 bp deletion pattern is shown from all of the four alleles is responsible for 57% (63/110) of the InDel pattern of the entire 110 edited plants (Table 5 and FIG. 11).

TABLE 5

Pattern of InDel mutation in 110 plants with edited StPPO2 gene				
InDel Pattern #	Deletion size (bp)	Number of plants	Deletion sites (Base position from PAM sequence)	Remarks
1	-2	25	5 th , 6 th	All 4 alleles were found to have -2 bp deletion pattern
2	-4	24	4 th , 5 th , 6 th , 7 th	All 4 alleles were found to have -4 bp deletion pattern
3	-2/-4	14	pattern 1 & 2	Among 4 alleles, each 2 alleles were found to have number 1 and number 2 deletion pattern, respectively
4	-1	47	4 th	Among 4 alleles, with number 1 and number 2 deletion pattern, an allele having -1 bp deletion pattern was found
	-5		4 th , 5 th , 6 th , 7 th , 8 th	Among 4 alleles, with number 1 and number 2 deletion pattern, an allele having -5 bp deletion pattern was found
	-10		5 th -14 th	Only from 1 allele of #205 line, -10 bp deletion pattern was found
Total		110		

[0063] To finally obtain a tuber from the potato lines having edited StPPO2 gene, young potato plants which have been found to have edited gene were cultured after transferring them to a greenhouse for cultivating seed potato. After 3 months or so, a tuber was obtained from some of the potato lines. PPO2 protein activity and suppression of browning were then observed using the tuber.

Example 5. Measurement of PPO2 Protein Activity in Plant with Edited StPPO2

[0064] PPO2 activity measurement from tissues of a potato plant which has been found to have edited StPPO2 gene (#14, 16, 25 and 30) and also a non-edited control (WT) potato was carried out according to the following method. First, 0.5 g of potato tuber was chopped to small pieces and added to a 1.5 ml tube. Cold sodium phosphate solution (pH 6.5, 1 ml) was added thereto and the tissues were homogenized using a pipette tip. After that, the mixture was centrifuged for 20 minutes at 13,000 rpm, 4° C. condition. While carrying out the centrifuge, sodium phosphate solution (pH 5.5) and 0.2 M pyrocatechol solution were admixed with each other in a 96-well plate and subjected to pre-incubation for 10 minutes. After the above centrifuge, the supernatant was added to the 96-well plate which has been added with a mixture solution of sodium phosphate and pyrocatechol followed by pre-incubation for 10 minutes, and the absorbance at 410 nm was immediately measured. The absorbance measurement was carried out up to 2 minutes with an interval of 30 seconds starting from see 0.

[0065] According to Bradford assay, protein concentration was measured by using the remaining supernatant. Based on the measured protein concentration and the absorbance value measured at 410 nm, units of PPO2 were calculated.

[0066] For a blind test to avoid any experimental bias, plant samples with edited gene were numbered, and then the

test was carried out as described in the above. According to the result of measuring the PPO2 protein activity, it was found that, in the plant line with edited gene, the PPO2 activity was suppressed at least by 28.6% and at most by 57.8% compared to the non-edited control (WT) as the result is illustrated in FIG. 12.

Example 6. Examination of Browning of Plant with Edited StPPO2

[0067] Observation of potato browning was made with the following 2 methods.

[0068] First, a potato tuber was sliced using the same method as the method employed in the previous report (Gonzalez M N et al., Front Plant Sci. (2020) 10: 1649), and used for the test. The result is shown in FIG. 13A. As a result of observing browning after slicing the potato tuber, it was found that the browning is suppressed compared to the control (WT). However, due to the moisture evaporation, there was no significant progress in the suppression of browning so that a potato tuber with the same size as above was simply peeled and used for the observation of browning. As a result, it was found that the tuber of a potato plant having edited StPPO2 (GE #14, #25) has significantly suppressed browning compared to the control (WT) (FIG. 13B). Consequently, it was possible to recognize that the browning of a plant line with edited StPPO2 gene is related to the activity of PPO2 protein.

REFERENCE TO AN ELECTRONIC SEQUENCE LISTING

[0069] A sequence listing electronically submitted on May 4, 2023 as an ASCII TXT file named 20230504_S12423GR06_TU_SEQ.TXT, created on May 3, 2023 and having a size of 7,888 bytes, is incorporated herein by reference in its entirety.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1
<211> LENGTH: 1958
<212> TYPE: DNA

-continued

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 1

```

tcttttgcgt tttgagcaat aatggcaagc ttgtgcaata gtagtagtac atctctcaaa    60
actcctttta cttcttcctc cacttcttta tcttccactc ctaagccctc tcaacttttc    120
atccatggaa aacgtaacca aatgttcaaa gtttcatgca aggttaccaa taataacgggt    180
gacaaaaacc aaaacgttga aacaaattct gttgatcgaa gaaatgttct tcttggtta    240
ggtggtcttt atggtgttgc taatgctata ccattagctg catccgctgc tccagctcca    300
cctctgato tctcgtcttg tagtatagcc aggattaacg aaaatcaggt ggtgccgtac    360
agttgttgcg cgcctaagcc tgatgatatg gagaaagttc cgtattacaa gttcccttct    420
atgactaagc tccgtgttcg tcagcctgct catgaagcta atgaggagta tattgccaag    480
tacaatctgg cgattagtcg aatgaagat cttgataaga cacaaccttt aaaccctatt    540
ggttttaagc aacaagctaa tatacattgt gcttattgta acggtgctta tagaattggt    600
ggcaaagagt tacaagttca taattcttgg cttttcttcc cgttccatag atggtaactg    660
tacttcacag agagaatcgt gggaaaattc attgatgato caactttcgc tttaccatat    720
tggaattggg accatccaaa aggtatgcgt tttcctgcca tgtatgatcg tgaagggact    780
tcccttttcg atgtaacacg tgaccaaagt caccgaaatg gagcagtaat cgatcttggg    840
tttttcggca atgaagtga aacaactcaa ctccagttga tgagcaataa tttaacacta    900
atgtaccgto aaatggtaac taatgctcca tgtcctcgga tgttctttgg cgggccttat    960
gatctcgggg ttaacactga actcccgga actatagaaa acatccctca cggctcctgtc   1020
cacatctggt ctggtacagt gagaggttca actttgccca atggtgcaat atcaaacggg   1080
gagaatatgg gtcattttta ctcagctggt ttggaccggg ttttcttttg ccatcacagc   1140
aatgtggatc ggatgtggag cgaatggaaa gcgacaggag ggaaaagaac ggatatacaca   1200
cataaagatt ggttgaactc cgagttcttt ttctatgatg aaaatgaaaa cccttaccgt   1260
gtgaaagtca gagactgttt ggacacgaag aagatgggat acgattacaa accaattgcc   1320
acaccatggc gtaacttcaa gcccttaaca aaggcttcag ctggaaaagt gaatacagct   1380
tcacttcgcg cagctagcaa tgtattccca ttggetaaac tcgacaaaagc aatttcgttt   1440
tccatcaata ggccgacttc gtcaaggact caacaagaga aaaatgcaca agaggagatg   1500
ttgacattca gtagcataag atatgataac agaggttaca taagggttca tgtgttttcg   1560
aacgtggaca ataatgtgaa tgcgaatgag cttgacaagg cggagtttgc ggggagttat   1620
acaagtttgc cacatgttca tagagctggt gagactaatc atatcgcgac tgttgatttc   1680
cagctggcga taacggaact gttggaggat attggtttgg aagatgaaga tactattgcy   1740
gtgactctgg tgccaaagag aggtggtgaa ggtatctcca ttgaaggtgc gacgatcagt   1800
cttgacagatt gttaattagt ctctattgaa tctgctgaga ttacactttg atggatgatg   1860
ctctgttttt gttttcttgt tctgtttttt cctctgttga aatcagcttt gttgcttgat   1920
ttcattgaag ttgttattca agaataaatc agttacaa                                1958

```

<210> SEQ ID NO 2

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: StPP02-1

<400> SEQUENCE: 2

tccatggatg aaaagttgag 20

<210> SEQ ID NO 3

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: StPP02-2

<400> SEQUENCE: 3

tgcataaaac tttgaacatt 20

<210> SEQ ID NO 4

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: StPP02-3

<400> SEQUENCE: 4

atactacaag acgagagatc 20

<210> SEQ ID NO 5

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: StPP02-4

<400> SEQUENCE: 5

ttaggcgcgc aacaactgta 20

<210> SEQ ID NO 6

<211> LENGTH: 72

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 6

ccacttcttt atcttccact cctaagccct ctcaactttt catccatgga aaacgtaacc 60

aaatgttcaa ag 72

<210> SEQ ID NO 7

<211> LENGTH: 71

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 1 del of StPP02

<400> SEQUENCE: 7

ccacttcttt atcttccact cctaagccct ctcaactttt atccatgga aacgtaacca 60

aatgttcaaa g 71

<210> SEQ ID NO 8

<211> LENGTH: 71

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 8

-continued

tctcaacttt tcatccatgg aaaacgtaac caaatgttca agtttcatg caaggttacc	60
aataataacg g	71

<210> SEQ ID NO 9
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 1 del of StPP02

<400> SEQUENCE: 9	
tctcaacttt tactccatgg aaaacgtaac caatgttcaa agtttcatgc aaggttacca	60
ataataacgg	70

<210> SEQ ID NO 10
 <211> LENGTH: 72
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 10	
gtgcatccg ctgctccagc tccacctcct gatctctcgt cttgtagtat agccaggatt	60
aacgaaaatc ag	72

<210> SEQ ID NO 11
 <211> LENGTH: 72
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 11	
ccaggattaa cgaaaatcag gtggtgccgt acagttgttg cgcgccctaa cctgatgata	60
tggagaaagt tc	72

<210> SEQ ID NO 12
 <211> LENGTH: 68
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 4 del of StPP02

<400> SEQUENCE: 12	
ccaggattaa cgaaaatcag gtggtgccgt acgttgccgc cctaagcctg atgatatgga	60
gaaagtcc	68

<210> SEQ ID NO 13
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 2 del of StPP02

<400> SEQUENCE: 13	
ccaggattaa cgaaaatcag gtggtgccgt acatgttgcg cgccctaaagcc tgatgatatg	60
gagaaagtcc	70

<210> SEQ ID NO 14
 <211> LENGTH: 67
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: 5 del of StPP02

<400> SEQUENCE: 14

ccaggattaa cgaaaatcag gtggtgccgt acatgcgcgc ctaagcctga tgatatggag 60

aaagttc 67

<210> SEQ ID NO 15

<211> LENGTH: 66

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 6 del of StPP02

<400> SEQUENCE: 15

ccaggattaa cgaaaatcag gtggtgccgt acagcgcgcc taagcctgat gatatggaga 60

aagttc 66

<210> SEQ ID NO 16

<211> LENGTH: 69

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 16

ggattaacga aaatcagggtg gtgccgtaca gttgttgcgc gcctaagcct gatgatatgg 60

agaaagttc 69

<210> SEQ ID NO 17

<211> LENGTH: 67

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 2bp del of StPP02

<400> SEQUENCE: 17

ggattaacga aaatcagggtg gtgccgtaca ttttgcgcgc ctaagcctga tgatatggag 60

aaagttc 67

<210> SEQ ID NO 18

<211> LENGTH: 68

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 1bp del of StPP02

<400> SEQUENCE: 18

ggattaacga aaatcagggtg gtgccgtacg ttgttgcgcg cctaagcctg atgatatgga 60

gaaagttc 68

<210> SEQ ID NO 19

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: 4bp del of StPP02

<400> SEQUENCE: 19

ggattaacga aaatcagggtg gtgccgtacg ttgcgcgcct aagcctgatg atatggagaa 60

agttc 65

-continued

```

<210> SEQ ID NO 20
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 10bp del of StPPO2

<400> SEQUENCE: 20

ggattaacga aaatcagggtg gtgccgtaca gcctaagcct gatgatattg agaaagttc      59

<210> SEQ ID NO 21
<211> LENGTH: 64
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5bp del of StPPO2

<400> SEQUENCE: 21

ggattaacga aaatcagggtg gtgccgtact tgcgcgccta agcctgatga tatggagaaa      60
gttc                                              64

<210> SEQ ID NO 22
<211> LENGTH: 66
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3bp del of StPPO2

<400> SEQUENCE: 22

ggattaacga aaatcagggtg gtgccgtact gttgcgcgcc taagcctgat gatattggaga      60
aagttc                                              66

```

1: A composition comprising:
a ribonucleoprotein complex of a guide RNA specific to a target nucleotide sequence in *Solanum tuberosum* Polyphenol Oxidase 2 (StPPO2) gene derived from potato and an endonuclease protein; or
a recombination vector comprising a DNA encoding a guide RNA specific to a target nucleotide sequence in StPPO2 gene derived from potato and a nucleic acid sequence encoding an endonuclease protein.

2: The composition according to claim 1, wherein the target nucleotide sequence in StPPO2 gene consists of the nucleotide sequence of SEQ ID NO: 5.

3: A guide nucleic acid as an RNA sequence specific to a target region in nucleotide sequence constituting *Solanum tuberosum* Polyphenol Oxidase 2 (StPPO2) gene in which the target region is the nucleotide sequence of SEQ ID NO: 5.

4: A method of producing a genome-edited potato plant with suppressed browning, the method comprising:
introducing a guide RNA specific to a target nucleotide sequence in StPPO2 gene derived from potato and an endonuclease protein to a potato plant cell to have genome editing; and
regenerating a potato plant from the potato plant cell that is genome-edited.

5: The method according to claim 4, wherein the introduction comprises:
using an ribonucleoprotein complex of the guide RNA and the endonuclease protein; or
using a recombination vector comprising a DNA encoding the guide RNA and a nucleic acid sequence encoding the endonuclease protein.

6: The method according to claim 4, wherein the target nucleotide sequence in StPPO2 gene consists of the nucleotide sequence of SEQ ID NO: 5.

7: The method according to claim 4, wherein the potato plant cell that is genome-edited of the regenerating has been induced to have deletion mutation of 1 to 10 bp in size in the target nucleotide sequence in StPPO2 gene.

8: A genome-edited potato plant with suppressed browning which is produced by the method of claim 4.

9: The genome-edited potato plant according to claim 8, wherein the genome-edited potato plant with suppressed browning has been induced to have deletion mutation of 1 to 10 bp in size in a target nucleotide sequence in StPPO2 gene.

10: Genome-edited seed potato of the potato plant according to claim 8.

* * * * *