

## LetsGetChecked Blood Sample ImPress IFU 120 SPECS AND INSTRUCTIONS

Colour Reference: CMYK

····

Size:

108.5mm wide by 152.5mm high folded

Paper:

. 135gsm silk

Fold type:

4 page accordion

Font:

Proxima Nova

(Regular, Semibold and Bold)

Part Number:

QR-IFU-120-R0 Mar-2025



\*Visual reference for how to fold down the sheet so the front page is the first one to see. Artwork and page count may vary.

## **FOLD GUIDE**

----- Fold

## Accordion fold





### Inside

	Contents pg	Steps pg 1	Steps	
		!		

## Outside

	Steps pg 4	Warnings	Front page
--	---------------	----------	------------

# Before you start

Estimated time needed: 10-15 minutes.



Read all the instructions including the warnings.



An instructional video can be viewed at LetsGetChecked.com/instructions-imp-collection



This is a single use device. Once you have removed the tabs, do not press the device until it is positioned on the arm and you are ready to begin sample collection.



Short sleeves are recommended for sample collection.




Have a timer ready to set for 10 minutes.



Contact customer support at help.LetsGetChecked.com if the device does not function as described.

# Contents

Remove all the contents from the box and make sure you have the following items:



## Also required but not provided:

- Timer
- Bandages
- Sterile Gauze
- Alcohol swab
- ☐ Medical gloves (if collecting the sample from someone else)

# How to collect a blood sample



Wash and dry your hands. If you are collecting the sample from someone else, the use of medical gloves is recommended.



Remove the device and tube cap from the packaging. You will use the cap later to seal the sample in the blood collection tube.



Clean the upper arm with an alcohol swab and allow to air-dry.



Take note of the two fill lines on the tube. You will aim to fill to the top line with blood. The top fill line may be hard to see once the device is placed on your arm.



Twist and remove the black tabs from the device.



Peel off the clear plastic strip using the tab at the top of the device.



Keep the arm straight and fingers pointing down towards the floor. Perform blood collection in an upright seated position.

Please continue overleaf.



LGC BLOOD SAMPLE\_IMPRESS\_INVESTIGATION\_QR\_IFU\_120\_R0\_4\_PG.indd 3-6 04/03/2025 09:57



Stick the device on the upper arm with the collection tube pointing downwards.



Using your palm, firmly press the device against the arm until you hear a click. Release to start the collection process.



Start a timer for 10 minutes. It can take up to two minutes to see the first drop of blood in the collection tube.



Follow the next step to remove the device when:

- Blood has filled to the top line
- 10 minutes have passed.



Peel the device from the arm while keeping the tube upright.



To remove the tube from the device, gently pull it downwards. Discard the device.



Place the cap on the blood collection tube and press down firmly until you hear a click. Clean arm with gauze and apply a bandage if needed.



Turn the tube upside down 10 times to mix the sample. Gently tap or flick the tube to get blood moving if needed. Wash and dry your hands.

## **⚠** WARNING:

- · You must be over the age of 18 to use this device to self-collect a blood sample.
- Single use only. Not intended for more than one use. For use only on a single patient.
- Not for use on infant heels.
- Do not use after the expiration date.
- · Do not resterilize.
- For external use only.
- Use personal protective equipment when handling biohazardous materials.
- The lancet is single use and automatically retracts into the device in a safe position after use.
- Improper use of blood lancets can increase the risk of inadvertent transmission of bloodborne pathogens, particularly in settings where multiple patients are tested.
- Minor bruising or residual marks may occur at the sample collection site. Consult with a healthcare provider if you are concerned or if this persists.
- Do not use the device on damaged, scarred, or infected skin. There is a small possibility of pain, nerve damage, scarring, bone damage, or infection. Consult a healthcare provider if you are concerned.
- Use while seated as fainting may occur with any blood sampling procedure. If you are concerned, consult a healthcare provider before collecting your sample. If you proceed, ensure you have someone with you. If you feel

faint or unwell during sample collection, stop sample collection. Consult a healthcare provider if you continue to feel faint or unwell.

- Do not use if any of the contents are damaged or missing. Do not use if device packaging has been opened or damaged. Do not tamper with the device Risk of serious injury.
- Discard the entire device after use. Follow local guidelines for discarding used sharps as a biohazard material.
- Laboratories must validate use of product for their specific diagnostic assay.
- Choking hazard Keep out of reach of children.
- Contact our team at +1 929-376-0056 before using this device if you have any questions or concerns.

### Do not use this device without consulting a healthcare provider if:

- You have a history of fainting or a medical condition that increases your risk of fainting.
- You have a bleeding disorder.
- You have a condition associated with an increased risk of infection, impaired wound healing, skin ulceration, or swelling in the arms.

#### Purpose/Intended use of device:

The ImPress is a single-use lancing device intended for obtaining capillary whole blood samples from the upper arm into a pre-installed compatible blood collection device.



PrivaPath Diagnostics Ltd., Unit 1, Northern Cross Business Park, North Road, Dublin 11. D11 XT26, Ireland.















LGC\_BLOOD\_SAMPLE\_IMPRESS\_INVESTIGATION\_QR\_IFU\_120\_R0\_4\_PG.indd 7-10 04/03/2025 09:57