

Notes from D&P and related books

Peter Mao

May 13, 2023

Contents

1	Dodson&Poston	8
1.1	Contents	8
1.2	0.2. Functions	8
1.2.1	typo: In definition of g ,	8
1.3	II. Affine Spaces	8
1.3.1	1.01. Definition: <i>affine space</i>	8
1.3.2	1.02. Tangent spaces, freeing and binding maps.	8
1.4	V.1.10. Notation.	8
1.5	VI. Topological Vector Spaces	8
1.5.1	1. Continuity	8
1.5.2	4. Compactness and Completeness	9
1.6	VII. Differentiation and Manifolds	9
1.7	Theorem (Inverse Function Theorem).	9
1.8	VII.3.02. Language: Df, df (the other d)	9
1.9	VII.3.03 Tangent bundle on a manifold, abbreviations	9
1.10	A tensor field of type $(0\ 0)^T$ is just a function,	9
1.11	VII.4.01. Covariant Vectors.	9
1.12	VII.4.02. Contravariant Vectors	10
1.13	VII.6 Vector fields and flows	10
1.13.1	6.02. Definition. A solution curve or integral curve	10
1.13.2	. Definition. A C^k local flow...	10
1.13.3	. Corollary $\phi_{t+s} = \phi_t \circ \phi_s$	10
1.13.4	6.07. Lemma. Let M be a manifold on an affine space X	10
1.14	VII.7 Lie Brackets	10
1.14.1	7.01. Definition. The Lie bracket or commutator	11
1.14.2	7.02. Theorem.	11
1.14.3	Fig. 7.1 is not referenced anywhere!	11

1.14.4	7.04. Theorem. ("the Chart theorem")	11
1.14.5	Exercises VII.7	12
1.15	VIII. Connections and Covariant Differentiation	13
1.16	VIII.2. Rolling Without Turning	13
1.17	VIII.3. Differentiating Sections	13
1.17.1	"directional derivative"	13
1.17.2	"we are free to decide what properties would be nice to have"	13
1.17.3	Typo! first f should be t	13
1.17.4	Need to understand the first term on the right	13
1.17.5	3.01. Definition. A connection (∇)	13
1.17.6	3.02. Coordinates – Christoffel symbols	13
1.17.7	3.03. Transformation Formula.	13
1.17.8	3.04. Definition: vector field along c	13
1.17.9	3.05. Differentiating, Along Curves, Fields Along Curves.	14
1.17.10	3.06. Definition: "vertical"	14
1.17.11	3.07. Definition: "vertical" and "horizontal" parts of a vector	14
1.17.12	3.08. Language	14
1.17.13	Exercises 3 & 4	14
1.18	VIII.4. Parallel Transport	14
1.18.1	4.01. Definition: parallel vector field	14
1.18.2	what does this even mean?	15
1.18.3	4.02. Theorem: uniqueness of the parallel vector field along c	15
1.18.4	4.03. Definition: <i>parallel transport</i> τ	15
1.19	Bibliography	16
1.20	Index of Notations	16
2	NeedhamT VDGF	16
2.1	Contents	17
2.2	1.3 Angular excess of a spherical triangle	17
2.3	1.5 Constructing geodesics with tape	17
2.4	2.1 Gaussian curvature in terms of excess angle	17
2.5	2.2 Circ/Area of a circle – reread this!	17
2.6	2.3 Local Gauss-Bonnet Theorem	17
2.7	Chapter 4 The Metric	17
2.7.1	4.3 Metric of a general surface	17
2.7.2	Gauss EFG coefficients of the first fundamental form	17
2.7.3	4.4 Metric Curvature Formula	17

2.7.4	4.5 Conformal Maps	17
2.8	Chapter 5 Pseudosphere and the Hyperbolic plane	17
2.8.1	5.2 Tractrix and the Psuedosphere	17
2.8.2	5.3 Conformal map of the pseudosphere	17
2.9	8.3 Newton's curvature formula	17
2.10	Eqn 8.6, curvature	17
2.11	Chapter 9 Curves in 3-Space	18
2.11.1	definition: Torsion	18
2.11.2	definition: binormal of a curve	18
2.12	Chapter 10 The Principal Curvatures of a Surface	18
2.12.1	(10.1) Euler's Curvature Formula:	18
2.12.2	Dupin's indicatrix	18
2.13	Chapter 11 Geodesics and Geodesic Curvature	18
2.13.1	Fig [11.2] $K = K_g + K_n$	18
2.14	Chapter 12 The Extrinsic Curvature of a Surface	19
2.14.1	Figure 12.5: normal vectors on principal and general curves	19
2.15	Chapter 15 The Shape Operator, S , and ∇_v	19
2.15.1	15.1 Directional Derivatives (∇_v)	19
2.15.2	15.2 The Shape Operator S	19
2.15.3	15.3 Geometric effect of S	19
2.15.4	(15.18) $\kappa(v) = v \cdot S(v)$	19
2.15.5	(15.7) sum of curvatures in perpendicular directions	19
2.15.6	(15.26) curvature and torsion	20
2.15.7	15.9 Classical Terminology and Notation: The Three Fundamental Forms	20
2.15.8	19.8 The Road Ahead	20
2.16	Chapter 21 An Historical Puzzle	20
2.16.1	GR and parallel transport	20
2.17	Chapter 22 Extrinsic Constructions	20
2.17.1	Projection (not rotation) of w into T_q	20
2.17.2	Rotation of w into T_q	20
2.18	Chapter 23 Intrinsic (covariant) Constructions	20
2.18.1	23.2 The Intrinsic (aka, "Covariant") Derivative	20
2.18.2	(Eqn 22.1) Intrinsic derivative, (D_v , the convariant derivative)	21
2.18.3	Fig 23.3	21
2.18.4	D_v is also called the Levi-Civita Connection.	21
2.18.5	intrinsic derivative $D_v w$	21

2.18.6	Here is an extrinsic way of looking at the intrinsic derivative.	21
2.18.7	Eqn 23.3 $D_v w$	21
2.18.8	Compare these with $C_i C_j$ in D&P p 217	21
2.19	1-forms used to be called covariant vectors or covectors	23
2.20	Further Reading	23
2.20.1	Intuitive Topology, by V. V. Prasolov.	23
2.20.2	Three-Dimensional Geometry and Topology, by William P. Thurston.	23
2.20.3	Differential Geometry: A Geometric Introduction, by David W. Henderson.	23
2.20.4	Gauge Fields, Knots and Gravity, by John Baez and Javier P. Muniain.	23
3	TuLW Differential Geometry	23
3.1	§1 Riemannian Manifolds	24
3.1.1	1.3 Riemannian Metrics	24
3.1.2	Example 1.11: Torus	24
3.1.3	partition of unity	24
3.1.4	Theorem 1.12. On every manifold M there is a Riemannian metric.	24
3.2	§2 Curves	24
3.2.1	2.3 Signed Curvature of a Plane Curve	24
3.2.2	Curvature formulae	24
3.3	§4 Directional Derivatives in Euclidean Space	24
3.3.1	Torsion of the directional derivative D	24
3.3.2	Curvature of the directional derivative	24
3.3.3	Definition 4.8. "On" vs "Along" a submanifold	24
3.3.4	Notation $X(M)$ and $\Gamma(TM _N)$	25
3.3.5	In a submanifold of \mathbb{R}^n , torsion "no longer makes sense"	25
3.4	§5 The Shape Operator	25
3.4.1	Defn: regular point – need a picture	25
3.4.2	5.2 The Shape Operator	25
3.4.3	Is the usage of "on" here the same as before?	25
3.4.4	Proposition 5.3. The shape operator is self-adjoint:	25
3.4.5	Proposition 5.5: Curvature is given by the second fundamental form	25
3.4.6	Proposition 5.6: Principal directions and curvatures	25
3.4.7	5.4 The First and Second Fundamental Forms	26
3.5	§6 Affine Connections	26

3.5.1	No canonical basis for $T_p M$ not embedded in a Euclidean space	26
3.5.2	Definition 6.1. An affine connection	26
3.5.3	6.2 Torsion and Curvature	26
3.5.4	Definition 6.4: Riemannian or Levi-Civita connection	26
3.5.5	Theorem 6.6. On a Riemannian manifold there is a unique Riemannian connection.	26
3.5.6	6.4 Orthogonal Projection on a Surface in \mathbb{R}^3	26
3.6	§7 Vector Bundles	27
3.6.1	7.1 Definition of a Vector Bundle	27
3.7	13.1 Theorem	27
3.8	14.5 Parallel Translation	27
4	Littlejohn Manifolds	27
4.1	Notes for page 3	27
4.2	(typo)	27
4.3	(typo)	27
4.4	8. Tangent Vectors	27
4.4.1	Eqn 11	27
4.5	9. Equivalence Classes of Curves	27
4.6	10. Tangent Vectors in Coordinates	27
4.6.1	Eqn 16	28
4.6.2	tangent vector at p	28
4.7	12. Covectors	28
4.7.1	Eqn 21 $df _p$	28
4.7.2	$df _p$ is an operator on vectors – it is not small or "infinitesimal"	28
4.7.3	13. Differential of Coordinates	28
5	BachmanD: A Geometric Approach to Differential Forms	28
5.1	1.2 Generalizing the integral	28
5.2	1.3 differential form in a nutshell	28
5.3	3 Forms	28
5.3.1	3.1 Coordinates for vectors	28
5.3.2	3.2 1-Forms	29
5.3.3	3.3 Multiplying 1-forms	29
5.3.4	3.4 2-Forms on $T_p \mathbb{R}^3$	30
5.3.5	3.5 2-Forms and 3-forms on $T_p \mathbb{R}^4$	30
5.3.6	3.6 n-Forms	30
5.3.7	3.7 Algebraic computation of products	30

5.4	4 Differential Forms	30
6	LeeJM _SmoothManifolds (+18)	30
6.1	Contents	30
6.2	Proposition 2.25 (Existence of Smooth Bump Functions). . .	30
6.3	Chapter 3 Tangent Vectors	30
6.4	Chapter 8 Vector Fields	30
6.4.1	Vector Fields on Manifolds	30
6.4.2	Example 8.12. cylindrical orthonormal frame on $\mathbb{R}^2 \setminus \{0\}$	30
6.4.3	Vector Fields and Smooth Maps	30
6.4.4	Lie Brackets	31
6.4.5	Lemma 8.25. Lie bracket of smooth vector fields. . . .	31
6.4.6	Proposition 8.26 (Coordinate Formula for the Lie Bracket). .	31
6.4.7	Example 8.27	31
6.4.8	see D&P 3	31
6.5	Chapter 9 Integral Curves and Flows	31
6.5.1	Flowouts	31
6.5.2	Lie Derivatives	31
6.5.3	Commuting Vector Fields	32
6.6	Chapter 11 The Cotangent Bundle	33
6.6.1	Prop 11.8 $V \cong V^{**}$	33
6.6.2	Tangent Covectors on Manifolds	33
6.6.3	contravariant transformation	33
6.6.4	covariant transformation	33
6.6.5	Prop 11.18 (coordinate covectors are dx^j 's)	33
6.6.6	Proposition 11.23 (Derivative of a Function Along a Curve).	34
6.7	Chapter 12 Tensors	34
7	TuLW Introduction to Manifolds	34
7.1	2.1 Directional derivative	34
7.2	Defn: germ	34
7.3	Defn: derivation	34
7.4	2.4 Vector fields	34
7.5	3.3 k-covectors	34
7.6	Lemma 3.11 $\tau(\sigma f) = (\tau\sigma)f$	34
7.7	3.7 Wedge product	35
7.8	Defn: graded algebra over field K	35
7.9	4 Differential forms on \mathbb{R}^n	35
7.10	4.6 Applications to Vector Calculus	35

7.11	4.7 Convention on Subscripts and Superscripts	35
7.12	Sec 5: Manifolds	35
7.13	5.1 Topological Manifolds	36
7.14	5.3 Smooth Manifolds - maximal atlas defn	36
7.15	Definition 5.9 smooth manifold	36
7.16	Smooth manifold conditions	36
7.17	5.4 Examples of Smooth Manifolds	36
7.18	§6 Smooth Maps on a Manifold	37
7.19	6.3 Diffeomorphisms	37
7.20	6.6 Partial Derivatives	37
7.21	6.7 The Inverse Function Theorem	37
7.22	Chapter 3: The tangent space	37
7.23	bump functions?	37
7.24	Theorem 20.4.	37
7.25	A.4 First and Second Countability	37
	7.25.1 Lemma A.10. Every open set in \mathbb{R}^n contains a rational point.	37
	7.25.2 Prop A.11 collection of open balls w/ rational centers/radii is a basis for \mathbb{R}^n	37
	7.25.3 Defn A.12 second countable \leq countable basis	37
	7.25.4 Defn A.15 basis of neighborhoods, first countable	37
7.26	List of Notations	38
8	Carroll - An Introduction to General Relativity	38
8.1	1. SR and flat Spacetime	38
8.2	2. Manifolds	38
8.3	3. Curvature	38
8.4	4. Gravitation	38
8.5	5. Schwarzschild's solution	38
8.6	6. More general black holes	38
8.7	7. Perturbation theory & Gravitational radiation	38
8.8	8. Cosmology	38
8.9	9. QFT in curved spacetime	38
8.10	A. Maps between manifolds	38
8.11	B. Diffeomorphisms and Lie derivatives	38
8.12	C. Submanifolds	38
8.13	D. Hypersurfaces	38
8.14	E. Stokes's Theorem	38
8.15	F. Geodesic congruences	38
8.16	G. Conformal transformations	38

8.17 H. Conformal diagrams	38
8.18 I. The parallel propagator	38
8.19 J. Noncoordinate bases	38
8.20 Bibliography	38
8.21 Index	38

1 Dodson&Poston

1.1 Contents

1.2 0.2. Functions

1.2.1 typo: In definition of g ,

\mathbb{N} should be \mathbb{Z}

1.3 II. Affine Spaces

1.3.1 1.01. Definition: *affine space*

with vector space T is a non-empty set of points and a *difference function*
 $d: X \times X \rightarrow T$

1.3.2 1.02. Tangent spaces, freeing and binding maps.

the difference function of 1.01 is related to the bijection $d_x: x \times X \rightarrow T$
 which is called a *freeing map*.

Its inverse, d^{\leftarrow} is called a *binding map*.

1.4 V.1.10. Notation.

Vectors in X^k_h are called tensors on X , covariant of degree h and
 contravariant of degree k , or of type $(k\ h)^T$ By convention
 $X^0_0 = \mathbb{R}$.

1.5 VI. Topological Vector Spaces

1.5.1 1. Continuity

1. VI.1.01 $\epsilon - \delta$ continuity
2. 1.06 continuity

- a cubic is a nice function to demonstrate that open \rightarrow open is not necessarily true for continuous functions.
- If your open set in the domain starts and ends between the local extrema AND includes them, then the range is the closed interval bound by the local extrema.

3. Hausdorff topology This one is important.

- Finite intersections of open sets, any unions of open sets
- Finite unions of closed sets, any intersections of closed sets.

1.5.2 4. Compactness and Completeness

1. 4.02 Bolzano-Weierstrauss See Stillwell NLT, Ex 8.4.1, 8.4.2
2. See Stillwell NLT exercises 8.4.3, 8.4.4
3. 4.10 theorem $f(C)$ is compact extents. I needed a reminder that functions are not continuous at infinity. see $\epsilon - \delta$ continuity

1.6 VII. Differentiation and Manifolds

1.7 Theorem (Inverse Function Theorem).

$D_x f$ is an isomorphism iff \exists neighborhoods of x and $f(x)$ where a local inverse f^{\leftarrow} exists.

1.8 VII.3.02. Language: Df , df (the other d)

if f is a real-valued function, then $df: TM \rightarrow \mathbb{R}$ is the "collection" of all $d_{f(q)} \circ D_q f$ where $D_q f: T_q M \rightarrow T_{f(q)} \mathbb{R}$ and $d_{f(q)}: T_{f(q)} \rightarrow \mathbb{R}$
this is NOT the same as $Df: TM \rightarrow T\mathbb{R}$.

1.9 VII.3.03 Tangent bundle on a manifold, abbreviations

$$(T_x^* M) = (T_x M)^*$$

1.10 A tensor field of type $(0, 0)^T$ is just a function,

1.11 VII.4.01. Covariant Vectors.

See Lee Prop 11.18

1.12 VII.4.02. Contravariant Vectors

"we can identify \vec{t} with the linear map:" $\partial_{\vec{t}}: f \mapsto df(\vec{t})$
so for any vector field \vec{v} and function f , we have $(\vec{v}f)(x) = df(\vec{v}_x)$

1.13 VII.6 Vector fields and flows

1.13.1 6.02. Definition. A solution curve or integral curve

of vector field v on manifold M is a curve $c: J \rightarrow M$ such that $c'(t) = v(c(t)) \forall t \in J$

1.13.2 . Definition. A C^k local flow...

- $\phi(U, t)M$ where $U \subset M$, open and t is a parameter in an open interval
- KEY POINT: by condition (ii), local flows are made up of solution curves.

1.13.3 . Corollary $\phi_{t+s} = \phi_t \circ \phi_s$

by definition, $\phi_t: U \rightarrow M: x \mapsto \phi(x, t)$, which returns the displacement from x along the flow ϕ .

1.13.4 6.07. Lemma. Let M be a manifold on an affine space X

1.14 VII.7 Lie Brackets

- typo: should be $\phi((x, y), t)$, not $\phi((x + y), t)$
- typo: $\psi_1\phi_1(0, 0) = (0, 1)$
- typo: $\phi_1\psi_1(0, 0) = (1, 1)$
- ϕ and ψ are local flows (see VII.6.03 p 197)
- There are many objects in the preface to this section:
 1. (x, y) : points on a manifold (or coordinates by abuse of language)
 2. (s, t) : parameters
 3. (ϕ, ψ) : flows
 4. (v, w) : vector fields (related to the flows)
 5. f : any function on the manifold
- claim (for this example only): $[\phi \circ \psi - \psi \circ \phi](x, y) = (1, 0)$

1.14.1 7.01. Definition. The Lie bracket or commutator

- act on vector fields, not flows.
- typo: (bottom of 200) $v(w(fg)) - w(v(fg))$
- $= g[v, w](f) - f[v, w](g)$

“...so we have a new derivation” (namely, $[v, w]$)

1.14.2 7.02. Theorem.

iff fields commute then composition of flows also commute.

1.14.3 Fig. 7.1 is not referenced anywhere!

1.14.4 7.04. Theorem. ("the Chart theorem")

If we have a linearly independent and commuting set of vector functions that span a manifold, then they can be "realized" as the basis vectors (∂_i 's) of some chart.

These flows have constant velocity in the neighborhood of x . (see example in preface to this section.)

1. On Randy's discomfort with this notation We want to justify the relationship $\phi_t^i(x^1, \dots, x^n) = (x^1, \dots, x^i + t, \dots, x^n)$.

Note that $M \ni \vec{x} = (x^1, \dots, x^n) = \theta(0, \dots, 0)$, so the LHS can be rewritten $\phi_t^i(\theta(0, \dots, 0)) = \phi_t^i \circ \theta(0, \dots, 0)$

Using 1.13.3 (and going through the proof backwards), this becomes $\theta(0, \dots, t, \dots, 0)$ with the t in the i slot, which is the same as $\vec{x} + t\vec{e}_i = (x^1, \dots, x^i + t, \dots, x^n)$

In the proof, this appears with parameter s instead of t .

2. $\theta(t^1 \dots t^n) \in M$ $\theta(t^1 \dots t^n)$ is a point in M . By varying one of the t 's, it describes a curve on M . $\theta(0, 0, \dots, 0) = x \in M$. As a map, $\theta: \mathbb{R}^n \rightarrow M$, so it **might** be the inverse of a chart function.
3. ...by the smoothness of the ϕ^1 why only ϕ^1 ? Each $\phi_t^i(x)$ moves the point along its flow line, so the composition of all these ϕ 's is a point in M . The fact that ϕ^1 is smooth means that through arbitrary points in M , ϕ^1 defines a smooth flow, so θ is also smooth through arbitrary points in M .

4. $D\theta$ takes the vector $e_1(t^1 \dots t^n) \dots$. This should be e_i , not e_1 . otherwise, how does $c(s)$ become a parameterized curve through $\theta(t^1 \dots t^n)$ along the i direction?
5. typo: 6.05 "6.50" should be 1.13.3
6. typo: should be θ so we can rewrite this as $\phi_s^i(p)$ for $p \in M$
7. How does this "so c is a solution curve" come about? See 1.13.2: a local flow is a solution curve by definition and $c = \phi_s$. so yes it is.
8. typo: remove θ should be $c^*(0) = v_i(c(0))$
9. $D\theta$ takes the standard basis... $D\theta: T_{(0, \dots, 0)}\mathbb{R}^n \rightarrow T_x M : e_j \rightarrow (v_j)_x$
10. What are they saying???

That is a linearly independent subset of the n -dimensional space $T_x M$, by assumption, so $D\theta(v_1, \dots, v_n)$ is an isomorphism.

What is "that"? "That" is $\{(v_1)_x, \dots, (v_n)_x\}$

if we're in Bachman, then $\langle e_1, \dots, e_n \rangle$ is a general-ish vector in $T_{0 \dots} \mathbb{R}^n$

1.14.5 Exercises VII.7

1. Ex VII.7.2 parts a and c are covered by 6.4.6
2. Ex VII.7.4: Typos abound! THIS IS THE LIE DERIVATIVE!! yes, very important. See Lee. :NOTER_{PAGE}: (220 0.42246376811594205 . 0.13516483516483516)

The subscript to the first v should be $\phi_h(x)$: $[u, v]_x = \lim_{h \rightarrow 0} \frac{(D_x \phi_h)^{\leftarrow} v_{\phi_h(x)} - v_x}{h}$

v_x , and $[u, v]_x$ are both tangent vectors in $T_x M$ $v_{\phi_h(x)}$ is a tangent vector in $T_{\phi_h(x)} M$

$\phi_h: M \rightarrow M$, so

$$D_x \phi_h: T_x M \rightarrow T_{\phi_h(x)} M: v_x \mapsto \lim_{h \rightarrow 0} \frac{v_{\phi_h(x)} - v_x}{h}$$

3. Typo in missing f , should be at least: $[\vec{u}, f\vec{v}] = \vec{u}(f)\vec{v} + f[\vec{u}, \vec{v}]$

Note: this agrees with Lee p 188 because $\vec{v}(1) = d(1)\vec{v} = 0$

1.15 VIII. Connections and Covariant Differentiation

1.16 VIII.2. Rolling Without Turning

1.17 VIII.3. Differentiating Sections

1.17.1 "directional derivative"

In D&P-speak, "vertical" is vertical in their picture of TM, meaning that the vector lies in $T_p M$ for some $p \in M$.

1.17.2 "we are free to decide what properties would be nice to have"

See Tu(DG) Sec 6, p 30. Without saying so, D&P appear to be leading us to the **Riemannian** or **Levi-Civita** connection. Tu says this is the unique, torsion-free affine connection on a Riemannian manifold.

1.17.3 Typo! first f should be t

1.17.4 Need to understand the first term on the right

See Tu(DG) Prop 4.9.ii p. 27

1.17.5 3.01. Definition. A connection (∇)

Koszul connection (same as Levi-Civita connection?) See Tu, Sec 6 (tangent vector, vector field) \rightarrow (tangent vector)

Ci: linear in the input tangent vector space Cii: linear in the vector field space Ciii: scaling t is the same as scaling w Civ: Liebniz property Cv: continuous vector fields have continous connections

1.17.6 3.02. Coordinates – Christoffel symbols

First, let us see what a connection looks like in coordinates, as some proofs will be easiest that way.

1. Typo? shold be "by Ciii" "by Ci)" is wrong... maybe

1.17.7 3.03. Transformation Formula.

1.17.8 3.04. Definition: vector field along c

a vector field **along** curve c assigns a vector tangent to M at each point of $c(t)$ for $t \in [a, b]$.

From the examples given, this appears to be in concordance with Tu's usage of **along**, in that the vectors do not necessarily have to be tangent to the curve, which we interpret as a submanifold of M when we read Tu.

1.17.9 3.05. Differentiating, Along Curves, Fields Along Curves.

D&P's funny ∇ is DV/dt in Tu – the "covariant derivative"? In IX.1.01, they call Dc^*/dt the acceleration.

In Needham, it is the *intrinsic derivative* (23.2).

Note – I'm using $\tilde{\nabla}$ over ∇ instead of $|$ through ∇ .

We denote the resulting linear map, taking vector fields along c (not vector fields on M) to vector fields along c , by $\tilde{\nabla}_{c*}$. not ∇_{c*} .

1.17.10 3.06. Definition: "vertical"

a vector \vec{v} in the space of derivatives of tangent vectors is *vertical* if its projection into the tangent space is zero.

1.17.11 3.07. Definition: "vertical" and "horizontal" parts of a vector

The vertical part of a vector \vec{v} in the space of derivatives of tangent vectors is the part normal to the tangent space, while the horizontal part of the vector is the part in the tangent space.

The projection P_w defined in this section is the projection to the "vertical." It corresponds to $\langle \vec{v}, \hat{n} \rangle \hat{n}$ in Tu Sec 6.4, Prop 6.8, p 47.

1.17.12 3.08. Language

Koszul connection - section 3.01 Ehresmann connection - split $T_w(TM)$ into horizontal and vertical parts

1.17.13 Exercises 3 & 4

∇ and the funny $\nabla \dots$

1.18 VIII.4. Parallel Transport

1.18.1 4.01. Definition: parallel vector field

A vector field v along $c \in M$ is *parallel* if the horizontal component of its derivative along c^* is zero. In other words, at each point along the curve,

the corresponding vector from the field must maintain the same angle with respect to the tangent to the curve.

This corresponds to Tu (DG) Sec 14.5 Defn 14.13 p 110. $DV/dt = 0$.

1.18.2 what does this even mean?

" v , considered as a curve in TM , has $c^*(t)$ horizontal for all t "

Just like the "rolling" business in this book, I'm not sure the "vertical"/"horizontal" distinction is pedagogically useful. In order to picture nontrivial parallel transport, we need at minimum, a 2D surface embedded in 3D. M is 2D, so $v \in TM$ is also 2D, but $c^*(t) \in T(TM)$ is 4D – hard to picture. Are there 2 horizontal dimensions and 2 vertical dimensions? This is **much** easier to follow in Needham.

See the discussion around Eqn (23.5), p 244 in Needham.

1.18.3 4.02. Theorem: uniqueness of the parallel vector field along c

for any v and c , the parallel vector field is unique.


1.18.4 4.03. Definition: *parallel transport* τ

See Needham 23.2, p 243.

1.19 Bibliography

1.20 Index of Notations

2 NeedhamT VDGF



2.1 Contents

2.2 1.3 Angular excess of a spherical triangle

excess = (sum of angles) - π = A/R^2

2.3 1.5 Constructing geodesics with tape

2.4 2.1 Gaussian curvature in terms of excess angle

2.5 2.2 Circ/Area of a circle – reread this!

2.6 2.3 Local Gauss-Bonnet Theorem

angular excess is the total (area integrated) curvature.

2.7 Chapter 4 The Metric

2.7.1 4.3 Metric of a general surface

2.7.2 Gauss EFG coefficients of the first fundamental form

$E = A^2$ $F = AB\cos\omega$ $G = B^2$

2.7.3 4.4 Metric Curvature Formula

2.7.4 4.5 Conformal Maps

2.8 Chapter 5 Pseudosphere and the Hyperbolic plane

2.8.1 5.2 Tractrix and the Psuedosphere

Tractrix: object dragged along Y axis by a chord of length R. σ = distance along path of object.

2.8.2 5.3 Conformal map of the pseudosphere

$(x,\sigma) \rightarrow (x,y) = x + iy = z$ where x is an angle $[0,2\pi)$ of the revolution and $y \geq 1$ (see 5.4)

2.9 8.3 Newton's curvature formula

2.10 Eqn 8.6, curvature

The area of the shaded sector in fig 8.6 explains the numerator of κ , but it doesn't really explain why the numerator should be that area. Kepler's law?

$$\kappa = (\text{area of swept velocity vector})/|v|^3$$

In Kepler's law, the position vector sweeps out equal areas in equal time.

2.11 Chapter 9 Curves in 3-Space

2.11.1 definition: Torsion

"rate of rotation of the osculating plane is called the torsion, denoted τ ."

2.11.2 definition: binormal of a curve

normal vector to the osculating plane

2.12 Chapter 10 The Principal Curvatures of a Surface

2.12.1 (10.1) Euler's Curvature Formula:

$$\kappa(\theta) = \kappa_1 \cos^2 \theta + \kappa_2 \sin^2 \theta$$

2.12.2 Dupin's indicatrix

nearby slices parallel to the tangent plane are, to first order, conic sections.


2.13 Chapter 11 Geodesics and Geodesic Curvature

2.13.1 Fig [11.2] $K = K_g + K_n$

K_g is "geodesic curvature" IN the surface K_n is "normal curvature" OF the surface

2.14 Chapter 12 The Extrinsic Curvature of a Surface

2.14.1 Figure 12.5: normal vectors on principal and general curves



[12.5] [a] If we move in a principal direction then the normal vector tips in that direction and initially stays within the normal plane; [b] If we move in a general direction then the normal vector immediately tips out of the normal plane.

2.15 Chapter 15 The Shape Operator, S , and ∇_v

2.15.1 15.1 Directional Derivatives (∇_v)

2.15.2 15.2 The Shape Operator S

“is simply defined to be the negative of the directional derivative of n along v ” $S(v) = -\nabla_v n$

2.15.3 15.3 Geometric effect of S

The principal directions are the eigenvectors of the Shape Operator S , and the principal curvatures are the corresponding eigenvalues: $S(e_i) = \kappa_i e_i$.

2.15.4 (15.18) $\kappa(v) = v \cdot S(v)$

where v is a unit tangent vector.

2.15.5 (15.7) sum of curvatures in perpendicular directions

The sum of the curvatures in any two perpendicular directions is equal to the sum of the principal curvatures.

This is a surprising result, given that the curvatures are represented in earlier parts of this book by ellipses.

2.15.6 (15.26) curvature and torsion

$$K_{\text{ext}} = ||S|| = -\tau^2$$

2.15.7 15.9 Classical Terminology and Notation: The Three Fundamental Forms

Note that the fundamental forms are not proper forms (see Act V, Chapter 32 and on.)

- $I(u, v) = u \cdot v$
- $II(u, v) = S(u) \cdot v$
- $III(u, v) = S(u) \cdot S(v)$

2.15.8 19.8 The Road Ahead

In the 3 chapters before this, Needham gives proofs of th

2.16 Chapter 21 An Historical Puzzle

2.16.1 GR and parallel transport

Einstein's success was all the more remarkable, and remains all the more puzzling, because he achieved it before Levi-Civita—pictured in [21.1]—discovered the concept of parallel transport, which did not occur until 1917!

2.17 Chapter 22 Extrinsic Constructions

2.17.1 Projection (not rotation) of w into T_q

$$w_{||} = P[w] = w - (w \cdot n)n$$

2.17.2 Rotation of w into T_q

\tilde{w} . But in the limit as $\epsilon \rightarrow \infty$, rotation and projection are the same.

2.18 Chapter 23 Intrinsic (covariant) Constructions

2.18.1 23.2 The Intrinsic (aka, “Covariant”) Derivative

aka Levi-Civita connection (Koszul connection?)

2.18.2 (Eqn 22.1) Intrinsic derivative, (D_v , the covariant derivative)

$\epsilon D_v w = w(q) - w_{||}(p \rightarrow q)$ both w 's on the RHS are in T_q

2.18.3 Fig 23.3

“here w is growing in length and rotating counterclockwise as it moves along G .”

2.18.4 D_v is also called the Levi-Civita Connection.

2.18.5 intrinsic derivative $D_v w$

vs the definition on the previous page, here $w(q)$ is parallel transported back to p [$w_{||}(q \rightarrow p)$] and then $w(p)$ is subtracted. (then $\lim \epsilon \rightarrow \infty$)

2.18.6 Here is an extrinsic way of looking at the intrinsic derivative.

2.18.7 Eqn 23.3 $D_v w$

same as Eqn 22.1, with $\nabla_v w$ dropped in for w

In other words, to obtain $D_v w$ we take the full rate of change $\nabla_v w$ in R^3 , then subtract out the part that is not tangent to the surface, thereby leaving behind the part that is intrinsic to the surface.

2.18.8 Compare these with Ci-Cv in D&P p 217

it is much simpler to think of **flattening onto the tabletop** the strip surrounding [curve] K , together with the vector fields x, y, z , for then D_v simply is ∇_v .

2.19 1-forms used to be called covariant vectors or covectors

2.20 Further Reading


2.20.1 Intuitive Topology, by V. V. Prasolov.

2.20.2 Three-Dimensional Geometry and Topology, by William P. Thurston.

2.20.3 Differential Geometry: A Geometric Introduction, by David W. Henderson.

2.20.4 Gauge Fields, Knots and Gravity, by John Baez and Javier P. Muniain.

3 TuLW DifferentialGeometry



3.1 §1 Riemannian Manifolds

3.1.1 1.3 Riemannian Metrics

A Riemannian metric maps a continuous vector field onto a continuous function.

3.1.2 Example 1.11: Torus

On a torus in \mathbb{R}^3 vs a torus as “the quotient space of a square with the opposite edges identified”

We will show later that there is no isometry between these two Riemannian manifolds with the same underlying torus.

3.1.3 partition of unity

What is this? The definition is unhelpful in describing the meaning of this term.

3.1.4 Theorem 1.12. On every manifold M there is a Riemannian metric.

3.2 §2 Curves

3.2.1 2.3 Signed Curvature of a Plane Curve

$$T' = \gamma''(s) = \kappa \vec{n}, \text{ so } \kappa = \langle T', \vec{n} \rangle = \langle \gamma'', \vec{n} \rangle$$

3.2.2 Curvature formulae

see Needham,

3.3 §4 Directional Derivatives in Euclidean Space

3.3.1 Torsion of the directional derivative D

If $D_X Y - D_Y X = [X, Y]$, then the torsion $T(X, Y) = D_X Y - D_Y X - [X, Y]$ is 0.

This only applies to \mathbb{R}^n .

3.3.2 Curvature of the directional derivative

3.3.3 Definition 4.8. "On" vs "Along" a submanifold

Along includes On, where On is tangent to the submanifold.

3.3.4 Notation $X(M)$ and $\Gamma(TM|_N)$

- $\mathfrak{X}(M)$ is the set of all C^∞ vector fields **on** manifold M
- $\Gamma(TM|_N)$ is the set of all C^∞ vector fields **along** a submanifold N in a manifold M .
- $\mathcal{F} = C^\infty(M)$ is the ring of C^∞ functions on M

3.3.5 In a submanifold of \mathbb{R}^n , torsion "no longer makes sense"

because: X is a vector field **on** M Y is a vector field **along** M

3.4 §5 The Shape Operator

See Needham Chapter 15.

3.4.1 Defn: regular point – need a picture

3.4.2 5.2 The Shape Operator

$L_p(X_p) = -D_{X_p}N$, which is in the tangent plane T_pM and points opposite the direction in which N changes wrt X_p .

3.4.3 Is the usage of "on" here the same as before?

"let N be a C^∞ unit normal vector field on M (Figure 5.1)."

3.4.4 Proposition 5.3. The shape operator is self-adjoint:

3.4.5 Proposition 5.5: Curvature is given by the second fundamental form

$$\kappa(X_p) = \langle L(X_p), X_p \rangle = II(X_p, X_p)$$

3.4.6 Proposition 5.6: Principal directions and curvatures

- Principal directions: eigenvectors of L
- Principal curvatures: eigenvalues of L

3.4.7 5.4 The First and Second Fundamental Forms

Defined for smooth surfaces in \mathbb{R}^3 .

- first fundamental form: inner product $\langle X, Y \rangle_E := \langle e_1, e_1 \rangle_F := \langle e_1, e_2 \rangle_G := \langle e_2, e_2 \rangle$
- second fundamental form: $II(X_p, Y_p) = \langle L(X_p), Y_p \rangle_e := II(e_1, e_1) f := II(e_1, e_2) g := II(e_2, e_2)$

3.5 §6 Affine Connections

We will see in a later section that there are infinitely many affine connections on any manifold. On a Riemannian manifold, however, there is a unique torsion-free affine connection compatible with the metric, called the **Riemannian** or **Levi-Civita** connection.

3.5.1 No canonical basis for $T_p M$ not embedded in a Euclidean space

Formula 4.2 here is the directional derivative.

3.5.2 Definition 6.1. An affine connection

3.5.3 6.2 Torsion and Curvature

“There does not seem to be a good reason for calling $T(X, Y)$ the torsion” Is this not the same torsion that appears in Needham Chapter 9?

3.5.4 Definition 6.4: Riemannian or Levi-Civita connection

“...is an affine connection that is torsion-free and compatible with the metric.”

3.5.5 Theorem 6.6. On a Riemannian manifold there is a unique Riemannian connection.

3.5.6 6.4 Orthogonal Projection on a Surface in \mathbb{R}^3

pr_p is the projection from $T_p \mathbb{R}^3$ to the tangent space of M at p .

3.6 §7 Vector Bundles

“Thus the set $X(M)$ has two module structures, over R and over F .”

“We will try to understand F -linear maps from the point of view of vector bundles. The main result (Theorem 7.26) asserts the existence of a one-to-one correspondence between F -linear maps $\rho : (E) \rightarrow (F)$ of sections of vector bundles and bundle maps $\rho : E \rightarrow F$.”

3.6.1 7.1 Definition of a Vector Bundle

“A vector bundle, intuitively speaking, is a family of vector spaces that locally “looks” like $U \times \mathbb{R}^r$.”

We need a picture of the terms defined here.

3.7 13.1 Theorem

We call DV/dt the covariant derivative (associated to ∇) of the vector field V along the curve $c(t)$ in M .

3.8 14.5 Parallel Translation

4 Littlejohn Manifolds

4.1 Notes for page 3

4.2 (typo)

should be: $x'^\mu = \psi_{ij}(x^\nu)$

4.3 (typo)

should be: $x^\nu = \psi_{ij}^{\leftarrow}(x'^\mu)$

4.4 8. Tangent Vectors

4.4.1 Eqn 11

4.5 9. Equivalence Classes of Curves

4.6 10. Tangent Vectors in Coordinates

"convective derivative" (see 4.4.1) "scalar" == "scalar field" in this context?

4.6.1 Eqn 16

4.6.2 tangent vector at p

Tangent vector as a =first order, linear, partial differential operator=

$$\vec{X} = \Sigma_i X^i \frac{\partial}{\partial x^i} |_p$$

4.7 12. Covectors

4.7.1 Eqn 21 $df|_p$

$df|_p: T_p M \rightarrow \mathbb{R}: X \mapsto Xf$ where Xf is defined in 4.6.1

4.7.2 $df|_p$ is an operator on vectors – it is not small or "infinitesimal"

"The most confusing thing for novices about this definition is that there is nothing small or "infinitesimal" about $df|_p$. In traditional theoretical physics the notation df usually denotes a small increment in the function f ."

4.7.3 13. Differential of Coordinates

5 BachmanD: A Geometric Approach to Differential Forms

5.1 1.2 Generalizing the integral

5.2 1.3 differential form in a nutshell

"A differential form is precisely a linear function which eats vectors, spits out numbers and is used in integration."

5.3 3 Forms

5.3.1 3.1 Coordinates for vectors

"The key to understanding the difference between L and $T_p L$ is their coordinate systems."

1. notation $\langle \cdot, \cdot \rangle$

We have switched to the notation " $\langle \cdot, \cdot \rangle$ " to indicate that dt we are not talking about points of P anymore, but rather vectors in $T_p P$.

5.3.2 3.2 1-Forms

Evaluating a 1-form on a vector is the same as projecting onto some line and then multiplying by some constant.

Evaluating a 1-form on a vector is the same as projecting onto each coordinate axis, scaling each by some constant and adding the results.

5.3.3 3.3 Multiplying 1-forms

The wedge product is a product operation on 1-forms that is closed, ie, produces another 1-form.

“we will use the symbol “ \wedge ” (pronounced “wedge”) to denote multiplication”

1. Notation $(\cdot, \langle \cdot, \cdot \rangle, \llbracket \cdot \rrbracket)$: x,y plane $\langle \cdot, \cdot \rangle$: dx,dy plane $\llbracket \cdot \rrbracket$: ω, ν plane
2. wedge product justification or reasoning

Do we know of a way to take these vectors and get a number?
 Actually, we know several, but the most useful one turns out to be the area of the parallelogram that the vectors span.
 This is precisely what we define to be the value of $(V1, V2)$

Still – why the skew-symmetric product vs any other one? What is special about a/the skew-symmetric operator?

3. 1-form in a nutshell

Evaluating ω on the pair of vectors $(V1, V2)$ gives the area of parallelogram spanned by $V1$ and $V2$ projected onto the plane containing the vectors $\langle \cdot \rangle$ and $\langle \cdot \rangle$, and multiplied by the area of the parallelogram spanned by $\langle \cdot \rangle$ and $\langle \cdot \rangle$.

4. 2-form in a nutshell

Every 2-form projects the parallelogram spanned by $V1$ and $V2$ onto each of the (2-dimensional) coordinate planes, computes the resulting (signed) areas, multiplies each by some constant, and adds the results.

Henceforth, we will define a 2-form to be a bilinear, skew-symmetric, real-valued function on $T_p \mathbb{R}^n \times T_p \mathbb{R}^n$.

5.3.4 3.4 2-Forms on $T_p\mathbb{R}^3$

5.3.5 3.5 2-Forms and 3-forms on $T_p\mathbb{R}^4$

5.3.6 3.6 n-Forms

length, area and volume are all signed quantities?

5.3.7 3.7 Algebraic computation of products

5.4 4 Differential Forms

6 LeeJM _SmoothManifolds (+18)

6.1 Contents

6.2 Proposition 2.25 (Existence of Smooth Bump Functions).

These are mentioned in Tu. Maybe the exposition here is useful (but I haven't read it yet.)

6.3 Chapter 3 Tangent Vectors

6.4 Chapter 8 Vector Fields

In the [third] section we introduce the Lie bracket operation, which is a way of combining two smooth vector fields to obtain another. Then we describe the most important application of Lie brackets: the set of all smooth vector fields on a Lie group that are invariant under left multiplication is closed under Lie brackets, and thus forms an algebraic object naturally associated with the group, called the Lie algebra of the Lie group.

6.4.1 Vector Fields on Manifolds

6.4.2 Example 8.12. cylindrical orthonormal frame on $\mathbb{R}^2\{0\}$

6.4.3 Vector Fields and Smooth Maps

first mention of "-related".

Suppose $F : M \rightarrow N$ is smooth and X is a vector field on M ; and suppose there happens to be a vector field Y on N with the property that for each $p \in M$; $dF_p(X_p) = Y_{F(p)}$. In this case, we say the vector fields X and Y are **F-related** (see Fig. 8.3).

The next proposition shows how F-related vector fields act on smooth functions.

6.4.4 Lie Brackets

6.4.5 Lemma 8.25. Lie bracket of smooth vector fields...

“Lemma 8.25. The Lie bracket of any pair of smooth vector fields is a smooth vector field.”

Geometric interpretation comes in Chapter 9

“of limited usefulness for computations”: $[X, Y]_p f = X_p(Yf) - Y_p(Xf)$

6.4.6 Proposition 8.26 (Coordinate Formula for the Lie Bracket).

“an extremely useful coordinate formula for the Lie bracket”: $[X, Y] = \left(X^i \frac{\partial Y^j}{\partial x^i} - Y^i \frac{\partial X^j}{\partial x^i} \right) \frac{\partial}{\partial x^j}$ or more concisely, $[X, Y] = (XY^j - YX^j) \frac{\partial}{\partial x^j}$

6.4.7 Example 8.27

Decent concrete example.

6.4.8 see D&P 3

$$[fX, gY] = fg[X, Y] + (fXg)Y - (gYf)X$$

$$fXgY - gYfX = fXgY + fgXY - gYfX - gfYX = (fXg)Y - (gYf)X + (fgXY - gfYX)$$

6.5 Chapter 9 Integral Curves and Flows

6.5.1 Flowouts

1. Theorem 9.22 (Canonical Form Near a Regular Point). Randy, read this.

6.5.2 Lie Derivatives

1. Fig. 9.13 The Lie derivative of a vector field
2. Theorem 9.38 Lie derivative == Lie bracket

6.5.3 Commuting Vector Fields

1. Theorem 9.42. Equivalent statements on vector fields

For smooth vector fields V and W on a smooth manifold M ; the following are equivalent: (a) V and W commute. (b) W is invariant under the flow of V . (c) V is invariant under the flow of W .

Consider $\rho = x/r \mathbf{i} + y/r \mathbf{j}$ # radial vec field "E1" p179 $\theta = -y/r \mathbf{i} + x/r \mathbf{j}$ # unit rotation "E2" p179 $\phi = -y \mathbf{i} + x \mathbf{j}$ # disc rotation "W" p207

$$[E1, E2] = -E2 \quad [E1, W] = 0$$

- (a) follow E1 out in r for E2, the vector at positions in the flow of E1 are constant Q : Is E2 invariant under the flow of E1 ?? [A: no]

for W , the vector at positions in the flow of E1 are increasing in magnitude. Q : Is W invariant under the flow of E1 ?? [A: yes]

- (b) follow a streamline in E2 or W The change in ρ is tangential to the flow. In W , $d\rho$ is constant, but in E2, $d\rho$ is smaller at larger radial positions.

- useful derivatives $\mathbf{i}(x/r) = y^2/r^3$ $\mathbf{i}(y/r) = -xy/r^3 = \mathbf{j}(x/r)$ $\mathbf{j}(y/r) = x^2/r^3$

$$\begin{aligned} [E1, W] &= (x/r \mathbf{i} + y/r \mathbf{j})(-y \mathbf{i} + x \mathbf{j}) - (-y \mathbf{i} + x \mathbf{j})(x/r \mathbf{i} + y/r \mathbf{j}) \\ &= (x/r) \mathbf{i}(x) \mathbf{j} - (y/r) \mathbf{j}(y) \mathbf{i} - [-y \mathbf{i}(x/r) \mathbf{i} - y \mathbf{i}(y/r) \mathbf{j} + x \mathbf{j}(x/r) \mathbf{i} + x \mathbf{j}(y/r) \mathbf{j}] \\ &= [(x/r) \mathbf{j} - (y/r) \mathbf{i}] - [-(y/r)^3 - x^2/r^3] \mathbf{i} - [xy^2/r^3 + (x/r)^3] \mathbf{j} \\ &= [(x/r) \mathbf{j} - (y/r) \mathbf{i}] - (y/r)[- (y/r)^2 - x^2/r^2] \mathbf{i} - (x/r)[y^2/r^2 + (x/r)^2] \mathbf{j} \\ &= [(x/r) \mathbf{j} - (y/r) \mathbf{i}] - [(x/r) \mathbf{j} + (y/r) \mathbf{i}] = E2 - E1 \end{aligned}$$

$$\begin{aligned} [E1, E2] &= 0 - [-y/r \mathbf{i} + x/r \mathbf{j}][x/r \mathbf{i} + y/r \mathbf{j}] = 0 - [(-y/r) \mathbf{i}(x/r) + (x/r) \mathbf{j}(x/r) \mathbf{i} - (-y/r) \mathbf{i}(y/r) + (x/r) \mathbf{j}(y/r) \mathbf{j}] \\ &= 0 - [(-y/r)y^2/r^3 \mathbf{i} - (x/r)xy/r^3 \mathbf{i} + xy^2/r^4 \mathbf{j} + x^3/r^4 \mathbf{j}] = 0 - [-y^3/r^3 \mathbf{i} - x^2y/r^3 \mathbf{i} + xy^2/r^3 \mathbf{j} + x^3/r^3 \mathbf{j}] / r = 0 - [-y/r \mathbf{i} + x/r \mathbf{j}] / r = -E2/r \end{aligned}$$

2. Theorem 9.44. Smooth vector fields commute if and only if their flows commute. D&P VII.7.02

3. Example 9.45 (Commuting and Noncommuting Frames).

Example 8.12. The standard coordinate frame is a global orthonormal frame on \mathbb{R}^n . For a less obvious example, consider the smooth vector fields defined on $\mathbb{R}^2 \setminus \{0\}$ by

$$E_1 = \frac{x}{r} \frac{\partial}{\partial x} + \frac{y}{r} \frac{\partial}{\partial y}, \quad E_2 = -\frac{y}{r} \frac{\partial}{\partial x} + \frac{x}{r} \frac{\partial}{\partial y}, \quad (8.3)$$

where $r = \sqrt{x^2 + y^2}$. A straightforward computation shows that (E_1, E_2) is an orthonormal frame for \mathbb{R}^2 over the open subset $\mathbb{R}^2 \setminus \{0\}$. Geometrically, E_1 and E_2 are unit vector fields tangent to radial lines and circles centered at the origin, respectively. //

4. Theorem 9.46 (Canonical Form for Commuting Vector Fields). D&P Thm VII.7.04

6.6 Chapter 11 The Cotangent Bundle

we define the differential of a real-valued function as a covector field (a smooth section of the cotangent bundle); it is a coordinate-independent analogue of the gradient.

6.6.1 Prop 11.8 $V \cong V^{**}$

“there is no canonical isomorphism $V \cong V^*$.”

6.6.2 Tangent Covectors on Manifolds

Thus it became customary to call tangent covectors **covariant vectors** because their components transform in the same way as (“vary with”) the coordinate partial derivatives, the Jacobian matrix multiplying the objects associated with the “new” coordinates to obtain those associated with the “old” coordinates. Analogously, tangent vectors were called **contravariant vectors**, because their components transform in the opposite way.

6.6.3 contravariant transformation

6.6.4 covariant transformation

6.6.5 Prop 11.18 (coordinate covectors are dx^j 's)

In other words, the coordinate covector field λ^j is none other than the differential dx^j .

See D&P VII.4.01

6.6.6 Proposition 11.23 (Derivative of a Function Along a Curve).

6.7 Chapter 12 Tensors

We deal primarily with covariant tensors, but we also give a brief introduction to contravariant tensors and tensors of mixed variance.

7 TuLW Introduction to Manifolds

7.1 2.1 Directional derivative

In D&P, this is denoted $D_{\text{pf}}(v)$ for the directional derivative in direction v at point p of function f .

7.2 Defn: germ

Equivalence class of functions on an neighborhood C^{inf}_p is the set of all germs of C^{inf} functions on \mathbb{R}^n at p .

7.3 Defn: derivation

Linear map $C^{\text{inf}}_p \rightarrow \mathbb{R}$ satisfying the Leibniz rule is a derivation

7.4 2.4 Vector fields

See D&P VII.4.02 Contravariant vectors. The 2nd paragraph above encapsulates the corresponding section in D&P in a nutshell.

7.5 3.3 k-covectors

"A 1-covector is simply a covector". A k -covector appears to be a tensor of covariant degree k

7.6 Lemma 3.11 $\tau(\sigma f) = (\tau\sigma)f$

$(\sigma f)(v_1, \dots, v_k) = f(v_{\sigma(1)}, \dots, v_{\sigma(k)})$ if $\sigma = (1, m, n)$, then $\sigma(1) = m$, $\sigma(m) = n$, $\sigma(n) = 1$

- concretely, if $\sigma = (1, 3, 2)(4, 5)$ and $\tau = (1, 5, 3, 2)$ then $(\sigma f)(v_1, v_2, v_3, v_4, v_5) = f(v_3, v_1, v_2, v_5, v_4)$
and $\tau(\sigma f)(v_1, v_2, v_3, v_4, v_5) = (\sigma f)(v_5, v_1, v_2, v_4, v_3) = f(v_2, v_5, v_1, v_3, v_4)$

$$\tau \sigma = (1,5,3,2) (1,3,2) (4,5) = (1,2,5,4,3)$$

so this checks out.

7.7 3.7 Wedge product

Tu motivates the definition of the wedge product by wanting to have a product that preserves the property of being alternating on alternating functions (on a vector space). Does this make the wedge product the group multiplication on alternating functions? No – the identity is not alternating, so the wedge can't get you there.

7.8 Defn: graded algebra over field K

The algebra "can be written as a direct sum of vector spaces over K such that $A^k \times A^l \mapsto A^{k+l}$.

7.9 4 Differential forms on \mathbb{R}^n

7.10 4.6 Applications to Vector Calculus

related to exact sequences

grad: scalar \rightarrow vector curl: vector \rightarrow vector div : vector \rightarrow scalar

7.11 4.7 Convention on Subscripts and Superscripts

vector fields: subscripts	e_1, \dots, e_n	contravariant
differential forms: superscripts	$\omega^1, \dots, \omega^n$	covariant
coordinate functions (0-forms):	x^1, \dots, x^n	covariant
differentials of coord functs (1-forms):	dx^1, \dots, dx^n	covariant
coordinate vector fields	$d/dx^1, \dots, d/dx^n$	contravariant
coordinate vector fields (alt form)	$\delta_1, \dots, \delta_n$	contravariant

- coefficient functions

of a vector field a^i
of a differential form b_j

7.12 Sec 5: Manifolds

smooth manifolds are the focus of this book

"maximal C^∞ atlas" make a topological manifold into a smooth manifold.

7.13 5.1 Topological Manifolds

defn: **second countable** A topological space with a countable basis.

7.14 5.3 Smooth Manifolds - maximal atlas defn

A maximal atlas on a locally Euclidean space is not contained in any larger atlas.

7.15 Definition 5.9 smooth manifold

A smooth manifold is a topological manifold with a maximal atlas. A maximal atlas is also called a **differentiable structure** "In practice, to check that a topological manifold M is a smooth manifold, it is not necessary to exhibit a maximal atlas. The existence of any atlas on M will do"

7.16 Smooth manifold conditions

to show that a topological space M is a C manifold, it suffices to check that:

1. M is Hausdorff and **second countable**,
2. M has a C atlas (not necessarily maximal).

7.17 5.4 Examples of Smooth Manifolds

- Euclidean space
- Open subset of a manifold
- Manifolds of dimension zero
- Graph of a smooth function
- General linear groups (nonzero determinant matrices)
- Unit circle in the (x,y) plane
- Product manifold

7.18 §6 Smooth Maps on a Manifold

7.19 6.3 Diffeomorphisms

7.20 6.6 Partial Derivatives

7.21 6.7 The Inverse Function Theorem

A function on an affine space or manifold is invertible at p iff the Jacobian determinant at p is nonzero

7.22 Chapter 3: The tangent space

7.23 bump functions?

These are covered in section 13, p 140

7.24 Theorem 20.4.

If X and Y are C vector fields on a manifold M , then the Lie derivative $L_X Y$ coincides with the Lie bracket $[X, Y]$.

7.25 A.4 First and Second Countability

7.25.1 Lemma A.10. Every open set in \mathbb{R}^n contains a rational point.

7.25.2 Prop A.11 collection of open balls w/ rational centers/radii is a basis for \mathbb{R}^n

7.25.3 Defn A.12 second countable \iff countable basis

7.25.4 Defn A.15 basis of neighborhoods, first countable

- A **basis of neighborhoods at p** is a collection of neighborhoods of p such that $p \in B_\alpha \subset U$
- Topological space S is **first countable** if it has a countable basis of neighborhoods at every point $p \in S$.
- Every second countable space is first countable.

7.26 List of Notations

8 Carroll - An Introduction to General Relativity

- 8.1 1. SR and flat Spacetime
- 8.2 2. Manifolds
- 8.3 3. Curvature
- 8.4 4. Gravitation
- 8.5 5. Schwarzschild's solution
- 8.6 6. More general black holes
- 8.7 7. Perturbation theory & Gravitational radiation
- 8.8 8. Cosmology
- 8.9 9. QFT in curved spacetime
- 8.10 A. Maps between manifolds
- 8.11 B. Diffeomorphisms and Lie derivatives
- 8.12 C. Submanifolds
- 8.13 D. Hypersurfaces
- 8.14 E. Stokes's Theorem
- 8.15 F. Geodesic congruences
- 8.16 G. Conformal transformations
- 8.17 H. Conformal diagrams
- 8.18 I. The parallel propagator
- 8.19 J. Noncoordinate bases
- 8.20 Bibliography
- 8.21 Index