





n=13 seeds: learn -0.068 (p=1.2e-04), extinguish +0.124 (p=2.4e-04)
recovery frac. 1.84 [1.38,2.25] (>1 = overshoot past baseline)
mean peak distance to reward A
0.65
0.60
0.55
0.50
0.45
0.40
pre-learning
(pred)
learned
(postLearn)
extinguished
(postExt)


Reliability drop is concentrated in FAR cells
(divergence from Krishnan's zone-specific drop)
split-half reliability (r)
1.0
0.8
0.6
0.4
0.2
0.0
p=0.008
p=0.893
p=0.002
postLearn
postExt
all
(n=13)
near
(n=13)
far
(n=13)


n=13 seeds: near/far contrast collapses
median 19.3x -> 1.4x (p=0.017)
head output (proximity readout)
1.0
0.8
0.6
0.4
0.2
0.0
postLearn
postExt
analytic target
(same both periods)
0.1
0.2
0.3
0.4
0.5
0.6
0.7
0.8
distance to reward A


Reliability drop under extinction
reliability change (%)
0
-10
-20
-30
-40
model
(n=13 seeds)
Krishnan 2022
(n=3 mice)


Fraction of enrichment lost
(model 110%: full loss + overshoot;
Krishnan 81%: substantial, partial)
% of reward-related
enrichment lost
100
80
60
40
20
0
model
(near-A fraction)
Krishnan 2022
(field-density
enrichment)