


STAT7203 & STAT2203
Week 6 Exercises — Practice Mid-semester exam

1. Let (A_1, \dots, A_4) be a random permutation of the integers $\{1, 2, 3, 4\}$ with all permutations equally likely. For $i < j$ define the random variables

$$X_{ij} = \begin{cases} 1, & \text{if } A_i > A_j \\ 0, & \text{otherwise.} \end{cases}$$

Then $\mathbb{P}(X_{12} = 0, X_{13} = 0)$ equals

- (a) $\frac{1}{24}$
 - (b) $\frac{1}{8}$
 - (c) $\frac{1}{6}$
 - (d) $\frac{1}{4}$
 - (e) $\frac{1}{3}$
2. Consider the system below comprised of three components. The system is working if there is a path from left to right through working components. The components fail independently of one another. The probability that a given component has failed in six months is 0.4.



The probability that the system is working at the end of six months

- (a) 0.256
- (b) 0.36
- (c) 0.504
- (d) 0.64
- (e) 0.84

3. A bank uses machine learning to detect credit card fraud. Their method correctly identifies a fraudulent transaction with 0.9 probability and it correctly identifies non-fraudulent transactions with 0.975 probability. Among the bank's customers fraudulent transactions account for 0.5% of all credit card transactions. The probability that a randomly selected transaction is fraudulent given it has been identified as fraudulent using the machine learning algorithm is approximately

- (a) 0.0045
- (b) 0.0046
- (c) 0.153
- (d) 0.48
- (e) 0.9






4. Suppose that X has the below probability mass function

x	0	1	2	3
$\mathbb{P}(X = x)$	0.3	0.5	0.1	0.1

The variance of X is

- (a) 0.8
 - (b) 0.89
 - (c) 1
 - (d) 1.34
 - (e) 1.8
5. Suppose that X_1, X_2, \dots, X_{20} are independent random variables each having a $\text{Geometric}(\frac{1}{4})$ distribution. Let $Y = \frac{1}{20} \sum_{i=1}^{20} X_i$. The standard deviation of Y is approximately
- (a) 0.173
 - (b) 0.6
 - (c) 0.775
 - (d) 2.683
 - (e) 12

6. Which of the following graphs depicts the cumulative distribution function of a $\text{Binomial}(3, \frac{1}{2})$ distribution?



7. Suppose X_1 and X_2 are two independent random variables each having a uniform distribution on the integers $1, 2, \dots, 6$. The probability that $X_1 - X_2 = 0$ given that $X_1 + X_2 = 6$ is
- (a) $\frac{1}{36}$
 - (b) $\frac{1}{12}$
 - (c) $\frac{1}{6}$
 - (d) $\frac{1}{5}$
 - (e) $\frac{1}{4}$
8. Consider the pair of random variables (X, Y) . Suppose that marginally $X \sim \text{Binomial}(2, \frac{1}{2})$ and $Y \sim \text{Binomial}(2, \frac{3}{4})$. If $\mathbb{P}(X > Y) = 0$ and $\mathbb{P}(X = 0, Y = 2) = \frac{1}{16}$, then $\mathbb{P}(X = 1, Y = 1)$ equals
- (a) $\frac{1}{16}$
 - (b) $\frac{1}{8}$
 - (c) $\frac{3}{16}$
 - (d) $\frac{1}{4}$
 - (e) $\frac{5}{16}$
9. Suppose X_1, X_2 and X_3 independent random variables, each having a $\text{Uniform}(\{1, 2, 3\})$ distribution. Let $Y_1 = X_1 + X_2$ and $Y_2 = X_1 + X_3$. The covariance between Y_1 and Y_2 is
- (a) 0
 - (b) 0.667
 - (c) 2
 - (d) 4
 - (e) 4.667
10. Suppose $X_1 \sim \text{Poisson}(\frac{3}{2})$ and $X_2 \sim \text{Poisson}(\frac{5}{2})$ are two independent random variables. Let $Y = X_1 + X_2$. The conditional expectation $\mathbb{E}[X_1 | Y = 3]$ is
- (a) 1
 - (b) 1.125
 - (c) 1.25
 - (d) 1.375
 - (e) 1.5

11. Suppose X and Y are two independent random variables such that $X \sim \text{Bernoulli}(p)$ and $Y \sim \text{Poisson}(\lambda)$. Define $Z = XY$. Then the moment generating function of Z is:

- (a) $M_Z(s) = \exp(p\lambda(e^s - 1))$
- (b) $M_Z(s) = 1 - p + p \exp(\lambda(e^s - 1))$
- (c) $M_Z(s) = (1 - p + ps) \exp(\lambda(e^s - 1))$
- (d) $M_Z(s) = (1 - p + pe^s)^\lambda$
- (e) $M_Z(s) = \left(\frac{pe^s}{1 - p + pe^s} \right)^\lambda$

12. Suppose the random variable X has probability generating function

$$M_X(s) = \frac{1 + e^s}{4 - 2e^s}.$$

Then the expected value of X is:

- (a) -1
- (b) 0.375
- (c) 1
- (d) 1.5
- (e) 3