

1 Introduction

Modal logic extends classical propositional logic by introducing modal operators, most notably \Box ("necessarily") and \Diamond ("possibly"). In the standard Kripke semantics, these operators are interpreted over a set of possible worlds connected by an accessibility relation. In the following, we give an example for \Box .

Example 1.0.1. Let $\phi = \Box p$ and $M = (W, R, V)$ with $W = \{w_1, w_2, w_3, w_4, w_5\}$, $R = \{(w_1, w_2), (w_2, w_3), (w_2, w_4), (w_2, w_5)\}$ and $V(p) = \{w_3, w_4, w_5\}$. (See figure 1.)




Figure 1: Intuitively, $\Box p$ is interpreted as "In all accessible worlds, p is true". Here we can see $\Box p$ is true at w_2 , because all accessible worlds from w_2 (in blue lines), make p true.

However, the term 'modal logic' is often used more broadly for a family of related systems. This include for example temporal or epistemic logic. Due to its relational nature, modal logic is a powerful tool for analyzing structured systems, where the relationships between states or agents are essential. It has wide-ranging applications—for instance, in formal verification. For example in model checking, temporal logic is used to describe the behavior of systems over time. In multi-agent systems, epistemic logic plays a key role in reasoning about knowledge and beliefs.

1.1 Topological semantics

An idea to generalize Kripke semantics was topology. The study of topological semantics in modal logic was initiated by McKinsey and Tarski in 1944 [3]. To assign meaning to modal formulas, we use topological space, which is a pair consisting of a domain X and a collection of open sets, where each set describes which points in X are considered "nearby" to each other. A topological model is a topological space with a valuation V , where we assign subsets of the domain to the variables. \Box and \Diamond are interpreted differently in topological context; they behave more like the interior and closure operators like in topology [16].

Example 1.1.1. Let $W = \{1, 2, 3\}$, $\tau = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, W\}$ (where τ is the collection of open sets) and $V(p) = W$. Furthermore, let $\phi = \Box p$.




Figure 2: Each ellipse represents an open set. Intuitively, $\Box p$ is satisfied at a point w , if we can find an open set U s.t. $w \in U$ and $U \subseteq V(p)$. This is true for 1 and 2 because we can find $U = \{1, 2\} \subseteq V(p)$ (where $V(p)$ is marked as the blue).

It is well known that the logic for any topological space is $S4$, where $S4$ is the modal logic with the transitivity ($\Box p \rightarrow \Box \Box p$) and reflexivity axiom ($\Box p \rightarrow p$). This holds because, under topological semantics, the interpretation of \Box naturally satisfies both reflexivity and transitivity. So, if we assume a topological space satisfies $\Box p$ at point w with some valuation $V(p)$, then we can find an open set U around w where $w \in U$. Therefore, $w \in V(p)$ which shows reflexivity. On the other hand, for any point in U , we can again find an open set (U itself) witnessing that $\Box p$ holds there. Hence, transitivity is also satisfied.

Topological semantics provides a rich, and intuitive semantics for $S4$ and related logics. It has a variety of applications e.g. spatial reasoning or program verification. Despite its rich expressiveness, it can not fully capture Kripke semantics.

1.2 Neighborhood frames

That is why Neighbourhood semantics (see [11] for an introduction) as a generalization of Kripke semantics for modal logic was introduced (independently) by Dana Scott [9] and Richard Montague [10]. Neighborhood frames are similar to topological spaces. Instead of collection of open sets, we have a neighborhood function, which assigns to each point a set of subsets of the domain. These subsets are also called neighbourhoods. The Neighborhood semantics only differs slightly in \Box and \Diamond . The following example demonstrates it.

Example 1.2.1. Let $W = \{1, 2, 3\}$,

$$\tau(x) = \begin{cases} 1 \rightarrow \{\{1\}, \{3\}, W\} \\ 2 \rightarrow \{\{2\}, \{1, 2\}, W\} \\ 3 \rightarrow \{\{3\}, \{1, 3\}\} \end{cases}$$

and $V(p) = \{1, 2\}$. Assume $\phi = \Box p$.




Figure 3: The green ellipses represents neighbourhoods from 1, the blue ellipses those of 2 and red ellipses those of 3. $\Box p$ is satisfied at a point w , if there exists a neighborhood $U \in \tau(w)$ s.t. $U \subseteq V(p)$. This is only true for 2.

Remark : For truth, we don't require that w has to be in U . Therefore, it is not necessary that all neighborhood frames validate reflexivity and hence S4.

Neighborhood semantics is more general than Kripke semantics and in the case of normal reflexive and transitive logics coincides with topological semantics. The original motivation for introducing these general models was to provide a semantics for non-normal modal logics. Beyond their theoretical motivation, interest in topological semantics and neighborhood frames has grown considerably, partly due to its applications in artificial intelligence.

1.3 Content

Oftentimes, it is necessary to combine frames for different modal logics into a complex frame. The natural way of doing that is a product construction. For Kripke frames, the resulting product is the Cartesian product of the two frames with two accessibility relations. For topological semantics, the product of topological spaces as bi-topological spaces with so-called horizontal and vertical topologies have been considered. In a similar fashion, the product of neighborhood frames was introduced by Sano in [8].

For our main question, we consider the neighborhood product additionally with the standard neighborhood product. We will answer whether the logic with three copies of T with the interaction axiom $\Box p \rightarrow \Box_1 p \wedge \Box_2 p$ can be fully characterized as the logic a product of neighbourhood frames with standard topology, where each frame validates T . The question was originally motivated by two papers where the authors showed in topological semantics a similar version of the question but for $S4$. In the other, it was proven that for any $L_1, L_2 \in \{D, D4, T, S4\}$, that their logic together can be characterized as the logic of a product of neighbourhood frames but without standard neighborhood product. In this work, we start with a brief recap of necessary notation and theorems in Kripke semantics and then introduce topological spaces and neighborhood frames. We will reprove the theorems from the authors in a more detailed way and then use similar techniques to answer our main question. Finally, we conclude with a

discussion of potential directions for future research related to the topic. This thesis primarily builds upon the works "Multimodal Logics of Products of Topologies" [1] and "Modal Logic of Products of Neighborhood Frames" [2].

2 Preliminaries

This chapter is primarily devoted to the semantics of Kripke. In addition, we define the notions of product frames and multimodal logic. We also present fundamental results in modal logic, such as Filtration Theorem and Sahlqvist Theorem which play a role in our thesis. The content is mainly based on the Modal Logic book [7]. Remark: natural numbers \mathbb{N} are without 0.

Definition 2.0.1. *Let $prop$ be a set of variables. Then a formula ϕ is defined as follows:*

$$\phi ::= p \mid \perp \mid \neg\phi \mid \phi \vee \psi \mid \Box_i \phi$$

where $p \in Prop$ and \Box_i is a modal operator. Other connectives are expressed through \neg and \vee and dual modal operators \Diamond_i as $\Diamond_i \phi = \neg \Box_i \neg \phi$

Definition 2.0.2. *Let $M = (W, R_1, \dots, R_n, V)$ be a model and $w \in W$ a state in M . The notion of a formula being true at w is inductively defined as follows :*

$M, w \Vdash p$	<i>iff</i> $w \in V(p)$
$M, w \Vdash \perp$	<i>never</i>
$M, w \Vdash \neg\phi$	<i>iff not</i> $M, w \Vdash \phi$
$M, w \Vdash \phi \vee \psi$	<i>iff</i> $M, w \Vdash \phi \vee M, w \Vdash \psi$
$M, w \Vdash \Box_i \phi$	<i>iff</i> $\forall v \in W : w R_i v \rightarrow M, v \Vdash \phi$

Definition 2.0.3. *Let $F = (W, R_1, R_2, \dots)$ and $F' = (W', R'_1, R'_2, \dots)$ be two frames. A bounded morphism from F to F' is a function $f : W \rightarrow W'$ satisfying the following conditions:*

$$\text{If } (u, v) \in R_i \text{ then } (f(u), f(v)) \in R'_i$$

$$\text{If } (f(w), v') \in R'_i \text{ then } \exists v \in W \text{ s.t. } (w, v) \in R_i \text{ and } f(v) = v'$$

We say F' is a bounded morphic image of F , if there is a surjective bounded morphism from F to F' .

Proposition 2.0.4. *Let ϕ be a formula in $L_{\Box_1, \dots, \Box_n}$, $F = (W, R_1, \dots, R_n)$ and $F' = (W', R'_1, \dots, R'_n)$ be two frames and F to F' a surjective bounded morphism. Then the following holds :*

$$\text{If } F \Vdash \phi \text{ then } F' \Vdash \phi$$

Proof. This can be shown by structural induction on the length of the formula. □

Corollary 2.0.5. *If F' is a bounded morphic image of F , then we have $Log(F) \subseteq Log(F')$*

Proof. The proof is by structural induction. □

In this work, we deal with products of frames and multimodal logics of product of frames. The study of products of Kripke frames and their modal logics was first initiated by Segerberg [4] and Shehtman [5]. In the following, we will define necessary notions for such logics.

Definition 2.0.6. Let $F = (W, R_1)$ and $G = (W, R_2)$. Then we define the Kripke product on $W \times V$ as follows :

$$(w, v)R'_1(w', v') \text{ iff } wR_1w' \text{ and } v = v'$$

$$(w, v)R'_2(w', v') \text{ iff } w = w' \text{ and } vR_2v'$$

R'_1 is also called horizontal and R'_2 vertical relation.

Definition 2.0.7. A normal modal logic is a set of modal formulas containing all propositional tautologies, closed under Substitution ($\frac{\phi(p_i)}{\phi(\psi)}$), Modus Ponens ($\frac{\phi, \phi \rightarrow \psi}{\psi}$), Generalization rules ($\frac{\phi}{\Box_i \phi}$) and the following axioms

$$\Box_i(p \rightarrow q) \rightarrow (\Box_i p \rightarrow \Box_i q)$$

K_n denotes the minimal normal modal logic with n modalities and $K = K_1$. Let L be a logic and let Γ be a set of formulas. Then $L + \Gamma$ denotes the minimal logic containing L and Γ

Definition 2.0.8. Let L_1 and L_2 be two modal logic with one modality \Box . Then the fusion of these logics are defined as follows :

$$L_1 \otimes L_2 = K_2 + L_1(\Box \rightarrow \Box_1) L_2(\Box \rightarrow \Box_2)$$

The following logics may be important

$$D = K + \Box p \rightarrow \Diamond p$$

$$T = K + \Box p \rightarrow p$$

$$D4 = D + \Box p \rightarrow \Box \Box p$$

$$S4 = T + \Box p \rightarrow \Box \Box p$$

We now introduce important notation to prove the multimodal version of the Filtration Theorem.

Definition 2.0.9. A set Σ is closed under subformulas, if for all formulas ϕ and ϕ' the following holds :

1. if $\neg \phi \in \Sigma$ then $\phi \in \Sigma$
2. if $\phi \vee \phi' \in \Sigma$ then $\phi, \phi' \in \Sigma$
3. if $\Box_i \phi \in \Sigma$ then $\phi \in \Sigma$

Example : Suppose we have a multimodal logic with \Box, \Box_1 called $\mathcal{L}_{\Box, \Box_1}$ where $\phi = \Box p \rightarrow \Box_1 q$ and $\phi \in \Sigma$. Then $\Sigma = \{\phi, \neg \Box p, \Box_1 q, \Box p, p, q\}$ is closed under subformulas.

Definition 2.0.10. Let $M = (W, R_1, \dots, R_n, V)$ be a model and suppose Σ is a set of formulas. We define a relation \equiv on W as follows :

$$w \equiv v \text{ iff } \forall \phi \in \Sigma : M, w \Vdash \phi \Leftrightarrow M, v \Vdash \phi$$

It is well known that the \equiv -relation is an equivalence relation. We denote the equivalence class of a state $w \in W$ as $[w]_\Sigma = \{v \mid v \equiv w\}$. Furthermore W_Σ is the set of all equivalence classes, i.e $W_\Sigma = \{[w]_\Sigma \mid w \in W\}$.

In the following, we define what is known as a "filtration." The definition is similar to the one found in the Modal Logic book, but we consider a multimodal version.

Definition 2.0.11. Let $M = (W, R_1, \dots, R_n, V)$ be a model, Σ is closed under subformulas and W_Σ the set of equivalence classes induced by \equiv . A model $M_\Sigma^f = (W^f, R_1^f, \dots, R_n^f, V^f)$ is called filtration of M through Σ if the following holds :

1. $W^f = W_\Sigma$
2. If $(w, v) \in R_i$ then $([w], [v]) \in R_i^f$
3. If $([w], [v]) \in R_i^f$ then for any $\Box_i \phi \in \Sigma$: if $M, w \Vdash \Box_i \phi$ then $M, v \Vdash \phi$
4. $V^f = \{[w] \mid M, w \Vdash p\}$, for all propositional variables $p \in \Sigma$

Theorem 2.0.12. Consider $L_{\Box_1 \dots \Box_n}$. Let $M^f = (W_\Sigma, R_1^f, \dots, R_n^f, V)$ be a filtration of M through a subformula closed set Σ . Then for all formulas $\phi \in \Sigma$, and all nodes $w \in M$, we have

$$M, w \Vdash \phi \text{ iff } M^f, [w] \Vdash \phi$$

Proof. By induction on ϕ .

Case $\phi = p$: Left to right follows immediately from filtration definition. Conversely, suppose $M^f, [w] \Vdash p$. This means $[w] \in V^f(p)$. But this means $V(p)$ can not be empty. Pick any $v \in V(p)$. Obviously, $w \equiv v$ and $M, v \Vdash p$. Hence, $M, w \Vdash p$.

Case $\phi = \neg\psi$: Suppose ψ holds. Then we have : $M, w \Vdash \phi$ iff $M, w \not\Vdash \psi$. Applying induction hypothesis, we get : $M^f, [w] \not\Vdash \psi$. But then, we have $M^f, [w] \Vdash \phi$. Right to left is the same.

Case $\phi = \phi_1 \vee \phi_2$: Suppose ϕ_1, ϕ_2 holds. Let $M, w \Vdash \phi$. That means $M, w \Vdash \phi_1$ or $M, w \Vdash \phi_2$. Applying induction hypothesis, we get $M^f, [w] \Vdash \phi_1$ or $M^f, [w] \Vdash \phi_2$. But then, $M^f, [w] \Vdash \phi_1 \vee \phi_2 = \phi$. Right to left is similar.

Case $\phi = \Box_i \psi$: Left to right. Suppose ψ holds and $M, w \Vdash \Box_i \psi$. We need to show $M^f, [w] \Vdash \Box_i \psi$, this means $\forall [v] \in W_\Sigma : [w] R_i [v] \rightarrow M^f, [v] \Vdash \Box_i \psi$. Pick any $[v] \in W_\Sigma$ s.t $[w] R_i [v]$. By condition 3, w.r.t to the modal operator, we have $M, v \Vdash \psi$. By induction hypothesis, we get $M^f, [v] \Vdash \psi$. Because $[v]$ was arbitrary it follows that $M^f, [w] \Vdash \Box_i \psi$.

Right to left. Suppose ψ holds and $M^f, [w] \Vdash \Box_i \psi$. Pick $v \in W$ s.t $w R_i v$. By condition 2, w.r.t to the modal operator, we have $[w] R_i^f [v]$. So, $M^f, [v] \Vdash \psi$. By induction hypothesis, we get $M, v \Vdash \psi$. Because v was arbitrary, we have $M, w \Vdash \Box_i \psi$. \square

After showing that filtration preserves truth, we now define a special relation known as the "smallest filtration."

Definition 2.0.13. Let $M = (W, R_1, \dots, R_n, V)$ be a model, Σ is closed under subformulas and W_Σ the set of equivalence classes. We define :

$$R^s = \{[w], [v] \mid \exists w' \in [w], \exists v' \in [v] : w' R_i v'\}$$

Lemma 2.0.14. Let $M = (W, R_1, \dots, R_n, V)$ be a model, Σ is closed under subformulas and W_Σ the set of equivalence classes induced by \equiv and V^f the standard valuation on W_Σ . Then $(W_\Sigma, R_1^s, \dots, R_n^s, V^f)$ is a filtration of M through Σ .

Proof. It suffices to show R_i^s fullfills the condition 2 and 3 w.r.t to the corresponding modal operator \Box_i . But R_i^s already satisfies condition 2. Let's check the other condition. Let $\Box_i \phi \in \Sigma$, $[w] R_i^s [v]$ and $M, w \Vdash \Box_i \phi$. Because of $[w] R_i^s [v]$ we pick a $w' \in [w]$ and $v' \in [v]$. By definition, we have $w' R_i v'$. Because $w' \equiv w$, we get $M, w' \Vdash \Box_i \phi$. Hence, $M, v' \Vdash \phi$ and by $v' \equiv v$, we get $M, v \Vdash \phi$. \square

Proposition 2.0.15. Let Σ be a finite subformula closed set of $L_{\Box_1, \dots, \Box_n}$. For any model M , if M^f is a filtration through Σ , then M^f contains at most 2^n nodes (where n denotes the size of Σ).

Proof. The states of M^f are the equivalence classes in W_Σ . Let $g : W_\Sigma \rightarrow P(\Sigma)$ defined by $g([w]) = \{\phi \in \Sigma \mid M, w \Vdash \phi\}$. g is well defined. Pick any u and v s.t $u \equiv v$. But then by definition of \equiv , they fullfill the same subformulas. This means $g([v]) = g([u])$. g is also injective. Pick any $[u], [v] \in W_\Sigma$ s.t $g([u]) = g([v])$. We show $[u] \subseteq [v]$. The other inclusion is similar. By assumption we have $u \equiv v$. Pick any $u' \in [u]$. Then we have $u' \equiv u \equiv v$. Hence, $u' \in [v]$. At the end, this means M^f contains at most 2^n nodes. \square

Theorem 2.0.16. Let ϕ be a formula of $L_{\Box_1, \dots, \Box_n}$. If ϕ is satisfiable, then it is satisfiable on a finite model containing at most 2^v nodes, where v is the number of subformulas of ϕ .

Proof. Assume that ϕ is satisfiable on a model on M . Take any filtration of M through the set of subformulas of ϕ . By Filtration Theorem, we get that ϕ is satisfied in the filtration model M^f . Furthermore, it is bounded by 2^v . \square

Now we define special kind of formulas called Sahlqvist formulas.

Definition 2.0.17. A modal formula ϕ is positive if all variables occurs in the scope of an even number of negations. In the other hand, a formula is negative, if all variables occurs in the scope of an odd number of negations. A boxed atom is a modal formula of the form $\Box^n p$ for some $n \in \mathbb{N}$, where p is a propositional variable and $\Box^n p$ is defined as follows : $\Box^0 p = p$, $\Box^1 p = \Box p$, $\Box^{n+1} p = \Box(\Box^n p)$.

Sahlqvist formulas can be build by the following inductive definition :

$$\phi ::= \neg \Box^d p \mid \pi \mid \phi_1 \wedge \phi_2 \mid \phi_1 \vee \phi_2 \mid \Box \phi$$

where p is an atom, $d \geq 0$ and π is a positive formula.

Examples for Sahlqvist formulas:

$$\begin{aligned} & \neg \Box \Box p \vee \Box p \\ & \neg \Box p \vee p \\ & \Box \neg \Box \Box \Box \Box p \vee \Box \Diamond \Box \Diamond p \\ & \Box \Box \Box \Box (\Box \neg \Box p \vee p) \wedge \neg \Box \Box p \vee \Box p \end{aligned}$$

Non Sahlqvist Formulas :

$$\begin{aligned} & \neg \Box \Diamond p \vee \Diamond \Box p \\ & \neg \Diamond \Box p \vee \Box \neg p \end{aligned}$$

We can extend this definition for multimodal logics. We say a boxed atom can be $\Box_i^n p$.

Sahlqvist formulas possess important properties, which are guaranteed by the Sahlqvist Theorem. It says that, when given a normal modal logic K and a set of Sahlqvist formulas, the resulting logic is complete w.r.t to the class of frames, which satisfies the corresponding first-order formula of the Sahlqvist formulas. This also holds for multimodal logic. We will not prove it here, but we will use it later in this work. The main idea is to use Sahlqvist to show that our logic is sound and complete w.r.t. a certain class of frames. Using the smallest filtration and Filtration Theorem, we can guarantee the finite model property while preserving the relevant frame properties.

3 Product of Topological spaces and Neighborhood frames

Topology is a fundamental branch in mathematics that explores the properties of space that remain invariant under continuous transformations. Unlike geometry, where we rely on precise measurements, topology focuses on the more flexible notions of closeness, boundary and countinuity without relying on specific distances.

One particularly rich and fruitful area of study is the interaction between modal logic and topology. Topological spaces provide an intuitive semantic framework for modal logic by interpreting modal operators in terms of spatial notions like interior, closure, and neighbourhoods. It is also well known, that the modal logic of any topological space is $S4$.

Neighborhood frames generalizes the idea of topology by loosen the conditions that define open sets in topological space. Instead of relying on a global systems of open sets, there exists a neighborhood function, which assigns to each point a collection of sets — its neighbourhoods. This provides us a more flexible framework to work with.

Oftentimes, it is in theoretical and practical interest to combine topological spaces or neighborhood frames. These product constructions become particularly relevant in the context of multimodal logic, where we aim to reason with multiple modal operators simultaneously. Each modality can be interpreted in its own topological or neighbourhood dimension, allowing us to model systems with distinct but interacting notions of necessity — such as time and space or knowledge and belief.

In this chapter, we introduce the notation for product of topological spaces and the semantics for multimodal logic in topology. We will also do it for neighborhood frames.

3.1 Topological semantics and product of topological spaces

Definition 3.1.1. *A topological space is a pair (X, τ) where τ is a collection of subsets of X (elements of τ are also called open sets) such that :*

1. *the empty set \emptyset and X are open*
2. *the union of an arbitrary collection of open sets is open*
3. *the intersection of finite collection of open sets is open*

A topological model is a structure $M = (X, \tau, v)$ where (X, τ) is a topological space and v is a valuation assigning subsets of X to propositional variables.

Remark 3.1.2. *There is an equivalent definition for open sets. Let (X, τ) be a topological space and U a set. U is open iff $\forall x \in U \exists V \subseteq U : V$ is open and $x \in V$. This is true because, the union of open sets is an open set.*

Definition 3.1.3. *Let (X, τ) be a topological space. Then the space is called Alexandroff, if arbitrary intersection of open sets are open. An equivalent definition is that every point has a smallest open set. That means for each $x \in X$ we have $U_x = \bigcap \{U \subseteq X \mid U \text{ is open and } x \in U\} \in \tau$.*

Example 3.1.4. Let $X = \{1, 2, 3\}$. We consider different topologies on X :

- If $\tau = \{\emptyset, X\}$, then (X, τ) is called the trivial topological space.
- If $\tau = \mathcal{P}(X)$, then (X, τ) is called the discrete topological space.
- If $\tau = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, X\}$, then (X, τ) is topological space.

These spaces are Alexandroff. In the following we will see a non-Alexandroff space.

Example 3.1.5. Let $X = \mathbb{R}$ with the standard topology. That means the basis for open sets is $\mathcal{B} = \{(a, b) \subseteq \mathbb{R} \mid a < b\}$ with $a, b \in \mathbb{R}$. Now, we consider a family of open sets $U_n = \{-\frac{1}{n}, \frac{1}{n}\}$ for $n \in \mathbb{N}$. But, it is the case that $\bigcap_{n=1}^{\infty} U_n = \{0\}$ is not open.

Definition 3.1.6. Let $M = (X, \tau, v)$ be a topological model and $x \in X$. The satisfaction of a formula at the point x in M is defined inductively as follows:

$$\begin{aligned}
M, x \models p & \quad \text{iff } x \in v(p) \\
M, x \models \perp & \quad \text{never} \\
M, x \models \neg\phi & \quad \text{iff } M, x \not\models \phi \\
M, x \models \phi \vee \psi & \quad \text{iff } M, x \models \phi \text{ or } M, x \models \psi \\
M, x \models \Box\phi & \quad \text{iff there exists } U \in \tau \text{ such that } x \in U \text{ and } \forall u \in U, M, u \models \phi
\end{aligned}$$

Definition 3.1.7. Let $\mathcal{X} = (X, \chi)$ and $\mathcal{Y} = (Y, v)$ be topological spaces and $N \subseteq X \times Y$.

We define

Horizontally open: N is horizontally open iff

$$\forall (x, y) \in N \exists U \in \chi \text{ such that } x \in U \text{ and } U \times \{y\} \subseteq N.$$

Vertically open: N is vertically open iff

$$\forall (x, y) \in N \exists V \in v \text{ such that } y \in V \text{ and } \{x\} \times V \subseteq N.$$




Figure 4: Each rectangle represents N and the redlines are subsets of N

If N is H -open and V -open, then we call it HV -open.

We denote τ_1 is the set of all H -open subsets of $X \times Y$ and τ_2 is the set of all V -open subsets of $X \times Y$. We say τ is the standard product of subsets $X \times Y$ s.t. $X \in \chi$ and $Y \in v$.

Example 3.1.8. Let $\mathcal{X} = (\mathbb{R}, \chi)$ and $Y = (\mathbb{R}, v)$ with standard topology, that means the basis $\mathcal{B} = \{(a, b) \subseteq \mathbb{R} \mid a < b\}$. Now consider $\mathcal{X} \times \mathcal{Y}$. The horizontal topology is generated by $\mathcal{B}_1 = \{U \times \{y\} \mid U \in \chi \text{ and } y \in \mathbb{R}\}$ and vertical topology by $\mathcal{B}_2 = \{\{x\} \times V \mid x \in \mathbb{R} \text{ and } V \in v\}$.

Definition 3.1.9. Let $\mathcal{X} = (X, \chi)$ and $\mathcal{Y} = (Y, v)$ be topological spaces and $(x, y) \in X \times Y$. The truth in $M = (X \times Y, \tau_1, \tau_2, \tau, v)$ is similar as in Definition 2.1.6. For \Box , \Box_1 and \Box_2 we interpret this as follows :

$$\begin{aligned} M, (x, y) \models \Box_1 \phi & \text{ iff } \exists U \in \tau_1 : (x, y) \in U \text{ and } \forall (x', y') \in U : (x', y') \models \phi \\ M, (x, y) \models \Box_2 \phi & \text{ iff } \exists V \in \tau_2 : (x, y) \in V \text{ and } \forall (x', y') \in V : (x', y') \models \phi \\ M, (x, y) \models \Box \phi & \text{ iff } \exists U \in \chi \exists V \in v : (x, y) \in U \times V \text{ and } \forall (x', y') \in U \times V : (x', y') \models \phi \end{aligned}$$

Definition 3.1.10. Let $\mathcal{X} = (X, \chi)$ and $\mathcal{Y} = (Y, v)$ be two topological spaces. Then the full product of these spaces is defined as follows :

$$\mathcal{X} \times_t^+ \mathcal{Y} = (X \times Y, \tau_1, \tau_2, \tau)$$

with horizontal, vertical and standard topologies.

Definition 3.1.11. For two unimodal logics L_1 and L_2 we define the full t -product of them as follows :

$$L_1 \times_t^+ L_2 = \text{Log}(\{\mathcal{X} \times_t^+ \mathcal{Y} \mid \mathcal{X} \models L_1 \text{ and } \mathcal{Y} \models L_2\})$$

where \mathcal{X}, \mathcal{Y} are topological spaces.

Definition 3.1.12. A topo-bisimulation between two topological models $M = (X, \tau, v)$ and $M' = (X', \tau', v')$ is a nonempty relation $\vartriangleleftharpoons \subseteq X \times X'$ s.t. if $x \vartriangleleftharpoons x'$ then

1. Base: $x \in V(p)$ iff $x' \in v'(p)$ for any propositional variable p
2. Forth: $x \in U \in \tau$ implies that there exists $U' \in \tau'$ s.t. $x' \in U'$ and for every $y' \in U'$ there is $y \in U$ with $y \vartriangleleftharpoons y'$
3. Back: $x' \in U' \in \tau'$ implies that there exists $U \in \tau$ s.t. $x \in U$ and for every $y \in U$ there is $y' \in U'$ with $y \vartriangleleftharpoons y'$

Proposition 3.1.13. Let $M = (X, \tau, v)$ and $M' = (X', \tau', v')$. Assume $w \in M'$ and $w' \in M'$ are topo-bisimilar points. Then for each formula ϕ we have

$$M, w \models \phi \text{ iff } M', w' \models \phi$$

Proof. The proof is by structural induction. □

Definition 3.1.14. Let X and Y be topological spaces and $f : X \rightarrow Y$ a function. We call f continuous if for each open set $U \subseteq Y$ the set $f^{-1}(U)$ is open in X . We say f is open if for each open set $V \subseteq X$ the set $f[V]$ is open in Y .

3.2 Product and semantics of neighborhood frames

Neighborhood frames are more flexible than topological space. Its logic is not $S4$ in general. It is well known, that neighborhood frames fully captures the semantics of Kripke. Moreover, neighborhood frames are expressive enough to characterize non-normal modal logics, which makes them a powerful tool.

Definition 3.2.1. Let X be a non-empty set. A function $\tau : X \rightarrow 2^{2^X}$ is called a neighbourhood function. A pair $F = (X, \tau)$ is called a neighbourhood frame (or n -frame). A model based on F is a tuple (X, τ, v) , where v assigns a subset of X to a variable.

Definition 3.2.2. Let $M = (X, \tau, v)$ be a neighbourhood model and $x \in X$. The truth of a formula is defined inductively as follows :

$M, x \Vdash p$	iff $x \in V(p)$
$M, x \Vdash \perp$	never
$M, x \Vdash \neg\phi$	iff $M, x \not\Vdash \phi$
$M, x \Vdash \phi \vee \psi$	iff $M, x \Vdash \phi \vee M, x \Vdash \psi$
$M, x \Vdash \Box\phi$	iff $\exists V \in N(x) \forall y \in V : M, y \Vdash \phi$

A formula is valid in a n -model M if it is valid at all points of M ($M \models \phi$). Formula is valid in a n -frame F if it is valid in all models based on F (notation $F \models \phi$). We write $F \models L$ if for any $\phi \in L$, $F \models \phi$. Logic of a class of n -frames \mathcal{C} as $\text{Log}(\mathcal{C}) = \{\phi \mid F \models \phi \text{ for all } F \in \mathcal{C}\}$. We define $nV(L) = \{F \mid F \text{ is an } n\text{-frame and } F \models \phi\}$.

Definition 3.2.3. Let X be a non-empty set and τ neighborhood function. We call τ is a filter if for each $x \in X$ the collection $\tau(x)$ satisfies the following conditions :

1. $\emptyset \notin \tau(x)$
2. If $U \in \tau(x)$ and $U \subseteq V$ then $V \in \tau(x)$ (upward closed)
3. If $U, V \in \tau(x)$, then $U \cap V \in \tau(x)$

Definition 3.2.4. Let $X = (X, \tau_1, \dots)$ and $Y = (Y, \sigma_1, \dots)$ be n -frames. Then the function $f : X \rightarrow Y$ is called bounded morphism if

1. f is surjective
2. $\forall x \in X \forall U \in \tau_i(x) : f(U) \in \sigma_i(f(x))$
3. $\forall x \in X \forall V \in \sigma_i(f(x)) \exists U \in \tau_i(x) : f(U) \subseteq V$

Corollary 3.2.5. Let $X = (X, \tau_1, \dots)$ and $Y = (Y, \sigma_1, \dots)$ be n -frames and $f : X \rightarrow Y$ a bounded morphism. Then

$$\text{Log}(X) \subseteq \text{Log}(Y)$$

Proof. The proof is by structural induction. □

In topology, we defined horizontal and vertical topologies for products of topological spaces. In the following, we will define a similar construction for neighborhood frames.

Definition 3.2.6. Let $\mathcal{X} = (X, \tau_1)$ and $\mathcal{Y} = (Y, \tau_2)$ be two n -frames. Then the product of these two frames is a n -2-frame and is defined as follows :

$$\begin{aligned}\mathcal{X} \times_n \mathcal{Y} &= (X \times Y, \tau'_1, \tau'_2) \\ \tau'_1(x, y) &= \{U \subseteq X \times Y \mid \exists V \in \tau_1(x) : V \times \{y\} \subseteq U\} \\ \tau'_2(x, y) &= \{U \subseteq X \times Y \mid \exists V \in \tau_2(y) : \{x\} \times V \subseteq U\}\end{aligned}$$

Additionally, we say the full product of n -frames $\mathcal{X} \times_n^+ \mathcal{Y}$ is :

$$\begin{aligned}\mathcal{X} \times_n^+ \mathcal{Y} &= (X \times Y, \tau'_1, \tau'_2, \tau) \text{ where} \\ \tau(x, y) &= \{U \subseteq X \times Y \mid \exists W \in \tau_1(x) \exists V \in \tau_2(y) : W \times V \subseteq U\}\end{aligned}$$

Definition 3.2.7. For two unimodal logics L_1 and L_2 we define the n -product of them as follows :

$$L_1 \times_n L_2 = \text{Log}(\{\mathcal{X} \times \mathcal{Y} \mid \mathcal{X} \in nV(L_1) \text{ and } \mathcal{Y} \in nV(L_2)\})$$

In similar way, we can define $L_1 \times_n^+ L_2$ with three topologies.

Proposition 3.2.8 ([8]). For two unimodal logics L_1 and L_2 it holds :

$$L_1 \otimes L_2 \subseteq L_1 \times_n L_2$$

Definition 3.2.9. Let $\mathcal{L}_{\Box, \Box_1, \Box_2}$ be a modal language with three modal operators \Box, \Box_1 and \Box_2 . We define the T -neighborhood product logic TNL as the least set of formulas in $\mathcal{L}_{\Box, \Box_1, \Box_2}$ containing all axioms $T \otimes T \otimes T + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$, and closed under modus ponens, substitution and $\Box-$, \Box_1- and \Box_2- neccessitation.

After introducing the necessary notation, we can now formally state our main research question as follows: Does the following equivalence hold?

$$TNL = T \times_n^+ T$$

The main idea is to introduce an infinite depth and branching tree called $T_{\omega, \omega, \omega[rn]}$ and show $\text{Log}(T_{\omega, \omega, \omega[rn]}) = TNL$. Then we use the idea in [2] to use the pseudo-infinite paths construction to build an n -frame called $N_\omega(T_{\omega[rn]})$ and construct a p -morphism-like map from $N_\omega(T_{\omega[rn]}) \times_n^+ N_\omega(T_{\omega[rn]})$ to $T_{\omega, \omega, \omega[rn]}$. We will introduce the notation later.

4 Completeness result for TNL

We introduce a frame called $T_{\omega, \omega, \omega[rn]}$. We will show TNL is sound and complete w.r.t $T_{\omega, \omega, \omega[rn]}$. The idea is to pick a class of frame \mathcal{C} s.t $Log(\mathcal{C}) = Log(T_{\omega, \omega, \omega[rn]})$ and then show the class has FMP. In the end, we will use an unravelling technique to show completeness.

Definition 4.0.1. *We say for a modal logic Λ has the finite model property (FMP) if for every formula ϕ that is not provable in Λ , is falsifiable in a finite model.*

Proposition 4.0.2. *The logic $T \otimes T \otimes T + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$ has FMP.*

Proof. The idea is to pick a class of frame \mathcal{F} , where TNL is sound and complete with respect to and then show by filtration the FMP.

Let $\mathcal{C} = \{F \mid F \models TNL\}$. Obviously, TNL is sound w.r.t \mathcal{C} . Completeness can be shown by using the Sahlqvist Theorem. We remember $T = K + \Box_i p \rightarrow p$ for $i \in \{1, 2, \epsilon\}$. These axioms and $\Box p \rightarrow \Box_1 p \wedge \Box_2 p$ are Sahlqvist formulas. By Sahlqvist, we have TNL is complete w.r.t $\{F \mid F \models \forall x R_i(x, x) \text{ and } F \models \forall x \forall y (R_1(x, y) \vee R_2(x, y)) \rightarrow R(x, y)\}$.

Now assume a formula ϕ is not derivable from TNL . By completeness we get ϕ is falsifiable in a model M and a world w . Hence, $M, w \models \neg\phi$. Now we build the set $\text{subf}(\neg\phi)$ which denotes the closed subformulas set of $\neg\phi$. Let $M^s = (W_\Sigma, R^s, R_1^s, R_2^s, V^s)$ be the smallest filtration of M through $\text{subf}(\neg\phi)$ and V^s is the standard valuation on W_Σ . By Filtration theorem, they preserve truth. It remains to show $F^s = (W_\Sigma, R^s, R_1^s, R_2^s) \in \mathcal{C}$. For that, every point must have an edge to itself in every relation and it must hold that $R_1^s, R_2^s \subseteq R^s$. For the first one, we show this for R^s because the rest is similar. Pick $w' \in [w] \in W_\Sigma$. Because M is based on a frame $F \in \mathcal{C}$, we have $w' R w'$. By the definition of smallest filtration, we have that $[w'] R^s [w']$. For the second one, we show only for R_1^s because it is the same for R_2^s . Pick $[w], [v]$ s.t $[w] R_1^s [v]$. By definition of smallest filtration there are points $w' \in [w]$ and $v' \in [v]$ s.t $w' R_1 v'$. Furthermore, it holds $R_1 \subseteq R$, so $w' R v'$. By smallest filtration, we get $[w'] R^s [v']$. Because $w' \equiv w$ and $v' \equiv v$, we have $[w'] = [w]$ and $[v'] = [v]$. It follows $[w] R^s [v]$. □

Definition 4.0.3. *Let $T_{\omega[rn]}$ ($r = \text{reflexive}$, $n = \text{non-transitive}$) denote the infinite branching and infinite depth tree, which is reflexive and non-transitive. Formally the tree can be defined as : $T_{\omega[rn]} = (W, R)$ where $W = \mathbb{N}^*$,*

$$s R t \text{ iff } \exists u \in \mathbb{N} \cup \{\epsilon\} : s \cdot u = t$$

(\cdot is the concatenation operator)

The $T_{\omega, \omega, \omega[rn]}$ tree has three relations with infinite branching, infinite depth and we have $R_1, R_2 \subseteq R$. Before characterizing it, we say \mathbb{N}_1^* is the set of finite number combinations which has subscript "1" to denote these numbers relate to R_1 . (examples : $2_1, 4231123_1, 32_1\epsilon 45_1\epsilon 9_1 = 32459_1$). The similar holds for \mathbb{N}_2^* .

Now let $T_{\omega, \omega, \omega[rn]} = (W, R, R_1, R_2)$ where $W = (\mathbb{N} \cup \mathbb{N}_1 \cup \mathbb{N}_2)^*$,

$$sRt \text{ iff } \exists u \in \mathbb{N} \cup \mathbb{N}_1 \cup \mathbb{N}_2 \cup \{\epsilon\} : s \cdot u = t$$

$$sR_1t \text{ iff } \exists u \in \mathbb{N}_1 \cup \{\epsilon\} : s \cdot u = t$$

$$sR_2t \text{ iff } \exists u \in \mathbb{N}_2 \cup \{\epsilon\} : s \cdot u = t$$

where $s, t \in W$. Again, the "." operator acts here as a concatenation operator.




Figure 5: This is $T_{\omega, \omega, \omega[rn]}$. The dotted lines on each node are from R_1 and the dashed lines represents R_2 . The whole tree is R .

Proposition 4.0.4. T is sound and complete w.r.t $T_{\omega[rn]}$.

Proof. Sound is clear. For completeness, we use the well known fact that T has FMP. This means, $T = \text{Log}\{F \mid F \models T\}$ where F is a finite frame. We can pick such a finite frame F and it suffices to find a surjective bounded morphism f from T_w to F . This would imply $\text{Log}(T_{w[rn]}) \subseteq T$.

Now, let $F = (W', R')$ be such a finite rooted frame with root w . We define inductively an assignment of nodes of F to the nodes of T_w . For the base case, we assign w to the root of T_w . The induction step looks like the following : Assume a point $x \in T_w$ has been assigned to a point $u \in F$ but the successors of x has no assignment. Let s_1, s_2, \dots, s_k be successors of u (k denotes amount of successors and $k \geq 1$ because reflexivity guarantees us at least one successor). For $n \geq 1, n \in \mathbb{N}, i \in \{1, \dots, k\}$ we assign s_i to the $(n * i)$ th-successor of x . This means we are assigning the successors alternately.

Now we check for f the conditions of bounded morphism. First condition : Let $x, y \in T_w$ s.t xRy and $f(x) = s$. But then, y will be assigned to a successor point of s . Hence, $f(x)R'f(y)$. Second condition : Suppose $f(x)R't$ and $f(x) = s$. Since t is a successor of s and $f(x) = s$, then a successor of x , say y , gets the assignment t .

□

Proposition 4.0.5. *TNL is sound and complete w.r.t $T_{\omega,\omega,\omega[rn]}$.*

Proof. For soundness, we have that $T_{\omega,\omega,\omega[rn]} \models \Box p \rightarrow \Box_1 p \wedge \Box_2 p$, because by definition we have $R_1, R_2 \subseteq R$. The rest is clear. For completeness, we use the fact *TNL* has FMP. Let $F = (W', R', R'_1, R'_2)$ be a finite rooted frame with root w and $F \models TNL$. We define inductively an assignment similar to Proposition 3.0.4. We assign w to the root of $T_{\omega,\omega,\omega[rn]}$. For induction step we start by only assigning points from R_1 to R'_1 and R_2 to R'_2 . After that, the remaining points will be assigned to the points in R' . The procedure works similar as described previously.

We will only check the conditions for R and R' . For R_1, R'_1 and R_2, R'_2 , we can argue as before. We pick any xRy with $f(x) = s$. If we also have xR_1y or xR_2y , then it follows $f(x)R'f(y)$ because $R_1, R_2 \subseteq R$ and $R'_1, R'_2 \subseteq R'$. Else, the successor of s was assigned to y , so $f(x)R'f(y)$.

Let $f(x)R't$ and $f(x) = s$. If we have $f(x)R'_1t$ or $f(x)R'_2t$, then the second condition follows because $R_1, R_2 \subseteq R$. Else, we have that t is a successor of $f(x) = s$, and a successor of x was assigned to t . \square

5 Logic of product of topological spaces with three topologies

In this section, we will reprove the logic of the product of two topological spaces with horizontal, vertical and classic product topology is $S4 \otimes S4 \otimes S4 + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$ [1]. First we will show a version for one modality because it is easier to understand and the proof idea is based on this version.

Definition 5.0.1. *Let T_2 be the infinite binary tree with reflexive and transitive descendant relation.*

Formally it is defined as follows : $T_2 = (W, R)$ where $W = \{0, 1\}^$,*

$$sRt \text{ iff } \exists u \in W^* : s \cdot u = t$$

The $T_{6,2,2}$ tree (similar to $T_{\omega, \omega, \omega[rn]}$) is the infinite six branching tree, where all nodes of $T_{6,2,2}$ is R -related, the first two R_1 -related and the last two R_2 -related. Formally we can define this tree as follows : $T_{6,2,2} = (W, R, R_1, R_2)$, where $W = \{0, 1, 2, 3, 4, 5\}^$,*

$$sRt \text{ iff } \exists u \in \{0, 1, 2, 3, 4, 5\}^* : s \cdot t = u$$

$$sR_1t \text{ iff } \exists u \in \{0, 1\}^* : s \cdot t = u$$

$$sR_2t \text{ iff } \exists u \in \{5, 6\}^* : s \cdot t = u$$

where $s, t \in W$ and \cdot is the concatenation-operator.

Theorem 5.0.2. *$S4$ is complete with respect to T_2 .*

Proof. The idea is to use the fact $S4$ has finite model property. By that we can pick a finite rooted $S4$ -frame and then show that this frame is the bounded morphic image of T_2 . For details see [12]. \square

Theorem 5.0.3. *(Cantor) Every countable dense linear ordering without endpoints is isomorphic to \mathbb{Q} .*

Proof. For a proof see e.g [13, Page 217, Theorem 2]. \square

Our strategy is as follows. We use completeness of $S4$ w.r.t. T_2 , view T_2 as an Alexandroff space, define a dense subset X of \mathbb{Q} without endpoints, and then establish a topo-bisimulation between X and T_2 . This will allow us to transfer counterexamples from T_2 to X , which by Cantor's theorem is order-isomorphic and hence homeomorphic to \mathbb{Q} .

Now let us define X as $X = \bigcup_{n \in \mathbb{N}} X_n$ where $X_0 = \{0\}$ and

$$X_{n+1} = X_n \cup \{x - \frac{1}{3^n}, x + \frac{1}{3^n} \mid x \in X_n\}$$

Lemma 5.0.4. For $n > 0$ and $x, y \in X_n, x \neq y$ we have : $|x - y| \geq \frac{1}{3^{n-1}}$

Proof. The proof is by induction on n . For the base case, if $n = 1$, then $X = 0, 1, -1$. For induction step, assume $u, v \in X_{n-1}$ with $u \neq v$. By induction hypothesis, it holds $|u - v| \geq \frac{1}{3^{n-2}}$. Suppose we pick $x = u + \frac{1}{3^{n-1}}$ and $y = v - \frac{1}{3^{n-1}}$. Hence, $x, y \in X_n$. But then, $|x - y| = |(u + \frac{1}{3^{n-1}}) - (v - \frac{1}{3^{n-1}})| = |u - v + \frac{2}{3^{n-1}}| \geq \frac{1}{3^{n-1}}$. If $u - v$ is positive, then the inequality follows immediately. Now assume $u - v$ is negative. By induction hypothesis we get $u - v \leq -\frac{1}{3^{n-2}}$. So the worst case is, when $u - v$ gets $-\frac{1}{3^{n-2}}$. But then, $|\frac{1}{3^{n-2}} - \frac{2}{3^{n-1}}| = \frac{1}{3^{n-1}}$. \square



Figure 6: This is an excerpt of X . Some points are visualized as brackets to distinguish them from nearby points at the same level.

In Figure 3, each colour represents a different level. The points in red are from X_1 , blue points from X_2 , green points from a part of X_3 and orange from a part of X_4 . We can observe that, the distance between a point and its source is getting exponentially smaller.

It follows from Lemma 4.0.4 that $(X, <)$ is a countable, dense, and linear ordered without endpoints. By Cantor's Theorem, X is homeomorphic to \mathbb{Q} . It also follows that for each $x \in X$ with $x \neq 0$ there exists an index n_x with $x \in X_{n_x}$ but $x \notin X_{n_x-1}$. But also, we can find a unique $y \in X_{n_x-1}$ with $x = y - \frac{1}{3^{n_x-1}}$ or $x = y + \frac{1}{3^{n_x-1}}$. Furthermore, the basis for the open intervals for any $x \in X$ will be $(x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})$ where n_x is the index where x gets produced.

Now we define f from X onto T_2 by recursion. If $x = 0$, then we let $f(0)$ be the root r of T_2 . If $x \neq 0$, then $x \in X_{n_x} - X_{n_x-1}$ and we let

$$f(x) = \begin{cases} \text{the left successor of } f(y) & \text{if } x = y - \frac{1}{3^{n_x-1}} \\ \text{the right successor of } f(y) & \text{if } x = y + \frac{1}{3^{n_x-1}} \end{cases}$$

Proposition 5.0.5. We claim, f is open and continuous.

Proof. We define the basis for the Alexandroff topology on T_2 as $\{B_t\}_{t \in T_2}$ where $B_t = \{s \in T_2 \mid tRs\}$. To show f is open, we pick an X -interval $(x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})$ and show $f((x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})) = B_{f(x)}$.

\subseteq : If we pick $x \in ((x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}}))$, then obviously $f(x)Rf(x)$. Assume we pick a point $y \in (x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})$ but $y \neq x$. Then we have $n_y > n_x$ and hence $f(x)Rf(y)$.

\supseteq : Assume $f(x)Rt$. We will show by induction on the length k between $f(x)$ and t s.t. there exists $y \in (x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})$ s.t. $f(y) = t$. Base case : If $k = 0$, then we have $t = f(x)$ and we can pick $y = x$. For induction step, assume it holds for $f(x)Rt'$ with length n , $f(y) = t'$ and $t'Rt$. For y , there must exist an index $n_y > n_x$ s.t. $y \in X_{n_y} - X_{n_y-1}$. Now

let $y' = y - \frac{1}{3^{n_y}}$ or $y' = y + \frac{1}{3^{n_y}}$. It holds that, $n'_y > n_y > n_x$, so $y' \in (x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})$. By definition of f , $f(y')$ is either left or right successor of $f(y)$ and by $f(y) = t'Rt$, we get $f(y') = t$. It follows f is open.

To show f is continuous, it suffices to show that for each $t \in T_2$, the f -inverse image of B_t is open. Let $x \in f^{-1}(B_t)$. Then $tRf(x)$. As argued before, we can find an open interval in X s.t. $f(x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}}) = B_{f(x)}$. By $tRf(x)$, we get $B_{f(x)} \subseteq B_t$. Hence, we found an open interval $I = (x - \frac{1}{3^{n_x}}, x + \frac{1}{3^{n_x}})$, s.t. $I = f^{-1}(B_{f(x)}) \subseteq f^{-1}(B_t)$. This holds because we can pick $y \in I$. By applying f , we get $f(y) \in B_{f(x)} \subseteq B_t$. But then $y \in f^{-1}(B_t)$. By that fact, we can follow with Remark 2.1.2, that $f^{-1}(B_t)$ is open and hence f is continuous. \square

To complete the proof, if $S4 \not\vdash \phi$, then by Theorem 4.0.2, there is a valuation v on T_2 s.t. $(T_2, v), r \not\models \phi$. We define a valuation ξ on X by $\xi(p) = f^{-1}(v(p))$. Since f is continuous and open, $f(0) = r$ and by the choice of the valuation, we have that 0 and r are topo-bisimilar. By Proposition 2.1.11 it follows $(X, \xi), 0 \not\models \phi$. Because X is homeomorphic to \mathbb{Q} , we obtain ϕ refutable on \mathbb{Q} .

Definition 5.0.6. Let $\mathcal{L}_{\Box, \Box_1, \Box_2}$ be a modal language with three modal operators \Box, \Box_1 and \Box_2 . We define the topological product logic TPL as the least set of formulas in $\mathcal{L}_{\Box, \Box_1, \Box_2}$ containing all axioms $S4 \otimes S4 \otimes S4 + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$, and closed under modus ponens, substitution and \Box -, \Box_1 - and \Box_2 -necessitation.

Theorem 5.0.7. TPL is complete w.r.t. $T_{6,2,2}$.

Proof. A generalization of Theorem 4.0.2 can be seen on [6]. \square

Now we are ready to show TPL is complete w.r.t. $\mathbb{Q} \times \mathbb{Q}$ with three topologies. The idea is similar as previous shown. We view $T_{6,2,2}$ as equipped with three Alexandroff topologies defined from R, R_1 and R_2 . It suffices to show that there exists a 3-topo-bisimulation between $T_{6,2,2}$ and $X \times X = X'$ where each X is as defined in Theorem 4.0.3.

Theorem 5.0.8 ([1]). TPL is complete w.r.t. $\mathbb{Q} \times \mathbb{Q}$.

Proof. We define h from $X \times X$ onto $T_{6,2,2}$ by recursion following the inductive definition of X . If $(x, y) = (0, 0)$, then $h(0, 0)$ is the root r of $T_{6,2,2}$. If $(x, y) \neq (0, 0)$, then as argued in Lemma 4.0.4, there exists $n_{(x,y)}$ with $(x, y) \in X'_{n_{(x,y)}} - X'_{n_{(x,y)}-1}$ and that there exists a unique $(u, v) \in X'_{n_{(x,y)}-1}$ such that $(x, y) = (u \pm \frac{1}{3^{n_{(x,y)}-1}}, v)$ or $(x, y) = (u, v \pm \frac{1}{3^{n_{(x,y)}-1}})$ or $(x, y) = (u \pm \frac{1}{3^{n_{(x,y)}-1}}, v \pm \frac{1}{3^{n_{(x,y)}-1}})$.

As visualized in Figure 4, we can see as an example X'_0, X'_1 and an excerpt of X'_2 . For clarity, we illustrated some points of X'_1 as a line. We clearly can see, that any point has his own "square", where the points within it can be only build from its center point. This is because, the "square size" gets exponentially smaller with each level, due to its construction. That means, in each iteration they can not intersect.

Now we define h as

$$h(x, y) = \begin{cases} \text{The left } R_1\text{-successor of } h(u, v) & \text{if } (x, y) = (u - \frac{1}{3^{n(x,y)-1}}, v) \\ \text{The right } R_1\text{-successor of } h(u, v) & \text{if } (x, y) = (u + \frac{1}{3^{n(x,y)-1}}, v) \\ \text{The left } R_2\text{-successor of } h(u, v) & \text{if } (x, y) = (u, v - \frac{1}{3^{n(x,y)-1}}) \\ \text{The right } R_2\text{-successor of } h(u, v) & \text{if } (x, y) = (u, v + \frac{1}{3^{n(x,y)-1}}) \\ \text{The first remaining successor of } h(u, v) & \text{if } (x, y) = (u + \frac{1}{3^{n(x,y)-1}}, v + \frac{1}{3^{n(x,y)-1}}) \\ & \text{or } (x, y) = (u - \frac{1}{3^{n(x,y)-1}}, v - \frac{1}{3^{n(x,y)-1}}) \\ \text{The last remaining successor of } h(u, v) & \text{if } (x, y) = (u + \frac{1}{3^{n(x,y)-1}}, v - \frac{1}{3^{n(x,y)-1}}) \\ & \text{or } (x, y) = (u - \frac{1}{3^{n(x,y)-1}}, v + \frac{1}{3^{n(x,y)-1}}) \end{cases}$$

It is still left to prove, that h is open and continuous w.r.t. all three topologies. We




Figure 7: This is an excerpt of $X \times X$.

will show it for τ_1 and τ . For τ_2 it is similar as for τ_1 .

First we observe the basis for τ_1 is

$$\{(x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times \{y\} \mid (x, y) \in X \times X\}$$

and for τ is

$$\{(x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times (y - \frac{1}{3^{n(x,y)}}, y + \frac{1}{3^{n(x,y)}}) \mid (x, y) \in X \times X\}$$

We also observe that a basis for the topology on $T_{6,2,2}$ from R_1 is $\{B_t^1\}_{t \in T_{6,2,2}}$ where $B_t^1 = \{s \in T_{6,2,2} \mid tR_1s\}$ and from R is $\{B_t\}_{t \in T_{6,2,2}}$ where $B_t = \{s \in T_{6,2,2} \mid tRs\}$. The

following arguments are carried over from Proposition 4.0.5.

Now, to see that g is open w.r.t. τ_1 , we pick an open set $(x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times \{y\} \in \tau_1$. In similar way, we can show $g((x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times \{y\}) = B_{g(x,y)}^1$. Thus g is open. To show g is continuous, it suffices to show that for each $t \in T_{6,2,2}$, the g -inverse image of B_t^1 belongs to τ_1 . Let $(x, y) \in g^{-1}(B_t^1)$. Then $tR_1g(x, y)$. Hence, it holds for (x, y) that $g((x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times \{y\}) = B_{g(x,y)}^1$. By $tR_1g(x, y)$, we get $B_{g(x,y)}^1 \subseteq B_t^1$. Thus, we found a neighborhood U of (x, y) s.t. $U = ((x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times \{y\}) \subseteq g^{-1}(B_t^1)$, implying g is continuous.

To see g is open w.r.t. τ , we pick an open set $(x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times (y - \frac{1}{3^{n(x,y)}}, y + \frac{1}{3^{n(x,y)}})$. A similar argument as in Proposition 4.0.5, yields us $g((x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times (y - \frac{1}{3^{n(x,y)}}, y + \frac{1}{3^{n(x,y)}})) = B_{g(x,y)}$.

To show g is continuous, it suffices to show for each $t \in T_{6,2,2}$ that $g^{-1}(B_t) \in \tau$. Let $(x, y) \in g^{-1}(B_t)$. Then $tRg(x, y)$. But then, it holds that $g((x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times (y - \frac{1}{3^{n(x,y)}}, y + \frac{1}{3^{n(x,y)}})) = B_{g(x,y)} \subseteq B_t$. Thus, we found a $U \in \tau$ s.t. $U = (x - \frac{1}{3^{n(x,y)}}, x + \frac{1}{3^{n(x,y)}}) \times (y - \frac{1}{3^{n(x,y)}}, y + \frac{1}{3^{n(x,y)}}) \subseteq g^{-1}(B_t)$. □

To complete the proof, if $TPL \not\models \phi$, then by Theorem 4.0.8, there is a valuation v on $T_{6,2,2}$ such that $(T_{6,2,2}, v), r \not\models \phi$. Now we define a valuation ξ on $X \times X$ by $\xi(p) = h^{-1}(v(p))$. Since h is continuous and open w.r.t. all three topologies and $h(0, 0) = r$, we have that $(0, 0)$ and r are 3-topo-bisimilar. Therefore, $(X \times X, \xi), (0, 0) \not\models \phi$. Similar as in Theorem 4.0.3, we argue that $X \times X$ is homeomorphic to $\mathbb{Q} \times \mathbb{Q}$ w.r.t. all three topologies, it follows that ϕ is refutable on $\mathbb{Q} \times \mathbb{Q}$.

Corollary 5.0.9. *In the language $L_{\square, \square_1, \square_2}$, TPL is the logic of products of arbitrary topologies.*

Proof. Let X and Y be arbitrary topological spaces. By the interpretation of \square , it follows $Log(X), Log(Y) \supseteq S4$. Let's consider the full product of the spaces $X \times_t^+ Y$. Hence, with theorem 4.0.8, it is easy to see that $Log(X \times_t^+ Y) \supseteq S4 \times_t^+ S4 = TPL$. □

6 Products of neighbourhood frames, completeness results

We now turn our focus to neighbourhood semantics, and in particular, we revisit the main theorem from the paper "Modal Logic of Products of Neighbourhood Frames" [2]. This work established that, for any $L_1, L_2 \in \{D, D4, T, S4\}$ we have $L_1 \otimes L_2 = L_1 \times_n L_2$. Our goal is to reprove this theorem within our framework and then adapt its key ideas to prove our theorem.

6.1 Kripke tree frames, natural neighborhood version and their logic

We introduce some trees as Kripke frames, which are essential for the proof.

Definition 6.1.1. *Let A be a nonempty set.*

$$A^* = \{a_1 \dots a_k \mid a_i \in A\}$$

is the set of all finite sequences of elements from A , including the empty sequence Λ . Elements from A^ will be denoted as \vec{a} . The length of a sequence $\vec{a} = a_1 \dots a_k$ is k (also $l(\vec{a}) = k$) and the length of Λ is 0 ($l(\Lambda) = 0$). Concatenation is denoted by " \cdot " : $(a_1 \dots a_k) \cdot (b_1 \dots b_l) = \vec{a} \cdot \vec{b} = a_1 \dots a_k b_1 \dots b_l$.*

Definition 6.1.2. *Let A be a nonempty set. We define an infinite frame $F_{in}[A] = (A^*, R)$ s.t for $\vec{a}, \vec{b} \in A^*$*

$$\vec{a} R \vec{b} \Leftrightarrow \exists x \in A (\vec{b} = \vec{a} \cdot x)$$

Furthermore we define :

$$F_{rn}[A] = (A^*, R^r), \text{ where } R^r = R \cup Id \text{ (reflexive closure)}$$

$$F_{it}[A] = (A^*, R^*), \text{ where } R^* = \bigcup_{i=1}^{\infty} R^i \text{ (transitive closure)}$$

$$F_{rt}[A] = (A^*, R^{r*})$$

where "t" stands for transitive, "n" for non-transitive, "r" for reflexive and "i" for ir-reflexive.

For now, we will use the following notion to generalize : $F_{\xi\eta}$ where $\xi \in \{i, r\}$ and $\eta \in \{t, n\}$

Definition 6.1.3. *Let $F_1 = F_{\xi_1\eta_1}[A] = (A^*, R_1)$ and $F_2 = F_{\xi_2\eta_2}[B] = (B^*, R_2)$, where $\xi_1, \xi_2 \in \{i, r\}$ and $\eta_1, \eta_2 \in \{t, n\}$. Furthermore, we assume $A = \{a_1, a_2, \dots\}$ and $B = \{b_1, b_2, \dots\}$ with $A \cap B = \emptyset$. Then we define the frame $F_1 \otimes F_2 = (W, R'_1, R'_2)$ as follows :*

$$W = (A \cup B)^*$$

$$\vec{x}R'_1\vec{y} \Leftrightarrow \vec{y} = \vec{x} \cdot \vec{z} \text{ for some } \vec{z} \in A^* \text{ such that } \Lambda R_1\vec{z}$$

$$\vec{x}R'_2\vec{y} \Leftrightarrow \vec{y} = \vec{x} \cdot \vec{z} \text{ for some } \vec{z} \in B^* \text{ such that } \Lambda R_2\vec{z}$$

Proposition 6.1.4 ([14], [15]). *Let F_1 and F_2 be as in Definition 5.1.3 . Then*

$$\text{Log}(F_1 \otimes F_2) = \text{Log}(F_1) \otimes \text{Log}(F_2)$$

Proposition 6.1.5 ([17]). *Let $F_{in} = F_{in}[\mathbb{N}]$, $F_{rn} = F_{rn}[\mathbb{N}]$, $F_{it} = F_{it}[\mathbb{N}]$ and $F_{rt} = F_{rt}[\mathbb{N}]$. Then the following holds:*

$$\text{Log}(F_{in}) = D$$

$$\text{Log}(F_{rn}) = T$$

$$\text{Log}(F_{it}) = D4$$

$$\text{Log}(F_{rt}) = S4$$

Definition 6.1.6. *Let $F = (W, R)$ be a Kripke frame. We define an n -frame $N(F) = (W, \tau)$ as follows. For any $w \in W$ we have :*

$$\tau(w) = \{U \mid R(w) \subseteq U \subseteq W\}$$

Lemma 6.1.7. *Let $F = (W, R)$ be a Kripke frame. Then*

$$\text{Log}(F) = \text{Log}(N(F))$$

Proof. The proof is by structural induction. □

6.2 Main Construction

In the following, we construct a useful neighborhood frame called $N_\omega[F]$ based on a tree frame F . We will show, that the construction has the same logic as the original frame. At the end, we construct a bounded morphism from the product of $N_\omega[F]$ frames to the fusion of frames F .

Definition 6.2.1. *Let $F = (A^*, R) = F_{\xi\eta}[A]$ and $0 \notin A$. We define "pseudo-infinite" sequences*

$$X = \{a_1a_2a_3\ldots \mid a_i \in A \cup \{0\} \text{ and } \exists N \forall k \geq N : a_k = 0\}$$

Furthermore, we define $f_F : X \rightarrow A^$ to be the function, that deletes all zeros.*

Example : Say $12034002340^\omega \in X$ (0^ω denotes infinitely many zeros). Then $f_F(12034002340^\omega) = 1234234$

Definition 6.2.2. Let $F = (A^*, R) = F_{\xi\eta}[A]$ and $0 \notin A$. Assume the function f_F and the set X as defined before. For $\alpha \in X$ such that $\alpha = a_1 a_2 \dots$ we define

$$st(a) = \min\{N \mid \forall k \geq N : a_k = 0\}$$

$$a \upharpoonright_k = a_1 a_2 \dots a_k$$

$$U_k(\alpha) = \{\beta \mid f_F(\alpha) R f_F(\beta) \text{ and } \alpha \upharpoonright_m = \beta \upharpoonright_m, \text{ where } m = \max(k, st(\alpha))\}$$

Remark : Let $\alpha \in X$ with $st(\alpha) = n$. Then we have that $U_n(\alpha) = U_j(\alpha)$ for any $j \leq n$

Lemma 6.2.3. $U_k(\alpha) \subseteq U_m(\alpha)$, whenever $k \geq m$.

Proof. Let $\beta \in U_k(\alpha)$. Since $\alpha \upharpoonright_k = \beta \upharpoonright_k$ and $k \geq m$, we have $\alpha \upharpoonright_m = \beta \upharpoonright_m$. It follows, $\beta \in U_m(\alpha)$. \square

Definition 6.2.4. Due to Lemma 5.2.3 the sets $U_n(\alpha)$ forms a filter base. So we can define :

$$\tau(\alpha) \text{ is a filter with base } \{U_n(\alpha) \mid n \in \mathbb{N}\}$$

$$N_\omega = (X, \tau) \text{ is the } n\text{-frame based on } F$$

Lemma 6.2.5. Let $F = (A^*, R) = F_{\xi\eta}[A]$. Based on that, let $N_\omega(F) = (X, \tau)$, $N(F) = (A^*, \sigma)$ and $f_F : N_\omega(F) \rightarrow N(F)$. Then for any $m \in \mathbb{N}$ and $x \in X$ with $x = a_1 a_2 \dots$ we have

$$f_F(U_m(x)) = R(f_F(x))$$

Proof. \subseteq : Let $f_F(\alpha) \in f_F(U_m(x))$ with $\alpha \in U_m(x)$. By definition of $U_m(x)$, we get $f_F(\alpha) \in R(f_F(x))$.

\supseteq : We pick $\vec{a} \in R(f_F(x))$. We have to find $\beta \in U_m(x)$ s.t $f_F(\beta) = \vec{a}$. We assume R is irreflexive and non-transitive. The other cases are similar.

Because $\vec{a} \in R(f_F(x))$, there must exists $c \in A$ such that $\vec{a} = f_F(x) \cdot c$. We construct $\beta = x \upharpoonright_m \cdot c 0^\omega$. Hence, $f(\beta) = f_F(x \upharpoonright_m) \cdot f_F(c)$ and because $0 \notin A$ we get $f_F(x \upharpoonright_m) \cdot c = \vec{a}$. \square

Lemma 6.2.6. Let $F = (A^*, R) = F_{\xi\eta}[A]$. Then $f_F : N_\omega(F) \rightarrow N(F)$ is a bounded morphism.

Proof. From now on this proof we will omit the subindex in f_F . Let $N_\omega(F) = (X, \tau)$ and $N(F) = (A^*, \sigma)$.

For surjectivity, we pick any $\vec{x} \in A^*$. But then, $\vec{x} 0^\omega \in X$. Hence, $f(\vec{x} 0^\omega) = \vec{x}$.

For the next condition, assume that $x \in X$ and $U \in \tau(x)$. We need to prove that $f(U) \in \sigma(f(x))$. That means $R(f(x)) \subseteq f(U)$. Because $U \in \tau(x)$, there is a m such that $U_m(x) \subseteq U$. By Lemma 5.2.5 we have $f(U_m(x)) = R(f(x))$. It follows,

$$R(f(x)) = f(U_m(x)) \subseteq f(U)$$

Assume $x \in X$ and V is a neighborhood of x , i.e $R(f(x)) \subseteq V$. We need to prove that there exists $U \in \tau(x)$, such that $f(U) \subseteq V$. As U we pick $U_m(x)$ for any $m \in \mathbb{N}$. By Lemma 5.2.5 we get $f(U_m(x)) = R(f(x))$. Hence,

$$f(U_m(x)) = R(f(x)) \subseteq V$$

□

Corollary 6.2.7. *For frame $F = F_{\xi\eta}[A]$ we have $\text{Log}(N_\omega(F)) \subseteq \text{Log}(F)$.*

Proof. It follows from Corollary 2.2.5, Lemma 5.1.7 and Lemma 5.2.6

$$\text{Log}(N_\omega(F)) \subseteq \text{Log}(N(F)) = \text{Log}(F)$$

□

Proposition 6.2.8. *Let $F_{in} = F_{in}[\mathbb{N}]$, $F_{rn} = F_{rn}[\mathbb{N}]$, $F_{it} = F_{it}[\mathbb{N}]$ and $F_{rt} = F_{rt}[\mathbb{N}]$. Then*

$$\text{Log}(N_\omega(F_{in})) = D$$

$$\text{Log}(N_\omega(F_{rn})) = T$$

$$\text{Log}(N_\omega(F_{it})) = D4$$

$$\text{Log}(N_\omega(F_{rt})) = S4$$

Proof. In all these cases, the inclusion from left to right follows from Proposition 5.1.5 and Corollary 5.2.7. Now the converse direction. Assume $X = (X, \tau)$.

It is easy to check that $X \models D$ iff for each $x \in X : \emptyset \notin \tau(x)$. For $N_\omega(F_{in})$ and $N_\omega(F_{it})$ this holds.

It is easy to check that $X \models T$ iff we have $x \in U \in \tau(x)$ for any x and U . For $N_\omega(F_{rn})$ and $N_\omega(F_{rt})$ this holds.

Now we check $X \models 4$ iff for each $U \in \tau(x) : \{y \mid U \in \tau(y)\} \in \tau(x)$.

\supseteq : Let $x \in X$ and assume $X, x \models \Box p$. That means there exists $U \in \tau(x)$ s.t $U \subseteq V(p)$. By assumption we have $S = \{y \mid U \in \tau(y)\} \in \tau(x)$. But then $X, x \models \Box \Box p$ because we can just pick the set S .

\subseteq : By contradiction, assume there exists a U s.t $\{y \mid U \in \tau(y)\} \notin \tau(x)$. Let $X, x \models \Box p$ where $V(p) = U$. If $X, x \models \Box \Box p$ then it must be the case that

$S = \{y \in X \mid X, y \models \Box p\} \in \tau(x)$. $X, y \models \Box p$ means $U \in \tau(y)$. That means $S = \{y \in X \mid U \in \tau(y)\}$. But by assumption $S \notin \tau(x)$. Hence, $X, x \not\models \Box \Box p$. But that's a contradiction. This also holds for $N_\omega(F_{it})$ and $N_\omega(F_{rt})$ because we have for any $y \in U_m(x)$ and $k \geq m : U_k(y) \subseteq U_m(x)$.

□

Assume $F_1 = (A^*, R_1) = F_{\eta_1 \xi_1}[A]$ and $F_2 = (B^*, R_2) = F_{\eta_2 \xi_2}[B]$ with $A \cap B = \emptyset$, $A = \{a_1, a_2, a_3, \dots\}$ and $B = \{b_1, b_2, b_3, \dots\}$. Consider the product of n-frames $F'_1 = (X_1, \tau_1) = N_\omega(F_1)$ and $F'_2 = (X_2, \tau_2) = N_\omega(F_2)$ is

$$X = (X_1 \times X_2, \tau'_1, \tau'_2) = N_\omega(F_1) \times_n N_\omega(F_2)$$

Furthermore, we have $F_1 \otimes F_2 = ((A \cup B)^*, R'_1, R'_2)$ as defined in Definition 5.1.3. We consider the neighborhood version

$$N(F_1 \otimes F_2) = ((A \cup B)^*, \sigma'_1, \sigma'_2)$$

Now we define $g : X_1 \times X_2 \rightarrow (A \cup B)^*$ as follows. For $(\alpha, \beta) \in X_1 \times X_2$ with $\alpha = a_1 a_2 \dots$ and $\beta = b_1 b_2 \dots$ we define $g(\alpha, \beta)$ to be the finite sequence which we get after eliminating all zeros from the infinite sequence $a_1 b_1 a_2 b_2 \dots$.

Example : Let $\alpha = 012340^\omega$ and $\beta = 0ab00e0^\omega$. Then $g(\alpha, \beta) = 1a2b34e$.

Lemma 6.2.9. *Let X and $N(F_1 \otimes F_2)$ be as defined before and $(\alpha, \beta) \in X_1 \times X_2$. Then for any $m > \max\{st(\alpha), st(\beta)\}$ we have*

$$R'_1(g(\alpha, \beta)) = g(U_m(\alpha) \times \{\beta\})$$

.

Proof. We will show both direction by assuming $F_1 = F_{in}[A]$. For $F_{it}[A], F_{rt}[A]$ and $F_{rn}[A]$ it is similar.

\subseteq : Let $\vec{w} \in R'_1(g(\alpha, \beta))$. By definition we get, there exists $\vec{c} \in A^*$ where $\vec{w} = g(\alpha, \beta) \cdot \vec{c}$ and $\Lambda R_1 \vec{c}$. Because R_1 is irreflexive and non-transitive, we get $\vec{c} \in A$. We construct (ζ, β) where $\zeta \in U_m(\alpha)$ and $g(\zeta, \beta) = \vec{w}$. For that, we can take $\zeta = \alpha \upharpoonright_m \cdot \vec{c} 0^\omega$. Obviously, $\zeta \in U_m(\alpha)$. Because $m \in \max\{st(\alpha), st(\beta)\}$, we have $g(\alpha \upharpoonright_m, \beta \upharpoonright_m) = g(\alpha, \beta)$. Hence, $g(\zeta, \beta) = g(\alpha \upharpoonright_m, \beta \upharpoonright_m) \cdot g(\vec{c} 0^\omega, 0^\omega) = g(\alpha \upharpoonright_m, \beta \upharpoonright_m) \cdot \vec{c} = \vec{w}$.

\supseteq : Assume $\zeta \in U_m(\alpha)$. We have to show $g(\zeta, \beta) \in R'_1(g(\alpha, \beta))$. Because R_1 is irreflexive and non-transitive, it suffices to find a $\vec{d} \in A$ s.t. $g(\alpha, \beta) \cdot \vec{d} = g(\zeta, \beta)$. By choosing m is maximal and $\zeta \upharpoonright_m = \alpha \upharpoonright_m$, we have $g(\zeta_m, \beta_m) = g(\alpha, \beta)$. We also know $f_F(\alpha) R_1 f_F(\zeta)$, that means there exists a $\vec{d} \in A$ s.t. $f_F(\alpha) \cdot \vec{d} = f_F(\zeta)$. This \vec{d} must appear at a point after ζ_m . We can follow $g(\zeta, \beta) = g(\alpha, \beta) \cdot \vec{d}$. Hence, $g(\zeta, \beta) \in R'_1(g(\alpha, \beta))$.

Remark : If $m \leq \max\{st(\alpha), st(\beta)\}$, then we cannot guarantee the equality. Assume R_1 as before and $\alpha = 123000^\omega$ and $\beta = d0b0a0^\omega$. Lets pick $m = 3$ and $\zeta = 123100^\omega$. Then we have $g(\alpha, \beta) = 1d23ba$ and $g(\zeta, \beta) = 1d23b1a$. Obviously, we don't have $g(\zeta, \beta) \in R'_1(g(\alpha, \beta))$.

Furthermore, we can show $R'_2(g(\alpha, \beta)) = g(\alpha \times U_m(\beta))$ similar as above.

□

Lemma 6.2.10. *Function $g : X \rightarrow N(F_1 \otimes F_2)$ is a bounded morphism.*

Proof. Let $\vec{z} = z_1 z_2 \dots z_n \in (A \cup B)^*$. Define for $i \leq n$:

$$x_i = \begin{cases} z_i, & \text{if } z_i \in A; \\ 0, & \text{if } z_i \notin A. \end{cases} \quad y_i = \begin{cases} z_i, & \text{if } z_i \in B; \\ 0, & \text{if } z_i \notin B. \end{cases}$$

Let $\alpha = x_1 x_2 \dots x_n 0^\omega$ and $\beta = y_1 y_2 \dots y_n 0^\omega$. Then $g(\alpha, \beta) = \vec{z}$. Hence, g is surjective.

For the next conditions we check only for τ'_1 and σ_1 . The other case is similar. Assume $(\alpha, \beta) \in X_1 \times X_2$ and $U \in \tau'_1(\alpha, \beta)$. We have to show $g(U) \in \sigma_1(g(\alpha, \beta))$. That means $R'_1(g(\alpha, \beta)) \subseteq g(U)$. Pick a $m > \max\{st(\alpha), st(\beta)\}$ s.t. $U_m(\alpha) \times \{\beta\} \subseteq U$. We can pick such m , because $U \in \tau'_1(\alpha, \beta)$. So there exists $U_k(\alpha) \in \tau(\alpha)$ s.t. $U_k(\alpha) \times \{\beta\} = U$. If $k > \max\{st(\alpha), st(\beta)\}$ then we are done. Else, by Lemma 5.2.3, we have for any $n \geq k : U_n(\alpha) \subseteq U_k(\alpha)$. So we can lift the k til we reach m . Then we use Lemma 5.2.9 to get the following :

$$R'_1(g(\alpha, \beta)) = g(U_m(\alpha) \times \{\beta\}) \subseteq g(U)$$

For the last condition we assume $(\alpha, \beta) \in X_1 \times X_2$ and $V \in \sigma_1(g(\alpha, \beta))$ (or rather $R'_1(g(\alpha, \beta)) \subseteq V$). We need to prove there exists $U \in \tau'_1(\alpha, \beta)$, such that $g(U) \subseteq V$. As U we take $U_m(\alpha) \times \{\beta\}$ for some $m > \max\{st(\alpha), st(\beta)\}$. Hence, by Lemma 5.2.9 we get :

$$g(U_m(\alpha) \times \{\beta\}) = R'_1(g(\alpha, \beta)) \subseteq V$$

□

6.3 Adapting the key ideas

Now we will show $g : N_\omega(T_{\omega[rn]}) \times_n^+ N_\omega(T_{\omega[rn]}) \rightarrow N(T_{\omega, \omega, \omega[rn]})$ is a bounded morphism. Let $(T_{\omega[rn]})_1 = (\mathbb{N}_1, R_1)$ and $(T_{\omega[rn]})_2 = (\mathbb{N}_2, R_2)$ as defined in Definition 3.0.3. Then we take the n -frames $N_\omega(T_{\omega[rn]})_1 = (X_1, \tau_1)$ and $N_\omega(T_{\omega[rn]})_2 = (X_2, \tau_2)$. For the proof, we will omit the subscripts after the frame.

Let's consider the full product of the n -frames :

$$N_\omega(T_{\omega[rn]}) \times_n^+ N_\omega(T_{\omega[rn]}) = (X_1 \times X_2, \tau'_1, \tau'_2, \tau)$$

We say $N(T_{\omega, \omega, \omega[rn]}) = ((\mathbb{N}_1 \cup \mathbb{N}_2 \cup \mathbb{N})^*, \sigma_1, \sigma_2, \sigma)$ where the tree $T_{\omega, \omega, \omega[rn]} = ((\mathbb{N}_1 \cup \mathbb{N}_2 \cup \mathbb{N})^* R'_1, R'_2, R)$ is as defined in Definition 3.0.3.

In order to define the bounded morphism, we have to fix a bijection first. Let $h : \mathbb{N}_1 \times \mathbb{N}_2 \rightarrow \mathbb{N}$ be a bijection.

Next, we define function $k : (\mathbb{N}_1 \cup \{0\}) \times (\mathbb{N}_2 \cup \{0\}) \rightarrow \mathbb{N}_1 \cup \mathbb{N}_2 \cup \mathbb{N} \cup \{0\}$ as follows :

$$f(a, b) = \begin{cases} a, & \text{if } b = 0; \\ b, & \text{if } a = 0; \\ h(a, b), & \text{otherwise.} \end{cases}$$

Let $(\alpha, \beta) \in X_1 \times X_2$ with $\alpha = a_1 a_2 \dots$ and $\beta = b_1 b_2 \dots$. We define

$$g'(\alpha, \beta) = f(a_1, b_1) f(a_2, b_2) \dots$$

At last, we define $g(\alpha, \beta)$ as $g'(\alpha, \beta)$ but removing all zeros.

Example 6.3.1. Assume h is the Cantor pairing function $h(m, n) = \frac{(m+n)(m+n+1)}{2} + n$ and $(\alpha, \beta) = (12_{\mathbb{N}_1} 021_{\mathbb{N}_1} 0^\omega, 1_{\mathbb{N}_2} 047_{\mathbb{N}_2} 0^\omega) \in X_1 \times X_2$.

Then $g'(\alpha, \beta) = f(1, 1) f(2, 0) f(0, 4) f(2, 7) f(1, 0) f(0, 0) \dots = h(1, 1) 24 h(2, 7) 10^\omega = 424 52 10^\omega$. Hence, $g(\alpha, \beta) = 424 52 1$.

Lemma 6.3.2. Function $g : N_\omega(T_{\omega[rn]}) \times_n^+ N_\omega(T_{\omega[rn]}) \rightarrow N(T_{\omega, \omega, \omega[rn]})$ is a bounded morphism.

Proof. Let $\vec{z} = z_1 z_2 \dots z_n \in (\mathbb{N}_1 \cup \mathbb{N}_2 \cup \mathbb{N})^*$. We define for $i \leq n$

$$x_i = \begin{cases} z_i, & \text{if } z_i \in \mathbb{N}_1; \\ a_1, & \text{if } z_i = h(a_1, b_2); \\ 0, & \text{other} \end{cases} \quad y_i = \begin{cases} z_i, & \text{if } z_i \in \mathbb{N}_2; \\ b_2, & \text{if } z_i = h(a_1, b_2); \\ 0, & \text{other} \end{cases}$$

Let $\alpha = x_1 x_2 \dots x_n 0^\omega$ and $\alpha = y_1 y_2 \dots y_n 0^\omega$. But then $g(\alpha, \beta) = \vec{z}$. Hence, g is surjective. For the next conditions we show it for τ and σ . For τ'_1, σ_1 and τ'_2, σ_2 we can argue similar as in Lemma 5.2.9 and 5.2.10.

First we need to show, $R(g(\alpha, \beta)) = g(U_m(\alpha) \times U_m(\beta))$ for $m > \max\{st(\alpha), st(\beta)\}$.

\supseteq : Pick $\zeta \in U_m(\alpha), v \in U_m(\beta)$. Show $g(\zeta, v) \in R(g(\alpha, \beta))$. By definition, we need to find a $\vec{c} \in \mathbb{N}_1 \cup \mathbb{N}_2 \cup \mathbb{N} \cup \{\epsilon\}$ s.t. $g(\alpha, \beta) \cdot \vec{c} = g(\zeta, v)$. By choosing $m > \max\{st(\alpha), st(\beta)\}$, $\alpha \upharpoonright_m = \zeta \upharpoonright_m$ and $\beta \upharpoonright_m = v \upharpoonright_m$, we get $g(\alpha, \beta) = g(\zeta_m, v_m)$. Furthermore, we have $f_F(\alpha) R_1 f_F(\zeta)$ and $f_F(\beta) R_2 f_F(v)$, that means $\exists \vec{d} \in \mathbb{N}_1 \cup \{\epsilon\}$ and $\exists \vec{e} \in \mathbb{N}_2 \cup \{\epsilon\}$ s.t. $f_F(\alpha) \cdot \vec{d} = f_F(\zeta)$ and $f_F(\beta) \cdot \vec{e} = f_F(v)$. So, \vec{d} must appear at a point after $\zeta \upharpoonright_m$ and \vec{e} in $v \upharpoonright_m$. But then we have $g(\zeta, v) = g(\alpha, \beta) \cdot g(\vec{d}, \vec{e})$.

Now there are four cases :

$$\begin{cases} g(\vec{d}, \vec{e}) \in \mathbb{N}_1, & \text{if } \vec{d} \in \mathbb{N}_1 \text{ \& } \vec{e} = \epsilon \\ g(\vec{d}, \vec{e}) \in \mathbb{N}_2, & \text{if } \vec{d} = \epsilon \text{ \& } \vec{e} \in \mathbb{N}_2 \\ g(\vec{d}, \vec{e}) \in \{\epsilon\}, & \text{if } \vec{d} = \vec{e} = \epsilon \\ g(\vec{d}, \vec{e}) \in \mathbb{N}, & \text{other} \end{cases}$$

Hence, $g(\zeta, v) \in R(g(\alpha, \beta))$.

\subseteq : Pick $\vec{w} \in R(g(\alpha, \beta))$. We will construct $\zeta \in U_m(\alpha)$ and $v \in U_m(\beta)$ s.t. $g(\zeta, v) = \vec{w}$. By definition there exists a $\vec{c} \in \mathbb{N}_1 \cup \mathbb{N}_2 \cup \mathbb{N} \cup \{\epsilon\}$ s.t. $g(\alpha, \beta) \cdot \vec{c} = \vec{w}$. Depending on the \vec{c} , we build $\zeta = \alpha \mid_m \cdot \vec{d} 0^\omega$ and $v = \beta \mid_m \cdot \vec{e} 0^\omega$ where

$$\begin{cases} \vec{d} = \vec{c}, \vec{e} = 0, & \text{if } \vec{c} \in \mathbb{N}_1; \\ \vec{d} = 0, \vec{e} = \vec{c}, & \text{if } \vec{c} \in \mathbb{N}_2; \\ \vec{d} = \vec{e} = 0 & \text{if } \vec{c} \in \{\epsilon\} \\ \vec{d} = \vec{\alpha}, \vec{e} = \vec{b} \text{ where } h(\vec{\alpha}, \vec{\beta}) = \vec{c} & \text{if } \vec{c} \in \mathbb{N} (\vec{\alpha} \in \mathbb{N}_1, \vec{\beta} \in \mathbb{N}_2); \end{cases}$$

We have $\zeta \in U_m(\alpha)$ and $v \in U_m(\beta)$. Furthermore, because of $g(\zeta \mid_m, v \mid_m) = g(\alpha, \beta)$, it holds that $g(\zeta, v) = g(\alpha, \beta) \cdot g(\vec{d}, \vec{e}) = g(\alpha, \beta) \cdot \vec{c} = \vec{w}$.

Now to check the conditions, we assume $x \in X_1$ and $y \in X_2$ and $U \in \tau(x, y)$. We need to prove $g(U) \in \sigma(g(x, y))$. In other words, show $R(g(x, y)) \subseteq g(U)$. We pick a $m > \max\{st(x), st(y)\}$ s.t. $U_m(x) \times U_m(y) \subseteq U$. But then we get

$$R(g(x, y)) = g(U_m(x) \times U_m(y)) \subseteq g(U)$$

Assume $x \in X_1$ and $x \in X_2$ and $R(g(x, y)) \subseteq V$. We need to find a $U \in \tau(x, y)$, s.t. $f(U) \subseteq V$. For U we pick $U_m(x) \times U_m(y)$ for some $m > \{st(x), st(y)\}$. But then

$$g(U_m(x) \times U_m(y)) = R(g(x, y)) \subseteq V$$

□

6.4 Completeness Results

Lemma 6.4.1. *Let $T_{\omega, \omega, \omega[rn]}$ be as in Definition 3.0.3. Then*

$$\text{Log}(T_{\omega, \omega, \omega[rn]}) = \text{Log}(N(T_{\omega, \omega, \omega[rn]}))$$

Proof. The proof is by structural induction. □

Corollary 6.4.2. *Let $F_1 = (A^*, R_1) = F_{\xi_1, \eta_1}[A]$ and $F_2 = (B^*, R_2) = F_{\xi_2, \eta_2}[B]$. Then*

$$\text{Log}(N_\omega(F_1) \times_n N_\omega(F_2)) \subseteq \text{Log}(F_1) \otimes \text{Log}(F_2)$$

. It follows from Corollary 2.2.5, Proposition 5.1.4 and Lemma 5.1.10.

Corollary 6.4.3. *Let $F_1, F_2 \in \{F_{in}, F_{rn}, F_{it}, F_{rt}\}$. Then*

$$\text{Log}(N_\omega(F_1) \times_n N_\omega(F_2)) = \text{Log}(F_1) \otimes \text{Log}(F_2)$$

Proof. The left to right inclusion follows from Corollary 5.4.3

To prove right to left, due to Proposition 5.1.5 and Proposition 5.2.8 we have $\text{Log}(N_\omega(F_i)) = \text{Log}(F_i)$ ($i \in \{1, 2\}$). Due to Proposition 2.2.8 we get $\text{Log}(N_\omega(F_1)) \otimes \text{Log}(N_\omega(F_2)) \subseteq \text{Log}(N_\omega(F_1)) \times_n \text{Log}(N_\omega(F_2))$. Because $N_\omega(F_1) \times_n N_\omega(F_2)$ is a frame in the n-product, we get $\text{Log}(N_\omega(F_1)) \times_n \text{Log}(N_\omega(F_2)) \subseteq \text{Log}(N_\omega(F_1) \times_n N_\omega(F_2))$. \square

Corollary 6.4.4. *Let $N_\omega(T_{\omega[rn]})_1 = F_1$ and $N_\omega(T_{\omega[rn]})_2 = F_2$ as defined before. Then*

$$T \times_n^+ T \subseteq \text{Log}(T_{\omega, \omega, \omega[rn]})$$

It follows from Corollary 2.2.5, Proposition 5.2.8, Lemma 5.3.2, Lemma 5.4.1.

Proof. First, it is the case that $\text{Log}(F_1) \times_n^+ \text{Log}(F_2) \subseteq \text{Log}(F_1 \times_n^+ F_2)$. We can argue as before. By Lemma 5.3.2 and Lemma 5.4.1 it holds $\text{Log}(F_1 \times_n^+ F_2) \subseteq \text{Log}(N(T_{\omega, \omega, \omega[rn]})) = \text{Log}(T_{\omega, \omega, \omega[rn]})$. But by Proposition 5.2.8 we have $\text{Log}(F_1) = \text{Log}(F_2) = T$. Hence, $\text{Log}(F_1) \times_n^+ \text{Log}(F_2) = T \times_n^+ T \subseteq \text{Log}(T_{\omega, \omega, \omega[rn]})$. \square

Theorem 6.4.5. *Let $L_1, L_2 \in \{S4, D4, D, T\}$. Then*

$$L_1 \times_n L_2 = L_1 \otimes L_2$$

Proof. Right to left is by Proposition 2.2.8.

For left to right, assume $L_1 = \text{Log}(F_1)$ and $L_2 = \text{Log}(F_2)$ for some $F_1, F_2 \in \{F_{in}, F_{rn}, F_{it}, F_{rt}\}$. It follows from Lemma 5.2.9 :

$$L_1 \times_n L_2 = \text{Log}(N_\omega(F_1)) \times_n^+ \text{Log}(N_\omega(F_2)) \subseteq \text{Log}(N_\omega(F_1) \times_n^+ N_\omega(F_2)) \subseteq \text{Log}(F_1 \otimes F_2) = \text{Log}(F_1) \otimes \text{Log}(F_2) = L_1 \otimes L_2 \quad \square$$

Theorem 6.4.6. *Let $N_\omega(T_{\omega[rn]})_1 = F'_1$ and $N_\omega(T_{\omega[rn]})_2 = F'_2$ as defined before. Then*

$$T \times_n^+ T = T \otimes T \otimes T + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$$

Proof. For left to right it is easy to check the axioms. We pick any frame F_1 and F_2 s.t. $F_1 \models T$ and $F_2 \models T$. We consider $F_1 \times_n^+ F_2$ the full product and let $(x, y) \in F_1 \times_n^+ F_2$.

Assume $F_1 \times_n^+ F_2, (x, y) \models \Box_1 p$. By definition, there exists a $U \in \tau'_1(x, y)$ s.t. $U \subseteq V(p)$. But we have $U \supseteq V \times \{y\}$ for some $V \in \tau(x)$. By assumption, $F_1 \models T$, so we can follow $x \in V$. Hence, $(x, y) \in V(p)$. For $\Box_2 p \rightarrow p$ it's done similar.

Let's check $\Box p \rightarrow p$. Assume $F_1 \times_n^+ F_2, (x, y) \models \Box p$. By definition there exists a $U \in \tau(x, y)$ s.t. $U \subseteq V(p)$. But $U \supseteq V \times W$ for some $V \in \tau_1(x)$ and $W \in \tau_2(y)$. Because of the assumption, we must have $x \in V$ and $y \in W$. Hence, $(x, y) \in V(p)$. It follows $F_1 \times_n^+ F_2, (x, y) \models p$.

Now we check the extra axiom. Assume $F_1 \times_n^+ F_2, (x, y) \Vdash \Box p$. By definition, there exists a $U \subseteq V(p)$, where $U \supseteq V \times W$ with $V \in \tau_1(x)$ and $W \in \tau_2(y)$. By assumption, we get $y \in W$. That means $U \in \tau'_1(x, y)$ because $U \supseteq V \times \{y\}$. The same argument we can apply to $\tau'_2(x, y)$. Hence, $F_1 \times_n^+ F_2, (x, y) \models \Box_1 p \wedge \Box_2 p$. The rest is clear.

Now for the other direction, we apply Corollary 5.4.4 to get $T \times_n^+ T \subseteq \text{Log}(T_{\omega, \omega, \omega[rn]})$. By Proposition 3.0.5 we have $\text{Log}(T_{\omega, \omega, \omega[rn]}) = TNL$. Hence,

$$T \times_n^+ T \subseteq T \otimes T \otimes T + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$$

□

Remark 6.4.7. Let $L \in \{D, D4\}$. Then it is not guaranteed that

$$L \times_n^+ L = L \otimes L \otimes L + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$$

This can be shown by a counterexample. We will only do it for D because for $D4$ it's done similar. Assume $F_1 = N_\omega(T_{\omega[in]})_1 = (X, \tau_1)$ and $F_2 = N_\omega(T_{\omega[in]})_2 = (Y, \tau_2)$. By Proposition 5.2.8, it follows $\text{Log}(F_1) = \text{Log}(F_2) = D$. Let's consider the full product $X = F_1 \times_n^+ F_2 = (W \times V, \tau'_1, \tau'_2, \tau)$. Then we get $\text{Log}(X) \supseteq D \times_n^+ D$. Now let $(x, y) \in X$, $(x, y) \models \Box p$ and $V(p) = U \times V$ where $U \in \tau_1(x)$ and $V \in \tau_2(y)$. By construction of $N_\omega(T_{\omega[rn]})$, $x \notin U$ and $y \notin V$, because $f_F(x)$ and $f_F(y)$ don't have a relation to itself in $T_{\omega[rn]}$. In order to satisfy $\Box_1 p$ (similar for $\Box_2 p$), we have to find $U' \in \tau'_1(x, y)$ s.t. $U' \subseteq V(p)$. But this is not possible because any U' contains only elements, where the second coordinate is y . It follows $X, (x, y) \not\models \Box p \rightarrow \Box_1 p \wedge \Box_2 p$. Hence, the extra axiom is not in $\text{Log}(X) \supseteq D \times_n^+ D$.

7 Conclusion

In this work, we reproved the theorems from "Multimodal Logics of Products of Topologies" and "Modal Logic of products of neighborhood frames". We showed the logic of the product of two arbitrary topological spaces is $S4 \otimes S4 \otimes S4 + \Box p \rightarrow \Box_1 p \wedge \Box_2 p$. The idea was to find a 3-topo-bisimulation between $T_{6,2,2}$ and $X \times X$ which is homeomorphic to $\mathbb{Q} \times \mathbb{Q}$. Additionally, by switching to neighborhood frames, it holds that for any $L_1, L_2 \in \{D, D4, T, S4\}$: $L_1 \otimes L_2 = L_1 \times_n L_2$. We constructed from a Kripke frame F a special neighborhood frame called $N_\omega(F)$ and then found a bounded morphism from the product of these constructed frame to the fusion of two frames. Inspired by these ideas, we first introduced a Kripke frame called $T_{\omega, \omega, \omega[rn]}$ where we proved FMP and used unravelling to show $\text{Log}(T_{\omega, \omega, \omega[rn]}) = TNL$. Afterwards, we constructed a bounded morphism from $N_\omega(T_{\omega[rn]}) \times_n^+ N_\omega(T_{\omega[rn]})$ onto $N(T_{\omega, \omega, \omega[rn]})$ to finish the proof.

There are tons of ways to continue the research. We may consider the logic K and ask what is the logic of $K \times_n^+ K$? What's about different combinations like : $D \times_n^+ T$, $T \times_n^+ K$, $S4 \times_n^+ D4$...? We can also try to find out whether for logic Λ with $T \subseteq \Lambda \subseteq S4$ the following holds :

$$\Lambda \otimes \Lambda \otimes \Lambda + \Box p \rightarrow \Box_1 p \wedge \Box_2 p = \Lambda \times_n^+ \Lambda$$

References

- [1] Johan van Benthem, Guram Bezhanishvili, Balder ten Cate, and Darko Sarenac. Multimodal logics of products of topologies. *Studia Logica*, 84:369–392, 2006.
- [2] Andrei Kudinov. Modal logic of some products of neighbourhood frames. In *Advances in Modal Logic, Volume 9*, pages 286–294, London, 2012. College Publications.
- [3] J.C.C. McKinsey and Alfred Tarski. The algebra of topology. *Annals of Mathematics*, 45:141–191, 1944.
- [4] K. Segerberg. Two-dimensional modal logic. *Journal of Philosophical Logic*, 2:77–96, 1973.
- [5] V.B. Shehtman. Two-dimensional modal logics. *Mathematical Notices of the USSR Academy of Sciences*, 23:417–424, 1978. (Translated from Russian.)
- [6] D. Sarenac. Modal logic and topological products. PhD Thesis, Stanford University, 2005.
- [7] Patrick Blackburn, Maarten de Rijke, and Yde Venema. *Modal Logic*. Cambridge University Press, August 2002.
- [8] Katsuhiko Sano. Axiomatizing hybrid products of monotone neighborhood frames. *Electronic Notes in Theoretical Computer Science*, 273:51–67, 2011.
- [9] Dana Scott. Advice on modal logic. In *Philosophical Problems in Logic: Some Recent Developments*, pages 143–173. D. Reidel, 1970.
- [10] Richard Montague. Universal grammar. *Theoria*, 36:373–398, 1970.
- [11] Eric Pacuit. *Neighborhood Semantics for Modal Logic*. Short Textbooks in Logic. Springer Cham, 2017.
- [12] R. Goldblatt. Diodorean modality in Minkowski spacetime. *Studia Logica*, 39:219–237, 1980.
- [13] K. Kuratowski and A. Mostowski. *Set Theory*. Studies in Logic and the Foundations of Mathematics, Vol. 86. North-Holland Publishing Co., Amsterdam–New York–Oxford; PWN—Polish Scientific Publishers, Warsaw, 1976.
- [14] M. Kracht and F. Wolter. Properties of independently axiomatizable bimodal logics. *Journal of Symbolic Logic*, 56(4):1469–1485, 1991.

- [15] K. Fine and G. Schurz. Transfer theorems for multimodal logics.
In J. Copeland, editor, *Logic and Reality: Proceedings of the Arthur Prior Memorial Conference*, pages 169–213. Oxford University Press, 1996.
- [16] G. Bezhanishvili and J. van Benthem. Modal Logics of Space.
In ILLC Prepublication Series PP-2006-08, Universiteit van Amsterdam, 2006.