

hello

IP = PSPACE

CONFERENCE, DATE

$\text{PSPACE} \subseteq \text{IP}$

For this inclusion, we use a well known PSPACE-complete problem, namely True-QBF

$\text{PSPACE} \subseteq \text{IP}$

For this inclusion, we use a well known PSPACE-complete problem, namely True-QBF

- QBF-Truth is the set of all valid quantified boolean formulas without free variables and for any variable p we have $p \in \{0, 1\}$

PSPACE \subseteq IP

For this inclusion, we use a well known PSPACE-complete problem, namely True-QBF

- QBF-Truth is the set of all valid quantified boolean formulas without free variables and for any variable p we have $p \in \{0, 1\}$
- $\forall x \exists y (x \vee y), \exists x \exists y \neg (x \wedge y)$

PSPACE \subseteq IP

For this inclusion, we use a well known PSPACE-complete problem, namely True-QBF

- QBF-Truth is the set of all valid quantified boolean formulas without free variables and for any variable p we have $p \in \{0, 1\}$
- $\forall x \exists y (x \vee y), \exists x \exists y \neg (x \wedge y)$
- QBF-Truth_{NNF} (abbrev. with QBF') is QBF-Truth but negations are only applied on variables
- $\exists x \exists y \neg (x \wedge y)$ is not in NNF but $\exists x \exists y (\neg x \vee \neg y)$

PSPACE \subseteq IP

For this inclusion, we use a well known PSPACE-complete problem, namely True-QBF

- QBF-Truth is the set of all valid quantified boolean formulas without free variables and for any variable p we have $p \in \{0, 1\}$
- $\forall x \exists y (x \vee y), \exists x \exists y \neg (x \wedge y)$
- QBF-Truth_{NNF} (abbrev. with QBF') is QBF-Truth but negations are only applied on variables
- $\exists x \exists y \neg (x \wedge y)$ is not in NNF but $\exists x \exists y (\neg x \vee \neg y)$
- $\text{QBF} \leq_m^P \text{QBF-Truth}_{\text{NNF}}$ can be easily done by the verifier
- it suffices to show $\text{QBF}' \in \text{IP}$ because we can reduce any problem in PSPACE to QBF in polynomial time

How do we find an algorithm for QBF' s.t it satisfies the IP conditions?

Arithmetization

The prover has to convince the verifier that the formula is valid but in case of an invalid formula it should reject with high probability (for all prover)

Arithmetization

The prover has to convince the verifier that the formula is valid but in case of an invalid formula it should reject with high probability (for all prover)

- The idea is to arithmetize the formula

Arithmetization

The prover has to convince the verifier that the formula is valid but in case of an invalid formula it should reject with high probability (for all prover)

- The idea is to arithmetize the formula
- $x \wedge y$ becomes $x*y$
- $x \vee y$ becomes $x+y$
- $\neg x$ becomes $1-x$

Arithmetization

The prover has to convince the verifier that the formula is valid but in case of an invalid formula it should reject with high probability (for all prover)

- The idea is to arithmetize the formula
- $x \wedge y$ becomes $x * y$
- $x \vee y$ becomes $x + y$
- $\neg x$ becomes $1 - x$

x	y	$x \wedge y = x * y$	$x \vee y = x + y$	$\neg(x \wedge y) = 1 - (x * y)$
0	0	0	0	1
0	1	0	1	1
1	0	0	1	1
1	1	1	2	0

$$\phi(x_1, x_2, \dots, x_n) = 1 \Leftrightarrow \phi_{arith}(x_1, x_2, \dots, x_n) > 0$$

Arithmetization

How do we arithmetize \forall and \exists ?

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$
- $\exists x \phi$ becomes $a_0 + a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$
- $\exists x \phi$ becomes $a_0 + a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Example : $\phi = \forall x \exists y \neg(x \wedge y)$

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$
- $\exists x \phi$ becomes $a_0 + a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Example : $\phi = \forall x \exists y \neg(x \wedge y)$

Arithmetize : $\neg(x \wedge y) \xrightarrow{\text{arith.}} 1 - (x * y) \xrightarrow{\exists \text{arith.}} \sum_{y \in \{0,1\}} 1 - (x * y) \xrightarrow{\forall \text{arith.}}$

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$
- $\exists x \phi$ becomes $a_0 + a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Example : $\phi = \forall x \exists y \neg(x \wedge y)$

Arithmetize : $\neg(x \wedge y) \xrightarrow{\text{arith.}} 1 - (x * y) \xrightarrow{\exists \text{arith.}} \sum_{y \in \{0,1\}} 1 - (x * y) \xrightarrow{\forall \text{arith.}}$

$$\prod_{x \in \{0,1\}} \sum_{y \in \{0,1\}} 1 - (x * y) = \phi_{\text{arith}}$$

- $\phi_{\text{arith}} = 2$

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$
- $\exists x \phi$ becomes $a_0 + a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Example : $\phi = \forall x \exists y \neg(x \wedge y)$

Arithmetize : $\neg(x \wedge y) \xrightarrow{\text{arith.}} 1 - (x * y) \xrightarrow{\exists \text{arith.}} \sum_{y \in \{0,1\}} 1 - (x * y) \xrightarrow{\forall \text{arith.}}$

$$\prod_{x \in \{0,1\}} \sum_{y \in \{0,1\}} 1 - (x * y) = \phi_{\text{arith}}$$

- $\phi_{\text{arith}} = 2$
- If ϕ is true then $\phi_{\text{arith}} > 0$
- If ϕ is false then $\phi_{\text{arith}} = 0$

Arithmetization

How do we arithmetize \forall and \exists ?

- $\forall x \phi$ becomes $a_0 * a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$
- $\exists x \phi$ becomes $a_0 + a_1$ where $a_0 = \phi[x := 0]$ and $a_1 = \phi[x := 1]$

Example : $\phi = \forall x \exists y \neg(x \wedge y)$

Arithmetize : $\neg(x \wedge y) \xrightarrow{\text{arith.}} 1 - (x * y) \xrightarrow{\exists \text{arith.}} \sum_{y \in \{0,1\}} 1 - (x * y) \xrightarrow{\forall \text{arith.}}$

$$\prod_{x \in \{0,1\}} \sum_{y \in \{0,1\}} 1 - (x * y) = \phi_{\text{arith}}$$

- $\phi_{\text{arith}} = 2$
- If ϕ is true then $\phi_{\text{arith}} > 0$
- If ϕ is false then $\phi_{\text{arith}} = 0$
- This can be shown by structural induction

Arithmetization

- $\phi = x$. If ϕ is true then $\phi_{arith} = 1$ and if ϕ is false then $\phi_{arith} = 0$
- $\phi = \neg x$ is the same but turned around

Arithmetization

- $\phi = x$. If ϕ is true then $\phi_{arith} = 1$ and if ϕ is false then $\phi_{arith} = 0$
- $\phi = \neg x$ is the same but turned around
- Suppose ϕ_1 and ϕ_2 are true, that means $\phi_{1arith} > 0$ and $\phi_{2arith} > 0$
(it's similar for the other case)

Arithmetization

- $\phi = x$. If ϕ is true then $\phi_{arith} = 1$ and if ϕ is false then $\phi_{arith} = 0$
- $\phi = \neg x$ is the same but turned around
- Suppose ϕ_1 and ϕ_2 are true, that means $\phi_{1arith} > 0$ and $\phi_{2arith} > 0$ (it's similar for the other case)
- $\phi_1 \wedge \phi_2, \phi_1 \vee \phi_2$ also holds
- For $\forall \phi_1$, we have by induction that $\phi_1[x := 0], \phi_1[x := 1]$ is true, so the multiplication of two positive value is positive (same for $\exists \phi_1$)

Protocol-Problem

How do we start the communication between prover and verifier?

Protocol-Problem

How do we start the communication between prover and verifier?

- On input $\langle \phi \rangle$, the prover sends a value $c > 0$ to the verifier and tries to convince that c is the arithmetic value of ϕ
(Remember : the verifier can not calculate its value by itself because it could be double exponential)

Protocol-Problem

How do we start the communication between prover and verifier?

- On input $\langle \phi \rangle$, the prover sends a value $c > 0$ to the verifier and tries to convince that c is the arithmetic value of ϕ
(Remember : the verifier can not calculate its value by itself because it could be double exponential)
- Problem : the value could be exponential but the verifier has only polynomial time

Protocol-Problem

How do we start the communication between prover and verifier?

- On input $\langle \phi \rangle$, the prover sends a value $c > 0$ to the verifier and tries to convince that c is the arithmetic value of ϕ
(Remember : the verifier can not calculate its value by itself because it could be double exponential)
- Problem : the value could be exponential but the verifier has only polynomial time

$\phi = \forall x_1 \dots \forall x_m \exists y \exists z. (y \vee z)$. What is ϕ_{arith} ? We calculate it step by step.

Protocol-Problem

How do we start the communication between prover and verifier?

- On input $\langle \phi \rangle$, the prover sends a value $c > 0$ to the verifier and tries to convince that c is the arithmetic value of ϕ
(Remember : the verifier can not calculate its value by itself because it could be double exponential)
- Problem : the value could be exponential but the verifier has only polynomial time

$\phi = \forall x_1 \dots \forall x_m \exists y \exists z. (y \vee z)$. What is ϕ_{arith} ? We calculate it step by step. Let $\phi' = \exists y \exists z (y \vee z)$. Then $\phi'_{arith} = \sum_{y \in \{0,1\}} \sum_{z \in \{0,1\}} y + z = 4$

Protocol-Problem

How do we start the communication between prover and verifier?

- On input $\langle \phi \rangle$, the prover sends a value $c > 0$ to the verifier and tries to convince that c is the arithmetic value of ϕ
(Remember : the verifier can not calculate its value by itself because it could be double exponential)
- Problem : the value could be exponential but the verifier has only polynomial time

$\phi = \forall x_1 \dots \forall x_m \exists y \exists z. (y \vee z)$. What is ϕ_{arith} ? We calculate it step by step. Let

$\phi' = \exists y \exists z (y \vee z)$. Then $\phi'_{arith} = \sum_{y \in \{0,1\}} \sum_{z \in \{0,1\}} y + z = 4$

$\phi_{arith} = \prod_{x_1 \in \{0,1\}} \dots \prod_{x_m \in \{0,1\}} \phi'_{arith} = 4^{2^m}$

- It holds that for formula ϕ with string length n : $\phi_{arith} \leq 2^{2^n}$
- We solve this problem by using modulo with a suitable value

Protocol-Problem

- Pick a value $k \geq 2^n$ with two conditions :
- k must be presentable in linear many bits

Protocol-Problem

- Pick a value $k \geq 2^n$ with two conditions :
- k must be presentable in linear many bits
- the calculation mod k must preserve ">0" for valid and "=0" for invalid formulas
- Number Theory Theorem : for any $a \leq 2^{2^n}$, $a > 0$, there exist a prime number $k \in [2^n, 2^{3n}]$ s.t $a \not\equiv 0 \pmod{k}$

Protocol continued

- Prover sends value c , prime number k and a proof b for the prime number property (it is possible to give a polynomial proof)

Protocol continued

- Prover sends value c , prime number k and a proof b for the prime number property (it is possible to give a polynomial proof)
- Verifier check $c > 0$, $k \in [2^n, 2^{3n}]$ and b is a correct proof for prime property

Even if k and b are correct, the verifier stays sceptical about c .

Protocol continued

- Prover sends value c , prime number k and a proof b for the prime number property (it is possible to give a polynomial proof)
- Verifier check $c > 0$, $k \in [2^n, 2^{3n}]$ and b is a correct proof for prime property

Even if k and b are correct, the verifier stays sceptical about c .

- If $\phi = \phi_1 \wedge \phi_2$, then ask prover to send a_1 and a_2 and check $c = a_1 * a_2$. If it's true then ask the prover to prove that the of ϕ_1 is a_1 and ϕ_2 is a_2
- For $\phi = \phi_1 \vee \phi_2$, we ask for a_1 and a_2 s.t $c = a_1 + a_2$

Protocol continued

- Prover sends value c , prime number k and a proof b for the prime number property (it is possible to give a polynomial proof)
- Verifier check $c > 0$, $k \in [2^n, 2^{3n}]$ and b is a correct proof for prime property

Even if k and b are correct, the verifier stays sceptical about c .

- If $\phi = \phi_1 \wedge \phi_2$, then ask prover to send a_1 and a_2 and check $c = a_1 * a_2$. If it's true then ask the prover to prove that the of ϕ_1 is a_1 and ϕ_2 is a_2
- For $\phi = \phi_1 \vee \phi_2$, we ask for a_1 and a_2 s.t $c = a_1 + a_2$
- In case $\phi = \forall x \phi_1$ we asked for a polynomial $p(x)$ that represents the arithmetic presentation of ϕ_1 where x is free and we check $c = p(0) * p(1)$
- If it is true, the verifier sends randomly a number d between $\{0, \dots, k-1\} = GF(K)$ and calculate $p(d)$. Now the verifier expects the prover to prove the value of $\phi_1[x := d]$ is $p(d)$
- The same process happens when we have $\phi = \exists \phi_1$, but we check $c = p(0) + p(1)$

Protocol continue

- When every variable got a number in $\text{GF}(K)$, say y_1, \dots, y_n the verifier calculates $\phi_{arith}(y_1, \dots, y_n)$ and accept if its equal to $q(y_1, \dots, y_n)$ (last polynomial sent by prover) else reject

Example

$$\phi = \forall x \exists y (\neg x \vee y) \wedge \exists z \exists w (z \vee w)$$

$$\phi_{arith} = \underbrace{\left(\prod_x \prod_y (1 - x) + y \right)}_{\phi_{1arith}} * \underbrace{\left(\sum_z \sum_w z + w \right)}_{\phi_{2arith}} \quad x, y, z, w \in \{0, 1\}$$


Example

$$\phi = \forall x \exists y (\neg x \vee y) \wedge \exists z \exists w (z \vee w)$$

$$\phi_{arith} = \underbrace{\left(\prod_x \prod_y (1 - x) + y \right)}_{\phi_{1arith}} * \underbrace{\left(\sum_z \sum_w z + w \right)}_{\phi_{2arith}} \quad x, y, z, w \in \{0, 1\}$$

Prover

$c=8, k=11, b$



Verifier

check $c > 0, 8, b$

Example

$$\phi = \forall x \exists y (\neg x \vee y) \wedge \exists z \exists w (z \vee w)$$

$$\phi_{arith} = \underbrace{\left(\prod_x \prod_y (1 - x) + y \right)}_{\phi_{1arith}} * \underbrace{\left(\sum_z \sum_w z + w \right)}_{\phi_{2arith}} \quad x, y, z, w \in \{0, 1\}$$

Prover

Verifier

$c=8, k=11, b$

check $c > 0, 8, b$

asks for $a_1 = \phi_1$ and $a_2 = \phi_2$

sends $a_1 = 2, a_2 = 4$

check $c = a_1 * a_2$

prove $a_1 = \phi_1, a_2 = \phi_2$

Example

$$\phi = \forall x \exists y (\neg x \vee y) \wedge \exists z \exists w (z \vee w)$$

$$\phi_{arith} = \underbrace{\left(\prod_x \prod_y (1 - x) + y \right)}_{\phi_{1arith}} * \underbrace{\left(\sum_z \sum_w z + w \right)}_{\phi_{2arith}} \quad x, y, z, w \in \{0, 1\}$$

Prover

Verifier

$c=8, k=11, b$

check $c > 0, 8, b$

asks for $a_1 = \phi_1$ and $a_2 = \phi_2$

sends $a_1 = 2, a_2 = 4$

check $c = a_1 * a_2$

prove $a_1 = \phi_1, a_2 = \phi_2$

send me $p_1(x)$ for ϕ_1 and $p_2(x)$ for ϕ_2

Example continue

Prover

calculate

$\phi_{1arith}(x),$

$\phi_{2arith}(z)$

sends $p_1(x) = \phi_{1arith}(x), \phi_{2arith}(z)$

$$\frac{\text{sends } p_1(x) = \phi_{1arith}(x), \phi_{2arith}(z)}{p_1(x) = \prod_y (1 - x) + y = x^2 - 3x + 2} \rightarrow$$

Verifier

check $a_1 = p_1(0) * p_1(1)$

check $a_2 = p_2(0) + p_2(1)$

Example continue

Prover

calculate

$\phi_{1arith}(x),$

$\phi_{2arith}(z)$

sends $p_1(x) = \phi_{1arith}(x), \phi_{2arith}(z)$

$$p_1(x) = \prod_y (1 - x) + y = x^2 - 3x + 2$$

Verifier

check $a_1 = p_1(0) * p_1(1)$

check $a_2 = p_2(0) + p_2(1)$

sends $p_1(d) = 0, p_2(g) = 7$

← ask for $p'_1(d, y), p'_2(g, w)$

Choose randomly $d, g \in$

$GF(k)$, say $d=2, g=3$

Example continue

Prover

calculate

$\phi_{1arith}(x),$

$\phi_{2arith}(z)$

Verifier

sends $p_1(x) = \phi_{1arith}(x), \phi_{2arith}(z)$ → check $a_1 = p_1(0) * p_1(1)$
 $p_1(x) = \prod_y (1 - x) + y = x^2 - 3x + 2$ check $a_2 = p_2(0) + p_2(1)$

sends $p_1(d) = 0, p_2(g) = 7$ ← Choose randomly $d, g \in GF(k)$, say $d=2, g=3$
ask for $p'_1(d, y), p'_2(g, w)$

sends $p'_1(d, y) = \phi_{1arith}(d, y)$ → $p'_1(d, 0) * p'_1(d, 1) = p_1(d)$
sends $p'_2(g, w) = \phi_{2arith}(g, w)$ $p'_2(g, 0) * p'_2(g, 1) = p_2(g)$
Choose $c, k \in GF(k)$

Example continue

Prover

calculate

$\phi_{1arith}(x),$

$\phi_{2arith}(z)$

Verifier

sends $p_1(x) = \phi_{1arith}(x), \phi_{2arith}(z)$ → check $a_1 = p_1(0) * p_1(1)$
 $p_1(x) = \prod_y (1 - x) + y = x^2 - 3x + 2$ check $a_2 = p_2(0) + p_2(1)$

sends $p_1(d) = 0, p_2(g) = 7$ Choose randomly $d, g \in GF(k)$, say $d=2, g=3$
← ask for $p'_1(d, y), p'_2(g, w)$

sends $p'_1(d, y) = \phi_{1arith}(d, y)$ → $p'_1(d, 0) * p'_1(d, 1) = p_1(d)$
sends $p'_2(g, w) = \phi_{2arith}(g, w)$ $p'_2(g, 0) * p'_2(g, 1) = p_2(g)$
Choose $c, k \in GF(k)$

Example continue

Prover

calculate

$\phi_{1arith}(x),$

$\phi_{2arith}(z)$

Verifier

sends $p_1(x) = \phi_{1arith}(x), \phi_{2arith}(z)$
—————→
 $p_1(x) = \prod_y (1 - x) + y = x^2 - 3x + 2$ check $a_1 = p_1(0) * p_1(1)$
check $a_2 = p_2(0) + p_2(1)$

sends $p_1(d) = 0, p_2(g) = 7$
←—————
ask for $p'_1(d, y), p'_2(g, w)$ Choose randomly $d, g \in GF(k), \text{say } d=2, g=3$

sends $p'_1(d, y) = \phi_{1arith}(d, y)$
—————→
sends $p'_2(g, w) = \phi_{2arith}(g, w)$ $p'_1(d, 0) * p'_1(d, 1) = p_1(d)$
 $p'_2(g, 0) * p'_2(g, 1) = p_2(g)$
Choose $c, k \in GF(k)$

The verifier check $\phi_{arith}(d, c, g, k) = p'_1(d, c) * p'_2(g, k)$. It accepts.

Next Example

$$\phi = \forall x \exists y \, x \wedge y \xrightarrow{\text{arith.}} \phi_{\text{arith}} = \prod_x \sum_y x * y$$

Next Example

$$\phi = \forall x \exists y \, x \wedge y \xrightarrow{\text{arith.}} \phi_{\text{arith}} = \prod_x \sum_y x * y$$

Prover can not tell the truth because the verifier would reject instantly.


Prover

Verifier

Next Example

$$\phi = \forall x \exists y x \wedge y \xrightarrow{\text{arith.}} \phi_{\text{arith}} = \prod_x \sum_y x * y$$


Prover can not tell the truth because the verifier would reject instantly.



Next Example

$$\phi = \forall x \exists y x \wedge y \xrightarrow{\text{arith.}} \phi_{\text{arith}} = \prod_x \sum_y x * y$$


Prover can not tell the truth because the verifier would reject instantly.



Next Example

$$\phi = \forall x \exists y x \wedge y \xrightarrow{\text{arith.}} \phi_{\text{arith}} = \prod_x \sum_y x * y$$

Prover can not tell the truth because the verifier would reject instantly.




The verifier check $\phi_{\text{arith}}(d, g) = p'_1(d, g)$. $12 \neq 28$. Verifier rejects.

Next Example

$$\phi = \forall x \exists y x \wedge y \xrightarrow{\text{arith.}} \phi_{\text{arith}} = \prod_x \sum_y x * y$$

Prover can not tell the truth because the verifier would reject instantly.



The verifier check $\phi_{\text{arith}}(d, g) = p'_1(d, g)$. $12 \neq 28$. Verifier rejects.

Simple QBF

- Problem : Polynomial could be exponential. The verifier can not verify it in polynomial time.

Simple QBF

- Problem : Polynomial could be exponential. The verifier can not verify it in polynomial time.
- Example : $\phi = \forall x_1, \dots, \forall x_m (x_1 \vee \dots \vee x_m) \xrightarrow{\text{arith}\phi(x_1)} \phi_{\text{arith}}(x) = \prod_{x_2} \dots \prod_{x_m} (x_1 + x_2 + \dots + x_m) \rightarrow \deg(\phi_{\text{arith}}(x)) \leq 2^{m-1}$
-

Correctness

- If ϕ is true, a truthful Prover can ensure that V accepts

Correctness

- If ϕ is true, a truthful Prover can ensure that V accepts
- If ϕ is false, the chance that V accepts is very small

Correctness

- If ϕ is true, a truthful Prover can ensure that V accepts
- If ϕ is false, the chance that V accepts is very small
- We use "Schwartz-Zippel" lemma. Let p be a non-zero multivariate polynomial $p(x_1, \dots, x_m)$ with degree $\leq d$ and S a finite set of integers. If a_1, \dots, a_m are chosen randomly independently and uniformly from S , then

$$\Pr[p(a_1, \dots, a_m) = 0] \leq \frac{d}{|S|}$$

- Prover sends wrong polynomial $p \neq h = \phi_{arith}$. Verifier chooses randomly $c \in GF(p)$ where $p \geq 2^n$. Furthermore we have $\deg(g - h) \leq 2n$.

Correctness

- If ϕ is true, a truthful Prover can ensure that V accepts
- If ϕ is false, the chance that V accepts is very small
- We use "Schwartz-Zippel" lemma. Let p be a non-zero multivariate polynomial $p(x_1, \dots, x_m)$ with degree $\leq d$ and S a finite set of integers. If a_1, \dots, a_m are chosen randomly independently and uniformly from S , then

$$\Pr[p(a_1, \dots, a_m) = 0] \leq \frac{d}{|S|}$$

- Prover sends wrong polynomial $p \neq h = \phi_{arith}$. Verifier chooses randomly $c \in GF(p)$ where $p \geq 2^n$. Furthermore we have $\deg(g - h) \leq 2n$. Then $\Pr[p(c) = h(c)] = \Pr[p(c) - h(c) = 0] \leq \frac{2n}{2^n}$.
-