

# Assignment 4

Maharaj Teertha Deb, 40227747

2024-02-27

## Section 3.1

### Problem 1 :


Consider the islands vector discussed in this section.

- (a) Compare the histograms that result when using breaks based on Sturges' and Scott's rules. Make this comparison on the log scale and on the original scale.

(a) **Answer:**

```
data(islands)


par(mfrow=c(2,2))
hist(islands, breaks = "Sturges", main = "Sturges")
hist(islands, breaks = "Scott", main = "Scott")
hist(log(islands), breaks = "Sturges", main = "Sturges on log scale")
hist(log(islands), breaks = "Scott", main = "Scott on log scale")
```



(b) Construct a normal QQ plot, and compare the result with the plots in Figure 3.13; which one is most similar, and what does this tell you about this data set?

(b) **Answer:**


```
qqnorm(islands)
qqline(islands, col = "red")
```



(c) Construct a boxplot for these data on the log scale as well as the original scale.

(c) Answer:


```
par(mfrow=c(1,2))
boxplot(islands, main = "Original scale")
boxplot(log(islands), main = "Log scale")
```



(d) Construct a dot chart of the areas. Is a log transformation needed here?

(d) Answer:


```
par(mfrow=c(1,2))
dotchart(islands,main="Original scale")
dotchart(log(islands),main="Log scale")
```



(e) Which form of graphic do you think is most appropriate for displaying these data?

(e) Answer:

```
dotchart(sort(log(islands)), main = "Log scale, sorted")
```



### Problem 3:


Consider the pressure data frame. There are two columns temperature and pressure

(a)

Construct a scatterplot with pressure on the vertical axis and temperature on the horizontal axis. Are the variables related linearly or nonlinearly?

(a) **Answer:**

```
plot(pressure ~ temperature, data = pressure) # Calculate correlation coefficient
```



```
correlation <- cor(pressure$temperature, pressure$pressure)

# Check for linearity
if(abs(correlation) < 0.005) {
  cat("The variables are related linearly (correlation =", correlation, ").\n")
} else {
  cat("The variables are related nonlinearly (correlation =", correlation, ").\n")
}
```

```
## The variables are related nonlinearly (correlation = 0.7577923 ).
```

(b)

The graph of the following function passes through the plotted points reasonably well:  $y = (0.168 + 0.007x)^{\frac{20}{3}}$ . The differences between the pressure values predicted by the curve and the observed pressure values are called residuals. Here is a way to calculate them:


```
residuals <- -with(pressure , pressure - (0.168 + 0.007 × temperature)20/3)
```

Construct a normal QQ plot of these residuals and decide whether they are normally distributed or whether they follow a skewed distribution.

(b) Answer:

```
# Calculate residuals
predicted_pressure <- (0.168 + 0.007 * pressure$temperature)^(20/3)
residuals <- pressure$pressure - predicted_pressure

# Construct a normal QQ plot
qqnorm(residuals)
qqline(residuals, col = 2)
```



```
# Assess normality
shapiro.test(residuals)
```

```
##
##  Shapiro-Wilk normality test
##
## data:  residuals
## W = 0.55893, p-value = 1.751e-06
```

(c)


Now, apply the power transformation  $y^{\frac{3}{20}}$  to the pressure data values. Plot these transformed values against temperature. Is a linear or nonlinear relationship evident now?



(c) Answer:

```
# Apply the power transformation  $y^{(3/20)}$  to pressure data values
transformed_pressure <- pressure$pressure^(3/20)

# Plot transformed values against temperature
plot(pressure$temperature, transformed_pressure,
     xlab = "Temperature", ylab = "Transformed Pressure",
     main = "Transformed Pressure vs Temperature")
```



```
# Calculate correlation coefficient between transformed pressure and temperature
correlation_transformed <- cor(pressure$temperature, transformed_pressure)

# Check for linearity
if(abs(correlation_transformed) < 0.005) {
  cat("The relationship between transformed pressure and temperature appears to
      be linear (correlation =", correlation_transformed, ").\n")
} else {
  cat("The relationship between transformed pressure and temperature appears to
      be nonlinear (correlation =", correlation_transformed, ").\n")
}
```

```
## The relationship between transformed pressure and temperature appears to
##      be nonlinear (correlation = 0.9984827 ).
```

(d)

Calculate residuals for the difference between transformed pressure values and those predicted by the straight line. Obtain a normal QQ plot, and decide whether the residuals follow a normal distribution or not.


(d) Answer:

```
# Fit a linear regression model to transformed pressure values and temperature
lm_model_transformed <- lm(transformed_pressure ~ pressure$temperature)

# Predict transformed pressure values using the linear model
predicted_transformed_pressure <- predict(lm_model_transformed)

# Calculate residuals
residuals_transformed <- transformed_pressure - predicted_transformed_pressure

# Construct a normal QQ plot of residuals
qqnorm(residuals_transformed)
qqline(residuals_transformed, col = 2)
```



```
# Assess normality
shapiro.test(residuals_transformed)
```

```
##
```

```
## Shapiro-Wilk normality test
##
## data: residuals_transformed
## W = 0.92153, p-value = 0.1208
```