

# Assignment-2

U.S.M.M TEJA  
cs21btech11059

## Question 1-(xi)[ICSE 2017]:

**Problem 1.** if  $a+ib = \frac{x+iy}{x-iy}$   
prove that  $a^2 + b^2 = 1$  and  $\frac{b}{a} = \frac{2xy}{x^2 - y^2}$

**Solution:**

$$a + ib = \frac{x + iy}{x - iy} \quad (1)$$

let r be the magnitude of the given complex number a+ib

$$\text{so } r = \sqrt{a^2 + b^2}$$

$$\tan \theta = \frac{b}{a} \quad (2)$$

$$x + iy = r(\cos \alpha + i \sin \alpha) \quad (3)$$

$$\tan \alpha = \frac{y}{x} \quad (4)$$

$$r(\cos \theta + i \sin \theta) = \frac{\cos \alpha + i \sin \alpha}{\cos \alpha - i \sin \alpha}$$

multiplying denominator on both numerator and denominator we get

$$r(\cos \theta + i \sin \theta) = \cos^2 \alpha - \sin^2 \alpha + 2i \sin \alpha \cdot \cos \alpha$$

$$r(\cos \theta + i \sin \theta) = \cos 2\alpha + i \sin 2\alpha \quad (5)$$

$$\implies r = 1 \quad (6)$$

magnitude of complex number on RHS is 1 so definitely magnitude on LHS is 1

so comparing both LHS and RHS  $\theta = 2\alpha$

applying tan on both sides, we get  $\tan \theta = \tan 2\alpha$

$$\frac{b}{a} = \frac{2 \tan \alpha}{1 - \tan^2 \alpha} \quad (7)$$

$$\frac{b}{a} = \frac{2y/x}{1 - y^2/x^2} \quad (8)$$

$$\implies \frac{b}{a} = \frac{2xy}{x^2 - y^2} \quad (9)$$