




# Response of Warm-Season Vegetables to FUSN™ in the Desert Southwest

Dr. Charlie Sanchez, University of Arizona

2015



## Introduction

Due to threats of terrorism and accidents, the fertilizer ammonium nitrate is facing increasing scrutiny and regulation. FUSN™ is a Honeywell proprietary technology currently marketed by J.R. Simplot Company whereby ammonium nitrate is fused with ammonium sulfate. This product is easier to handle and store and has a lower detonation potential than ammonium nitrate. FUSN is classified as a non-oxidizer and non-explosive by the U.S. Department of Transportation. The objective of these experiments was to evaluate FUSN and a nitrogen (N) fertilizer source for warm-season vegetables.

## Materials and Methods

Studies were conducted during the spring and summer of 2015 to evaluate the response of sweet corn (*Zea mays*), cantaloupes (*Cucumis melo*), and chili peppers (*Capsicum annuum*) to FUSN fertilizer. The soil used in all experiments was a Superstition loamy sand (sandy, mixed, hyperthermic Typic Calciorthid). Sweet corn (cv. BS5082) was seeded on 42-inch beds and cantaloupes (cv. Oro Rico) were seeded on 84-inch beds on March 31, 2015. Chili pepper (cv. Big Jim) transplants previously grown in the greenhouse were set out in 42-inch beds using a mechanical transplanter on April 7, 2015. The N rates were 0, 100, 200, and 300 lbs N/ac as FUSN and urea. We applied 50% of the N at planting and the remaining 50% midseason. The plots were irrigated by sub-surface drip.

Sweet corn yields were collected June 27, 2015, by harvesting all marketable ears. Cantaloupe yields were collected July 2, 2015, by harvesting all marketable fruit. Chili pepper yields were collected June 22 and June 28, 2015, by harvesting all mature fruit. All marketable yields were determined after grading.

## Results

Sweet corn yields were not high due to heat stress and the impact on pollination. Soil tests showed nitrate-N increased to N rate and FUSN had soil nitrate-N slightly higher than urea (Table 1). There were no significant differences in soil ammonium-N, although the urea plots had higher residual ammonium. There were few consistent or meaningful differences in leaf N concentration. Yields significantly increased by N rate. Although not statistically significant at  $P < .05$ , there was a trend for FUSN to produce higher yields compared to urea that would be significant at  $P < .10$ .

In the melon experiment, soil nitrate-N levels increased with N rate (Table 2). Yields were also significantly increased by N rate. Although FUSN yields averaged 18% greater than those with urea, the differences were not statistically significant. We had bird damage in these plots. We tried to select harvest areas of uniform stand but we did not entirely overcome the variation.

Although pepper leaf tissue N was lower for the control compared to all treatments receiving N fertilizer, the effect was not statistically significant (Table 3). Chili pepper yields were increased by N rate during the first harvest but yields seemed to compensate during the second harvest and total yields were similar regardless of N rate or source at  $P < .05$ . For the final harvest, yield effects to rate and source would be significant at  $P < .10$ .

| Treatment | N Rate<br>(kg/ha) | N<br>Source    | Soil NH <sub>4</sub> -N<br>(mg/kg) | Soil NO <sub>3</sub> -N<br>(mg/kg) | Leaf N<br>(%) | Yield<br>(Mt/ha) |
|-----------|-------------------|----------------|------------------------------------|------------------------------------|---------------|------------------|
| 1         | 0                 | -              | 3.2                                | 7.0                                | 1.1           | 0.2              |
| 2         | 100               | Urea           | 4.1                                | 14.3                               | 1.1           | 1.4              |
| 3         | 200               | Urea           | 13.7                               | 30.0                               | 1.0           | 0.9              |
| 4         | 300               | Urea           | 10.2                               | 34.0                               | 1.1           | 2.1              |
| 5         | 100               | F $\bar{U}$ SN | 4.8                                | 38.6                               | 1.0           | 1.1              |
| 6         | 200               | F $\bar{U}$ SN | 7.4                                | 42.8                               | 1.0           | 1.8              |
| 7         | 300               | F $\bar{U}$ SN | 4.1                                | 37.7                               | 1.2           | 4.3              |
| Stat.     | N Rate            |                | NS                                 | L**                                | Q*            | L**              |
|           | N Source          |                | NS                                 | *                                  | NS            | NS               |

**Table 1.** Sweet corn response to F $\bar{U}$ SN and urea during spring and summer 2015  
(\* = P < .10, \*\* = P < .05).

| Treatment | N Rate<br>(kg/ha) | N<br>Source    | Soil<br>NH <sub>4</sub> -N<br>(mg/kg) | Soil<br>NO <sub>3</sub> -N<br>(mg/kg) | Midrib nitrate-<br>N (mg/kg) | Yield<br>(Mt/ha) |
|-----------|-------------------|----------------|---------------------------------------|---------------------------------------|------------------------------|------------------|
| 1         | 0                 | -              | 3.8                                   | 35.5                                  | 30207                        | 9.7              |
| 2         | 100               | Urea           | 4.2                                   | 32.7                                  | 36593                        | 10.2             |
| 3         | 200               | Urea           | 4.3                                   | 37.7                                  | 34363                        | 8.2              |
| 4         | 300               | Urea           | 5.0                                   | 43.8                                  | 45735                        | 8.1              |
| 5         | 100               | F $\bar{U}$ SN | 4.1                                   | 32.3                                  | 44188                        | 9.1              |
| 6         | 200               | F $\bar{U}$ SN | 4.6                                   | 30.3                                  | 26365                        | 11.2             |
| 7         | 300               | F $\bar{U}$ SN | 1.0                                   | 50.0                                  | 28967                        | 10.7             |
| Stat.     | N Rate            |                | NS                                    | L*Q*                                  | NS                           | NS               |
|           | N Source          |                | NS                                    | NS                                    | NS                           | NS               |

**Table 2.** Cantaloupe response to F $\bar{U}$ SN and urea during spring and summer 2015 (\* = P < .10).

|       | N Rate<br>(kg/ha) | N<br>Source    | Soil<br>NH <sub>4</sub> -N<br>(mg/kg) | Soil<br>NO <sub>3</sub> -N<br>(mg/kg) | Leaf N<br>(%) | Yield (Mt/ha) |      |       |
|-------|-------------------|----------------|---------------------------------------|---------------------------------------|---------------|---------------|------|-------|
|       |                   |                |                                       |                                       |               | 6/22          | 6/28 | Total |
| 1     | 0                 | -              | 3.8                                   | 2.4                                   | 2.18          | 3.1           | 3.3  | 6.3   |
| 2     | 100               | Urea           | 3.7                                   | 3.2                                   | 2.37          | 2.7           | 4.0  | 6.7   |
| 3     | 200               | Urea           | 8.7                                   | 15.1                                  | 2.41          | 5.3           | 2.6  | 7.9   |
| 4     | 300               | Urea           | 4.1                                   | 5.3                                   | 2.32          | 3.2           | 3.3  | 6.5   |
| 5     | 100               | F $\bar{U}$ SN | 3.8                                   | 3.8                                   | 2.25          | 4.3           | 3.4  | 7.7   |
| 6     | 200               | F $\bar{U}$ SN | 4.1                                   | 3.7                                   | 2.25          | 4.3           | 2.9  | 7.3   |
| 7     | 300               | F $\bar{U}$ SN | 4.4                                   | 4.9                                   | 2.25          | 4.6           | 3.5  | 8.1   |
| Stat. | N Rate            |                | NS                                    | NS                                    | NS            | L*            | NS   | NS    |
|       | N<br>Source       |                | NS                                    | NS                                    | NS            | NS            | NS   | NS    |

**Table 3.** Chili pepper response to F $\bar{U}$ SN and urea during spring and summer 2015 (\* = P < .10).



Simplot® is a registered trademark of J.R. Simplot Company. FUSN™ is a trademark of J.R. Simplot Company. 2015-15-0616