



**KOLEJ MATRIKULASI NEGERI SEMBILAN**

***PROGRAM SCORE***  
**SET 1**

**PROGRAM KECEMERLANGAN AKADEMIK (PKA)**  
**KOLEJ MATRIKULASI NEGERI SEMBILAN**  
**MINGGU KE 17**

---

---

**FIZIK (SP015)**  
**2 jam**

---

---

NO	MARKS
1	/2
2	/10
3	/13
4	/8
5	/5
6	/23
7	/8
8	/11
TOTAL	/80

## LIST OF SELECTED CONSTANT VALUES

Speed of light in a vacuum	$c$	$= 3.00 \times 10^8 \text{ m s}^{-1}$
Permeability constant	$\mu_0$	$= 4\pi \times 10^{-7} \text{ H m}^{-1}$
Permittivity constant	$\epsilon_0$	$= 8.85 \times 10^{-12} \text{ F m}^{-1}$
Elementary charge	$e$	$= 1.60 \times 10^{-19} \text{ C}$
Planck's constant	$h$	$= 6.63 \times 10^{-34} \text{ J s}$
Electron mass	$m_e$	$= 9.11 \times 10^{-31} \text{ kg}$ $= 5.49 \times 10^{-4} \text{ u}$
Neutron mass	$m_n$	$= 1.674 \times 10^{-27} \text{ kg}$ $= 1.008665 \text{ u}$
Proton mass	$m_p$	$= 1.672 \times 10^{-27} \text{ kg}$ $= 1.007277 \text{ u}$
Deuteron mass	$m_d$	$= 3.34 \times 10^{-27} \text{ kg}$ $= 2.014102 \text{ u}$
Universal gas constant	$R$	$= 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Rydberg's constant	$R_H$	$= 1.097 \times 10^7 \text{ m}^{-1}$
Avogadro constant	$N_A$	$= 6.02 \times 10^{23} \text{ mol}^{-1}$
Boltzmann's constant	$k$	$= 1.38 \times 10^{-23} \text{ J K}^{-1}$
Gravitational constant	$G$	$= 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
Free-fall acceleration	$g$	$= 9.81 \text{ m s}^{-2}$
Atomic mass constant	$1 \text{ u}$	$= 1.66 \times 10^{-27} \text{ kg}$ $= 931.5 \frac{\text{MeV}}{c^2}$
Electron Volt	$1 \text{ eV}$	$= 1.6 \times 10^{-19} \text{ J}$
Constant of proportionality for Coulomb's law, $k = \frac{1}{4\pi\epsilon_0}$		$= 9.0 \times 10^9 \text{ m}^2 \text{ C}^{-2}$
Atmospheric Pressure	$1 \text{ atm}$	$= 1.013 \times 10^5 \text{ Pa}$
Density of water	$\rho_w$	$= 1000 \text{ kg m}^{-3}$

**LIST OF SELECTED FORMULAE**  
**SENARAI RUMUS TERPILIH**

- |  |  |
|--|--|
| 1. $v = u + at$                                    | 19. $v = r\omega$                                      |
| 2. $s = ut + \frac{1}{2}at^2$                      | 20. $a_t = r\alpha$                                    |
| 3. $v^2 = u^2 + 2as$                               | 21. $\omega = \omega_o + \alpha t$                     |
| 4. $s = \frac{1}{2}(u + v)t$                       | 22. $\theta = \omega_o t + \frac{1}{2}\alpha t^2$      |
| 5. $p = mv$  | 23. $\theta = \frac{1}{2}(\omega_o + \omega)t$         |
| 6. $J = F\Delta t$                                 | 24. $\omega^2 = \omega_o^2 + 2\alpha\theta$            |
| 7. $J = \Delta p = mv - mu$                        | 25. $\tau = rF \sin \theta$                            |
| 8. $f = \mu N$                                     | 26. $I = \sum mr^2$                                    |
| 9. $W = \vec{F} \cdot \vec{s} = F_s \cos \theta$   | 27. $I_{\text{solid sphere}} = \frac{2}{5}MR^2$        |
| 10. $K = \frac{1}{2}mv^2$                          | 28. $I_{\text{solid cylinder/disc}} = \frac{1}{2}MR^2$ |
| 11. $U = mgh$                                      | 29. $I_{\text{ring}} = MR^2$                           |
| 12. $U_s = \frac{1}{2}kx^2 = \frac{1}{2}Fx$        | 30. $I_{\text{rod}} = \frac{1}{12}ML^2$                |
| 13. $W = \Delta K$                                 | 31. $\sum \tau = I\alpha$                              |
| 14. $P_{\text{av}} = \frac{\Delta W}{\Delta t}$    |  |
| 15. $P = \vec{F} \cdot \vec{v} = Fv \cos \theta$   |  |
| 16. $a_c = \frac{v^2}{r} = r\omega^2 = v\omega$    |  |
| 17. $F_c = \frac{mv^2}{r} = mr\omega^2 = mv\omega$ |  |
| 18. $s = r\theta$                                  |  |

**LIST OF SELECTED FORMULAE**  
**SENARAI RUMUS TERPILIH**

$$32. \quad L = I\omega$$

$$33. \quad y = A \sin \omega t$$

$$34. \quad v = \omega A \cos \omega t = \pm \omega \sqrt{A^2 - y^2}$$

$$35. \quad a = -\omega^2 A \sin \omega t = -\omega^2 y$$

$$36. \quad K = \frac{1}{2} m \omega^2 (A^2 - y^2)$$

$$37. \quad U = \frac{1}{2} m \omega^2 y^2$$

$$38. \quad E = \frac{1}{2} m \omega^2 A^2$$

$$39. \quad \omega = \frac{2\pi}{T} = 2\pi f$$

$$40. \quad T = 2\pi \sqrt{\frac{l}{g}}$$

$$41. \quad T = 2\pi \sqrt{\frac{m}{k}}$$

$$42. \quad k = \frac{2\pi}{\lambda}$$

$$43. \quad v = f\lambda$$

$$44. \quad y(x, t) = A \sin(\omega t \pm kx)$$

$$45. \quad v_y = A\omega \cos(\omega t \pm kx)$$

$$46. \quad y = 2A \cos kx \sin \omega t$$

$$47. \quad f_n = \frac{nv}{2L}$$

$$48. \quad f_n = \frac{n}{2L} \sqrt{\frac{T}{\mu}}$$

$$49. \quad f_n = \frac{nv}{4L}$$

$$50. \quad v = \sqrt{\frac{T}{\mu}}$$

$$51. \quad \mu = \frac{m}{L}$$

$$52. \quad f_a = \left( \frac{v \pm v_o}{v \mp v_s} \right) f$$

$$53. \quad \sigma = \frac{F}{A}$$

$$54. \quad \varepsilon = \frac{\Delta L}{L_o}$$

$$55. \quad Y = \frac{\sigma}{\varepsilon}$$

$$56. \quad U = \frac{1}{2} F \Delta L$$

$$57. \quad \frac{U}{V} = \frac{1}{2} \sigma \varepsilon$$

$$58. \quad \frac{Q}{t} = -kA \left( \frac{\Delta T}{L} \right)$$

$$59. \quad \Delta L = \alpha L_o \Delta T$$

$$60. \quad \Delta A = \beta A_o \Delta T$$

$$61. \quad \Delta V = \gamma V_o \Delta T$$

$$62. \quad \beta = 2\alpha$$

**LIST OF SELECTED FORMULAE**  
**SENARAI RUMUS TERPILIH**

$$63. \quad \gamma = 3\alpha \qquad 69. \quad K_{\text{tr}} = \frac{3}{2} \left( \frac{R}{N_A} \right) T = \frac{3}{2} kT$$

$$64. \quad n = \frac{m}{M} = \frac{N}{N_A} \qquad 70. \quad U = \frac{1}{2} f N k T = \frac{1}{2} f n R T$$

$$65. \quad v_{\text{rms}} = \sqrt{\langle v^2 \rangle} \qquad 71. \quad \Delta U = Q - W$$

$$66. \quad v_{\text{rms}} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}} \qquad 72. \quad W = nRT \ln \frac{V_f}{V_i} = nRT \ln \frac{P_i}{P_f}$$


$$67. \quad PV = \frac{1}{3} N m v_{\text{rms}}^2 \qquad 73. \quad W = \int P dV = P(V_f - V_i)$$

$$68. \quad P = \frac{1}{3} \rho v_{\text{rms}}^2 \qquad 74. \quad W = \int P dV = 0$$

Answer **ALL** questions.

- 1 The Young's Modulus of a material is given by equation  $Y = \frac{\sigma}{\epsilon}$  where  $\sigma$  and  $\epsilon$  are refer to the tensile stress and tensile strain respectively. What is the dimension of Young's Modulus?  
[2 marks]
- 2 (a) The speed of a car travelling along a straight road decrease uniformly from  $16 \text{ m s}^{-1}$  to  $12 \text{ ms}^{-1}$  over 80 m. Calculate  
(i) the deceleration of the car  
(ii) the time taken for the speed to decrease from  $16 \text{ ms}^{-1}$  to  $12 \text{ ms}^{-1}$   
[3 marks]
- (b) An object is thrown horizontally from the edge of a table with an initial velocity of  $3 \text{ m s}^{-1}$ . If the height of the table is 2.0 m, calculate  
(i) the final velocity of object before it reach the floor.  
(ii) the horizontal displacement of the object.  
[7 marks]
- 3 (a) A fisherman on a stationary boat jumps off onto a jetty with a velocity of  $1.5 \text{ m s}^{-1}$  causing the boat to moves backwards.  
(i) If the mass of the fisherman and boat are 65 kg and 450 kg, determine the velocity of the boat.  
(ii) If the fisherman's feet are in contact with the jetty for 10 ms, determine the magnitude of the average force exerted on his feet.  
[6 marks]

(b)



**FIGURE 1**

**FIGURE 1** shows an object of mass 2.0 kg placed on a rough plane inclined at  $30^\circ$  with the horizontal. The coefficient of kinetic friction between the object and the plane surface is 0.25. A constant horizontal force,  $F = 50 \text{ N}$  acts on the object and pushes it along the inclined plane with acceleration,  $a$ .

- (i) Sketch a free body diagram showing all the forces acting on the object.  
(ii) Calculate the acceleration of the object.

[7 marks]

4

(a)




FIGURE 2

An object of mass 2.0 kg travels along horizontal floor under the action of force,  $F$ . **FIGURE 2** shows the graph of force,  $F$  against displacement,  $s$ . The speed of the object at  $s = 0$  is  $10 \text{ m s}^{-1}$ . Determine the kinetic energy of an object at  $s = 10 \text{ m}$

[3 marks]

(b)




FIGURE 3

An object of mass 2.0 kg is placed 30 cm directly above the top end of a vertical spring as shown in **FIGURE 3**. It is then released from that height. The spring constant  $k = 20 \text{ N m}^{-1}$ .

- Calculate the speed of the object just before it strikes the spring.
- Determine the maximum compression,  $x$ .

[5 marks]

5

A 900 kg car moving at  $10 \text{ m s}^{-1}$  takes a turn around a circle with a radius of 25.0 m. Determine


- the angular velocity of the car.
- the period of motion of the car.
- the net force acting upon the car.

[5 marks]

- 6 (a) A mass is suspended vertically on the end of spring and undergoes simple harmonic motion of frequency 4 Hz and amplitude 2 cm. At displacement,  $x = 1$  cm, determine the
- velocity of the mass.
  - acceleration of the mass.

[4 marks]

(b)

**FIGURE 4**

**FIGURE 4** shows a displacement-time graph of a simple harmonic motion for an object with mass 25 g. Determine

- the angular frequency.
- the equation for the simple harmonic motion.
- the total energy.

[6 marks]

- (c) The equation of progressive wave is given as

$$y = 6 \sin 2\pi(2t - x)$$

where  $x$  and  $y$  are in meter and  $t$  in second. Determine the

- direction of the wave propagation.
- wavelength,  $\lambda$
- frequency,  $f$
- wave speed.
- particle velocity at  $x = 2$  m and  $t = 0.1$  s.

[6 marks]

- (d) Two waves in a long string are given by

$$y_1 = 8 \sin(2\pi t - 2x) \quad y_2 = 8 \sin(2\pi t + 2x)$$

where  $y_1$ ,  $y_2$  and  $x$  in meter and  $t$  in second.

- Write an expression for the new wave when both waves are superimposed.
- Determine the maximum amplitude,  $A$  for stationary wave.

[2 marks]



- (e) A 50 cm closed pipe makes a humming sound when the wind blows across the opened end. If the speed of sound is  $340 \text{ m s}^{-1}$ , calculate
- second overtone of the pipe.
  - its fundamental frequency if the length of pipe is halved.

[2 marks]


- (f) A fire engine is coming towards a station emitting siren at frequency 800 Hz. What is the possible frequency heard by a boy sitting in the station if the incoming fire engine is moving at  $65 \text{ m s}^{-1}$ ? The speed of sound is  $340 \text{ m s}^{-1}$ .

[3 marks]

- 7 (a) An aluminium wire with diameter 1.5 mm is suspended from a rigid support with a load of 150 kg attached to its lower end. If the wire becomes 5.5 m long, what is the
- initial length of the wire.
  - extension of the wire.
- (Young's modulus of aluminium is  $7 \times 10^{10} \text{ Nm}^{-2}$ )

[3 marks]

- (b)

**FIGURE 5**

Two bars of the same diameter but different materials are joined end to end as shown in **FIGURE 5**. One end of the bars is kept at  $80^\circ\text{C}$  while the other end is at  $5^\circ\text{C}$ . Determine the steady state temperature at the interface of the bars if the length of the first bar is same as the second bar.

(Given thermal conductivity of the first bar is  $240 \text{ W m}^{-1}^\circ\text{C}^{-1}$  and the second one is  $110 \text{ W m}^{-1}^\circ\text{C}^{-1}$ )

[3 marks]

- (c) A block of gold has dimensions of  $12 \text{ cm} \times 10 \text{ cm} \times 8 \text{ cm}$  at temperature of  $30^\circ\text{C}$ . If the temperature increases to  $120^\circ\text{C}$ , determine the change in the volume of the gold block. The coefficient of linear expansion for gold =  $1.4 \times 10^{-6}^\circ\text{C}^{-1}$ .

[2 marks]


- 8 (a) A tank contains  $1 \text{ m}^3$  of helium at a pressure of  $1.3 \times 10^5 \text{ Pa}$  and temperature of  $350 \text{ K}$ . The temperature of the gas increases from  $350 \text{ K}$  to  $370 \text{ K}$ . What is the change in the internal energy of the gas?

[4 marks]

- (b) The rms speed of hydrogen molecules at a particular temperature is  $1330 \text{ m s}^{-1}$ . What is the rms speed of nitrogen molecules at the same temperature? Molar mass of hydrogen =  $2 \text{ g mol}^{-1}$  and molar mass of nitrogen =  $28 \text{ g mol}^{-1}$ .

[2 marks]

(c)

**FIGURE 6**

An ideal gas has undergone two processes AB and BC as shown in **FIGURE 6**. 4000 J of heat is dissipated from the gas in the process BC. If  $P_0$  is  $1.0 \times 10^5$  Pa and  $V_0$  is  $0.02 \text{ m}^3$ , calculate the

- (i) work done for AB and BC.
- (ii) changes in internal energy for BC.

[5 marks]