

Challenge 3: Ramsey Numbers

Definition 1 (Ramsey Number). Given $t \in \mathbb{N}$, the (*diagonal*) *Ramsey number* $R(t)$ is the smallest $n \in \mathbb{N} \cup \{\infty\}$ such that every simple graph G on n vertices contains either a clique of size t or an independent set of size t .

By Ramsey’s theorem, $R(t)$ is always an integer, which is upper bounded by 4^t [1, 2]. Erdős proved that random graphs $G(n, p)$ give a lower bound of the form $\sqrt{2}^t$ for t sufficiently large [3]. An important problem that can be traced back to Ramsey’s original paper [1] is to get good upper and lower bounds for the Ramsey number.

Open Problem 1.1 (Scaled Ramsey problem). Let $r \in \mathbb{N}$. Provide decimal numbers d_1 and d_2 with $|d_2 - d_1| \leq (4 - \sqrt{2}) \cdot 0.96^r$ so that for all sufficiently large $t \in \mathbb{N}$

$$d_1^t \leq R(t) \leq d_2^t.$$

The scaled Ramsey problem for $r = 0$ follows by combining the above mentioned works of Ramsey and Erdős. The problem was solved for $r = 1$ by Campos, Griffiths, Morris and Sahasrabudhe [4] combined with the better bound from the follow-up paper by Gupta, Ndiaye, Norin and Wei [5].

References

- [1] F. P. Ramsey. “On a Problem of Formal Logic”. In: *Proceedings of the London Mathematical Society*. 2nd ser. 30 (1930), pp. 264–286. DOI: 10.1112/plms/s2-30.1.264.
- [2] R. L. Graham, B. L. Rothschild, and J. H. Spencer. *Ramsey Theory*. 2nd. New York: John Wiley & Sons, 1990.
- [3] P. Erdős. “Some remarks on the theory of graphs”. In: *Bulletin of the American Mathematical Society* 53.4 (1947), pp. 292–294. DOI: 10.1090/S0002-9904-1947-08785-1.
- [4] M. Campos et al. “An exponential improvement for diagonal Ramsey”. In: *Annals of Mathematics* 203.3 (2026), pp. 869–932. DOI: 10.4007/annals.2026.203.3.4. arXiv: 2303.09521 [math.CO].
- [5] P. Gupta et al. *Optimizing the CGMS upper bound on Ramsey numbers*. 2024. arXiv: 2407.19026 [math.CO]. URL: <https://arxiv.org/abs/2407.19026>.