

Geometric Class Field Theory

Assaf Marzan

July 9, 2026

Abstract

We calculate the ramification of the symmetric product $\mathcal{F}^{[\deg \mathfrak{m}]}$ of a local system $\mathcal{F}$ on a curve $C \setminus \mathfrak{m}$. Assuming the ramification of $\mathcal{F}$ is bounded by $\mathfrak{m}$, we find that the symmetric product $\mathcal{F}^{[\deg \mathfrak{m}]}$ is at most tamely ramified at the generic point of the exceptional divisor $E_{\mathfrak{m}}$ of the blowup of $C^{(\deg \mathfrak{m})}$ at $\mathfrak{m}$. As a primary application, we utilize this result to prove Geometric Class Field Theory. Our approach builds upon the geometric framework for the unramified case originally established by Deligne for the rank-one Langlands correspondence, following the subsequent extensions to the ramified case developed by Takeuchi and Guignard.

Contents

1	Introduction	4
2	Preliminaries	6
2.1	Scheme Quotients in general	6
2.2	Torsors	6
2.3	Torsors over the Étale's Sites	8
2.4	Symmetric Powers of Schemes and Torsors	10
2.5	Algebraic Preliminaries on Ramification	16
2.6	Algebraic Geometry	19
3	Ramification After Blowup Is Tame	21
3.1	Ramification of G -Torsors	22
3.2	Proof of Theorem 3	26
3.3	Extending To Product of Blowups	35
4	Geometric Class Field Theory	38
4.1	Étale Fundamental Groups and Tame Fundamental Groups	38
4.2	Generalized Picard Scheme	38
4.3	The Abel-Jacobi Morphism and its Fibers	40
4.4	Compactification of Blowup of Symmetric Powers of a Curve	41
4.5	Proof of Theorem 54	42
4.5.1	Reduction Lemmas	42
4.6	Proof of Theorem 54	44
5	Proof of Theorem 2	45
6	Setup and notation	46
7	The master diagram	46
8	Unramifiedness of $\mathcal{P}^{[d]G}$ at η_C	47
8.1	Proof Attempts	47
8.1.1	Attempt 1	47
9	A Wrong Path	49
9.1	Setup and notation	49

9.2	The master diagram	50
9.3	Unramifiedness of $\mathcal{P}^{[d]G}$ at η_C	51
9.4	Difficulties in Proving the Theorem	51

1 Introduction

In this thesis, we give an elementary proof of a certain important geometric theorem occurring in Deligne's approach to geometric class field theory. We (usually) work over a perfect field k , C is a projective smooth geometrically connected curve over k , with genus g . One of the main geometric ingredients in the approach, is showing why a local system $\mathcal{F}$ with ramification bounded by a modulus $\mathfrak{m} = \sum n_i P_i$ on $U = C \setminus \mathfrak{m}$ descends via the Abel-Jacobi $\Phi : U \rightarrow \text{Pic}_{C,\mathfrak{m}}$ to $\text{Pic}_{C,\mathfrak{m}}$. The approach, innovated by Deligne, relies on analyzing the symmetric powers $\mathcal{F}^{[d]}$ of $\mathcal{F}$ on the symmetric powers $U^{(d)}$ of U , and showing that for sufficiently large d , $\mathcal{F}^{[d]}$ descends to $\text{Pic}_{C,\mathfrak{m}}^d$ via the degree d Abel-Jacobi map $\Phi_d : U^{(d)} \rightarrow \text{Pic}_{C,\mathfrak{m}}^d$. The geometric-fibers of Φ_d (for $d \geq \deg \mathfrak{m} + 2g - 1$) over any point are isomorphic to

$$\begin{cases} \mathbb{A}_{k^{sep}}^{d-\deg \mathfrak{m}-g+1} & \text{if } \mathfrak{m} > 0 \\ \mathbb{P}_{k^{sep}}^{d-g} & \text{if } \mathfrak{m} = 0 \end{cases}$$

Where g is the genus of the curve C . The unramified case ($\mathfrak{m} = 0$) is relatively simple, as the Abel-Jacobi map is proper, surjective with geometrically connected fibers, which follows from the fact that it is a fibration in projective spaces. Thus, by using the homotopy exact sequence for the étale fundamental group, one gets an isomorphism between the étale fundamental group of $U^{(d)} (= C^{(d)})$ and that of $\text{Pic}_{C,\mathfrak{m}}^d (= \text{Pic}_C^d)$.

The ramified case ($\mathfrak{m} > 0$) is more subtle, as the Abel-Jacobi map is not proper anymore, and one needs to analyze the ramification of $\mathcal{F}^{[d]}$ "along the boundary" of $U^{(d)}$ in $C^{(d)}$.

Previous work has generalized Deligne's approach to the ramified case, most notably by Guignard [Gui19] and Takeuchi [Tak19]. Their approaches differ. To descend, Guignard proves that the restriction of $\mathcal{F}^{[d]}$ to any line in the fiber of the degree d Abel-Jacobi map is a constant étale sheaf. He achieves this by demonstrating that the restriction is at most tamely ramified and invoking the triviality of the tame fundamental group of $\mathbb{A}_k^1$. His analysis relies on local geometric class field theory. It is also worth noting that Guignard's method generalizes to relative curves over arbitrary base schemes. Takeuchi, on the other hand, constructs a compactification of $U^{(d)}$ by blowing up $C^{(d)}$ along certain well-chosen centers. This compactification, denoted by $\tilde{C}_{\mathfrak{m}}^{(d)}$, has $U^{(d)}$ as an open subscheme with a codimension 1 closed subscheme H as complement. He then shows that the Abel-Jacobi map extends to a proper morphism from $\tilde{C}_{\mathfrak{m}}^{(d)}$ to $\text{Pic}_{C,\mathfrak{m}}^d$, which is a fibration in projective spaces. Thus, by the homotopy exact sequence for the étale fundamental group, one gets an isomorphism between the étale fundamental group of $\tilde{C}_{\mathfrak{m}}^{(d)}$ and that of $\text{Pic}_{C,\mathfrak{m}}^d$. To conclude the descent, Takeuchi analyzes the ramification of $\mathcal{F}^{[d]}$ along the boundary H of $\tilde{C}_{\mathfrak{m}}^{(d)}$, showing that it is tamely ramified there, which suffices. His methods relies on the theory of Witt vectors and refined Swan conductors [incorporate [Rab14] here].

For an account of these approaches, see [Gui19] and [Tak19]. For a full approach following Deligne's method in the unramified case, and the tamely ramified case see [Ten15], and [Töt11].

In this thesis, we calculate the ramification of $\mathcal{F}^{[d]}$ directly to show that it is at most tame at the generic point of H , avoiding the use of Swan conductors. To prove Geometric Class Field Theory, we utilize Takeuchi's construction of the compactification $\tilde{C}_{\mathfrak{m}}^{(d)}$ of $U^{(d)}$ via the blowing up of $C^{(d)}$.

In the rest of the introduction, we state the main theorem of geometric class field theory Theorem 1, and its reductions to Theorem 2 and Theorem 3, which we prove in this thesis.

Let k be a perfect field, and let C be a projective smooth geometrically connected curve over k ,

with genus g . Geometric class field theory gives a geometric description of abelian coverings of C by relating it to isogenies of the generalized Picard schemes.

Fix a modulus $\mathfrak{m}$, i.e. an effective Cartier divisor of C and let U be its complement in C . The pairs $(\mathcal{L}, \alpha)$, where $\mathcal{L}$ is an invertible $\mathcal{O}_C$ -module and α is a rigidification of $\mathcal{L}$ along $\mathfrak{m}$, are parametrized by a k -group scheme $\text{Pic}_{C, \mathfrak{m}}$, called the rigidified Picard scheme. The Abel-Jacobi morphism

$$\Phi : U \rightarrow \text{Pic}_{C, \mathfrak{m}}$$

is the morphism which sends a section x of U to the pair $(\mathcal{O}(x), 1)$. The fundamental result of GCFT can be formulated as:

Theorem 1 (Geometric Class Field Theory). *Let Λ be a finite ring of cardinality invertible in k , and let $\mathcal{F}$ be an étale sheaf of Λ -modules, locally free of rank 1 on U , with ramification bounded by $\mathfrak{m}$. Then, there exists a unique (up to isomorphism) multiplicative¹ étale sheaf of Λ -modules $\mathcal{G}$ on $\text{Pic}_{C, \mathfrak{m}}$, locally free of rank 1, such that the pullback of $\mathcal{G}$ by Φ is isomorphic to $\mathcal{F}$.*

Let d be a positive integer. We denote by $U^{(d)}$ the d -th symmetric power of U over k . For an étale sheaf $\mathcal{F}$ on $U_{\text{ét}}$, we denote by $\mathcal{F}^{[d]}$ the d -th symmetric power of $\mathcal{F}$ on $U^{(d)}$. The degree d Abel-Jacobi morphism is defined as the map

$$\Phi_d : U^{(d)} \rightarrow \text{Pic}_{C, \mathfrak{m}}^d$$

which sends a section $x_1 + \cdots + x_d$ of $U^{(d)}$ to the pair $(\mathcal{O}(x_1 + \cdots + x_d), 1)$.

A method of descent shows that to prove [Theorem 1](#), it suffices to prove the following reduced version²:

Theorem 2. *Let Λ be a finite ring of cardinality invertible in k , and let $\mathcal{F}$ be an étale sheaf of Λ -modules, locally free of rank 1 on $U_{\text{ét}}$, with ramification bounded by $\mathfrak{m}$. Then, for sufficiently large integer d , there exists a unique (up to isomorphism) étale sheaf of Λ -modules $\mathcal{G}_d$ on $\text{Pic}_{C, \mathfrak{m}}^d$, locally free of rank 1, such that the pullback of $\mathcal{G}_d$ by Φ_d is isomorphic to $\mathcal{F}^{[d]}$.*

To prove [Theorem 2](#) we follow the work of [\[Tak19\]](#) (and similiary done in [\[T6t11\]](#)), analyzing the ramification of $\mathcal{F}^{[d]}$ after blowing up $C^{(d)}$. We analyze this ramification using elementary methods. We prove the following Theorem:

Theorem 3. *Let Λ be a finite ring of cardinality invertible in k , and let $\mathcal{F}$ be an étale sheaf of Λ -modules, locally free of rank 1 on $U_{\text{ét}}$, with ramification bounded by $\mathfrak{m} = \sum n_i P_i$. Let $\mathfrak{n} = n_i P_i \subset \mathfrak{m}$ be a sub modulus of $\mathfrak{m}$. Then $\mathcal{F}^{[\deg \mathfrak{n}]}$ is tamely ramified at the generic point $\eta_{\mathfrak{n}}$ of the exceptional divisor $E_{\mathfrak{n}}$ of the blowup $X_{\mathfrak{n}}$ of $C^{(\deg \mathfrak{n})}$ at the closed point $\mathfrak{n}$.*

The thesis is organized as follows:

Chapter 1 - Preliminaries provides the necessary preliminaries and covers the foundational material upon which this work is based, generally without providing proofs.

Chapter 2 - Ramification After Blowup Is Tame is devoted to the proof of [Theorem 3](#), along with several corollaries that will be instrumental in the proof of [Theorem 2](#).

¹The notion of a multiplicative locally free Λ -module of rank 1 is due to [\[Gui19\]](#) and corresponds to isogenies $G \rightarrow \text{Pic}_{C, \mathfrak{m}}$ with constant kernel $\Lambda^\times$. This concept corresponds to multiplicative characters of $H^1(\text{Pic}_{C, \mathfrak{m}}, \mathbb{Q}/\mathbb{Z})$ in the formulation of [\[Tak19\]](#), and generalizes Hecke eigensheaves in the context of [\[Ten15\]](#).

²See the last page of [\[Gui19\]](#), Section 8.3 of [\[Ten15\]](#), or the proof of Theorem 1.2 in [\[Tak19\]](#) for details on this reduction.

Chapter 3 - Geometric Class Field Theory presents the proof of [Theorem 2](#).

Acknowledgements. First and foremost, I would like to express my sincere gratitude to Professor Yakov Varshavsky for introducing me to this subject, and for his mentoring, invaluable guidance, insightful meetings, and patience throughout this process. Secondly, I am also deeply grateful to my family for their support throughout my studies, and to my friends who motivated me to complete this work. Finally, I would like to express my gratitude to all the individuals without whom this work would not have been possible. This includes colleagues, researchers, as well as the broader global community of individuals whose passion and dedication drive us all forward.

2 Preliminaries

This section establishes the foundational definitions and theorems necessary for the remainder of this work. We focus specifically on the theory of G -torsors, which are central to our study due to their correspondence with locally free sheaves of rank 1. Subsequently, we review the relevant background on ramification theory from the existing literature. Finally, we conclude with several algebraic geometric remarks and notational conventions that will be employed implicitly throughout this thesis.

2.1 Scheme Quotients in general

basic definitions, categorical quotient, when it exists, etc, properties, universal properties, etc... prop 14 below

2.2 Torsors

In what follows, we largely adhere to the treatment of torsors and group objects found in published notes of Alex Youcis [\[You20\]](#). Let $\mathcal{C} = (\mathcal{C}, J)$ be a site and let $\mathcal{E} = Sh(\mathcal{C})$ be the associated topos. Let $\mathcal{G}$ be a group object in $\mathcal{E}$. We denote by $\mathcal{G}\mathcal{E}$ the category of objects in $\mathcal{E}$ endowed with a left $\mathcal{G}$ -action.

Definition 4. A $\mathcal{G}$ -torsor in $\mathcal{E}$ is an object $\mathcal{P}$ of $\mathcal{G}\mathcal{E}$ satisfying the following conditions:

1. The structural morphism $\mathcal{P} \rightarrow 1$ is an epimorphism in $\mathcal{E}$ (i.e., $\mathcal{P}$ is locally non-empty).
2. The map $\mathcal{G} \times \mathcal{P} \rightarrow \mathcal{P} \times \mathcal{P}$ defined by $(g, p) \mapsto (g \cdot p, p)$ is an isomorphism in $\mathcal{E}$ (i.e., $\mathcal{G}$ acts simply transitively on $\mathcal{P}$).

Since $\mathcal{E}$ is the topos of sheaves on a site $\mathcal{C}$, the definition can be reformulated in terms of covers. A $\mathcal{G}$ -sheaf $\mathcal{P}$ is a $\mathcal{G}$ -torsor if:

1. For every object $X \in \mathcal{C}$, there exists a covering $\{U_i \rightarrow X\} \in J$ such that $\mathcal{P}(U_i) \neq \emptyset$ for all i .
2. For any $X \in \mathcal{C}$ where $\mathcal{P}(X)$ is non-empty, the action of $\mathcal{G}(X)$ on $\mathcal{P}(X)$ is simply transitively.

A fundamental property of torsors is their local triviality: a $\mathcal{G}$ -sheaf $\mathcal{P}$ is a $\mathcal{G}$ -torsor if and only if it is locally isomorphic to the trivial torsor. Specifically, for every $X \in \mathcal{C}$, there must exist a cover

$\{U_i \rightarrow X\}$ such that the restriction $\mathcal{P}|_{U_i}$ is isomorphic, as a $\mathcal{G}|_{U_i}$ -sheaf, to $\mathcal{G}|_{U_i}$ acting on itself by left multiplication.

A **morphism of $\mathcal{G}$ -torsors** $f : \mathcal{P}_1 \rightarrow \mathcal{P}_2$ is a morphism of sheaves that is equivariant with respect to the $\mathcal{G}$ -action.

It is a standard result that every morphism of $\mathcal{G}$ -torsors is an isomorphism. Consequently, the category of $\mathcal{G}$ -torsors in $\mathcal{E}$ forms a groupoid.

Definition 5. We denote the groupoid of $\mathcal{G}$ -torsors in $\mathcal{E}$ by $\mathbf{Tors}(\mathcal{E}, \mathcal{G})$. The set of isomorphism classes of $\mathcal{G}$ -torsors is denoted by $\text{Tors}(\mathcal{E}, \mathcal{G})$.

Torsors exhibit functoriality with respect to the group object:

Definition 6. Let $\varphi : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ be a morphism of group sheaves on $\mathcal{C}$, and let $\mathcal{P}$ be a $\mathcal{G}_1$ -torsor. We define the **contracted product** $\mathcal{G}_2 \times^{\mathcal{G}_1} \mathcal{P}$ as the quotient sheaf $\mathcal{G}_1 \backslash (\mathcal{G}_2 \times \mathcal{P})$, where $\mathcal{G}_1$ acts on the product by:

$$g_1 \cdot (g_2, p) = (g_2 \varphi(g_1)^{-1}, g_1 \cdot p)$$

The contracted product inherits a natural left $\mathcal{G}_2$ -action given on local sections by $h \cdot [g_2, p] = [hg_2, p]$, which endows it with the structure of a $\mathcal{G}_2$ -torsor.

This construction yields a functor:

$$\varphi_* : \mathbf{Tors}(\mathcal{E}, \mathcal{G}_1) \rightarrow \mathbf{Tors}(\mathcal{E}, \mathcal{G}_2), \quad \mathcal{P} \mapsto \mathcal{G}_2 \times^{\mathcal{G}_1} \mathcal{P}$$

On the level of isomorphism classes, φ_* induces a map of pointed sets $\text{Tors}(\mathcal{E}, \mathcal{G}_1) \rightarrow \text{Tors}(\mathcal{E}, \mathcal{G}_2)$, sending the class of the trivial $\mathcal{G}_1$ -torsor to the class of the trivial $\mathcal{G}_2$ -torsor.

When $\mathcal{G}$ is a **sheaf of abelian groups** (an *abelian sheaf*), the pointed set $\text{Tors}(\mathcal{E}, \mathcal{G})$ inherits the structure of an abelian group:

Definition 7. Assume $\mathcal{G}$ is abelian sheaf. Let $\mathcal{P}_1$ and $\mathcal{P}_2$ be objects of $\mathbf{Tors}(\mathcal{E}, \mathcal{G})$. We define the **product** $\mathcal{P}_1 \otimes^{\mathcal{G}} \mathcal{P}_2$ to be the quotient sheaf $(\mathcal{P}_1 \times \mathcal{P}_2)/\mathcal{G}$, where $\mathcal{G}$ acts on T -points by having:

$$g \cdot (f_1, f_2) := (gf_1, g^{-1}f_2)$$

The group object $\mathcal{G}$ then acts on the resulting quotient via its action on the presheaf quotient, which is given on classes by:

$$g \cdot [(f_1, f_2)] = [(gf_1, f_2)] = [(f_1, gf_2)]$$

where the square brackets denote the class in the quotient set.³

Let $[\mathcal{P}_1]$ and $[\mathcal{P}_2]$ be the isomorphism classes of $\mathcal{P}_1$ and $\mathcal{P}_2$ in $\text{Tors}(\mathcal{E}, \mathcal{G})$. We define the sum $[\mathcal{P}_1] + [\mathcal{P}_2]$ to be the class $[\mathcal{P}_1 \otimes^{\mathcal{G}} \mathcal{P}_2]$. This structure turns $\text{Tors}(\mathcal{E}, \mathcal{G})$ into an abelian group, where the identity is the class of the trivial torsor and the inverse is obtained by the opposite action.

Let Λ be a commutative ring object in the topos $\mathcal{E}$. We denote by $\mathbf{Pic}(\mathcal{E}, \Lambda)$ the groupoid of locally free Λ -modules of rank 1 in $\mathcal{E}$ (i.e., invertible modules and their isomorphisms). For the multiplicative group object $G = \Lambda^\times$, there exists a classical correspondence between such Λ -modules and G -torsors. Indeed, both categories are symmetric monoidal categories, $\mathbf{Pic}(\mathcal{E}, \Lambda)$ under the tensor product $\otimes_\Lambda$, with unit object Λ . And $\mathbf{Tors}(\mathcal{E}, G)$ under the product $\otimes^G$, with unit object G acting on itself by translation.

³Equivantly, the sum is obtained as the contracted product of the $\mathcal{G} \times \mathcal{G}$ -torsor $\mathcal{P}_1 \times \mathcal{P}_2$ along the multiplication map $m : \mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$.

Throughout this thesis, we shall primarily adopt the language of G -torsors, with the understanding that the associated Λ -module structure is implicitly inferred.

The following proposition formalizes this dictionary:

Proposition 8. *The associated module functor $\Phi : \mathcal{P} \mapsto \mathcal{P} \otimes^G \Lambda$ induces a canonical equivalence of monoidal categories:*

$$\Phi : (\mathbf{Tors}(\mathcal{E}, G), \otimes^G, G) \xrightarrow{\sim} (\mathbf{Pic}(\mathcal{E}, \Lambda), \otimes_\Lambda, \Lambda)$$

with quasi-inverse functor given by $\Psi : \mathcal{L} \mapsto \underline{\text{Isom}}_\Lambda(\Lambda, \mathcal{L})$.

Lastly, for any object $\mathcal{X} \in \mathcal{E}$, there is a canonical identification between the slice category $(\mathcal{GE})/\mathcal{X}$ and the category of group objects $\mathcal{G}(\mathcal{E}/\mathcal{X})$, where $\mathcal{X}$ is viewed as having the trivial $\mathcal{G}$ -action.

We denote by $\mathbf{Tors}(\mathcal{X}, \mathcal{G})$ the category of G -torsors over $\mathcal{X}$ in $\mathcal{GE}/\mathcal{X}$.

2.3 Torsors over the Étale's Sites

Fix a scheme X , and let G be a smooth affine X -group scheme. Denote by $X_{\text{ét}}$ and $X_{\text{ét}}$ the big and small étale sites on X , respectively. The functor of points of G , which we denote by $G(\cdot) = \text{Hom}_X(-, G)$, defines a group sheaf on $X_{\text{ét}}$.

Definition 9. A *principal G -bundle* (or *principal homogeneous space*) for G is a scheme $f : Y \rightarrow X$ equipped with a left G -action $\rho : G \times_X Y \rightarrow Y$ satisfying:

1. The morphism $G \times_X Y \rightarrow Y \times_X Y$ sending $(g, y) \mapsto (gy, y)$ is an isomorphism of X -schemes.
2. There exists an étale covering $\{U_i \rightarrow X\}$ such that $Y_{U_i} \cong G_{U_i}$ as G -schemes, where G acts on itself by left translation.

Remark. Since $G \rightarrow X$ is smooth and Y is étale-locally isomorphic to G , the morphism $f : Y \rightarrow X$ is smooth and affine. Note that Y is generally not étale over X unless G is an étale group scheme (e.g., a finite constant group). Similarly, if G is finite then $Y \rightarrow X$ is finite.

Similarly to the case of G -torsors of a topos, we define a morphism of principal G -bundles to be a morphism of X -schemes commuting with the G -action. We now have:

Theorem 10. *The morphism sending $Y \mapsto \text{Hom}_X(-, Y)$ is an equivalence of categories from the category of principal G -bundles to the category of G -torsors on $X_{\text{ét}}$. Similarly, the morphism sending Y to $\text{Hom}_X(-, Y)$ to the category of G -torsors on $X_{\text{ét}}$ is an equivalence.*

Corollary 11. *There is a natural equivalence $\mathbf{Tors}(X_{\text{ét}}, G) \cong \mathbf{Tors}(X_{\text{ét}}, G)$ inducing a bijection of pointed sets $\mathbf{Tors}(X_{\text{ét}}, G) \xrightarrow{\cong} \mathbf{Tors}(X_{\text{ét}}, G)$ which is an isomorphism of abelian groups if G is abelian. And every G -torsor, in each of the above sites can be realized as a scheme, which is a principal G -bundle.*

Constant Finite Group Torsors

Let G be a finite group, within this section, we denote the associated constant group scheme over X by $\underline{G}$ to maintain a clear distinction between the group as a set and the group as a scheme. In subsequent sections, we shall follow the standard practice of identifying G with its constant group scheme and omit the underline for brevity.

$\underline{G}$ is given by $\coprod_{g \in G} X$ with the action shuffling the X 's according to multiplication.

By following the definitions, one sees that if X be a connected scheme and $f : Y \rightarrow X$ is a finite Galois cover with Galois group G . Then, $f : Y \rightarrow X$ is a principal $\underline{G}$ -bundle. (Recall that a *finite Galois cover* is a finite étale surjection $Y \rightarrow X$ with Y connected and such that $G = \text{Aut}(Y/X)$ acts transitively on the geometric points of Y lying over any geometric point of X).

On the otherhand if $f : Y \rightarrow X$ is a principal $\underline{G}$ -bundle with Y connected, then Y is a finite Galois cover with automorphism group G .

However, not all $\underline{G}$ -torsors are connected. If $H \subset G$ is a proper subgroup then any connected finite étale cover $f : Y \rightarrow X$ with Galois group H gives rise to a non-connected $\underline{G}$ -torsor by looking at the induced $\underline{G}$ -torsor $\varphi_*(Y)$ under the inclusion $\varphi : \underline{H} \rightarrow \underline{G}$. On the otherhand, if we fix a geometric point $\bar{x} \rightarrow X$, then to give an homomorphism $\rho \in \text{Hom}_{\text{cont}}(\pi_1^{\text{ét}}(X, \bar{x}), G)$ is equivalent to give a connected pointed Galois cover $(Y, \bar{y}) \rightarrow (X, \bar{x})$ with Galois group $H = \rho(\pi_1^{\text{ét}}(X, \bar{x})) \subset G$. Thus, pushing forward to $\underline{G}$ we get a principal $\underline{G}$ -bundle. The choice of a different geometric point $\bar{x}' \rightarrow X$ differ the homomorphism by an inner automorphism, thus we have:

Theorem 12. *Let X be a connected scheme and $\bar{x}$ a geometric point of X . Suppose in addition that G is a finite abstract group. Define a map*

$$\text{Hom}_{\text{cont}}(\pi_1^{\text{ét}}(X, \bar{x}), G) / \text{Inn}(G) \rightarrow \text{Tors}(X_{\text{ét}}, \underline{G}) \quad (1)$$

by sending a homomorphism $\rho : \pi_1^{\text{ét}}(X, \bar{x}) \rightarrow G$ to the principal $\underline{G}$ -bundle $\varphi_(Y)$ where Y is the principal $\rho(\pi_1^{\text{ét}}(X, \bar{x}))$ -bundle obtained above and φ is the inclusion $\rho(\pi_1^{\text{ét}}(X, \bar{x})) \hookrightarrow G$. Then, the map is a bijection of pointed sets where the trivial homomorphisms (which is the only element of its $\text{Inn}(G)$ -orbit) is the distinguished element of the left hand side.*

If G is abelian then $\text{Inn}(G)$ is trivial, and we obtain:

Corollary 13. *Let G be a finite abelian group, X a connected scheme, and $\bar{x}$ a geometric point of X . Then, the map from [Theorem 12](#) induces an isomorphism of abelian groups*

$$\text{Hom}_{\text{cont}}(\pi_1^{\text{ét}}(X, \bar{x}), G) \xrightarrow{\cong} \text{Tors}(X_{\text{ét}}, \underline{G}) \quad (2)$$

Decomposition of Torsors

because the contracted product is also P/H , we should add this!!! say this. etc... maybe have a section about quotient of schemes. In the case of finite groups, the functoriality of the contracted product has a concrete geometric interpretation as a tower of covers. Let G be a finite abelian group and $H \subseteq G$ a subgroup. Let $\pi : G \rightarrow G/H$ be the natural quotient map.

Proposition 14. *Let $\mathcal{P} \rightarrow X$ be a G -torsor in $X_{\text{ét}}$. There exists a factorization of the morphism $\mathcal{P} \rightarrow X$:*

$$\mathcal{P} \xrightarrow{\phi} \mathcal{P}_{G/H} \xrightarrow{\psi} X$$

that is unique up to a unique isomorphism commuting with the projections, where:

1. $\mathcal{P}_{G/H} := \mathcal{P} \times^G (G/H)$ is a G/H -torsor over X .
2. $\mathcal{P}$ is an H -torsor over the scheme $\mathcal{P}_{G/H}$.

Proof. The first assertion follows directly from the definition of the contracted product functor $\pi_* : \text{Tors}(X, G) \rightarrow \text{Tors}(X, G/H)$. To see the second assertion, we note that the map $\phi : \mathcal{P} \rightarrow \mathcal{P}_{G/H}$ is

surjective. Locally on X , we may choose a cover $U \rightarrow X$ such that $\mathcal{P}|_U \cong G \times U$. Over this cover, the map ϕ is identified with the product of the quotient map and the identity:

$$\pi \times \text{id}_U : G \times U \rightarrow (G/H) \times U$$

Since $G \rightarrow G/H$ is a trivial H -torsor (the fiber over any element $\bar{g} \in G/H$ is a coset gH , which is an H -set isomorphic to H acting on itself), $\mathcal{P}$ is locally an H -torsor over $\mathcal{P}_{G/H}$. By the descent of torsors, $\mathcal{P}$ is an H -torsor over $\mathcal{P}_{G/H}$ globally.

For uniqueness, suppose $\mathcal{P} \rightarrow \mathcal{Q} \rightarrow X$ is another such factorization. Because $\mathcal{P} \rightarrow \mathcal{Q}$ is an H -torsor, $\mathcal{Q}$ represents the categorical quotient $\mathcal{P}/H$ in $X_{\text{ét}}$. Since $\mathcal{P}_{G/H}$ also represents this quotient, the universal property of quotients yields a unique isomorphism $\mathcal{Q} \xrightarrow{\sim} \mathcal{P}_{G/H}$ making all associated diagrams commute.

□

2.4 Symmetric Powers of Schemes and Torsors

This section reviews the construction of quotients for schemes and torsors under finite group actions, specifically focusing on symmetric powers. To ensure these quotients exist as schemes, we utilize the framework of admissible actions from [SGA1]. Our treatment here closely follows the exposition in [Gui19]. The definitions and results presented below are adapted from their work, the difference being that we ask Γ to act on the left, to make the later construction of symmetric powers more natural. This foundation provides the necessary criteria for admissibility and base change required to define the symmetric powers of a scheme X and a G -torsor $\mathcal{P}$ over X .

Let S be a scheme.

Definition 15 ([SGA1], V.1.7.).

- Let T be an object of a category $\mathcal{C}$ endowed with a right action of a group Γ . We say that **the quotient T/Γ exists** in $\mathcal{C}$ if the covariant functor

$$\begin{aligned} \mathcal{C} &\rightarrow \text{Sets} \\ U &\mapsto \text{Hom}_{\mathcal{C}}(T, U)^{\Gamma} \end{aligned}$$

is representable by an object of $\mathcal{C}$.

- Let T be an S -scheme. An action of a finite group Γ on T is **admissible** if there exists an affine Γ -invariant morphism $f : T \rightarrow T'$ such that the canonical morphism $\mathcal{O}_{T'} \rightarrow f_*\mathcal{O}_T$ induces an isomorphism from $\mathcal{O}_{T'}$ to $(f_*\mathcal{O}_T)^{\Gamma}$.

Proposition 16. *The following holds:*

1. ([SGA1] V.1.3). *Let T be an S -scheme endowed with an admissible right action of a finite group Γ . If $f : T \rightarrow T'$ is an affine Γ -invariant morphism such that the canonical morphism $\mathcal{O}_{T'} \rightarrow f_*\mathcal{O}_T$ induces an isomorphism from $\mathcal{O}_{T'}$ to $(f_*\mathcal{O}_T)^{\Gamma}$, then the quotient T/Γ exists and is isomorphic to T' .*
2. ([SGA1], V.1.8). *Let T be an S -scheme endowed with a right action of a finite group Γ . Then, the action of Γ on T is admissible if and only if T is covered by Γ -invariant affine open subsets.*

3. ([SGA1], V.1.9). Let T be an S -scheme endowed with an admissible right action of a finite group Γ , and let S' be a flat S -scheme. Then, the action of Γ on the S' -scheme $T \times_S S'$ is admissible, and the canonical morphism

$$(T \times_S S')/\Gamma \rightarrow (T/\Gamma) \times_S S'$$

is an isomorphism.

Now let X be an S -scheme and let $d \geq 0$ be an integer. The group S_d of permutations of $\llbracket 1, d \rrbracket$ acts on the right on the S -scheme $X^{\times_{S^d}} = X \times_S \cdots \times_S X$ by the formula

$$(x_i)_{i \in \llbracket 1, d \rrbracket} \cdot \sigma = (x_{\sigma(i)})_{i \in \llbracket 1, d \rrbracket}.$$

Proposition 17 ([Gui19] Proposition 2.27). *If X is a scheme, Zariski locally quasi-projective over S , then the right action of the symmetric group S_d on the d -fold fiber product $X^{\times_{S^d}}$ is admissible. Consequently, the quotient $\mathrm{Sym}_S^d(X) = X^{\times_{S^d}}/S_d$ exists as a scheme over S .*

Definition 18. Under the hypotheses of the proposition above, we define the **relative symmetric product** of X over S of degree d as the quotient

$$\mathrm{Sym}_S^d(X) := X^{\times_{S^d}}/S_d.$$

When the base scheme S is clear from the context, we shall denote this quotient by $X^{(d)}$.

Guingard shows that when $X = \mathrm{Spec}(B)$ and $S = \mathrm{Spec}(A)$ then $\mathrm{Sym}_S^d(X)$ is representable by an affine S -scheme (See [Gui19] Remark 2.28).

Proposition 19 ([Gui19] Proposition 2.28). *If X is flat and Zariski-locally quasi-projective over S , then $\mathrm{Sym}_S^d(X)$ is flat over S . Moreover, for any S -scheme S' , the canonical morphism*

$$\mathrm{Sym}_{S'}^d(X \times_S S') \rightarrow \mathrm{Sym}_S^d(X) \times_S S'$$

is an isomorphism.

Symmetric Powers of Torsors

Let G be a finite abelian group, let $\mathcal{P}$ be a G -torsor over an S -scheme X in $S_{\mathrm{\acute{e}t}}$.

Proposition 20 ([SGA1], IX.5.8). *Assume that $\mathcal{P}$ and X are endowed with right actions from a finite group Γ such that the morphism $\mathcal{P} \rightarrow X$ is Γ -equivariant, and that the following properties hold:*

- (a) *The right Γ -action on $\mathcal{P}$ commutes with the left G -action.*
- (b) *The right Γ -action on X is admissible, and the quotient morphism $X \rightarrow X/\Gamma$ is finite.*
- (c) *For any geometric point $\bar{x}$ of X , the action of the stabilizer $\Gamma_{\bar{x}}$ of $\bar{x}$ in Γ on the fiber $\mathcal{P}_{\bar{x}}$ of $\mathcal{P}$ at $\bar{x}$ is trivial.*

Then the action of Γ on $\mathcal{P}$ is admissible, and $\mathcal{P}/\Gamma$ is a G -torsor over X/Γ in $S_{\mathrm{\acute{e}t}}$.

By [Theorem 10](#), $\mathcal{P}$ is representable by a finite étale X -scheme. For each $i \in \llbracket 1, d \rrbracket$ let $p_i : X^{\times sd} \rightarrow X$ be the projection on i -th factor, and let us consider the G -torsor

$$p_1^{-1}\mathcal{P} \otimes^G \dots \otimes^G p_d^{-1}\mathcal{P} = K_G \backslash \mathcal{P}^{\times sd}$$

over $X^{\times sd}$, where $K_G \subseteq G^d$ is the kernel of the multiplication morphism $G^d \rightarrow G$ (the dependence on d is suppressed in the notation).⁴ The object $K_G \backslash \mathcal{P}^{\times sd}$ of $S_{\text{ét}}$ is too representable by an S -scheme which is finite étale over $X^{\times sd}$. The group S_d acts on the right on $K_G \backslash \mathcal{P}^{\times sd}$ by the formula

$$(p_i)_{i \in \llbracket 1, d \rrbracket} \cdot \sigma = (p_{\sigma(i)})_{i \in \llbracket 1, d \rrbracket}.$$

This action of S_d commutes with the left action of G on $K_G \backslash \mathcal{P}^{\times sd}$.

Proposition 21 ([Gui19] Proposition 2.32.). *If X is Zariski-locally quasi-projective on S , then the right action of S_d on $K_G \backslash \mathcal{P}^{\times sd}$ is admissible, so that the quotient $\mathcal{P}^{[d]}$ of $K_G \backslash \mathcal{P}^{\times sd}$ by S_d exists as a scheme over S . Moreover, the canonical morphism $\mathcal{P}^{[d]} \rightarrow \text{Sym}_S^d(X)$ is a G -torsor, and the morphism*

$$p_1^{-1}\mathcal{P} \otimes^G \dots \otimes^G p_d^{-1}\mathcal{P} \rightarrow r^{-1}\mathcal{P}^{[d]}$$

where $r : X^{\times sd} \rightarrow \text{Sym}_S^d(X)$ is the canonical projection, is an isomorphism of G -torsors over $X^{\times sd}$.

When G is abelian, and letting the S_d -action be a left action (via the inverse), then the condition $\prod_{i=1}^d g_i = 1$ defining K_G is invariant under permuting the coordinates. Therefore, the left action of S_d normalizes the action of K_G . The group generated by these two actions is the semidirect product $\Xi = K_G \rtimes S_d$, which is a finite group acting on the left on $\mathcal{P}^d$. Then the double quotient $(K_G \backslash \mathcal{P}^{\times sd})/S_d \cong S_d \backslash (K_G \backslash \mathcal{P}^{\times sd})$ **add here, explain why the quotienting by the double quotient exists, should the action be free or something?** is the same as the quotient $\Xi \backslash \mathcal{P}^{\times sd}$, where Ξ acts on $\mathcal{P}^{\times sd}$ by the formula:

$$((g_1, \dots, g_d), \sigma) \star (p_1, \dots, p_d) = (g_1 p_{\sigma^{-1}(1)}, g_2 p_{\sigma^{-1}(2)}, \dots, g_d p_{\sigma^{-1}(d)})$$

Corollary 22. *Under the assumptions of the above discussion, there is a natural canonical morphism*

$$q : S_d \backslash \mathcal{P}^{\times sd} \rightarrow \Xi \backslash \mathcal{P}^{\times sd}$$

or equivalently

$$q : S_d \backslash \mathcal{P}^{\times sd} \rightarrow (K_G \backslash \mathcal{P}^{\times sd})/S_d$$

between the symmetric power of $\mathcal{P}$ as a scheme and the symmetric power of $\mathcal{P}$ as a G -torsor. We make it a convention to denote the symmetric power of $\mathcal{P}$ as a scheme by $\mathcal{P}^{(d)}$, and the symmetric power of $\mathcal{P}$ as a G -torsor by $\mathcal{P}^{[d]}$. So that the above quotient map is also written as: $q : \mathcal{P}^{(d)} \rightarrow \mathcal{P}^{[d]}$.

Proof. S_d is a subgroup of Ξ , hence any map that is Ξ -invariant is automatically S_d -invariant. $\square$

Remark. Note that the above discussion does not require that $\mathcal{P}$ is a G -torsor over X , only that it is a scheme with a left G -action.

⁴Equivantly, the sum is obtained as the contracted product of the G^d -torsor $p_1^{-1}\mathcal{P}_1 \times \dots \times p_d^{-1}\mathcal{P}_d$ along the multiplication map $m : G^d \rightarrow G$.

Quotient of Torsors Towers

We now study how the symmetric power construction of [Proposition 21](#) interacts with the canonical decomposition of [Proposition 14](#). Throughout this subsection, G denotes a finite abelian group, $H \subseteq G$ a subgroup, and $\mathcal{P} \rightarrow X$ a G -torsor over an S -scheme X that is Zariski-locally quasi-projective over S . Recall that $\mathcal{P} \rightarrow X$ factors canonically as

$$\mathcal{P} \xrightarrow{\phi} \mathcal{P}_{G/H} \xrightarrow{\psi} X,$$

where $\mathcal{P}_{G/H} = \mathcal{P} \times^G (G/H)$ is a G/H -torsor over X and ϕ is an H -torsor.

In addition to the kernel $K_G = \ker(G^d \rightarrow G)$ of the summation morphism appearing in [Proposition 21](#), we shall need the analogous *summation kernels*

$$K_H := \ker(H^d \rightarrow H), \quad K_{G/H} := \ker((G/H)^d \rightarrow G/H).$$

Applying the snake lemma to the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^d & \longrightarrow & G^d & \longrightarrow & (G/H)^d \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H & \longrightarrow & G & \longrightarrow & G/H \longrightarrow 0, \end{array}$$

in which the vertical maps are the surjective summation morphisms, yields a short exact sequence

$$0 \rightarrow K_H \rightarrow K_G \rightarrow K_{G/H} \rightarrow 0. \quad (3)$$

The induced isomorphism $K_G/K_H \cong K_{G/H}$ will play a central role below.

We define four quotient schemes

$$\begin{aligned} \mathcal{P}^{[d]_G} &:= (K_G \rtimes S_d) \backslash \mathcal{P}^{\times sd}, \\ \mathcal{P}^{[d]_H} &:= (K_H \rtimes S_d) \backslash \mathcal{P}^{\times sd}, \\ (\mathcal{P}_{G/H})^{(d)} &:= (H^d \rtimes S_d) \backslash \mathcal{P}^{\times sd}, \\ (\mathcal{P}_{G/H})^{[d]} &:= ((H^d + K_G) \rtimes S_d) \backslash \mathcal{P}^{\times sd}, \end{aligned}$$

where S_d acts on $\mathcal{P}^{\times sd}$ as in [Corollary 22](#) and the subgroups $K_H \subseteq K_G \subseteq H^d + K_G$ and $H^d \subseteq H^d + K_G$ of G^d act diagonally on $\mathcal{P}^{\times sd}$ via the G -action on $\mathcal{P}$. The first space is the symmetric G -power of $\mathcal{P}$ as in [Proposition 21](#); the remaining three are, respectively, the symmetric H -power of $\mathcal{P}$ relative to ϕ , the symmetric power of the scheme $\mathcal{P}_{G/H}$, and the symmetric G/H -power of $\mathcal{P}_{G/H}$. **The latter two identifications follow from $(\mathcal{P}_{G/H})^{\times sd} \cong H^d \backslash \mathcal{P}^{\times sd}$ together with the fact that the preimage of $K_{G/H} \subseteq (G/H)^d$ under $G^d \twoheadrightarrow (G/H)^d$ equals $H^d + K_G$.** the last isomorphism like that, is from the correspondence theorem, claude generated at the end of the article, add it somewhere

Proposition 23. *The four quotient schemes above exist and fit into a canonical commutative*

diagram

$$\begin{array}{ccc}
\mathcal{P}^{[d]_H} & \xrightarrow{H} & (\mathcal{P}_{G/H})^{(d)} \\
\downarrow & & \downarrow q \\
\mathcal{P}^{[d]_G} & \xrightarrow{H} & (\mathcal{P}_{G/H})^{[d]} \\
& \searrow G & \downarrow G/H \\
& & X^{(d)},
\end{array}$$

in which each labelled arrow is a torsor under the indicated constant group. The right-hand vertical q is the canonical projection of [Corollary 22](#), applied to the G/H -torsor $\mathcal{P}_{G/H} \rightarrow X$; in general it is not a torsor under any constant group scheme (see the remark below). The unlabelled left-hand vertical $\mathcal{P}^{[d]_H} \rightarrow \mathcal{P}^{[d]_G}$ is the canonical projection induced by the inclusion of S_d -stable subgroups $K_H \rtimes S_d \subseteq K_G \rtimes S_d$ in $G^d \rtimes S_d$; it is the base change of q along the structural map $\mathcal{P}^{[d]_G} \rightarrow (\mathcal{P}_{G/H})^{[d]}$ via the Cartesian property below, and likewise fails to be a torsor under a constant group scheme in general.

Moreover, the upper square is Cartesian:

$$\mathcal{P}^{[d]_H} \cong \mathcal{P}^{[d]_G} \times_{(\mathcal{P}_{G/H})^{[d]}} (\mathcal{P}_{G/H})^{(d)}.$$

Proof. The four labelled arrows are torsors under the indicated constant groups by direct application of [Proposition 21](#) and [Proposition 14](#):

- Applied to the G -torsor $\mathcal{P} \rightarrow X$, [Proposition 21](#) gives the G -torsor $\mathcal{P}^{[d]_G} \rightarrow X^{(d)}$ (the diagonal arrow).
- Applied to the H -torsor $\phi : \mathcal{P} \rightarrow \mathcal{P}_{G/H}$ (taking $\mathcal{P}_{G/H}$ in place of the base X), [Proposition 21](#) gives the H -torsor $\mathcal{P}^{[d]_H} \rightarrow (\mathcal{P}_{G/H})^{(d)}$ (the top arrow).
- Applied to the G/H -torsor $\psi : \mathcal{P}_{G/H} \rightarrow X$, [Proposition 21](#) gives the G/H -torsor $(\mathcal{P}_{G/H})^{[d]} \rightarrow X^{(d)}$ (the bottom-right vertical), and [Corollary 22](#) produces the canonical morphism $q : (\mathcal{P}_{G/H})^{(d)} \rightarrow (\mathcal{P}_{G/H})^{[d]}$.

The factorization $\mathcal{P}^{[d]_G} \xrightarrow{H} (\mathcal{P}_{G/H})^{[d]} \xrightarrow{G/H} X^{(d)}$ of the diagonal arrow into an H -torsor followed by a G/H -torsor is the canonical decomposition of [Proposition 14](#), applied to the G -torsor $\mathcal{P}^{[d]_G} \rightarrow X^{(d)}$ and the subgroup $H \subseteq G$. Indeed, both $\mathcal{P}^{[d]_G} \times^G (G/H)$ and $(\mathcal{P}_{G/H})^{[d]}$ are realized as the admissible quotient $((H^d + K_G) \rtimes S_d) \backslash \mathcal{P}^{\times S_d}$. **here maybe add more info, like isomorphism theorem for torsors and qoutients, or something like that, to make it more clear, from which it follows**

To prove the upper square is Cartesian, set $F := \mathcal{P}^{[d]_G} \times_{(\mathcal{P}_{G/H})^{[d]}} (\mathcal{P}_{G/H})^{(d)}$. Base-changing the H -torsor $\mathcal{P}^{[d]_G} \rightarrow (\mathcal{P}_{G/H})^{[d]}$ along q exhibits F as an H -torsor over $(\mathcal{P}_{G/H})^{(d)}$. The universal property of the fiber product furnishes a canonical morphism $\mathcal{P}^{[d]_H} \rightarrow F$ over $(\mathcal{P}_{G/H})^{(d)}$, which is H -equivariant since the H -actions on both source and target are induced by the diagonal H -action on $\mathcal{P}^{\times S_d}$. **add earlier in the section of torsors that in fact base changing an H -torsor along a morphism gives another H -torsor is a torsor.** Any morphism of H -torsors is an isomorphism, so $\mathcal{P}^{[d]_H} \cong F$ and the upper square is Cartesian. The left vertical $\mathcal{P}^{[d]_H} \rightarrow \mathcal{P}^{[d]_G}$ is then identified, via this Cartesian square, with the base change of q along $\mathcal{P}^{[d]_G} \rightarrow (\mathcal{P}_{G/H})^{[d]}$. $\square$

Remark. The two vertical morphisms $q : (\mathcal{P}_{G/H})^{(d)} \rightarrow (\mathcal{P}_{G/H})^{[d]}$ and $\mathcal{P}^{[d]H} \rightarrow \mathcal{P}^{[d]G}$ are not, in general, torsors under any constant group scheme. The obstruction is purely group-theoretic: in $K_G \rtimes S_d$ the subgroup $K_H \rtimes S_d$ is not normal, because the conjugation action of S_d on K_G by coordinate permutation is non-trivial — for $k \in K_G$ and $\sigma \in S_d$, the components of $k - \sigma \cdot k$ are $k_i - k_{\sigma^{-1}(i)} \in G$, which need not lie in H . Equivalently, $H^d \rtimes S_d$ is not normal in $(H^d + K_G) \rtimes S_d$. So the naive set-theoretic quotients $(K_G \rtimes S_d)/(K_H \rtimes S_d)$ and $((H^d + K_G) \rtimes S_d)/(H^d \rtimes S_d)$ are not groups, even though $K_G/K_H \cong K_{G/H}$ and $(H^d + K_G)/H^d \cong K_{G/H}$ as abstract groups.

Compatibility with Addition

The construction of symmetric powers is compatible with a natural addition law. For any integers $d_1, d_2 \geq 0$, there exists a canonical **addition morphism**

$$+_{d_1, d_2} : X^{(d_1)} \times_S X^{(d_2)} \rightarrow X^{(d_1+d_2)}$$

induced by the equivariant isomorphism $X^{\times_S d_1} \times_S X^{\times_S d_2} \cong X^{\times_S (d_1+d_2)}$ with respect to the inclusion of the product of symmetric groups $S_{d_1} \times S_{d_2} \subseteq S_{d_1+d_2}$.

Proposition 24. *Let $\mathcal{P}$ be a G -torsor over X . There is a canonical isomorphism of G -torsors over $X^{(d_1)} \times_S X^{(d_2)}$:*

$$+_{d_1, d_2}^{-1} \left(\mathcal{P}^{[d_1+d_2]} \right) \cong r_1^{-1} \mathcal{P}^{[d_1]} \otimes^G r_2^{-1} \mathcal{P}^{[d_2]}.$$

Proof. Let $\pi = r_{d_1} \times r_{d_2}$. By the commutativity of the following diagram

$$\begin{array}{ccc} X^{\times_S d_1} \times_S X^{\times_S d_2} & \xrightarrow{\sim} & X^{\times_S (d_1+d_2)} \\ \pi \downarrow & & \downarrow r_{d_1+d_2} \\ X^{(d_1)} \times_S X^{(d_2)} & \xrightarrow{+_{d_1, d_2}} & X^{(d_1+d_2)} \end{array}$$

and applying [Proposition 21](#), we obtain canonical isomorphisms:

$$\pi^{-1} \left(+_{d_1, d_2}^{-1} \mathcal{P}^{[d_1+d_2]} \right) \cong r_{d_1+d_2}^{-1} \mathcal{P}^{[d_1+d_2]} \cong \bigotimes_{i=1}^{d_1+d_2} p_i^{-1} \mathcal{P}$$

and

$$\pi^{-1} (r_1^{-1} \mathcal{P}^{[d_1]} \otimes^G r_2^{-1} \mathcal{P}^{[d_2]}) \cong \left(\bigotimes_{i=1}^{d_1} p_i^{-1} \mathcal{P} \right) \otimes^G \left(\bigotimes_{j=d_1+1}^{d_1+d_2} p_j^{-1} \mathcal{P} \right).$$

Since both pullbacks identify with

$$\bigotimes_{k=1}^{d_1+d_2} p_k^{-1} \mathcal{P}$$

in an $S_{d_1} \times S_{d_2}$ -equivariant manner, the isomorphism descends to the quotient $X^{(d_1)} \times_S X^{(d_2)}$ by [Proposition 19](#). $\square$

2.5 Algebraic Preliminaries on Ramification

This section reviews the ramification of discrete valuations, beginning with the general theory of *discrete valuation rings* and their integral closures in finite separable extensions. We then specialize to complete rings in Galois extensions to describe the ramification filtration via lower and upper numbering, following the treatments in [Stacks, Tag 0EXQ] and [Ser79].

Definition 25 ([Stacks, Tag 09E4]). We say that $A \rightarrow B$ or $A \subset B$ is an extension of discrete valuation rings if A and B are discrete valuation rings and $A \rightarrow B$ is injective and local. In particular, if π_A and π_B are uniformizers of A and B , then $\pi_A = u\pi_B^e$ for some $e \geq 1$ and unit u of B . The integer e does not depend on the choice of the uniformizers as it is also the unique integer ≥ 1 such that

$$\mathfrak{m}_A B = \mathfrak{m}_B^e$$

The integer e is called the *ramification index* of B over A . We say that B is *weakly unramified* over A if $e = 1$. If the extension of residue fields $\kappa_A = A/\mathfrak{m}_A \subset \kappa_B = B/\mathfrak{m}_B$ is finite, then we set $f = [\kappa_B : \kappa_A]$ and we call it the *residual degree* or residue degree of the extension $A \subset B$.

Note that we do not require the extension of fraction fields to be finite.

Now let A be a discrete valuation ring with fraction field K , let L/K be a finite separable field extension and let $B \subset L$ be the integral closure of A in L . Then the ring extension $A \subset B$ is finite, hence B is Noetherian. The dimension of B is 1, hence B is a Dedekind domain. Let $\mathfrak{m}_1, \dots, \mathfrak{m}_n$ be the maximal ideals of B (i.e., the primes lying over $\mathfrak{m}_A$). We obtain extensions of discrete valuation rings

$$A \subset B_{\mathfrak{m}_i}$$

and hence ramification indices e_i and residue degrees f_i . We have

$$[L : K] = \sum_{i=1, \dots, n} e_i f_i$$

Note that if A is henselian (e.g. A is complete) then $n = 1$.

Definition 26 ([Stacks, Tag 09E9]). Let A be a discrete valuation ring with fraction field K . Let L/K be a finite separable extension. With B and $\mathfrak{m}_i$, $i = 1, \dots, n$ as above, we say the extension L/K is

1. unramified with respect to A if $e_i = 1$ and the extension $\kappa(\mathfrak{m}_i)/\kappa_A$ is separable for all i ,
2. tamely ramified with respect to A if either the characteristic of κ_A is 0 or the characteristic of κ_A is $p > 0$, the field extensions $\kappa(\mathfrak{m}_i)/\kappa_A$ are separable, and the ramification indices e_i are prime to p , and
3. totally ramified with respect to A if $n = 1$ and the residue field extension $\kappa(\mathfrak{m}_1)/\kappa_A$ is trivial.

If the discrete valuation ring A is clear from context, then we sometimes say L/K is unramified, totally ramified, or tamely ramified for short.

Now and for the rest of the section assume A is a *complete discrete valuation ring* with uniformizer π and residue field κ , which we assume to be perfect. When A and κ are of the same characteristic

$p > 0$, then A contains a coefficient field $k \cong \kappa$ and a well known structure theorem holds: $A = k[[\pi]] \cong k[[t]]$. Let K be the fraction field of A , then $K = k((\pi))$.

For the remainder of this section, assume A is a *complete discrete valuation ring* with uniformizer π and a perfect residue field κ . In the equal characteristic case, where $\text{char}(A) = \text{char}(\kappa) = p > 0$, the Cohen Structure Theorem implies that A contains a coefficient field $k \cong \kappa$, such that $A \cong k[[\pi]] \cong k[[t]]$. Consequently, the fraction field is given by $K = k((\pi))$. Let L/K be a finite Galois extension with Galois group G . The integral closure B of A in L is itself a complete discrete valuation ring. Since the residue field κ is perfect, the extension of residue fields is separable, and there exists a uniformizer $\Pi \in B$ such that $B = A[\Pi]$.

The *lower numbering ramification filtration* of G , denoted by $(G_i)_{i \geq -1}$, is defined as:

$$G_i = \{\sigma \in G \mid v_B(\sigma(x) - x) \geq i + 1 \text{ for all } x \in B\}$$

where v_B is the valuation on L associated with B . In this indexing, $G_{-1} = G$ and G_0 corresponds to the *inertia group* of the extension L/K . Note that L/K is unramified if and only if $G_0 = \{1\}$, and it is tamely ramified if and only if $G_1 = \{1\}$.

A standard result in the theory of local fields (complete local fields with a discrete valuation and perfect residue field) ensures that the condition in the definition of G_i need only be checked for the uniformizer Π of B . Specifically, if we define the function

$$i_K^L(\sigma) = v_B(\sigma(\Pi) - \Pi)$$

for $\sigma \in G$, then $G_i = \{\sigma \in G \mid i_K^L(\sigma) \geq i + 1\}$. The subgroups G_i are normal in G and become trivial for sufficiently large i .

Consider a tower of fields $K \subset E \subset L$, and let $H = \text{Gal}(L/E) \subset G$. The lower numbering is compatible with subgroups:

$$G_i \cap H = H_i \quad \text{for all } i \geq -1$$

which follows from the identity $i_E^L = i_K^L|_H$. However, the lower numbering does not behave as simply with respect to quotients. If H is normal in G , the image of G_i in G/H is generally not $(G/H)_i$. Instead, we have $G_i H/H = (G/H)_j$, where the index j is given by the formula:

$$j = \frac{1}{e_{L/E}} \sum_{\tau \in H} \min(i_K^L(\tau), i + 1) - 1$$

In the literature, one reindexes the ramification groups by defining the Herbrand function $\phi_{L/K} : [-1, \infty) \rightarrow [-1, \infty)$:

$$\phi_{L/K}(i) = \frac{1}{e_{L/K}} \sum_{\sigma \in G} \min(i_K^L(\sigma), i + 1) - 1 = \int_0^i \frac{1}{[G_0 : G_t]} dt$$

This function is continuous, strictly increasing, and piecewise linear, making it a bijection. It satisfies the composition law $\phi_{L/K} = \phi_{E/K} \circ \phi_{L/E}$ for a tower of extensions $K \subset E \subset L$. Using this function, we define the *upper numbering ramification groups* as:

$$G^i = G_{\phi_{L/K}^{-1}(i)}$$

The primary advantage of this reindexing is its compatibility with quotients: for any normal subgroup $H \subset G$, we have:

$$G^i H/H = (G/H)^i$$

for all $i \geq -1$.

Proposition 27 ([Ser79], IV Corollary 3). *the quotients G_i/G_{i+1} , $i \geq 1$, are abelian groups, and are direct products of cyclic groups of order p . The group G_1 is a p -group.*

Proposition 28 ([Ser79], IV Corollary 4). *The inertia group G_0 has the following property: It is the semi-direct product of a cyclic group of order prime to p with a normal subgroup whose order is a power of p .*

Kummer and Artin-Schreier Theories

For cyclic extensions, Kummer and Artin-Schreier theories provide explicit ways to calculate ramification. These results show how the valuation of a defining element $a \in K$ determines the ramification index and the jumps in the upper numbering filtration. Throughout this section, let K be a discrete valuation field with a perfect residue field κ of characteristic $p > 0$.

Theorem 29 (Ramification in Kummer Extensions, [Koc97], Proposition 1.83). *Assume K contains the n -th roots of unity μ_n . Let L/K be the extension given by the equation $X^n = a$ for some $a \in K^\times$ and denote by G its Galois group. Then we have:*

1. *If $v_K(a) \in n\mathbb{Z}$ and the image of $a\pi^{-v_K(a)}$ in the residue field κ is an n -th power, the extension L/K is trivial.*
2. *If $v_K(a) \in n\mathbb{Z}$ and the image of $a\pi^{-v_K(a)}$ in the residue field κ is not an n -th power, the extension L/K is cyclic and unramified.*
3. *If $v_K(a) \notin n\mathbb{Z}$, the extension L/K is cyclic and ramified. Specifically, if $\gcd(|v_K(a)|, n) = 1$, the extension is totally ramified of degree n . Otherwise it has ramification index $\frac{n}{\gcd(v_K(a), n)}$.*

Conversly, Kummer theory ensures that every cyclic extension of degree n , prime to p of a field that contains n -th roots of unity, is of the above form. Moreover, in the above we can always take $a \in \mathcal{O}_K$

Note that in the case of total ramification the extension is tamely ramified.

Theorem 30 (Ramification in Artin-Schreier Extensions, [Tho05]). *Let $\wp(x) = x^p - x$ be the Artin-Schreier operator. Let L/K be the extension given by the equation $X^p - X = a$ for some $a \in K$ and denote by G its Galois group. Then we have:*

1. *If $v_K(a) > 0$ or if $v_K(a) = 0$ and $a \in \wp(K)$, the extension L/K is trivial.*
2. *If $v_K(a) = 0$ and if $a \notin \wp(K)$, the extension L/K is cyclic of degree p and unramified.*
3. *If $v_K(a) = -m < 0$ with $m \in \mathbb{Z}_{>0}$ and if m is prime to p , the extension L/K is cyclic of degree p again and totally ramified. Moreover, its ramification groups are given by:*

$$G = G^{(-1)} = \dots = G^{(m)} \quad \text{and} \quad G^{(m+1)} = 1.$$

Conversely, Artin-Schreier theory ensures that every cyclic extension of degree p takes this form. Moreover, in the above and under the isomorphism $K \cong k((t))$, one can always take a of the form $ct^{-m} + a_{-m+1}t^{-m+1} + \dots + a_{-1}t^{-1} + a_0$. If k is algebraically closed then there is a change of variables such that $a = u^{-m}$.

2.6 Algebraic Geometry

In this section we group together some general theorems in algebraic geometry that we will be employing throughout the text. All schemes are assumed to be locally of finite type.

Theorem 31. *Let $f : X \rightarrow Y$ be a finite flat map between integral schemes, of finite type over a field k . If $Z \subset X$ is a prime divisor with generic point η_Z , then $f(Z) \subset Y$ is a prime divisor with generic point $\eta_{f(Z)}$ satisfying $f(\eta_Z) = \eta_{f(Z)}$*

Proof. f is finite hence proper hence closed so $f(Z)$ is closed subset of Y , it is irreducible as the image of an irreducible. Since $Z = \{\eta_Z\}$ we get:

$$\{f(\eta_Z)\} \subset f(Z) = f(\overline{\{\eta_Z\}}) \subseteq \overline{f(\{\eta_Z\})} = \overline{\{f(\eta_Z)\}}$$

And since $f(Z)$ is closed we get $f(Z) = \overline{\{f(\eta_Z)\}}$.

For flat map of integral schemes we have for every $x \in X$, $y = f(x)$ the dimension formula:

$$\dim(\mathcal{O}_{X,x}) = \dim(\mathcal{O}_{Y,y}) + \dim(\mathcal{O}_{X_y,x})$$

And since $\dim(\mathcal{O}_{X_y,x}) = 0$ we get $\dim(\mathcal{O}_{X,x}) = \dim(\mathcal{O}_{Y,y})$ concluding that $f(Z)$ is a prime divisor as well. $\square$

A known theorem states that:

Theorem 32. *Let X, Y be two integral schemes over a field k . If X is geometrically integral then $X \times_k Y$ is integral. If both X, Y are geometrically integral, then $X \times_k Y$ is geometrically integral.*

Theorem 33. *Let C be smooth projective curve geometrically connected over a field k . Then:*

1. $C^{(d)}$ is smooth
2. For every d , $C^{(d)}$ is integral.
3. For every d , $C^{(d)}$ is geometrically integral.
4. The product of every finite number of $C^{(d)}$ is geometrically integral.

Proof. 1. Let t_i be a local parameter for C at P_i . The local ring of the product C^d at the point $(P_1, \dots, P_d)$ is isomorphic $k[[t_1, t_2, \dots, t_d]]$ and the local ring of the quotient at the divisor $D = \sum P_i$ is $k[[t_1, \dots, t_d]]^{S_d}$ which is isomorphic to $k[[t_1, \dots, t_d]]^{S_d} \cong k[[s_1, \dots, s_d]]$ where the s_i are the symmetric polynomials, hence this ring is regular local ring.

2. C is irreducible hence C^d is irreducible hence $C^{(d)}$ is irreducible. Since $C^{(d)}$ is smooth it is reduced.
3. By [Stacks, Tag 0366], C is geometrically integral, so it follows from the above.
4. Theorem 32

$\square$

Blowups

Theorem 34 ([Stacks, Tag 0805]). *Let $X_1 \rightarrow X_2$ be a flat morphism of schemes. Let $Z_2 \subset X_2$ be a closed subscheme. Let Z_1 be the inverse image of Z_2 in X_1 . Let X'_i be the blowup of Z_i in X_i . Then there exists a cartesian diagram*

$$\begin{array}{ccc} X'_1 & \longrightarrow & X'_2 \\ \downarrow & & \downarrow \\ X_1 & \longrightarrow & X_2 \end{array}$$

of schemes.

Theorem 35. *If X is integral then $Bl_Z(X)$ is integral.*

If X is a smooth curve, we have the following result regarding its symmetric power:

Proposition 36. *Let $f : X \rightarrow S$ be a smooth morphism of relative dimension 1. Suppose f is flat and Zariski-locally quasi-projective. Then the relative symmetric power $X_S^{(d)} = \text{Sym}_S^d(X)$ is smooth over S of relative dimension d .*

Proof. Since $f : X \rightarrow S$ is smooth, the d -fold product X_S^d is smooth over S . By Proposition 19, the quotient $X_S^{(d)}$ is flat over S . Smoothness is a fiberwise property for flat morphisms of finite presentation (cf. [Stacks, Tag 01V8]); thus, it suffices to verify the smoothness of the fibers. For every $s \in S$, the fiber of the symmetric power is given by:

$$(\text{Sym}_S^d(X))_s \cong \text{Sym}_{\kappa(s)}^d(X_s)$$

Consequently, we may reduce to the case where $S = \text{Spec}(k)$ for a field k . Since smoothness is preserved under base change to the algebraic closure, we may further assume $k = \bar{k}$. Let $z = \sum_{i=1}^r d_i \cdot x_i \in X^{(d)}$ be a point represented by an effective cycle of degree d , where the points $x_i \in X(k)$ are distinct and $\sum d_i = d$. Choose pairwise disjoint Zariski-open subsets $U_i \subset X$ such that $x_i \in U_i$. Let $W \subset X^{(d)}$ be the open subset consisting of cycles whose support is contained in $\bigcup U_i$ and which meet each U_i with degree exactly d_i . There is a canonical isomorphism:

$$W \cong \prod_{i=1}^r U_i^{(d_i)}$$

Thus, it suffices to show that each $U_i^{(d_i)}$ is smooth of dimension d_i at the point $d_i \cdot x_i$.

We may therefore restrict our attention to the "worst" case: the diagonal point $z = d \cdot x$. Let $R = \mathcal{O}_{X,x}$ be the local ring of the curve at x . Since X is a smooth curve over k , R is a regular local ring of dimension 1, i.e., a discrete valuation ring. Its completion is $\hat{R} \cong k[[t]]$. The completed local ring of the product X^d at the point $(x, \dots, x)$ is:

$$\hat{\mathcal{O}}_{X^d, (x, \dots, x)} \cong \hat{R} \hat{\otimes}_k \dots \hat{\otimes}_k \hat{R} \cong k[[t_1, \dots, t_d]]$$

where t_i is the uniformizer for the i -th copy of X . The symmetric group S_d acts on this power series ring by permuting the variables t_i . By the *Fundamental Theorem of Symmetric Polynomials*, the ring of invariants is:

$$(k[[t_1, \dots, t_d]])^{S_d} = k[[e_1, \dots, e_d]]$$

where e_j is the j -th elementary symmetric power series in the variables t_i .

The ring $k[[e_1, \dots, e_d]]$ is a formal power series ring in d variables. A fundamental theorem in commutative algebra states that a local ring is regular if and only if its completion is a formal power series ring over a field. Since $\hat{\mathcal{O}}_{X^{(d)}, z} \cong k[[e_1, \dots, e_d]]$, the local ring $\mathcal{O}_{X^{(d)}, z}$ is a regular local ring of dimension d . This implies that $X^{(d)}$ is smooth over k of dimension d , completing the proof. $\square$

A Base-Change Lemma for Galois Extensions

Finally, we record a standard base-change lemma for Galois field extensions, used when comparing a Galois extension with the function field of a base change.

Lemma 37. *Let E/F be a finite Galois extension with group H , and let K/F be an arbitrary field extension. Fix a common overfield $\Omega \supseteq E, K$ (for instance an algebraic closure of K together with an F -embedding $E \hookrightarrow \Omega$), and form inside it the intersection $M := E \cap K$, the compositum $E \cdot K$, and the subgroup $N := \text{Gal}(E/M) \leq H$. Then*

1. $E \cdot K/K$ is Galois, and restriction is an isomorphism $\text{Gal}(E \cdot K/K) \xrightarrow{\sim} \text{Gal}(E/M) = N$; and

$$2. \text{ as } K\text{-algebras, } E \otimes_F K \cong \prod_{i=1}^{[M:F]} E \cdot K.$$

In particular $E \otimes_F K$ is a field if and only if $M = F$, i.e. if and only if E and K are linearly disjoint over F .

Proof. Write $E = F[\alpha]$ and let $g \in F[x]$ be the minimal polynomial of α , so $\deg g = |H|$ and g splits completely in E because E/F is Galois. Part (1) is the classical translation theorem: $E \cdot K/K$ is Galois (it is generated over K by the roots of the separable g , which all lie in E), and $\sigma \mapsto \sigma|_E$ embeds $\text{Gal}(E \cdot K/K)$ into H with image the subgroup fixing $E \cap K = M$ pointwise, namely $\text{Gal}(E/M) = N$. For (2), factor $g = g_1 \cdots g_m$ into monic irreducibles over K . Adjoining any single root of g to K already produces all of $E \cdot K$ (since $E = F(\alpha)$ contains every conjugate of α), so each g_i has degree $[E \cdot K : K] = [E : M] = |N|$ by (1); hence $m = |H|/|N| = [M : F]$. Therefore

$$E \otimes_F K \cong K[x]/(g) \cong \prod_{i=1}^m K[x]/(g_i) \cong \prod_{i=1}^{[M:F]} E \cdot K,$$

the last step because $K[x]/(g_i) \cong K(\alpha) = E \cdot K$ for every i . This product is a single field exactly when $m = 1$, i.e. when $[M : F] = 1$, i.e. when $M = F$. $\square$

3 Ramification After Blowup Is Tame

Throughout this section let C be a projective smooth geometrically connected curve over k . Let $\mathfrak{m} = \sum n_i P_i$ be a fixed modulus. Let $U = C \setminus \mathfrak{m}$. For any modulus $\mathfrak{n} \subset \mathfrak{m}$, we define $Z_{\mathfrak{n}}$ as the closed subscheme of $C^{(\deg \mathfrak{n})} = \text{Sym}_k^{\deg \mathfrak{n}}(C)$ defined by $\mathfrak{n}$ as a point of $C^{(\deg \mathfrak{n})}$. We then define $X_{\mathfrak{n}}$ as the blowup of $C^{(\deg \mathfrak{n})}$ at $Z_{\mathfrak{n}}$, and we denote by $E_{\mathfrak{n}} = Z_{\mathfrak{n}} \times_{C^{(d)}} X_{\mathfrak{n}}$ the exceptional divisor of this blowup, it is irreducible of codimension 1. We denote by $\eta_{\mathfrak{n}}$ the generic point of $E_{\mathfrak{n}}$.

Diagrammatically:

$$\begin{array}{ccc} \overline{\{\eta_{\mathbf{n}}\}} = E_{\mathbf{n}} & \hookrightarrow & X_{\mathbf{n}} \\ \downarrow & & \downarrow \pi_{\mathbf{n}} \\ Z_{\mathbf{n}} & \hookrightarrow & C^{(\deg \mathbf{n})} \end{array} \quad (4)$$

Our main goal in this section is to prove [Theorem 3](#):

Theorem 38. *Let G be a finite abelian group. And let $\mathcal{P} \rightarrow U$ be a G -torsor in $U_{\text{ét}}$ with ramification bounded by $\mathbf{m} = \sum n_i P_i$. Let $\mathbf{n} = n_i P_i \subset \mathbf{m}$ be a sub modulus of $\mathbf{m}$. Then $\mathcal{P}^{[\deg \mathbf{n}]}$ is tamely ramified at $\eta_{\mathbf{n}}$.*

Our approach proceeds in three stages. First, we establish the result for the cyclic case $G = \mathbb{Z}/p\mathbb{Z}$ and for groups G where $\gcd(|G|, p) = 1$. Next, we extend the proof to cyclic p -groups $G = \mathbb{Z}/p^r\mathbb{Z}$. Finally, for a general finite abelian group G , we conclude by applying the structure theorem for finite abelian groups and [Proposition 14](#) to decompose the torsor into these constituent cases.

3.1 Ramification of G -Torsors

Generally speaking, assume we are given a locally Noetherian scheme X , a dense open $U \subset X$, a finite étale morphism $f : Y \rightarrow U$ and a prime divisor $Z \subset X$ such that $Z \cap U = \emptyset$ and the local ring $\mathcal{O}_{X,\xi}$ of X at the generic point ξ of Z is a discrete valuation ring. Setting $K_{\xi} = \text{Frac}(\mathcal{O}_{X,\xi})$ we obtain a cartesian square

$$\begin{array}{ccc} \text{Spec}(K_{\xi}) & \longrightarrow & U \\ \downarrow & & \downarrow \\ \text{Spec}(\mathcal{O}_{X,\xi}) & \longrightarrow & X \end{array}$$

of schemes. In particular, we see that $Y \times_U \text{Spec}(K_{\xi})$ is the spectrum of a finite separable algebra L_{ξ}/K_{ξ} .

Definition 39. We say:

1. Y is *unramified* over X at (Z, ξ) resp. Y is *tamely ramified* over X at (Z, ξ) if L_{ξ}/K_{ξ} is unramified, resp. tamely ramified with respect to $\mathcal{O}_{X,\xi}$ ([Definition 26](#))
2. Y is unramified over X in codimension 1, resp. Y is tamely ramified over X in codimension 1 if L_{ξ}/K_{ξ} is unramified, resp. tamely ramified with respect to $\mathcal{O}_{X,\xi}$ for every (Z, ξ) as above.

More precisely, we decompose L_{ξ} into a product of finite separable field extensions of K_{ξ} and we require each of these to be unramified, resp. tamely ramified with respect to $\mathcal{O}_{X,\xi}$.

Now back to our original assumptions, C is projective smooth geometrically connected curve over a field k and $U = C \setminus \mathbf{m}$. Every (Z, ξ) is now a closed point P and $\mathcal{O}_{C,P}$ is a discrete valuation ring. Assume $f : Y \rightarrow U$ is actually G -torsor $\mathcal{P} \rightarrow U$ in $U_{\text{ét}}$ for G finite abelian group.

By passing to the completion $\widehat{\mathcal{O}}_{C,P}$, we obtain the complete local field $\widehat{K}_P$. Restricting the torsor $\mathcal{P}$ to the punctured formal neighborhood $\text{Spec}(\widehat{K}_P)$ allows us to study the ramification in a strictly local context. This restriction yields a G -torsor over a field, which necessarily decomposes into a disjoint union of spectra of finite separable field extensions

$$\mathcal{P}|_{\text{Spec}(\widehat{K}_P)} \cong \bigsqcup \text{Spec}(F)$$

such that:

1. These extensions $F/\widehat{K}_P$ are all isomorphic and Galois.
2. The Galois group $H = \text{Gal}(F/\widehat{K}_P)$ identifies with the stabilizer of any connected component of the pullback $\mathcal{P}|_{\text{Spec}(\widehat{K}_P)}$, which is a subgroup of G .
3. The number of such components is then given by the index $[G : H]$.

Definition 40. We say that the extension $F/\widehat{K}_P$ has *ramification bounded by r* if the ramification group H^r is trivial. We say $\mathcal{P}$ has *ramification bounded by r at P* if the corresponding local field extensions $F/\widehat{K}_P$ satisfy this condition.

Definition 41. Let $\mathfrak{m} = \sum n_i P_i$ be a modulus on C . We say the G -torsor $\mathcal{P} \rightarrow U$ has **ramification bounded by $\mathfrak{m}$** if, for each point P_i , the local restriction $\mathcal{P}|_{\text{Spec}(\widehat{K}_{P_i})}$ has ramification bounded by the coefficient n_i .

This local property remains invariant under base change to the separable closure. Let $\bar{s} = \text{Spec}(\bar{k}) \rightarrow \text{Spec}(k)$ be a geometric point. The higher ramification groups for $\mathcal{P}|_{\text{Spec}(\widehat{K}_P)}$ are isomorphic to those of the geometric restriction $\mathcal{P}_{\bar{k}}$ at any point $\bar{x}$ lying over P . Thus an equivalent definition:

Definition 42. Let $\mathfrak{m} = \sum n_i P_i$ be a modulus on C . The G -torsor $\mathcal{P} \rightarrow U$ has **ramification bounded by $\mathfrak{m}$** if for every geometric point $\bar{x}$ of $\mathfrak{m}$ (with image $\bar{s}$ in $\text{Spec}(k)$), the restriction of $\mathcal{P}$ to

$$\text{Spec}(\widehat{\mathcal{O}}_{C_{\bar{k}}, \bar{x}}) \times_{C_{\bar{k}}} U_{\bar{k}}$$

has ramification bounded by the multiplicity of $\mathfrak{m}_{\bar{s}}$ at $\bar{x}$ in the sense of [Definition 40](#).

These definitions are equivalent because $\widehat{\mathcal{O}}_{C_{\bar{k}}, \bar{x}} \cong \widehat{\mathcal{O}}_{C, P} \otimes_k \bar{k}$. Furthermore, the notions of tame ramification and unramifiedness derived here coincide with the codimension-1 criteria in [Definition 39](#).

There is another equivalent formulation through the language of characters. The group G , being a finite abelian group over k , is étale. Consequently, the correspondence between G -torsors and the fundamental group allows us to describe G -torsors via characters. Specifically, there is an isomorphism: ([Corollary 13](#))

$$\text{Hom}_{\text{cont.}}(\pi_1^{\text{ét}}(U, \bar{x}), G) \xrightarrow{\cong} \text{Tors}(U_{\text{ét}}, G)$$

In our local setting, where we consider the restriction to the complete local field $\widehat{K}_P$, the fundamental group identifies with the absolute Galois group $G_{\widehat{K}_P} := \text{Gal}(\widehat{K}_P^{\text{sep}}/\widehat{K}_P)$. Thus, the restricted G -torsor $\mathcal{P}|_{\text{Spec}(\widehat{K}_P)}$ corresponds to a unique continuous homomorphism:

$$\rho : G_{\widehat{K}_P} \rightarrow G$$

Under this correspondence, the torsor $\mathcal{P}$ has ramification bounded by r at P if and only if the homomorphism ρ trivializes the r -th ramification group in the upper numbering:

$$\rho(G_{\widehat{K}_P}^r) = \{1\}$$

Basic Properties of Ramification of G -Torsors

We now prove some elementary lemmas regarding the ramification of G -torsors.

Lemma 43. *Let G be a finite abelian group and X be a locally Noetherian scheme over a field k . Let $U \subset X$ be a dense open subset and let (Z, ξ) be a prime divisor in the complement $X \setminus U$. Assume $\mathcal{P}_1$ and $\mathcal{P}_2$ are two G -torsors on $U_{\text{ét}}$, Such that $\mathcal{P}_1$ has ramification bounded by r_1 at (Z, ξ) , and $\mathcal{P}_2$ has ramification bounded by r_2 at (Z, ξ) . Then their product $\mathcal{P}_1 \otimes^G \mathcal{P}_2$ has ramification bounded by $\max(r_1, r_2)$ at (Z, ξ) .*

Proof. Let $A = \widehat{\mathcal{O}_{X, \xi}}$ and let $K = \text{Frac}(A)$. Let $\rho_1, \rho_2 : G_K \rightarrow G$ be the associated continuous homomorphisms corresponding to the G -torsors $\mathcal{P}_1|_{\text{Spec}(K)}$, $\mathcal{P}_2|_{\text{Spec}(K)}$. We finish by noticing that the associated character of $(\mathcal{P}_1 \otimes^G \mathcal{P}_2)|_{\text{Spec}(K)}$ is $\rho = \rho_1 + \rho_2$. $\square$

Lemma 44. *Let $f : X \rightarrow Y$ be a morphism of locally Noetherian schemes. Let $U_X \subset X$ and $U_Y \subset Y$ be dense open subschemes such that $f^{-1}(U_Y) \subset U_X$.*

Let (Z_X, ξ_X) and (Z_Y, ξ_Y) be prime divisors of X and Y such that $Z_X \cap U_X = \emptyset$ and $Z_Y \cap U_Y = \emptyset$. Suppose $f(\xi_X) = \xi_Y$ and that f is étale at ξ_X . Let $\mathcal{P}$ be a G -torsor over U_Y , and let $f^{-1}\mathcal{P}$ be its pullback to $f^{-1}(U_Y) \subset U_X$. Then, the ramification of $f^{-1}\mathcal{P}$ at ξ_X is bounded by r if and only if the ramification of $\mathcal{P}$ at ξ_Y is bounded by r .

Proof. The boundedness of ramification is determined by the behavior of the torsor over the completion of the local rings at the generic points. Let $A = \mathcal{O}_{Y, \xi}$ and $B = \mathcal{O}_{X, \xi_X}$ be the discrete valuation rings at the generic points, with fraction fields K and L respectively. Since f is étale at ξ_X , the map $A \rightarrow B$ is a flat, unramified (hence weakly unramified) local homomorphism. Consequently, the extension of completions $\widehat{L}/\widehat{K}$ is a finite unramified extension of complete discretely valued fields. We finish by noticing that the upper numbering filtration on the absolute Galois group is compatible with unramified base change.⁵ $\square$

Let C be a smooth, proper (projective) curve over a field k , and let $U \subset C$ be a dense open subset. Let G be a finite abelian group. and let $\pi_U : U' \rightarrow U$ be a G -torsor in $U_{\text{ét}}$. Let C' be the normal proper model of U' over k (i.e. the unique smooth proper curve which has U' as a dense open subset). Let $\pi : C' \rightarrow C$ be the induced morphism. (which is finite generically étale over U). Let $U'^{(d)}$ be the standard d -th symmetric product of U' as a scheme, and let $U'^{[d]}$ be the torsor-theoretic symmetric product, by [Corollary 22](#) we have a canonical morphism $q_U : U'^{(d)} \rightarrow U'^{[d]}$. Note that $C'^{(d)}$ is smooth and finite over $C^{(d)}$, hence it is the normalization of $C^{(d)}$ inside $K(C'^{(d)})$. $U'^{(d)}$ is a dense open subset of $C'^{(d)}$, hence $C'^{(d)}$ is a normal proper model of $U'^{(d)}$. Let $C'^{[d]}$ be the normalization of $C^{(d)}$ in the function field $K(U'^{[d]})$, then q_U extends to a map $q : C'^{(d)} \rightarrow C'^{[d]}$, and

⁵Specifically, let $G_K = \text{Gal}(K^{\text{sep}}/K)$ and $G_L = \text{Gal}(L^{\text{sep}}/L)$. For an unramified extension, the Herbrand function is the identity, which implies that for any $r \geq 0$:

$$G_L^r = G_K^r \cap G_L$$

we have the following diagrams:

$$\begin{array}{ccc}
U' & \hookrightarrow & C' \\
\pi_U \downarrow & & \downarrow \pi \\
U & \hookrightarrow & C
\end{array}
\qquad
\begin{array}{ccc}
U'^{(d)} & \hookrightarrow & C'^{(d)} \\
qu \downarrow & & \downarrow q \\
U'^{[d]} & \hookrightarrow & C'^{[d]} \\
\downarrow & & \downarrow \\
U^{(d)} & \hookrightarrow & C^{(d)}
\end{array}$$

Lemma 45. *In the context of the preceding discussion, the morphism $U'^{(d)} \rightarrow U'^{[d]} \hookrightarrow C'^{[d]}$ is unramified in codimension 1 (see [Definition 39](#)).*

Proof. Let $C' \setminus U' = \{x_1, \dots, x_m\}$. For each $x \in C' \setminus U'$, define the locus

$$Z_x = \{D \in C'^{(d)} \mid x \in \text{Supp}(D)\}.$$

Each Z_x is a prime divisor in the symmetric product $C'^{(d)}$, and the boundary $C'^{(d)} \setminus U'^{(d)}$ is precisely the finite union $\bigcup_{i=1}^m Z_{x_i}$.

Let η be the generic point of Z_x . It suffices to show that the local ring extension $\mathcal{O}_{C'^{[d]}, q(\eta)} \hookrightarrow \mathcal{O}_{C'^{(d)}, \eta}$ is weakly unramified ([Definition 25](#)).

Let $p : C'^d \rightarrow C'^{(d)}$ be the natural quotient. Consider the closed subvariety

$$H = \{x\} \times C' \times \dots \times C' \subset C'^d.$$

The subvariety H is irreducible, and its image under p is exactly Z_x . Consequently, p maps the generic point η_H of H to the generic point η of Z_x .

$$\begin{array}{ccccc}
\eta_H & \longrightarrow & H & \longrightarrow & C'^d \\
\downarrow & & \downarrow & & \downarrow p \\
\eta & \longrightarrow & Z_x & \longrightarrow & C'^{(d)} \\
\downarrow & & \downarrow & & \downarrow q \\
q(\eta) & \longrightarrow & q(Z_x) & \longrightarrow & C'^{[d]}
\end{array}$$

It suffices to show that the local ring extension $\mathcal{O}_{C'^{[d]}, q(\eta)} \hookrightarrow \mathcal{O}_{C'^d, \eta_H}$ is weakly unramified. Let $R = \mathcal{O}_{C'^d, \eta_H}$. And let $A = \mathcal{O}_{C'^{[d]}, q(\eta)}$.

Recall that $C'^{[d]}$ is the quotient of C'^d by the finite group $\Xi = K_G \rtimes S_d$ (see [Corollary 22](#)), and the morphism $q \circ p : C'^d \rightarrow C'^{[d]}$ is *branched Galois Cover* (finite, surjective map of normal integral schemes $X \rightarrow Y$, with function field extension $K(X)/K(Y)$ Galois).

Hence the decomposition group of η_H in Ξ and the inertia group of η_H in Ξ are defined:

$$D_{\eta_H} = \{g \in \Xi \mid g(\eta_H) = \eta_H\}$$

$$I_{\eta_H} = \{g \in D_{\eta_H} \mid g^*(u) \equiv u \pmod{\mathfrak{m}_R} \text{ for all } u \in R\}$$

Claim: The inertia group I_{η_H} is trivial.

Let $K = K(C')$ be the function field of C' , then $K(C'^d) = K^{\otimes_k d} \cong \text{Frac}(K_1 \otimes_k K_2 \otimes_k \cdots \otimes_k K_d)$ Where K_i is the copy of K corresponding to the i -th factor of C'^d . The closed subscheme H corresponds to the prime ideal $\mathfrak{p}_H \subset \mathcal{O}_{C'^d}$, and $R = \mathcal{O}_{C'^d, \mathfrak{p}_H}$. The residue field of R is exactly $K(H) \cong \text{Frac}(k(x) \otimes_k K_2 \otimes_k \cdots \otimes_k K_d)$ where $k(x)$ is the residue field of x .

The action of $\gamma = (g_1, \dots, g_d, \sigma) \in \Xi$, via the pullback (γ^*) on $K(C'^d)$ does two things (in that order):

1. It permutes the tensor factors according to σ . A function strictly in K_i is mapped to a function in $K_{\sigma(i)}$.
2. It applies the field automorphism $g_i^* \in \text{Aut}(K/k)$ to the respective factors.

Let $\gamma \in I_{\eta_H} \subset \Xi$, Then the induced automorphism $\bar{\gamma}^*$ on $K(H)$ must be the identity map. This implies that σ must fix i for every $i \geq 2$ (The K_i are linearly independent), and hence $\sigma = id$. The trivial action of $\bar{\gamma}^*$ restricts to a trivial action of g_i^* on K_i for every $i \geq 2$, and hence $g_i = 1$ for every $i \geq 2$ (since the action of G on $K(C')$ is faithful). Finally, since $g_1 g_2 \cdots g_d = 1$, we also have $g_1 = 1$. Since the inertia group I_{η_H} is trivial, the local ring extension $A \hookrightarrow R$ is weakly unramified. (See for example [Stacks, Tag 09E3], [Stacks, Tag 0BTF])

□

3.2 Proof of Theorem 3

We argue that it is enough to prove Theorem 38 for $C = \mathbb{P}_k^1$, $\mathfrak{n} = d \cdot 0$, and $\mathcal{P} \rightarrow \mathbb{G}_m = \mathbb{P}_k^1 \setminus \{0, \infty\}$ a G -torsor given by an explicit global equation. Informally, this is because *the ramification of $\mathcal{P}^{[d]}$ at $\eta_{\mathfrak{n}}$ is a formal-local invariant of $\mathcal{P}$ at P* : it depends only on the completed local ring $\widehat{\mathcal{O}_{C,P}} \cong k[[x]]$ together with the restriction of $\mathcal{P}$ to the punctured formal disc $\text{Spec}(k((x)))$, since all of the constructions involved — self products, quotients by finite groups, and blowups — commute with the flat base change to the completion. We emphasize that this is not a general principle that can simply be quoted: it is a statement about the particular constructions at hand, and it is proved by checking each construction separately. This is the content of Lemma 46 and Corollary 47 below. The second ingredient of the reduction is Lemma 48: every local torsor class appearing in Theorem 38 is realized by an explicit torsor over $\mathbb{G}_m \subset \mathbb{P}_k^1$.

Throughout this subsection we assume that k is algebraically closed; in particular every closed point of C is k -rational, and $\mathfrak{n} = dP$ with $P \in C(k)$ and $d = \deg \mathfrak{n}$.

Remark. The assumption that k is algebraically closed is harmless in our situation, for the following reasons.

1. **(Base change to $\bar{k}$.)** Assume k is perfect, so that $\bar{k} = k^{sep}$. The formation of $C^{(d)}$ (a quotient by a finite group), of the blowup $X_{\mathfrak{n}}$ along $Z_{\mathfrak{n}}$, and of $\mathcal{P}^{[d]}$ (Corollary 22) commutes with the flat base change $\text{Spec}(\bar{k}) \rightarrow \text{Spec}(k)$. Since $E_{\mathfrak{n}} \cong \mathbb{P}_k^{\deg \mathfrak{n} - 1}$ is geometrically irreducible, $(E_{\mathfrak{n}})_{\bar{k}}$ is irreducible and its generic point $\bar{\eta}$ lies over $\eta_{\mathfrak{n}}$; moreover $\mathcal{O}_{X_{\mathfrak{n}}, \eta_{\mathfrak{n}}} \rightarrow \mathcal{O}_{(X_{\mathfrak{n}})_{\bar{k}}, \bar{\eta}}$ is a weakly unramified extension of discrete valuation rings with separable residue field extension. The proof of Lemma 44 uses only these two properties of the extension (there the extension was moreover finite étale), so it applies verbatim: the ramification of $(\mathcal{P}_{\bar{k}})^{[d]} = (\mathcal{P}^{[d]})_{\bar{k}}$ at $\bar{\eta}$ is bounded by r if and only if the ramification of $\mathcal{P}^{[d]}$ at $\eta_{\mathfrak{n}}$ is. Hence unramifiedness and tameness at $\eta_{\mathfrak{n}}$ may be checked after base change to $\bar{k}$.

2. **(Rationality of P — a caveat.)** If P is a closed point with $\kappa(P) \neq k$, then after base change to $\bar{k}$ the point Z_n corresponds to the divisor $\sum_{\bar{P} \mapsto P} n\bar{P}$ on $C_{\bar{k}}$, whose support consists of $[\kappa(P) : k]$ distinct points, whereas the computation below treats a modulus supported at a single rational point. We therefore prove [Theorem 38](#) under the standing assumption that k is algebraically closed; this suffices for our purposes, since in the proof of [Theorem 54](#) the field is assumed algebraically closed before the present theorem is invoked.

We now formulate the formal–local invariance precisely. Fix $P \in C(k)$ lying in the support of $\mathfrak{m}$, set $\mathfrak{n} = dP$, and fix a uniformizer x of $\mathcal{O}_{C,P}$; it induces an isomorphism $\widehat{\mathcal{O}_{C,P}} \cong k[[x]]$. The scheme $\mathrm{Spec}(k[[x]])$ has two points — the closed point, mapping to P , and the generic point, mapping to the generic point of C — so the punctured formal disc $\mathrm{Spec}(k((x)))$ maps to U , and we denote by

$$\mathcal{P}_x := \mathcal{P} \times_U \mathrm{Spec}(k((x)))$$

the corresponding restriction of $\mathcal{P}$; it is a G -torsor over $\mathrm{Spec}(k((x)))$. Let $e_1, \dots, e_d \in k[[x_1, \dots, x_d]]$ be the elementary symmetric polynomials in $x_1, \dots, x_d$, and set

$$V = \mathrm{Spec}\left(k[[e_1, \dots, e_d]][e_d^{-1}]\right), \quad W = \mathrm{Spec}\left(k[[x_1, \dots, x_d]][(x_1 \cdots x_d)^{-1}]\right).$$

The inclusion $k[[e_1, \dots, e_d]] \subset k[[x_1, \dots, x_d]]$ induces a finite flat morphism $\hat{r} : W \rightarrow V$ (note that inverting $e_d = x_1 \cdots x_d$ inverts each x_i), and for each i we let

$$q_i : W \rightarrow \mathrm{Spec}(k((x))), \quad x \mapsto x_i.$$

The group S_d acts on W over V by permuting $x_1, \dots, x_d$, and $q_{\sigma(i)} = q_i \circ \sigma$.

Lemma 46 (Formal–local invariance). *In the above notation:*

1. The choice of x induces an isomorphism $\widehat{\mathcal{O}_{C^{(d)}, Z_n}} \cong k[[e_1, \dots, e_d]]$ carrying the maximal ideal to $(e_1, \dots, e_d)$, such that the induced morphism $c : \mathrm{Spec}(k[[e_1, \dots, e_d]]) \rightarrow C^{(d)}$ is flat and satisfies $c^{-1}(U^{(d)}) = V$; we write $c_V : V \rightarrow U^{(d)}$ for the restriction.
2. There is an induced isomorphism $\widehat{\mathcal{O}_{X_n, \eta_n}} \cong \kappa[[e_d]]$, where $\kappa = k(\frac{e_1}{e_d}, \dots, \frac{e_{d-1}}{e_d})$. In particular $\widehat{K}_{\eta_n} := \mathrm{Frac}(\widehat{\mathcal{O}_{X_n, \eta_n}}) \cong \kappa((e_d))$, and the canonical morphism $\mathrm{Spec}(\widehat{K}_{\eta_n}) \rightarrow U^{(d)}$ factors as $\mathrm{Spec}(\kappa((e_d))) \rightarrow V \xrightarrow{c_V} U^{(d)}$, where the first map does not depend on $(C, \mathcal{P})$.
3. There is a canonical isomorphism of G -torsors over V

$$\mathcal{P}^{[d]} \times_{U^{(d)}} V \cong \Xi \backslash \left(q_1^{-1} \mathcal{P}_x \times_W \cdots \times_W q_d^{-1} \mathcal{P}_x \right).$$

Consequently, the G -torsor $\mathcal{P}^{[d]}|_{\mathrm{Spec}(\widehat{K}_{\eta_n})}$ — and hence whether the ramification of $\mathcal{P}^{[d]}$ at η_n is bounded by r (resp. tame, resp. unramified) — depends only on d and on the isomorphism class of the G -torsor $\mathcal{P}_x$ over $k((x))$, and not otherwise on the pair $(C, \mathcal{P})$.

Proof. (1) Since P is a k -rational point of the smooth curve C , the uniformizer x induces $\widehat{\mathcal{O}_{C,P}} \cong k[[x]]$. As completion at rational points takes products of k -varieties to completed tensor products, we get $\widehat{\mathcal{O}_{C^{(d)}, (P, \dots, P)}} \cong k[[x]] \widehat{\otimes}_k \cdots \widehat{\otimes}_k k[[x]] = k[[x_1, \dots, x_d]]$. Write $A = \mathcal{O}_{C^{(d)}, Z_n}$ and $B = \mathcal{O}_{C^{(d)}, (P, \dots, P)}$. Since k is algebraically closed, a closed point $(Q_1, \dots, Q_d) \in C^d$ maps to $Z_n = dP$

under the finite projection $r : C^d \rightarrow C^{(d)}$ if and only if $Q_1 = \dots = Q_d = P$; hence $(P, \dots, P)$ is the unique point of C^d over Z_n , $B = (r_* \mathcal{O}_{C^d})_{Z_n}$ is a finite A -module, and $A = B^{S_d}$ (formation of invariants commutes with localization at invariant primes). Since $\mathfrak{m}_B$ is the unique maximal ideal of B over $\mathfrak{m}_A$, we have $\mathfrak{m}_B^N \subset \mathfrak{m}_A B \subset \mathfrak{m}_B$ for some N , so the $\mathfrak{m}_A$ -adic and $\mathfrak{m}_B$ -adic topologies on B agree, and therefore $B \otimes_A \hat{A} \cong \hat{B} \cong k[[x_1, \dots, x_d]]$. Now $A = B^{S_d}$ is the kernel of $B \rightarrow \prod_{\sigma \in S_d} B$, $b \mapsto (\sigma(b) - b)_\sigma$; as $\hat{A}$ is flat over A , tensoring with $\hat{A}$ preserves this kernel, so

$$\hat{A} \cong (B \otimes_A \hat{A})^{S_d} \cong k[[x_1, \dots, x_d]]^{S_d}.$$

Finally, $k[[x_1, \dots, x_d]]^{S_d} = k[[e_1, \dots, e_d]]$: the ring $k[x_1, \dots, x_d]$ is a finite free module over $k[e_1, \dots, e_d]$, the same topological argument identifies $k[[x_1, \dots, x_d]]$ with $k[x_1, \dots, x_d] \otimes_{k[e_1, \dots, e_d]} k[[e_1, \dots, e_d]]$, and taking S_d -invariants — again a kernel, preserved by the flat base change $k[e_1, \dots, e_d] \rightarrow k[[e_1, \dots, e_d]]$ — yields $k[[e_1, \dots, e_d]]$. Under these identifications the maximal ideal of $\hat{A}$ is $(e_1, \dots, e_d)$, and c is flat as the composition of the flat morphisms $\text{Spec}(\hat{A}) \rightarrow \text{Spec}(A) \rightarrow C^{(d)}$.

It remains to check that $c^{-1}(C^{(d)} \setminus U^{(d)}) = V(e_d)$ as sets. We have $C^{(d)} \setminus U^{(d)} = \bigcup_{Q \in \text{Supp}(\mathfrak{m})} Z_Q$ with $Z_Q = \{D : Q \in \text{Supp}(D)\}$, as in the proof of [Lemma 45](#). For $Q \neq P$ the divisor Z_Q does not contain the point $Z_n = dP$, so its ideal is not contained in $\mathfrak{m}_A$, and $c^{-1}(Z_Q) = \emptyset$. For $Q = P$, pull back along the finite surjective morphism $\text{Spec}(\hat{B}) \rightarrow \text{Spec}(\hat{A})$: the preimage of Z_P in C^d is $\bigcup_i p_i^{-1}(P)$, whose trace on $\text{Spec}(\hat{B}) = \text{Spec}(k[[x_1, \dots, x_d]])$ is $V(x_1) \cup \dots \cup V(x_d) = V(x_1 \cdots x_d) = V(e_d)$; by surjectivity, $c^{-1}(Z_P) = V(e_d)$ as claimed. Hence $c^{-1}(U^{(d)}) = V$, and likewise $\text{Spec}(\hat{B}) \times_{C^d} U^d = W$.

(2) Blowing up commutes with flat base change, and the center Z_n pulls back under c to the closed point $V(e_1, \dots, e_d)$; hence

$$\hat{Y} := \text{Bl}_{(e_1, \dots, e_d)}(\text{Spec } k[[e_1, \dots, e_d]]) \cong X_n \times_{C^{(d)}} \text{Spec}(\hat{A}),$$

and the exceptional divisor $\hat{E} \subset \hat{Y}$ is the base change of E_n along $\text{Spec}(\hat{A}/\hat{\mathfrak{m}}) = \text{Spec}(\kappa(Z_n))$, i.e. the projection $\text{pr} : \hat{Y} \rightarrow X_n$ restricts to an isomorphism $\hat{E} \cong E_n \cong \mathbb{P}_k^{d-1}$. Let $\hat{\eta}$ be the generic point of $\hat{E}$. Then $\text{pr}(\hat{\eta}) = \eta_n$, and the induced local homomorphism of discrete valuation rings $\mathcal{O}_{X_n, \eta_n} \rightarrow \mathcal{O}_{\hat{Y}, \hat{\eta}}$ carries a uniformizer to a uniformizer (the ideal of E_n pulls back to the ideal of $\hat{E}$) and induces an isomorphism of residue fields (both are the function field of $\mathbb{P}_k^{d-1}$); hence it induces an isomorphism on completions. The completion of $\mathcal{O}_{\hat{Y}, \hat{\eta}}$ is computed exactly as for the blowup of affine space at the origin, which we carry out in detail below: on the chart $u_d \neq 0$ we have $e_i = \frac{u_i}{u_d} e_d$, the local ring at $\hat{\eta}$ is $\left(\hat{A} \left[\frac{u_1}{u_d}, \dots, \frac{u_{d-1}}{u_d} \right] \right)_{(e_d)}$, its residue field is $\kappa = k\left(\frac{u_1}{u_d}, \dots, \frac{u_{d-1}}{u_d}\right) = k\left(\frac{e_1}{e_d}, \dots, \frac{e_{d-1}}{e_d}\right)$, and its completion is $\kappa[[e_d]]$. Finally, e_d is invertible in $\hat{K}_{\eta_n} = \kappa((e_d))$, so the composite $\text{Spec}(\hat{K}_{\eta_n}) \rightarrow \text{Spec}(\mathcal{O}_{\hat{Y}, \hat{\eta}}) \rightarrow \hat{Y} \rightarrow \text{Spec}(\hat{A})$ factors through V , giving the asserted factorization of $\text{Spec}(\hat{K}_{\eta_n}) \rightarrow U^{(d)}$ through c_V .

(3) Let $h : \mathcal{P}^{\times_k d} \rightarrow U^{(d)}$ denote the canonical morphism; it is finite, being the composition of the finite étale morphism $\mathcal{P}^{\times_k d} \rightarrow U^d$ with the finite morphism $U^d \rightarrow U^{(d)}$. By [Corollary 22](#), $\mathcal{P}^{[d]} = \Xi \backslash \mathcal{P}^{\times_k d} = \underline{\text{Spec}}_{U^{(d)}} \left((h_* \mathcal{O}_{\mathcal{P}^{\times_k d}})^\Xi \right)$. Pushforward along the affine morphism h commutes with the flat base change $c_V : V \rightarrow U^{(d)}$, and the formation of Ξ -invariants is a kernel, hence also commutes with flat base change; therefore

$$\mathcal{P}^{[d]} \times_{U^{(d)}} V \cong \Xi \backslash \left(\mathcal{P}^{\times_k d} \times_{U^{(d)}} V \right).$$

By step (1), $U^d \times_{U^{(d)}} V \cong W$, so $\mathcal{P}^{\times_k d} \times_{U^{(d)}} V \cong \mathcal{P}^{\times_k d} \times_{U^d} W$. For each i , the composition $W \rightarrow \text{Spec}(k[[x_1, \dots, x_d]]) \rightarrow \text{Spec}(k[[x]]) \rightarrow C$ — the middle map being the completion of the projection

$p_i : C^d \rightarrow C$, i.e. $x \mapsto x_i$ — factors through $\mathrm{Spec}(k((x))) \rightarrow U$, since x_i is invertible on W ; this composite is precisely q_i , and it commutes with $p_i : U^d \rightarrow U$. Since $\mathcal{P}^{\times_k d} = p_1^{-1}\mathcal{P} \times_{U^d} \cdots \times_{U^d} p_d^{-1}\mathcal{P}$, base changing to W gives an isomorphism

$$\mathcal{P}^{\times_k d} \times_{U^d} W \cong q_1^{-1}\mathcal{P}_x \times_W \cdots \times_W q_d^{-1}\mathcal{P}_x,$$

which is equivariant for the actions of G^d (in particular of K_G) and of S_d — the latter acting on W by permuting the x_i , compatibly with $q_{\sigma(i)} = q_i \circ \sigma$. Passing to Ξ -quotients yields the displayed isomorphism.

For the final statement: the schemes V and W , the maps q_i and $\hat{r}$, the Ξ -action, and the point $\mathrm{Spec}(\kappa((e_d))) \rightarrow V$ of (2) are all constructed from d alone; by (3), the torsor $\mathcal{P}^{[d]}|_{\mathrm{Spec}(\hat{K}_{\eta_n})}$ is the pullback along $\mathrm{Spec}(\kappa((e_d))) \rightarrow V$ of a torsor constructed from $\mathcal{P}_x$ and these data. Since the ramification at η_n is by definition read off from $\mathcal{P}^{[d]}|_{\mathrm{Spec}(\hat{K}_{\eta_n})}$ together with the discrete valuation ring $\kappa[[e_d]]$ (Definition 39, Definition 40), it depends only on $(\mathcal{P}_x, d)$. $\square$

Corollary 47 (Reduction to $\mathbb{P}^1$). *Let C, C' be smooth projective geometrically connected curves over the algebraically closed field k , let $\mathcal{P} \rightarrow U = C \setminus \mathfrak{m}$ and $\mathcal{P}' \rightarrow U' = C' \setminus \mathfrak{m}'$ be G -torsors, let $P \in \mathrm{Supp}(\mathfrak{m})(k)$ and $P' \in \mathrm{Supp}(\mathfrak{m}')(k)$, and let x, x' be uniformizers at P, P' respectively. Assume that $\mathcal{P}_x \cong \mathcal{P}'_{x'}$ as G -torsors, compatibly with the isomorphism $k((x)) \cong k((x'))$, $x \mapsto x'$. Then for every $d \geq 1$ and every r , the ramification of $\mathcal{P}^{[d]}$ at η_{dP} is bounded by r if and only if the ramification of $\mathcal{P}'^{[d]}$ at $\eta_{dP'}$ is; in particular, the one is unramified (resp. tamely ramified) at η_{dP} if and only if the other is at $\eta_{dP'}$.*

Proof. Immediate from Lemma 46: both $\mathcal{P}^{[d]}|_{\mathrm{Spec}(\hat{K}_{\eta_{dP}})}$ and $\mathcal{P}'^{[d]}|_{\mathrm{Spec}(\hat{K}_{\eta_{dP'}})}$ are identified, compatibly with the discrete valuation ring $\kappa[[e_d]]$, with the pullback along $\mathrm{Spec}(\kappa((e_d))) \rightarrow V$ of $\Xi \setminus (q_1^{-1}\mathcal{P}_x \times_W \cdots \times_W q_d^{-1}\mathcal{P}_x)$. $\square$

Lemma 48 (Realization over $\mathbb{G}_m$). *Let k be algebraically closed of characteristic $p > 0$, and let $\mathcal{Q}$ be a G -torsor over $\mathrm{Spec}(k((x)))$.*

1. *If $\mathcal{Q}$ is tamely ramified, then there exist an integer e prime to p , an injective homomorphism $\iota : \mathbb{Z}/e\mathbb{Z} \hookrightarrow G$, and an integer a prime to e , such that*

$$\mathcal{P}' := \iota_* \left(\mathrm{Spec} \left(k[x, x^{-1}][T]/(T^e - x^a) \right) \right)$$

is a G -torsor over $\mathbb{G}_m$, tamely ramified at 0 and ∞ , with $\mathcal{P}'_x \cong \mathcal{Q}$. Here $T^e = x^a$ is regarded as a $\mathbb{Z}/e\mathbb{Z}$ -torsor via a fixed primitive e -th root of unity $\zeta_e \in k$, and $\iota_ \mathcal{Q}' := (\mathcal{Q}' \times G)/(\mathbb{Z}/e\mathbb{Z})$ denotes the induced G -torsor.*

2. *If $G = \mathbb{Z}/p\mathbb{Z}$ and $\mathcal{Q}$ has ramification bounded by d (Definition 40), then there exists $f \in x^{-1}k[x^{-1}]$, either $f = 0$ or $f = cx^{-m} + a_{-m+1}x^{-m+1} + \cdots + a_{-1}x^{-1}$ with $c \neq 0$, $p \nmid m$ and $m < d$, such that*

$$\mathcal{P}' := \mathrm{Spec} \left(k[x, x^{-1}][X]/(X^p - X - f) \right)$$

is a $\mathbb{Z}/p\mathbb{Z}$ -torsor over $\mathbb{G}_m$ with $\mathcal{P}'_x \cong \mathcal{Q}$. Moreover $\mathcal{P}'$ extends to a torsor over $\mathbb{P}_k^1 \setminus \{0\}$ — in particular it is unramified at ∞ — and it has ramification bounded by d at 0.

In both cases $\mathcal{P}'_x$ denotes the restriction of $\mathcal{P}'$ to $\mathrm{Spec}(k((x)))$, as before.

Proof. (1) Since k is algebraically closed, the residue field of $k((x))$ is algebraically closed, so $k((x))$ admits no non-trivial unramified extensions, and its tamely ramified extensions are exactly the Kummer extensions $k((x^{1/e}))$ with $p \nmid e$ (Theorem 29). Hence the tame quotient G^t of $\text{Gal}(k((x))^{sep}/k((x)))$ is procyclic, $G^t \cong \varprojlim_{p \nmid e} \mu_e(k) \cong \prod_{\ell \neq p} \mathbb{Z}_\ell$; fix a topological generator γ . Let $\rho : \text{Gal}(k((x))^{sep}/k((x))) \rightarrow G$ be the continuous homomorphism classifying $\mathcal{Q}$. Tame-ness means that ρ kills the wild inertia subgroup, so ρ factors through G^t and is determined by $g := \rho(\gamma)$, whose order e is prime to p . Let $\iota : \mathbb{Z}/e\mathbb{Z} \xrightarrow{\sim} \langle g \rangle \subset G$ be given by $1 \mapsto g$, so that $\rho = \iota \circ \chi$ for a surjective character $\chi : G^t \rightarrow \mathbb{Z}/e\mathbb{Z}$. On the other hand, for $a \in \mathbb{Z}$ the equation $T^e = x^a$ defines a $\mu_e \cong \mathbb{Z}/e\mathbb{Z}$ -torsor over $\text{Spec}(k((x)))$, and by Kummer theory the classes of x^a , $a \in \mathbb{Z}/e\mathbb{Z}$, exhaust $\text{Hom}_{\text{cont.}}(G^t, \mathbb{Z}/e\mathbb{Z}) \cong \mathbb{Z}/e\mathbb{Z}$, the class of x^a being surjective precisely when $\gcd(a, e) = 1$. Choose a with the class of x^a equal to χ ; then $\mathcal{Q} \cong \iota_*(T^e = x^a)$ over $k((x))$. The same equation over $k[x, x^{-1}]$ defines a $\mathbb{Z}/e\mathbb{Z}$ -torsor over $\mathbb{G}_m$ (étale since e is invertible in k), so $\mathcal{P}' := \iota_*(\text{Spec}(k[x, x^{-1}][T]/(T^e - x^a)))$ is a G -torsor over $\mathbb{G}_m$ with $\mathcal{P}'_x \cong \mathcal{Q}$; it is tamely ramified at 0 and at ∞ since its local characters there factor through the tame quotients.

(2) If $\mathcal{Q}$ is trivial, take $f = 0$. Otherwise, since $\mathbb{Z}/p\mathbb{Z}$ is simple and $k((x))$ admits no non-trivial unramified extensions, $\mathcal{Q} = \text{Spec}(F)$ for a cyclic extension $F/k((x))$ of degree p together with an isomorphism $\text{Gal}(F/k((x))) \cong \mathbb{Z}/p\mathbb{Z}$. By Artin-Schreier theory such data correspond to non-zero classes in $k((x))/\wp(k((x)))$, where $\wp(u) = u^p - u$, via $g \mapsto k((x))[X]/(X^p - X - g)$. We claim that every class admits a representative f as in the statement:

- $\wp$ is surjective on $k[[x]]$: given $h = \sum_{i \geq 0} b_i x^i$, solve $\wp(u) = h$ with $u = \sum_{i \geq 0} u_i x^i$ by choosing u_0 with $u_0^p - u_0 = b_0$ (possible as k is algebraically closed) and setting recursively $u_i = u_{i/p}^p - b_i$, with the convention $u_{i/p} = 0$ when $p \nmid i$. Hence, we may choose a representative in $x^{-1}k[x^{-1}]$.
- If the representative has pole order m divisible by p , with leading term cx^{-m} , then subtracting $\wp(c^{1/p}x^{-m/p}) = cx^{-m} - c^{1/p}x^{-m/p}$ strictly decreases the pole order. Iterating, we reach a representative f which is either 0 (impossible, as the class is non-zero) or has pole order m prime to p .

By Theorem 30, the unique ramification jump of $F = k((x))[X]/(X^p - X - f)$ is at m , so the ramification of $\mathcal{Q}$ is bounded by d if and only if $m < d$. Finally, $f \in k[x, x^{-1}]$, so the same equation defines a finite étale $\mathbb{Z}/p\mathbb{Z}$ -cover $\mathcal{P}' \rightarrow \mathbb{G}_m$ (a torsor via $X \mapsto X + 1$) with $\mathcal{P}'_x \cong \mathcal{Q}$; and writing $y = x^{-1}$, we have $f \in yk[y]$, so $X^p - X - f$ defines a finite étale cover of $\mathbb{A}_k^1 = \text{Spec}(k[y]) = \mathbb{P}_k^1 \setminus \{0\}$ extending $\mathcal{P}'$, whence $\mathcal{P}'$ is unramified at ∞ . $\square$

Combining Corollary 47 with Lemma 48, in order to prove Theorem 38 for a torsor which is tamely ramified at P , or for a $\mathbb{Z}/p\mathbb{Z}$ -torsor with ramification bounded by d at P , we may — and do — assume from now on that

$$C = \mathbb{P}_k^1, \quad \mathfrak{m} = d \cdot 0 + \infty, \quad \mathfrak{n} = d \cdot 0, \quad U = \mathbb{G}_m,$$

and that $\mathcal{P} \rightarrow \mathbb{G}_m$ is one of the explicit torsors of Lemma 48.

We start by analyzing the blowup $\tilde{X} = \text{Bl}_0(X)$ of $X = \mathbb{A}_k^n$ where $0 \in X$ be the origin. Recall that $\tilde{X} \subset X \times_k \mathbb{P}_k^{n-1}$ is defined by the equations $x_i u_j = x_j u_i$, where $[u_1 : \dots : u_n]$ are the homogeneous coordinates of $\mathbb{P}_k^{n-1}$. The exceptional divisor $E \subset \tilde{X}$ is the fiber over the origin, $E = \{(0, [u_1 : \dots : u_n])\}$, which is of codimension 1 in $\tilde{X}$. Let $\eta \in E$ be the generic point of E , and let $R = \mathcal{O}_{\tilde{X}, \eta}$ be the associated local ring. This ring R is a discrete valuation ring (DVR) with fraction field $K = K(X) = k(x_1, \dots, x_n)$.

On the affine chart U_1 where $u_1 \neq 0$, we have $x_i = \frac{u_i}{u_1}x_1$. The coordinate ring is:

$$\mathcal{O}_{\hat{X}}(U_1) = k\left[x_1, \frac{u_2}{u_1}, \dots, \frac{u_n}{u_1}\right]$$

In this chart, the generic point η corresponds to the prime ideal $\mathfrak{p}_1 = (x_1)$. Thus, the local ring is $R = k[x_1, \frac{u_2}{u_1}, \dots, \frac{u_n}{u_1}]_{(x_1)}$. The residue field is $\kappa(\eta) = k(\frac{u_2}{u_1}, \dots, \frac{u_n}{u_1})$, and the completion of R with respect to its maximal ideal is:

$$\hat{R} = \kappa(\eta)[[x_1]]$$

In this local ring, x_1 is a uniformizer. Note that any x_i (for $i > 1$) can also serve as a uniformizer, as $x_i = (\frac{u_i}{u_1})x_1$ and $\frac{u_i}{u_1}$ is a unit in R .

For any monomial $M = x_1^{a_1} \dots x_n^{a_n} \in K$, we can write:

$$M = x_1^{\sum a_i} \left(\frac{u_2}{u_1}\right)^{a_2} \dots \left(\frac{u_n}{u_1}\right)^{a_n}$$

Since the term in parentheses is a unit in R , the valuation ν_E associated with E satisfies:

$$\nu_E(M) = \sum a_i = \deg M$$

Consequently, for any polynomial $f = f_d + f_{d+1} + \dots + f_l$, where f_i is the homogeneous part of degree i , we have $\nu_E(f) = d$ (the order of vanishing at the origin).

Our first result is:

Theorem 49. *Let $\mathcal{P} \rightarrow \mathbb{G}_m \subset \mathbb{P}_k^1$ be a G torsor which is either*

1. *tamely ramified at 0*
2. *totally ramified at 0 with $G = \mathbb{Z}/p\mathbb{Z}$ and ramification bounded by d .*

Then the ramification of the G -torsor $\mathcal{P}^{[d]} \rightarrow \mathbb{G}_m^{(d)}$ at $\eta_{\mathfrak{m}}$ the generic point of $E_{\mathfrak{m}} \subset X_{\mathfrak{m}}$ is

1. *tamely ramified if $\mathcal{P}$ was tamely ramified*
2. *unramified if $\mathcal{P}$ was totally ramified with $G = \mathbb{Z}/p\mathbb{Z}$ and ramification bounded by d*

Proof. The local ring at the generic point of the exceptional divisor of the blowup of the affine space at 0 point is $R = k[x_d, \frac{u_1}{u_d}, \dots, \frac{u_{d-1}}{u_d}]_{(x_d)}$. Where for every $i < d$, we have $x_i = \frac{u_i}{u_d}x_d$ are all uniformizers. The residue field is $\kappa(\eta) = k(\frac{u_1}{u_d}, \dots, \frac{u_{d-1}}{u_d})$ and the completion of R with respect to its maximal ideal is:

$$\hat{R} = \kappa(\eta)[[x_d]]$$

In our situation, when we take symmetric product of the affine space, the situation is similar with different coordinates if we let $e_1, \dots, e_d$ be the symmetric polynomials in $x_1, \dots, x_d$ then: The local ring is $R = k[e_d, \frac{u_2}{u_d}, \dots, \frac{u_{d-1}}{u_d}]_{(e_d)}$. $e_i = \frac{u_i}{u_d}e_d$ are all uniformizers. The residue field being: $\kappa(\eta) = k(\frac{u_1}{u_d}, \dots, \frac{u_{d-1}}{u_d})$ and the completion: $\hat{R} = \kappa(\eta)[[s_d]]$ Note that from $e_i = \frac{u_i}{u_d}e_d$ We get $\frac{e_i}{e_d} = \frac{u_i}{u_d}$ in the fraction field. hence

$$\hat{K} = k\left(\frac{e_1}{e_d}, \dots, \frac{e_{d-1}}{e_d}\right)((e_d)) \quad (5)$$

We compute directly the extesntion of complete valued fields over the complete valued field at the generic point. Note that by [Corollary 47](#) and [Lemma 48](#) we can assume $\mathcal{P} = \text{Spec } k[x, x^{-1}][X]/(X^n - a)$ for $a \in k[x, x^{-1}]$ or $\mathcal{P} = \text{Spec } k[x, x^{-1}][X]/(X^p + X - f(x, x^{-1}))$ where $f(x, x^{-1}) = cx^{-m} + a_{-m+1}x^{-m+1} + \dots + a_{-1}x^{-1} + a_0 = cx^{-m} + f_{-m+1}(x^{-1})$ where $m < d$.

We deal with each case separately.

Artin-Schreier Extensions:

Set $R = k[x, x^{-1}]$, and $S = \text{Spec } k[x, x^{-1}][X]/(X^p + X - f(x^{-1}))$ we have

$$\begin{aligned} R^{\otimes_k d} &= k[x_1, x_1^{-1}, \dots, x_d, x_d^{-1}] \\ S^{\otimes_k d} &= k[x_1, x_1^{-1}, \dots, x_d, x_d^{-1}][X_1, \dots, X_d]/(X_1^p - X_1 - f(x_1), \dots, X_d^p - X_d - f(x_d)) \\ &= R^{\otimes_k d}[X_1, \dots, X_d]/(X_1^p - X_1 - f(x_1), \dots, X_d^p - X_d - f(x_d)) \end{aligned}$$

Next, we want to understand the ring corresponding to $p_1^{-1}(\mathcal{P}) \otimes \dots \otimes p_d^{-1}(\mathcal{P})$ on C^d - the d 'th-product of the $G = \mathbb{Z}/p\mathbb{Z}$ -torsors $p_1^{-1}(\mathcal{P}), \dots, p_d^{-1}(\mathcal{P})$ on U^d . It corresponds to qoutient: $(p_1^{-1}(\mathcal{P}) \times \dots \times p_d^{-1}(\mathcal{P}))/G^{d-1}$ where the action of G^{d-1} on the product is:

$$(g_1, \dots, g_{d-1}) \cdot (p_1, p_2, \dots, p_{d-1}, p_d) = (g_1(p_1), g_1^{-1}g_2(p_2), \dots, g_{d-2}^{-1}g_{d-1}(p_{d-1}), g_{d-1}^{-1}(p_d))$$

The affine ring corrsponding to the torsor product is $(S^{\otimes_k d})^{G^{d-1}}$

Recall that the action of $g \in G = \mathbb{Z}/p\mathbb{Z}$ on X is $g(X) = X + g$ (g correspond to a number $0 \leq g \leq p-1$). So, the action of $(g_1, \dots, g_{d-1})$ on the generators $(X_1, X_2, \dots, X_{d-1}, X_d)$ Is $X_1 \mapsto X_1 + g_1, X_i \mapsto X_i - g_{i-1} + g_i$ for $1 < i < d$ and $X_d \mapsto X_d - g_{d-1}$. So we see that $Y = X_1 + \dots + X_d$ is invariant. Moreover $Y^p - Y - \sum_{i=1}^d f(x_i) = 0$ is irreducible degree p equation for Y , Since we are quotienting a rank p^d extesntion by a group of order p^{d-1} the resulting invaraint subbrng must have rank p over $R^{\otimes_k d}$, So we conclude:

$$(S^{\otimes_k d})^{G^{d-1}} \cong R^{\otimes_k d}[Y]/(Y^p - Y - \sum_{i=1}^d f(x_i))$$

The group S_d acts on $R^{\otimes_k d} = k[x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_d^{\pm 1}]$ by permuting the variables $\{x_i\}_{i=1}^d$. And since $Y = \sum_{i=1}^d X_i$ it leaves Y invaraint. The invaraint subbrng $(R^{\otimes_k d})^{S_d}$ is simply $k[e_1, e_2, \dots, e_d, e_d^{-1}]$ where $\{e_i\}$ are the symmetric polynomials in $x_1, \dots, x_d$:

$$e_k = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq d} x_{j_1} x_{j_2} \dots x_{j_k}$$

To find $(R^{\otimes_k d}[Y]/(Y^p - Y - \sum_{i=1}^d f(x_i)))^{S_d}$ its enough to express $\sum_{i=1}^d f(x_i)$ in $e_1, \dots, e_d$, this can be done with the newton polynomials, moreover, we claim the following:

Lemma 50. *Let $f(x) = cx^{-m} + a_{-m+1}x^{-m+1} + \dots + a_{-1}x^{-1} + a_0$, define $\alpha(x_1, \dots, x_d) = \sum_{i=1}^d f(x_i)$, and deonte by $e_1, \dots, e_d$ the elementary symmetric polynomials in $x_1, \dots, x_d$. If $m < d$ then $\alpha(x_1, \dots, x_d) \in k(e_1/e_d, e_2/e_d, \dots, e_{d-1}/e_d)$*

Proof. Changing variables $y_i = x_i^{-1}$ for each $i \in \{1, \dots, d\}$ We get

$$\alpha = \sum_{i=1}^d f(x_i) = \sum_{i=1}^d \left(cy_i^m + a_{-m+1}y_i^{m-1} + \dots + a_{-1}y_i + a_0 \right)$$

Rearranging the sums, we get

$$\alpha = c \sum_{i=1}^d y_i^m + a_{-m+1} \sum_{i=1}^d y_i^{m-1} + \dots + a_{-1} \sum_{i=1}^d y_i + da_0$$

Let $p_k(y_1, \dots, y_d) = \sum_{i=1}^d y_i^k$ be the k -th power sum symmetric polynomial. The expression for α is a linear combination of these power sums:

$$\alpha = cp_m(y) + a_{-m+1}p_{m-1}(y) + \dots + a_{-1}p_1(y) + da_0$$

According to the *Fundamental Theorem of Symmetric Polynomials*, any symmetric polynomial in $y_1, \dots, y_d$ can be expressed as a polynomial in the elementary symmetric polynomials $e_k(y_1, \dots, y_d)$. Since $m < d$, α is a polynomial in $e_1(y), e_2(y), \dots, e_m(y)$. ($y = (y_1, \dots, y_d)$) The elementary symmetric polynomials in $y_i = 1/x_i$ are related to the elementary symmetric polynomials in x_i as follows:

$$e_k(y_1, \dots, y_d) = \sum_{1 \leq i_1 < \dots < i_k \leq d} \frac{1}{x_{i_1} \dots x_{i_k}} = \frac{\sum_{1 \leq j_1 < \dots < j_{d-k} \leq d} x_{j_1} \dots x_{j_{d-k}}}{x_1 x_2 \dots x_d}$$

Thus,

$$e_k(y_1, \dots, y_d) = \frac{e_{d-k}(x_1, \dots, x_d)}{e_d(x_1, \dots, x_d)}$$

which concludes the proof. $\square$

Finally, restricting $\mathcal{P}^{[d]}$ to $\text{spec } \hat{K}$ we get by (5) and Theorem 30 the result. (that $\mathcal{P}$ is unramified at the generic point of the exceptional divisor of the blowup).

Kummer Extensions: Few things are different in that case,

Set $R = k[x, x^{-1}]$, and $S = R[X]/(X^n - f)$ where $f = f(x, 1/x) \in R$ In this case we have $\text{char } k = p$ and $\gcd(p, n) = 1$.

We have

$$\begin{aligned} R^{\otimes kd} &= k[x_1, x_1^{-1}, \dots, x_d, x_d^{-1}] \\ S^{\otimes kd} &= k[x_1, x_1^{-1}, \dots, x_d, x_d^{-1}][X_1, \dots, X_d]/(X_1^n - f_1, \dots, X_d^n - f_d) \\ &= R^{\otimes kd}[X_1, \dots, X_d]/(X_1^n - f_1, \dots, X_d^n - f_d) \end{aligned}$$

Where $f_i = f(x_i, x_i^{-1})$

Next, we want to figure out $(S^{\otimes kd})^{G^{d-1}}$

Recall that the action of $g \in G = \mathbb{Z}/n\mathbb{Z}$ on X is $g(X) = \zeta^g X$ (g correspond to a number $0 \leq g \leq n-1$). So, the action of $(g_1, \dots, g_{d-1})$ on the generators $(X_1, X_2, \dots, X_{d-1}, X_d)$ Is $X_1 \mapsto \zeta^{g_1} X_1$, $X_i \mapsto \zeta^{g_i - g_{i-1}} X_i$ for $1 < i < d$ and $X_d \mapsto \zeta^{-g_{d-1}} X_d$.

So we see that $Y = X_1 X_2 \dots X_d$ is invariant. And $Y^n - \prod_{i=1}^d f_i$ is irreducible degree n equation for Y , So, like before, we conclude:

$$\left(S^{\otimes_k d}\right)^{G^{d-1}} \cong R^{\otimes_k d}[Y]/(Y^n - \prod_{i=1}^d f_i)$$

The group S_d acts on $R^{\otimes_k d}[Y]/(Y^n - \prod_{i=1}^d f_i)$ by permuting the indices. on the variables x_i , On $Y = \prod_{i=1}^d X_i$ it is invaraint. The invaraint subring $(R^{\otimes_k d})^{S_d}$ is simply $k[e_1, e_2, \dots, e_d, e_d^{-1}]$ like before.

The polynomial $F = \prod_{i=1}^d f_i$ is symmetric in $\{x_i\}_1^d$ so it can be expressed as a polynomial $\tilde{F}(e_1, \dots, e_d)$ in the elementary symmetric variables. Hence the qoutient ring is:

$$\left(\frac{k[x_1^{\pm 1}, \dots, t_d^{\pm 1}][Y]}{(Y^n - \prod_{i=1}^d f_i)}\right)^{S_d} \cong \frac{k[e_1, \dots, e_d, e_d^{-1}][Y]}{(Y^n - \tilde{F}(e_1, \dots, e_d))}$$

So we see again, that restricting $\mathcal{P}^{[d]}$ to $\text{spec } \hat{K}$ we get by (5) and Theorem 29, that $\mathcal{P}$ is tamely ramified at the generic point of the exceptional divisor of the blowup. $\square$

So, we conclude

Corollary 51. *In the general settings of a curve C , and $\mathcal{P} \rightarrow U = C \setminus \mathfrak{m}$ a G -torsor with ramification bounded by $\mathfrak{n} = nP \subset \mathfrak{m}$ at P , we have:*

1. *If $G = \mathbb{Z}/p\mathbb{Z}$ then $\mathcal{P}^{[\deg \mathfrak{n}]}$ is unramified at $\eta_{\mathfrak{n}}$*
2. *If $\mathcal{P}$ is tamely ramified at $\mathfrak{n}$ then $\mathcal{P}^{[\deg \mathfrak{n}]}$ is tamely ramified at $\eta_{\mathfrak{n}}$*

Proof. By the remark opening this subsection we may assume k is algebraically closed, so that $P \in C(k)$ and $\deg \mathfrak{n} = n$. The restriction $\mathcal{P}_x$ of $\mathcal{P}$ to the punctured formal disc at P is tamely ramified, resp. a $\mathbb{Z}/p\mathbb{Z}$ -torsor with ramification bounded by n , so by Lemma 48 there is a torsor $\mathcal{P}' \rightarrow \mathbb{G}_m$ of the corresponding explicit form with $\mathcal{P}'_x \cong \mathcal{P}_x$. By Corollary 47, the ramification of $\mathcal{P}^{[\deg \mathfrak{n}]}$ at $\eta_{\mathfrak{n}}$ agrees with that of $\mathcal{P}'^{[\deg \mathfrak{n}]}$ at $\eta_{n,0}$, which was computed in Theorem 49. $\square$

Next, we generalize by induction to the case of $G = \mathbb{Z}/p^r\mathbb{Z}$.

Theorem 52. *Assume $G = \mathbb{Z}/p^r\mathbb{Z}$. Then $\mathcal{P}^{[\deg(\mathfrak{n})]}$ is unramified at $\eta_{\mathfrak{n}}$*

Proof. Again, we can assume k is algebraically closed. $\mathcal{P}^{[\deg \mathfrak{n}]}$ has bounded ramification at nP by n . Let $R = \widehat{\mathcal{O}_{C,P}}$ and let $K = \text{Frac}(R)$ Then $\mathcal{P}^{[\deg \mathfrak{n}]}|_{\text{Spec } K}$ correspond to $\rho : \text{Gal}(K^{sep}/K) \rightarrow G$. Let $H = \rho(G^0)$ be the image of the inertia subgroup. Decompose $\mathcal{P} \rightarrow U$ to $\mathcal{P} \rightarrow \mathcal{P}_{G/H} \rightarrow U$ as in Proposition 14, where $\mathcal{P} \rightarrow \mathcal{P}_{G/H}$ is an H -torsor and $\mathcal{P}_{G/H} \rightarrow U$ is a G/H torsor. Then $\mathcal{P}_{G/H}$ is unramified at P . Thus $\mathcal{P}_{G/H}$ extends to $U' = U \cup \{P\}$, and taking any point $Q \in \mathcal{P}_{G/H}$ over P , results in isomorphism $\widehat{\mathcal{O}_{\mathcal{P}_{G/H},Q}} \cong \widehat{\mathcal{O}_{C,P}}$. But because k is algebraically closed, we even have equality, and deduce that locally the G/H action on $\widehat{\mathcal{O}_{C,P}}$ is trivial. Thus, we can assume $G = H$ and that $\mathcal{P} \rightarrow U$ is totally ramified at P .

We proceed by induction on r . The base case $r = 1$ is done by the previous theorem. Decompose $\mathcal{P} \rightarrow U$ as $\mathcal{P} \rightarrow \mathcal{P}' \rightarrow U$ where $\mathcal{P} \rightarrow \mathcal{P}'$ is a $\mathbb{Z}/p\mathbb{Z}$ -torsor and $\mathcal{P}' \rightarrow U$ is a $\mathbb{Z}/p^{r-1}\mathbb{Z}$ -torsor.

- (1) By the induction hypothesis, $\mathcal{P}'^{[\deg \mathfrak{n}]}$ is unramified at $\eta_{\mathfrak{n}}$.

Let C' be the normal proper model of $\mathcal{P}'$ viewed as a scheme, and let $\pi : C' \rightarrow C$ be the extension of $\mathcal{P}' \rightarrow U$. Let $\mathfrak{n}' = \pi^{-1}(\mathfrak{n})$ be the pullback of $\mathfrak{n}$ to C' . Let $\tilde{\mathfrak{n}}$ be the pullback of $\mathfrak{n}$ to C' .

Then, following [Lemma 45](#), and the discussion surrounding it, with U' being $\mathcal{P}'$, we get that

(2) $\mathcal{P}'^{(\deg \mathfrak{n})} \rightarrow \mathcal{P}'^{[\deg \mathfrak{n}]} \hookrightarrow C'^{[\deg \mathfrak{n}]}$ is unramified at codimension one points.

And since $\mathcal{P} \rightarrow \mathcal{P}'$ is a $\mathbb{Z}/p\mathbb{Z}$ -torsor, we can apply the previous theorem to deduce that $\mathcal{P}^{[\deg \mathfrak{n}]} \rightarrow \mathcal{P}'^{(\deg \mathfrak{n})}$ is unramified at $\eta_{\mathfrak{n}}$.

Not clear how to finish here without analyzing the ramification even further. Thus, take a subgroup $H \subset G$ such that $G/H \cong \mathbb{Z}/p\mathbb{Z}$ and decompose $\mathcal{P} \rightarrow U$ as: $\square$

Finally, **Proof of Theorem [theorem 38](#)**: Idea: decompose G be the structure theorem, and finish by the above.

3.3 Extending To Product of Blowups

We conclude this section by proving some auxiliary results that would be useful in [Section 4](#).

Let X and Y be smooth schemes over a field k , and let $x \in X$ and $y \in Y$ be closed points. We denote the blowups of these schemes at the given points by $\pi_X : \text{Bl}_x(X) \rightarrow X$ and $\pi_Y : \text{Bl}_y(Y) \rightarrow Y$. Furthermore, let $\pi_{X \times Y} : \text{Bl}_{(x,y)}(X \times_k Y) \rightarrow X \times_k Y$ be the blowup of the product scheme at the point (x, y) . We denote by E_X, E_Y , and $E_{X \times Y}$ the respective exceptional divisors, and let η_X, η_Y , and $\eta_{X \times Y}$ be their generic points.

In this section, we establish the following result concerning the stability of ramification bounds under the external product of torsors.

Proposition 53. *Let G be a finite abelian group. Suppose $\mathcal{G}_X$ and $\mathcal{G}_Y$ are G -torsors defined on open subsets $U_X \subset \text{Bl}_x(X)$ and $U_Y \subset \text{Bl}_y(Y)$ that are disjoint from the exceptional divisors. If the ramification of $\mathcal{G}_X$ at η_X and $\mathcal{G}_Y$ at η_Y is bounded by r , then the external product torsors*

$$\mathcal{G}_{X \times Y} := pr_1^{-1} \mathcal{G}_X \otimes pr_2^{-1} \mathcal{G}_Y$$

has ramification bounded by r at the generic point $\eta_{X \times Y}$ of the exceptional divisor in the product blowup.

The proposition is purely local in nature, it suffices to consider the case where X and Y are affine. More precisely, by the smoothness of X and Y , we may restrict our attention to open neighborhoods of x and y that are étale over affine spaces. The rest of this section treats that case.

The Affine Case

Let $X = \mathbb{A}_k^n$ be the affine n -space over a field k , and let $0 \in X$ be the origin. Let $\tilde{X} = \text{Bl}_0(X)$ be the blowup of X at the origin. Recall that $\tilde{X} \subset X \times_k \mathbb{P}_k^{n-1}$ is defined by the equations $x_i u_j = x_j u_i$, where $[u_1 : \dots : u_n]$ are the homogeneous coordinates of $\mathbb{P}_k^{n-1}$. The exceptional divisor $E \subset \tilde{X}$ is the fiber over the origin, $E = \{(0, [u_1 : \dots : u_n])\}$, which is of codimension 1 in $\tilde{X}$. Let $\eta \in E$ be the generic point of E , and let $R = \mathcal{O}_{\tilde{X}, \eta}$ be the associated local ring. This ring R is a discrete valuation ring (DVR) with fraction field $K = K(X) = k(x_1, \dots, x_n)$.

On the affine chart U_1 where $u_1 \neq 0$, we have $x_i = \frac{u_i}{u_1}x_1$. The coordinate ring is:

$$\mathcal{O}_{\tilde{X}}(U_1) = k\left[x_1, \frac{u_2}{u_1}, \dots, \frac{u_n}{u_1}\right]$$

In this chart, the generic point η corresponds to the prime ideal $\mathfrak{p}_1 = (x_1)$. Thus, the local ring is $R = k[x_1, \frac{u_2}{u_1}, \dots, \frac{u_n}{u_1}]_{(x_1)}$. The residue field is $\kappa(\eta) = k(\frac{u_2}{u_1}, \dots, \frac{u_n}{u_1})$, and the completion of R with respect to its maximal ideal is:

$$\hat{R} = \kappa(\eta)[[x_1]]$$

In this local ring, x_1 is a uniformizer. Note that any x_i (for $i > 1$) can also serve as a uniformizer, as $x_i = (\frac{u_i}{u_1})x_1$ and $\frac{u_i}{u_1}$ is a unit in R .

For any monomial $M = x_1^{a_1} \dots x_n^{a_n} \in K$, we can write:

$$M = x_1^{\sum a_i} \left(\frac{u_2}{u_1}\right)^{a_2} \dots \left(\frac{u_n}{u_1}\right)^{a_n}$$

Since the term in parentheses is a unit in R , the valuation ν_E associated with E satisfies:

$$\nu_E(M) = \sum a_i = \deg M$$

Consequently, for any polynomial $f = f_d + f_{d+1} + \dots + f_l$, where f_i is the homogeneous part of degree i , we have $\nu_E(f) = d$ (the order of vanishing at the origin).

The Product Case

Now, let $X = \mathbb{A}^n$ and $Y = \mathbb{A}^m$ with origins $x = 0$ and $y = 0$. As before, $\text{Bl}_0(X) \subset X \times \mathbb{P}^{n-1}$ and $\text{Bl}_0(Y) \subset Y \times \mathbb{P}^{m-1}$ have exceptional divisors E_X and E_Y respectively. Consider the product $X \times_k Y \cong \mathbb{A}_k^{n+m}$. The blowup of the product at the origin $(0,0)$, denoted $\text{Bl}_{(0,0)}(X \times_k Y)$, is a subscheme of $(X \times Y) \times \mathbb{P}^{n+m-1}$ defined by:

$$\begin{cases} x_i w_j = x_j w_i & 1 \leq i, j \leq n \\ y_k w_{n+l} = y_l w_{n+k} & 1 \leq k, l \leq m \\ x_i w_{n+j} = y_j w_i & 1 \leq i \leq n, 1 \leq j \leq m \end{cases}$$

where $[w_1 : \dots : w_{n+m}]$ are the homogeneous coordinates of $\mathbb{P}^{n+m-1}$. The exceptional divisor $E_{X \times Y}$ is isomorphic to $\mathbb{P}^{n+m-1}$.

Comparison of Blowups

Both $\text{Bl}_0(X) \times_k \text{Bl}_0(Y)$ and $\text{Bl}_{(0,0)}(X \times_k Y)$ are birational to $X \times_k Y$. They share a common dense open set $\tilde{U}$ defined by the condition that neither the X -coordinates nor the Y -coordinates vanish simultaneously in the projective space:

$$\tilde{U} = \{((x, y), [w_1 : \dots : w_{n+m}]) \mid (w_1, \dots, w_n) \neq 0 \text{ and } (w_{n+1}, \dots, w_{n+m}) \neq 0\}$$

This yields a diagram of open immersions:

$$\begin{array}{ccc} & \tilde{U} & \\ f_1 \swarrow & & \searrow f_2 \\ \mathrm{Bl}_0(X) \times_k \mathrm{Bl}_0(Y) & & \mathrm{Bl}_{(0,0)}(X \times_k Y) \end{array}$$

where f_1 maps the coordinates to the respective projectivizations $[w_1 : \dots : w_n]$ and $[w_{n+1} : \dots : w_{n+m}]$. Also note that the generic point $\eta_{X \times Y}$ of $E_{X \times_k Y}$ is inside $\tilde{U}$

Extensions of DVRs

Let S be the local ring of the generic point $\eta_{X \times Y}$ of $E_{X \times Y}$ in $\tilde{U}$. On the chart where $w_1 \neq 0$ and $w_{n+1} \neq 0$, we have $x_1 = (\frac{w_1}{w_{n+1}})y_1$. Since $\frac{w_1}{w_{n+1}}$ is a unit in this chart, x_1 and y_1 are equivalent as uniformizers. We have:

$$S = k \left[x_1, \frac{w_2}{w_1} \dots, \frac{w_n}{w_1}, \frac{w_{n+1}}{w_1} \dots \frac{w_{n+m}}{w_1} \right]_{(x_1)} = k \left[\frac{w_1}{w_{n+1}}, \frac{w_2}{w_{n+1}} \dots \frac{w_n}{w_{n+1}}, y_1 \dots \frac{w_{n+m}}{w_{n+1}} \right]_{(y_1)}$$

$$k(\eta_{X \times Y}) = k \left(\frac{w_2}{w_1} \dots, \frac{w_n}{w_1}, \frac{w_{n+1}}{w_1} \dots \frac{w_{n+m}}{w_1} \right)$$

Let R_X be the local ring of the exceptional divisor E_X in $\mathrm{Bl}_0(X)$. The pullback of $E_X \times_k \mathrm{Bl}_0(Y)$ along f_1 induces an extension of DVRs $R_X \hookrightarrow S$. Which is:

1. Weakly Unramified: x_1 is a uniformizer in both R_X and S , so the ramification index is $e = 1$.
2. Residually Transcendental: The residue field extension $\kappa(\eta_X) \subset \kappa(\eta)$ is:

$$k \left(\frac{w_2}{w_1}, \dots, \frac{w_n}{w_1} \right) \subset k \left(\frac{w_2}{w_1}, \dots, \frac{w_n}{w_1}, \frac{w_{n+1}}{w_1}, \dots, \frac{w_{n+m}}{w_1} \right)$$

Hence separable.

Since this extension is generated by transcendental elements, it is separable and formally smooth at the maximal ideal ([Stacks, Tag 09E7]).

Ramification of G -Torsors

Let G be a finite abelian group. Let $\mathcal{Q}$ be a G -torsor on an open $U \subset \mathrm{Bl}_0(X)$ disjoint from E_X . Let $V \subset \mathrm{Bl}_0(Y)$ be an open subscheme, and let $\pi_X : \mathrm{Bl}_0(X) \times_k V \rightarrow \mathrm{Bl}_0(X)$ be the projection onto the first factor. By restricting this projection to $U \times_k V$, we obtain the pullback G -torsor:

$$\pi_X^{-1}(\mathcal{Q}) \cong \mathcal{Q} \times_k V$$

which is defined on the open subset $U \times_k V \subset \mathrm{Bl}_0(X) \times_k \mathrm{Bl}_0(Y)$. The extension of local rings $R_X \rightarrow S$ is weakly unramified (the ramification index $e = 1$) and residually transcendental with separable residue field extension. Under these conditions the ramification filtration is preserved. Therefore, the pullback $\pi_X^{-1}(\mathcal{Q})$ has ramification bounded by r at the generic point $\eta_{X \times Y}$ of the exceptional divisor $E_{X \times Y}$ if and only if the original torsor $\mathcal{Q}$ has ramification bounded by r at the generic point η_X of E_X

And we finish by [Lemma 43](#).

4 Geometric Class Field Theory

The main theorem of this section is

Theorem 54. *Let Λ be a finite ring of cardinality invertible in k , and let $\mathcal{F}$ be an étale sheaf of Λ -modules, locally free of rank 1 on U , with ramification bounded by $\mathfrak{m}$. Considering $U^{(d)}$ as an open subscheme of the blowup $\tilde{C}_{\mathfrak{m}}^{(d)}$ of $C^{(d)}$, we have that for sufficiently large integer d , $\mathcal{F}^{[d]}$ is tamely ramified on $H = \tilde{C}_{\mathfrak{m}}^{(d)} \setminus U^{(d)} = E_0 \times_{C^{(d)}} \tilde{C}_{\mathfrak{m}}^{(d)}$.*

From which we will deduce [Theorem 2](#).

We start by recalling the necessarily material, and then move to proving this theorem.

4.1 Etale Fundamental Groups and Tame Fundamental Groups

We recall the definition and basic properties of the étale fundamental group, following stacks project [\[Stacks, Tag 0BQ6\]](#)

Proposition 55 ([\[Stacks, Tag 0C0J\]](#)). *Let $f : X \rightarrow S$ be a flat proper morphism of finite presentation whose geometric fibres are connected and reduced. Assume S is connected and let $\bar{s}$ be a geometric point of S . Then there is an exact sequence*

$$\pi_1(X_{\bar{s}}) \rightarrow \pi_1(X) \rightarrow \pi_1(S) \rightarrow 1$$

of fundamental groups.

Corollary 56. *Let $f : X \rightarrow S$ be a proper smooth morphism of finite presentation whose geometric fibres are connected. Assume S is connected and let $\bar{s}$ be a geometric point of S . Then there is an exact sequence*

$$\pi_1(X_{\bar{s}}) \rightarrow \pi_1(X) \rightarrow \pi_1(S) \rightarrow 1$$

of fundamental groups.

4.2 Generalized Picard Scheme

We recall the notion of generalized Jacobian varieties and study their fundamental properties. The material presented here is primarily adapted from [\[Gui19\]](#) and [\[Tak19\]](#). For further background on the general theory of abelian varieties and Jacobians, the reader may also consult [\[Mil08\]](#). Let S be a scheme and let C be a projective smooth S -scheme whose geometric fibers are connected and of dimension 1. Let $\mathfrak{m}$ be a modulus on C , defined as an effective Cartier divisor of C/S (i.e., a closed subscheme of C which is finite flat of finite presentation over S). We denote the projection $C \times_S T \rightarrow T$ by pr for any S -scheme T .

The Functor of Points

Let d be an integer. For an S -scheme T , we consider the set of data $(\mathcal{L}, \psi)$ where:

- $\mathcal{L}$ is an invertible sheaf of degree d on C_T .

- $\psi : \mathcal{O}_{\mathfrak{m}_T} \xrightarrow{\sim} \mathcal{L}|_{\mathfrak{m}_T}$ is a trivialization of $\mathcal{L}$ along the modulus.

Two such pairs $(\mathcal{L}, \psi)$ and $(\mathcal{L}', \psi')$ are said to be isomorphic if there exists an isomorphism of invertible sheaves $f : \mathcal{L} \rightarrow \mathcal{L}'$ such that the following diagram commutes:

$$\begin{array}{ccc} & \mathcal{O}_{\mathfrak{m}_T} & \\ \psi' \swarrow & & \searrow \psi \\ \mathcal{L}'|_{\mathfrak{m}_T} & \xrightarrow{f|_{\mathfrak{m}_T}} & \mathcal{L}|_{\mathfrak{m}_T} \end{array}$$

We define the presheaf $\text{Pic}_{C, \mathfrak{m}}^{d, \text{pre}}$ on Sch/S by assigning to T the set of isomorphism classes of such pairs. Let $\text{Pic}_{C, \mathfrak{m}}^d$ denote the étale sheafification of this presheaf.

Representability and Structure

The fundamental properties of this functor are as follows:

1. $\text{Pic}_{C, \mathfrak{m}}^d$ is represented by an S -scheme. (Note: If $\mathfrak{m}$ is faithfully flat over S , the presheaf is already a étale sheaf).
2. $\text{Pic}_{C, \mathfrak{m}}^0$ is a smooth commutative group S -scheme with geometrically connected fibers, referred to as the *generalized Jacobian variety* of C with modulus $\mathfrak{m}$.
3. For any d , $\text{Pic}_{C, \mathfrak{m}}^d$ is a $\text{Pic}_{C, \mathfrak{m}}^0$ -torsor.

In the case where $\mathfrak{m} = 0$, we recover the standard Jacobian variety, denoted simply as Pic_C^d .

Relation to the Standard Jacobian

We now examine the behavior of the generalized Picard scheme under the variation of the modulus. By viewing the structure along the modulus as an additional rigidification, we obtain natural transition maps corresponding to the inclusion of moduli.

Let $\mathfrak{m}_1$ and $\mathfrak{m}_2$ be moduli such that $\mathfrak{m}_1 \subset \mathfrak{m}_2$. There exists a natural map

$$\text{Pic}_{C, \mathfrak{m}_2}^d \rightarrow \text{Pic}_{C, \mathfrak{m}_1}^d$$

obtained by restricting the isomorphism ψ . Since $\mathfrak{m}_2$ is a finite S -scheme, this map is a surjection as a morphism of étale sheaves. In particular, for any modulus $\mathfrak{m}$, there is a natural surjective morphism of étale sheaves:

$$\text{Pic}_{C, \mathfrak{m}}^d \rightarrow \text{Pic}_C^d.$$

Local Freeness and Base Change

Let $\mathfrak{m}$ be a modulus which is everywhere strictly positive. Let g denote the genus of C , which is a locally constant function on S . We restrict our attention to degrees d satisfying the condition:

$$d \geq \max\{2g - 1 + \deg \mathfrak{m}, \deg \mathfrak{m}\}. \quad (6)$$

Assuming S is quasi-compact, such a d always exists.

Fix an integer d satisfying the condition above. Let T be an S -scheme and let $\mathcal{L}$ be an invertible sheaf of degree d on C_T . One can show that the pushforwards $\mathrm{pr}_*\mathcal{L}$ and $\mathrm{pr}_*\mathcal{L}(-\mathfrak{m})$ are locally free sheaves and their formations commute with any base change. Explicitly, for any morphism of S -schemes $f : T' \rightarrow T$, the base change morphisms are isomorphisms:

$$f^*\mathrm{pr}_*\mathcal{L} \xrightarrow{\sim} \mathrm{pr}_*f^*\mathcal{L}$$

and

$$f^*\mathrm{pr}_*(\mathcal{L}(-\mathfrak{m})) \xrightarrow{\sim} \mathrm{pr}_*f^*(\mathcal{L}(-\mathfrak{m})).$$

In particular, following [Gui19], if $\mathcal{L}$ is invertible $\mathcal{O}_C$ -module with degree d satisfying 6 on each fiber of f then, $\mathrm{pr}_*\mathcal{L}$ is a locally free $\mathcal{O}_S$ -module of rank $d - g + 1$.

For further background and verification of these constructions, we refer the reader to Milne's notes on abelian Varieties ([Mil08]).

4.3 The Abel-Jacobi Morphism and its Fibers

Let $U = C \setminus \mathfrak{m}$ be the complement of the modulus in C . The effective cartier divisors of degree d which are prime to $\mathfrak{m}$ are parameterized by the symmetric power $\mathrm{Sym}_S^d(U) = U^{(d)}$ over S (See [Gui19] Proposition 4.12, [Mil08] Theorem 3.13). For any such divisor $D \in U^{(d)}$, the associated line bundle $\mathcal{O}_C(D)$ admits a canonical trivialization along $\mathfrak{m}$. Specifically, the canonical section 1_D is regular and non-vanishing on $\mathfrak{m}$ because $\mathrm{supp}(D) \cap \mathrm{supp}(\mathfrak{m}) = \emptyset$. This section restricts to a nowhere-vanishing section on the subscheme $\mathfrak{m}$, thereby determining a trivialization $\psi_D^{-1} : \mathcal{O}_C(D)|_{\mathfrak{m}} \xrightarrow{\sim} \mathcal{O}_{\mathfrak{m}}$. This is done functorially in families, yielding a morphism from the symmetric power to the generalized Picard scheme (over S):

$$\Phi_d : U^{(d)} \rightarrow \mathrm{Pic}_{C,\mathfrak{m}}^d, \quad D \mapsto [(\mathcal{O}_C(D), \psi_D)], \quad (7)$$

When $\mathfrak{m} = 0$, $d \geq \max\{2g - 1, 0\}$ and C admits a section over S , $C^{(d)}$ is a projective space bundle over Pic_C^d . It is proper, surjective with geometrically connected fibers.

Guignard ([Gui19] Theorem 4.14) proves that for $\mathfrak{m} > 0$ and d satisfying (6), the Abel-Jacobi morphism Φ_d is surjective smooth of relative dimension $d - \deg \mathfrak{m} - g + 1$, with geometrically connected fibers.

When $S = \mathrm{Spec}(k)$, the geometric-fibers of Φ_d are well understood:

Theorem 57. *Assuming $S = \mathrm{Spec}(k)$ and $d \geq \max\{2g - 1 + \deg \mathfrak{m}, \deg \mathfrak{m}\}$. Then, the geometric-fibers of the Abel-Jacobi morphism*

$$\Phi_d : U^{(d)} \rightarrow \mathrm{Pic}_{C,\mathfrak{m}}^d$$

over any point are isomorphic to

$$\begin{cases} \mathbb{A}_{k^{sep}}^{d-\deg \mathfrak{m}-g+1} & \text{if } m > 0 \\ \mathbb{P}_{k^{sep}}^{d-g} & \text{if } m = 0 \end{cases}$$

In both cases Φ_d is a fibration in affine spaces or projective spaces, depending on whether $\mathfrak{m}$ is non-zero or zero.

Proof. see [Ten15] Propositions 3.13-3.14, or [T6t11] Prop 2.1.4:

□

4.4 Compactification of Blowup of Symmetric Powers of a Curve

We recall that our objective is to descend the local system $\mathcal{F}^{[d]}$ from $U^{(d)}$ to $\text{Pic}_{C,\mathfrak{m}}^d$ along the Abel-Jacobi map Φ_d :

$$\begin{array}{ccc} \mathcal{F}^{[d]} & & \\ \downarrow & & \\ U^{(d)} & \xrightarrow{\Phi_d} & \text{Pic}_{C,\mathfrak{m}}^d \end{array}$$

(Here, the purple arrow emphasizes that the morphism is of sheaves on the étale site).

However, we encounter an obstruction: in the case we are considering ($\mathfrak{m} > 0$), the fibers of Φ_d are affine spaces (of the same degree) rather than the better-behaved projective spaces. This hint that a solution to this problem is to compactify the morphism to yield projective fibers.

This section describes the result of the compactification constructed by [Tak19] via the method of blowup.

Let $\mathfrak{m} = \sum_{i=1}^n k_P P$ with $\deg P = d_P$ be a modulus on C , and let d satisfy (6). Takeuchi ([Tak19]) defines $Z_0 = Z_0(\mathfrak{m}, d)$ as the closed subscheme of $C^{(d)}$ defined by the map $C^{(d-\deg \mathfrak{m})} \rightarrow C^{(d)}$ adding $\mathfrak{m}$. He also defines $X_{\mathfrak{m},d}$ as the blowup of $C^{(d)}$ along Z_0 . Let $E_0 = E_{\mathfrak{m},d} = Z_0(\mathfrak{m}, d) \times_{C^{(d)}} X_{\mathfrak{m},d}$ be the exceptional divisor of the blowup. It is irreducible of codimension 1, and we let $\eta_0 = \eta_{\mathfrak{m},d}$ be its generic point.

Diagrammatically:

$$\begin{array}{ccc} \overline{\{\eta_0\}} = E_0 & \longrightarrow & X_{\mathfrak{m},d} \\ \downarrow & & \downarrow \pi \\ Z_0 & \xrightarrow{c.i} & C^{(d)} \end{array}$$

Incorporating $U^{(d)}$, the local system $\mathcal{F}^{[d]}$ and the Abel-Jacobi map, we have:

$$\begin{array}{ccccc} & & & \mathcal{F}^{[d]} & \\ & & & \downarrow & \\ \overline{\{\eta_0\}} = E_0 & \longrightarrow & X_{\mathfrak{m},d} & \longleftrightarrow & U^{(d)} \xrightarrow{\Phi_d} \text{Pic}_{C,\mathfrak{m}}^d \\ \downarrow & & \downarrow \pi & \swarrow & \\ Z_0 & \xrightarrow{c.i} & C^{(d)} & & \end{array}$$

In Section 3 of [Tak19] Takeuchi constructs, for d satisfying (6) a compactification denoted by $\tilde{C}_{\mathfrak{m}}^{(d)}$ and proves the following:

Theorem 58 (Takeuchi). *The scheme $\tilde{C}_{\mathfrak{m}}^{(d)}$ is an open subscheme of $X_{\mathfrak{m},d}$ containing $U^{(d)}$. The morphism $\Phi_d : U^{(d)} \rightarrow \text{Pic}_{C,\mathfrak{m}}^d$ extends to a morphism $\tilde{\Phi}_d : \tilde{C}_{\mathfrak{m}}^{(d)} \rightarrow \text{Pic}_{C,\mathfrak{m}}^d$ which makes $\tilde{C}_{\mathfrak{m}}^{(d)}$ a projective space bundle over $\text{Pic}_{C,\mathfrak{m}}^d$. Furthermore, the complement of $U^{(d)}$ in $\tilde{C}_{\mathfrak{m}}^{(d)}$ is isomorphic to the fiber product $E_0 \times_{C^{(d)}} \tilde{C}_{\mathfrak{m}}^{(d)}$.*

Diagrammatically we have:

$$\begin{array}{ccccc}
& & & & \mathcal{F}^{[d]} \\
& & & & \downarrow \\
E_0 \times_{C^{(d)}} \tilde{C}_m^{(d)} & \longrightarrow & \tilde{C}_m^{(d)} & \longleftarrow & U^{(d)} \\
\downarrow & & \downarrow & \searrow \tilde{\Phi}_d & \downarrow \Phi_d \\
\overline{\{\eta_0\}} = E_0 & \longrightarrow & X_{m,d} & & \text{Pic}_{C,m}^d \\
\downarrow & & \downarrow \pi & & \\
Z_0 & \xrightarrow{c.i} & C^{(d)} & &
\end{array}$$

Also note that $E_0 \times_{C^{(d)}} \tilde{C}_m^{(d)} = Z_0 \times_{C^{(d)}} \tilde{C}_m^{(d)}$

4.5 Proof of Theorem 54

In what follows we reduce Theorem 54 to Theorem 38

4.5.1 Reduction Lemmas

The following lemma is adapted from [Tak19] (Lemma 4.1)

Lemma 59. *Let C be a projective, smooth, and geometrically connected curve over a perfect field k . Let $\mathfrak{m} = \sum_{i=1}^r k_i P_i$ be an effective divisor where $P_1, \dots, P_r$ are distinct closed points. Let $U = C \setminus \mathfrak{m}$ and let $d \geq \deg \mathfrak{m}$.*

Suppose $\mathfrak{n}_1, \dots, \mathfrak{n}_l$ are pairwise coprime submoduli of $\mathfrak{m}$ such that $\mathfrak{m} = \sum_{j=1}^l \mathfrak{n}_j$. Consider the summation morphism:

$$\pi : C^{(\deg \mathfrak{n}_1)} \times_k \dots \times_k C^{(\deg \mathfrak{n}_l)} \times_k C^{(d - \deg \mathfrak{m})} \longrightarrow C^{(d)}$$

defined by $(D_1, \dots, D_l, D_{\text{extra}}) \mapsto \sum_{j=1}^l D_j + D_{\text{extra}}$.

Then π is étale at the generic point of the closed subvariety

$$V = \{\mathfrak{n}_1\} \times_k \dots \times_k \{\mathfrak{n}_l\} \times_k C^{(d - \deg \mathfrak{m})}$$

inside the domain $C^{(\deg \mathfrak{n}_1)} \times_k \dots \times_k C^{(\deg \mathfrak{n}_l)} \times_k C^{(d - \deg \mathfrak{m})}$.

Proof. We may assume that k is algebraically closed (hence $\deg P_i = 1$ for all i). By miracle flatness π is flat, it is quasi-finite and projective as a map between projective spaces. so we conclude π is finite and flat. It is enough to show that there exists a closed point Q of $\mathfrak{n}_1 + \dots + \mathfrak{n}_l + C^{(d - \deg \mathfrak{m})} \subset C^{(d)}$ over which there are $\deg \pi$ points on $C^{(\mathfrak{n}_1)} \times_k \dots \times_k C^{(\mathfrak{n}_l)} \times_k C^{(d - \deg \mathfrak{m})}$. (Because it will be unramified at this point and thus also at the generic point of V .) Choose Q as a point corresponding to a divisor $\mathfrak{n}_1 + \dots + \mathfrak{n}_l + P_{r+1} + \dots + P_{r+d - \deg \mathfrak{m}}$, where $P_1, \dots, P_{r+d - \deg \mathfrak{m}}$ are distinct points of $U(k)$. $\square$

Corollary 60. *The morphism $C^{(\deg \mathfrak{m})} \times_k C^{(d - \deg \mathfrak{m})} \xrightarrow{\pi} C^{(d)}$ is finite flat everywhere, and étale at the generic point of the closed subvariety $Z_{\mathfrak{m}} \times_k C^{(d - \deg \mathfrak{m})} \subset C^{(\deg \mathfrak{m})} \times_k C^{(d - \deg \mathfrak{m})}$.*

Following from this, we look at the following diagram, coming from the flat base change $C^{(d-\deg \mathfrak{m})} \rightarrow \text{Spec}(k)$ (Proposition 19) of (4):

$$\begin{array}{ccccc}
E_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})} & \longrightarrow & X_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})} & & \mathcal{F}^{[\deg \mathfrak{m}]} \times_k C^{(d-\deg \mathfrak{m})} \\
\downarrow & & \downarrow & \nwarrow & \downarrow \\
Z_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})} & \longrightarrow & C^{(\deg \mathfrak{m})} \times_k C^{(d-\deg \mathfrak{m})} & & U^{(\deg \mathfrak{m})} \times_k C^{(d-\deg \mathfrak{m})}
\end{array}$$

Note that $U^{(\deg \mathfrak{m})} \times_k C^{(d-\deg \mathfrak{m})}$ is dense open subscheme of $X_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})}$, And $E_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})}$ is a prime divisor of $X_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})}$. Hence it is well defined question according to Definition 39 to ask whether $\mathcal{F}^{[\deg \mathfrak{m}]} \times_k C^{(d-\deg \mathfrak{m})}$ is tamely ramified at the generic point θ of $E_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})}$.

Lemma 61. *If $\mathcal{F}^{[\deg \mathfrak{m}]}$ is tamely ramified at $\eta_{\mathfrak{m}}$, then $\mathcal{F}^{[\deg \mathfrak{m}]} \times_k C^{(d-\deg \mathfrak{m})}$ is tamely ramified at the generic point θ of $E_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})}$.*

Proof. This follows from [Stacks, Tag 0EYD] □

Replacing $C^{(d-\deg \mathfrak{m})}$ with the dense open subscheme $U^{(d-\deg \mathfrak{m})} \subset C^{(d-\deg \mathfrak{m})}$, we get that the G -torsor $p_1^{-1}\mathcal{P}^{[\deg \mathfrak{m}]}$ ($\mathcal{P}$ corresponds to $\mathcal{F}$ under Proposition 8) is tamely ramified at θ the generic point of $E_{\mathfrak{m}} \times_k U^{(d-\deg \mathfrak{m})} \subset U^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})}$, where $p_1 : U^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})} \rightarrow U^{(\deg \mathfrak{m})}$ is the projection to the first factor.

Looking at the second projection $p_2 : X_{\mathfrak{m}} \times_k U^{(d-\deg \mathfrak{m})} \rightarrow U^{(d-\deg \mathfrak{m})}$, and the fact that $\mathcal{P}^{[d-\deg \mathfrak{m}]}$ is étale on $U^{(d-\deg \mathfrak{m})}$ we get that $p_2^{-1}\mathcal{P}^{[d-\deg \mathfrak{m}]} = X_{\mathfrak{m}} \times_k \mathcal{P}^{[d-\deg \mathfrak{m}]}$ is étale on $X_{\mathfrak{m}} \times_k U^{(d-\deg \mathfrak{m})}$. Hence, its restriction to $U^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})}$ is unramified at θ .

Thus, by the following lemma, we conclude that $\mathcal{P}^{[\deg \mathfrak{m}]} \boxtimes \mathcal{P}^{[d-\deg \mathfrak{m}]} = p_1^{-1}\mathcal{P}^{[\deg \mathfrak{m}]} \otimes^G p_2^{-1}\mathcal{P}^{[d-\deg \mathfrak{m}]}$ is tamely ramified at θ the generic point of $E_{\mathfrak{m}} \times_k U^{(d-\deg \mathfrak{m})}$.

Lemma 62. *Let $X \rightarrow \text{Spec } k$ be a scheme over a field k , and let $\mathcal{P}_1, \mathcal{P}_2$ be two G -torsors on U_{et} . Let ξ be the generic point of a prime divisor $D \subset X$. If $\mathcal{P}_1$ is tamely ramified at ξ , and $\mathcal{P}_2$ is unramified at ξ , then the product $\mathcal{P}_1 \otimes^G \mathcal{P}_2$ is tamely ramified at ξ .*

Proof. This follows from Lemma 43. □

Combining this with Corollary 60 we get

Corollary 63. *Let $\mathfrak{m}$ be a modulus as above, and let $\eta_{\mathfrak{m}}$ be the generic point of $E_{\mathfrak{m}}$. Let $\mathcal{P}$ be a G -torsor on U_{et} with ramification bounded by $\mathfrak{m}$. Assume $\mathcal{P}^{[\deg \mathfrak{m}]}$ is tamely ramified at $\eta_{\mathfrak{m}}$. Then $\mathcal{P}^{[\deg \mathfrak{m}]} \boxtimes \mathcal{P}^{[d-\deg \mathfrak{m}]}$ is tamely ramified at the generic point θ of $E_{\mathfrak{m}} \times_k C^{(d-\deg \mathfrak{m})} \subset C^{(\deg \mathfrak{m})} \times_k C^{(d-\deg \mathfrak{m})}$, and $\mathcal{P}^{[d]}$ is tamely ramified at the generic point $\eta_0 = \eta_{\mathfrak{m},d}$ of $E_0 = E_{\mathfrak{m},d}$.*

Proof. The first assertion, that $\mathcal{P}^{[\deg \mathfrak{m}]} \boxtimes \mathcal{P}^{[d-\deg \mathfrak{m}]}$ is tamely ramified at θ , follows from the preceding discussion. Thus, it remains to show that $\mathcal{P}^{[d]}$ is tamely ramified at η_0 .

Consider the blowup diagram defining $X_{\mathfrak{m},d}$:

$$\begin{array}{ccc}
\overline{\{\eta_0\}} = E_0 & \longrightarrow & X_{\mathfrak{m},d} \\
\downarrow & & \downarrow \pi \\
Z_0 & \xrightarrow{c.i} & C^{(d)}
\end{array}$$

By performing a base change along the flat addition map $+: C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})} \rightarrow C^{(d)}$, we obtain the following commutative diagram:

$$\begin{array}{ccccc}
& & (C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})}) \times_{C^{(d)}} E_0 & \longrightarrow & \overline{\{\eta_0\}} = E_0 \\
& & \downarrow & & \downarrow \\
& & (C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})}) \times_{C^{(d)}} X_{\mathfrak{m},d} & \longrightarrow & X_{\mathfrak{m},d} \\
& & \downarrow & & \downarrow \pi \\
\mathfrak{m} \times_k U^{(d-\deg \mathfrak{m})} & \xrightarrow{c.i} & C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})} & \xrightarrow{+} & C^{(d)}
\end{array}$$

Since blowups commute with flat base change, and the inverse image of the center Z_0 under the map $+$ is $\mathfrak{m} \times_k U^{(d-\deg \mathfrak{m})}$, the scheme $(C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})}) \times_{C^{(d)}} X_{\mathfrak{m},d}$ is the blowup of $C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})}$ along $\mathfrak{m} \times_k U^{(d-\deg \mathfrak{m})}$. This, in turn, is isomorphic to the base change of the blowup $X_{\mathfrak{m}}$ (of $C^{(\deg \mathfrak{m})}$ along $\mathfrak{m}$) via the (flat) projection $C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})} \rightarrow C^{(\deg \mathfrak{m})}$.

Assembling these facts, we obtain the following Cartesian square:

$$\begin{array}{ccccc}
E_{\mathfrak{m}} \times U^{(d-\deg \mathfrak{m})} & \xrightarrow{\tilde{+}} & \overline{\{\eta_0\}} = E_0 \\
\downarrow & & \downarrow \\
X_{\mathfrak{m}} \times_k U^{(d-\deg \mathfrak{m})} & \xrightarrow{\tilde{+}} & X_{\mathfrak{m},d} \\
\downarrow & & \downarrow \pi \\
\mathfrak{m} \times_k U^{(d-\deg \mathfrak{m})} & \xrightarrow{c.i} & C^{(\deg \mathfrak{m})} \times_k U^{(d-\deg \mathfrak{m})} & \xrightarrow{+} & C^{(d)}
\end{array}$$

The point θ defined in the Corollary is the generic point of $E_{\mathfrak{m}} \times_k U^{(d-\deg \mathfrak{m})}$. Let η be the generic point of $\mathfrak{m} \times_k U^{(d-\deg \mathfrak{m})}$. By [Corollary 60](#), the map $+$ is étale at η . Consequently, the lifted map $\tilde{+}$ is étale at θ . Given the isomorphism $(+^{-1})(\mathcal{P}^{[d]}) \cong \mathcal{P}^{[\deg \mathfrak{m}]} \boxtimes \mathcal{P}^{[d-\deg \mathfrak{m}]}$ from [Proposition 24](#), the tame ramification of the box product at θ descends to the tame ramification of $\mathcal{P}^{[d]}$ at η_0 by applying [Lemma 44](#). □

Lemma 64. *Let $\mathfrak{n}_1, \mathfrak{n}_2 \subset \mathfrak{m}$ be two coprime sub moduli of $\mathfrak{m}$. Assume $\mathcal{P}^{[\deg \mathfrak{n}_1]}, \mathcal{P}^{[\deg \mathfrak{n}_2]}$ are at most tamely ramified at $\eta_{\mathfrak{n}_1}, \eta_{\mathfrak{n}_2}$ respectively. Then $\mathcal{P}^{[\deg \mathfrak{n}_1 + \deg \mathfrak{n}_2]}$ is at most tamely tamified at $\eta_{\mathfrak{n}_1 + \mathfrak{n}_2}$.*

Proof. It follows from [Proposition 53](#) and [Proposition 24](#) □

4.6 Proof of Theorem 54

Proof of Theorem 54. Let $\mathcal{F}$ be as in [Theorem 54](#), $\mathfrak{m} = \sum_{i=1}^n k_P P$ with $\deg P = d_P$. Then by [Theorem 3](#) for every $\mathfrak{n} \subset \mathfrak{m}$ of the form $\mathfrak{n} = k_P P$, $\mathcal{F}^{[\deg \mathfrak{n}]}$ is at most tamely ramified at $\eta_{\mathfrak{n}}$. By [Lemma 64](#), $\mathcal{F}^{[\deg \mathfrak{m}]}$ is then at most tamely ramified at $\eta_{\mathfrak{m}}$. And thus by [Corollary 63](#) $\mathcal{F}^{[d]}$ is tamely ramified at the generic point η_0 of E_0 □

5 Proof of Theorem 2

We work over $S = \text{Spec } k$, for k perfect.

By Proposition 8, its equivalent to prove:

Theorem 65. *Let $G = \Lambda^\times$ be a finite abelian group (Λ as before), and let $\mathcal{P}$ be a G -torsor on U , with ramification bounded by $\mathfrak{m}$. Then, for sufficiently large integer d , there exists a unique (up to isomorphism) G -torsor $\mathcal{Q}_d$ on $\text{Pic}_{C,\mathfrak{m}}^d$, such that the pullback of $\mathcal{Q}_d$ by Φ_d is isomorphic to $\mathcal{P}^{[d]}$.*

Proof. We divide the proof into two cases, when $\mathfrak{m} = 0$ and when $\mathfrak{m} > 0$.

Case 1: $\mathfrak{m} = 0$. By Section 4.3, for d large enough, the Abel-Jacobi morphism $\Phi_d : C^{(d)} \rightarrow \text{Pic}_C^d$ is proper surjective and smooth, with geometrically connected fibers, each isomorphic to $\mathbb{P}_{k^{sep}}^{d-g}$. Hence by Corollary 56 it induces an exact sequence of etale fundamental groups:

$$\pi_1^{et}(\mathbb{P}_{k^{sep}}^{d-g}) \rightarrow \pi_1^{et}(C^{(d)}) \rightarrow \pi_1^{et}(\text{Pic}_C^d) \rightarrow 1$$

But $\mathbb{P}_{k^{sep}}^{d-g}$ is simply connected ([Ten15] Example 4.9, [Töt11] Example 1.4.12), hence its etale fundamental group is trivial, and we get an isomorphism of etale fundamental groups:

$$\pi_1^{et}(C^{(d)}) \cong \pi_1^{et}(\text{Pic}_C^d)$$

Implying the theorem in this case.

Case 2: $\mathfrak{m} > 0$. In this case by Theorem 58, for d large enough, the Abel-Jacobi morphism $\Phi_d : U^{(d)} \rightarrow \text{Pic}_{C,\mathfrak{m}}^d$ extends to a proper surjective and smooth map, with geometrically connected fibers isomorphic to projective spaces,

$$\tilde{\Phi}_d : \tilde{C}_{\mathfrak{m}}^{(d)} \rightarrow \text{Pic}_{C,\mathfrak{m}}^d$$

Hence we get an isomorphism of etale fundamental groups:

$$\pi_1^{et}(\tilde{C}_{\mathfrak{m}}^{(d)}) \cong \pi_1^{et}(\text{Pic}_{C,\mathfrak{m}}^d)$$

By Theorem 54, $\mathcal{F}^{[d]}$ is tamely ramified on the boundary divisor $H = \tilde{C}_{\mathfrak{m}}^{(d)} \setminus U^{(d)}$.

Thus, by Lemma 66 below, we have: $\mathcal{F}^{[d]}$ extends to a locally constant sheaf $\tilde{\mathcal{F}}^{[d]}$ on $\tilde{C}_{\mathfrak{m}}^{(d)}$, which by the isomorphism of etale fundamental groups above, corresponds to a unique locally constant sheaf $\mathcal{G}_d$ on $\text{Pic}_{C,\mathfrak{m}}^d$, such that $\tilde{\Phi}_d^* \mathcal{G}_d \cong \tilde{\mathcal{F}}^{[d]}$. Restricting back to $U^{(d)}$, we get $\Phi_d^* \mathcal{G}_d \cong \mathcal{F}^{[d]}$, as required. $\square$

Lemma 66. *If $\mathcal{F}^{[d]}$ is a locally constant sheaf on $U^{(d)}$ which is tamely ramified along the boundary divisor $H = \tilde{C}_{\mathfrak{m}}^{(d)} \setminus U^{(d)}$, then $\mathcal{F}^{[d]}$ extends to a locally constant sheaf $\tilde{\mathcal{F}}^{[d]}$ on $\tilde{C}_{\mathfrak{m}}^{(d)}$.*

Proof. The lemma we are referencing above can be proved in two routes:

Route 1 - Showing $\ker\{\pi_1^{ab}(U^{(d)}) \rightarrow \pi_1^{ab}(\text{Pic}_{C,\mathfrak{m}}^d)\}$ is pro- p group.

Route 2 - Showing

$$\pi_1^{t,ab}(U^{(d)}) \rightarrow \pi_1^{t,ab}(\text{Pic}_{C,\mathfrak{m}}^d)$$

is isomorphism to its image. here one needs to be precise.

$\square$

6 Setup and notation

Let $\mathcal{P} \rightarrow U$ be a G -torsor over $U = C \setminus \mathfrak{m}$, where G is a finite abelian group, $H \trianglelefteq G$, and C is a smooth projective curve. Fix $p_\infty \in \mathfrak{m}$ and an integer d . Assume the G torsor $\mathcal{P} \rightarrow U$ has ramification strictly bounded by d at p_∞ . (i.e. $< d$) according to the definition in [Definition 40](#) (i.e. the higher d 'th-ramification group of the extension of DVRs at p_∞ is trivial).

Recall the four quotient schemes, defined in: [Section 2.4](#)

$$\mathcal{P}^{[d]}_H, \quad \mathcal{P}^{[d]}_G, \quad (\mathcal{P}_{G/H})^{(d)}, \quad (\mathcal{P}_{G/H})^{[d]},$$

together with the canonical Cartesian square

$$\begin{array}{ccc} \mathcal{P}^{[d]}_H & \xrightarrow{H} & (\mathcal{P}_{G/H})^{(d)} \\ \downarrow & & \downarrow q \\ \mathcal{P}^{[d]}_G & \xrightarrow{H} & (\mathcal{P}_{G/H})^{[d]} \\ & \searrow G & \downarrow G/H \\ & & U^{(d)}, \end{array}$$

in which the arrows labelled H , G/H and G are torsors under the indicated finite constant groups (in particular, *étale*), and the upper square is Cartesian ([Proposition 23](#)).

The G/H -torsor $\mathcal{P}_{G/H} \rightarrow U$ compactifies to a finite map of smooth projective curves $f: C' \rightarrow C$. Pick $p'_\infty \in f^{-1}(p_\infty)$

$$f^{(d)}: C'^{(d)} \rightarrow C^{(d)}$$

for the induced map of d -fold symmetric powers; it sends $d \cdot p'_\infty \mapsto d \cdot p_\infty$. Let

$$\pi_C: X_C \rightarrow C^{(d)}, \quad \pi_{C'}: X_{C'} \rightarrow C'^{(d)}$$

be the blowups at $d \cdot p_\infty$ and $d \cdot p'_\infty$ respectively, with exceptional divisors $E_C, E_{C'}$ and generic points $\eta_C, \eta_{C'}$. The DVRs of these generic points are

$$V_C := \mathcal{O}_{X_C, \eta_C} \subset K(C^{(d)}), \quad V_{C'} := \mathcal{O}_{X_{C'}, \eta_{C'}} \subset K(C'^{(d)}).$$

7 The master diagram

The torsor tower over the open base $U^{(d)}$ embeds into the projective geometry of $C^{(d)}$ and $C'^{(d)}$, which in turn admit canonical blowups at the points at infinity. We assemble all of this into a single commutative diagram.

$$\begin{array}{ccccccc}
\mathcal{P}^{[d]}_H & \xrightarrow{H} & (\mathcal{P}_{G/H})^{(d)} & \xrightarrow{\iota'} & C'^{(d)} & \xleftarrow{\pi_{C'}} & X_{C'} \\
\downarrow & & \downarrow q & & \downarrow f^{(d)} & & \downarrow f_X \\
\mathcal{P}^{[d]}_G & \xrightarrow{H} & (\mathcal{P}_{G/H})^{[d]} & & & & \\
& \searrow G & \downarrow G/H & & & & \\
& & U^{(d)} & \xrightarrow{\iota} & C^{(d)} & \xleftarrow{\pi_C} & X_C
\end{array}$$

Reading the diagram:

- The **left block** (columns 1–2) is the torsor tower of [Proposition 23](#): H -torsors horizontally, the G/H -torsor on the lower right, and the composite G -torsor $\mathcal{P}^{[d]}_G \rightarrow U^{(d)}$.
- The **middle block** (column 3) records the compactifications. The map ι is the canonical open immersion $U^{(d)} \hookrightarrow C^{(d)}$. The map ι' is the open immersion $(\mathcal{P}_{G/H})^{(d)} \hookrightarrow C'^{(d)}$ obtained from compactifying $\mathcal{P}_{G/H} \subset C'$. The vertical $f^{(d)}$ is the finite morphism of symmetric powers; restricting along ι' recovers the canonical map $(\mathcal{P}_{G/H})^{(d)} \rightarrow U^{(d)}$, which is the composite of q with the G/H -torsor.
- The **right block** (column 4) records the blowups at the diagonal points at infinity. The dashed arrow $f_X: X_{C'} \dashrightarrow X_C$ is the rational lift of $f^{(d)}$, defined on the strict transform; on generic points of the exceptional divisors it sends $\eta_{C'} \mapsto \eta_C$ so we have induced extension of DVRs $V_C \hookrightarrow V_{C'}$.

8 Unramifiedness of $\mathcal{P}^{[d]}_G$ at η_C

Theorem 67. *Suppose*

(H1) $(\mathcal{P}_{G/H})^{[d]} \rightarrow U^{(d)}$ *is unramified at η_C ; and*

(H2) $\mathcal{P}^{[d]}_H \rightarrow (\mathcal{P}_{G/H})^{(d)}$ *is unramified at $\eta_{C'}$.*

(H3) $G = \mathbb{Z}/p^r\mathbb{Z}$ *is cyclic of prime power order.*

Then $\mathcal{P}^{[d]}_G \rightarrow U^{(d)}$ is unramified at η_C .

8.1 Proof Attempts

8.1.1 Attempt 1

Proof attempt (Galois-closure version). Setup. Write

$$K := K(U^{(d)}) = K(C^{(d)}), \quad M := K((\mathcal{P}_{G/H})^{[d]}), \quad K' := K((\mathcal{P}_{G/H})^{(d)}) = K(C'^{(d)}),$$

$$L := K(\mathcal{P}^{[d]}_G), \quad L_H := K(\mathcal{P}^{[d]}_H).$$

The Galois closure of all these fields over K is $K(\mathcal{P}^{\times d})$. Indeed, $\mathcal{P}^{[d]_G}, \mathcal{P}^{[d]_H}, (\mathcal{P}_{G/H})^{(d)}, (\mathcal{P}_{G/H})^{[d]}$ are defined in [Section 2.4](#) as the quotients of $\mathcal{P}^{\times d}$ by the subgroups

$$K_G \rtimes S_d, \quad K_H \rtimes S_d, \quad H^d \rtimes S_d, \quad (H^d + K_G) \rtimes S_d$$

of the big group $\Gamma := G^d \rtimes S_d$ acting on $\mathcal{P}^{\times d}$; and $U^{(d)}$ is the quotient by $G^d \rtimes S_d$. So

$$N := K(\mathcal{P}^{\times d})/K \quad \text{is Galois with group } \Gamma,$$

and the various subfields correspond to the subgroups above ([Proposition 23](#)). All five intermediate fields above are fixed by the respective subgroups; in particular, the bottom map $U^{(d)} \leftarrow N$ has Galois group Γ , with normal subgroups giving the Galois subextensions

$$\text{Gal}(L/K) = G^d \rtimes S_d / K_G \rtimes S_d \cong G, \quad \text{Gal}(M/K) \cong G/H,$$

$$\text{Gal}(L_H/K') \cong H, \quad \text{Gal}(K'/K) \cong (G/H)^d \rtimes S_d / S_d \text{ (not a group quotient if } K'/K \text{ is non-Galois),}$$

together with the canonical surjections $\text{Gal}(L/K) \twoheadrightarrow \text{Gal}(M/K)$ (summation modulo H) and $\text{Gal}(K(\mathcal{P}^{\times d})/K') \twoheadrightarrow \text{Gal}(L_H/K')$ (summation).

Goal in inertia language. Fix a discrete valuation $\widetilde{W}$ on N extending V_C , chosen so that $\widetilde{W}|_{K'} = V_{C'}$ — this is possible because both $V_{C'}$ and any extension of V_C to N restrict to V_C on K , and the Galois group Γ acts transitively on the extensions. Let

$$J := I(\widetilde{W}/V_C) \subseteq \Gamma = G^d \rtimes S_d$$

be the inertia group of $\widetilde{W}$ over V_C . Writing elements of Γ as $((g_1, \dots, g_d), \sigma)$, the surjection $\Gamma \rightarrow G$ inducing $\text{Gal}(L/K) = G$ is the summation $((g_i), \sigma) \mapsto \sum_i g_i$. We must show

$$\sum_{i=1}^d g_i = 0 \quad \text{in } G, \quad \text{for every } ((g_i), \sigma) \in J. \quad (8)$$

Equivalently, $J \subseteq K_G \rtimes S_d$. (Then the residue extension at the image of $\widetilde{W}$ in L is also separable: this is automatic from the analogous separability statements in (H1) and (H2), which we suppress.)

What the hypotheses give.

(H1) says M/K is unramified at the restriction $\widetilde{W}|_M$. Under the surjection $\Gamma \rightarrow G/H$, $((g_i), \sigma) \mapsto \sum_i g_i \bmod H$, inertia maps to inertia, so

$$\sum_{i=1}^d g_i \in H \quad \text{for every } ((g_i), \sigma) \in J. \quad (9)$$

(H2) says L_H/K' is unramified at $V_{C'} = \widetilde{W}|_{K'}$. Now $\text{Gal}(N/K') = H^d \rtimes S_d$, and the surjection $H^d \rtimes S_d \twoheadrightarrow H$ is again summation. The inertia of $\widetilde{W}$ over $V_{C'}$ in $\text{Gal}(N/K')$ equals $J \cap (H^d \rtimes S_d)$. So (H2) reads

$$J \cap (H^d \rtimes S_d) \subseteq K_H \rtimes S_d \quad \text{i.e.} \quad \sum_i h_i = 0 \text{ whenever } ((h_i), \sigma) \in J, \quad h_i \in H. \quad (10)$$

Combining equation (??), an element $((g_i), \sigma) \in J$ has $\sum g_i \in H$, hence $(g_i) \in H^d + K_G$. So we can decompose $g_i = h_i + k_i$ with $(h_i) \in H^d$ and $(k_i) \in K_G$ (the decomposition is unique modulo

$(k_0, -k_0)$ for $k_0 \in K_H$, but $\sum h_i$ is well-defined modulo K_H , hence $\sum h_i$ as an element of H is canonical). Goal equation (??) becomes

$$\sum_i h_i = 0 \quad \text{for every } ((g_i), \sigma) \in J, \quad \text{where } g_i = h_i + k_i \text{ as above.} \quad (11)$$

The case $(k_i) = 0$ (purely H -part). If $((g_i), \sigma) \in J$ with $g_i \in H$, i.e. $j \in J \cap (H^d \rtimes S_d)$, then equation (??) gives $\sum g_i = 0$, hence $\sum h_i = 0$. ✓

Where the argument stalls. For a general $((g_i), \sigma) \in J$ with nonzero $(k_i) \in K_G$, hypotheses (H1) and (H2) do *not* suffice to force $\sum h_i = 0$. Concretely, the failure mode is the following: a subgroup

$$J \subseteq (H^d + K_G) \rtimes S_d$$

satisfying equation (??) is precisely a subgroup all of whose elements “purely in H^d ” lie in K_H . But J can still contain elements $((g_i), \sigma)$ where $g_i = h_i + k_i$ with $(k_i) \neq 0$ and $\sum h_i \neq 0$ in H — e.g. a “diagonal” subgroup $\{(h, k(h)) : h \in H_0\} \subseteq H \oplus K_{G/H}$ projecting non-trivially to both factors.

Missing input. The hypotheses (H1) and (H2) alone are insufficient. To close equation (??) we need an additional input that constrains the “ $K_{G/H}$ -part” of J , equivalently the inertia of the non-Galois comparison map $q : (\mathcal{P}_{G/H})^{(d)} \rightarrow (\mathcal{P}_{G/H})^{[d]}$ at $\widetilde{W}|_{K'} = V_{C'}$ over $\widetilde{W}|_M$. A sufficient hypothesis would be (H3): the extension K'/M is unramified at $V_{C'}$ over its restriction to M , i.e. $J \cap \text{Gal}(N/M) \subseteq H^d \rtimes S_d$. Combined with (H1) and (H2), (H3) forces $J \subseteq K_G \rtimes S_d$: any $((g_i), \sigma) \in J$ has $g_i = h_i + k_i$ and (H3) gives $(k_i) = 0$, so $(h_i) \in K_H$ by the case above, so $\sum g_i = 0$. However, (H3) is *not* automatic — Lemma 45 establishes unramifiedness of q only at codimension-1 valuations of $C'^{(d)}$ supported in the boundary, whereas $V_{C'}$ is the blowup-divisor valuation at the maximally degenerate diagonal point $d \cdot p'_\infty \in C'^{(d)}$ (codimension d). In the tame cyclic test case $G/H = \mathbb{Z}/n$, a direct computation in the formal model at $d \cdot p'_\infty$ shows that the inertia of $V_{C'}$ in K'/M has order $\gcd(d, n)$ — generally non-trivial — so (H3) fails for tame compactifications when $\gcd(d, n) > 1$. The truth of Theorem 67 in this regime therefore relies on a more subtle interaction between the inertia structure at V_C and the ramification bound on $\mathcal{P} \rightarrow U$ at p_∞ , not exploited above. $\square$

9 A Wrong Path

In this section we want to explain why Theorem 52 is difficult to prove by induction on r , using the previously proven case of $G = \mathbb{Z}/p\mathbb{Z}$. (Corollary 51).

9.1 Setup and notation

Let $\mathcal{P} \rightarrow U$ be a G -torsor over $U = C \setminus \mathfrak{m}$, where G is a finite abelian group, $H \trianglelefteq G$, and C is a smooth projective curve. Fix $p_\infty \in \mathfrak{m}$ and an integer d . Assume the G torsor $\mathcal{P} \rightarrow U$ has ramification strictly bounded by d at p_∞ . (i.e. $< d$) according to the definition in Definition 40 (i.e. the higher d 'th-ramification group of the extension of DVRs at p_∞ is trivial).

Recall the four quotient schemes, defined in: Section 2.4

$$\mathcal{P}^{[d]_H}, \quad \mathcal{P}^{[d]_G}, \quad (\mathcal{P}_{G/H})^{(d)}, \quad (\mathcal{P}_{G/H})^{[d]},$$

together with the canonical Cartesian square

$$\begin{array}{ccc}
\mathcal{P}^{[d]}_H & \xrightarrow{H} & (\mathcal{P}_{G/H})^{(d)} \\
\downarrow & & \downarrow q \\
\mathcal{P}^{[d]}_G & \xrightarrow{H} & (\mathcal{P}_{G/H})^{[d]} \\
& \searrow G & \downarrow G/H \\
& & U^{(d)},
\end{array}$$

in which the arrows labelled H , G/H and G are torsors under the indicated finite constant groups (in particular, *étale*), and the upper square is Cartesian ([Proposition 23](#)).

The G/H -torsor $\mathcal{P}_{G/H} \rightarrow U$ compactifies to a finite map of smooth projective curves $f: C' \rightarrow C$. Pick $p'_\infty \in f^{-1}(p_\infty)$

$$f^{(d)}: C'^{(d)} \rightarrow C^{(d)}$$

for the induced map of d -fold symmetric powers; it sends $d \cdot p'_\infty \mapsto d \cdot p_\infty$. Let

$$\pi_C: X_C \rightarrow C^{(d)}, \quad \pi_{C'}: X_{C'} \rightarrow C'^{(d)}$$

be the blowups at $d \cdot p_\infty$ and $d \cdot p'_\infty$ respectively, with exceptional divisors $E_C, E_{C'}$ and generic points $\eta_C, \eta_{C'}$. The DVRs of these generic points are

$$V_C := \mathcal{O}_{X_C, \eta_C} \subset K(C^{(d)}), \quad V_{C'} := \mathcal{O}_{X_{C'}, \eta_{C'}} \subset K(C'^{(d)}).$$

9.2 The master diagram

The torsor tower over the open base $U^{(d)}$ embeds into the projective geometry of $C^{(d)}$ and $C'^{(d)}$, which in turn admit canonical blowups at the points at infinity. We assemble all of this into a single commutative diagram.

$$\begin{array}{ccccccc}
\mathcal{P}^{[d]}_H & \xrightarrow{H} & (\mathcal{P}_{G/H})^{(d)} & \xhookrightarrow{\iota'} & C'^{(d)} & \xleftarrow{\pi_{C'}} & X_{C'} \\
\downarrow & & \downarrow q & & \downarrow f^{(d)} & & \downarrow f_X \\
\mathcal{P}^{[d]}_G & \xrightarrow{H} & (\mathcal{P}_{G/H})^{[d]} & & & & \\
& \searrow G & \downarrow G/H & & & & \\
& & U^{(d)} & \xhookrightarrow{\iota} & C^{(d)} & \xleftarrow{\pi_C} & X_C
\end{array}$$

Reading the diagram:

- The **left block** (columns 1–2) is the torsor tower of [Proposition 23](#): H -torsors horizontally, the G/H -torsor on the lower right, and the composite G -torsor $\mathcal{P}^{[d]}_G \rightarrow U^{(d)}$.

- The **middle block** (column 3) records the compactifications. The map ι is the canonical open immersion $U^{(d)} \hookrightarrow C^{(d)}$. The map ι' is the open immersion $(\mathcal{P}_{G/H})^{(d)} \hookrightarrow C'^{(d)}$ obtained from compactifying $\mathcal{P}_{G/H} \subset C'$. The vertical $f^{(d)}$ is the finite morphism of symmetric powers; restricting along ι' recovers the canonical map $(\mathcal{P}_{G/H})^{(d)} \rightarrow U^{(d)}$, which is the composite of q with the G/H -torsor.
- The **right block** (column 4) records the blowups at the diagonal points at infinity. The dashed arrow $f_X: X_{C'} \dashrightarrow X_C$ is the rational lift of $f^{(d)}$, defined on the strict transform; on generic points of the exceptional divisors it sends $\eta_{C'} \mapsto \eta_C$ so we have induced extension of DVRs $V_C \hookrightarrow V_{C'}$.

9.3 Unramifiedness of $\mathcal{P}^{[d]G}$ at η_C

So, going back to the begining of this section, a natural way to prove [Theorem 52](#) from [Corollary 51](#) by induction on r , comes down to proving the following theorem.

Theorem 68. *Suppose*

- (H1) $(\mathcal{P}_{G/H})^{[d]} \rightarrow U^{(d)}$ *is unramified at η_C ; and*
- (H2) $\mathcal{P}^{[d]H} \rightarrow (\mathcal{P}_{G/H})^{(d)}$ *is unramified at $\eta_{C'}$.*
- (H3) $G = \mathbb{Z}/p^r\mathbb{Z}$ *is cyclic of prime power order.*

Then $\mathcal{P}^{[d]G} \rightarrow U^{(d)}$ is unramified at η_C .

And the next section is devoted to explaining why this is not an easy task, and in fact, we were not able to prove it.

9.4 Difficulties in Proving the Theorem

First, note that we can assume $\mathcal{P} \rightarrow U$ is connected, and that H is non-trivial.

- (R1) $\mathcal{P} \rightarrow U$ *is connected.* Let $\rho: \pi_1(U) \rightarrow G$ be the monodromy of $\mathcal{P}$ and put $G_0 := \text{Im}(\rho)$, $H_0 := H \cap G_0$. Pick a connected component $\mathcal{P}_0 \subseteq \mathcal{P}$; it is a connected G_0 -torsor, and $\mathcal{P}$ is the disjoint union of $[G : G_0]$ copies of $\mathcal{P}_0$ permuted by G . Since the symmetric-power quotients of [Section 2.4](#) commute with such disjoint unions, each of the four schemes in [Proposition 23](#) for $(\mathcal{P}, G, H)$ decomposes as a disjoint union of copies of the corresponding scheme for $(\mathcal{P}_0, G_0, H_0)$. Unramifiedness at η_C (resp. $\eta_{C'}$) is detected componentwise, so (H1) and (H2) are inherited by $(\mathcal{P}_0, G_0, H_0)$; (H3) is inherited because subgroups of cyclic p -groups are cyclic p -groups. Replacing $(\mathcal{P}, G, H)$ by $(\mathcal{P}_0, G_0, H_0)$ we may assume $\mathcal{P}$ is connected and ρ is surjective.
- (R2) $H \neq 0$. If $H = 0$ then $\mathcal{P}_{G/H} = \mathcal{P}$ and $(\mathcal{P}_{G/H})^{[d]} = \mathcal{P}^{[d]G}$, so (H1) is literally the conclusion.

We can read a decomposition of fields directly from the canonical Cartesian square of [Proposition 23](#). After the reduction (R1) all four corners are integral: the three corners $\mathcal{P}^{[d]G}$, $(\mathcal{P}_{G/H})^{[d]}$ and $(\mathcal{P}_{G/H})^{(d)}$ are integral, and the fourth, $\mathcal{P}^{[d]H} = (K_H \rtimes S_d) \backslash \mathcal{P}^{\times S^d}$, is integral as a quotient

of the integral scheme $\mathcal{P}^{\times sd}$ (Theorem 32). So we may take function fields; using the open immersions ι, ι' of the master diagram together with $K_C = \text{Frac } V_C = K(C^{(d)}) = K(U^{(d)})$ and $K_{C'} = \text{Frac } V_{C'} = K(C'^{(d)}) = K((\mathcal{P}_{G/H})^{(d)})$, we identify

$$K(U^{(d)}) = K_C, \quad K((\mathcal{P}_{G/H})^{[d]}) = F, \quad K((\mathcal{P}_{G/H})^{(d)}) = K_{C'}, \quad K(\mathcal{P}^{[d]G}) = E,$$


with E/F Galois of group H , F/K_C Galois of group G/H , and the composite E/K_C Galois of group G . The fourth corner $\mathcal{P}^{[d]H}$ is the fibre product $\mathcal{P}^{[d]G} \times_{(\mathcal{P}_{G/H})^{[d]}} (\mathcal{P}_{G/H})^{(d)}$; being integral, its single function field is the compositum $E \cdot K_{C'}$. In particular its generic fibre $E \otimes_F K_{C'}$ over $(\mathcal{P}_{G/H})^{(d)}$ is a field, so by Lemma 37 the extensions E and $K_{C'}$ are linearly disjoint over F ; that is,

$$E \cap K_{C'} = F,$$

and $E \cdot K_{C'}/K_{C'}$ is Galois with group H . We denote:

1. $K_C := \text{Frac } V_C$, $K_{C'} := \text{Frac } V_{C'}$;
2. $F := K((\mathcal{P}_{G/H})^{[d]})$, $E := K(\mathcal{P}^{[d]G})$;
3. $V_F := V_{C'}|_F$;
4. a place v_E of E above V_F , and a place $v_\star$ of $E \cdot K_{C'}$ above both v_E and $V_{C'}$.

Thus we have:



(lines denote field inclusion / place lying-over, read upwards; in the left diagram a label is the Galois group of the extension where it is Galois). In addition E/K_C is Galois with group G .

Read through this dictionary, the hypotheses of the theorem become statements about the ramification of the places above:

- (H1) $(\mathcal{P}_{G/H})^{[d]} \rightarrow U^{(d)}$ is unramified at η_C means that the G/H -extension F/K_C is unramified at V_C . In particular V_F is unramified over V_C , with $e(V_F | V_C) = 1$.
- (H2) $\mathcal{P}^{[d]H} \rightarrow (\mathcal{P}_{G/H})^{(d)}$ is unramified at $\eta_{C'}$ means that the H -extension $E \cdot K_{C'}/K_{C'}$ is unramified at $V_{C'}$; equivalently the inertia group $I(v_\star | V_{C'}) \leq H$ is trivial.
- (H3) $G = \mathbb{Z}/p^r\mathbb{Z}$ makes G , hence H and every inertia group, a cyclic p -group; as the residue characteristic is p , the ramification of E/F is potentially *wild*.

The conclusion we are after — that $\mathcal{P}^{[d]G} \rightarrow U^{(d)}$ is unramified at η_C — is the statement that the G -extension E/K_C is unramified at V_C , i.e. that the inertia $I(v_E | V_C) \leq G$ is trivial. By (H1) the

lower layer F/K_C is already unramified at V_C , so this inertia lies in $H = \text{Gal}(E/F)$ and coincides with the inertia of the top layer,

$$I(v_E | V_C) = I(v_E | V_F) \leq H.$$

The theorem therefore reduces to the single assertion

$$E/F \text{ is unramified at } V_F, \quad \text{i.e.} \quad I(v_E | V_F) = 1.$$

Now $E \cdot K_{C'}/K_{C'}$ is the base change of E/F along $F \hookrightarrow K_{C'}$, and under the restriction isomorphism $\text{Gal}(E \cdot K_{C'}/K_{C'}) \cong \text{Gal}(E/F) = H$ the inertia injects

$$I(v_\star | V_{C'}) \hookrightarrow I(v_E | V_F).$$

Hypothesis (H2) says the source is trivial; the goal is that the target is trivial. These need *not* coincide: the two differ precisely by the ramification of $K_{C'}/F$ at $V_{C'} | V_F$ — the ramification that the symmetric-power base change q introduces at the point at infinity — which can be wild and can absorb the ramification of E/F . It is exactly this discrepancy that obstructs deducing the goal from (H1)–(H3).

Put in other words: The base change $K_{C'}/F$ is wildly ramified, so the inertia comparison is not onto .

The inertia injection, and why (H2) is not enough

The injection used above is an instance of the following general fact.

Lemma 69. *Let E/F be a finite Galois extension, linearly disjoint over F from a field extension K/F , with compositum $E \cdot K$; thus restriction is an isomorphism $r: \text{Gal}(E \cdot K/K) \xrightarrow{\sim} \text{Gal}(E/F)$, $\sigma \mapsto \sigma|_E$. Let w be a place of $E \cdot K$, and write $w_K := w|_K$, $w_E := w|_E$, $w_F := w|_F$ for its restrictions. Then r carries the inertia group of w over w_K into that of w_E over w_F :*

$$r(I(w | w_K)) \subseteq I(w_E | w_F),$$

so that $I(w | w_K) \hookrightarrow I(w_E | w_F)$ is injective. Moreover, completing at w identifies $I(w | w_K)$ with the inertia group of the local compositum $\widehat{E} \cdot \widehat{K}$ over $\widehat{K}$; in particular, if $\widehat{E} \subseteq \widehat{K}$ then $I(w | w_K) = 1$, no matter how ramified E/F is at w_F .

Proof. That r is an isomorphism is the translation theorem for the Galois extension E/F and the linearly disjoint K/F (Lemma 37(1) with $M = F$). For the inclusion, let $\sigma \in I(w | w_K)$. Since E/F is Galois and σ fixes $F \subseteq K$ pointwise, $\sigma(E) = E$ and $\sigma|_E \in \text{Gal}(E/F)$ fixes w_F . As σ lies in the decomposition group of w we have $w \circ \sigma = w$; restricting to E gives $w_E \circ \sigma|_E = w_E$, so $\sigma|_E \in D(w_E | w_F)$. Finally the residue extension $\kappa(w_E) \hookrightarrow \kappa(w)$ is equivariant and σ acts trivially on $\kappa(w)$, hence $\sigma|_E$ acts trivially on $\kappa(w_E)$, i.e. $\sigma|_E \in I(w_E | w_F)$. Injectivity is inherited from that of r . For the last assertion, the completion of $E \cdot K$ at w is the local compositum $\widehat{E}\widehat{K}$, with $\text{Gal}(\widehat{E}\widehat{K}/\widehat{K}) = D(w | w_K)$ and inertia subgroup $I(w | w_K)$; if $\widehat{E} \subseteq \widehat{K}$ then $\widehat{E}\widehat{K} = \widehat{K}$ and this group is trivial. $\square$

Taking $w = v_\star$ (so $w_K = V_{C'}$, $w_E = v_E$, $w_F = V_F$) recovers the injection $I(v_\star | V_{C'}) \hookrightarrow I(v_E | V_F)$. The implication “(H2) $\Rightarrow$ goal” would follow if this were always onto; the next example shows it need not be, already for $G = \mathbb{Z}/p^2\mathbb{Z}$ over $\mathbb{F}_p$.

Example 70. Take $k = \mathbb{F}_p$, $G = \mathbb{Z}/p^2\mathbb{Z}$, and $H = p\mathbb{Z}/p^2\mathbb{Z} \cong \mathbb{Z}/p\mathbb{Z}$ the unique subgroup of order p , so $G/H \cong \mathbb{Z}/p\mathbb{Z}$. We describe the local fields at the place $V_{C'} \mid V_F$ — the only place where the obstruction lives — writing $\widehat{(\cdot)}$ for completions.

The base and the G -tower. Let $\widehat{K_C} = \mathbb{F}_p((t))$, and present the local G -extension $\widehat{E}/\widehat{K_C}$ (an Artin–Schreier–Witt extension) by its two layers:

- lower layer $\widehat{F}/\widehat{K_C}$ (group G/H): *split*, so $\widehat{F} = \mathbb{F}_p((t))$. Then F/K_C is unramified at V_C — this is (H1) — and $\widehat{V_F} = \widehat{V_C}$;
- upper layer $\widehat{E}/\widehat{F}$ (group H): the Artin–Schreier extension

$$\widehat{E} = \widehat{F}(y), \quad y^p - y = t^{-1}.$$

By [Theorem 30\(3\)](#) (with $m = 1$, prime to p) this is totally, wildly ramified of degree p , with $I(v_E \mid V_F) = H \cong \mathbb{Z}/p\mathbb{Z} \neq 1$.

Hence $I(v_E \mid V_C) = I(v_E \mid V_F) = H \neq 1$: the extension E/K_C is ramified at V_C , so *the conclusion of the theorem fails*.

The base change. The places of $K_{C'}$ over V_F are governed by the symmetric-power cover q . Suppose at $V_{C'}$ this cover reproduces the *same* Artin–Schreier extension,

$$\widehat{K_{C'}} = \widehat{F}(y), \quad y^p - y = t^{-1}, \quad \text{i.e.} \quad \widehat{K_{C'}} = \widehat{E}.$$

This is compatible with the global linear disjointness $E \cap K_{C'} = F$: two *distinct* degree- p extensions of the global field F may have isomorphic completions at a place. Now the local compositum is $\widehat{E} \cdot \widehat{K_{C'}} = \widehat{E} \cdot \widehat{K_{C'}} = \widehat{E} = \widehat{K_{C'}}$, so $E \cdot K_{C'}/K_{C'}$ is split, hence unramified, at $V_{C'}$:

$$I(v_\star \mid V_{C'}) = \text{Gal}(\widehat{E}/\widehat{K_{C'}}) = 1.$$

Thus (H2) holds, and (H3) holds by construction.

Outcome. All of (H1)–(H3) hold, yet E/K_C is wildly ramified at V_C with inertia $H \cong \mathbb{Z}/p\mathbb{Z}$. The inertia injection of [Lemma 69](#) is here

$$\underbrace{I(v_\star \mid V_{C'})}_{=1} \hookrightarrow \underbrace{I(v_E \mid V_F)}_{=\mathbb{Z}/p\mathbb{Z}},$$

the inclusion of the trivial group: (H2) carries no information about the ramification of E/F , which has been entirely *absorbed* by that of $K_{C'}/F$ — the wild ramification introduced by q at the point at infinity.

Remark. This is globally consistent with everything above. Since $\widehat{E} \otimes_{\widehat{F}} \widehat{K_{C'}} = \widehat{E} \otimes_{\widehat{F}} \widehat{E}$ is not a field, $\mathcal{P}^{[d]_H}$ (function field $E \cdot K_{C'}$) *splits* at $V_{C'}$ into several places and is therefore unramified at $\eta_{C'}$, as (H2) demands; meanwhile $\mathcal{P}^{[d]_G}$ stays ramified at η_C . The hypotheses simply do not see the local coincidence $\widehat{K_{C'}} = \widehat{E}$, and ruling out such coincidences for a given $(\mathcal{P}, d)$ is exactly the global input the inductive argument lacks — which is why this is a wrong path.

Absorption made explicit: two local fields

Example 70 posited the local coincidence $\widehat{K_{C'}} = \widehat{E}$ abstractly. For completeness we exhibit it in the smallest possible setting — two honest local fields and a one-line Artin–Schreier reduction — to make the absorption phenomenon fully visible.

Example 71 (absorption of a wild break, explicit fields). Let $k = \bar{\mathbb{F}}_p$ and let $F = k((t))$. Define a second local field $K = k((\pi))$, and embed $F \hookrightarrow K$ by

$$t = \frac{\pi^p}{1 - \pi^{p-1}} = \pi^p(1 + \pi^{p-1} + \pi^{2(p-1)} + \cdots), \quad \text{so} \quad v_\pi(t) = p.$$

Since $t = \pi^p \cdot (\text{unit})$ and $dt/d\pi = -\pi^{2p-2}(1 - \pi^{p-1})^{-2} \neq 0$, the extension K/F is separable, *totally (wildly) ramified of degree p* ; equivalently it is the Artin–Schreier extension $\sigma^p - \sigma = t^{-1}$ with $\sigma = \pi^{-1}$, of break 1 ([Theorem 30\(3\)](#)).

Now let E/F be the Artin–Schreier extension

$$E = F(y), \quad y^p - y = t^{-1}.$$

Over F this is ramified: $v_t(t^{-1}) = -1$ is prime to p , so by [Theorem 30\(3\)](#) it is totally wildly ramified of degree p , break 1, with $I(v_E | v_F) = \mathbb{Z}/p\mathbb{Z} \neq 1$.

Yet E is absorbed by K . Computing t^{-1} inside K ,

$$t^{-1} = \pi^{-p}(1 - \pi^{p-1}) = \pi^{-p} - \pi^{-1} = \wp(\pi^{-1}), \quad \wp(x) := x^p - x,$$

so $y = \pi^{-1} \in K$ already solves $y^p - y = t^{-1}$. Hence $E = F(\pi^{-1}) \subseteq K$ (indeed $E = K$, both being degree p over F), the compositum is $E \cdot K = K$, and

$$E \cdot K/K \text{ is trivial, hence unramified: } I(v_\star | v_K) = 1,$$

while E/F is ramified. The ramification of E/F has been *entirely absorbed* by the wild base change to K — exactly the failure of surjectivity in the inertia injection $I(v_\star | v_K) \hookrightarrow I(v_E | v_F)$, here $1 \hookrightarrow \mathbb{Z}/p\mathbb{Z}$.

Remark. The mechanism, and why F cannot do it. Read over K , the datum t^{-1} has pole order $v_\pi(t^{-1}) = p$, divisible by p ; this is what licenses the Artin–Schreier reduction $y \mapsto y - \pi^{-1}$, which removes the pole completely because $\wp(\pi^{-1})$ matches t^{-1} to all orders. The substitution is available over K but *not* over F : the reducing element π^{-1} has valuation -1 , whereas $v_\pi(F^\times) = p\mathbb{Z}$ contains no element of valuation -1 at all. Thus the wild base change does not “fight” the ramification of E/F ; it *supplies the one element of intermediate valuation* that the Artin–Schreier reduction needs. The ramification index $e(K/F) = p$ is the gate that turns the prime-to- p break 1 into the divisible-by- p pole order p over K .

General r . For $G = \mathbb{Z}/p^r$ the faithful local model has $\widehat{K_{C'}}/\widehat{F}$ totally ramified of degree p^{r-1} (the residue field k being algebraically closed, there is no residual part), and absorption is the proper inclusion $\widehat{E} \subsetneq \widehat{K_{C'}}$ of the top-layer $\mathbb{Z}/p$ -extension. The explicit fields are obtained by iterating the recipe above: with $E = k((\pi))$, $\pi = y^{-1}$ as here, set $\pi = \rho^p(1 - \rho^{p-1})^{-1}$ to get a degree- p overfield $k((\rho)) \supsetneq E$, and so on $r - 1$ times. The pole order of the top datum is multiplied to $p^{r-1} \cdot 1 \equiv 0 \pmod{p}$ and reduced away by the same one-line substitution. Over an algebraically closed residue field, “ $E \cdot K/K$ unramified with E/F ramified” is *equivalent* to $\widehat{E} \subseteq \widehat{K_{C'}}$ — there are no nontrivial unramified extensions to hide in — so absorption is precisely a containment of local fields, never a softer phenomenon.

References

- [SGA1] Alexander Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*. Vol. 224. Lecture notes in mathematics. Springer-Verlag, 1971.
- [Ser79] Jean-Pierre Serre. *Local Fields*. Springer, 1979.
- [Koc97] Helmut Koch. *Algebraic Number Theory*. Vol. 62. Encyclopaedia of Mathematical Sciences. Berlin, Heidelberg: Springer-Verlag, 1997. ISBN: 978-3-540-63003-6.
- [Tho05] Lara Thomas. “Ramification groups in Artin-Schreier-Witt extensions”. In: *Journal de théorie des nombres de Bordeaux* 17.2 (2005), pp. 689–720. DOI: [10.5802/jtnb.514](https://doi.org/10.5802/jtnb.514). URL: <https://jtnb.centre-mersenne.org/articles/10.5802/jtnb.514/>.
- [Mil08] James S. Milne. *Abelian Varieties (v2.00)*. Available at www.jmilne.org/math/. 2008.
- [Tót11] Péter Tóth. “Geometric Abelian Class Field Theory”. In: (May 2011). Master of Science thesis. URL: <https://math.bu.edu/people/rmagner/Seminar/GCFTthesis.pdf>.
- [Rab14] Joseph Rabinoff. *The Theory of Witt Vectors*. 2014. arXiv: [1409.7445](https://arxiv.org/abs/1409.7445) [math.RA]. URL: <https://arxiv.org/abs/1409.7445>.
- [Ten15] Avichai Tendler. *Geometric Class Field Theory*. 2015. arXiv: [1507.00104](https://arxiv.org/abs/1507.00104) [math.AG]. URL: <https://arxiv.org/abs/1507.00104>.
- [Stacks] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>. 2018.
- [Gui19] Quentin Guignard. “On the ramified class field theory of relative curves”. In: *Algebra & Number Theory* 13 (May 2019). Revised version submitted on 29 May 2019, pp. 1299–1326. DOI: [10.2140/ant.2019.13.1299](https://doi.org/10.2140/ant.2019.13.1299). arXiv: [1804.02243](https://arxiv.org/abs/1804.02243) [math.AG].
- [Tak19] Daichi Takeuchi. “Blow-ups and the class field theory for curves”. In: *Algebraic Number Theory* 13.6 (2019), pp. 1327–1351. DOI: [10.2140/ant.2019.13.1327](https://doi.org/10.2140/ant.2019.13.1327). URL: <https://doi.org/10.2140/ant.2019.13.1327>.
- [You20] Alex Youcis. “Notes on Torsors”. Available at <https://alex-youcis.github.io/torsors.pdf>. 2020. URL: <https://alex-youcis.github.io/torsors.pdf>.