

# Clinical Archetypes Overview — NM\_false\_alarm

Top 3 highest-significance patient subgroups

## #1 COAGULATION DERANGEMENT

### Clinical Conditions

- Coagulation derangement present
- Synthetic liver dysfunction

#### Key Statistics

|             |              |
|-------------|--------------|
| Patients:   | 837          |
| Coverage:   | 17.2%        |
| Outcome:    | 8.6% vs 3.4% |
| Enrichment: | 2.51×        |
| p-value:    | 8.07e-14     |

## #2 ACUTE KIDNEY INJURY

### Clinical Conditions

- Acute kidney injury present

#### Key Statistics

|             |              |
|-------------|--------------|
| Patients:   | 1,132        |
| Coverage:   | 22.8%        |
| Outcome:    | 6.4% vs 2.8% |
| Enrichment: | 2.25×        |
| p-value:    | 2.17e-13     |

## #3 COAGULATION DERANGEMENT

### Clinical Conditions

- Coagulation derangement present
- Acute kidney injury present

#### Key Statistics

|             |              |
|-------------|--------------|
| Patients:   | 729          |
| Coverage:   | 15.0%        |
| Outcome:    | 8.4% vs 3.4% |
| Enrichment: | 2.44×        |
| p-value:    | 5.38e-11     |