

# Clinical Archetypes Overview — SM\_false\_alarm

Top 3 highest-significance patient subgroups

#1

## ACUTE KIDNEY INJURY

### Clinical Conditions

- Coagulation derangement present
- Acute kidney injury present

### Key Statistics

|             |              |
|-------------|--------------|
| Patients:   | 741          |
| Coverage:   | 15.3%        |
| Outcome:    | 9.9% vs 2.9% |
| Enrichment: | 3.45×        |
| p-value:    | 4.59e-24     |

#2

## ACUTE KIDNEY INJURY

### Clinical Conditions

- Acute kidney injury present

### Key Statistics

|             |              |
|-------------|--------------|
| Patients:   | 1,074        |
| Coverage:   | 22.2%        |
| Outcome:    | 7.4% vs 2.9% |
| Enrichment: | 2.61×        |
| p-value:    | 1.25e-19     |

#3

## NEUTROPHILIA

### Clinical Conditions

- Ionized hypocalcemia present
- Neutrophilia present

### Key Statistics

|             |              |
|-------------|--------------|
| Patients:   | 795          |
| Coverage:   | 16.4%        |
| Outcome:    | 7.8% vs 2.9% |
| Enrichment: | 2.73×        |
| p-value:    | 2.60e-15     |