

Clinical Archetypes Overview — SM_false_alarm

Top 8 highest-significance patient subgroups

#1 COAGULATION DERANGEMENT

Clinical Conditions

- Prolonged coagulation present
- Synthetic hepatic dysfunction

Key Statistics

Patients:	846
Coverage:	17.5%
Outcome:	9.6% vs 2.9%
Enrichment:	3.36×
p-value:	4.68e-26

#2 COAGULATION DERANGEMENT

Clinical Conditions

- Low albumin present
- Prolonged coagulation present

Key Statistics

Patients:	716
Coverage:	14.8%
Outcome:	10.3% vs 2.9%
Enrichment:	3.62×
p-value:	2.21e-25

#3 COAGULATION DERANGEMENT

Clinical Conditions

- Creatinine elevation present
- Prolonged coagulation present

Key Statistics

Patients:	741
Coverage:	15.3%
Outcome:	9.9% vs 2.9%
Enrichment:	3.45×
p-value:	7.54e-24

#4 NEUTROPHILIA

Clinical Conditions

- High neutrophil count present
- Synthetic hepatic dysfunction

Key Statistics

Patients:	544
Coverage:	11.2%
Outcome:	11.6% vs 2.9%
Enrichment:	4.06×
p-value:	1.14e-23

#5 NEUTROPHILIA

Clinical Conditions

- High neutrophil count present
- Prolonged coagulation present

Key Statistics

Patients:	1,432
Coverage:	29.6%
Outcome:	6.6% vs 2.9%
Enrichment:	2.33×
p-value:	1.40e-20

#6 LOW ALBUMIN

Clinical Conditions

- Low albumin present

Key Statistics

Patients:	956
Coverage:	19.8%
Outcome:	8.1% vs 2.9%
Enrichment:	2.82×
p-value:	2.56e-20

#7 COAGULATION DERANGEMENT

Clinical Conditions

- Mechanical ventilation data present
- Prolonged coagulation present

Key Statistics

Patients:	1,011
Coverage:	20.9%
Outcome:	7.7% vs 2.9%
Enrichment:	2.70×
p-value:	1.45e-19

#8 CREATININE ELEVATION

Clinical Conditions

- Creatinine elevation present

Key Statistics

Patients:	1,074
Coverage:	22.2%
Outcome:	7.4% vs 2.9%
Enrichment:	2.61×
p-value:	2.75e-19