

Clinical Archetypes Overview — SM_false_alarm

Top 8 highest-significance patient subgroups

#1 LOW ALBUMIN

Clinical Conditions

- Low albumin present

Key Statistics

Patients:	753
Coverage:	18.5%
Outcome:	19.4% vs 11.9%
Enrichment:	1.63×
p-value:	3.20e-09

#2 LOW ALBUMIN

Clinical Conditions

- Ionized hypocalcemia present
- Low albumin present

Key Statistics

Patients:	412
Coverage:	10.1%
Outcome:	22.6% vs 11.9%
Enrichment:	1.90×
p-value:	4.56e-09

#3 NEUTROPHILIA

Clinical Conditions

- High neutrophil count present
- Low albumin present
- Synthetic hepatic dysfunction

Key Statistics

Patients:	301
Coverage:	7.4%
Outcome:	24.6% vs 11.9%
Enrichment:	2.06×
p-value:	7.21e-09

#4 NEUTROPHILIA

Clinical Conditions

- High neutrophil count present
- Synthetic hepatic dysfunction

Key Statistics

Patients:	439
Coverage:	10.8%
Outcome:	21.9% vs 11.9%
Enrichment:	1.84×
p-value:	9.98e-09

#5 NEUTROPHILIA

Clinical Conditions

- High neutrophil count present
- Low albumin present

Key Statistics

Patients:	494
Coverage:	12.1%
Outcome:	21.1% vs 11.9%
Enrichment:	1.77×
p-value:	9.98e-09

#6 COAGULATION DERANGEMENT

Clinical Conditions

- Low albumin present
- Prolonged coagulation present

Key Statistics

Patients:	557
Coverage:	13.7%
Outcome:	20.1% vs 11.9%
Enrichment:	1.69×
p-value:	2.61e-08

#7 LIVER DYSFUNCTION

Clinical Conditions

- Low albumin present
- Synthetic hepatic dysfunction

Key Statistics

Patients:	456
Coverage:	11.2%
Outcome:	21.1% vs 11.9%
Enrichment:	1.77×
p-value:	4.79e-08

#8 LOW ALBUMIN

Clinical Conditions

- Low albumin present
- SIRS heart-rate criterion present

Key Statistics

Patients:	320
Coverage:	7.9%
Outcome:	23.1% vs 11.9%
Enrichment:	1.94×
p-value:	5.30e-08