

Conjugate Gradient Method

Introduction

Gram-Schmidt

$$R = r_i \otimes e_i$$

$$R^T \cdot R = I$$

Foundation of Conjugate Gradient Method

(P, P^{-T}) are a pair of covariant & contravariant transformations, and (p_i, p^i) are the vectors that represent:

$$\begin{cases} P = p_i \otimes e_i \\ P^{-T} = p^i \otimes e_i \\ p^i \cdot p_j = \delta_{ij} \end{cases}$$

The transformation that transforms covariant space to contravariant is:

$$\Phi = p^i \otimes p_i = P^{-T} \cdot P^{-1}$$

This transformation Φ is positive semi-definite.

Lemma 1: for any covariant transformation P , the covariant-to-contravariant transformation $\Phi = P^{-T} \cdot P^{-1}$ is also the covariant-to-contravariant transformation for $Q = P \cdot R$.

Proof 1: If the covariant transformation happens after a unitary transformation, then the covariant-to-contravariant transformation is independent of the unitary transformation:

$$Q = P \cdot R$$

$$\Psi = Q^{-T} \cdot Q^{-1} = P^{-T} \cdot R^{-T} \cdot R^{-1} \cdot P^{-1} = \Phi = P^{-T} \cdot P^{-1}$$

Lemma 2: any positive semi-definite tensor can be represented as a tensor multiply (dot product) by its transpose.

Proof 2: A is positive semi-definite tensor, by spectrum decomposition:

$$A = A_{ij} e_i \otimes e_j = \sum_i \lambda_i n_i \otimes n_i \quad \text{where } A_{ij} = A_{ji}, \lambda_i \geq 0$$

$$\mathbf{A} = \mathbf{B}^T \cdot \mathbf{B} \quad \text{where } \mathbf{B} = \mathbf{A}^{\frac{1}{2}} = \sum_i \lambda_i^{\frac{1}{2}} \mathbf{n}_i \otimes \mathbf{n}_i$$

Now if we claim $\mathbf{A} = \mathbf{P}^{-T} \cdot \mathbf{P}^{-1}$, by the same procedure we find:

$$\mathbf{P} = \mathbf{A}^{-\frac{1}{2}} = \lambda_i^{-\frac{1}{2}} \mathbf{n}_i \otimes \mathbf{n}_i$$

Then with Lemma 1:

$$\mathbf{A} = \mathbf{Q}^{-T} \cdot \mathbf{Q}^{-1} \quad \text{where } \mathbf{Q} = \mathbf{P} \cdot \mathbf{R}$$

$$\det(\mathbf{A}) = \det(\mathbf{Q})$$

This has guarantee that, by selecting any initial directions within \mathbf{A} space, we can always generate a series of conjugate vectors with respect to \mathbf{A} (\mathbf{A} – **orthonormal** basis) that span the entire \mathbf{A} space through the Generalized Gram-Schmidt Process.

Implementation of Conjugate Gradient Method

Relationship with Krylov Subspace