

CLINICAL MULTIPLE CHOICE EVALUATION

LECTURE: ROLES & RESPONSIBILITIES OF NURSES DURING DRUG ADMINISTRATION



Candidate Instructions

- This formal testing paper contains **25 strict, text-grounded Multiple Choice Questions**.
- All items are directly derived from the core syllabus, frameworks, and verified statistical metrics.
- Select the single best answer choice (**A, B, C, or D**) for each item.
- The full verification matrix key is appended to the absolute end of this paper booklet.

QUESTION 1

According to the lecture text, which of the following perfectly defines a "Medication Error"?

- A) Any unpreventable physiological complication directly triggered by a correctly processed medication
- B) Any Patient Safety Incident (PSI) where there has been an error in prescribing, preparing, dispensing, administering, monitoring, or providing advice on medicines
- C) A purely administrative discrepancy that does not involve clinical delivery or monitoring steps
- D) An unavoidable chemical interaction occurring exclusively during multi-drug clinical trials

QUESTION 2

The lecture explicitly highlights that medication errors are structurally distinct from which of the following?

- A) Patient Safety Incidents (PSI)
- B) Adverse drug reactions
- C) Human factor variations
- D) Weak organizational medication systems

QUESTION 3

Which list comprises the three specific "Human Factors" explicitly listed in the curriculum slides to consider when processes are affected?

- A) Inexperience, poor motivation, lack of attention
- B) Fatigue, environmental conditions, staffing levels
- C) Age, cognitive background, language barriers
- D) Technical baseline, communication layout, ward history

QUESTION 4

How many medication errors are estimated to be made every single year in England?

- A) Over 50 million
- B) Over 110 million
- C) Over 180 million
- D) Over 237 million

QUESTION 5

The avoidable consequences of medication errors cost the NHS upwards of how much money every year?

- A) £12 million
- B) £45 million
- C) £71 million
- D) £98 million

QUESTION 6

The avoidable consequences of medication errors are estimated to cost how many lives every year in England?

- A) 500+ lives
- B) 1,000+ lives
- C) 1,700+ lives
- D) 2,500+ lives

QUESTION 7

According to research statistics presented in the slides, over half (54%) of all medication errors occur at which stage of the process?

- A) The point of prescribing
- B) The point of dispensing
- C) The point of administration
- D) The point of preparing

QUESTION 8

What percentage of total medication errors are made during the prescribing stage?

- A) 16%
- B) 21%
- C) 38%
- D) 42%

QUESTION 9

Despite covering fewer patients than other sectors, which sector exhibits the highest error rates at 42%?

- A) Primary care
- B) Hospitals
- C) Care homes
- D) Outpatient clinics

QUESTION 10

Researchers estimate that nearly 3 out of 4 medication errors (72%) fall under which severity classification?

- A) Minor
- B) Moderate harm potential
- C) Serious harm potential
- D) Fatal potential

QUESTION 11

What percentage of medication errors are estimated to have the explicit potential to cause serious harm?

- A) 2%
- B) 16%
- C) 21%
- D) 26%

QUESTION 12

Which of the following medications is listed first as most often implicated in hospital admissions attributable to medication errors?

- A) Inhaled corticosteroids
- B) Non-steroidal anti-inflammatories (NSAIDs)
- C) Water tablets (diuretics)
- D) Cardiac glycosides

QUESTION 13

According to NHS Resolution initial data for litigation claims, which specific medication class was implicated in the highest number of claims (104)?

- A) Antimicrobials
- B) Opiates
- C) Anticoagulants
- D) Insulin

QUESTION 14

The Nursing and Midwifery Council's (NMC) (2018) Code explicitly mandates that nurses must execute which primary action?

- A) Prioritize the completion of unit audit records over external patient interactions
- B) Put the interests of people using or needing nursing services first
- C) Delegate all medication preparation steps to student tiers under all circumstances
- D) Decline medication administration if any human factor variations are present on the ward

QUESTION 15

Based on the framework provided by Ofosu and Jarrett (2015), which three elements are grouped together as core causes behind medication errors?

- A) Weak infrastructure, poor illumination, system failures
- B) Interruptions and distractions, poor drug calculations, inadequate education and compliance
- C) Incompetence, callousness, system negligence
- D) Unclear prescriptions, wrong frequency, omitted drugs

QUESTION 16

The lecture presents a tragic case example where a patient died following an incorrect route error. What specifically occurred?

- A) An intravenous solution was connected to an enteral feeding tube line
- B) A patient received an oral antibiotic tablet crushed in water via an intravenous infusion
- C) An intramuscular injection was given sub-dermally into the patient's heel zone
- D) A nebulizer drug was completely replaced with an oral syrup mix

QUESTION 17

What are the strict supervision parameters defined by the NMC for student nurses regarding medication administration or supply?

- A) Independent execution once the second year of study is fully completed
- B) Indirect supervision via telephone check-ins by a qualified staff tier
- C) Student nurses must NEVER administer or supply medicinal products without DIRECT supervision
- D) Peer-to-peer supervision alongside another student nurse from the same cohort block

QUESTION 18

What is the immediate initial step that must be taken the moment a medication error is identified?

- A) Complete a full local Datix event log before approaching the patient bed
- B) Stop the medication and revert the effect (if possible!) and monitor the patient's status
- C) Contact the family immediately to gain explicit informed consent to proceed
- D) Review the Yellow Card scheme logs to determine if it requires national alert escalation

QUESTION 19

Under the "10 Rs of Drug Administration" framework, which of the following is highlighted as a critical action under "1. Right Patient"?

- A) Check the name on the prescription and wristband, using 2 or more identifiers and asking the patient to identify themselves
- B) Verify the room and bay configuration number exclusively
- C) Rely completely on verbal confirmation from a peer staff tier during handover
- D) Cross-reference the patient chart against the shift entry index log numbers only

QUESTION 20

Under "2. Right Drug", what explicit instruction is given regarding medication naming conventions?

- A) Brand names should be preferred because they prevent formatting errors
- B) Brand names should be avoided, and the generic name of the medication must be verified
- C) Only the localized abbreviated slang title should be verified across drug cups
- D) The chemical formula string must be manually transcribed into the chart

QUESTION 21

What is the study of how the body interacts with administered substances for the entire duration of exposure?

- A) Pharmacodynamics
- B) Pharmacokinetics
- C) Biotransformation
- D) Capillary permeability

QUESTION 22

According to the pharmacokinetics layout slide, which organ serves as the "Main" excretory organ?

- A) Lung
- B) Skin
- C) Liver
- D) Kidneys

QUESTION 23

What is the mandatory timeline rule regarding chart verification under "10. Right Response or Outcome (Documentation)"?

- A) Ensure you have signed for the medication immediately BEFORE approaching the bay zone
- B) Ensure you have signed for the medication AFTER it has been administered
- C) Sign for all scheduled medications in a single block task at the end of the full shift window
- D) Sign the chart only after the next shift nurse co-signs the verification log

QUESTION 24

A study carried out in 2010 by McMullan et al. demonstrated that what percentage of nursing students failed the initial drug calculation test?

- A) 45%
- B) 55%
- C) 89%
- D) 92%

QUESTION 25

The 2010 calculation error study noted that independent of status or experience, which demographic group was statistically significantly more able to perform numerical calculations?

- A) Older participants (≥ 35 years)
- B) Younger participants (≤ 21 years)
- C) Participants with less than one year of practice experience
- D) Student nurses who prepared primarily via online formula modules

OFFICIAL EXAM ANSWER KEY MATRIX

Verify all candidate choice selections against this matrix block.

Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10
B	B	B	D	D	C	C	B	C	A

Q11	Q12	Q13	Q14	Q15	Q16	Q17	Q18	Q19	Q20
A	B	C	B	B	B	C	B	A	B

Q21	Q22	Q23	Q24	Q25
B	D	B	D	A