

Clinical Multiple Choice

Evaluation: Risk Assessment in Adult Nursing

1. Which of the following is NOT explicitly identified as a purpose of conducting risk assessments in the clinical environment?

- A) Ensuring compliance with clinical guidelines and legislation
- B) Protects healthcare staff from liability
- C) Eliminating all clinical hazards entirely with zero margin for error
- D) Preventing adverse events

2. Which of the following describes the fundamental 'Key Principles' of risk assessment?

- A) Ad-hoc, isolated process, non-participatory, and task-oriented
- B) Systematic approach, continuous process, involves all healthcare team members, patient-centered
- C) Manager-led, fixed-point annual review, senior-tier only, audit-driven
- D) Reactive-based, budget-restricted, equipment-focused, and time-saving

3. What are the three categories of risks in adult nursing explicitly outlined on Slide 11?

- A) Clinical Risks, Environmental Risks, and Psychosocial Risks
- B) Legal Risks, Financial Risks, and Staffing Risks
- C) Surgical Risks, Pharmacological Risks, and Hygiene Risks
- D) Equipment Risks, Documentation Risks, and Management Risks

4. In the context of 'Clinical Risks' for patient deterioration, which specific tools are cited as primary screening methods?

- A) Waterlow index and Morse scale
- B) ABCDE Framework and Sepsis screen tool
- C) Braden Scale and NEWS2 Threshold
- D) Bedrail assessment and physical space mapping

5. When assessing 'Falls risk', what are the two distinct types of factors that must be evaluated?

- A) Inpatient and outpatient factors
- B) Intrinsic factors and extrinsic factors
- C) Primary factors and secondary factors
- D) Cognitive factors and physical factors

Official Answer Key

1: C | 2: B | 3: A | 4: B | 5: B