

A HIGH-ORDER DISCONTINUOUS GALERKIN METHOD FOR THE BIDOMAIN PROBLEM OF CARDIAC ELECTROPHYSIOLOGY

Project N° 2

Supervisors: Christian Vergara, Paola Antonietti

Federica Botta, Matteo Calafà

Course of Numerical Analysis for Partial Differential Equations

A. Y. 2020/2021

Table of Contents

- 1 Introduction
- 2 Dubiner basis implementation

The physical problem


Mechanical contraction of the human heart



Electrical activation of the cardiac cells



Continuous electrical diffusion over the entire cardiac surface.



The mathematical model

Bidomain model + FitzHugh-Nagumo with Neumann B.C.

$$\left\{ \begin{array}{ll} \chi_m C_m \frac{\partial V_m}{\partial t} - \nabla \cdot (\Sigma_i \nabla \phi_i) + \chi_m I_{ion}(V_m, w) = I_i^{ext}, & \text{in } \Omega_{mus} \times (0, T], \\ -\chi_m C_m \frac{\partial V_m}{\partial t} - \nabla \cdot (\Sigma_e \nabla \phi_e) - \chi_m I_{ion}(V_m, w) = -I_e^{ext}, & \text{in } \Omega_{mus} \times (0, T], \\ I_{ion}(V_m, w) = k V_m (V_m - a)(V_m - 1) + w, & \text{in } \Omega_{mus} \times (0, T], \\ \frac{\partial w}{\partial t} = \epsilon (V_m - \gamma w), & \text{in } \Omega_{mus} \times (0, T], \\ \Sigma_i \nabla \phi_i \cdot n = b_i, & \text{on } \partial \Omega_{mus} \times (0, T], \\ \Sigma_e \nabla \phi_e \cdot n = b_e, & \text{on } \partial \Omega_{mus} \times (0, T], \\ \text{Initial conditions for } \phi_i, \phi_e, w, & \text{in } \Omega_{mus} \times \{t = 0\}. \end{array} \right.$$

Unknowns: $\phi_i, \phi_e, V_m = \phi_i - \phi_e, w$

Our objectives

What had already been done:

- Implementation of a Discontinuous Galerkin with FEM basis for the Bidomain problem.
- Implementation of a Semi-Implicit temporal scheme.

What we did:

- Implementation of a Discontinuous Galerkin with **Dubiner** basis for the Bidomain problem.
- Implementation of further temporal schemes.
- Bugs corrections and optimizations.
- Pseudo-realistic simulations.

Analytical definition

Definition (Dubiner basis)

The Dubiner basis that generates the space $\mathbb{P}^p(\hat{K})$ of polynomials of degree p over the reference triangle is the set of functions:

$$\phi_{ij} : \hat{K} \rightarrow \mathbb{R},$$


$$\phi_{ij}(\xi, \eta) = c_{ij} 2^j (1 - \eta)^j J_i^{0,0} \left(\frac{2\xi}{1 - \eta} - 1 \right) J_j^{2i+1,0}(2\eta - 1),$$

for $i, j = 0, \dots, p$ and $i + j \leq p$, where $c_{ij} := \sqrt{\frac{2(2i+1)(i+j+1)}{4^i}}$
and $J_i^{\alpha,\beta}(\cdot)$ is the i -th Jacobi polynomial.

Properties

- They consist in a pseudo tensor-product of Jacobi polynomials if the following transformation is then applied:

$$\xi = \frac{(1+a)(1-b)}{4}, \eta = \frac{(1+b)}{2}.$$



- They are $L^2(\hat{K})$ orthonormal (\hat{K} is the reference triangle).

Main works

Remark

*Dubiner basis coefficients of a discretized function have **modal** meaning instead of a nodal meaning.*

Then, our main works regarded:

- Methods for the evaluation of the Dubiner functions and gradients in the reference points.
- Methods for the evaluation of the FEM coefficients of a discretized function starting from its Dubiner coefficients and viceversa.
 - 1 FEM \rightarrow Dubiner is needed when we use the initial solution data into the Dubiner system.
 - 2 FEM \leftarrow Dubiner is needed when we want to get the solution obtained from the Dubiner system in a comprehensible form.

FEM-Dubiner conversion strategies

Consider:

- An element $\mathcal{K} \in \tau_h$
- $\{\psi_i\}_{i=1}^p, \{\varphi_j\}_{j=1}^q$ as the FEM and Dubiner functions with support in \mathcal{K} .
- $\{\hat{u}_i\}_{i=1}^p, \{\tilde{u}_j\}_{j=1}^q$ as the FEM and Dubiner coefficients of a function u_h .

FEM \leftarrow Dubiner

Exploiting the nodal meaning of FEM, we compute its value in a point:

$$\hat{u}_i = \sum_{j=1}^q \tilde{u}_j \phi_j(x_i),$$

FEM \rightarrow Dubiner

Exploiting the L^2 -orthonormality of Dubiner, we compute its Fourier coeff.:

$$\tilde{u}_j = \int_{\mathcal{K}} u_h(x) \varphi_j(x) dx = \int_{\mathcal{K}} \sum_{i=1}^p \hat{u}_i \psi_i(x) \varphi_j(x) dx = \sum_{i=1}^p \left(\int_{\mathcal{K}} \psi_i(x) \varphi_j(x) dx \right) \hat{u}_i.$$