

A HIGH-ORDER DISCONTINUOUS GALERKIN METHOD FOR THE BIDOMAIN PROBLEM OF CARDIAC ELECTROPHYSIOLOGY

Project N° 2

Supervisors: Christian Vergara, Paola Antonietti

Federica Botta, Matteo Calafà

Course of Numerical Analysis for Partial Differential Equations

A.Y. 2020/2021

Table of Contents

- 1 Introduction
- 2 Dubiner basis implementation
- 3 Semi-discrete Discontinuous Galerkin
- 4 Temporal discretization
- 5 Uniqueness of the potentials

The physical problem


Mechanical contraction of the human heart



Electrical activation of the cardiac cells



Continuous electrical diffusion over the entire cardiac surface.



The mathematical model

Bidomain model + FitzHugh-Nagumo with Neumann B.C.

$$\left\{ \begin{array}{ll} \chi_m C_m \frac{\partial V_m}{\partial t} - \nabla \cdot (\Sigma_i \nabla \phi_i) + \chi_m l_{ion}(V_m, w) = I_i^{ext}, & \text{in } \Omega_{mus} \times (0, T], \\ -\chi_m C_m \frac{\partial V_m}{\partial t} - \nabla \cdot (\Sigma_e \nabla \phi_e) - \chi_m l_{ion}(V_m, w) = -I_e^{ext}, & \text{in } \Omega_{mus} \times (0, T], \\ l_{ion}(V_m, w) = k V_m (V_m - a)(V_m - 1) + w, & \text{in } \Omega_{mus} \times (0, T], \\ \frac{\partial w}{\partial t} = \epsilon (V_m - \gamma w), & \text{in } \Omega_{mus} \times (0, T], \\ \Sigma_i \nabla \phi_i \cdot n = b_i, & \text{on } \partial \Omega_{mus} \times (0, T], \\ \Sigma_e \nabla \phi_e \cdot n = b_e, & \text{on } \partial \Omega_{mus} \times (0, T], \\ \text{Initial conditions for } \phi_i, \phi_e, w, & \text{in } \Omega_{mus} \times \{t = 0\}. \end{array} \right.$$

Unknowns: $\phi_i, \phi_e, V_m = \phi_i - \phi_e, w$

Our objectives

What had already been done:

- Implementation of a Discontinuous Galerkin with FEM basis for the Bidomain problem.
- Implementation of a Semi-Implicit temporal scheme.

What we did:

- Implementation of a Discontinuous Galerkin with **Dubiner** basis for the Bidomain problem.
- Implementation of further temporal schemes.
- Bugs corrections and optimizations.
- Pseudo-realistic simulations.



Analytical definition

Definition (Dubiner basis)

The Dubiner basis that generates the space $\mathbb{P}^p(\hat{K})$ of polynomials of degree p over the reference triangle is the set of functions:

$$\phi_{ij} : \hat{K} \rightarrow \mathbb{R},$$


$$\phi_{ij}(\xi, \eta) = c_{ij} 2^j (1 - \eta)^j J_i^{0,0} \left(\frac{2\xi}{1 - \eta} - 1 \right) J_j^{2i+1,0}(2\eta - 1),$$

for $i, j = 0, \dots, p$ and $i + j \leq p$, where $c_{ij} := \sqrt{\frac{2(2i+1)(i+j+1)}{4^i}}$
 and $J_i^{\alpha,\beta}(\cdot)$ is the i -th Jacobi polynomial.

Properties

- They consist in a pseudo tensor-product of Jacobi polynomials if the following transformation is then applied:

$$\xi = \frac{(1+a)(1-b)}{4}, \eta = \frac{(1+b)}{2}.$$



- They are $L^2(\hat{K})$ orthonormal (\hat{K} is the reference triangle).

Main works

Remark

*Dubiner basis coefficients of a discretized function have **modal** meaning instead of a nodal meaning.*

Then, our main works regarded:

- Methods for the evaluation of the Dubiner functions and gradients in the reference points.
- Methods for the evaluation of the FEM coefficients of a discretized function starting from its Dubiner coefficients and viceversa.
 - 1 FEM \rightarrow Dubiner is needed when we use the initial solution data into the Dubiner system.
 - 2 FEM \leftarrow Dubiner is needed when we want to get the solution obtained from the Dubiner system in a comprehensible form.

FEM-Dubiner conversion strategies

Consider:

- An element $\mathcal{K} \in \tau_h$
- $\{\psi_i\}_{i=1}^p, \{\varphi_j\}_{j=1}^q$ as the FEM and Dubiner functions with support in \mathcal{K} .
- $\{\hat{u}_i\}_{i=1}^p, \{\tilde{u}_j\}_{j=1}^q$ as the FEM and Dubiner coefficients of a function u_h .

FEM \leftarrow Dubiner

Exploiting the nodal meaning of FEM, we compute its value in a point:

$$\hat{u}_i = \sum_{j=1}^q \tilde{u}_j \phi_j(x_i),$$

FEM \rightarrow Dubiner

Exploiting the L^2 -orthonormality of Dubiner, we compute its Fourier coeff.:

$$\tilde{u}_j = \int_{\mathcal{K}} u_h(x) \varphi_j(x) dx = \int_{\mathcal{K}} \sum_{i=1}^p \hat{u}_i \psi_i(x) \varphi_j(x) dx = \sum_{i=1}^p \left(\int_{\mathcal{K}} \psi_i(x) \varphi_j(x) dx \right) \hat{u}_i.$$

Discretization

space-dependent: Discontinuous Galerkin method

Bidomain problem

```
graph TD; A[Bidomain problem] --> B[space-dependent: Discontinuous Galerkin method]; A --> C[time-dependent: Semi-implicit, Godunov operator-splitting and quasi-implicit operator-splitting]
```

time-dependent: Semi-implicit, Godunov operator-splitting and quasi-implicit operator-splitting

Semi-discrete Discontinuous Galerkin formulation

For any $t \in [0, T]$ find $\Phi_h(t) = [\phi_i^h(t), \phi_e^h(t)]^T \in [V_h^p]^2$ and $w_h(t) \in V_h^p$ such that:

$$\left\{ \begin{array}{l} \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \chi_m c_m \frac{\partial V_m^h}{\partial t} v_h d\omega + a_i(\phi_i^h, v_h) + \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \chi_m k (V_m^h - 1)(V_m^h - a) V_m^h v_h d\omega + \\ \quad + \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \chi_m w_h v_h d\omega = (I_i^{ext}, v_h), \quad \forall v_h \in V_h^p, \\ - \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \chi_m c_m \frac{\partial V_m^h}{\partial t} v_h d\omega + a_e(\phi_e^h, v_h) - \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \chi_m k (V_m^h - 1)(V_m^h - a) V_m^h v_h d\omega + \\ \quad - \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \chi_m w_h v_h d\omega = (-I_e^{ext}, v_h), \quad \forall v_h \in V_h^p, \\ \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \frac{\partial w_h}{\partial t} v_h d\omega = \sum_{\mathcal{K} \in \tau_h} \int_{\mathcal{K}} \epsilon (V_m^h - \gamma w_h) v_h d\omega, \quad \forall v_h \in V_h^p, \end{array} \right.$$

where:

- $$a_l(\phi_l^h, v_h) = \sum_{K \in \tau_h} \int_K (\Sigma_l \nabla_h \phi_l^h) \cdot \nabla_h v_h d\omega - \sum_{F \in \mathcal{F}_h^I} \int_F \{ \{ \Sigma_l \nabla_h \phi_l^h \} \} \cdot [[v_h]] d\sigma +$$

$$- \delta \sum_{F \in \mathcal{F}_h^I} \int_F \{ \{ \Sigma_l \nabla_h v_h \} \} \cdot [[\phi_l^h]] d\sigma + \sum_{F \in \mathcal{F}_h^I} \int_F \Gamma [[\phi_l^h]] \cdot [[v_h]] d\sigma \quad l = i, e,$$
- $$(I_i^{ext}, v_h) = \sum_{K \in \tau_h} \int_K I_i^{ext} v_h d\omega + \int_{\partial\Omega} b_i v_h d\sigma,$$
- $$(-I_e^{ext}, v_h) = - \sum_{K \in \tau_h} \int_K I_e^{ext} v_h d\omega + \int_{\partial\Omega} b_e v_h d\sigma.$$

Semi-implicit scheme

Idea:

- treat most of the terms of the PDE implicitly,
- treat the non-linear term semi-implicitly,
- treat the ODE implicitly with the exception of the term V_m .

Semi-implicit discretized system

Find $\Phi^{n+1} = [\phi_i^{n+1} \phi_e^{n+1}]^T$ and $w^{n+1} \forall n = 0, \dots, N-1$ such that:

$$\begin{cases} (\frac{1}{\Delta t} + \epsilon\gamma)Mw^{n+1} = \epsilon MV_m^n + \frac{M}{\Delta t} w^n, \\ (B + C_{nl}(V_m^n))\Phi^{n+1} = r^{n+1}. \end{cases}$$

Godunov operator-splitting scheme

The main feature is the sub-division of the problem into two different problems to be solved sequentially, such that $L(u) = L_1(u) + L_2(u)$. In our case:

1:

$$\begin{cases} \chi_m C_m M \frac{\hat{V}_m^{n+1} - V_m^n}{\Delta t} + C(V_m^n) V_m^n + \chi_m M w^n = 0, \\ \frac{w^{n+1} - w^n}{\Delta t} = \epsilon(V_m^n - \gamma w^n). \end{cases}$$

2:

$$\begin{cases} \chi_m C_m M \frac{V_m^{n+1} - \hat{V}_m^{n+1}}{\Delta t} + A_i \phi_i^{n+1} = F_i^{n+1}, \\ -\chi_m C_m M \frac{V_m^{n+1} - \hat{V}_m^{n+1}}{\Delta t} + A_e \phi_e^{n+1} = F_e^{n+1}. \end{cases}$$

Godunov operator-splitting discretized system

Find $\Phi^{n+1} = [\phi_i^{n+1} \phi_e^{n+1}]^T$ and $w^{n+1} \quad \forall n = 0, \dots, N-1$ such that:

$$\begin{cases} \left(\frac{\chi_m C_m}{\Delta t} \begin{bmatrix} M & -M \\ M & -M \end{bmatrix} + \begin{bmatrix} A_i & 0 \\ 0 & -A_e \end{bmatrix} \right) \begin{bmatrix} \phi_i^{n+1} \\ \phi_e^{n+1} \end{bmatrix} = \begin{bmatrix} F_i^{n+1} \\ -F_e^{n+1} \end{bmatrix} + \\ -\chi_m \begin{bmatrix} M & 0 \\ 0 & M \end{bmatrix} \begin{bmatrix} w^n \\ w^n \end{bmatrix} + \left(\frac{\chi_m C_m}{\Delta t} \begin{bmatrix} M & 0 \\ 0 & M \end{bmatrix} - \begin{bmatrix} C(V_m^n) & 0 \\ 0 & C(V_m^n) \end{bmatrix} \right) \begin{bmatrix} V_m^n \\ V_m^n \end{bmatrix}, \\ w^{n+1} = (1 - \epsilon \gamma \Delta t) w^n + \epsilon \Delta t V_m^n. \end{cases}$$

Quasi-implicit operator-splitting scheme

Idea:

- sub-division of the operator as Godunov operator-splitting
- treat implicitly all the terms except the cubic one

In this case:

1:

$$\begin{cases} \chi_m C_m M \frac{\tilde{V}_m^{n+1} - V_m^n}{\Delta t} + C(V_m^n) V_m^{n+1} + \chi_m M w^{n+1} = 0, \\ \frac{w^{n+1} - w^n}{\Delta t} = \epsilon (V_m^{n+1} - \gamma w^{n+1}). \end{cases}$$

2:

$$\begin{cases} \chi_m C_m M \frac{V_m^{n+1} - \tilde{V}_m^{n+1}}{\Delta t} + A_i \phi_i^{n+1} = F_i^{n+1}, \\ -\chi_m C_m M \frac{V_m^{n+1} - \tilde{V}_m^{n+1}}{\Delta t} + A_e \phi_e^{n+1} = F_e^{n+1}. \end{cases}$$

Quasi-implicit operator-splitting discretized system

Find $\Phi^{n+1} = [\phi_i^{n+1} \phi_e^{n+1}]^T$ and $w^{n+1} \quad \forall n = 0, \dots, N-1$ such that:

$$\begin{cases} \left(\begin{bmatrix} Q_n & -Q_n \\ Q_n & -Q_n \end{bmatrix} + \begin{bmatrix} A_i & 0 \\ 0 & -A_e \end{bmatrix} \right) \begin{bmatrix} \phi_i^{n+1} \\ \phi_e^{n+1} \end{bmatrix} = \begin{bmatrix} R_n \\ R_n \end{bmatrix} + \begin{bmatrix} F_i^{n+1} \\ -F_e^{n+1} \end{bmatrix}, \\ w^{n+1} = \frac{w^n + \epsilon \Delta t (\phi_i^{n+1} - \phi_e^{n+1})}{1 + \epsilon \gamma \Delta t}. \end{cases}$$

Uniqueness

About uniqueness of the unknowns:

- V_m, w proved in *Existence and uniqueness of the solution for the bidomain model used in cardiac electrophysiology* by Y. Bourgault, Y. Coudière, and C. Pierre.
- ϕ_i, ϕ_e appear only through their difference V_m or their gradient. This means that there cannot be uniqueness.

Uniqueness of potentials

Theorem

The classical solutions ϕ_i, ϕ_e are unique up to a constant depending only on time.

Namely:

Suppose now there exist two couples $(\phi_i^1, \phi_e^1), (\phi_i^2, \phi_e^2)$ of potentials solutions of the Bidomain problem.

$$\exists \tilde{\varphi} : [0, T] \rightarrow \mathbb{R} \text{ such that } \phi_i^1(x, t) - \phi_i^2(x, t) = \phi_e^1(x, t) - \phi_e^2(x, t) = \tilde{\varphi}(t)$$

$$\forall x \in \Omega, \forall t \in [0, T].$$

SO HOW TO RESOLVE IT?

- 1 Imposition of the value of the function in a specific point.
- 2 Imposition of the function mean value.

Imposition of a value in a specific point

Focus on only one of the potentials ϕ_i .

$$\phi_i(\bar{x}, t) = \varphi(t) \quad \forall t \in [0, T] \quad \xrightarrow[\text{with } \{u_j\} \text{ as vector of solution}]{\text{Numerical version}} \quad u_l^n = \varphi(t^n) \quad \forall n \in \{1, N\}$$