

Complex Geometry Project

April 24, 2022

Akshunna S. Dogra^{a1, 2}

¹Department of Mathematics, Imperial College London

²EPSRC CDT in Mathematics of Random Systems: Analysis, Modelling and Simulation

Imperial College London

^aadogra@nyu.edu

Abstract

In this essay we go over the proof of the isomorphism between the Čech cohomology $\check{H}(M; \mathbb{R})$ and the de Rham cohomology $H_{\text{dR}}(M; \mathbb{R})$ of a smooth manifold M .

Definition 0.1 • \mathcal{A}^p sheaf of smooth p -forms on M

• \mathcal{Z}^p sheaf of closed p -forms on M

Remark 0.2 For $p = 0$, \mathcal{Z}^0 is simply the constant sheaf $\underline{\mathbb{R}}$.

Lemma 0.3 For $p > 0$ and all $q \geq 0$, the Čech cohomology groups of the sheaves \mathcal{A}^q are trivial, i.e. $\check{H}^p(M, \mathcal{A}^q) = 0$.

Proof. It is enough to prove that this holds for any locally finite cover \mathcal{U} of M . Let $\mathcal{U} = (U_\alpha)_{\alpha \in I}$ be such a cover and let $(\varrho_\alpha)_{\alpha \in I}$ be a corresponding subordinate, i.e. $\text{supp}(\varrho_\alpha) \subset U_\alpha$, partition of unity.

Let $\omega \in Z^p(\mathcal{U}, \mathcal{A}^q)$ we have to find a $\psi \in C^{p-1}(\mathcal{U}, \mathcal{A}^q)$ such that $d\psi = \omega$. ω consists of a collection of maps

$$\omega_{\alpha_1 \dots \alpha_p} : \bigcap_{i=1}^n U_{\alpha_i} \longrightarrow \mathcal{A}^q \left(\bigcap_{i=1}^n U_{\alpha_i} \right)$$

for any combination of p distinct indices $\alpha_1, \dots, \alpha_p \in I$. For indices $\alpha_1, \dots, \alpha_{p-1} \in I$ we define ψ to be

$$\psi_{\alpha_1 \dots \alpha_{p-1}} = \sum_{\beta \in I} \varrho_\beta \omega_{\beta \alpha_1 \dots \alpha_{p-1}}.$$

We have to prove that $\omega = d\psi$. On $\bigcap_{i=1}^p U_{\alpha_i}$

$$\begin{aligned}
(d\psi)_{\alpha_1 \dots \alpha_p} &= - \sum_{i=1}^p (-1)^i \psi_{\alpha_1 \dots \widehat{\alpha_i} \dots \alpha_p} |_{U_{\alpha_i}} \\
&= - \sum_{i=1}^p \sum_{\beta \in I} (-1)^i \varrho_\beta |_{U_{\alpha_i}} \omega_{\beta \alpha_1 \dots \widehat{\alpha_i} \dots \alpha_p} |_{U_{\alpha_i}} \\
&= - \sum_{i=1}^p \sum_{\beta \in I} (-1)^i \varrho_\beta |_{\bigcap_{j=1}^p U_{\alpha_j}} \omega_{\beta \alpha_1 \dots \widehat{\alpha_i} \dots \alpha_p} |_{U_{\alpha_i}} \\
&= - \sum_{\beta \in I} \varrho_\beta |_{\bigcap_{j=1}^p U_{\alpha_j}} \sum_{i=1}^p (-1)^i \omega_{\beta \alpha_1 \dots \widehat{\alpha_i} \dots \alpha_p} |_{U_{\alpha_i}} \\
&= \sum_{\beta \in I} \varrho_\beta |_{\bigcap_{j=1}^p U_{\alpha_j}} \omega_{\alpha_1 \dots \alpha_p} |_{U_\beta} \\
&= \sum_{\beta \in I} \varrho_\beta \omega_{\alpha_1 \dots \alpha_p} = \omega_{\alpha_1 \dots \alpha_p}.
\end{aligned}$$

where in the third line we used that on $\bigcap_{i=1}^n U_{\alpha_i}$ and all $i \in \{1, \dots, p\}$

$$\varrho_\beta |_{\bigcap_{j=1}^p U_{\alpha_j}} \equiv \varrho_\beta |_{U_{\alpha_i}},$$

in the forth the fact that $\omega \in Z^p$ and thus

$$\omega_{\alpha_1 \dots \alpha_p} |_{U_\beta} + \sum_{i=1}^p (-1)^i \omega_{\beta \alpha_1 \dots \widehat{\alpha_i} \dots \alpha_p} |_{U_{\alpha_i}} = 0,$$

in the last line that on $\bigcap_{i=1}^n U_{\alpha_i}$

$$\varrho_\beta |_{\bigcap_{j=1}^p U_{\alpha_j}} \omega_{\alpha_1 \dots \alpha_p} |_{U_\beta} \equiv \varrho_\beta \omega_{\alpha_1 \dots \alpha_p},$$

and that $(\varrho_\beta)_\beta$ is a partition of unity. □

Lemma 0.4 *The sheaf cochain complex*

$$0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{A}^0 \xrightarrow{d} \mathcal{A}^1 \xrightarrow{d} \mathcal{A}^2 \xrightarrow{d} \dots$$

is exact, that is it splits into short exact sequences

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathbb{R} & \longrightarrow & \mathcal{A}^0 & \xrightarrow{d} & \mathcal{A}^1 & \longrightarrow & 0 \\
& & & & \vdots & & & & \\
0 & \longrightarrow & \mathcal{A}^p & \hookrightarrow & \mathcal{A}^p & \xrightarrow{d} & \mathcal{A}^{p+1} & \longrightarrow & 0 \\
& & & & \vdots & & & &
\end{array}$$

Proof. By definition, the cochain complex is exact iff it is exact on stalks. Thus, it suffices to prove that for every $x \in M$ there exists $U \subset M$ open and sufficiently small, s.t.

$$\cdots \longrightarrow \mathcal{A}^{p-1}(U) \xrightarrow{d} \mathcal{A}^p(U) \xrightarrow{d} \mathcal{A}^{p+1}(U) \xrightarrow{d} \cdots$$

is exact as a cochain complex of Abelian groups. However, this is exactly the content of the classical Poincaré Lemma, see for example [Lee12, Theorem 17.14]. If U is diffeomorphic to a star-shaped open subset of \mathbb{R}^n , e.g. diffeomorphic via a chart to an open ball, then every closed differential form is exact. The second statement follows then as $\ker(d^p|_U) = \mathcal{Z}^p(U) = \operatorname{im}(d^{p+1}|_U)$. \square

Theorem 0.5 (De Rham's Theorem) *There exists an isomorphism*

$$\check{H}(M; \mathbb{R}) \cong H_{\text{dR}}(M; \mathbb{R}).$$

Proof. From the short exact sequences in 0.4 we can deduce the existence of the following corresponding long exact sequences on cohomology: for all $p, q \in \mathbb{N}$

$$\begin{aligned} \cdots \longrightarrow H^{p-1}(M, \mathcal{A}^{q-1}) &\longrightarrow H^{p-1}(M, \mathcal{Z}^q) \longrightarrow \\ &\longrightarrow H^p(M, \mathcal{Z}^{q-1}) \longrightarrow H^p(M, \mathcal{A}^{q-1}) \longrightarrow \cdots \end{aligned}$$

For $p > 1$ the leftmost and rightmost terms are zero thus

$$H^{p-1}(M, \mathcal{Z}^q) \cong H^p(M, \mathcal{Z}^{q-1}).$$

This gives us the following sequence of isomorphisms

$$\begin{aligned} \check{H}^p(M, \mathbb{R}) &:= H^p(M, \mathbb{R}) \cong H^{p-1}(M, \mathcal{Z}^1) \cong \\ &\cong H^{p-2}(M, \mathcal{Z}^2) \cong \cdots \cong H^1(M, \mathcal{Z}^{p-1}). \end{aligned}$$

For $p = 1$ (in the above sense) we get the following exact sequence

$$H^0(M, \mathcal{A}^{p-1}) \longrightarrow H^0(M, \mathcal{Z}^p) \longrightarrow H^1(M, \mathcal{Z}^{p-1}) \longrightarrow 0.$$

Since the 0th order cohomology is simply the space of global sections the above is isomorphic

$$\mathcal{A}^{p-1}(M) \xrightarrow{d} \mathcal{Z}^p(M) \longrightarrow H^1(M, \mathcal{Z}^{p-1}) \longrightarrow 0.$$

By exactness it follows therefore that

$$H^1(M, \mathcal{Z}^{p-1}) \cong \mathcal{Z}^p(M) / d\mathcal{A}^{p-1}(M) =: H_{\text{dR}}^p(M, \mathbb{R}).$$

This proves the assertion. \square

References

- [Lee12] J. LEE. *Introduction to Smooth Manifolds*. Graduate Texts in Mathematics, 218. Springer New York, New York, NY, 2nd ed., 2012.