

A From established ADL difficulty to the next observed state
Age ≥60 years at biomarker baseline; five harmonised ADL items
HRS primary cohort
2016 Venous Blood Study
Function: 2016–2022
Index visit: ≥1 ADL difficulty
3,712 adjacent-wave intervals
346 unknown outcomes excluded
Eligible analysis risk set
2,014 participants · 3,239 intervals
Recovery / Persistent difficulty / Death
127 further exclusions: biomarkers, covariates or weights
CHARLS
exploratory comparison
2011 blood panel
Function: 2011, 2013,
2015, 2018
Index visit: ≥1 ADL difficulty
3,469 adjacent-wave intervals
205 unknown outcomes excluded
Eligible analysis risk set
1,463 participants · 2,306 intervals
Recovery / Persistent difficulty / Death
958 further exclusions: biomarkers, covariates or weights
B Observed states at the next wave
Eligible intervals (%)
n = 3,239
n = 547
n = 1,776
n = 916
n = 2,306
n = 265
n = 1,183
n = 858
HRS
CHARLS
Recovery
Persistent difficulty
Death
C Distribution of the signed difference
Density
HRS
CHARLS
P10
P50
P90
P10
P50
P90
Signed eGFRcys − eGFRcr difference
(mL/min/1.73 m²)