



NUMBER SETS

Adam Klepáč

November 20, 2023

CONTENTS

Natural Numbers

Unpacking The Axioms

Operations On Natural Numbers

Integers

Digression



Mathematical Structures

Destructive & Symmetric Transformations

NATURAL NUMBERS

NATURAL NUMBERS – INTUITION

Natural numbers are intuitively objects which represent a **quantity**.



NATURAL NUMBERS – INTUITION

Natural numbers are intuitively objects which represent a **quantity**.
They're the following set:


$$\mathbb{N} = \{0, 1, 2, 3, \dots\}.$$

NATURAL NUMBERS – INTUITION

Natural numbers are intuitively objects which represent a **quantity**.
They're the following set:

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}.$$

A good way to think about them is to view them as '*collections of blocks*'. You get the next natural number by adding another block on top of the previous collection.



NATURAL NUMBERS – DEFINITION


There are many ways to define natural numbers.



NATURAL NUMBERS – DEFINITION

There are many ways to define natural numbers.


One of the popular ones is using a **successor** function, denoted s .



NATURAL NUMBERS – DEFINITION

There are many ways to define natural numbers.

One of the popular ones is using a **successor** function, denoted s . If n is a natural number, then $s(n)$ basically means ‘add another block on top of n ’.



NATURAL NUMBERS – DEFINITION

There are many ways to define natural numbers.

One of the popular ones is using a **successor** function, denoted s . If n is a natural number, then $s(n)$ basically means ‘add another block on top of n ’.


One would be of course tempted to write

$$s(n) = n + 1$$

but that **doesn't make any sense**. We **don't have addition yet!** In fact, you need the successor function to define addition in the first place.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:



NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.
3. The number 0 is not the successor of any natural number.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.
3. The number 0 is not the successor of any natural number.
4. If $s(n) = s(m)$, then $n = m$.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.
3. The number 0 is not the successor of any natural number.
4. If $s(n) = s(m)$, then $n = m$.
5. (Induction Axiom) If a statement is true for 0 and it being true for n also implies that it is true for $s(n)$, then it is true for all natural numbers.

1

UNPACKING THE AXIOMS



NATURAL NUMBERS – AXIOM 1

There exists the natural number 0.

NATURAL NUMBERS – AXIOM 1

There exists the natural number 0.

Hopefully obvious.

NATURAL NUMBERS – AXIOM 2

Every natural number has a successor which is also natural.

NATURAL NUMBERS – AXIOM 2

Every natural number has a successor which is also natural.

Basically means that the natural numbers are an infinite set. You can add another block atop any collection of blocks.

NATURAL NUMBERS – AXIOM 3

The number 0 is not the successor of any natural number.

NATURAL NUMBERS – AXIOM 3

The number 0 is not the successor of any natural number.

Basically means that the natural numbers are infinite only ‘in one direction’. There is a **first** natural number.

NATURAL NUMBERS – AXIOM 4

If $s(n) = s(m)$, then $n = m$.

NATURAL NUMBERS – AXIOM 4

If $s(n) = s(m)$, then $n = m$.

This means that the successor function is **injective** – each natural number has a different successor.

NATURAL NUMBERS – AXIOM 5

If a statement is true for 0 and it being true for n also implies that it is true for $s(n)$, then it is true for all natural numbers.

NATURAL NUMBERS – AXIOM 5

If a statement is true for 0 and it being true for n also implies that it is true for $s(n)$, then it is true for all natural numbers.

This means that any feature of the natural numbers ‘propagates’ via the successor function. Basically, if something is true for 0 and we know that it is true for the next natural number if it is true for the previous one, then it is true for 1 as well. Because it is true for 1, it is true for 2 as well, etc.

2

OPERATIONS ON NATURAL NUMBERS



WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.



WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.

For example, $+$ and \cdot are operations because they take **two** natural numbers and produce **one**.

WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.

For example, $+$ and \cdot are operations because they take **two** natural numbers and produce **one**.

We don't often see them as functions because we don't write them as such. We write $n + m$ instead of $+(n, m)$ and $n \cdot m$ instead of $\cdot(n, m)$.

WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.

For example, $+$ and \cdot are operations because they take **two** natural numbers and produce **one**.

We don't often see them as functions because we don't write them as such. We write $n + m$ instead of $+(n, m)$ and $n \cdot m$ instead of $\cdot(n, m)$.

In this sense, subtraction and division **are not operations!** They take two natural numbers but they **do not produce a natural number**.

ADDITION

We define **addition** on natural numbers by the following two formulae:

- $n + 0 = n$,
- $n + s(m) = s(n + m)$.

ADDITION

We define **addition** on natural numbers by the following two formulae:

- $n + 0 = n$,
- $n + s(m) = s(n + m)$.

We can imagine addition as ‘adding blocks *to the side*’ and the successor function as ‘adding one block *on top*’.

ADDITION

We define **addition** on natural numbers by the following two formulae:

- $n + 0 = n$,
- $n + s(m) = s(n + m)$.

We can imagine addition as ‘adding blocks *to the side*’ and the successor function as ‘adding one block *on top*’.

In this sense, $n + s(m) = s(n + m)$ only means that if you add one block atop m blocks and then n blocks to the side you have the same number of blocks as if you add n blocks next to m blocks and then another on top of that.

ADDITION



ADDITION


$$2 + s(3)$$

ADDITION

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

2 3

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

$2 + s(3)$

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \\ \text{dark blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

$s(2 + 3)$

ADDITION

$$\begin{matrix} \text{dark blue} \\ \text{light blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

2 3

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \\ \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$


$2 + s(3)$

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \\ \text{dark blue} \\ \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

$s(2 + 3)$

Using the formula $n + s(m) = s(n + m)$, one calculates $n + m$ by taking the successor of n , m times.

ADDITION



Using the formula $n + s(m) = s(n + m)$, one calculates $n + m$ by taking the successor of n , m times. Like this:

$$n + 0 = n,$$

$$n + 1 = n + s(0) = s(n + 0) = s(n),$$

$$n + 2 = n + s(1) = s(n + 1) = s(n + s(0)) = s(s(n + 0)) = s(s(n)),$$

⋮

ADDITION – PROPERTIES

Addition of natural numbers satisfies these two properties:

- **Commutativity:**

$$n + m = m + n.$$

ADDITION – PROPERTIES

Addition of natural numbers satisfies these two properties:

- **Commutativity:**


$$n + m = m + n.$$

- **Associativity:**

$$n + (m + k) = (n + m) + k.$$

ADDITION – PROPERTIES

Using blocks, **commutativity** just means that putting n blocks next to m blocks is the same as putting m blocks next to n blocks.



ADDITION – PROPERTIES

Using blocks, **commutativity** just means that putting n blocks next to m blocks is the same as putting m blocks next to n blocks.


$$2 + 3$$

ADDITION – PROPERTIES

Using blocks, **commutativity** just means that putting n blocks next to m blocks is the same as putting m blocks next to n blocks.




$$2 + 3$$



$$3 + 2$$

ADDITION – PROPERTIES

Using blocks, **associativity** just means that putting m blocks next to k blocks and then n more blocks next to those is the same as putting m blocks next to n blocks and then k more blocks next to those.


$$4 + (2 + 3)$$

ADDITION – PROPERTIES

Using blocks, **associativity** just means that putting m blocks next to k blocks and then n more blocks next to those is the same as putting m blocks next to n blocks and then k more blocks next to those.


$$4 + (2 + 3)$$


$$(4 + 2) + 3$$

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m,$

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m$,
- $m \cdot s(n) = m \cdot n + m$.

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m$,
- $m \cdot s(n) = m \cdot n + m$.

We can imagine multiplication $m \cdot n$ by adding a collections of n blocks for every one block in the collection of m blocks.

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m$,
- $m \cdot s(n) = m \cdot n + m$.

We can imagine multiplication $m \cdot n$ by adding a collection of n blocks for every one block in the collection of m blocks.

If we write $s(n) = n + 1$, then the second formula just means that

$$m \cdot s(n) = m \cdot (n + 1) = m \cdot n + m.$$

MULTIPLICATION

$$\begin{array}{c} \blacksquare \\ \blacksquare \end{array} \quad \begin{array}{c} \square \\ \square \\ \square \end{array}$$


2 3

A diagram illustrating multiplication. On the left, there is a network of five nodes connected by lines. To the right of this network, two sets of colored squares are shown vertically. The first set, labeled '2', contains two dark blue squares. The second set, labeled '3', contains three light blue squares. This visual representation likely corresponds to the multiplication problem $2 \times 3 = 6$.

MULTIPLICATION

$$2 \cdot s(3)$$



The diagram shows two separate groups of three blue squares each. The first group is labeled '2' below it, and the second group is labeled '3' below it. This visual representation corresponds to the mathematical expression $2 \cdot s(3)$.


$$2 \cdot s(3)$$


The diagram shows a single 2x3 grid of six blue squares. This visual representation corresponds to the mathematical expression $2 \cdot s(3)$.

MULTIPLICATION


$$\begin{matrix} & \\ \text{2} & \text{3} \end{matrix}$$


$$2 \cdot s(3)$$


$$2 \cdot 3 + 2$$

MULTIPLICATION

$$\begin{matrix} \text{■} \\ \text{■} \end{matrix} \quad \begin{matrix} \text{□} \\ \text{□} \\ \text{□} \end{matrix}$$

2 3

$$\begin{matrix} \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \end{matrix}$$

$2 \cdot s(3)$

$$\begin{matrix} \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \end{matrix} \quad \begin{matrix} \text{□} \\ \text{□} \\ \text{□} \end{matrix} \quad \text{■}$$

$2 \cdot 3 + 2$

The formula $m \cdot s(n) = m \cdot n + m$ allows us to compute $m \cdot n$ by applying it n times.

MULTIPLICATION

$$\begin{matrix} \text{■} \\ \text{■} \end{matrix} \quad \begin{matrix} \text{□} \\ \text{□} \\ \text{□} \end{matrix}$$

2 3

$$\begin{matrix} \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \end{matrix}$$

$2 \cdot s(3)$

$$\begin{matrix} \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \end{matrix} \quad \text{■}$$

$2 \cdot 3 + 2$

The formula $m \cdot s(n) = m \cdot n + m$ allows us to compute $m \cdot n$ by applying it n times. More precisely,

$$m \cdot 1 = m$$

$$m \cdot 2 = m \cdot s(1) = m \cdot 1 + m = m + m$$

$$m \cdot 3 = m \cdot s(2) = m \cdot 2 + m = m \cdot s(1) + m = m \cdot 1 + m + m = m + m + m$$

⋮

MULTIPLICATION – PROPERTIES

- Commutativity:

$$m \cdot n = n \cdot m.$$



MULTIPLICATION – PROPERTIES

- Commutativity:

$$m \cdot n = n \cdot m.$$

- Associativity:

$$m \cdot (n \cdot k) = (m \cdot n) \cdot k.$$

MULTIPLICATION – PROPERTIES

- Commutativity:

$$m \cdot n = n \cdot m.$$

- Associativity:

$$m \cdot (n \cdot k) = (m \cdot n) \cdot k.$$

- Distributivity:


$$m \cdot (n + k) = m \cdot n + m \cdot k.$$



INTEGERS

INTEGERS

Integers or whole numbers come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.



INTEGERS

Integers or **whole numbers** come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.

In mathematical terms, this means introducing **inverses** to numbers **with respect to addition**.

INTEGERS

Integers or whole numbers come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.

In mathematical terms, this means introducing **inverses** to numbers **with respect to addition**.

We typically denote these inverses by the symbol $-$.

INTEGERS

Integers or whole numbers come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.

In mathematical terms, this means introducing **inverses** to numbers **with respect to addition**.

We typically denote these inverses by the symbol $-$.

Beware! ‘Taking inverse’ is a **unary** operation (meaning it acts on **one element**), not a binary one.

1

DIGRESSION



MATHEMATICAL STRUCTURES

DESTRUCTIVE & SYMMETRIC TRANSFORMATIONS



MATHEMATICAL STRUCTURES

MATHEMATICAL STRUCTURE (ALSO ‘UNIVERSE’)

A **mathematical** structure is a **set** with **operations**. If X is the set and op_1, \dots, op_n the operations on X , we write the resulting structure as

$$(X, op_1, \dots, op_n).$$

MATHEMATICAL STRUCTURES

MATHEMATICAL STRUCTURE (ALSO ‘UNIVERSE’)

A **mathematical** structure is a **set** with **operations**. If X is the set and op_1, \dots, op_n the operations on X , we write the resulting structure as

$$(X, op_1, \dots, op_n).$$

The reason mathematical structures are called by some enthusiasts as mathematical ‘universes’ is that they really *are* universes in the broadest sense possible – a bunch of elements with prescribed rules of interaction.

OPERATION ON A SET

OPERATION

An **operation** on a set X is really just a **rule of interaction** between its elements. In symbols, it is a **function**

$$op : X^n \rightarrow X$$

where $X^n \rightarrow X$ just means ‘Take n elements of X and give me back one.’

OPERATION ON A SET

Examples:

- $(\mathbb{N}, +, \cdot)$ is a structure where $+$ and \cdot are **binary** operations (meaning they take two elements and return one). They can be seen as functions $\mathbb{N}^2 \rightarrow \mathbb{N}$.


OPERATION ON A SET

Examples:

- $(\mathbb{N}, +, \cdot)$ is a structure where $+$ and \cdot are **binary** operations (meaning they take two elements and return one). They can be seen as functions $\mathbb{N}^2 \rightarrow \mathbb{N}$.
- $(\{\text{orderings of vertices of a regular } n\text{-gon}\}, r, r^2, \dots, r^n, s_1, \dots, s_n)$ is a structure where r is the rotation by $360^\circ/n$ and s_1, \dots, s_n are all the reflections. They are all **unary** operations (they take one element and return one).

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.



OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.



OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements.

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements. When I multiply $3 \cdot 5 = 15$, I have no way to get back the 3 or the 5.

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements. When I multiply $3 \cdot 5 = 15$, I have no way to get back the 3 or the 5.
- The operation r on the vertices of a regular polygon is **symmetric** – it can be reversed or **inverted**.

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements. When I multiply $3 \cdot 5 = 15$, I have no way to get back the 3 or the 5.
- The operation r on the vertices of a regular polygon is **symmetric** – it can be reversed or **inverted**. If, for example, vertex A is sent to G by this rotation, then rotation in the opposite direction sends G back to A .

OPERATION AS A TRANSFORMATION

All destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.



OPERATION AS A TRANSFORMATION

All destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an (commutative) operation op on a set X and two elements $x, y \in X$.



OPERATION AS A TRANSFORMATION

All destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an (commutative) operation op on a set X and two elements $x, y \in X$.

- Suppose $op(x, y)$ is some element $z \in X$. For the operation op to be **symmetric**, we need two things:

OPERATION AS A TRANSFORMATION

All destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an (commutative) operation op on a set X and two elements $x, y \in X$.

- Suppose $op(x, y)$ is some element $z \in X$. For the operation op to be **symmetric**, we need two things:
 - A special element $e \in X$ (the **identity** element).

OPERATION AS A TRANSFORMATION

All destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an (commutative) operation op on a set X and two elements $x, y \in X$.

- Suppose $op(x, y)$ is some element $z \in X$. For the operation op to be **symmetric**, we need two things:
 - A special element $e \in X$ (the **identity** element).
 - Elements x^* and y^* (the **inverse** elements to x and y) such that

$$op(x, x^*) = op(y, y^*) = e.$$

OPERATION AS A TRANSFORMATION

All destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an (commutative) operation op on a set X and two elements $x, y \in X$.

- Suppose $op(x, y)$ is some element $z \in X$. For the operation op to be **symmetric**, we need two things:
 - A special element $e \in X$ (the **identity** element).
 - Elements x^* and y^* (the **inverse** elements to x and y) such that

$$op(x, x^*) = op(y, y^*) = e.$$

- It's possible X doesn't have those things – which means the operation op cannot be symmetric **on X** , but it can be symmetric on a larger set which does contain those elements.

INVERSE AS A FUNCTION

Inverse can be thought of as a **unary** operation – it takes an element x and gives back its inverse x^* .

INVERSE AS A FUNCTION

Inverse can be thought of as a **unary** operation – it takes an element x and gives back its inverse x^* .

Clearly, the inverse to x^* is x , again.

OPERATION AS A TRANSFORMATION

Motivating examples:

- The operation $+$ on \mathbb{N} **cannot be made symmetric**. Natural numbers do not contain inverses to $+$.

OPERATION AS A TRANSFORMATION

Motivating examples:

- The operation $+$ on \mathbb{N} **cannot be made symmetric**. Natural numbers do not contain inverses to $+$. We fix this by introducing **integers**, basically forcefully adding to each number n its inverse with respect to $+$, that is, the negative number $-n$. The new set is called the **integers**, denoted \mathbb{Z} . The identity element is 0 in this case.

OPERATION AS A TRANSFORMATION

Motivating examples:

- The operation $+$ on \mathbb{N} **cannot be made symmetric**. Natural numbers do not contain inverses to $+$. We fix this by introducing **integers**, basically forcefully adding to each number n its inverse with respect to $+$, that is, the negative number $-n$. The new set is called the **integers**, denoted \mathbb{Z} . The identity element is 0 in this case.
- The operation \cdot on \mathbb{N} cannot be made symmetric. We fix this by introducing **fractions/reciprocals** to elements. The inverse with respect to \cdot of an element z is $\frac{1}{z}$. The identity element is 1.