



NUMBER SETS

Adam Klepáč

December 11, 2023

CONTENTS

Natural Numbers

Unpacking The Axioms

Operations On Natural Numbers

Integers

Digression



Mathematical Structures

Destructive & Symmetric Transformations

Rational Numbers

NATURAL NUMBERS

NATURAL NUMBERS – INTUITION

Natural numbers are intuitively objects which represent a **quantity**.



NATURAL NUMBERS – INTUITION

Natural numbers are intuitively objects which represent a **quantity**.
They're the following set:


$$\mathbb{N} = \{0, 1, 2, 3, \dots\}.$$

NATURAL NUMBERS – INTUITION

Natural numbers are intuitively objects which represent a **quantity**.
They're the following set:

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}.$$

A good way to think about them is to view them as '*collections of blocks*'. You get the next natural number by adding another block on top of the previous collection.



NATURAL NUMBERS – DEFINITION


There are many ways to define natural numbers.



NATURAL NUMBERS – DEFINITION

There are many ways to define natural numbers.


One of the popular ones is using a **successor** function, denoted s .



NATURAL NUMBERS – DEFINITION

There are many ways to define natural numbers.

One of the popular ones is using a **successor** function, denoted s . If n is a natural number, then $s(n)$ basically means ‘add another block on top of n ’.



NATURAL NUMBERS – DEFINITION

There are many ways to define natural numbers.

One of the popular ones is using a **successor** function, denoted s . If n is a natural number, then $s(n)$ basically means ‘add another block on top of n ’.


One would be of course tempted to write

$$s(n) = n + 1$$

but that **doesn't make any sense**. We **don't have addition yet!** In fact, you need the successor function to define addition in the first place.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:



NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.
3. The number 0 is not the successor of any natural number.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.
3. The number 0 is not the successor of any natural number.
4. If $s(n) = s(m)$, then $n = m$.

NATURAL NUMBERS – DEFINITION

The following **five axioms** (often called *Peano axioms*) constitute the definition of natural numbers:

1. There exists the natural number 0.
2. Every natural number has a successor which is also natural.
3. The number 0 is not the successor of any natural number.
4. If $s(n) = s(m)$, then $n = m$.
5. (Induction Axiom) If a statement is true for 0 and it being true for n also implies that it is true for $s(n)$, then it is true for all natural numbers.

1

UNPACKING THE AXIOMS



NATURAL NUMBERS – AXIOM 1

There exists the natural number 0.



NATURAL NUMBERS – AXIOM 1

There exists the natural number 0.

Hopefully obvious.

NATURAL NUMBERS – AXIOM 2

Every natural number has a successor which is also natural.

NATURAL NUMBERS – AXIOM 2

Every natural number has a successor which is also natural.

Basically means that the natural numbers are an infinite set. You can add another block atop any collection of blocks.

NATURAL NUMBERS – AXIOM 3

The number 0 is not the successor of any natural number.

NATURAL NUMBERS – AXIOM 3

The number 0 is not the successor of any natural number.

Basically means that the natural numbers are infinite only ‘in one direction’. There is a **first** natural number.

NATURAL NUMBERS – AXIOM 4

If $s(n) = s(m)$, then $n = m$.

NATURAL NUMBERS – AXIOM 4

If $s(n) = s(m)$, then $n = m$.

This means that the successor function is **injective** – each natural number has a different successor.

NATURAL NUMBERS – AXIOM 5

If a statement is true for 0 and it being true for n also implies that it is true for $s(n)$, then it is true for all natural numbers.

NATURAL NUMBERS – AXIOM 5

If a statement is true for 0 and it being true for n also implies that it is true for $s(n)$, then it is true for all natural numbers.

This means that any feature of the natural numbers ‘propagates’ via the successor function. Basically, if something is true for 0 and we know that it is true for the next natural number if it is true for the previous one, then it is true for 1 as well. Because it is true for 1, it is true for 2 as well, etc.

2

OPERATIONS ON NATURAL NUMBERS



WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.



WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.

For example, $+$ and \cdot are operations because they take **two** natural numbers and produce **one**.

WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.

For example, $+$ and \cdot are operations because they take **two** natural numbers and produce **one**.

We don't often see them as functions because we don't write them as such. We write $n + m$ instead of $+(n, m)$ and $n \cdot m$ instead of $\cdot(n, m)$.

WHAT IS AN OPERATION?

By **operation**, we mean a function which takes **one or multiple** natural numbers and produces **one** natural number.

For example, $+$ and \cdot are operations because they take **two** natural numbers and produce **one**.

We don't often see them as functions because we don't write them as such. We write $n + m$ instead of $+(n, m)$ and $n \cdot m$ instead of $\cdot(n, m)$.

In this sense, subtraction and division **are not operations!** They take two natural numbers but they **do not produce a natural number**.

ADDITION

We define **addition** on natural numbers by the following two formulae:

- $n + 0 = n$,
- $n + s(m) = s(n + m)$.

ADDITION

We define **addition** on natural numbers by the following two formulae:

- $n + 0 = n$,
- $n + s(m) = s(n + m)$.

We can imagine addition as ‘adding blocks *to the side*’ and the successor function as ‘adding one block *on top*’.

ADDITION

We define **addition** on natural numbers by the following two formulae:

- $n + 0 = n$,
- $n + s(m) = s(n + m)$.

We can imagine addition as ‘adding blocks *to the side*’ and the successor function as ‘adding one block *on top*’.

In this sense, $n + s(m) = s(n + m)$ only means that if you add one block atop m blocks and then n blocks to the side you have the same number of blocks as if you add n blocks next to m blocks and then another on top of that.

ADDITION



ADDITION


$$2 + s(3)$$

ADDITION

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

2 3

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

$2 + s(3)$

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \\ \text{dark blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

$s(2 + 3)$

ADDITION

$$\begin{matrix} \text{dark blue} \\ \text{light blue} \end{matrix} \quad \begin{matrix} \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

2 3

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \\ \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$


$2 + s(3)$

$$\begin{matrix} \text{dark blue} \\ \text{dark blue} \\ \text{dark blue} \\ \text{light blue} \\ \text{light blue} \\ \text{light blue} \end{matrix}$$

$s(2 + 3)$

Using the formula $n + s(m) = s(n + m)$, one calculates $n + m$ by taking the successor of n , m times.

ADDITION



Using the formula $n + s(m) = s(n + m)$, one calculates $n + m$ by taking the successor of n , m times. Like this:

$$n + 0 = n,$$

$$n + 1 = n + s(0) = s(n + 0) = s(n),$$

$$n + 2 = n + s(1) = s(n + 1) = s(n + s(0)) = s(s(n + 0)) = s(s(n)),$$

⋮

ADDITION – PROPERTIES

Addition of natural numbers satisfies these two properties:

- **Commutativity:**

$$n + m = m + n.$$

ADDITION – PROPERTIES

Addition of natural numbers satisfies these two properties:

- **Commutativity:**


$$n + m = m + n.$$

- **Associativity:**

$$n + (m + k) = (n + m) + k.$$

ADDITION – PROPERTIES

Using blocks, **commutativity** just means that putting n blocks next to m blocks is the same as putting m blocks next to n blocks.



ADDITION – PROPERTIES

Using blocks, **commutativity** just means that putting n blocks next to m blocks is the same as putting m blocks next to n blocks.


$$2 + 3$$

ADDITION – PROPERTIES

Using blocks, **commutativity** just means that putting n blocks next to m blocks is the same as putting m blocks next to n blocks.




$$2 + 3$$



$$3 + 2$$

ADDITION – PROPERTIES

Using blocks, **associativity** just means that putting m blocks next to k blocks and then n more blocks next to those is the same as putting m blocks next to n blocks and then k more blocks next to those.


$$4 + (2 + 3)$$

ADDITION – PROPERTIES

Using blocks, **associativity** just means that putting m blocks next to k blocks and then n more blocks next to those is the same as putting m blocks next to n blocks and then k more blocks next to those.


$$4 + (2 + 3)$$


$$(4 + 2) + 3$$

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m,$

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m$,
- $m \cdot s(n) = m \cdot n + m$.

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m$,
- $m \cdot s(n) = m \cdot n + m$.

We can imagine multiplication $m \cdot n$ by adding a collections of n blocks for every one block in the collection of m blocks.

MULTIPLICATION

We define **multiplication** on natural numbers by the following formulae:

- $m \cdot 1 = m$,
- $m \cdot s(n) = m \cdot n + m$.

We can imagine multiplication $m \cdot n$ by adding a collection of n blocks for every one block in the collection of m blocks.

If we write $s(n) = n + 1$, then the second formula just means that

$$m \cdot s(n) = m \cdot (n + 1) = m \cdot n + m.$$

MULTIPLICATION

$$\begin{array}{c} \blacksquare \\ \blacksquare \end{array} \quad \begin{array}{c} \square \\ \square \\ \square \end{array}$$

2 3

A diagram illustrating multiplication. On the left, there is a network of five nodes connected by lines. To the right of this network, two sets of colored squares are shown vertically. The first set, labeled '2', contains two dark blue squares. The second set, labeled '3', contains three light blue squares. This visual representation likely corresponds to the multiplication problem $2 \times 3 = 6$.


MULTIPLICATION


$$\begin{matrix} & \text{■} \\ & \text{■} \\ 2 & \end{matrix} \quad \begin{matrix} & \text{□} \\ & \text{□} \\ & \text{□} \\ 3 & \end{matrix}$$

$$\begin{matrix} & \text{□} & \text{□} \\ 2 \cdot s(3) & \end{matrix}$$

MULTIPLICATION


$$\begin{matrix} & \\ \text{2} & \text{3} \end{matrix}$$


$$2 \cdot s(3)$$


$$2 \cdot 3 + 2$$

MULTIPLICATION

$$\begin{matrix} \text{■} \\ \text{■} \end{matrix} \quad \begin{matrix} \text{□} \\ \text{□} \\ \text{□} \end{matrix}$$

2 3

$$\begin{matrix} \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \end{matrix}$$

$2 \cdot s(3)$

$$\begin{matrix} \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \\ \text{□} \end{matrix} \quad \begin{matrix} \text{□} \\ \text{□} \\ \text{□} \end{matrix} \quad \text{■}$$

$2 \cdot 3 + 2$

The formula $m \cdot s(n) = m \cdot n + m$ allows us to compute $m \cdot n$ by applying it n times.

MULTIPLICATION

$$\begin{matrix} \text{■} \\ \text{■} \end{matrix} \quad \begin{matrix} \text{□} \\ \text{□} \\ \text{□} \end{matrix}$$

2 3

$$\begin{matrix} \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \end{matrix}$$

$2 \cdot s(3)$

$$\begin{matrix} \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \\ \text{□} & \text{□} \end{matrix} \quad \text{■}$$

$2 \cdot 3 + 2$

The formula $m \cdot s(n) = m \cdot n + m$ allows us to compute $m \cdot n$ by applying it n times. More precisely,

$$m \cdot 1 = m$$

$$m \cdot 2 = m \cdot s(1) = m \cdot 1 + m = m + m$$

$$m \cdot 3 = m \cdot s(2) = m \cdot 2 + m = m \cdot s(1) + m = m \cdot 1 + m + m = m + m + m$$

⋮

MULTIPLICATION – PROPERTIES

- Commutativity:

$$m \cdot n = n \cdot m.$$



MULTIPLICATION – PROPERTIES

- Commutativity:

$$m \cdot n = n \cdot m.$$

- Associativity:

$$m \cdot (n \cdot k) = (m \cdot n) \cdot k.$$

MULTIPLICATION – PROPERTIES

- Commutativity:

$$m \cdot n = n \cdot m.$$

- Associativity:

$$m \cdot (n \cdot k) = (m \cdot n) \cdot k.$$

- Distributivity:


$$m \cdot (n + k) = m \cdot n + m \cdot k.$$



INTEGERS

INTEGERS

Integers or whole numbers come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.



INTEGERS

Integers or **whole numbers** come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.

In mathematical terms, this means introducing **inverses** to numbers **with respect to addition**.

INTEGERS

Integers or whole numbers come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.

In mathematical terms, this means introducing **inverses** to numbers **with respect to addition**.

We typically denote these inverses by the symbol $-$.

INTEGERS

Integers or whole numbers come about when we say we can not only add blocks on top of collections but we can also **remove** blocks.

In mathematical terms, this means introducing **inverses** to numbers **with respect to addition**.

We typically denote these inverses by the symbol $-$.

Beware! ‘Taking inverse’ is a **unary** operation (meaning it acts on **one element**), not a binary one.

1

DIGRESSION



MATHEMATICAL STRUCTURES

DESTRUCTIVE & SYMMETRIC TRANSFORMATIONS



MATHEMATICAL STRUCTURES

MATHEMATICAL STRUCTURE (ALSO ‘UNIVERSE’)

A **mathematical** structure is a **set** with **operations**. If X is the set and op_1, \dots, op_n the operations on X , we write the resulting structure as

$$(X, op_1, \dots, op_n).$$

MATHEMATICAL STRUCTURES

MATHEMATICAL STRUCTURE (ALSO ‘UNIVERSE’)

A **mathematical** structure is a **set** with **operations**. If X is the set and op_1, \dots, op_n the operations on X , we write the resulting structure as

$$(X, op_1, \dots, op_n).$$

The reason mathematical structures are called by some enthusiasts as mathematical ‘universes’ is that they really *are* universes in the broadest sense possible – a bunch of elements with prescribed rules of interaction.

OPERATION ON A SET

OPERATION

An **operation** on a set X is really just a **rule of interaction** between its elements. In symbols, it is a **function**

$$op : X^n \rightarrow X$$

where $X^n \rightarrow X$ just means ‘Take n elements of X and give me back one.’

OPERATION ON A SET

Examples:

- $(\mathbb{N}, +, \cdot)$ is a structure where $+$ and \cdot are **binary** operations (meaning they take two elements and return one). They can be seen as functions $\mathbb{N}^2 \rightarrow \mathbb{N}$.


OPERATION ON A SET

Examples:

- $(\mathbb{N}, +, \cdot)$ is a structure where $+$ and \cdot are **binary** operations (meaning they take two elements and return one). They can be seen as functions $\mathbb{N}^2 \rightarrow \mathbb{N}$.
- $(\{\text{orderings of vertices of a regular } n\text{-gon}\}, r, r^2, \dots, r^n, s_1, \dots, s_n)$ is a structure where r is the rotation by $360^\circ/n$ and s_1, \dots, s_n are all the reflections. They are all **unary** operations (they take one element and return one).

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.



OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.



OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements.

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements. When I multiply $3 \cdot 5 = 15$, I have no way to get back the 3 or the 5.

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements. When I multiply $3 \cdot 5 = 15$, I have no way to get back the 3 or the 5.
- The operation r on the vertices of a regular polygon is **symmetric** – it can be reversed or **inverted**.

OPERATION AS A TRANSFORMATION

Operations basically describe interactions between set elements – two or more elements combine or **transform** into another one.

This interaction can be **destructive** or **symmetric**.

- The operations $+$ and \cdot on \mathbb{N} are **destructive** – they *destroy* the elements. When I multiply $3 \cdot 5 = 15$, I have no way to get back the 3 or the 5.
- The operation r on the vertices of a regular polygon is **symmetric** – it can be reversed or **inverted**. If, for example, vertex A is sent to G by this rotation, then rotation in the opposite direction sends G back to A .

OPERATION AS A TRANSFORMATION

Most destructive operations on a set X can be made symmetric by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.



OPERATION AS A TRANSFORMATION

Most destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an commutative and associative **binary** operation Δ on a set X and two elements $x, y \in X$.

OPERATION AS A TRANSFORMATION

Most destructive operations on a set X **can be made symmetric** by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an commutative and associative **binary** operation Δ on a set X and two elements $x, y \in X$.

- For the operation Δ to be **symmetric**, we need two things:

OPERATION AS A TRANSFORMATION

Most destructive operations on a set X can be made symmetric by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an commutative and associative **binary** operation Δ on a set X and two elements $x, y \in X$.

- For the operation Δ to be **symmetric**, we need two things:
 - A special element $e \in X$ (the **identity** element) which satisfies

$$x \Delta e = x$$

for all elements $x \in X$.

OPERATION AS A TRANSFORMATION

Most destructive operations on a set X can be made symmetric by making the set larger and introducing **inverses** and **identity elements** with respect to the given operation.

Let's pick an commutative and associative **binary** operation Δ on a set X and two elements $x, y \in X$.

- For the operation Δ to be **symmetric**, we need two things:
 - A special element $e \in X$ (the **identity** element) which satisfies

$$x \Delta e = x$$

for all elements $x \in X$.

- Elements x^* and y^* (the **inverse** elements to x and y) such that

$$x \Delta x^* = y \Delta y^* = e.$$

OPERATION AS A TRANSFORMATION

Let's pick an (commutative) operation Δ on a set X and two elements $x, y \in X$.

- Suppose $x \Delta y$ is some element $z \in X$. For the operation Δ to be **symmetric**, we need two things:
 - A special element $e \in X$ (the **identity** element) which satisfies

$$x \Delta e = x$$

for all elements $x \in X$.

- Elements x^* and y^* (the **inverse** elements to x and y) such that

$$x \Delta x^* = y \Delta y^* = e.$$

- It's possible X doesn't have those things – which means the operation Δ cannot be symmetric **on X** , but it can be symmetric on a larger set which does contain those elements.

OPERATION AS A TRANSFORMATION

For an operation to be invertible, it really just means its **effects can be reversed**, that there's always a way back.



OPERATION AS A TRANSFORMATION

For an operation to be invertible, it really just means its **effects can be reversed**, that there's always a way back.

If x^*, y^* are the inverse to x, y with respect to Δ , then we can **recover** x and y after performing the operation Δ .

OPERATION AS A TRANSFORMATION

For an operation to be invertible, it really just means its **effects can be reversed**, that there's always a way back.

If x^*, y^* are the inverse to x, y with respect to Δ , then we can **recover** x and y after performing the operation Δ .

Suppose $x \Delta y = z$. Then, $z \Delta y^* = x$ and $z \Delta x^* = y$.

OPERATION AS A TRANSFORMATION

For an operation to be invertible, it really just means its **effects can be reversed**, that there's always a way back.

If x^*, y^* are the inverse to x, y with respect to Δ , then we can **recover** x and y after performing the operation Δ .

Suppose $x \Delta y = z$. Then, $z \Delta y^* = x$ and $z \Delta x^* = y$.

Why? We know that $x \Delta x^* = e = y \Delta y^*$.

OPERATION AS A TRANSFORMATION

For an operation to be invertible, it really just means its **effects can be reversed**, that there's always a way back.

If x^*, y^* are the inverse to x, y with respect to Δ , then we can **recover** x and y after performing the operation Δ .

Suppose $x \Delta y = z$. Then, $z \Delta y^* = x$ and $z \Delta x^* = y$.

Why? We know that $x \Delta x^* = e = y \Delta y^*$. Therefore, for example,

$$z \Delta y^* = (x \Delta y) \Delta y^* = x \Delta (y \Delta y^*) = x \Delta e = x.$$

INVERSE AS A FUNCTION

Inverse can be thought of as a **unary** operation – it takes an element x and gives back its inverse x^* .

INVERSE AS A FUNCTION

Inverse can be thought of as a **unary** operation – it takes an element x and gives back its inverse x^* .

Clearly, the inverse to x^* is x , again.

DRAWING STRUCTURES

Let's draw some mathematical structures.




DRAWING STRUCTURES

Let's draw some mathematical structures. Remember that a mathematical structure is just a set with a bunch of operations. If our set X is finite, we can draw its elements as dots.



DRAWING STRUCTURES


Let's draw some mathematical structures. Remember that a mathematical structure is just a set with a bunch of operations. If our set X is finite, we can draw its elements as dots. The simplest way to 'draw' a mathematical structure is a table.



DRAWING STRUCTURES

Let's draw some mathematical structures. Remember that a mathematical structure is just a set with a bunch of operations. If our set X is finite, we can draw its elements as dots. The simplest way to 'draw' a mathematical structure is a table.

Imagine our set has three points a , b and c . We can define a **unary** operation ${}^{\wedge} : X \rightarrow X$ (meaning it takes one element) on X by writing what it does into a table:



DRAWING STRUCTURES

Let's draw some mathematical structures. Remember that a mathematical structure is just a set with a bunch of operations. If our set X is finite, we can draw its elements as dots. The simplest way to 'draw' a mathematical structure is a table.

Imagine our set has three points a , b and c . We can define a **unary** operation ${}^{\wedge} : X \rightarrow X$ (meaning it takes one element) on X by writing what it does into a table:

\hat{a}	\hat{b}	\hat{c}
<hr/>		
c	a	b

DRAWING STRUCTURES

The table

\hat{a}	\hat{b}	\hat{c}
c	a	b


from the previous slide can be also drawn using dots and arrows for example like this:

DRAWING STRUCTURES

The table

\hat{a}	\hat{b}	\hat{c}
<hr/>		
c	a	b


from the previous slide can be also drawn using dots and arrows for example like this:



where the magenta arrows denote the operation \wedge .


DRAWING STRUCTURES

Let's think what could be an **inverse** to the operation \wedge .



DRAWING STRUCTURES


Let's think what could be an **inverse** to the operation \wedge .



What the operation \wedge does is basically changing a dot into another dot following the **arrows**.

DRAWING STRUCTURES

Let's think what could be an **inverse** to the operation \wedge .




What the operation \wedge does is basically changing a dot into another dot following the **arrows**.

As I have three dots, this means, that after traversing one **arrow**, I have to **traverse two arrows** to get back to where I started.

DRAWING STRUCTURES

Let's think what could be an **inverse** to the operation \wedge .




What the operation \wedge does is basically changing a dot into another dot following the **arrows**.

As I have three dots, this means, that after traversing one **arrow**, I have to **traverse two arrows** to get back to where I started.

However, traversing two **arrows** just means **applying the operation \wedge twice**.

DRAWING STRUCTURES


Let's think what could be an **inverse** to the operation \wedge .



What this means in symbols, is that if $\hat{c} = b$, then $\hat{b} = c$.

DRAWING STRUCTURES

Let's think what could be an **inverse** to the operation \wedge .



What this means in symbols, is that if $\hat{c} = b$, then $\hat{\hat{b}} = c$.

So, to get an inverse element with respect to \wedge , we just need to apply \wedge twice.

DRAWING STRUCTURES – EXAMPLES

Imagine a similar unary operation $\tilde{\cdot}$. It is given by the following table


\tilde{a}	\tilde{b}	\tilde{c}
b	b	b

DRAWING STRUCTURES – EXAMPLES

Imagine a similar unary operation $\tilde{\cdot}$. It is given by the following table

\tilde{a}	\tilde{b}	\tilde{c}
<hr/>		
b	b	b

and can be pictured like this




DRAWING STRUCTURES – EXAMPLES

Imagine a similar unary operation $\tilde{\cdot}$. It is given by the following table


\tilde{a}	\tilde{b}	\tilde{c}
<hr/>		
b	b	b

and can be pictured like this




How many elements do we need to add to make $\tilde{\cdot}$ symmetric?


DRAWING STRUCTURES – EXAMPLES



Can we make \sim symmetric?



DRAWING STRUCTURES – EXAMPLES



Can we make \sim symmetric?

The answer is **no** because I can never get back after reaching b . This happens since all the arrows end in b .

DRAWING STRUCTURES – EXAMPLES

Let's now imagine a structure with a binary operation Δ instead. Let's upgrade the number of elements to 4 and label them a , b , c and d .



DRAWING STRUCTURES – EXAMPLES

Let's now imagine a structure with a binary operation Δ instead. Let's upgrade the number of elements to 4 and label them a , b , c and d .

To represent a binary operation using a table, we need to write each element to each row and each column.



DRAWING STRUCTURES – EXAMPLES

Let's now imagine a structure with a binary operation \triangle instead. Let's upgrade the number of elements to 4 and label them a , b , c and d .

To represent a binary operation using a table, we need to write each element to each row and each column.

Suppose our operation \triangle acts like this:

\triangle	a	b	c	d
a	c	a	d	b
b	a	b	c	d
c	d	c	b	a
d	b	d	a	c

DRAWING STRUCTURES – EXAMPLES

Let's now imagine a structure with a binary operation \triangle instead. Let's upgrade the number of elements to 4 and label them a , b , c and d .

To represent a binary operation using a table, we need to write each element to each row and each column.


Suppose our operation \triangle acts like this:

\triangle	a	b	c	d
a	c	a	d	b
b	a	b	c	d
c	d	c	b	a
d	b	d	a	c

Is this operation **symmetric** or **destructive**?

DRAWING STRUCTURES – EXAMPLES

Drawing binary operations as arrows between elements would be a nightmare. Just watch what such a diagram would look like for the operation from the previous slide.



DRAWING STRUCTURES – EXAMPLES

Let's change the operation Δ a little.



DRAWING STRUCTURES – EXAMPLES

Let's change the operation \triangle a little.

\triangle	a	b	c	d
a	c	d	d	b
b	d	b	c	d
c	d	c	b	d
d	b	d	d	c

DRAWING STRUCTURES – EXAMPLES

Let's change the operation \triangle a little.

\triangle	a	b	c	d
a	c	d	d	b
b	d	b	c	d
c	d	c	b	d
d	b	d	d	c

Is it still **symmetric**?

OPERATION AS A TRANSFORMATION

Motivating examples:

- The operation $+$ on \mathbb{N} **cannot be made symmetric**. Natural numbers do not contain inverses to $+$.

OPERATION AS A TRANSFORMATION

Motivating examples:

- The operation $+$ on \mathbb{N} **cannot be made symmetric**. Natural numbers do not contain inverses to $+$. We fix this by introducing **integers**, basically forcefully adding to each number n its inverse with respect to $+$, that is, the negative number $-n$. The new set is called the **integers**, denoted \mathbb{Z} . The identity element is 0 in this case.

OPERATION AS A TRANSFORMATION

Motivating examples:

- The operation $+$ on \mathbb{N} **cannot be made symmetric**. Natural numbers do not contain inverses to $+$. We fix this by introducing **integers**, basically forcefully adding to each number n its inverse with respect to $+$, that is, the negative number $-n$. The new set is called the **integers**, denoted \mathbb{Z} . The identity element is 0 in this case.
- The operation \cdot on \mathbb{Z} cannot be made symmetric. We fix this by introducing **fractions/reciprocals** to elements. The inverse with respect to \cdot of an element z is $\frac{1}{z}$. The identity element is 1.

DEFINING INTEGERS

We define the set of integers (or whole numbers) \mathbb{Z} as the set of natural numbers \mathbb{N} plus the inverses to all natural numbers with respect to addition.



DEFINING INTEGERS

We define the set of integers (or whole numbers) \mathbb{Z} as the set of natural numbers \mathbb{N} plus the inverses to all natural numbers with respect to addition.

We define addition on integers by the formulae

- $n + (-m) = x$ where $x + m = n$ for $n \geq m$ and
- $n + (-m) = -(m + (-n))$ for $n \leq m$.

DEFINING INTEGERS

We define the set of integers (or whole numbers) \mathbb{Z} as the set of natural numbers \mathbb{N} plus the inverses to all natural numbers with respect to addition.

We define addition on integers by the formulae

- $n + (-m) = x$ where $x + m = n$ for $n \geq m$ and
- $n + (-m) = -(m + (-n))$ for $n \leq m$.

and multiplication simply by

$$m \cdot (-n) = -(n \cdot m).$$

DEFINING INTEGERS

We define the set of integers (or whole numbers) \mathbb{Z} as the set of natural numbers \mathbb{N} plus the inverses to all natural numbers with respect to addition.

We define addition on integers by the formulae

- $n + (-m) = x$ where $x + m = n$ for $n \geq m$ and
- $n + (-m) = -(m + (-n))$ for $n \leq m$.

and multiplication simply by

$$m \cdot (-n) = -(n \cdot m).$$

The operation $+$ on \mathbb{Z} is symmetric, while \cdot is still **destructive**. This leads us to the rationals.

RATIONAL NUMBERS

DEFINING RATIONAL NUMBERS


The rational numbers are an extension of the integers when we add all the **reciprocals** – fractions $1/z$ for $z \in \mathbb{Z}$.



DEFINING RATIONAL NUMBERS

The rational numbers are an extension of the integers when we add all the **reciprocals** – fractions $1/z$ for $z \in \mathbb{Z}$.

However, rational numbers also have a strong *intuitive* meaning – they represent **parts of a whole**.



DEFINING RATIONAL NUMBERS

The rational numbers are an extension of the integers when we add all the **reciprocals** – fractions $1/z$ for $z \in \mathbb{Z}$.

However, rational numbers also have a strong *intuitive* meaning – they represent **parts of a whole**.

In this sense, it's logical to write things like

$$\frac{3}{6} = \frac{1}{2}$$

because both fractions represent the same amount, essentially.

DEFINING RATIONAL NUMBERS

The rational numbers are an extension of the integers when we add all the **reciprocals** – fractions $1/z$ for $z \in \mathbb{Z}$.

However, rational numbers also have a strong *intuitive* meaning – they represent **parts of a whole**.

In this sense, it's logical to write things like

$$\frac{3}{6} = \frac{1}{2}$$

because both fractions represent the same amount, essentially.

Nonetheless, as *numbers*, $3 \neq 1$ and $6 \neq 2$. We would like to formalize the notion of *sameness* for two fractions.


DEFINING RATIONAL NUMBERS

Let's try to figure out when fractions a/b and c/d are 'the same'.



DEFINING RATIONAL NUMBERS

Let's try to figure out when fractions a/b and c/d are 'the same'.
This happens if the two fractions represent the same ratio.



DEFINING RATIONAL NUMBERS

Let's try to figure out when fractions a/b and c/d are 'the same'.

This happens if the two fractions represent the same ratio. Notice that this is true **exactly when**

$$a \cdot d = b \cdot c.$$

DEFINING RATIONAL NUMBERS

Let's try to figure out when fractions a/b and c/d are 'the same'.

This happens if the two fractions represent the same ratio. Notice that this is true **exactly when**

$$a \cdot d = b \cdot c.$$

Examples:

- $3/4 = 6/8$ because $3 \cdot 8 = 4 \cdot 6$.

DEFINING RATIONAL NUMBERS

Let's try to figure out when fractions a/b and c/d are 'the same'.

This happens if the two fractions represent the same ratio. Notice that this is true **exactly when**

$$a \cdot d = b \cdot c.$$

Examples:

- $3/4 = 6/8$ because $3 \cdot 8 = 4 \cdot 6$.
- $1/2 = 10/20$ because $1 \cdot 20 = 2 \cdot 10$.

MULTIPLICATION OF RATIONAL NUMBERS

Defining multiplication on rational numbers is easy.

MULTIPLICATION OF RATIONAL NUMBERS

Defining multiplication on rational numbers is easy. We just set

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{a \cdot c}{b \cdot d}.$$

ADDITION OF RATIONAL NUMBERS

Addition is a little trickier.



ADDITION OF RATIONAL NUMBERS

Addition is a little trickier.

The trick is to use the ‘ratio representation’ intuitive view of rational numbers.



ADDITION OF RATIONAL NUMBERS

Addition is a little trickier.

The trick is to use the ‘ratio representation’ intuitive view of rational numbers.

If we want to add a/b to c/d , we choose a common denominator (a common ‘part of a whole’), that is $b \cdot d$, then write a/b as $a \cdot d/b \cdot d$ and c/d as $b \cdot c/b \cdot d$ and add those.

ADDITION OF RATIONAL NUMBERS

Addition is a little trickier.

The trick is to use the ‘ratio representation’ intuitive view of rational numbers.

If we want to add a/b to c/d , we choose a common denominator (a common ‘part of a whole’), that is $b \cdot d$, then write a/b as $a \cdot d/b \cdot d$ and c/d as $b \cdot c/b \cdot d$ and add those.

The result is of course the formula

$$\frac{a}{b} + \frac{c}{d} = \frac{a \cdot d + b \cdot c}{b \cdot d}.$$