



SYSTEMS OF LINEAR EQUATIONS

Adam Klepáč

March 4, 2024

CONTENTS

Functions

Function Composition

Real Functions

Linear Functions

FUNCTIONS

WHAT IS A FUNCTION?


Intuitively, a function is a **box** which receives data and gives some data back.



We'll call the data that a **function receives**, **inputs** and the **data it gives back**, **outputs**. **Inputs** and **outputs** need not necessarily be just 'one object', they can be for example lists of numbers.

FUNCTIONS – EXAMPLE

A function which returns the **average** of a given set of numbers receives the numbers and also their count as **input** and returns the **average** as **output**.



FUNCTIONS – EXAMPLE

We can also consider ‘non-mathematical’ functions. Like a function which receives a type of meal and returns the ingredients.



1

FUNCTION COMPOSITION

FUNCTION COMPOSITION

If we have **two functions**, we can **in certain cases** ‘compose’ them.


Composition simply means that **one function follows the other** – in other words, the **output** of the first function is the **input** of the second.

Of course, **composition** is only possible if the **output** of the first function is a valid **input** for the second.

For instance, you could hardly compose the **ingredients** function with the **average** function.

FUNCTION COMPOSITION


Considering two functions



their composition can look like this




What would the output of this composition look like



FUNCTION COMPOSITION

So, the **order of the composition matters!** Here are a few examples of compositions which make sense:



```
graph LR; A["x1, x2, ..., xn  
n"] --> B[average]; B --> C[times two]; C --> D["2 · (x1 + x2 + ... + xn) / n"]
```

The diagram illustrates the composition of two functions: a list of numbers x_1, x_2, \dots, x_n divided by n leads to the "average", which then leads to "times two", resulting in the formula $2 \cdot \frac{x_1 + x_2 + \dots + x_n}{n}$.

We can of course compose **as many functions** as we like. An example of this:



FUNCTIONS – NOTATION

Drawing pictures like this would be cumbersome. Instead of



we simply write **function(input) = output**.


You're probably used to seeing function written like $f(x) = y$. The picture corresponding to this is



FUNCTIONS – NOTATION

The symbol used for function composition is \circ . It is however a little confusing because it is written in order ‘from right to left’.

For example, if f and g are two functions, their composition $f \circ g$ corresponds to this picture



that is, first g , then f .

2

REAL FUNCTIONS




REAL FUNCTION

A **real function** is simply a function whose **input** and **output** are both real numbers.

Examples of such functions are

- $f(x) = 0$,
- $g(x) = \tan^6(\log^{\sin(x^2+4)}(\frac{5x^3-2}{9x^7}))$,

where $x \in \mathbb{R}$. Or, using pictures,



OPERATIONS ON REAL FUNCTIONS

As both the **input** and the **output** of a real function is a real number, **we can always compose real functions.**

However, that **doesn't mean that the order doesn't matter!** Different order of composition gives different functions.

For example, take $f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$. Then,

- $(f \circ g)(x) = 2 \left(\frac{1}{1+x} \right)^2 + 7$ and
- $(g \circ f)(x) = \frac{1}{1+(2x^2+7)}$.

OPERATIONS ON REAL FUNCTIONS

Real functions can also be **added** and **multiplied**, just like real numbers.

This involves simply adding or multiplying their respective outputs. For the functions

$$f(x) = 2x^2 + 7 \text{ and } g(x) = \frac{1}{1+x}, \text{ their}$$

- **sum** is the function with output

$$(f + g)(x) = f(x) + g(x) = 2x^2 + 7 + \frac{1}{1+x}.$$

- **product** is the function with output

$$(f \cdot g)(x) = f(x) \cdot g(x) = (2x^2 + 7) \cdot \left(\frac{1}{1+x}\right) = \frac{2x^2 + 7}{1+x}.$$

GRAPHS

As real functions have real numbers as inputs and outputs, they can be easily **graphed**.



Graphing a real function f simply means drawing the points $(x, f(x))$ or $(\text{input}, \text{output})$ in some chosen coordinate system.

We typically use the **Cartesian** coordinate system with two axes (one for **input** and one for **output**) that are mutually perpendicular. These are often called the x -axis and the y -axis.

However, later, we'll also use the **polar** coordinate system where every point is instead determined by its angle and distance from the origin of the system.



GRAPHS

The functions $f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$ have the following (parts of) graphs:



GRAPHS

Just to better drive home the idea that the order of function composition is important, look at the graphs of $f \circ g$ and $g \circ f$.



3

LINEAR FUNCTIONS

LINEAR FUNCTION

Linear functions are a special type of real functions whose output is always of the form $a \cdot (\text{input}) + b$ for fixed numbers $a, b \in \mathbb{R}$.

In other words, linear functions are those real functions that only scale the input and add something to it.

LINEAR FUNCTION

A real function f is linear if

$$f(x) = ax + b$$

for some $a, b \in \mathbb{R}$.

LINEAR FUNCTIONS – PROPERTIES

Linear functions have some unique properties:

- If f and g are linear, so are $f \circ g$ and $g \circ f$. Indeed, we can see this easily. Suppose $f(x) = ax + b$ and $g(x) = cx + d$, then

$$(f \circ g)(x) = a(cx + d) + b = (ac)x + (ad + b),$$

$$(g \circ f)(x) = c(ax + b) + d = (ac)x + (cb + d).$$

- If f and g are linear, so is $f + g$. If we just compute the sum, we get

$$(f + g)(x) = (ax + b) + (cx + d) = (a + c)x + (b + d).$$

LINEAR FUNCTIONS – GRAPHS

Graphs of linear functions are straight lines.

Let us see why this is true. Choose a linear function

$$f(x) = ax + b.$$

Pick a number – x . One way to show that the graph of f is a line is to move by two different distances from x and see that we get two similar triangles.

If we move by 1, from x to $x + 1$, then on the y -axis we move from $ax + b$ to $a(x + 1) + b$, that is, we move by

$$a(x + 1) + b - (ax + b) = a.$$

If we move by 2, from x to $x + 2$, on the y -axis, we move

by

$$a(x + 2) + b - (ax + b) = 2a.$$

