



PRIME FACTORIZATIONS GCD & LCM

Adam Klepáč

January 8, 2024

CONTENTS

Divisibility

Chinese Remainder Theorem


Prime Factorization

DIVISIBILITY

The background of the slide features a minimalist design with two large triangles meeting at a point in the center. The triangle on the left is a dark, rich brown, and the triangle on the right is a warm, golden-brown color. The top half of the slide is a solid, very light cream or off-white color, creating a clean, modern aesthetic.

DIVISIBILITY

We say that a number $n \in \mathbb{N}$ **is divisible** by $m \in \mathbb{N}$ or that m **divides** n



DIVISIBILITY

We say that a number $n \in \mathbb{N}$ **is divisible** by $m \in \mathbb{N}$ or that m **divides** n if there exists a number $q \in \mathbb{N}$ (called the **quotient**) such that

$$n = m \cdot q.$$

DIVISIBILITY


We say that a number $n \in \mathbb{N}$ **is divisible** by $m \in \mathbb{N}$ or that m **divides** n if there exists a number $q \in \mathbb{N}$ (called the **quotient**) such that

$$n = m \cdot q.$$

We write the fact that m divides n as $m \mid n$.

DIVISIBILITY

Even if m doesn't divide n , we can always at least **divide with remainder**.



DIVISIBILITY

Even if m doesn't divide n , we can always at least **divide with remainder**.

Formally, for every two numbers $n, m \in \mathbb{N}$, there always exist number $q \in \mathbb{N}$ and $r \leq n$ (called the **remainder**) such that

$$n = m \cdot q + r.$$

INTEGER DIVISION & MODULUS

- The operation which to a pair (n, m) assigns the **quotient** of the division of n by m is called **integer division** and denoted by div .

INTEGER DIVISION & MODULUS

- The operation which to a pair (n, m) assigns the **quotient** of the division of n by m is called **integer division** and denoted by div .
 - For example, $7 \text{ div } 2 = 3$ and $5 \text{ div } 9 = 0$.

INTEGER DIVISION & MODULUS

- The operation which to a pair (n, m) assigns the **quotient** of the division of n by m is called **integer division** and denoted by div .
 - For example, $7 \text{ div } 2 = 3$ and $5 \text{ div } 9 = 0$.
- The operation which to a pair (n, m) assigns the **remainder** of the division of n by m is called **modulus** and denoted mod .

INTEGER DIVISION & MODULUS

- The operation which to a pair (n, m) assigns the **quotient** of the division of n by m is called **integer division** and denoted by div .
 - For example, $7 \text{ div } 2 = 3$ and $5 \text{ div } 9 = 0$.
- The operation which to a pair (n, m) assigns the **remainder** of the division of n by m is called **modulus** and denoted mod .
 - For example, $7 \text{ mod } 2 = 1$ and $5 \text{ mod } 9 = 5$.

INTEGER DIVISION & MODULUS

Using integer division and modulus, we can write the operation of division with remainder like this:

$$n = m \cdot (n \operatorname{div} m) + n \operatorname{mod} m.$$

CHINESE REMAINDER THEOREM

The slide features a white background with two large, overlapping geometric shapes at the bottom. On the left, a dark brown triangle points upwards towards the center. On the right, a tan triangle points downwards towards the center. The two triangles overlap in the middle, creating a darker brown triangular region at the bottom center.

COUNTING SOLDIERS

The famous Chinese military general, strategist and philosopher 孙子 describes in his book 孙子兵法 an efficient method to count the number of soldiers.



COUNTING SOLDIERS

The famous Chinese military general, strategist and philosopher 孙子 describes in his book 孙子兵法 an efficient method to count the number of soldiers.

He instructs the soldiers to arrange into a grid with

- 21 soldiers in each row. He counts the number of soldiers remaining in the last row to be 3.

COUNTING SOLDIERS

The famous Chinese military general, strategist and philosopher 孙子 describes in his book 孙子兵法 an efficient method to count the number of soldiers.

He instructs the soldiers to arrange into a grid with

- 21 soldiers in each row. He counts the number of soldiers remaining in the last row to be 3.
- 25 soldiers in each row. He counts the number of soldiers remaining in the last row to be 6.

COUNTING SOLDIERS

The famous Chinese military general, strategist and philosopher 孙子 describes in his book 孙子兵法 an efficient method to count the number of soldiers.

He instructs the soldiers to arrange into a grid with

- 21 soldiers in each row. He counts the number of soldiers remaining in the last row to be 3.
- 25 soldiers in each row. He counts the number of soldiers remaining in the last row to be 6.
- 32 soldiers in each row. He counts the number of soldiers remaining in the last row to be 23.

COUNTING SOLDIERS

The famous Chinese military general, strategist and philosopher 孙子 describes in his book 孙子兵法 an efficient method to count the number of soldiers.


He instructs the soldiers to arrange into a grid with

- 21 soldiers in each row. He counts the number of soldiers remaining in the last row to be 3.
- 25 soldiers in each row. He counts the number of soldiers remaining in the last row to be 6.
- 32 soldiers in each row. He counts the number of soldiers remaining in the last row to be 23.

孙子 then determines, that there are 1431 soldiers standing before him.

1

PRIME FACTORIZATION



PRIME NUMBER

PRIME NUMBER

A number $n \in \mathbb{N}$ is **prime** if it has **exactly two divisors** – 1 and itself.
We denote the set of all prime numbers by \mathbb{P} .

DECOMPOSITION INTO PRIMES

Prime numbers are somewhat of a building blocks of other numbers.



DECOMPOSITION INTO PRIMES

Prime numbers are somewhat of a building blocks of other numbers.
Each natural number decomposes **uniquely** into a product of prime numbers.



DECOMPOSITION INTO PRIMES

Prime numbers are somewhat of a building blocks of other numbers.

Each natural number decomposes **uniquely** into a product of prime numbers.

Formally,

PRIME FACTORIZATION

For each number $n \in \mathbb{N}$, there exist prime numbers $p_1, \dots, p_m \in \mathbb{P}$ and powers $k_1, \dots, k_m \in \mathbb{N}$ such that

$$n = p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m}.$$

DECOMPOSITION INTO PRIMES

Prime numbers are somewhat of a building blocks of other numbers.
Each natural number decomposes **uniquely** into a product of prime numbers.
Formally,

PRIME FACTORIZATION


For each number $n \in \mathbb{N}$, there exist prime numbers $p_1, \dots, p_m \in \mathbb{P}$ and powers $k_1, \dots, k_m \in \mathbb{N}$ such that

$$n = p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m}.$$

The numbers p_1, \dots, p_m are called the **prime factors** of n .

DECOMPOSITION INTO PRIMES

Currently, there's no known fast algorithm how to decompose a number into its prime factors.



DECOMPOSITION INTO PRIMES

Currently, there's no known fast algorithm how to decompose a number into its prime factors.

The best we can do is to try each prime number from 2 up to \sqrt{n} .

DECOMPOSITION INTO PRIMES

Currently, there's no known fast algorithm how to decompose a number into its prime factors.

The best we can do is to try each prime number from 2 up to \sqrt{n} .

We divide the number in question as many times as possible and proceed to find the next prime factor of the resulting quotient.

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.



DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.



DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445
3	1815

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445
3	1815
3	605

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121
11	11

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121
11	11
11	1

DECOMPOSITION INTO PRIMES – EXAMPLE

Let's decompose the number 16335 into its prime factors.

It's as easy as keeping a table of all the prime factors we've divided the number by.

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121
11	11
11	1

This means that $16335 = 3^3 \cdot 5 \cdot 11^2$.

GREATEST COMMON DIVISOR

GREATEST COMMON DIVISOR

Given two numbers $n, m \in \mathbb{N}$, their **Greatest Common Divisor** (GCD) or Highest Common Factor (HCF) is the largest natural number $k \in \mathbb{N}$ such that $k \mid n$ and $k \mid m$.

GREATEST COMMON DIVISOR

The simplest (and most inefficient) way how to compute the GCD is to decompose the two numbers into prime factors and take all those that the two numbers have in common.



GREATEST COMMON DIVISOR

The simplest (and most inefficient) way how to compute the GCD is to decompose the two numbers into prime factors and take all those that the two numbers have in common.

For example, let's take a look at the decompositions of 16335 and 17325.

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121
11	11
11	1

Prime Factors	Quotients
3	5775
3	1925
5	385
5	77
7	11
11	1

GREATEST COMMON DIVISOR

If we highlight the prime factors the two numbers share

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121
11	11
11	1

Prime Factors	Quotients
3	5775
3	1925
5	385
5	77
7	11
11	1

GREATEST COMMON DIVISOR

If we highlight the prime factors the two numbers share

Prime Factors	Quotients	Prime Factors	Quotients
3	5445	3	5775
3	1815	3	1925
3	605	5	385
5	121	5	77
11	11	7	11
11	1	11	1

we deduce that the GCD of 16335 and 17325 is

$$3^2 \cdot 5 \cdot 11 = 495.$$

GREATEST COMMON DIVISOR

If we highlight the prime factors the two numbers share

Prime Factors	Quotients	Prime Factors	Quotients
3	5445	3	5775
3	1815	3	1925
3	605	5	385
5	121	5	77
11	11	7	11
11	1	11	1

we deduce that the GCD of 16335 and 17325 is

$$3^2 \cdot 5 \cdot 11 = 495.$$

We'll denote the GCD of n and m as $\gcd(n, m)$ or simply as (n, m) .

LEAST COMMON MULTIPLE

LEAST COMMON MULTIPLE

Given $n, m \in \mathbb{N}$ their **Least Common Multiple** (LCM) is the **smallest** number $k \in \mathbb{N}$ such that $n \mid k$ and $m \mid k$.

LEAST COMMON MULTIPLE

LEAST COMMON MULTIPLE

Given $n, m \in \mathbb{N}$ their **Least Common Multiple** (LCM) is the **smallest** number $k \in \mathbb{N}$ such that $n \mid k$ and $m \mid k$. It is given by the formula

$$\text{lcm}(n, m) = \frac{n \cdot m}{\text{gcd}(n, m)}.$$

LEAST COMMON MULTIPLE

The other option how to calculate the least common multiple of two numbers is to take their prime factors elevated on the highest power that appears in at least one of the numbers.

LEAST COMMON MULTIPLE

Explicitly, from the prime decompositions of 16335 and 17325

Prime Factors	Quotients
3	5445
3	1815
3	605
5	121
11	11
11	1

Prime Factors	Quotients
3	5775
3	1925
5	385
5	77
7	11
11	1

LEAST COMMON MULTIPLE

Explicitly, from the prime decompositions of 16335 and 17325

Prime Factors	Quotients	Prime Factors	Quotients
3	5445	3	5775
3	1815	3	1925
3	605	5	385
5	121	5	77
11	11	7	11
11	1	11	1

we see that their LCM is

$$3^3 \cdot 5^2 \cdot 7 \cdot 11^2 = 571725.$$