


PROBABILISTIC INTUITION



Imagine you have 9 balls of different colours.



Imagine you have 9 balls of different colours.





Imagine you have 9 balls of different colours.



• If you pick a ball at random, what colour is it most likely to be?



Imagine you have 9 balls of different colours.

- •
- • •
- • •
- If you pick a ball at random, what colour is it most likely to be?
- How many times more likely is picking a red ball than picking a green ball?



Imagine you have 9 balls of different colours.

- •
- • •
- • •
- If you pick a ball at random, what colour is it most likely to be?
- How many times more likely is picking a red ball than picking a green ball?


QUANTIFYING PROBABILITY



PROBABILITY

A probability is a number between 0 and 1 measuring how likely is something to happen.









what is the probability of picking a ball of a specific colour?



In our example of 9 balls







what is the probability of picking a ball of a specific colour?

- For red, it's 4/9.
- For blue, it's 3/9.
- For green, it's 2/9.



In our example of 9 balls







what is the probability of picking a ball of a specific colour?

- For red, it's 4/9.
- For blue, it's 3/9.
- For green, it's 2/9.

The probabilities above sum up to 1 because I am certain to pick some ball.



We'll all the outcome of a random choice, a random variable and typically write it as X.



We'll all the outcome of a random choice, a random variable and typically write it as *X*. A random variable always lies in the set of all possible outcomes.



We'll all the outcome of a random choice, a **random variable** and typically write it as *X*. A random variable always lies in the set of all possible outcomes. In this case, the variable *X* must lie in the set of possible colours, {red, blue, green}.

Л



We'll all the outcome of a random choice, a random variable and typically write it as X.

A random variable always lies in the set of all possible outcomes.

In this case, the variable *X* must lie in the set of possible colours, {red, blue, green}.

We'll write the probability that X is equal to one of the elements in the set as P(X = colour).



We'll all the outcome of a random choice, a random variable and typically write it as X.

A random variable always lies in the set of all possible outcomes.

In this case, the variable X must lie in the set of possible colours, {red, blue, green}.

We'll write the probability that X is equal to one of the elements in the set as P(X = colour).

So, for the 9-ball example from before, we would have

$$P(X = \text{red}) = \frac{4}{9}$$
, $P(X = \text{blue}) = \frac{3}{9}$, $P(X = \text{green}) = \frac{2}{9}$.

CALCULATING PROBABILITY



In the case the set of outcomes is **finite**, the probability of *X* being one of the possible outcomes is always

CALCULATING PROBABILITY



In the case the set of outcomes is **finite**, the probability of *X* being one of the possible outcomes is always

$$P(X \in S) = \frac{|S|}{|O|},$$

where *S* is a certain subset of *O* – all the possible outcomes.

CALCULATING PROBABILITY - EXAMPLE



We'll describe our 9-ball example more formally.

CALCULATING PROBABILITY - EXAMPLE



We'll describe our 9-ball example more formally.

We'll assign the balls number from 1 to 9. The set of all possible outcomes of picking a random ball is then

$$O = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}.$$

- 1 2
- 4 5
- 7 8 9





- 1 2 3
- 4 5 6
- 7 8 9

We'll form three subsets of O:

$$R = \{2, 6, 7, 9\},$$

$$B = \{3, 4, 5\},$$

$$G = \{1, 8\}.$$





- 1 2 3
- 4 5 6
- 7 8 9

We'll form three subsets of O:

$$R = \{2, 6, 7, 9\},\$$

$$B = \{3, 4, 5\},\$$

$$G = \{1, 8\}.$$

We can use the formula from before to calculate the probability that X will be a green ball:

$$P(X \in G) = \frac{|G|}{|O|} = \frac{2}{9}.$$

PROBABILITY EQUATIONS



What if I asked about the probability that the ball I pick is red or blue?



What if I asked about the probability that the ball I pick is red or blue?
We can literally use the same formula as before. Now, the set of outcomes we're interested in is $R \cup B$ and so



What if I asked about the probability that the ball I pick is red or blue?
We can literally use the same formula as before. Now, the set of outcomes we're interested in is RUB and so

$$P(X \in R \cup B) = \frac{|R \cup B|}{|O|} = \frac{|R| + |B|}{|O|} = \frac{4+3}{9} = \frac{7}{9}.$$



What if I asked about the probability that the ball I pick is red or blue?

We can literally use the same formula as before. Now, the set of outcomes we're interested in is $R \cup B$ and so

$$P(X \in R \cup B) = \frac{|R \cup B|}{|O|} = \frac{|R| + |B|}{|O|} = \frac{4+3}{9} = \frac{7}{9}.$$

However, this example cannot be easily generalized. We'll see why.

SUMS OF PROBABILITIES - COUNTEREXAMPLE

Suppose we're instead choosing from a set of numbers between 1 and 20.





SUMS OF PROBABILITIES - COUNTEREXAMPLE

Suppose we're instead choosing from a set of numbers between 1 and 20. We want to calculate the probability that a randomly picked number is even or divisible by 5.





Suppose we're instead choosing from a set of numbers between 1 and 20.

We want to calculate the probability that a randomly picked number is even or divisible by 5.

So, we have

$$O = \{1, 2, \dots, 20\},$$

 $E = \{2, 4, 6, \dots, 20\},$
 $F = \{5, 10, 15, 20\}.$





Suppose we're instead choosing from a set of numbers between 1 and 20.

We want to calculate the probability that a randomly picked number is even or divisible by 5.

So, we have

$$O = \{1, 2, \dots, 20\},$$

 $E = \{2, 4, 6, \dots, 20\},$
 $F = \{5, 10, 15, 20\}.$

and we want to figure out the probability $P(X \in E \cup F)$.

SUMS OF PROBABILITIES – COUNTEREXAMPLE



Let's try to use the same formula as before:

$$P(X \in E \cup F) = \frac{|E \cup F|}{|O|} \stackrel{??}{=} \frac{|E| + |F|}{|O|} = \frac{10 + 4}{20} = \frac{14}{20}.$$



SUMS OF PROBABILITIES - COUNTEREXAMPLE

Let's try to use the same formula as before:

$$P(X \in E \cup F) = \frac{|E \cup F|}{|O|} \stackrel{??}{=} \frac{|E| + |F|}{|O|} = \frac{10 + 4}{20} = \frac{14}{20}.$$

This doesn't quite add up.





Let's try to use the same formula as before:

$$P(X \in E \cup F) = \frac{|E \cup F|}{|O|} \stackrel{??}{=} \frac{|E| + |F|}{|O|} = \frac{10 + 4}{20} = \frac{14}{20}.$$

This doesn't quite add up.

If we count such numbers by hand, we get the set

$${2,4,5,6,8,10,12,14,15,16,18,20}.$$





Let's try to use the same formula as before:

$$P(X \in E \cup F) = \frac{|E \cup F|}{|O|} \stackrel{??}{=} \frac{|E| + |F|}{|O|} = \frac{10 + 4}{20} = \frac{14}{20}.$$

This doesn't quite add up.

If we count such numbers by hand, we get the set

$${2,4,5,6,8,10,12,14,15,16,18,20}.$$

There's only 12 of them.





Let's try to use the same formula as before:

$$P(X \in E \cup F) = \frac{|E \cup F|}{|O|} \stackrel{??}{=} \frac{|E| + |F|}{|O|} = \frac{10 + 4}{20} = \frac{14}{20}.$$

This doesn't quite add up.

If we count such numbers by hand, we get the set

$${2,4,5,6,8,10,12,14,15,16,18,20}.$$

There's only 12 of them.

The problem is that we counted the numbers 10 and 20 twice!

SUMS OF PROBABILITIES - COUNTEREXAMPLE



Let's try to use the same formula as before:

$$P(X \in E \cup F) = \frac{|E \cup F|}{|O|} \stackrel{??}{=} \frac{|E| + |F|}{|O|} = \frac{10 + 4}{20} = \frac{14}{20}.$$

This doesn't quite add up.

If we count such numbers by hand, we get the set

$${2,4,5,6,8,10,12,14,15,16,18,20}.$$

There's only 12 of them.

The problem is that we counted the numbers 10 and 20 twice!

So, to get the size of $E \cup F$, we cannot just add the size of E to the size of F but we also have to subtract the elements that appear twice – the size of $E \cap F$.

SUMS OF PROBABILITIES - FORMULA



The previous example applies in general. If A, B are two subsets of the set of outcomes, O, then

SUMS OF PROBABILITIES - FORMULA



The previous example applies in general. If A, B are two subsets of the set of outcomes, O, then

$$P(X \in A \cup B) = \frac{|A \cup B|}{|O|} = \frac{|A| + |B| - |A \cap B|}{|O|}.$$