

CONTENTS



Cryptography on Regular Polygons

GENERAL POLYGONS

GENERAL POLYGONS - DEFINITION



POLYGON

A polygon is a closed 2D shape made of only segments.

GENERAL POLYGONS - DEFINITION



POLYGON

A polygon is a closed 2D shape made of only segments.

The endpoints of those segments are called vertices.

GENERAL POLYGONS - DEFINITION



POLYGON


A polygon is a closed 2D shape made of only segments.

The endpoints of those segments are called vertices.

The segments themselves are called edges.


GENERAL POLYGONS – EXAMPLES





GENERAL POLYGONS - EXAMPLES






A polygon with $n \in \mathbb{N}$ sides is called an n-gon.

GENERAL POLYGONS - EXAMPLES






A polygon with $n \in \mathbb{N}$ sides is called an n-gon.

For example a polygon with 123456 sides is called a 123456-gon or decadismyriatrischilliatetrahectapentacontakaihexagon.

GENERAL POLYGONS - COUNTEREXAMPLES





GENERAL POLYGONS - CONVEXITY



CONVEX POLYGON


A polygon is called **convex** if it has no internal angle greater than 180°.


GENERAL POLYGONS - CONVEXITY



CONVEX POLYGON

A polygon is called convex if it has no internal angle greater than 180°.






CONVEX POLYGONS

CONVEX POLYGONS - SPECIAL TYPES






Trapezoid/Trapezium

A convex quadrilateral with at least two parallel sides.


CONVEX POLYGONS - SPECIAL TYPES





A convex quadrilateral with at least A convex quadrilateral with two

two parallel sides.




Parallelogram

pairs of parallel sides.


CONVEX POLYGONS - SPECIAL TYPES






Trapezoid/Trapezium

A convex guadrilateral with at least A convex guadrilateral with two two parallel sides.



Parallelogram

pairs of parallel sides.



Rhombus


An equilateral (all sides of the same length) parallelogram.

CONVEX POLYGONS - DIAGONALS



DIAGONAL IN A CONVEX POLYGON

A diagonal of a **convex** polygon is a segment connecting two of its non-adjacent vertices.




Diagonal in a convex hexagon.

CONVEX POLYGONS - DIAGONALS



DIAGONAL IN A CONVEX POLYGON

A diagonal of a **convex** polygon is a segment connecting two of its non-adjacent vertices.



Diagonal in a convex hexagon.

Voluntary HW: How many different diagonals does a convex *n*-gon have?



TRIANGULATION OF A CONVEX POLYGON

A triangulation of a convex polygon is its division into triangles by non-intersecting diagonals.



TRIANGULATION OF A CONVEX POLYGON

A triangulation of a convex polygon is its division into triangles by non-intersecting diagonals.







Examples of triangulations.



TRIANGULATION OF A CONVEX POLYGON

A triangulation of a convex polygon is its division into triangles by non-intersecting diagonals.







Examples of triangulations.

Voluntary HW: How many different triangulations of an *n*-gon are there?



TRIANGULATION OF A CONVEX POLYGON

A triangulation of a convex polygon is its division into triangles by non-intersecting diagonals.







Examples of triangulations.

Voluntary HW: Find a **non-convex** polygon which **cannot** be triangulated.





Internal angles of a pentagon.

Question: What is the sum of internal angles of a convex *n*-gon?





Internal angles of a pentagon.

Question: What is the sum of internal angles of a convex *n*-gon?

For a triangle, it's 180°.





Internal angles of a pentagon.

Question: What is the sum of internal angles of a convex *n*-gon?

- For a triangle, it's 180°.
- For a square, it's 360°.





Internal angles of a pentagon.

Question: What is the sum of internal angles of a convex *n*-gon?

- For a triangle, it's 180°.
- For a square, it's 360°.
- For a pentagon, it's 540°.



We can count internal angles using triangulations.





We can count internal angles using triangulations. Into how many triangles is a convex *n*-gon divided?



We can count internal angles using triangulations. Into how many triangles is a convex *n*-gon divided? Each triangle shares two vertices with an adjacent one.



We can count internal angles using triangulations. Into how many triangles is a convex *n*-gon divided? Each triangle shares two vertices with an adjacent one. We choose the first triangle – it covers 3 vertices.



We can count internal angles using triangulations. Into how many triangles is a convex *n*-gon divided? Each triangle shares two vertices with an adjacent one. We choose the first triangle – it covers 3 vertices. After that, each triangle covers only one more vertex.



We can count internal angles using triangulations. Into how many triangles is a convex n-gon divided? Each triangle shares two vertices with an adjacent one. We choose the first triangle – it covers 3 vertices. After that, each triangle covers only one more vertex. This means, that an n-gon is divided into n-2 triangles.



We can count internal angles using triangulations. Into how many triangles is a convex n-gon divided? Each triangle shares two vertices with an adjacent one. We choose the first triangle – it covers 3 vertices. After that, each triangle covers only one more vertex. This means, that an n-gon is divided into n-2 triangles.









Construction of a triangulation of a hexagon.



A convex n-gon is divided into n-2 triangles.



A convex n-gon is divided into n-2 triangles.

The sum of all internal angles in a triangle is 180°.



A convex n-gon is divided into n-2 triangles.

The sum of all internal angles in a triangle is 180° .

SUM OF INTERNAL ANGLES IN A CONVEX POLYGON

The sum of all internal angles of a convex n-gon is $(n-2) \cdot 180^{\circ}$.

REGULAR POLYGONS

DEFINITION



REGULAR POLYGON

A regular polygon is a convex polygon whose sides all have the same length and whose internal angles all have the same size.

DEFINITION



REGULAR POLYGON

A regular polygon is a convex polygon whose sides all have the same length and whose internal angles all have the same size.



Equilateral triangle (regular trigon)



Square (regular tetragon)



Regular pentagon



Regular hexagon




ROTATION

Rotation of a polygon consists of well ... rotating each of its points by a fixed angle around a fixed point (called *anchor*).



ROTATION

Rotation of a polygon consists of well ... rotating each of its points by a fixed angle around a fixed point (called *anchor*).






REFLECTION

Reflection of a polygon consists of 'mirroring' each of its points through a given line (called *axis of reflection*).



REFLECTION

Reflection of a polygon consists of 'mirroring' each of its points through a given line (called axis of reflection).






POINT SYMMETRY

Point symmetry of a polygon consists of 'mirroring' each of its points through a given point (called *center of symmetry*).



POINT SYMMETRY

Point symmetry of a polygon consists of 'mirroring' each of its points through a given point (called *center of symmetry*).



Examples of point symmetries.





- rotational symmetries
 - \circ rotation by $\frac{360^{\circ}}{n}$



- rotational symmetries
 - \circ rotation by $\frac{360^{\circ}}{n}$
- reflection (line) symmetries



- rotational symmetries
 - \circ rotation by $\frac{360^{\circ}}{n}$
- reflection (line) symmetries
 - o for *n* even reflections over lines passing through centres of opposite sides
 - o for *n* even over lines passing through opposite vertices
 - o for *n* odd over lines passing through the centre of a side and an opposite vertex




- rotational symmetries
 - \circ rotation by $\frac{360^{\circ}}{n}$
- reflection (line) symmetries
 - o for *n* even reflections over lines passing through centres of opposite sides
 - o for *n* even over lines passing through opposite vertices
 - o for *n* odd over lines passing through the centre of a side and an opposite vertex
- point symmetries




- rotational symmetries
 - \circ rotation by $\frac{360^{\circ}}{n}$
- reflection (line) symmetries
 - o for *n* even reflections over lines passing through centres of opposite sides
 - o for *n* even over lines passing through opposite vertices
 - o for *n* odd over lines passing through the centre of a side and an opposite vertex
- point symmetries
 - o only through the 'centre' the point where its axes of symmetry intersect in case *n* is even














Examples of regular polygon symmetries

CRYPTOGRAPHY ON REGULAR POLYGONS

CHAINING SYMMETRIES



Given two symmetries, s_1 and s_2 of a regular polygon, one can apply them one after the other ('compose' them, like functions).


CHAINING SYMMETRIES




Given two symmetries, s_1 and s_2 of a regular polygon, one can apply them one after the other ('compose' them, like functions).

We'll denote this composition simply by s_1s_2 .





















Discounting point symmetry, an *n*-gon has 2*n* symmetries.



Discounting point symmetry, an n-gon has 2n symmetries. Two symmetries can 'combine' to create a different symmetry.





Discounting point symmetry, an *n*-gon has 2*n* symmetries.

Two symmetries can 'combine' to create a different symmetry.

Natural question: How many (and which) symmetries of a regular polygon do I need to get all the others?





Discounting point symmetry, an *n*-gon has 2*n* symmetries.

Two symmetries can 'combine' to create a different symmetry.

Natural question: How many (and which) symmetries of a regular polygon do I need to get all the others?

For example,

• if s_1 is any line symmetry and s_2 is a rotation by 60° counter-clockwise, then $s_2^3 s_1$ (s_2^3 means $s_2 s_2 s_2$) reflects a hexagon through a line perpendicular to the line of s_1 .



Discounting point symmetry, an *n*-gon has 2*n* symmetries.

Two symmetries can 'combine' to create a different symmetry.

Natural question: How many (and which) symmetries of a regular polygon do I need to get all the others?

For example,

- if s_1 is any line symmetry and s_2 is a rotation by 60° counter-clockwise, then $s_2^3 s_1$ (s_2^3 means $s_2 s_2 s_2$) reflects a hexagon through a line perpendicular to the line of s_1 .
- if s_1 is a rotation by 120° clockwise and s_2 is a reflection through a vertical line passing through the top vertex, then s_1s_2 is a reflection through the line given by the rotation of the line of s_2 60° clockwise.



Actually, for a general *n*-gon, we need only two:



Actually, for a general *n*-gon, we need only two:

• rotation by $360^{\circ}/n$ in any direction (we'll denote it r),





Actually, for a general *n*-gon, we need only two:

- rotation by $360^{\circ}/n$ in any direction (we'll denote it r),
- any reflection (we'll denote it s).

CHAINING SYMMETRIES - TRIANGLE



Let r be the rotation by 120° and s any line symmetry.

CHAINING SYMMETRIES - TRIANGLE



Let r be the rotation by 120° and s any line symmetry.

• The other two rotational symmetries are r^2 and r^3 .

CHAINING SYMMETRIES - TRIANGLE



Let r be the rotation by 120° and s any line symmetry.

- The other two rotational symmetries are r^2 and r^3 .
- The other two line symmetries are rs and r^2s .

CHAINING SYMMETRIES - TRIANGLE



Let r be the rotation by 120° and s any line symmetry.

- The other two rotational symmetries are r^2 and r^3 .
- The other two line symmetries are rs and r^2s .
- Therefore, all the symmetries of an equilateral triangle are

$$\{r, r^2, r^3, s, rs, r^2s\}.$$



In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^{\circ}/n$ where k doesn't share a prime factor with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.







In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^{\circ}/n$ where k doesn't share a prime factor with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.

Why?

• If k shares factors with n, then you can never get rotation by $360^{\circ}/n$.





In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^{\circ}/n$ where k doesn't share a prime factor with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.

Why?

- If k shares factors with n, then you can never get rotation by $360^{\circ}/n$.
- Two symmetries can never produce a rotation.





In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^{\circ}/n$ where k doesn't share a prime factor with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.

Why?

- If k shares factors with n, then you can never get rotation by $360^{\circ}/n$.
- Two symmetries can never produce a rotation.
- Two rotations can never produce a symmetry.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.





You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a rotation, then

1. First measure the angle counter-clockwise.





You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a rotation, then

- 1. First measure the angle **counter-clockwise**.
- 2. Find a such that r^a is the rotation by $360^{\circ}/n$.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a rotation, then

- 1. First measure the angle counter-clockwise.
- 2. Find a such that r^a is the rotation by $360^{\circ}/n$.
- 3. Then, find *b* such that $(r^a)^b = r^{ab}$ is your desired rotation.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a reflection, then

1. Find a such that r^a is the rotation by $360^{\circ}/n$.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a reflection, then

- 1. Find a such that r^a is the rotation by $360^{\circ}/n$.
- 2. Determine the angle **in any direction** between your given line of symmetry **s** and the reflection you want.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a reflection, then

- 1. Find a such that r^a is the rotation by $360^{\circ}/n$.
- 2. Determine the angle **in any direction** between your given line of symmetry **s** and the reflection you want.
- 3. Find b such that r^{ab} is a rotation in the opposite direction by twice the angle from the previous step.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a reflection, then

- 1. Find a such that r^a is the rotation by $360^{\circ}/n$.
- 2. Determine the angle **in any direction** between your given line of symmetry **s** and the reflection you want.
- 3. Find b such that r^{ab} is a rotation in the opposite direction by twice the angle from the previous step.
- 4. r^{ab}s is your desired reflection.



You're given a rotation r by $k \cdot 360^{\circ}/n$ such that k doesn't share factors with n and a line symmetry s.

If you need to calculate a reflection, then


- 1. Find a such that r^a is the rotation by $360^{\circ}/n$.
- 2. Determine the angle **in any direction** between your given line of symmetry **s** and the reflection you want.
- 3. Find b such that r^{ab} is a rotation in the opposite direction by twice the angle from the previous step.
- 4. rabs is your desired reflection.

Voluntary HW: Why does this algorithm work?



We're given two symmetries of the square:





and want to produce



reflection over the second diagonal



We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?



We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?

- We use the algorithm.
 - 1. Repeating r three times gives the rotation by 90° counter-clockwise, that is, a = 3.



We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?

We use the algorithm.

- 1. Repeating r three times gives the rotation by 90° counter-clockwise, that is, a = 3.
- 2. The angle between the two diagonals is 90° in any direction.



We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?

We use the algorithm.

- 1. Repeating r three times gives the rotation by 90° counter-clockwise, that is, a = 3.
- 2. The angle between the two diagonals is 90° in any direction.
- 3. Repeating the rotation from step 1 two times (that is, b=2) and then using s gives the desired symmetry in this case it's $(r^3)^2s=r^6s$. Of course, r^4 is rotation by 360° which does nothing, so the final symmetry is r^2s .