

logo.png

title.jpg

POLYGONS

Adam Klepáč

September 18, 2023

CONTENTS



Cryptography on Regular Polygons


CRYPTOGRAPHY ON REGULAR POLYGONS

The background of the slide is white. It features three large, solid-colored triangles that meet at a central point near the bottom. A yellow triangle is on the left, a cyan triangle is on the right, and a green triangle is at the bottom center, pointing upwards. The title text is centered in the upper half of the slide.

CHAINING SYMMETRIES

Given two symmetries, s_1 and s_2 of a regular polygon, one can apply them one after the other ('compose' them, like functions).


CHAINING SYMMETRIES



Given two symmetries, s_1 and s_2 of a regular polygon, one can apply them one after the other ('compose' them, like functions).


We'll denote this composition simply by s_1s_2 .

CHAINING SYMMETRIES – EXAMPLE




Example of chains of symmetries.

CHAINING SYMMETRIES – EXAMPLE




Example of chains of symmetries.

CHAINING SYMMETRIES – EXAMPLE



Example of chains of symmetries.

CHAINING SYMMETRIES – EXAMPLE



Example of chains of symmetries.

CHAINING SYMMETRIES – HOW MANY DO WE NEED?


Discounting point symmetry, an n -gon has $2n$ symmetries.



CHAINING SYMMETRIES – HOW MANY DO WE NEED?

Discounting point symmetry, an n -gon has $2n$ symmetries.

Two symmetries can 'combine' to create a different symmetry.




CHAINING SYMMETRIES – HOW MANY DO WE NEED?

Discounting point symmetry, an n -gon has $2n$ symmetries.

Two symmetries can 'combine' to create a different symmetry.

Natural question: How many (and which) symmetries of a regular polygon do I need to get all the others?



CHAINING SYMMETRIES – HOW MANY DO WE NEED?

Discounting point symmetry, an n -gon has $2n$ symmetries.

Two symmetries can 'combine' to create a different symmetry.

Natural question: How many (and which) symmetries of a regular polygon do I need to get all the others?

For example,

- if s_1 is any line symmetry and s_2 is a rotation by 60° counter-clockwise, then $s_2^3 s_1$ (s_2^3 means $s_2 s_2 s_2$) reflects a hexagon through a line perpendicular to the line of s_1 .

CHAINING SYMMETRIES – HOW MANY DO WE NEED?

Discounting point symmetry, an n -gon has $2n$ symmetries.

Two symmetries can 'combine' to create a different symmetry.


Natural question: How many (and which) symmetries of a regular polygon do I need to get all the others?

For example,


- if s_1 is any line symmetry and s_2 is a rotation by 60° counter-clockwise, then $s_2^3 s_1$ (s_2^3 means $s_2 s_2 s_2$) reflects a hexagon through a line perpendicular to the line of s_1 .
- if s_1 is a rotation by 120° clockwise and s_2 is a reflection through a vertical line passing through the top vertex, then $s_1 s_2$ is a reflection through the line given by the rotation of the line of s_2 60° clockwise.

CHAINING SYMMETRIES – HOW MANY DO WE NEED?

Actually, for a general n -gon, we need only **two**:



CHAINING SYMMETRIES – HOW MANY DO WE NEED?

- 
- Actually, for a general n -gon, we need only **two**:
 - rotation by $360^\circ/n$ in any direction (we'll denote it r),


CHAINING SYMMETRIES – HOW MANY DO WE NEED?

Actually, for a general n -gon, we need only **two**:

- rotation by $360^\circ/n$ in any direction (we'll denote it r),
- any reflection (we'll denote it s).

CHAINING SYMMETRIES – TRIANGLE


Let r be the rotation by 120° and s any line symmetry.



CHAINING SYMMETRIES – TRIANGLE

Let r be the rotation by 120° and s any line symmetry.


- The other two rotational symmetries are r^2 and r^3 .



CHAINING SYMMETRIES – TRIANGLE

Let r be the rotation by 120° and s any line symmetry.

- The other two rotational symmetries are r^2 and r^3 .
- The other two line symmetries are rs and r^2s .



CHAINING SYMMETRIES – TRIANGLE


Let r be the rotation by 120° and s any line symmetry.

- The other two rotational symmetries are r^2 and r^3 .
- The other two line symmetries are rs and r^2s .
- Therefore, all the symmetries of an equilateral triangle are

$$\{r, r^2, r^3, s, rs, r^2s\}.$$

CHAINING SYMMETRIES – GENERAL ALGORITHM

In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^\circ / n$ where k **doesn't share a prime factor** with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.



CHAINING SYMMETRIES – GENERAL ALGORITHM

In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^\circ / n$ where k **doesn't share a prime factor** with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.

Why?

- If k shares factors with n , then you can never get rotation by $360^\circ / n$.

CHAINING SYMMETRIES – GENERAL ALGORITHM

In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^\circ / n$ where k **doesn't share a prime factor** with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.

Why?

- If k shares factors with n , then you can never get rotation by $360^\circ / n$.
- Two symmetries can never produce a rotation.

CHAINING SYMMETRIES – GENERAL ALGORITHM


In general, to create all symmetries, one needs a rotation by an angle $k \cdot 360^\circ / n$ where k **doesn't share a prime factor** with n (in other words, the fraction $\frac{k}{n}$ cannot be simplified) and any one line symmetry.

Why?

- If k shares factors with n , then you can never get rotation by $360^\circ / n$.
- Two symmetries can never produce a rotation.
- Two rotations can never produce a symmetry.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ / n$ such that k doesn't share factors with n and a line symmetry s .



CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ / n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a rotation, then

1. First measure the angle **counter-clockwise**.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ / n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a rotation, then

1. First measure the angle **counter-clockwise**.
2. Find a such that r^a is the rotation by $360^\circ / n$.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ / n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a rotation, then

1. First measure the angle **counter-clockwise**.
2. Find a such that r^a is the rotation by $360^\circ / n$.
3. Then, find b such that $(r^a)^b = r^{ab}$ is your desired rotation.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ / n$ such that k doesn't share factors with n and a line symmetry s .



CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ/n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a reflection, then

1. Find a such that r^a is the rotation by $360^\circ/n$.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ/n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a reflection, then

1. Find a such that r^a is the rotation by $360^\circ/n$.
2. Determine the angle **in any direction** between your given line of symmetry s and the reflection you want.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ/n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a reflection, then

1. Find a such that r^a is the rotation by $360^\circ/n$.
2. Determine the angle **in any direction** between your given line of symmetry s and the reflection you want.
3. Find b such that r^{ab} is a rotation **in the opposite direction** by **twice** the angle from the previous step.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ/n$ such that k doesn't share factors with n and a line symmetry s .

If you need to calculate a reflection, then

1. Find a such that r^a is the rotation by $360^\circ/n$.
2. Determine the angle **in any direction** between your given line of symmetry s and the reflection you want.
3. Find b such that r^{ab} is a rotation **in the opposite direction** by **twice** the angle from the previous step.
4. $r^{ab}s$ is your desired reflection.

CHAINING SYMMETRIES – GENERAL ALGORITHM

You're given a rotation r by $k \cdot 360^\circ/n$ such that k doesn't share factors with n and a line symmetry s .


If you need to calculate a reflection, then

1. Find a such that r^a is the rotation by $360^\circ/n$.
2. Determine the angle **in any direction** between your given line of symmetry s and the reflection you want.
3. Find b such that r^{ab} is a rotation **in the opposite direction** by **twice** the angle from the previous step.
4. $r^{ab}s$ is your desired reflection.


Voluntary HW: *Why does this algorithm work?*

CHAINING SYMMETRIES – ALGORITHM EXAMPLE

We're given two symmetries of the square:




and want to produce



CHAINING SYMMETRIES – ALGORITHM EXAMPLE

We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?



CHAINING SYMMETRIES – ALGORITHM EXAMPLE

We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?

We use the algorithm.

1. Repeating r three times gives the rotation by 90° counter-clockwise, that is, $a = 3$.

CHAINING SYMMETRIES – ALGORITHM EXAMPLE

We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?

We use the algorithm.

1. Repeating r three times gives the rotation by 90° counter-clockwise, that is, $a = 3$.
2. The angle between the two diagonals is 90° in any direction.

CHAINING SYMMETRIES – ALGORITHM EXAMPLE

We're given two symmetries of the square: rotation r by 270° counter-clockwise and reflection s over the first diagonal.

How to produce reflection over the other diagonal?

We use the algorithm.

1. Repeating r three times gives the rotation by 90° counter-clockwise, that is, $a = 3$.
2. The angle between the two diagonals is 90° in any direction.
3. Repeating the rotation from step 1 two times (that is, $b = 2$) and then using s gives the desired symmetry – in this case it's $(r^3)^2s = r^6s$. Of course, r^4 is rotation by 360° which does nothing, so the final symmetry is r^2s .