



SYSTEMS OF LINEAR EQUATIONS

Adam Klepáč

March 25, 2024

CONTENTS



FUNCTIONS

The bottom of the slide features two large, overlapping triangular shapes. The triangle on the left is a light blue color, and the triangle on the right is a darker blue color. They meet at a point in the center, creating a V-shape that points downwards.


WHAT IS A FUNCTION?

Intuitively, a function is a **box** which receives data and gives some data back.



WHAT IS A FUNCTION?

Intuitively, a function is a **box** which receives data and gives some data back.



WHAT IS A FUNCTION?

Intuitively, a function is a **box** which receives data and gives some data back.



We'll call the data that **a function receives**, **inputs** and the **data it gives back**, **outputs**.

WHAT IS A FUNCTION?

Intuitively, a function is a **box** which receives data and gives some data back.



We'll call the data that **a function receives**, **inputs** and the **data it gives back**, **outputs**.

Inputs and **outputs** need not necessarily be just 'one object', they can be for example lists of numbers.


FUNCTIONS – EXAMPLE

A function which returns the **average** of a given set of numbers receives the numbers and also their count as **input** and returns the **average** as **output**.

FUNCTIONS – EXAMPLE

A function which returns the **average** of a given set of numbers receives the numbers and also their count as **input** and returns the **average** as **output**.


x_1, x_2, \dots, x_n
 n



$$\frac{x_1 + x_2 + \dots + x_n}{n}$$

FUNCTIONS – EXAMPLE

We can also consider ‘non-mathematical’ functions. Like a function which receives a type of meal and returns the ingredients.



FUNCTIONS – EXAMPLE

We can also consider ‘non-mathematical’ functions. Like a function which receives a type of meal and returns the ingredients.




1

FUNCTION COMPOSITION



FUNCTION COMPOSITION

If we have **two functions**, we can **in certain cases** 'compose' them.



FUNCTION COMPOSITION

If we have **two functions**, we can **in certain cases** 'compose' them.

Composition simply means that **one function follows the other** – in other words, the **output** of the first function is the **input** of the second.

FUNCTION COMPOSITION

If we have **two functions**, we can **in certain cases** 'compose' them.

Composition simply means that **one function follows the other** – in other words, the **output** of the first function is the **input** of the second.

Of course, **composition** is only possible if the **output** of the first function is a valid **input** for the second.

FUNCTION COMPOSITION

If we have **two functions**, we can **in certain cases** 'compose' them.


Composition simply means that **one function follows the other** – in other words, the **output** of the first function is the **input** of the second.

Of course, **composition** is only possible if the **output** of the first function is a valid **input** for the second.

For instance, you could hardly compose the **ingredients** function with the **average** function.


FUNCTION COMPOSITION

Considering two functions



FUNCTION COMPOSITION

Considering two functions




their composition can look like this



FUNCTION COMPOSITION


Considering two functions



their composition can look like this




What would the output of this composition look like



FUNCTION COMPOSITION

So, the **order of the composition matters!** Here are a few examples of compositions which make sense:



FUNCTION COMPOSITION


So, the **order of the composition matters!** Here are a few examples of compositions which make sense:

omelette \longrightarrow ingredients \longrightarrow total price \longrightarrow \$4.37

x_1, x_2, \dots, x_n
 n \longrightarrow average \longrightarrow times two \longrightarrow $2 \cdot \frac{x_1 + x_2 + \dots + x_n}{n}$

FUNCTION COMPOSITION

So, the **order of the composition matters!** Here are a few examples of compositions which make sense:



We can of course compose **as many functions** as we like. An example of this:



FUNCTIONS – NOTATION

Drawing pictures like this would be cumbersome.



FUNCTIONS – NOTATION

Drawing pictures like this would be cumbersome. Instead of



we simply write `function(input) = output`.

FUNCTIONS – NOTATION

Drawing pictures like this would be cumbersome. Instead of




we simply write **function(input) = output**.

You're probably used to seeing function written like $f(x) = y$. The picture corresponding to this is



FUNCTIONS – NOTATION


The symbol used for function composition is \circ . It is however a little confusing because it is written in order ‘from right to left’.



FUNCTIONS – NOTATION

The symbol used for function composition is \circ . It is however a little confusing because it is written in order ‘from right to left’.

For example, if f and g are two functions, their composition $f \circ g$ corresponds to this picture



that is, **first g , then f** .

2

REAL FUNCTIONS



REAL FUNCTION

A **real function** is simply a function whose **input** and **output** are both real numbers.



REAL FUNCTION

A **real function** is simply a function whose **input** and **output** are both real numbers.

Examples of such functions are

- $f(x) = 0$,
- $g(x) = \tan^6(\log^{\sin(x^2+4)}(\frac{5x^3-2}{9x^7}))$,

where $x \in \mathbb{R}$.


REAL FUNCTION

A **real function** is simply a function whose **input** and **output** are both real numbers.

Examples of such functions are


- $f(x) = 0$,
- $g(x) = \tan^6(\log^{\sin(x^2+4)}(\frac{5x^3-2}{9x^7}))$,

where $x \in \mathbb{R}$. Or, using pictures,



OPERATIONS ON REAL FUNCTIONS

As both the **input** and the **output** of a real function is a real number, **we can always compose real functions.**



OPERATIONS ON REAL FUNCTIONS

As both the **input** and the **output** of a real function is a real number, **we can always compose real functions.**

However, that **doesn't mean that the order doesn't matter!** Different order of composition gives different functions.

OPERATIONS ON REAL FUNCTIONS

As both the **input** and the **output** of a real function is a real number, **we can always compose real functions.**

However, that **doesn't mean that the order doesn't matter!** Different order of composition gives different functions.

For example, take $f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$. Then,

OPERATIONS ON REAL FUNCTIONS

As both the **input** and the **output** of a real function is a real number, **we can always compose real functions.**

However, that **doesn't mean that the order doesn't matter!** Different order of composition gives different functions.

For example, take $f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$. Then,

- $(f \circ g)(x) = 2 \left(\frac{1}{1+x} \right)^2 + 7$ and

OPERATIONS ON REAL FUNCTIONS

As both the **input** and the **output** of a real function is a real number, **we can always compose real functions.**

However, that **doesn't mean that the order doesn't matter!** Different order of composition gives different functions.

For example, take $f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$. Then,

- $(f \circ g)(x) = 2 \left(\frac{1}{1+x} \right)^2 + 7$ and
- $(g \circ f)(x) = \frac{1}{1+(2x^2+7)}.$

OPERATIONS ON REAL FUNCTIONS

Real functions can also be **added** and **multiplied**, just like real numbers.



OPERATIONS ON REAL FUNCTIONS

Real functions can also be **added** and **multiplied**, just like real numbers.
This involves simply adding or multiplying their respective outputs.



OPERATIONS ON REAL FUNCTIONS

Real functions can also be **added** and **multiplied**, just like real numbers.

This involves simply adding or multiplying their respective outputs. For the functions

$f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$, their

- **sum** is the function with output

$$(f + g)(x) = f(x) + g(x) = 2x^2 + 7 + \frac{1}{1+x}.$$

OPERATIONS ON REAL FUNCTIONS

Real functions can also be **added** and **multiplied**, just like real numbers.

This involves simply adding or multiplying their respective outputs. For the functions

$f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$, their

- **sum** is the function with output

$$(f + g)(x) = f(x) + g(x) = 2x^2 + 7 + \frac{1}{1+x}.$$

- **product** is the function with output

$$(f \cdot g)(x) = f(x) \cdot g(x) = (2x^2 + 7) \cdot \left(\frac{1}{1+x} \right) = \frac{2x^2 + 7}{1+x}.$$

GRAPHS



As real functions have real numbers as inputs and outputs, they can be easily **graphed**.



GRAPHS

As real functions have real numbers as inputs and outputs, they can be easily **graphed**. **Graphing** a real function f simply means drawing the points $(x, f(x))$ or (*input*, *output*) in some chosen coordinate system.

GRAPHS

As real functions have real numbers as inputs and outputs, they can be easily **graphed**.

Graphing a real function f simply means drawing the points $(x, f(x))$ or (*input*, *output*) in some chosen coordinate system.

We typically use the **Cartesian** coordinate system with two axes (one for *input* and one for *output*) that are mutually perpendicular.

GRAPHS

As real functions have real numbers as inputs and outputs, they can be easily **graphed**.

Graphing a real function f simply means drawing the points $(x, f(x))$ or (*input*, *output*) in some chosen coordinate system.

We typically use the **Cartesian** coordinate system with two axes (one for *input* and one for *output*) that are mutually perpendicular. These are often called the *x*-axis and the *y*-axis.

GRAPHS

As real functions have real numbers as inputs and outputs, they can be easily **graphed**.



Graphing a real function f simply means drawing the points $(x, f(x))$ or (*input*, *output*) in some chosen coordinate system.

We typically use the **Cartesian** coordinate system with two axes (one for *input* and one for *output*) that are mutually perpendicular. These are often called the *x*-axis and the *y*-axis.

However, later, we'll also use the **polar** coordinate system where every point is instead determined by its angle and distance from the origin of the system.



GRAPHS

The functions $f(x) = 2x^2 + 7$ and $g(x) = \frac{1}{1+x}$ have the following (parts of) graphs:



GRAPHS

Just to better drive home the idea that the order of function composition is important, look at the graphs of $f \circ g$ and $g \circ f$.



3

LINEAR FUNCTIONS



LINEAR FUNCTION

Linear functions are a special type of real functions **whose output** is always **of the form** $a \cdot (\textit{input}) + b$ for fixed numbers $a, b \in \mathbb{R}$.



LINEAR FUNCTION

Linear functions are a special type of real functions **whose output** is always **of the form** $a \cdot (\text{input}) + b$ for fixed numbers $a, b \in \mathbb{R}$.

In other words, linear functions are those real functions **that only scale the input and add something to it.**

LINEAR FUNCTION

Linear functions are a special type of real functions **whose output** is always **of the form** $a \cdot (\text{input}) + b$ for fixed numbers $a, b \in \mathbb{R}$.

In other words, linear functions are those real functions **that only scale the input and add something to it**.

LINEAR FUNCTION

A real function f is **linear** if

$$f(x) = ax + b$$

for some $a, b \in \mathbb{R}$.

LINEAR FUNCTIONS – PROPERTIES

Linear functions have some unique properties:

- If f and g are linear, so are $f \circ g$ and $g \circ f$.

LINEAR FUNCTIONS – PROPERTIES

Linear functions have some unique properties:

- If f and g are linear, so are $f \circ g$ and $g \circ f$. Indeed, we can see this easily. Suppose $f(x) = ax + b$ and $g(x) = cx + d$, then

$$(f \circ g)(x) = a(cx + d) + b = (ac)x + (ad + b),$$

$$(g \circ f)(x) = c(ax + b) + d = (ac)x + (cb + d).$$

LINEAR FUNCTIONS – PROPERTIES

Linear functions have some unique properties:

- If f and g are linear, so are $f \circ g$ and $g \circ f$. Indeed, we can see this easily. Suppose $f(x) = ax + b$ and $g(x) = cx + d$, then

$$(f \circ g)(x) = a(cx + d) + b = (ac)x + (ad + b),$$

$$(g \circ f)(x) = c(ax + b) + d = (ac)x + (cb + d).$$

- If f and g are linear, so is $f + g$.

LINEAR FUNCTIONS – PROPERTIES

Linear functions have some unique properties:

- If f and g are linear, so are $f \circ g$ and $g \circ f$. Indeed, we can see this easily. Suppose $f(x) = ax + b$ and $g(x) = cx + d$, then

$$(f \circ g)(x) = a(cx + d) + b = (ac)x + (ad + b),$$

$$(g \circ f)(x) = c(ax + b) + d = (ac)x + (cb + d).$$

- If f and g are linear, so is $f + g$. If we just compute the sum, we get

$$(f + g)(x) = (ax + b) + (cx + d) = (a + c)x + (b + d).$$

LINEAR FUNCTIONS – GRAPHS

Graphs of linear functions **are straight lines**.



LINEAR FUNCTIONS – GRAPHS

Graphs of linear functions **are straight lines**.

Let us see why this is true. Choose a linear function

$$f(x) = ax + b.$$



LINEAR FUNCTIONS – GRAPHS

Graphs of linear functions **are straight lines**.

Let us see why this is true. Choose a linear function

$$f(x) = ax + b.$$

Pick a number – x . One way to show that the graph of f is a line is to move by two different distances from x and see that we get two similar triangles.

LINEAR FUNCTIONS – GRAPHS

Graphs of linear functions **are straight lines**.

Let us see why this is true. Choose a linear function

$$f(x) = ax + b.$$

Pick a number – x . One way to show that the graph of f is a line is to move by two different distances from x and see that we get two similar triangles.

If we move by 1, from x to $x + 1$, then on the y -axis we move from $ax + b$ to $a(x + 1) + b$, that is, we move by

$$a(x + 1) + b - (ax + b) = a.$$

LINEAR FUNCTIONS – GRAPHS

Graphs of linear functions **are straight lines**.

Let us see why this is true. Choose a linear function

$$f(x) = ax + b.$$


Pick a number – x . One way to show that the graph of f is a line is to move by two different distances from x and see that we get two similar triangles.

If we move by 1, from x to $x + 1$, then on the y -axis we move from $ax + b$ to $a(x + 1) + b$, that is, we move by

$$a(x + 1) + b - (ax + b) = a.$$

If we move by 2, from x to $x + 2$, on the y -axis, we move by

$$a(x + 2) + b - (ax + b) = 2a.$$



LINEAR FUNCTIONS – INTERSECTIONS

As two non-parallel lines **always intersect**, we should expect the equation $f(x) = g(x)$ to always **have a solution** assuming the graphs of f and g are not parallel.

LINEAR FUNCTIONS – INTERSECTIONS

As two non-parallel lines **always intersect**, we should expect the equation $f(x) = g(x)$ to always **have a solution** assuming the graphs of f and g are not parallel.

The question is, when are the graphs of f and g parallel?

LINEAR FUNCTIONS – INTERSECTIONS

As two non-parallel lines **always intersect**, we should expect the equation $f(x) = g(x)$ to always **have a solution** assuming the graphs of f and g are not parallel.

The question is, when are the graphs of f and g parallel?

That happens exactly when their **rates growth** are identical.

LINEAR FUNCTIONS – INTERSECTIONS



As two non-parallel lines **always intersect**, we should expect the equation $f(x) = g(x)$ to always **have a solution** assuming the graphs of f and g are not parallel.

The question is, when are the graphs of f and g parallel?

That happens exactly when their **rates growth** are identical.

In symbols, if $f(x) = ax + b$ and $g(x) = cx + d$, then the graphs of f and g are parallel if $a = c$.

LINEAR FUNCTIONS – INTERSECTIONS



LINEAR EQUATIONS

The bottom of the slide features two large, overlapping triangular shapes. The shape on the left is a light blue triangle pointing downwards. The shape on the right is a darker blue triangle pointing upwards. They overlap in the center, creating a darker blue triangular region at the bottom.

LINEAR EQUATION

LINEAR EQUATION

If f, g are linear functions, the equation

$$f(x) = g(x)$$

is called a **linear equation** (in one variable).

LINEAR EQUATION – SOLUTION

Suppose $f(x) = ax + b$ and $g(x) = cx + d$. What are the possible solutions to $f(x) = g(x)$?



LINEAR EQUATION – SOLUTION

Suppose $f(x) = ax + b$ and $g(x) = cx + d$. What are the possible solutions to $f(x) = g(x)$?

Three things can happen:

- $a \neq c$. In this case, we can divide the equation by $a - c$ and get

$$ax + b = cx + d$$

$$(a - c)x = d - b$$

$$x = \frac{d - b}{a - c}.$$

LINEAR EQUATION – SOLUTION

Suppose $f(x) = ax + b$ and $g(x) = cx + d$. What are the possible solutions to $f(x) = g(x)$?

Three things can happen:

- $a \neq c$. In this case, we can divide the equation by $a - c$ and get

$$ax + b = cx + d$$

$$(a - c)x = d - b$$

$$x = \frac{d - b}{a - c}.$$

- $a = c$ and $b \neq d$. In this case the graphs of the two functions are parallel lines – there is no solution.

LINEAR EQUATION – SOLUTION

Suppose $f(x) = ax + b$ and $g(x) = cx + d$. What are the possible solutions to $f(x) = g(x)$?

Three things can happen:

- $a \neq c$. In this case, we can divide the equation by $a - c$ and get

$$ax + b = cx + d$$

$$(a - c)x = d - b$$

$$x = \frac{d - b}{a - c}.$$

- $a = c$ and $b \neq d$. In this case the graphs of the two functions are parallel lines – there is no solution.
- $a = c$ and $b = d$. In this case, the functions are one and the same and every number is a solution.

LINEAR EQUATIONS & OPERATIONS

For two linear functions f, g , when does

$$(f \circ g)(x) = (g \circ f)(x)$$

have a solution?

LINEAR EQUATIONS & OPERATIONS

For two linear functions f , g , when does

$$(f \circ g)(x) = (g \circ f)(x)$$

have a solution?

We can calculate that easily. If $f(x) = ax + b$ and $g(x) = cx + d$, then

$$(f \circ g)(x) = a(cx + d) + b = (ac)x + (ad + b),$$

$$(g \circ f)(x) = c(ax + b) + d = (ac)x + (bc + d).$$

LINEAR EQUATIONS & OPERATIONS

For two linear functions f, g , when does

$$(f \circ g)(x) = (g \circ f)(x)$$

have a solution?

We can calculate that easily. If $f(x) = ax + b$ and $g(x) = cx + d$, then

$$(f \circ g)(x) = a(cx + d) + b = (ac)x + (ad + b),$$

$$(g \circ f)(x) = c(ax + b) + d = (ac)x + (bc + d).$$

We see that the graphs of $f \circ g$ and $g \circ f$ are parallel, so this equation has a solution only in the case that $ad + b = bc + d$.

1

LINEAR EQUATIONS IN TWO VARIABLES

LINEAR EQUATIONS IN TWO VARIABLES

SYSTEM OF LINEAR EQUATIONS

A pair of equations

$$ax + by = c,$$

$$dx + ey = f,$$

that have to be **simultaneously** satisfied, is called a **system of (two) linear equations** (in two variables).

LINEAR EQUATIONS IN TWO VARIABLES

Linear equations in two variables can be **reduced to** linear equations **in one variable**.



LINEAR EQUATIONS IN TWO VARIABLES

Linear equations in two variables can be **reduced to** linear equations **in one variable**. This is easily done. We can simply **isolate** one of the variables and make the other variable **into a function**.



LINEAR EQUATIONS IN TWO VARIABLES

Linear equations in two variables can be **reduced to** linear equations **in one variable**. This is easily done. We can simply **isolate** one of the variables and make the other variable **into a function**.

For example, imagine the equation

$$4x - y = 2.$$

LINEAR EQUATIONS IN TWO VARIABLES

Linear equations in two variables can be **reduced to** linear equations **in one variable**. This is easily done. We can simply **isolate** one of the variables and make the other variable **into a function**.

For example, imagine the equation

$$4x - y = 2.$$

We can rewrite this as

$$y = 4x - 2,$$

basically making y into a linear function $f(x) = 4x - 2$.

LINEAR EQUATIONS IN TWO VARIABLES

Linear equations in two variables can be **reduced to** linear equations **in one variable**. This is easily done. We can simply **isolate** one of the variables and make the other variable **into a function**.

For example, imagine the equation

$$4x - y = 2.$$

Similarly, we can write

$$x = \frac{1}{4}y + \frac{2}{4}$$

and turn x into a linear function $g(y) = \frac{1}{4}y + \frac{2}{4}$.

LINEAR EQUATIONS IN TWO VARIABLES

You probably know this reduction under the name of **substitution**.



LINEAR EQUATIONS IN TWO VARIABLES

You probably know this reduction under the name of **substitution**.
Let us see it in practice.



LINEAR EQUATIONS IN TWO VARIABLES

You probably know this reduction under the name of **substitution**.

Let us see it in practice.

Suppose we want to solve the system

$$3x + y = 4,$$

$$x - 2y = 6.$$

LINEAR EQUATIONS IN TWO VARIABLES

You probably know this reduction under the name of **substitution**.

Let us see it in practice.

Suppose we want to solve the system

$$3x + y = 4,$$

$$x - 2y = 6.$$

From the first equation, we see that

$$y = 3x + 6,$$

LINEAR EQUATIONS IN TWO VARIABLES

You probably know this reduction under the name of **substitution**.

Let us see it in practice.

Suppose we want to solve the system

$$3x + y = 4,$$

$$x - 2y = 6.$$

From the first equation, we see that

$$y = 3x + 6,$$

and from the second that

$$y = \frac{1}{2}x - 3.$$

LINEAR EQUATIONS IN TWO VARIABLES

From the first equation, we see that

$$y = 3x + 6,$$

and from the second that

$$y = \frac{1}{2}x - 3.$$

Therefore, we get the linear equation **in one variable**

$$3x + 6 = \frac{1}{2}x - 3.$$

LINEAR EQUATIONS IN TWO VARIABLES

From the first equation, we see that

$$y = 3x + 6,$$

and from the second that

$$y = \frac{1}{2}x - 3.$$

Therefore, we get the linear equation **in one variable**

$$3x + 6 = \frac{1}{2}x - 3.$$

whose solution is $x = -\frac{18}{5}$.

LINEAR EQUATIONS IN TWO VARIABLES

From the first equation, we see that

$$y = 3x + 6,$$

and from the second that

$$y = \frac{1}{2}x - 3.$$

The functions $f(x) = 3x + 6$ and $g(x) = \frac{1}{2}x - 3$ are linear, so we can draw this equation as an intersection of two lines.

LINEAR EQUATIONS IN TWO VARIABLES

