

Problem Collection for Introduction to Mathematical Reasoning

By Dana C. Ernst and Nándor Sieben
Northern Arizona University

Problem 1. Three strangers meet at a taxi stand and decide to share a cab to cut down the cost. Each has a different destination but all are heading in more-or-less the same direction. Bob is traveling 10 miles, Sally is traveling 20 miles, and Mike is traveling 30 miles. If the taxi costs \$2 per mile, how much should each contribute to the total fare? What do you think is the most common answer to this question?

Problem 2. Christine wants to take yoga classes to increase her strength and flexibility. In her neighborhood, there are two yoga studios: Namaste Yoga and Yoga Spirit. At Namaste Yoga, a student's first class costs \$12, and additional classes cost \$10 each. At Yoga Spirit, a student's first class costs \$24, and additional classes cost \$8 each. Because Christine wants to save money, she is interested in comparing the costs of the two studios. For what number of yoga classes do the two studios cost the same amount?

Problem 3. Imagine a hallway with 1000 doors numbered consecutively 1 through 1000. Suppose all of the doors are closed to start with. Then some dude with nothing better to do walks down the hallway and opens all of the doors. Because the dude is still bored, he decides to close every other door starting with door number 2. Then he walks down the hall and changes (i.e., if open, he closes it; if closed, he opens it) every third door starting with door 3. Then he walks down the hall and changes every fourth door starting with door 4. He continues this way, making a total of 1000 passes down the hallway, so that on the 1000th pass, he changes door 1000. At the end of this process, which doors are open and which doors are closed?

Problem 4. The Sunny Day Juice Stand sells freshly squeezed lemonade and orange juice at the farmers' market. The juices are ladled out of large glass jars, each holding exactly the same amount of juice. Linda and Julie set up their stand early one Saturday morning. The first customer of the day ordered orange juice and Linda carefully ladled out 8 ounces into a paper cup. As she was about to hand the cup to the customer, he changed his mind and asked for lemonade instead. Accidentally, Linda dumped the cup of orange juice into the jar of lemonade. She quickly mixed up the juices, ladled out a cup of the mixture (mostly lemonade) and turned to hand it to the customer. "I've decided I don't want anything to drink right now," he said, and frazzled, Linda dumped the cupful of juice mixture into the orange juice jar. Linda's assistant, Julie, watched all of this with amusement. As the man walked away, she wondered aloud, "Now is there more orange juice in the lemonade or more lemonade in the orange juice?"

Problem 5. Imagine you have 25 pebbles, each occupying one square on a 5 by 5 chess board. Tackle each of the following variations of a puzzle.

- (a) Variation 1: Suppose that each pebble must move to an adjacent square by only moving up, down, left, or right. If this is possible, describe a solution. If this is impossible, explain why.
- (b) Variation 2: Suppose that all but one pebble (your choice which one) must move to an adjacent square by only moving up, down, left, or right. If this is possible, describe a solution. If this is impossible, explain why.
- (c) Variation 3: Consider Variation 1 again, but this time also allow diagonal moves to adjacent squares. If this is possible, describe a solution. If this is impossible, explain why.

Problem 6. Consider an $n \times n$ chess board and variation 1 of the pebble puzzle from above. For what values of n is the puzzle solvable? For what values of n is the puzzle unsolvable? Justify your answers by either providing a method for a solution or an explanation for why a solution is not possible.

Problem 7. Consider an $n \times n$ chess board and variation 2 of the pebble puzzle from above. For what values of n is the puzzle solvable? For what values of n is the puzzle unsolvable? Justify your answers by either providing a method for a solution or an explanation for why a solution is not possible.

Problem 8. Describe where on Earth from which you can travel one mile south, then one mile east, and then one mile north and arrive at your original location. There is more than one such location. Find them all.

Problem 9. I have 10 sticks in my bag. The length of each stick is an integer. No matter which 3 sticks I try to use, I cannot make a triangle out of those sticks. What is the minimum length of the longest stick?

Problem 10. Suppose there are two bags of candy containing 8 pieces and 6 pieces, respectively. You and your friend are going to play a game and the winner gets to eat all of the candy. Here are the rules for the game:

1. You and your friend will alternate removing pieces of candy from the bags. Let's assume that you go first.
2. On each turn, the designated player selects a bag that still has candy in it and then removes at least one piece of candy. The designated player can only remove candy from a single bag and he/she must remove at least one piece.
3. The winner is the one that removes all the candy from the last remaining bag.

Does one of you have a guaranteed winning strategy? If so, describe that strategy. Can you generalize to handle any number of pieces of candy in either of the two bags?

Problem 11. Suppose you have 6 toothpicks that are exactly the same length. Can you arrange the toothpicks so that exactly 4 identical triangles are formed? You cannot cut, break, or bend the toothpicks. Moreover, each vertex of a triangle must be formed when the tips of two toothpicks meet.

Problem 12. An ant is crawling along the edges of a unit cube. What is the maximum distance it can cover starting from a corner so that it does not cover any edge twice?

Problem 13. The grid below has 12 boxes and 15 edges connecting boxes. In each box, place one of the six integers from 1 to 6 such that the following conditions hold:

- For each possible pair of distinct numbers from 1 to 6, there is exactly one edge connecting two boxes with that pair of numbers.
- If an edge has an arrow, then it points from a box with a smaller number to a box with a larger number.

You do not need to prove that your configuration is the only one possible; you merely need to find a configuration that satisfies the constraints above.

