

INSERT TITLE

TEVA ILAN

A SENIOR THESIS
PRESENTED TO THE FACULTY
OF PRINCETON UNIVERSITY
IN CANDIDACY FOR THE DEGREE
OF BACHELOR OF THE ARTS

RECOMMENDED FOR ACCEPTANCE
BY THE DEPARTMENT OF
ASTROPHYSICAL SCIENCES

ADVISER: YOUR ADVISOR

STAR WARS DAY

I hereby declare that I am the sole author of this thesis.

I authorize Princeton University to lend this thesis to other institutions or individuals for the purpose of scholarly research.

Teva Ilan

I further authorize Princeton University to reproduce this thesis by photocopying or by other means, in total or in part, at the request of other institutions or individuals for the purpose of scholarly research.

Teva Ilan

Abstract

Your Abstract

Contents

Abstract	iii
1 Introduction	1
1.1 Background	1
2 Data	2
2.1 CMB Maps	2
2.2 Galaxy Cluster Catalogs	2
3 Theory and Methods	4
3.1 The tSZ Effect	4
3.2 Aperture Photometry	4
3.3 Stacking	5
4 Results	8
5 Discussion and Conclusion	11
6 Future Research	12
7 References	13

Chapter 1

Introduction

1.1 Background

The Sunyaev–Zel’dovich (SZ) effect results from inverse Compton scattering of CMB photons off of the hot gas in galaxy clusters. The dominant component of this effect results from interactions between Cosmic Microwave Background (CMB) photons and electrons in the gas that have high energies as a result of their temperature. This produces a frequency dependent shift in the observed CMB temperature along the Line-of-Sight to the galaxy cluster and is known as the thermal SZ (tSZ) effect. The shape of the distortion of the observed CMB due to the tSZ effect is constant to first order but there is a higher order dependency on the temperature of the gas due to relativistic corrections. This is called the relativistic SZ effect (rSZ) (Hincks et al, 2018). There is also a second order effect that results from interactions between the CMB photons and electrons in the gas that have high energies as a result of their bulk motion. This effect depends on the peculiar velocity of the cluster, has no frequency dependence, and is known as the kinetic SZ (kSZ) effect (Hincks et al., 2018).

Thus the tSZ effect can be used as a proxy for the mass or the amount of gas in clusters, the kSZ effect can be used to measure their peculiar velocity, and the rSZ effect can be used to measure the gas temperature. Understanding the tSZ signal and thus understanding the amount of gas in clusters is important because it informs the astrophysics of galaxy cluster formation. The kSZ effect is important as a measurement of bulk kSZ can be used to constrain radial inhomogeneity in the peculiar velocities of galaxy clusters. An alternative explanation for the apparent accelerating expansion of the universe is that our galaxy is located in a giant under-dense void and the observed acceleration is due to the peculiar velocities of galaxy clusters rather than a change in the rate of cosmic expansion (Bull et al., 2011). One of the goals of this work was to put a constraint on bulk kSZ, which is important given the fact that our understanding of cosmology is based on homogeneity on large scales. A detection of bulk kSZ would derail this foundational assumption of cosmology.

Trying to measure rSZ is important because if measurable, it would provide an additional means of measuring the gas temperature in clusters, which is traditionally done via X-ray spectroscopy (Hincks et al, 2018). Though rSZ is a higher order effect it can be on the order of a few percent at certain frequencies and ignoring it can lead to an underestimation of the Compton- y parameter which in turn results in a biased Y – M relation, which is used to estimate cosmological parameters (Hincks et al, 2018). Additionally, apart from a recent claimed detection by Hincks et al., rSZ has never been detected. Thus, constraining or detecting rSZ is itself a topic of interest. In this work, we were able to constrain rSZ but failed to make a detection with the available data.

Chapter 2

Data

2.1 CMB Maps

We used CMB maps at 5 frequencies: 90 GHz, 150 GHz, 217 GHz, 353 GHz, 545 GHz, and 857 GHz. The 90 GHz and 150 GHz maps are from the Atacama Cosmology Telescope (ACT) while the high frequency maps are from the Planck satellite. All of the maps go from -44 degrees to 46 degrees in right ascension and -9 degrees to 5 degrees in declination. Using maps from multiple frequencies allows us to disentangle different components of the observed flux from each other, specifically the dust component, the thermal Sunyaev-Zel'dovich (tSZ) effect component, and the kinetic Sunyaev-Zel'dovich (kSZ).

2.2 Galaxy Cluster Catalogs

Our work involved two sets of galaxy cluster catalogues: the redMaPPer and an ACT confirmed cluster catalog. RedMaPPer is a red-sequence cluster finder, which means that it finds galaxy cluster candidates by looking for over densities of galaxies that form a red sequence.




Figure 2.1: This plot shows a histogram of the distribution of redshifts of the clusters in the RedMaPPer catalog. Evidently, the clusters in the RedMaPPer catalog range from $z \approx 0.05$ to $z \approx 0.6$. Note that this plot only includes clusters from RedMaPPer that were used in our analysis and thus does not include clusters that fell outside the range of our maps or were too close to point sources.




Figure 2.2: This plot shows a histogram of the distribution of the richnesses of the clusters in the RedMaPPer catalog. Evidently, the vast majority of clusters in the RedMaPPer catalog fall between richnesses of 25 and 100. Note that this plot only includes clusters from RedMaPPer that were used in our analysis and thus does not include clusters that fell outside the range of our maps or were too close to point sources.

Chapter 3

Theory and Methods

3.1 The tSZ Effect

The tSZ effect is the result of inverse Compton scattering of photons from the CMB off of high energy electrons in hot gas. When CMB photons go through hot cluster gas they gain energy from the high energy electrons. Then when we measure the CMB these photons produce a frequency-dependent change in the observed CMB temperature along the Line-of-Sight (LOS) to the galaxy cluster (Greco et al.). The tSZ effect is quantified by the Compton-y parameter. The observationally relevant Compton-y parameter is the Comptonization parameter integrated along the full LOS within some aperture radius,

$$Y_c^{cyl} = d_A^{-2}(z) \frac{\sigma_T}{m_e c^2} \int_0^{R_c} 2\pi R dR \int_{-\inf}^{\inf} dl P_e(r, M, z) \quad (3.1)$$

Where Y_c^{cyl} is the observed Compton-y parameter, $d_A(z)$ is the angular diameter distance to the cluster, σ_T is the Thomson scattering cross-section, $m_e c^2$ is the rest mass of the electron, R is the projected radius which we integrate up to R_c , and $P_e(r, M, z)$ is the electron pressure profile (Greco et al.).

3.2 Aperture Photometry

To extract the Compton-y parameter from the CMB maps and differentiate it from effects of the dust, and bulk kSZ, we perform aperture photometry on the CMB maps at the locations of clusters indicated by the cluster catalogs. To perform aperture photometry we define a circular aperture of some radius (which we determined empirically to correspond to the typical radius of the tSZ effect at each frequency) surrounded by an annulus. We then subtract the mean pixel value in the annulus from the circular aperture and define the observed signal, S_i , to be the sum of the pixel values inside the circular aperture after this subtraction. The mean value of the annulus is subtracted to correct for large-scale foreground contamination, assuming it is roughly constant over the extracted aperture (Greco et al.). This signal, S_i , is then modeled just as described in Greco et al.,

$$\begin{aligned} S_i &= a_i Y_c^{cyl} + b_i D_c + \delta T_{CMB} \\ a_i &= g(\nu_i) T_{CMB} \\ b_i &= \left(\frac{\nu_i(1+z)}{\nu_0} \right)^\beta B(\nu_i(1+z), T_{dust}) \left[\frac{\partial B(\nu_i, T)}{\partial T} \right]_{T_{CMB}}^{-1}, \end{aligned} \quad (3.2)$$

where $g(\nu_i) = h\nu_i/k_B T_{CMB} \coth(h\nu_i/2k_B T_{CMB})$, $T_{CMB} = 2.7255K$, D_c is the amplitude of the dust emission integrated along the full LOS within the aperture radius, $\nu_0 = 353$ GHz is the reference frequency, $\beta = 1.78$ is the dust spectral emissivity index, $T_{dust} = 20K$ is the dust temperature, $B(\nu_i, T)$ is the Planck function, and δT_{CMB} provides a constraint on bulk kSZ.

We do a weighted sum of the aperture photometry values for all the clusters at each frequency in which the weight of each value is inversely proportional to the amount of noise in that area of the map. Then we fit these weighted sums to the model above using emcee.

We use a similar methodology to fit for tSZ, dust, and rSZ (rather than bulk kSZ). When fitting for these parameters, we use a slightly different model.

$$\begin{aligned}
 S_i &= a_i Y_c^{cyl} + b_i D_c + Y_c^{cyl} g_{rel}(\nu_i, T_{eff}) \\
 a_i &= g_{rel}(\nu_i, T_{eff}) T_{CMB} \\
 b_i &= \left(\frac{\nu_i(1+z)}{\nu_0} \right)^\beta B(\nu_i(1+z), T_{dust}) \left[\frac{\partial B(\nu_i, T)}{\partial T} \right]_{T_{CMB}}^{-1},
 \end{aligned} \tag{3.3}$$

where $g_{rel}(\nu, T_{eff})$ is $g(\nu)$ modified with relativistic corrections thus introducing a dependency on the effective temperature (T_{eff}). In the case $T_{eff} = 0$, $g_{rel}(\nu, T_{eff} = 0) = g(\nu)$. A plot of $g_{rel}(\nu, T_{eff})$ for different values of T_{eff} and a plot of the dust coefficient are shown below.




Figure 3.1: This plot shows $g(\nu)$ as a function of frequency and for different values of T_{eff} .




Figure 3.2: This plot shows the dust coefficient as a function of frequency at a constant value of $z=0.36315244$, which is the average z value for our sample of galaxy clusters.

3.3 Stacking

In addition to doing this aperture photometry, we also use rectangular cutouts of the maps centered on the clusters as the to create unweighted and weighted stacks. These stacks provide a visual

representation of the SZ effect at the different frequencies. Additionally ,the mean of the aperture photometry values of all the clusters at a given frequency should be the same as the aperture photometry values of the corresponding stacks. Thus, we were able to use the stacks to confirm our aperture photometry methodology.




Figure 3.3: This shows the weighted stack stamp plots the RedMaPPer catalog. The 90 GHz and 150 GHz (subfigures a and b respectively) are on the ACT maps while the 217, 353, 545, and 857 GHz maps (subfigures c, d, e, and f respectively) are on the Planck maps.

Chapter 4

Results

Before doing the mcmc fitting, we performed a linear fit to the data by minimizing chi-squared. Here we only fit for two parameters, the Compton-y parameter and the integrated amplitude of the dust emission. Using this fit we found that $Y_c^{cyl} = 7.96871772 * 10^{-17}$ and $D_c = 5.74833793 * 10^{-18}$. This fit is plotted in the figure below.




Figure 4.1: This plot shows a linear fit of the data using only two fit parameters: Y (i.e. the Compton-y parameter) and D (the dust parameter). Note that the y-axis is on a log scale.

However, the linear fit algorithm was not effective at fitting for bulk kSZ or rSZ (in addition to Compton-y and dust). In order to effectively fit for these parameters we needed to include priors which constrained the possible values of the parameters. Without such priors, the fit algorithm resulted in physically nonsensical values (e.g. $T_{eff} < 0$ and bulk kSZ significantly deviating from 0). Thus, we used mcmc to fit with priors. For the mcmc fit with parameters Compton-y, dust, and effective temperature, we found that $Y = 6.3e - 17^{+2.1e-18}_{-2.3e-18}$, $D = 5.7e - 18^{+3.3e-19}_{-3.3e-19}$, and $T_{eff} < 1.4e8$. For the mcmc fit with parameters Compton-y, dust, and bulk kSZ, we found that $Y = 7.9e - 17^{+2.1e-18}_{-2.1e-18}$, $D = 5.8e - 18^{+3.3e-19}_{-3.4e-19}$, and $kSZ < -3.6e - 18$. The results of these fits are shown in the plots below.




Figure 4.2: This plot shows the corner plot of the results of the mcmc fit for Compton Y , dust, and the effective temperature. The priors used were $1e-17 < Y < 1e-14$, $1e-19 < D < 1e-16$ and $0.0 < T_{eff} < 1e10$. The resulting best fit parameters were $Y = 6.3e-17^{+2.1e-18}_{-2.3e-18}$, $D = 5.7e-18^{+3.3e-19}_{-3.3e-19}$, and $T_{eff} < 1.4e8$, where the errors on Y and D are 1 sigma and the constraint on T_{eff} is 2 sigma.




Figure 4.3: This plot shows the results of the mcmc fit for Compton Y , dust, and the effective temperature on a logarithmic y-scale. The priors used were $1e-17 < Y < 1e-14$, $1e-19 < D < 1e-16$ and $0.0 < T_{eff} < 1e10$. The resulting best fit parameters were $Y = 6.3e-17^{+2.1e-18}_{-2.3e-18}$, $D = 5.7e-18^{+3.3e-19}_{-3.3e-19}$, and $T_{eff} < 1.4e8$, where the errors on Y and D are 1 sigma and the constraint on T_{eff} is 2 sigma.




Figure 4.4: This plot shows the corner plot of the results of the mcmc fit for Compton Y , dust, and the bulk kSZ . The priors used were $1e-17 < Y < 1e-14$, $1e-19 < D < 1e-16$ and $-1e-17 < kSZ < 1e-17$. The resulting best fit parameters were $Y = 7.9e-17^{+2.1e-18}_{-2.1e-18}$, $D = 5.8e-18^{+3.3e-19}_{-3.4e-19}$, and $kSZ < 3.0e-18$, where the errors on Y and D are 1 sigma and the constraint on kSZ is 2 sigma.




Figure 4.5: This plot shows the results of the mcmc fit for Compton Y , dust, and the bulk kSZ on a logarithmic y-scale. The priors used were $1e-17 < Y < 1e-14$, $1e-19 < D < 1e-16$ and $-1e-17 < kSZ < 1e-17$. The resulting best fit parameters were $Y = 7.9e-17^{+2.1e-18}_{-2.1e-18}$, $D = 5.8e-18^{+3.3e-19}_{-3.4e-19}$, and $kSZ < -3.6e-18$, where the errors on Y and D are 1 sigma and the constraint on kSZ is 2 sigma.

Chapter 5

Discussion and Conclusion

Chapter 6

Future Research

Chapter 7

References

- J. P. Greco, J. C. Hill, D. N. Spergel, and N. Battaglia, *Astrophys. J.* 808, 151 (2015).
A. D. Hincks, R. Genova-Santos, G. Luzzia and E. S. Battistellia. Submitted to JCAP March 8th, 2018.
P. Bull, T. Clifton, and P. G. Ferreira *Phys. Rev. D* 85, 024002 (2012).