Initial-state matching and stability diagnostics for a counter-streaming electron-fluid closure

Taylor Kalin

taylorkalin.com teylersf@gmail.com

ORCID: 0009-0003-9000-9315

28 October 2026

Abstract

This diagnostic comment examines initial-state controls in arXiv:2607.07106v1 using the published decks, archived PIC moments and reloaded initial fluid output. The central two-population moment identities are correct. However, the PIC input specifies proper momentum where the fluid input specifies ordinary velocity. The saved fluid states corroborate their initializers to roundoff. Relative to either fluid model, the initial PIC moment-current profile differs by 9.87% in the full-domain, x-averaged relative ℓ^2 norm; the two lower-sheet central cells differ by about 11.6%. Their Cartesian velocity-covariance pressures are about 20% lower in p_{xx} and 38.4% lower in p_{zz}. A conditional public-loader calculation explains the finite-temperature drift reduction, while separating verified archived moments from an unverified custom diagnostic implementation. No native deposited-current file is identifiable in the complete posted PIC inventory. Separately, an auxiliary homogeneous longitudinal benchmark has a finite-isotropization fluid mode with growth $0.00952\omega_{pe}$ where its collisionless kinetic reference is Penrose-stable. This is not a sheet eigenanalysis or a refutation of the source's application-specific instability claim. We also qualify the symmetry-plane density explanation and give a cancellation-conditioned relative heat-flux bound. No new reconnection run is claimed. A matched-state rerun is required to establish dynamical significance; the source's later nonlinear conclusions are not disproved here.

1 Introduction and scope

Counter-streaming electron populations provide a useful representation of the strongly non-Maxwellian distributions generated during collisionless magnetic reconnection. McGrae-Menge and collaborators [1] report that tagging electrons by their initial side of a current sheet isolates sub-populations whose intrinsic third moments are small compared with their combined heat flux. Their exact moment decomposition explains how relative motion and population asymmetry can generate a substantial aggregate third moment even when each population has little intrinsic heat flux. Evolving additional population moments is therefore physically motivated.

The principal contribution of this diagnostic comment is an initial-state audit of the public inputs and archived moments. Equal nominal drift and temperature parameters need not specify equal laboratory-frame moments in a relativistic particle initializer and a nonrelativistic fluid initializer. This identifies controls needed to interpret the comparison; it does not determine their effect on its later nonlinear outcome.

Our first result uses the authors' publicly archived initial moments and input decks [2]. Identical nominal drift expressions enter a proper-momentum initializer in PIC and an ordinary-velocity

initializer in the fluid model. The recorded initial moments show a current and pressure mismatch consistent with that distinction. This is a reproducible control issue for interpreting the nonlinear comparison; its eventual dynamical effect cannot be determined from initial-state analysis alone.

An auxiliary result is a controlled longitudinal stability test of the stated continuum isotropization closure. A positive two-Maxwellian equilibrium is stable to longitudinal electrostatic perturbations under the established Penrose criterion [3], but its finite-relaxation fluid counterpart has an overstable mode. We give an explicit linear system and check both its eigenvalue residual and Gauss constraint. The test is homogeneous and nonrelativistic. It does not replace the source's inhomogeneous reconnection calculation, refute its application-specific instability-capability statement, or show that the identified mode caused any of its plotted features. The source populations initially share a velocity; the counter-streaming equilibrium used in this calibration test is different.

Two secondary observations make the interpretation more precise. Continuity depends on the divergence of the mass flux, so a zero normal velocity at a symmetry plane does not prohibit density accumulation there. A conditioned third-moment error bound also shows why a thermal-to-drift speed ratio alone does not give a universal relative-error guarantee when population contributions cancel. Neither result disputes the empirical parameter trend in the published scan.

1.1 Relation to established work

Heat-flux closure and its effect on collisionless response are established subjects. Hammett and Perkins [4] developed fluid moment models for Landau damping, while Wang and collaborators [5] compared multifluid moment models with PIC reconnection. Goldman and collaborators [6] studied multibeam energy moments and the distinction between component and aggregate centering. The mixture algebra used below belongs to this established moment framework.

Closure-dependent instability spectra are also known. Ng and collaborators [7] demonstrate why a pressure-tensor model must be checked against kinetic response rather than judged solely by its retained variables. Our finite-relaxation example supplies a concrete test associated with the closure and reference rate in the present source; we do not claim the general phenomenon of closure-dependent unstable modes as new.

Work on heat-flux closure for nonlinear electrostatic two-stream systems [8, 9] further limits a literal reading of broad claims that no fluid closure can capture non-Maxwellian counter-streaming. It does not establish prior use of the specific tagged-population closure in electron-only reconnection. Our analysis therefore addresses explicit initialization, continuity and stability statements, and preserves the source's application-specific empirical evidence.

1.2 Audited version and available evidence

The supplied ten-page PDF is byte-identical to a fresh versioned download of arXiv:2607.07106v1: 7,847,791 bytes, SHA-256 `fc5344e2ba8f4db98baf90d805bb6b7ce 8835c32ea7bf54bba4d62f405b955a3`. The official history checked on 28 October 2026 lists the 8 July v1 and no later revision or journal reference. The PDF displays 9 July, while the experimental HTML displays a different compilation date. The original TeX uses `\date{\today}`; those compilation dates do not identify additional scientific revisions. Appendix A records source and data locators. No original author has been contacted, and no author confirmation or private simulation-build access is claimed.

2 Initial-state matching in the supplied input decks

The source compares a relativistic PIC calculation with nonrelativistic ten-moment fluid calculations and describes their common initialization in Appendices A and C [1]. The deposited reference-case decks and initial moments make the physical matching of those states directly testable [2]. Equal numerical drift and temperature parameters need not specify equal laboratory-frame velocity moments in the two codes. This issue concerns the controlled comparison; the mixture identities do not depend on either code's initialization convention.

2.1 A proper-momentum parameter is used as a fluid velocity

The PIC deck sets `use_spatial_ufl=.true.` and supplies the curl-derived drift expressions as `spatial_ufl`. The official OSIRIS reference defines these inputs as proper velocities, or momentum divided by particle mass, with relativistic boosting as the default [10]. No classical-addition override appears in the posted deck. The Lua decks instead set nonrelativistic conserved momentum density to $\rho\mathbf{v}$ using the same expressions. This is the ordinary velocity variable of the ten-moment equations [5].

Writing $\mathbf{U} = \mathbf{p}/(m_e c)$, a cold population with proper drift $\mathbf{U}$ has

$$\frac{\mathbf{v}}{c} = \frac{\mathbf{U}}{\sqrt{1+|\mathbf{U}|^2}}. \tag{1}$$

Thus the same numerical vector used for $\mathbf{U}$ in PIC and $\mathbf{v}/c$ in the fluid initializer does not match the physical cold drift. At the unperturbed lower sheet center, the reference expressions give approximately $(0, 0, 0.5)$: the cold PIC speed is $0.4472136c$, whereas the prescribed fluid speed is $0.5c$. The fractional difference is 10.56%. It persists when the thermal spread tends to zero at fixed $B_{\text{Reconn.}} = 0.5$ and sheet width d_e; low thermal beta alone does not remove it. Equation (1) is a cold-limit explanation. A finite-temperature mean must be measured after loading rather than identified with that limit.

2.2 A conditional finite-temperature loading calculation

The complete 155-line posted PIC deck contains two thermal-distribution blocks with $\sigma = 0.111803$, and no classical-addition override or include directive. The inspected public OSIRIS loader draws independent Gaussian thermal proper-momentum components and boosts each particle without changing its weight [11]. With $c = 1$, for thermal momentum $\mathbf{w} \sim \mathcal{N}(0, \sigma^2 I)$ and proper drift $\mathbf{U}$, this rule is

$$\gamma_0 = \sqrt{1+|\mathbf{w}|^2}, \quad G = \sqrt{1+|\mathbf{U}|^2}, \quad \mathbf{p} = \mathbf{w} + \mathbf{U}\left(\gamma_0 + \frac{\mathbf{w}\cdot\mathbf{U}}{G+1}\right), \quad \mathbf{v} = \frac{\mathbf{p}}{G\gamma_0 + \mathbf{w}\cdot\mathbf{U}}.$$

This is a prediction conditional on the public loading rule, not an identification of the original executable. With $\beta = |\mathbf{U}|/G$ and $z = w_\parallel/\gamma_0$, pairing opposite Gaussian momenta gives

$$\mathbb{E}[v_\parallel] = \beta\left[1 - (1-\beta^2)\mathbb{E}\left(\frac{z^2}{1-\beta^2 z^2}\right)\right] < \beta$$

for positive drift and nonzero thermal spread. Thus the mean can fall below the cold conversion without an additional force. An invariant scalar distribution sampled at equal laboratory time would require a different Jacobian-weighted measure; it must not be silently substituted for this unweighted sample-boost rule.

At $\mathbf{U} = (0, 0, 0.5)$ the finite-temperature mean is $0.44297014c$, below the cold $0.44721360c$. Evaluating the full spatial drift expressions and perturbations at the two lower diagnostic rows gives an x-averaged zero-shape prediction $0.44197416c$. A separate, unfitted sensitivity uses the declared quadratic interpolation, the recorded moment-cell offset and the posted 25×25 regular particle lattice. Summing the standard quadratic weights over their support gives the following conditional values:

Quantity	$Y = 15.95d_e$	$Y = 16.05d_e$
Archived $\overline{v}_z/c$	0.44098722	0.44087669
Pointwise loading prediction	0.44197416	0.44197416
Quadratic-shape prediction	0.44098868	0.44098868
Archived P_{xx}	0.00994868	0.00997543
Quadratic-shape P_{xx}	0.00996490	0.00996490
Archived P_{zz}	0.00769653	0.00769303
Quadratic-shape P_{zz}	0.00770632	0.00770632

These pressure predictions use expected raw moments centered at each node before averaging, in the infinite-particle limit. No temperature, drift, shape width or perturbation is fitted to the archived values. Quadrature orders 12, 20, 32 and 48 agree in the quadratic mean to about $5 \times 10^{-16}c$; an independent Cartesian quadrature checks the reduced moment calculation to 1.73×10^{-15} in the tested quantities. This explains the scale of the additional reduction below the cold value. The standard public deposition kernel is not verified as the original custom velocity diagnostic's kernel, so the pointwise prediction and shape sensitivity remain distinct. Neither is a new PIC run or a confidence statement about the unidentified original build.

2.3 Archived initial current and staggered magnetic curl

The original data authors explicitly define the archived diagnostic `vfl_i` as $(1/n) \int f v_i \, \mathrm{d}^3 v$ [2]. It is therefore an ordinary mean velocity, even though the input parameter `ufl` is a proper-momentum parameter. With the source normalization $q_e = -1$, the physical moment current is

$$J_z^{\mathrm{mom}} = -\left(n_1 \overline{v}_{1z} + n_2 \overline{v}_{2z}\right). \tag{2}$$

The selected initial density diagnostics give $n_1 + n_2 = 1$ to within 1.5×10^{-8}. No particle tracking or new time integration is needed to evaluate this current.

The HDF5 files also record their grid offsets: density and velocity moments have $(0.5, 0.5)$, B_x has $(0.5, 1)$ and B_y has $(1, 0.5)$. The backward staggered differences therefore place

$$J^B_{z,i,j} = \frac{B_{y,i,j} - B_{y,i-1,j}}{\Delta x} - \frac{B_{x,i,j} - B_{x,i,j-1}}{\Delta y}, \qquad \Delta x = \Delta y = 0.1 d_e, \tag{3}$$

on the moment-cell centers. Periodic differencing makes the first term's x average exactly zero. The raw data use $y \in [-32, 32] d_e$; the paper's coordinate is $Y = y + 32 d_e$, so its lower sheet is at $Y = 16 d_e$.

The supplied fluid initializer prescribes an x-averaged ordinary velocity $0.49875256c$ in these same cells. The archived PIC means are $0.44098722c$ and $0.44087669c$, approximately 11.6% smaller. The small difference between this continuum prescription and Table 1's staggered curl does not explain the observed deficit. The adjacent cells at the other sheet show the same effect with reversed current sign.

Table 1: Initial x-averaged currents in the cells adjoining the lower sheet. The deficit is $1 - \langle J_z^{\mathrm{mom}}\rangle_x / \langle J_z^B\rangle_x$. Currents use the source's normalized units.

Y/d_e	$\langle J_z^{\mathrm{mom}}\rangle_x$	$\langle J_z^B\rangle_x$	Deficit
15.95	−0.44098723	−0.49833998	11.5088%
16.05	−0.44087669	−0.49833998	11.5309%

The result is not restricted to these four illustrative cells. Across all 640 rows, the relative ℓ^2 discrepancy between the x-averaged moment-current and Yee-curl profiles is 9.8265%. Restricting to $14 \le Y/d_e \le 18$ or $46 \le Y/d_e \le 50$ gives 9.8287% and 9.8408%, respectively. These are norms of the one-dimensional averaged profiles, not norms of the full two-dimensional fields. Figure 2 displays both sheets and the full domain. The input-prescribed current differs from the Yee profile by only 0.0630% in the full-profile relative norm.

As a deterministic discretization sensitivity, averaging the staggered fields onto cell centers before a centered derivative changes the curl profile by 0.3755% in that norm. This is a deliberately smoothed representation, not an alternative reading of the recorded Yee offsets. The central-cell moment-current deficits remain about 11.05–11.10% against that smoothed profile. Eight contiguous 60-column blocks at $Y = 15.95 d_e$ have moment-current means between −0.4417104 and −0.4404351, compared with the Yee mean −0.4983400. The per-column standard deviation is 0.0019906. These spatial summaries are descriptive; correlated particle deposition and the imposed perturbation preclude treating them as independent-sample confidence intervals.

Here J_z^{mom} is a moment-based physical diagnostic. We have not reloaded OSIRIS's native charge-conserving deposited current, which can differ from a simple qv deposition. These numbers therefore do not specify the exact discrete electric-field update of the original executable. In the normalized continuum equations, $\partial_t E_z = (\nabla \times \mathbf{B})_z - J_z$. The comparison is a continuum magnetostatic diagnostic. It does not establish an initial transient in the native numerical electric-field update.

2.4 The laboratory-frame pressure moments also differ

The record defines its second-moment diagnostics without a density normalization: $M_{s,ii} = \int f_s v_i^2 \, \mathrm{d}^3 v$ [2]. Following the source's own pressure definition, at $m_e = 1$ we compute

$$P_{ii} = M_{1,ii} + M_{2,ii} - n\bar{v}_i^2. \tag{4}$$

The posted fluid initializers prescribe an isotropic thermal pressure

$$p_{\mathfrak{fl}} = n_0 (0.111803)^2 = 0.012499910809.$$

The archived initial fluid `.bp` fields independently confirm this pressure in both fluid models, as detailed below. The measured ratios to the archived fluid pressure are

Y/d_e	$\langle P_{xx}\rangle_x / p_{\mathfrak{fl}}$	$\langle P_{zz}\rangle_x / p_{\mathfrak{fl}}$
15.95	0.795900	0.615727
16.05	0.798040	0.615447

For example, at $Y = 15.95 d_e$ the archived values are $\langle P_{xx}\rangle_x = 0.00994868$ and $\langle P_{zz}\rangle_x = 0.00769653$. The latter is about 38.4% below the archived fluid pressure. These are the ordinary-velocity covariance moments defined in the source, not components of a relativistic energy–momentum tensor. A

common rest-frame thermal-momentum parameter does not imply equal laboratory-frame velocity covariance after a boost. Centering the pressure after averaging in x instead adds the density-weighted variance of the resolved mean flow. At the four central rows this changes either diagonal component by approximately 4×10^{-6}, far smaller than the reported pressure mismatch; the source definition and our principal result center each cell first. Figure 1 shows the local x-averaged profiles.

2.5 Corroboration against the saved fluid state

We reloaded five initial BP snapshots: the one-fluid electron state, both two-fluid electron states, and each model's electromagnetic field. They record time and frame zero, a single available step, the same 480×640 grid and bounds, zero ghost cells and polynomial order zero. Each file's embedded base64 input deck decodes byte-for-byte to the posted Lua file. This checks the input used in the saved output, rather than assuming that the separate prescription was executed unchanged.

The ten stored components are ρ, the three ρu_i, and the six symmetric raw second moments. Their order is checked against every returned initializer component over all 307,200 cells, including off-diagonal moments and the field components. The largest one-fluid component error is 4.38×10^{-16}; the summed two-fluid state differs from the one-fluid state by at most 1.12×10^{-16}. Aggregate pressure is reconstructed after summing the conservative population moments. Thus the earlier comparison with the prescribed initial state is corroborated by the actual archived state.

Against either reloaded fluid model, the full 640-row, x-averaged PIC profiles have relative ℓ^2 differences of 9.8675% for J_z, 8.3169% for P_{xx} and 9.5440% for P_{zz}. At the two lower central rows the actual fluid current is -0.4987525573, giving moment-current deficits 11.5820% and 11.6041%. The local pressure ratios reported above are unchanged. These comparisons concern physical velocity moments, not native PIC current deposition or later-time dynamics.

All selected PIC files have `TIME=0` and `ITER=0`; the fluid files have time and frame zero. No explicit time-averaging attribute appears. This verifies the posted snapshot metadata, not the impossibility of an undocumented operation in the unidentified PIC build. A complete inventory of all 223 PIC ZIP entries, including 94 scientific files, identifies no native deposited-current diagnostic. We therefore cannot make the requested native-current comparison from this deposit and do not substitute the `UDIST` moment current for it.

2.6 What a controlled comparison requires

The data establish an initial-state mismatch in the posted comparison. They do not establish how much of any later reconnection difference it causes. That requires reruns with controlled initial moments and numerical refinement. One constructive test is to initialize the fluid models from the measured PIC n, $\overline{\mathbf{v}}$ and P_{ij}, preserving the intended population labels. Another is to calibrate the PIC loading to a prescribed physical fluid state and then verify the deposited moments. The cold conversion in Eq. (1) is necessary for a cold drift match, but by itself does not guarantee a finite-temperature pressure or current match.

The fluid BP metadata records Gkeyll changeset `9663ea594e80+`; its plus suffix does not identify a clean exact source tree or executable. The original PIC executable remains unidentified. The current public OSIRIS code inspected for documentation does not expose the authors' custom `vfl`/third-moment diagnostic implementation. Our diagnostic interpretation follows their explicit dataset definitions and file metadata. All selected files are at $t = 0$. The present calculation is a reproducible reanalysis of those files, with no new PIC or Gkeyll run and no causal attribution of the source's later nonlinear results.




Figure 1: Reanalysis of the archived initial state near the lower sheet. Left: the ordinary-velocity moment current differs from both the staggered magnetic curl and the posted fluid prescription. Right: the local centered pressure components, averaged in x after centering, differ from the isotropic fluid pressure. The profiles use the authors' archived diagnostics and their published definitions [2]. The drawn fluid curves use the initializer; the reloaded initial fluid output independently corroborates them to roundoff. No new simulation is shown.




Figure 2: Full-domain, x-averaged archived PIC profiles at $t = 0$, with both current sheets visible. Left: physical moment current and the correctly staggered magnetic curl. Right: velocity-covariance pressures divided by the fluid pressure confirmed in the archived output. The figure extends the local view; it is neither an independent simulation nor a native deposited-current diagnostic.

3 An auxiliary homogeneous linear-stability test

The source argues that its two-population description can capture linear growth of counter-streaming instabilities in reconnection [1]. Our calculation does not test those evolving distributions. It supplies a separate calibration example for the implemented local-isotropization closure: population drifts alone do not guarantee a correct collisionless kinetic spectrum. Closure-dependent differences in such spectra are established issues [7, 5]; the example below makes this validation issue explicit at the stated reference relaxation-rate scale.

Consider a homogeneous nonrelativistic electron plasma with immobile neutralizing ions. Two equal-density populations have ordinary velocities $\pm U e_x$ and isotropic thermal variance σ^2. Their longitudinal velocity marginal, normalized to unit density, is

$$f_0(v) = \frac{e^{-(v-U)^2/(2\sigma^2)} + e^{-(v+U)^2/(2\sigma^2)}}{2\sqrt{2\pi}\sigma}. \tag{5}$$

A constant magnetic field parallel to e_x leaves the longitudinal, gyrotropic perturbation subspace used below invariant. In the kinetic equation its gyrophase-rotation operator vanishes on gyrotropic perturbations; a longitudinal electric perturbation preserves that subspace. In the fluid equations, parallel velocity perturbations and a diagonal pressure perturbation with equal transverse entries have vanishing magnetic rotation terms. This justification concerns a uniform parallel field, not the source's rotating field, transverse gradients or relativistic response.

The collisionless kinetic reference. For $U > \sigma$, the only local minimum of f_0 is at zero. Indeed, $f_0'/f_0 = [-v + U\tanh(Uv/\sigma^2)]/\sigma^2$, and $U\tanh(Uv/\sigma^2) - v$ is strictly concave on $v > 0$, with exactly one positive zero. The Penrose criterion [3] therefore tests

$$I = \int_{\mathbb{R}} \frac{f_0'(v)}{v}\,\mathrm{d}v = \frac{2\zeta D(\zeta) - 1}{\sigma^2}, \qquad \zeta = \frac{U}{\sqrt{2}\sigma}, \quad D(z) = e^{-z^2}\int_0^z e^{t^2}\,\mathrm{d}t. \tag{6}$$

This integrand has a removable singularity at zero. At $U/\sigma = 6/5$, $\zeta^2 = 18/25$. The alternating Dawson series has decreasing term magnitudes and gives the rational upper bound

$$2\zeta D(\zeta) \le \frac{36}{25} - \frac{432}{625} + \frac{15552}{78125} = \frac{74052}{78125} < 1. \tag{7}$$

Consequently $I < 0$, and this kinetic equilibrium has no exponentially growing longitudinal electrostatic mode for any real wavenumber. Direct quadrature gives $\sigma^2 I = -0.08730405848\ldots$; the sign does not depend on that numerical evaluation.

Linearizing the finite-relaxation fluid model. The source retains $\partial_m q_{sijm} = \nu(p_{sij} - \delta_{ij}\,\mathrm{tr}\,p_s/3)$, with $\nu = \kappa\sigma > 0$. Normalize the total plasma frequency, density, electron mass, charge magnitude and permittivity to one. For each beam $s = \pm$, set $U_s = sU$, $n_s = 1/2$, and equilibrium diagonal pressure $p_s^0 = n_s\sigma^2$. Write its perturbations as $N_s = \delta n_s$, $V_s = \delta u_{sx}$, $P_s = \delta p_{sxx}$ and $H_s = \delta\,\mathrm{tr}\,p_s$. For spatial factor e^{ikx}, the invariant nine-variable system is

$$\begin{aligned}
\dot N_s &= -ikU_sN_s - ikn_sV_s,\\
\dot V_s &= -ikU_sV_s - ikP_s/n_s - E,\\
\dot P_s &= -ikU_sP_s - 3ikp_s^0V_s - \nu(P_s - H_s/3),\\
\dot H_s &= -ikU_sH_s - 5ikp_s^0V_s,\\
\dot E &= \sum_{s=\pm}(U_sN_s + n_sV_s).
\end{aligned} \tag{8}$$

Here $\delta p_{syy} = \delta p_{szz} = (H_s - P_s)/2$. A perturbation of ν multiplies the zero equilibrium deviator and thus contributes nothing at linear order, even when thermal speed is computed locally. Eliminating these variables independently, with temporal convention $e^{-i\omega t}$, gives

$$\epsilon_{\mathrm{fl}}(\omega, k) = 1 - \frac{1}{2}\sum_{s=\pm} \frac{1}{(\omega - kU_s)^2 - k^2\sigma^2 g_s},$$
$$g_s = \frac{5}{3} + \frac{4D_s}{3(D_s + \nu)}, \qquad D_s = -i(\omega - kU_s). \tag{9}$$

Proposition 3.1 (A finite-relaxation false positive)**.** *At $U/\sigma = 6/5$ and $k\sigma/\omega_{pe} = 1/5$, the collisionless kinetic equilibrium is stable to longitudinal electrostatic perturbations, while the local-isotropization fluid model is overstable for all sufficiently small positive ν.*

Proof. Kinetic stability follows from (7). At $\nu = 0$ the longitudinal fluid pressure response has index three. Putting $W = \omega^2$, $a = k^2U^2$, $b = 3k^2\sigma^2$, its dispersion equation is

$$W^2 - [2(a + b) + 1]W + (a - b)(a - b - 1) = 0. \tag{10}$$

In units $\sigma = \omega_{pe} = 1$, its simple lower positive frequency is $\omega_0 = [(847 - 5\sqrt{24553})/1250]^{1/2} = 0.2254420613\ldots$. Let $\Omega_s = \omega_0 - kU_s$ and $A_s = \Omega_s^2 - 3k^2$. Implicit differentiation of (9) yields

$$\left.\frac{\mathrm{d}\ \mathrm{Im}\,\omega}{\mathrm{d}\nu}\right|_{\nu=0} = -\frac{2k^2}{3}\frac{\sum_s 1/(\Omega_s A_s^2)}{\sum_s \Omega_s/A_s^2} = 2.488846777\ldots > 0. \tag{11}$$

The sign can also be checked without the decimal value: the rational bounds $0.225 < \omega_0 < 0.226$ make the numerator sum negative and the denominator sum positive. The root is simple, so its continuation has positive imaginary part for sufficiently small positive relaxation. □

At a reference-scale rate obtained from $\kappa = 0.1/d_e$ and $\sigma = c/\sqrt{80}$, $\nu/\omega_{pe} = 1/(40\sqrt{5}) = 0.01118033989\ldots$, the matrix in (8) numerically gives

$$\frac{\lambda}{\omega_{pe}} = 0.00951849926 \ \pm\ 0.21649095264\,i, \qquad \lambda = -i\omega. \tag{12}$$

The eigen-equation, Gauss constraint $ikE + N_+ + N_- = 0$, and independent dielectric residual are all below 2×10^{-10} in the recorded checks; actual residuals are much smaller. The zero-relaxation matrix has no positive growth beyond roundoff. The literal fluid deck rounds its thermal speed to $0.111803c$; substituting the resulting $\nu/\omega_{pe} = 0.0111803$ also leaves positive growth, with a change smaller than $10^{-6}\omega_{pe}$. A periodic-box check using $k_x = 2\pi\,14/L_x$ for $L_x = 48d_e$, with $\sigma = c/\sqrt{80}$, gives growth $0.00951299405\omega_{pe}$ and a wavelength resolved by about 34 cells at the reported grid spacing. It is an x-periodic mode of the homogeneous system, not a transverse k_y eigenmode of the inhomogeneous current sheet.

These are continuum model diagnostics. They establish a false positive relative to the specified collisionless kinetic reference, including at the source's positive relaxation-rate scale. They do not identify a mode measured in the original current sheet, replace a kinetic stability analysis of that sheet, or show that a particular nonlinear plot is contaminated. The source's initial populations have a common velocity and its equilibrium varies across y. No survival of this mode in that equilibrium is established. The benchmark is retained as an auxiliary closure test, rather than a counterexample to the source's narrower application-specific capability claim.




Figure 3: Homogeneous longitudinal test with $U/\sigma = 1.2$ and $k\sigma/\omega_{pe} = 0.2$. Left: nonzero eigenvalues of the ideal and finite-rate fluid systems; the static Gauss-constraint mode is excluded. Right: the largest fluid growth rate as relaxation varies, with $\nu_0/\omega_{pe} = 1/(40\sqrt{5})$ marked. The collisionless kinetic reference has no growing longitudinal electrostatic mode by the Penrose proof above. This comparison concerns the stated continuum closure, not an observed mode in the original reconnection calculation.

4 A secondary continuity clarification

On PDF p. 5, the source attributes its one-fluid density deficit to the vanishing of u_y at the symmetric mid-plane, describing this as blocking transport into the sheet [1]. The value of a flux at one plane does not determine its divergence:

$$\partial_t n = -\partial_x(nu_x) - \partial_y(nu_y). \tag{13}$$

In particular, $\partial_y(nu_y)$ can be nonzero where $u_y = 0$. This is standard continuum kinematics, not a new plasma mechanism. Its role here is to qualify an explanation based on symmetry alone; the observed one-fluid density deficit and its dynamical cause remain separate questions.

Proposition 4.1 (A periodic density-accumulation example)**.** *For $a, n_0 > 0$, on a periodic coordinate $y \in [0, 2\pi)$,*

$$u_y(y) = -a \sin y, \qquad n(y,t) = \frac{n_0}{\cosh(at) - \cos y \, \sinh(at)} \tag{14}$$

is a smooth positive solution of the one-dimensional continuity equation. Its total mass is $2\pi n_0$, and $u_y(0) = 0$ while $n(0,t) = n_0 e^{at}$.

Proof. The denominator is positive for every finite t. Differentiating (14) gives $\partial_t n + \partial_y(nu_y) = 0$ directly. The standard integral $\int_0^{2\pi}(A - B\cos y)^{-1}\mathrm{d}y = 2\pi/(A^2 - B^2)^{1/2}$, with $A = \cosh(at)$ and $B = \sinh(at)$, gives the conserved mass. Evaluation at zero gives the stated increasing central density. □

This is an exact kinematic transport example, not a self-consistent plasma–Maxwell solution or a substitute reconnection run. It refutes the inference from symmetry alone. The two-population

model's total density and density-weighted velocity obey exactly the same continuity equation after the population equations are summed. A one-fluid model loses the separate population-crossing information, but it is not geometrically forbidden from supplying or accumulating total density within a finite-width layer. Differences in the evolution of aggregate pressure and higher moments can still explain a particular benchmark failure.

5 Conditioning a relative heat-flux accuracy claim

The source's Eqs. (3)–(4) and (B1)–(B2) are correct. For $n_1, n_2 > 0$, put $n = n_1 + n_2$, $w_s = n_s/n$, $\Delta u = u_1 - u_2$ and $T_s = p_s/n_s$. Recentring each population on the aggregate mean gives

$$\begin{aligned} p_{ij} &= p_{1ij} + p_{2ij} + mnw_1w_2\Delta u_i\Delta u_j, \\ q_{ijk} &= q_{1ijk} + q_{2ijk} + nw_1w_2(\Delta u_i\Delta T_{jk} + \Delta u_j\Delta T_{ik} + \Delta u_k\Delta T_{ij}) \\ &\quad - mnw_1w_2(w_1 - w_2)\Delta u_i\Delta u_j\Delta u_k. \end{aligned} \tag{15}$$

The source explicitly defines the unnormalized three-term symmetrization, so its tensor formula has the required scalar factor three. Direct positive finite-mixture calculations, including nonunit mass, verify all entries.

For the scalar xxx component and $\Delta u \neq 0$, define $\theta_s = T_{sxx}/[m(\Delta u)^2]$. Its counter-streaming contribution is

$$q_{\mathrm{CS}} = mn(\Delta u)^3 C, \qquad C = w_1w_2\{3(\theta_1 - \theta_2) - (w_1 - w_2)\}. \tag{16}$$

Thus the source's cubic-drift explanation of the empirical parameter collapse requires qualification. Equation (16) also contains a temperature-contrast term linear in drift and can vanish by symmetry or cancellation.

Proposition 5.1 (A sufficient conditioned relative-error bound)**.** *Suppose* $|q_{sxxx}| \le mn_s a_s v_{\mathrm{th},s}^3$, *with* $v_{\mathrm{th},s} = (T_{sxx}/m)^{1/2}$ *and stated nonnegative bounds* a_s. *Set* $E = \sum_s w_s a_s (v_{\mathrm{th},s}/|\Delta u|)^3$. *If* $E < |C|$, *then*

$$\frac{|q_{xxx} - q_{\mathrm{CS}}|}{|q_{xxx}|} \le \frac{E}{|C| - E}. \tag{17}$$

Proof. The exact residual is $r = q_{1xxx} + q_{2xxx}$, so $|r| \le mn|\Delta u|^3 E$. The reverse triangle inequality gives $|q_{xxx}| \ge mn|\Delta u|^3(|C| - E) > 0$. Division gives (17). □

The bound is sufficient rather than necessary. Its imbalance/cancellation factor cannot be dropped. Equal population densities and temperatures give $q_{\mathrm{CS}} = 0$ at any drift. With equal densities but unequal temperatures, $q_{\mathrm{CS}} = 3n\Delta u\,\Delta T_{xx}/4$, which is linear in drift.

A near-Maxwellian counterexample to a speed-ratio-only guarantee. Let W take values $-1/\sqrt{2}$ and $\sqrt{2}$ with probabilities 2/3 and 1/3. Then $\mathbb{E}[W] = 0$, $\mathbb{E}[W^2] = 1$ and $\mathbb{E}[W^3] = 1/\sqrt{2}$. For $\eta > 0$, let $S = (W + \eta G)/\sqrt{1 + \eta^2}$ with G a standard Gaussian. Let $H = (S, G_y, G_z)$, with independent standard Gaussian transverse components, and define X_ϵ to have law $(1 - \epsilon)\mathcal{N}(0, I) + \epsilon\mathcal{L}(H)$, $0 < \epsilon < 1$. It has zero mean, covariance I, and third moment $\mathbb{E}\left[X_{\epsilon,x}^3\right] = \epsilon/[\sqrt{2}(1 + \eta^2)^{3/2}]$. Its total-variation distance from $\mathcal{N}(0, I)$ is at most ϵ. Give the populations equal densities and velocity laws

$$V_1 = Ue_x + \sigma X_\epsilon, \qquad V_2 = -Ue_x + \sigma X_\epsilon. \tag{18}$$

Each population density is a positive smooth Gaussian mixture, within ϵ in total variation of its matching Maxwellian. Their intrinsic temperature tensors are both $m\sigma^2 I$, so $q_{\mathrm{CS}} = 0$, whereas

$q_{xxx} = mn\sigma^3\epsilon/[\sqrt{2}(1+\eta^2)^{3/2}] \neq 0$. Consequently the relative closure error is one even when both ϵ and the thermal-to-beam speed ratio σ/U are arbitrarily small. The absolute residual does tend to zero relative to mnU^3. The recorded calculations use $\epsilon = 0.01$ and integrate the smooth laws with positive Gauss–Hermite quadrature.

This construction does not disprove the source's measured collapse in its particular family of reconnection simulations. It identifies assumptions needed to extend that empirical observation to a relative-error guarantee. The source scan uses initial speeds; inserting those into a local bound would also require estimates of their subsequent evolution.

The valid limitation of a memoryless one-fluid closure. Additional population information can be genuinely necessary. For example, let $n = m = 1$, and let each population be an isotropic Gaussian of variance $3I/4$. Choose a weight w, set $\Delta = [4w(1-w)]^{-1/2}$, and assign means $(1-w)\Delta e_x$ and $-w\Delta e_x$. For both $w = 1/2$ and $w = 1/4$, the aggregate density is one, the mean is zero, and the pressure is $\mathrm{diag}(1, 3/4, 3/4)$. The respective q_{xxx} values are zero and $1/(4\sqrt{3})$. Hence no arbitrary-distribution algebraic function of only n, u, p can recover both third moments. This supports the source's population-resolving motivation. It is not an impossibility statement for one-fluid descriptions with memory, extra variables, gradients or nonlocality.

6 Finite isotropization over a comparison interval

The source openly retains isotropization for numerical stability [1]. Its smallness relative to a fast local heat-flux scale does not establish closeness to zero-intrinsic-heat-flux evolution over an entire simulation. A homogeneous calculation makes the distinct finite-time parameter explicit.

At fixed density, let p be a positive-definite pressure tensor and $\bar{p} = \operatorname{tr} p/3$. The relaxation part alone satisfies

$$\dot{p} = -\nu(p - \bar{p}I), \qquad p(t) = \bar{p}I + e^{-\nu t}(p(0) - \bar{p}I). \tag{19}$$

Its trace, and thus its thermal energy, are constant. For the Gaussian entropy diagnostic per particle, $s_G = \frac{1}{2}\log\det p - \frac{5}{2}\log n + \text{constant}$, one obtains

$$\dot{s}_G = \frac{\nu}{2}(\bar{p}\,\operatorname{tr}(p^{-1}) - 3) \geq 0, \tag{20}$$

with equality only at isotropy. The inequality follows by applying the arithmetic–harmonic mean inequality to the three positive eigenvalues of p. This is an entropy diagnostic of the moment model, not a measurement of PIC entropy or evidence of external heat input.

The deviation from the zero-relaxation homogeneous pressure is controlled by $1 - e^{-\nu T}$, not by ν alone. At the ideal reference-scale rate $\nu/\omega_{pe} = 1/(40\sqrt{5})$ and $T = 300\omega_{pe}^{-1}$,

$$\nu T = 3.354101966\ldots, \qquad e^{-\nu T} = 0.0349407340\ldots. \tag{21}$$

Thus an initially present homogeneous anisotropy would retain about 3.49% of its amplitude over that interval. For the example $p(0) = \mathrm{diag}(2, 1/2, 1/2)$ at unit density, the entropy diagnostic increases by $0.3461208970\ldots$ while the trace remains three.

This example does not establish how much relaxation changes the evolving source current sheet, whose anisotropy is time-dependent. It shows why a finite-duration sensitivity study in κ, or a direct comparison of the retained pressure-source terms, is needed when interpreting the implementation as approximately zero-heat-flux evolution. It does not dispute the source's disclosed need for finite stabilization, nor identify irreversible adiabatic relaxation with external energy transfer.

Table 2: Selected verification outcomes. The code, data definitions and scope limits distinguish initial-state reanalysis from independent continuum tests.

Check	Outcome
Initial moment current at $Y = 16.05 d_e$	11.5309% below staggered curl
Initial P_{zz}/p_{fl} at that row	0.615446655
Exact mixture identities	All scalar and tensor checks pass
Symmetry-plane continuity	Zero normal velocity; increasing density
Conditioned relative-error example	$0.024134 \leq 0.036243$
Identical one-fluid lower moments	Different third moments
Finite-relaxation mode growth	$0.009518499\,\omega_{pe}$
Homogeneous relaxation over $300/\omega_{pe}$	3.4941% anisotropy remains

7 Reproducible calculations and evidence limits

The repository separates reanalysis of archived simulation output from analytical model tests. The initial-state analysis uses the authors' data dictionary, independently readable HDF5 attributes, published input decks, and explicitly aligned grid locations. The stability, continuity and moment calculations use specified equations and inputs and require no reconnection simulation executable.

The inspected Zenodo record is identified by the record-specific DOI 10.5281/zenodo.21114117; the source's concept DOI is 10.5281/zenodo.21114116. Both resolve to this dataset family. The registration agency is DataCite, whose API identifies the record DOI as findable; a Crossref-only lookup returns no record. The direct repository link is `https://zenodo.org/records/21114117`. These identifiers do not denote two different simulations or a replacement data source.

The data retriever requests only selected ZIP members from the public Zenodo archive. It requires bounded HTTP partial responses, checks member lengths and CRC32 against the ZIP directory, and records SHA-256 hashes of the extracted inputs. The published checksum of the complete archive is identified as publisher-supplied; a partial retrieval does not verify that whole-archive checksum. Small derived profiles and provenance are included with their attribution. The original paper and full simulation archives are not redistributed by this project.

Numerical checks preserve the exact mixture identities rather than treating them as errors. The linear calculation is checked independently through an eliminated dielectric relation and the Gauss constraint, in addition to the matrix eigenvalue residual. The kinetic comparison has an exact rational stability bound, so its conclusion does not depend on locating a numerical threshold near marginal stability. The continuity example satisfies the transport equation explicitly. These checks test the scientific statements used in the paper, not only whether a plotting script runs.

The execution record in Appendix B pins the code, input hashes, numerical results and environment. Its timestamps are author-environment execution evidence, not third-party certification or independent human peer review. The accompanying verification script reports actual failures instead of silently replacing absent inputs with synthetic data.

No OSIRIS or Gkeyll reconnection run was performed for this study. The published input decks and archived diagnostics do not uniquely identify the authors' exact executable builds. In particular, the documented custom velocity diagnostics are interpreted using the authors' explicit data dictionary; we do not assert that a present public source checkout is the private build that produced them. This limits claims about implementation internals while leaving the documented archived-moment comparison testable.

8 Consequences for controlled validation

An initialization audit should precede attributing every later difference to the heat-flux closure. The current test exposes a mismatch in physical moments despite matching algebraic drift expressions. A controlled repeat should match density, ordinary mean velocity and the selected pressure definition, and report the initial magnetic curl and moment current on their actual staggered locations. If relativistic particle initialization and a nonrelativistic fluid model are compared, the conversion and the target moment convention must be stated explicitly. The cold proper-velocity conversion alone is not a proof that finite-temperature moments have been matched.

The measured initial mismatch does not determine the sign or size of its effect at the reported $t = 300\omega_{pe}^{-1}$ snapshot. Reconnection may relax some initial differences, amplify others, or preserve the qualitative advantage of retaining two populations. A matched-initial-state comparison and resolution or relaxation-rate studies are needed to separate those possibilities. We therefore do not infer that correcting the drift would eliminate the reported single-fluid density deficit, reproduce all PIC heat flux with one fluid, or reverse the source's qualitative conclusion.

The auxiliary homogeneous stability test supplies a separate calibration. Retaining population drift creates the possibility of streaming modes; it does not ensure the correct kinetic susceptibility or growth rate. The implemented relaxation should be included in the dispersion calculation. Finite relaxation can change a spectrum even when the unperturbed pressure is isotropic and the relaxation source vanishes at equilibrium. In the stated example the resulting growing mode is a false positive relative to the collisionless kinetic reference. This is a test of a model's continuum response, not evidence that the original simulation visibly contains that mode. We have not performed the source-sheet eigenanalysis or a matched-state reconnection rerun requested for a causal assessment. The present contribution is a diagnostic comment whose nonlinear significance remains undetermined.

The pressure-source diagnostic should also distinguish a third-moment amplitude from its divergence. The scale proxy $p\,d_e/|q|$ used in the source is a useful estimate when d_e is an appropriate gradient length. It is not identical to $p/|\nabla\cdot q|$. A spatially constant nonzero heat-flux tensor has a finite amplitude proxy and zero divergence. The source explicitly notes that its isotropization closure constrains a divergence and omits the reconstructed fluid tensor from some plots; we do not characterize that disclosure as a hidden omission. Direct source-term comparisons and rate convergence would make the approximation more informative.

Finally, the source's time-maximum spatial score is a best-time diagnostic, not a uniform-in-time error bound or a forward prediction test at every scan point. The conditioned estimate above explains why both absolute residuals and a nondegenerate denominator are needed when assessing relative accuracy. These qualifications preserve the observed scan and suggest measurements needed before extending its interpretation to new current sheets, later island recirculation, or turbulence.

The valid component-to-aggregate moment decomposition remains the foundation of this analysis. Additional labels carry information absent from a generic state described only by total density, velocity and pressure. The corrections identify controls and assumptions needed for using that additional information reliably, rather than a general rejection of multicomponent fluid modeling.

9 Conclusion

Matching a nominal drift expression does not match a proper-momentum PIC initialization to an ordinary-velocity fluid initialization. The archived PIC moments and reloaded fluid output make that distinction quantitatively testable. The full-profile and finite-temperature checks strengthen that initial-state finding while leaving its nonlinear importance undetermined. The separate ho-

mogeneous benchmark shows why finite isotropization should be calibrated as part of a closure; it establishes no unstable eigenmode of the source's inhomogeneous sheet. The exact moment decomposition remains valid and useful. Correcting the density-transport explanation and conditioning relative-error statements provides a constructive basis for further comparison, without asserting that the reported nonlinear reconnection results have been disproved.

Author and methods disclosure

Taylor Kalin is the author and is responsible for this study. OpenAI Codex assisted with source inspection, derivations, code, data analysis, and manuscript preparation under his direction. Separate AI-agent checks were used during preparation; they are not represented as independent human expert review. Reported results are deterministic classical computations and reanalysis of public archived moments. No new reconnection simulation or experimental measurement is claimed.

Funding and competing interests. This study was self-funded by Taylor Kalin, including computing resources and AI tools. No external funding was received. No competing interests were reported for this study.

Prior dissemination and review sequencing

This is one study. GitHub and AiraXiv deposits are preprint versions of the same work. Agent-only editions, where used, identify Taylor Kalin as canonical author and human operator; they are not independent new findings. Review venues are sequenced to avoid concurrent consideration of the same manuscript. Actual identifiers and outcomes are recorded at https://github.com/Teylersf/counterstreaming-heatflux-audit in the publication-status record.

References

[1] Madox C. McGrae-Menge, Jacob R. Pierce, Maria Almanza, Alexander Velberg, Nathaniel Barbour, William D. Dorland, Nuno F. Loureiro, Frederico Fiuza, and E. Paulo Alves. Counter-streaming heat-flux closure for electron-only collisionless magnetic reconnection, 2026. Audited version 1, submitted 8 July 2026. https://arxiv.org/abs/2607.07106v1.

[2] Madox McGrae-Menge. Counter-streaming heat-flux closure for electron-only collisionless magnetic reconnection. Zenodo, record 21114117, July 2026. Published 2 July 2026. DataCite record DOI: 10.5281/zenodo.21114117; concept DOI: 10.5281/zenodo.21114116. Input decks and archived diagnostics. https://zenodo.org/records/21114117.

[3] Oliver Penrose. Electrostatic instabilities of a uniform non-Maxwellian plasma. *The Physics of Fluids*, 3(2):258–265, 1960. https://doi.org/10.1063/1.1706024.

[4] Gregory W. Hammett and Francis W. Perkins. Fluid moment models for Landau damping with application to the ion-temperature-gradient instability. *Physical Review Letters*, 64(25):3019–3022, 1990. https://doi.org/10.1103/PhysRevLett.64.3019.

[5] Liang Wang, Ammar H. Hakim, A. Bhattacharjee, and K. Germaschewski. Comparison of multi-fluid moment models with particle-in-cell simulations of collisionless magnetic reconnection. *Physics of Plasmas*, 22(1):012108, 2015. https://doi.org/10.1063/1.4906063.

[6] M. V. Goldman, D. L. Newman, J. P. Eastwood, and G. Lapenta. Multibeam energy moments of multibeam particle velocity distributions. *Journal of Geophysical Research: Space Physics*, 125(12):e2020JA028340, 2020. https://doi.org/10.1029/2020JA028340.

[7] Jonathan Ng, Ammar Hakim, J. Juno, and A. Bhattacharjee. Drift instabilities in thin current sheets using a two-fluid model with pressure tensor effects. *Journal of Geophysical Research: Space Physics*, 124(5):3331–3346, 2019. https://doi.org/10.1029/2018JA026313.

[8] Emil R. Ingelsten, Madox C. McGrae-Menge, E. Paulo Alves, and Istvan Pusztai. Data-driven discovery of a heat flux closure for electrostatic plasma phenomena. *Journal of Plasma Physics*, 91(2):E64, 2025. https://doi.org/10.1017/S0022377825000285.

[9] Emil R. Ingelsten, Madox C. McGrae-Menge, E. Paulo Alves, and Istvan Pusztai. Data-driven multi-species heat flux closures for two-stream-unstable plasmas with nonlinear sparse regression. *Physics of Plasmas*, 33(4):042106, 2026. https://doi.org/10.1063/5.0311956.

[10] OSIRIS Collaboration. Momentum distribution: `udist`. OSIRIS Reference Guide. https://osiris-code.github.io/osiris/reference/udist. Accessed 28 October 2026; current public documentation, not identification of the original simulation build.

[11] OSIRIS Collaboration. OSIRIS momentum-distribution and particle-loading source code. Public source, commit a858fe1da3d6dbf3652b16556af87f7dad7881eb. Accessed 28 October 2026. Conditional loader interpretation; not identification of the original executable. https://github.com/osiris-code/osiris/tree/a858fe1da3d6dbf3652b16556af87f7dad7881eb/source/spec.

A Source passages and artifact scope

All source equation and page locators refer to the PDF of arXiv:2607.07106v1 [1]. The supplied ten-page PDF is byte-identical to a fresh versioned arXiv download: 7,847,791 bytes, SHA-256

```
fc5344e2ba8f4db98baf90d805bb6b7c
e8835c32ea7bf54bba4d62f405b955a3.
```

The official history checked on 28 October 2026 lists v1 submitted 8 July 2026 at 07:43:25 UTC, with no later revision or journal reference. The PDF displays 9 July; the experimental HTML displays 24 August. The original `apssamp.tex`, line 95, uses `\date{\today}`, so separately compiled display dates are not version identifiers. No source author has been contacted, and no author confirmation is claimed.

The separately retrieved versioned source archive has 7,439,982 bytes and SHA-256

```
4f9abd5f41ffca0f1879981429165ff1
a989276792c812e860f57cb5038c3b2e.
```

The repository's selective excerpts contain six attributed files and 17 fragments, with original TeX line and character positions. A standalone verifier checks the archive and source hashes and reconstructs every excerpt without executing the source or making network requests. The full PDF and source archive are not redistributed.

The stated initialization. Appendix A, p. 6, states that the electron fluid velocities are initialized from

$$\epsilon_{ijk}\partial_j B_k = J_i = q_e n_0 \overline{v}_i.$$

Appendix C, p. 8, states that the fluid simulations inherit the PIC domain, grid, boundary conditions and magnetic-field initial condition exactly. It specifies the common initial drift

$$\overline{v}_{1i} = \overline{v}_{2i} = \overline{v}_i$$

and thermal speed $0.111803c$ for the electron fluids. These are the matching statements tested in the main text. Original TeX locators are lines 225, 244 and 280; the printed seed is Eq. (A3), lines 238–240.

The current public `udist` documentation describes `ufl` and `spatial_ufl` as fluid momentum in proper-velocity units and gives relativistic momentum addition as the default [10]. This documents the input-language interpretation; it does not identify the authors' exact historical executable. For the archived diagnostics, the definition used here comes directly from the authors' Zenodo deposit description [2]:

$$\mathrm{vfl}_i = \frac{1}{n}\int f v_i \,\mathrm{d}^3 v, \qquad (v_i v_j)_{\mathrm{diagnostic}} = \int f v_i v_j \,\mathrm{d}^3 v.$$

The deck and diagnostic checks, rather than an assumed private code revision, support the initial-moment comparison.

The density-transport explanation. The main text, p. 5, attributes the one-electron-fluid density deficit to the following mechanism:

> "at the mid-plane $\overline{v}_y = 0$ by symmetry, blocking 1-e-Fluid transport into the sheet".

The complete argument is in original TeX line 191. The symmetry-plane continuity calculation assesses this causal explanation; the plotted density deficit remains empirical evidence from the reported benchmark.

The implemented closure and instability claim. On p. 4 the source specifies the finite local model

$$\partial_m q_{ijm} = \kappa v_{th}(p_{ij} - \delta_{ij}p), \qquad p = \mathrm{tr}(p_{ij})/3, \qquad \kappa = 0.1 d_e^{-1},$$

with an isotropization timescale of about $90\omega_{pe}^{-1}$ (original TeX line 179). Figure 3's caption, p. 4, explicitly omits its contribution from the displayed fluid heat flux and notes that only its divergence is constrained (line 166). The finite term is therefore a disclosed part of the model. The longitudinal calculation in this study tests that finite model, rather than silently replacing it by its zero-coefficient limit. On p. 5 the source also asserts that the two-electron-fluid model can capture linear growth because the instabilities are due to counter-streaming (line 207). The general importance of closure choice for instability fidelity is established prior context [7].

Accuracy metric and retained qualifications. The color in source Fig. 4, p. 5, is the best-time score

$$\max_t \left[1 - \frac{\langle |q_{xxx} - q_{CSxxx}| \rangle_{xy}}{\langle |q_{xxx}| \rangle_{xy}}\right]$$

(original TeX line 198). Its proposed thermal/drift scaling is explained in line 202. The present audit distinguishes this empirical best-time diagnostic from a uniform error guarantee; it does not

claim to have rerun the parameter scan. The source acknowledges later stream folding (p. 5, line 207), separate fluid color scales and amplitude differences in the full-tensor comparison (Fig. 5, p. 7, line 250), and kinetic free-expansion limitations (Appendix C, p. 8, line 293). The exact mixture identities in source Eqs. (3)–(4) and (B1)–(B2) are preserved as valid algebra.

B Pinned execution record

The reported calculations use the immutable code commit:

0d07de76f3d0b56e49e9ef2b9de931b8567ad0c8

This pins the numerical code and inputs; the final release also contains the manuscript source.

```
git clone https://github.com/Teylersf/counterstreaming-heatflux-audit.git
cd counterstreaming-heatflux-audit
git checkout 0d07de76f3d0b56e49e9ef2b9de931b8567ad0c8
python -m pip install -r requirements.txt
python experiments/fetch_data.py
python experiments/data_checks.py
python -X utf8 experiments/verify.py
python -X utf8 experiments/make_figures.py
```

This is an author-environment execution record, not third-party certification or independent human peer review.

```
Started UTC: 2026-10-28T23:09:06.987425+00:00
Python: 3.12.14 (main, Aug 25 2026, 14:01:42) [MSC v.1944 64 bit (AMD64)]
NumPy: 2.3.5
h5py: 3.16.0
PASS: initial_current
PASS: initial_pressure
PASS: mixture_identities
PASS: continuity_symmetry
PASS: conditioning_counterexample
PASS: singlefluid_nonidentifiability
PASS: finite_relaxation_stability
PASS: isotropization_entropy
All assertions passed: true
Exit status: 0
```

File	SHA-256
results/verification.json	6e724e564d4ca23bf718d4add4eded9311aa6ebbeb161c4abc606ec750f01491
experiments/verify.py	23c64243de9e58d493fc87b18b4c47edf917fb67d11853898009e6e8824903e6
experiments/closure_checks.py	797fa1a2c85b36e09531fcff343ebed773674ecfe5a1caf8bd6518826ed3efbb
experiments/data_checks.py	fb2253e5f1970e80baea7a5349a8ac0cbf0e847b0cfe3f04eadb99b1e32e064b
experiments/fetch_data.py	9739bc225cda81134c984edb9c06a39a36a5c24403d6b67564e6de2d063a6fc0
experiments/profile_sensitivity.py	16f79b0c6c4aad933b2efedf8516dbca609a53993442bcccd78cd16be5c15e7e
experiments/revision_data.py	ad9f47607fecb68eb3a914783e3d88af8cef1e05c5ed557784286fea901db0db
experiments/thermal_loading.py	8ca81418bba26d8307337d7d0f1612bcf169186f91a00b1c6e84b5ea5748511d
results/profile-sensitivity.json	f74e9bc750cb661cda931045f7b471cb29271d525c15e08504b54f618c4de586
results/revision-data.json	0abafde815b680b75af3fa01efcfee8cf20aa1e140325199ce8fb9cc22dd2066
results/thermal-loading.json	f7c6c4918020ea9d7278f2c3bd299850f2869bf30cd29694f7b7075478a91161
data/provenance-revision.json	25b9abe520ffee744c3881759e60a3f2e88f4ca63834eec6c619e9804af5f3f9

The full log, input provenance, derived profiles and strict verification JSON are included in the repository.

Revision analyses. The original eight-group execution record is retained, with its four code files byte-identical. The pinned commit also contains the separate full-profile sensitivity, archived-fluid corroboration and conditional thermal-loading calculations.

```
python experiments/profile_sensitivity.py
python experiments/revision_data.py --inventory --fetch
# Decode BP in a separate Linux ADIOS2 environment:
python experiments/revision_data.py --decode-bp
python experiments/revision_data.py --compare
python experiments/thermal_loading.py
```

The BP reader used ADIOS2 2.13.0.1001 and NumPy 2.3.5 under Linux Python 3.12.3. Portable reader setup and exact member/export hashes are provided in the data documentation. No native deposited-current comparison, full-archive checksum verification, matched-state reconnection rerun or sheet eigenanalysis is claimed.