

# Analysis I - Hw 1

Daniel Yu

September 23, 2024

---

## 1 Problem 1

Let  $(X, \rho)$  be a metric space and  $E$ , a non-empty subset in  $X$ . Consider the new metric space,  $(E, \rho)$ . Prove that a set  $U \subseteq E$  is open in  $E \Leftrightarrow \exists$  an open set  $\tilde{U}$  in  $X$  such that  $U = \tilde{U} \cap E$ . Similarly, prove this for closed sets.

*Proof.*  $\rightarrow$  Assume that there is a set  $U \subseteq E$  that is open in  $(E, \rho)$  and  $E \subseteq X$  where  $X$  forms a metric space  $(X, \rho)$ . If the open set  $U$  is also open in  $(X, \rho)$  then  $\tilde{U} = U$  and we are done, so assume that  $U$  is not open in  $(X, \rho)$ . We can construct an open set  $\tilde{U}$  in  $(X, \rho)$  by taking the ball  $B_r(x)' \subseteq X$  that contains  $B_r(x) \subseteq U$ .  $B_r(x)' \supseteq B_r(x)$  since  $U \subseteq X$ . We can do this  $\forall x \in U$  with some  $r > 0$ . By this construction,  $\tilde{U}$  is an open set because for all  $x \in U$ ,  $\exists r > 0$  such that  $B_r(x)' \subseteq X$ . Then,  $B_r(x)' \cap E = B_r(x)$  since  $B_r(x)'$  contains  $B_r(x) \subseteq U \subseteq E$  and points in  $B_r(x)' \setminus E$  i.e.  $B_r(x)' = B_r(x) \cup B_r(x)' \setminus E$ . So,

$$(\cup_{x \in U} B_r(x)') \cap E = \cup_{x \in U} B_r(x).$$

and every  $x \in U$  has an open set  $B_r(x) \subseteq E$ , which is precisely the definition of the open set  $U$  in  $E$ !

$\leftarrow$  Assume that  $\exists$  an open set  $\tilde{U} \subseteq X$  such that  $U = \tilde{U} \cap E$ . By set theory,  $U \subseteq E$ . We know that  $\tilde{U}$  is open, so

$$\forall x \in \tilde{U}, \exists r > 0, \text{ such that } B_r(x) \subseteq \tilde{U}.$$

Consider the ball  $B_r(x) \cap E$ , the intersection of the ball  $B_r(x) \subseteq X$  with  $E$ . We know that this intersection must be open in  $E$  because we are taking an open set in  $X$  and intersecting it with  $E$  which is open with respect to itself. Since  $B_r(x) \subseteq \tilde{U}$ , then  $B_r(x) \cap E \subseteq \tilde{U} \cap E = U$ . So for every  $x \in U$ ,  $\exists r > 0$  such that an open set is formed in  $E$ :

$$B_r(x) \cap E \subseteq U.$$

$U \subseteq E$  is open in  $E$ .

Now to prove the same for closed sets. We just proved that a set  $U \subseteq E$  is open in  $E \Leftrightarrow \exists$  an open set  $\tilde{U}$  in  $X$  such that  $U = \tilde{U} \cap E$ . We know that the complement of an open set is a closed set. Thus, taking the complement, we know that there is a set  $C = E \setminus U$  where  $C \subseteq E$  that is closed in  $E$ . It follows that the open set  $\tilde{U}$  in  $X$  has a complement  $\tilde{C} = X \setminus \tilde{U}$  that is closed in  $X$ . Then

$$\tilde{C} \cap E = (X \setminus \tilde{U}) \cap E = (X \cap E) \setminus (\tilde{U} \cap E) = X \setminus U = C.$$

Thus, there exists a closed set  $C \subseteq E \Leftrightarrow \exists$  a closed set  $\tilde{C}$  in  $X$  such that  $C = \tilde{C} \cap E$ .  $\square$

**Note.** Intuition

Consider  $U \subseteq E$  an open set. This means that  $\forall x \in U \exists r > 0$  such that

$$B_r(x) \subseteq U.$$

If we consider  $U$  in  $X \supseteq E$ , then  $U$  may not be necessarily be open in  $X$  because there may  $x' \in X$  but  $x' \notin E$  such that  $\exists B_r(x) \forall r > 0$  such that  $x' \in B_r(x)$ . Since  $B_r(x) \subseteq U \subseteq E$ , if  $x' \in B_r(x)$ , then:

$$B_r(x) \not\subseteq U.$$

For example, let  $X = \mathbb{R}$ , the closed interval  $[0, 1)$  would not be open because there is no ball centered at  $B_r(0)$  of any radius greater than 0 that is a subset of  $[0, 1)$ . However, if we restrict  $X = [0, 1)$ , then  $[0, 1)$  becomes open since  $B_r(0) = [0, r) \subseteq [0, 1)$  when  $r < 1$  (now the  $x < 0$  don't exist). The idea is that we can find an open set  $\tilde{U}$  in  $X$  that is an analogue of  $U$  in  $E$ .

## 2 Problem 2

Given  $K \subseteq E$ , then prove  $K$  is compact in  $E \Leftrightarrow K$  is compact in  $X$ .

*Proof.*  $\rightarrow$  If  $K$  is compact in  $E \subseteq X$ , then for any open cover  $\{U_\alpha\}_{\alpha \in I}$  in  $E$  such that  $K \subseteq \cup_{\alpha \in I} U_\alpha$  that covers  $K$  and there is some finite subcover  $\{U_{\alpha_1}, U_{\alpha_2}, \dots, U_{\alpha_n}\} \subseteq E$  that covers  $K$ . Since  $E \subseteq X$ , we

---

can use the statement from problem 1. For each open subset  $U_{\alpha_i} \subseteq E$ , there exists open subset  $U_{\beta_i} \subseteq X$  such that  $U_{\alpha_i} = U_{\beta_i} \cap E \rightarrow U_{\alpha_i} \subseteq U_{\beta_i}$ . So,

$$K \subseteq \cup_{\alpha_i \in I_1} U_{\alpha_i} \subseteq \cup_{\beta_i \in I_1} U_{\beta_i}.$$

Thus, we can construct a finite subcover  $\{U_{\beta_1}, U_{\beta_2}, \dots, U_{\beta_n}\}$  of  $K$  in  $X$ . As any open cover of  $K$  in  $X$  can be restricted  $\{U_{\beta} \cap E\}_{\beta \in I}$  to be an open cover of  $K$  in  $E$  and any open cover of  $K$  in  $E$  can be augmented to be an open cover of  $K$  in  $X$ ,  $\{\{U_{\alpha}\}_{\alpha \in I}, X \setminus E\}$ , then any open cover in  $X$  can be mapped to some open cover in  $E$  such that we can follow the construction above to create a finite subcover of  $K$  in  $X$ . So,  $K$  is compact in  $X$ .

← Assume that  $K$  is compact in  $X$  and  $K \subseteq E$ . This means for any open cover of  $K$  in  $X$ :

$$\{U_{\alpha}\}_{\alpha \in I}.$$

there exists a finite subcover  $\{U_{\alpha_1}, \dots, U_{\alpha_n}\}$  such that

$$K \subseteq \cup_{\alpha \in I_1} U_{\alpha} \subseteq X.$$

Then since  $E \subseteq X$ , we can use the statement from problem 1. For each open subset  $U_{\alpha_i} \subseteq X$ , there exists open subset  $U_{\beta_i} \subseteq E$  such that  $U_{\beta_i} = U_{\alpha_i} \cap E \rightarrow U_{\beta_i} \subseteq E$ . And since,  $K \subseteq E$ ,

$$U_{\alpha_i} \cap K \subseteq U_{\alpha_i} \cap E = U_{\beta_i}.$$

so,

$$(\cup_{\alpha \in I_1} U_{\alpha_i}) \cap K = K \subseteq (\cup_{\alpha \in I_1} U_{\beta_i}) \cap E = \cup_{\beta \in I_1} U_{\beta_i}.$$

and  $\{U_{\beta_1}, \dots, U_{\beta_n}\}$  is an finite subcover of  $K$  in  $E$ . Then the open cover of  $K$  in  $E$  would just be  $\{U_{\beta}\}_{\beta \in I}$ . As any open cover of  $K$  in  $X$  can be restricted  $\{U_{\alpha} \cap E\}_{\alpha \in I}$  to be an open cover of  $K$  in  $E$  and any open cover of  $K$  in  $E$  can be augmented to be an open cover of  $K$  in  $X$ ,  $\{\{U_{\beta}\}_{\beta \in I}, X \setminus E\}$ , then for any open cover in  $E$  which can be mapped to some open cover in  $X$ , we can follow the construction above to create a finite subcover of  $K$  in  $E$  and  $K$  is compact.  $\square$

## Due October 2nd