



Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Numerical Analysis

Calculation of eigenvalues

by Csaba Gáspár

Széchenyi István University

2020, autumn semester



Motivations

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

It is well known that the eigenvalues of the matrix A are the solutions of the **characteristic equation**:

$$\det(A - \lambda I) = 0$$

However, in practice this can be hardly performed.

Numerical approximations are needed!

The simplest localization result is:

$$\rho(A) \leq \|A\|$$

with arbitrary induced matrix norm.



Motivations

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

It is well known that the eigenvalues of the matrix A are the solutions of the **characteristic equation**:

$$\det(A - \lambda I) = 0$$

However, in practice this can be hardly performed.

Numerical approximations are needed!

The simplest localization result is:

$$\rho(A) \leq \|A\|$$

with arbitrary induced matrix norm.



Motivations

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

It is well known that the eigenvalues of the matrix A are the solutions of the **characteristic equation**:

$$\det(A - \lambda I) = 0$$

However, in practice this can be hardly performed.

Numerical approximations are needed!

The simplest localization result is:

$$\rho(A) \leq \|A\|$$

with arbitrary induced matrix norm.



Motivations

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

It is well known that the eigenvalues of the matrix A are the solutions of the **characteristic equation**:

$$\det(A - \lambda I) = 0$$

However, in practice this can be hardly performed.

Numerical approximations are needed!

The simplest localization result is:

$$\rho(A) \leq \|A\|$$

with arbitrary induced matrix norm.

Bounds for the eigenvalues by Gershgorin circles

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a square matrix (with possibly complex entries).

Define the numbers $r_k := \sum_{j \neq k} |a_{kj}|$ ($k = 1, 2, \dots, N$), and denote by B_k the closed disk of the complex plane centered at a_{kk} with radius r_k (**Gershgorin disks**).

Gershgorin's theorem

All the eigenvalues of the matrix A are located in the union of the Gershgorin disks.

Bounds for the eigenvalues by Gershgorin circles

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a square matrix (with possibly complex entries).

Define the numbers $r_k := \sum_{j \neq k} |a_{kj}|$ ($k = 1, 2, \dots, N$), and denote by B_k the closed disk of the complex plane centered at a_{kk} with radius r_k (**Gershgorin disks**).

Gershgorin's theorem

All the eigenvalues of the matrix A are located in the union of the Gershgorin disks.



Bounds for the eigenvalues by Gershgorin circles

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a square matrix (with possibly complex entries).

Define the numbers $r_k := \sum_{j \neq k} |a_{kj}|$ ($k = 1, 2, \dots, N$), and denote by B_k the closed disk of the complex plane centered at a_{kk} with radius r_k (**Gershgorin disks**).

Gershgorin's theorem

All the eigenvalues of the matrix A are located in the union of the Gershgorin disks.



Bounds for the eigenvalues by Gershgorin circles

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a square matrix (with possibly complex entries).

Define the numbers $r_k := \sum_{j \neq k} |a_{kj}|$ ($k = 1, 2, \dots, N$), and denote by B_k the closed disk of the complex plane centered at a_{kk} with radius r_k (**Gershgorin disks**).

Gershgorin's theorem

All the eigenvalues of the matrix A are located in the union of the Gershgorin disks.



Normal matrices

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

The square matrix $A \in \mathbb{M}_{N \times N}$ is said to be a **normal matrix**, if $AA^* = A^*A$.

Obviously, every self-adjoint matrix is normal.

If $A \in \mathbb{M}_{N \times N}$ is a normal matrix, then A has a system of eigenvectors which form an **orthonormal system** (i.e. the Euclidean norm of any eigenvector is 1, and the different eigenvectors are orthogonal).



Normal matrices

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

The square matrix $A \in \mathbb{M}_{N \times N}$ is said to be a **normal matrix**, if $AA^* = A^*A$.

Obviously, every self-adjoint matrix is normal.

If $A \in \mathbb{M}_{N \times N}$ is a normal matrix, then A has a system of eigenvectors which form an **orthonormal system** (i.e. the Euclidean norm of any eigenvector is 1, and the different eigenvectors are orthogonal).



Normal matrices

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

The square matrix $A \in \mathbf{M}_{N \times N}$ is said to be a **normal matrix**, if $AA^* = A^*A$.

Obviously, every self-adjoint matrix is normal.

If $A \in \mathbf{M}_{N \times N}$ is a normal matrix, then A has a system of eigenvectors which form an **orthonormal system** (i.e. the Euclidean norm of any eigenvector is 1, and the different eigenvectors are orthogonal).



The power iteration

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal matrix with the eigenvalues:
 $|\lambda_1| \leq |\lambda_2| \leq \dots \leq |\lambda_{N-1}| < |\lambda_N|$, and denote by s_1, s_2, \dots, s_N
the orthonormal eigenvectors.

The power iteration

Let $x_0 \in \mathbf{R}^N$ be a vector such that $\langle x_0, s_N \rangle \neq 0$, and define:

$$x_{n+1} := Ax_n \quad (n = 0, 1, 2, \dots)$$

Then the sequence of **Rayleigh quotients** $\frac{\langle Ax_n, x_n \rangle}{\|x_n\|^2} = \frac{\langle x_{n+1}, x_n \rangle}{\|x_n\|^2}$
converges to λ_N . For the speed of convergence, we have the
following estimation:

$$\left| \frac{\langle Ax_n, x_n \rangle}{\|x_n\|^2} - \lambda_N \right| \leq \text{const.} \cdot \left| \frac{\lambda_{N-1}}{\lambda_N} \right|^{2n}$$



The inverse iteration

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal, regular matrix with the eigenvalues: $0 < |\lambda_1| < |\lambda_2| \leq |\lambda_3| \leq \dots \leq |\lambda_N|$, and denote by s_1, s_2, \dots, s_N the corresponding orthonormal eigenvectors.

The inverse iteration

Apply the power iteration to the matrix A^{-1} (its eigenvalues are $\frac{1}{\lambda_1}, \dots, \frac{1}{\lambda_N}$).

Let $x_0 \in \mathbf{R}^N$ be a vector such that $\langle x_0, s_1 \rangle \neq 0$, and define:

$$x_{n+1} := A^{-1}x_n \quad (n = 0, 1, 2, \dots)$$

Then the sequence of the **Rayleigh quotients**

$$\frac{\langle A^{-1}x_n, x_n \rangle}{\|x_n\|^2} = \frac{\langle x_{n+1}, x_n \rangle}{\|x_n\|^2} \text{ converges to } \frac{1}{\lambda_1}.$$

It is worth calculating the vector x_{n+1} by solving the equation $Ax_{n+1} = x_n$ and applying an *LU* decomposition.



The inverse iteration

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal, regular matrix with the eigenvalues: $0 < |\lambda_1| < |\lambda_2| \leq |\lambda_3| \leq \dots \leq |\lambda_N|$, and denote by s_1, s_2, \dots, s_N the corresponding orthonormal eigenvectors.

The inverse iteration

Apply the power iteration to the matrix A^{-1} (its eigenvalues are $\frac{1}{\lambda_1}, \dots, \frac{1}{\lambda_N}$).

Let $x_0 \in \mathbf{R}^N$ be a vector such that $\langle x_0, s_1 \rangle \neq 0$, and define:

$$x_{n+1} := A^{-1}x_n \quad (n = 0, 1, 2, \dots)$$

Then the sequence of the **Rayleigh quotients**

$$\frac{\langle A^{-1}x_n, x_n \rangle}{\|x_n\|^2} = \frac{\langle x_{n+1}, x_n \rangle}{\|x_n\|^2} \text{ converges to } \frac{1}{\lambda_1}.$$

It is worth calculating the vector x_{n+1} by solving the equation $Ax_{n+1} = x_n$ and applying an *LU* decomposition.



The inverse iteration

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal, regular matrix with the eigenvalues: $0 < |\lambda_1| < |\lambda_2| \leq |\lambda_3| \leq \dots \leq |\lambda_N|$, and denote by s_1, s_2, \dots, s_N the corresponding orthonormal eigenvectors.

The inverse iteration

Apply the power iteration to the matrix A^{-1} (its eigenvalues are $\frac{1}{\lambda_1}, \dots, \frac{1}{\lambda_N}$).

Let $x_0 \in \mathbf{R}^N$ be a vector such that $\langle x_0, s_1 \rangle \neq 0$, and define:

$$x_{n+1} := A^{-1}x_n \quad (n = 0, 1, 2, \dots)$$

Then the sequence of the **Rayleigh quotients**

$$\frac{\langle A^{-1}x_n, x_n \rangle}{\|x_n\|^2} = \frac{\langle x_{n+1}, x_n \rangle}{\|x_n\|^2} \text{ converges to } \frac{1}{\lambda_1}.$$

It is worth calculating the vector x_{n+1} by solving the equation $Ax_{n+1} = x_n$ and applying an *LU* decomposition.



Shifted inverse power method

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal matrix with eigenvalues: $\lambda_1, \lambda_2, \dots, \lambda_N$, and denote by s_1, s_2, \dots, s_N the corresponding orthonormal eigenvectors.

Assume that λ_k is a single eigenvalue, and λ_0 is a sufficiently good approximation of λ_k , such that the nearest eigenvalue to λ_0 is λ_k , i.e. for every $j \neq k$, the inequality $|\lambda_j - \lambda_0| > |\lambda_k - \lambda_0|$ holds.

Then the number $\lambda_k - \lambda_0$ is an eigenvalue of the matrix $(A - \lambda_0 I)$ with minimal absolute value, so that the inverse iteration is applicable.

Shifted inverse power method

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal matrix with eigenvalues: $\lambda_1, \lambda_2, \dots, \lambda_N$, and denote by s_1, s_2, \dots, s_N the corresponding orthonormal eigenvectors.

Assume that λ_k is a single eigenvalue, and λ_0 is a sufficiently good approximation of λ_k , such that the nearest eigenvalue to λ_0 is λ_k , i.e. for every $j \neq k$, the inequality $|\lambda_j - \lambda_0| > |\lambda_k - \lambda_0|$ holds.

Then the number $\lambda_k - \lambda_0$ is an eigenvalue of the matrix $(A - \lambda_0 I)$ with minimal absolute value, so that the inverse iteration is applicable.



Shifted inverse power method

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a normal matrix with eigenvalues: $\lambda_1, \lambda_2, \dots, \lambda_N$, and denote by s_1, s_2, \dots, s_N the corresponding orthonormal eigenvectors.

Assume that λ_k is a single eigenvalue, and λ_0 is a sufficiently good approximation of λ_k , such that the nearest eigenvalue to λ_0 is λ_k , i.e. for every $j \neq k$, the inequality $|\lambda_j - \lambda_0| > |\lambda_k - \lambda_0|$ holds.

Then the number $\lambda_k - \lambda_0$ is an eigenvalue of the matrix $(A - \lambda_0 I)$ with minimal absolute value, so that the inverse iteration is applicable.

Shifted inverse power method

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Shifted inverse iteration

Let $x_0 \in \mathbf{R}^N$ be a vector such that $\langle x_0, s_k \rangle \neq 0$, and define:

$$x_{n+1} := (A - \lambda_0 I)^{-1} x_n \quad (n = 0, 1, 2, \dots)$$

Then the sequence of **Rayleigh quotients** $\frac{\langle x_{n+1}, x_n \rangle}{\|x_n\|^2}$ converges to $\frac{1}{\lambda_k - \lambda_0}$.

Determination of all eigenvalues

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Jacobi's method: Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint matrix. Denote by $A_0 := A$. Define the pair of indices (p, q) ($p < q$), for which $|A_{pq}|$ is maximal above the main diagonal. Define

$$\cot 2t := \frac{A_{qq} - A_{pp}}{2A_{pq}},$$

and define the (orthogonal) matrix Q_n as follows:

$$Q_n := \begin{pmatrix} 1 & & & & & & \\ & 1 & & & & & \\ & & \cos t & \dots & \dots & \dots & \sin t \\ & & \dots & 1 & & & \dots \\ & & \dots & & \dots & & \dots \\ & & \dots & & & 1 & \dots \\ & & -\sin t & \dots & \dots & \dots & \cos t \\ & & & & & & 1 \\ & & & & & & & 1 \end{pmatrix}$$

Determination of all eigenvalues

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Method of Jacobi

Define the matrix sequence recursively:

$$A_{n+1} := Q_n^* A_n Q_n$$

If the eigenvalues of A are distinct, then the sequence of matrices (A_n) elementwise converges to a diagonal matrix, the diagonal entries of which are the eigenvalues of the matrix A .

Determination of all eigenvalues

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint positive definite matrix.

Method based of the Cholesky decomposition

Denote by $A_0 := A$, and for $n = 0, 1, 2, \dots$ define

$$A_{n+1} := L_n^* L_n,$$

where $L_n L_n^*$ is the Cholesky decomposition of A_n . Then the sequence of matrices (A_n) elementwise converges to a diagonal matrix, the diagonal entries of which are the eigenvalues of the matrix A .

Determination of all eigenvalues

Numerical
Analysis

by Csaba
Gáspár

Computation
of eigenvalues

Bounds for the
eigenvalues

Determination of
the eigenvalue
with maximal
absolute value

Determination of
the eigenvalue
with minimal
absolute value

Determination of
the intermediate
eigenvalues

Determination of
all eigenvalues

Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint positive definite matrix.

Method based of the QR decomposition

Denote by $A_0 := A$, and for $n = 0, 1, 2, \dots$ define

$$A_{n+1} := R_n Q_n,$$

where $Q_n R_n$ is the QR decomposition of A_n . Then the sequence of matrices (A_n) elementwise converges to a diagonal matrix, the diagonal entries of which are the eigenvalues of the matrix A .