



Numerical Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Numerical Analysis Introduction

by Csaba Gáspár

Széchenyi István University

2020, autumn semester



Motivation: ill-conditioned systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of equations:

$$1000x + 999y = 1$$

$$999x + 998y = 1$$

Solution: $x = 1$, $y = -1$.

$$1000x + 999y = 1$$

$$999x + 998y = 0.999$$

Solution: $x = 0.001$, $y = 0$.

Ill-conditioned systems of equations

Condition number of a self-adjoint matrix

$$\text{cond}(A) := \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Motivation: ill-conditioned systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of equations:

$$1000x + 999y = 1$$

$$999x + 998y = 1$$

Solution: $x = 1$, $y = -1$.

$$1000x + 999y = 1$$

$$999x + 998y = 0.999$$

Solution: $x = 0.001$, $y = 0$.

Ill-conditioned systems of equations

Condition number of a self-adjoint matrix

$$\text{cond}(A) := \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Motivation: ill-conditioned systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of equations:

$$1000x + 999y = 1$$

$$999x + 998y = 1$$

Solution: $x = 1$, $y = -1$.

$$1000x + 999y = 1$$

$$999x + 998y = 0.999$$

Solution: $x = 0.001$, $y = 0$.

Ill-conditioned systems of equations

Condition number of a self-adjoint matrix

$$\text{cond}(A) := \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Motivation: ill-conditioned systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of equations:

$$1000x + 999y = 1$$

$$999x + 998y = 1$$

Solution: $x = 1$, $y = -1$.

$$1000x + 999y = 1$$

$$999x + 998y = 0.999$$

Solution: $x = 0.001$, $y = 0$.

Ill-conditioned systems of equations

Condition number of a self-adjoint matrix

$$\text{cond}(A) := \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Motivation: ill-conditioned systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of equations:

$$1000x + 999y = 1$$

$$999x + 998y = 1$$

Solution: $x = 1$, $y = -1$.

$$1000x + 999y = 1$$

$$999x + 998y = 0.999$$

Solution: $x = 0.001$, $y = 0$.

Ill-conditioned systems of equations

Condition number of a self-adjoint matrix

$$\text{cond}(A) := \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Motivation: ill-conditioned systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of equations:

$$1000x + 999y = 1$$

$$999x + 998y = 1$$

Solution: $x = 1$, $y = -1$.

$$1000x + 999y = 1$$

$$999x + 998y = 0.999$$

Solution: $x = 0.001$, $y = 0$.

Ill-conditioned systems of equations

Condition number of a self-adjoint matrix

$$\text{cond}(A) := \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Motivation: computation of a determinant

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A \in \mathbf{M}_{200 \times 200}$ be an arbitrary matrix. What is the computational cost of the calculation of $\det(A)$ by definition (i.e. performing an expansion by minors)?

c_N := number of multiplications.

$$c_N = N \cdot c_{N-1}$$

$$c_{200} = 200 \cdot c_{199} = 200 \cdot 199 \cdot c_{198} = \dots = 200!$$

(impossible to perform)

Motivation: computation of a determinant

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A \in \mathbf{M}_{200 \times 200}$ be an arbitrary matrix. What is the computational cost of the calculation of $\det(A)$ by definition (i.e. performing an expansion by minors)?

$c_N :=$ number of multiplications.

$$c_N = N \cdot c_{N-1}$$

$$c_{200} = 200 \cdot c_{199} = 200 \cdot 199 \cdot c_{198} = \dots = 200!$$

(impossible to perform)



Motivation: computation of a determinant

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A \in \mathbf{M}_{200 \times 200}$ be an arbitrary matrix. What is the computational cost of the calculation of $\det(A)$ by definition (i.e. performing an expansion by minors)?

$c_N :=$ number of multiplications.

$$c_N = N \cdot c_{N-1}$$

$$c_{200} = 200 \cdot c_{199} = 200 \cdot 199 \cdot c_{198} = \dots = 200!$$

(impossible to perform)



Motivation: computation of a determinant

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A \in \mathbf{M}_{200 \times 200}$ be an arbitrary matrix. What is the computational cost of the calculation of $\det(A)$ by definition (i.e. performing an expansion by minors)?

$c_N :=$ number of multiplications.

$$c_N = N \cdot c_{N-1}$$

$$c_{200} = 200 \cdot c_{199} = 200 \cdot 199 \cdot c_{198} = \dots = 200!$$

(impossible to perform)



Vectors and matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

\mathbf{R}^N : the set of ordered N -tuples of real numbers.

\mathbf{R}^N is a vector space (with respect to the componentwise operations).

$\mathbf{M}_{N \times M}$: the set of N -by- M matrices.

$\mathbf{M}_{N \times M}$ is a vector space (with respect to the elementwise operations)



Vectors and matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

\mathbb{R}^N : the set of ordered N -tuples of real numbers.

\mathbb{R}^N is a vector space (with respect to the componentwise operations).

$\mathbb{M}_{N \times M}$: the set of N -by- M matrices.

$\mathbb{M}_{N \times M}$ is a vector space (with respect to the elementwise operations)



Vectors and matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

\mathbf{R}^N : the set of ordered N -tuples of real numbers.

\mathbf{R}^N is a vector space (with respect to the componentwise operations).

$\mathbf{M}_{N \times M}$: the set of N -by- M matrices.

$\mathbf{M}_{N \times M}$ is a vector space (with respect to the elementwise operations)



Vectors and matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

\mathbb{R}^N : the set of ordered N -tuples of real numbers.

\mathbb{R}^N is a vector space (with respect to the componentwise operations).

$\mathbb{M}_{N \times M}$: the set of N -by- M matrices.

$\mathbb{M}_{N \times M}$ is a vector space (with respect to the elementwise operations)



Matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Matrix multiplication

If $A \in \mathbf{M}_{N \times M}$, $B \in \mathbf{M}_{M \times K}$, then AB is defined by

$$(AB)_{kj} = \sum_{i=1}^M A_{ki} B_{ij} \quad (k = 1, \dots, N, \quad j = 1, \dots, K)$$

$AB \in \mathbf{M}_{N \times K}$. The multiplication is not commutative but associative.

- Unit matrix: I , diagonal elements = 1, off-diagonal elements = 0
- For arbitrary $A \in \mathbf{M}_{N \times N}$: $AI = IA = A$
- Inverse matrix of A : $A^{-1}A = I$



Matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Matrix multiplication

If $A \in \mathbf{M}_{N \times M}$, $B \in \mathbf{M}_{M \times K}$, then AB is defined by

$$(AB)_{kj} = \sum_{i=1}^M A_{ki} B_{ij} \quad (k = 1, \dots, N, \quad j = 1, \dots, K)$$

$AB \in \mathbf{M}_{N \times K}$. The multiplication is not commutative but associative.

- Unit matrix: I , diagonal elements = 1, off-diagonal elements = 0
- For arbitrary $A \in \mathbf{M}_{N \times N}$: $AI = IA = A$
- Inverse matrix of A : $A^{-1}A = I$



Matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Matrix multiplication

If $A \in \mathbf{M}_{N \times M}$, $B \in \mathbf{M}_{M \times K}$, then AB is defined by

$$(AB)_{kj} = \sum_{i=1}^M A_{ki} B_{ij} \quad (k = 1, \dots, N, \quad j = 1, \dots, K)$$

$AB \in \mathbf{M}_{N \times K}$. The multiplication is not commutative but associative.

- Unit matrix: I , diagonal elements = 1, off-diagonal elements = 0
- For arbitrary $A \in \mathbf{M}_{N \times N}$: $AI = IA = A$
- Inverse matrix of A : $A^{-1}A = I$



Matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

- Regular or invertible matrix: which has an inverse
- Adjoint matrix of A : $A_{kj}^* = A_{jk}$
- The matrix is called self-adjoint, if $A^* = A$

Eigenvalue equation: $As = \lambda s$ (λ : eigenvalue, s : eigenvector)

The eigenvalues satisfy the *characteristic equation*:

$$\det(A - \lambda I) = 0$$



Matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

- Regular or invertible matrix: which has an inverse
- Adjoint matrix of A : $A_{kj}^* = A_{jk}$
- The matrix is called self-adjoint, if $A^* = A$

Eigenvalue equation: $As = \lambda s$ (λ : eigenvalue, s : eigenvector)

The eigenvalues satisfy the *characteristic equation*:

$$\det(A - \lambda I) = 0$$



Matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

- Regular or invertible matrix: which has an inverse
- Adjoint matrix of A : $A_{kj}^* = A_{jk}$
- The matrix is called self-adjoint, if $A^* = A$

Eigenvalue equation: $As = \lambda s$ (λ : eigenvalue, s : eigenvector)

The eigenvalues satisfy the *characteristic equation*:

$$\det(A - \lambda I) = 0$$



Self-adjoint matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The self-adjoint matrix $A \in \mathbf{M}_{N \times N}$ is called

- *positive definite*, if $x^*Ax > 0$ for every vector $x \neq 0$;
- *positive semidefinite*, if $x^*Ax \geq 0$ for every vector x ;
- *negative definite*, if $x^*Ax < 0$ for every vector $x \neq 0$;
- *negative semidefinite*, if $x^*Ax \leq 0$ for every vector x ;
- *indefinite*, if x^*Ax takes both positive and negative values.



Self-adjoint matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The self-adjoint matrix $A \in \mathbf{M}_{N \times N}$ is

- positive definite, if all eigenvalues of A are positive;
- positive semidefinite, if all eigenvalues of A are nonnegative;
- negative definite, if all eigenvalues of A are negative;
- negative semidefinite, if all eigenvalues of A are nonpositive;
- indefinite, if A has both positive and negative eigenvalues.



Vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to define a vector (x_1, x_2, \dots, x_N) from \mathbf{R}^N to be 'great' or 'small'?

Let $x := (x_1, x_2, \dots, x_N) \in \mathbf{R}^N$ be an N -dimensional vector.
Then the 'greatness' of x can be characterized in several ways:



Vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to define a vector (x_1, x_2, \dots, x_N) from \mathbf{R}^N to be 'great' or 'small'?

Let $x := (x_1, x_2, \dots, x_N) \in \mathbf{R}^N$ be an N -dimensional vector. Then the 'greatness' of x can be characterized in several ways:

Maximum norm

$$\|x\|_{\max} := \max_{1 \leq k \leq N} |x_k|$$



Vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to define a vector (x_1, x_2, \dots, x_N) from \mathbf{R}^N to be 'great' or 'small'?

Let $x := (x_1, x_2, \dots, x_N) \in \mathbf{R}^N$ be an N -dimensional vector. Then the 'greatness' of x can be characterized in several ways:

Sum norm

$$\|x\|_1 := \sum_{k=1}^N |x_k|$$



Vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to define a vector (x_1, x_2, \dots, x_N) from \mathbf{R}^N to be 'great' or 'small'?

Let $x := (x_1, x_2, \dots, x_N) \in \mathbf{R}^N$ be an N -dimensional vector. Then the 'greatness' of x can be characterized in several ways:

Euclidean norm

$$\|x\|_2 := \|x\| := \sqrt{\sum_{k=1}^N |x_k|^2}$$

Properties of the vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations
Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product
Orthogonality
Matrix norms
Linear systems

For any vector norm:

$$||x|| \geq 0, \quad \text{and} \quad ||x|| = 0 \quad \text{if and only if} \quad x = (0, 0, \dots, 0)$$

$$||\alpha \cdot x|| = |\alpha| \cdot ||x|| \quad \text{for arbitrary } \alpha \in \mathbf{R}$$

$$||x + y|| \leq ||x|| + ||y|| \quad (\text{triangle inequality})$$

For the Euclidean norm: this is a consequence of the *Cauchy inequality* (see later).

Properties of the vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

For any vector norm:

$$\|x\| \geq 0, \quad \text{and} \quad \|x\| = 0 \quad \text{if and only if} \quad x = (0, 0, \dots, 0)$$

$$\|\alpha \cdot x\| = |\alpha| \cdot \|x\| \quad \text{for arbitrary } \alpha \in \mathbf{R}$$

$$\|x + y\| \leq \|x\| + \|y\| \quad (\text{triangle inequality})$$

For the Euclidean norm: this is a consequence of the *Cauchy inequality* (see later).

Properties of the vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

For any vector norm:

$$\|x\| \geq 0, \quad \text{and} \quad \|x\| = 0 \quad \text{if and only if} \quad x = (0, 0, \dots, 0)$$

$$\|\alpha \cdot x\| = |\alpha| \cdot \|x\| \quad \text{for arbitrary } \alpha \in \mathbf{R}$$

$$\|x + y\| \leq \|x\| + \|y\| \quad (\text{triangle inequality})$$

For the Euclidean norm: this is a consequence of the *Cauchy inequality* (see later).

Properties of the vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations
Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product
Orthogonality
Matrix norms
Linear systems

For any vector norm:

$$\|x\| \geq 0, \quad \text{and} \quad \|x\| = 0 \quad \text{if and only if} \quad x = (0, 0, \dots, 0)$$

$$\|\alpha \cdot x\| = |\alpha| \cdot \|x\| \quad \text{for arbitrary } \alpha \in \mathbf{R}$$

$$\|x + y\| \leq \|x\| + \|y\| \quad (\text{triangle inequality})$$

For the Euclidean norm: this is a consequence of the *Cauchy inequality* (see later).



Distance of vectors

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations
Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product
Orthogonality
Matrix norms
Linear systems

How to measure that two vectors are 'near' or 'far'?

Distance of vectors

The number $\|x - y\|$ is called the **distance of the vectors** x and y , where $\|\cdot\|$ is an arbitrary vector norm.



Distance of vectors

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to measure that two vectors are 'near' or 'far'?

Distance of vectors

The number $\|x - y\|$ is called the **distance of the vectors** x and y , where $\|\cdot\|$ is an arbitrary vector norm.

Limit of a vector sequence

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations
Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product
Orthogonality
Matrix norms
Linear systems

Limit of a vector sequence

The sequence of vectors $x^{(1)}, x^{(2)}, x^{(3)}, \dots$ is called to be **convergent**, and to tend to the vector x , if the distances of the vectors $x^{(n)}$ and x converges to 0, i.e.:

$$\|x^{(n)} - x\| \rightarrow 0 \quad (n \rightarrow +\infty),$$

where $\|\cdot\|$ is a given vector norm.

If the sequence of vectors $x^{(n)}$ from \mathbf{R}^N is convergent with respect to a vector norm, then the convergence remains valid with respect to any other vector norm.

Limit of a vector sequence

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Limit of a vector sequence

The sequence of vectors $x^{(1)}, x^{(2)}, x^{(3)}, \dots$ is called to be **convergent**, and to tend to the vector x , if the distances of the vectors $x^{(n)}$ and x converges to 0, i.e.:

$$\|x^{(n)} - x\| \rightarrow 0 \quad (n \rightarrow +\infty),$$

where $\|\cdot\|$ is a given vector norm.

If the sequence of vectors $x^{(n)}$ from \mathbf{R}^N is convergent with respect to a vector norm, then the convergence remains valid with respect to any other vector norm.

Inner product of vectors

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Inner product

Let $x = (x_1, x_2, \dots, x_N)$, $y = (y_1, y_2, \dots, y_N) \in \mathbf{R}^N$ be N -dimensional vectors. The following number is called their **inner product** (scalar product):

$$\langle x, y \rangle := \sum_{k=1}^N x_k \cdot y_k$$

The inner product generates the Euclidean norm in the following sense:

$$\|x\| = \sqrt{\langle x, x \rangle}$$



Inner product of vectors

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Inner product

Let $x = (x_1, x_2, \dots, x_N)$, $y = (y_1, y_2, \dots, y_N) \in \mathbf{R}^N$ be N -dimensional vectors. The following number is called their **inner product** (scalar product):

$$\langle x, y \rangle := \sum_{k=1}^N x_k \cdot y_k$$

The inner product generates the Euclidean norm in the following sense:

$$||x|| = \sqrt{\langle x, x \rangle}$$

Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $x, y, z \in \mathbf{R}^N$ be N -dimensional vectors, and let $\alpha \in \mathbf{R}$ be an arbitrary number. Then:

$$\langle x, x \rangle \geq 0, \text{ and } \langle x, x \rangle = 0 \text{ if and only if } x = (0, 0, \dots, 0)$$

$$\langle x, y \rangle = \langle y, x \rangle$$

$$\langle \alpha x, y \rangle = \alpha \cdot \langle x, y \rangle$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $x, y, z \in \mathbf{R}^N$ be N -dimensional vectors, and let $\alpha \in \mathbf{R}$ be an arbitrary number. Then:

$$\langle x, x \rangle \geq 0, \text{ and } \langle x, x \rangle = 0 \text{ if and only if } x = (0, 0, \dots, 0)$$

$$\langle x, y \rangle = \langle y, x \rangle$$

$$\langle \alpha x, y \rangle = \alpha \cdot \langle x, y \rangle$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$$

Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $x, y, z \in \mathbf{R}^N$ be N -dimensional vectors, and let $\alpha \in \mathbf{R}$ be an arbitrary number. Then:

$$\langle x, x \rangle \geq 0, \text{ and } \langle x, x \rangle = 0 \text{ if and only if } x = (0, 0, \dots, 0)$$

$$\langle x, y \rangle = \langle y, x \rangle$$

$$\langle \alpha x, y \rangle = \alpha \cdot \langle x, y \rangle$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $x, y, z \in \mathbf{R}^N$ be N -dimensional vectors, and let $\alpha \in \mathbf{R}$ be an arbitrary number. Then:

$$\langle x, x \rangle \geq 0, \text{ and } \langle x, x \rangle = 0 \text{ if and only if } x = (0, 0, \dots, 0)$$

$$\langle x, y \rangle = \langle y, x \rangle$$

$$\langle \alpha x, y \rangle = \alpha \cdot \langle x, y \rangle$$

$$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Useful equalities:

$$||x + y||^2 = ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$||x - y||^2 = ||x||^2 - 2\langle x, y \rangle + ||y||^2$$

$$\langle Ax, y \rangle = \langle x, A^*y \rangle$$

Cauchy inequality

$$|\langle x, y \rangle| \leq ||x|| \cdot ||y||$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Useful equalities:

$$||x + y||^2 = ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$||x - y||^2 = ||x||^2 - 2\langle x, y \rangle + ||y||^2$$

$$\langle Ax, y \rangle = \langle x, A^*y \rangle$$

Cauchy inequality

$$|\langle x, y \rangle| \leq ||x|| \cdot ||y||$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Useful equalities:

$$||x + y||^2 = ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$||x - y||^2 = ||x||^2 - 2\langle x, y \rangle + ||y||^2$$

$$\langle Ax, y \rangle = \langle x, A^*y \rangle$$

Cauchy inequality

$$|\langle x, y \rangle| \leq ||x|| \cdot ||y||$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Useful equalities:

$$||x + y||^2 = ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$||x - y||^2 = ||x||^2 - 2\langle x, y \rangle + ||y||^2$$

$$\langle Ax, y \rangle = \langle x, A^*y \rangle$$

Cauchy inequality

$$|\langle x, y \rangle| \leq ||x|| \cdot ||y||$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the Cauchy inequality: For arbitrary scalar $\alpha \in \mathbb{R}$, obviously $\|x - \alpha y\|^2 \geq 0$, therefore:

$$\|x\|^2 - 2\alpha\langle x, y \rangle + \alpha^2\|y\|^2 \geq 0$$

Define α by $\alpha := \frac{\|x\|}{\|y\|}$, then:

$$\|x\|^2 - 2\frac{\|x\|}{\|y\|}\langle x, y \rangle + \frac{\|x\|^2}{\|y\|^2} \cdot \|y\|^2 \geq 0.$$

whence

$$\langle x, y \rangle \leq \|x\| \cdot \|y\|$$

Substituting y by $(-y)$:

$$\langle x, -y \rangle \leq \|x\| \cdot \|-y\| = \|x\| \cdot \|y\|,$$

which implies the inequality.



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the Cauchy inequality: For arbitrary scalar $\alpha \in \mathbb{R}$, obviously $\|x - \alpha y\|^2 \geq 0$, therefore:

$$\|x\|^2 - 2\alpha \langle x, y \rangle + \alpha^2 \|y\|^2 \geq 0$$

Define α by $\alpha := \frac{\|x\|}{\|y\|}$, then:

$$\|x\|^2 - 2 \frac{\|x\|}{\|y\|} \langle x, y \rangle + \frac{\|x\|^2}{\|y\|^2} \cdot \|y\|^2 \geq 0.$$

whence

$$\langle x, y \rangle \leq \|x\| \cdot \|y\|$$

Substituting y by $(-y)$:

$$\langle x, -y \rangle \leq \|x\| \cdot \|-y\| = \|x\| \cdot \|y\|,$$

which implies the inequality.



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the Cauchy inequality: For arbitrary scalar $\alpha \in \mathbb{R}$, obviously $\|x - \alpha y\|^2 \geq 0$, therefore:

$$\|x\|^2 - 2\alpha \langle x, y \rangle + \alpha^2 \|y\|^2 \geq 0$$

Define α by $\alpha := \frac{\|x\|}{\|y\|}$, then:

$$\|x\|^2 - 2 \frac{\|x\|}{\|y\|} \langle x, y \rangle + \frac{\|x\|^2}{\|y\|^2} \cdot \|y\|^2 \geq 0.$$

whence

$$\langle x, y \rangle \leq \|x\| \cdot \|y\|$$

Substituting y by $(-y)$:

$$\langle x, -y \rangle \leq \|x\| \cdot \|-y\| = \|x\| \cdot \|y\|,$$

which implies the inequality.



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the Cauchy inequality: For arbitrary scalar $\alpha \in \mathbb{R}$, obviously $\|x - \alpha y\|^2 \geq 0$, therefore:

$$\|x\|^2 - 2\alpha \langle x, y \rangle + \alpha^2 \|y\|^2 \geq 0$$

Define α by $\alpha := \frac{\|x\|}{\|y\|}$, then:

$$\|x\|^2 - 2 \frac{\|x\|}{\|y\|} \langle x, y \rangle + \frac{\|x\|^2}{\|y\|^2} \cdot \|y\|^2 \geq 0.$$

whence

$$\langle x, y \rangle \leq \|x\| \cdot \|y\|$$

Substituting y by $(-y)$:

$$\langle x, -y \rangle \leq \|x\| \cdot \|-y\| = \|x\| \cdot \|y\|,$$

which implies the inequality.



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the Cauchy inequality: For arbitrary scalar $\alpha \in \mathbb{R}$, obviously $\|x - \alpha y\|^2 \geq 0$, therefore:

$$\|x\|^2 - 2\alpha \langle x, y \rangle + \alpha^2 \|y\|^2 \geq 0$$

Define α by $\alpha := \frac{\|x\|}{\|y\|}$, then:

$$\|x\|^2 - 2 \frac{\|x\|}{\|y\|} \langle x, y \rangle + \frac{\|x\|^2}{\|y\|^2} \cdot \|y\|^2 \geq 0.$$

whence

$$\langle x, y \rangle \leq \|x\| \cdot \|y\|$$

Substituting y by $(-y)$:

$$\langle x, -y \rangle \leq \|x\| \cdot \|-y\| = \|x\| \cdot \|y\|,$$

which implies the inequality.



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the triangle inequality:

Let $x, y \in \mathbf{R}^N$ be arbitrary vectors, then:

$$||x + y||^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$$

$$= ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$\leq ||x||^2 + 2 \cdot ||x|| \cdot ||y|| + ||y||^2 = (||x|| + ||y||)^2$$

Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the triangle inequality:

Let $x, y \in \mathbf{R}^N$ be arbitrary vectors, then:

$$||x + y||^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$$

$$= ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$\leq ||x||^2 + 2 \cdot ||x|| \cdot ||y|| + ||y||^2 = (||x|| + ||y||)^2$$

Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the triangle inequality:

Let $x, y \in \mathbf{R}^N$ be arbitrary vectors, then:

$$||x + y||^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$$

$$= ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$\leq ||x||^2 + 2 \cdot ||x|| \cdot ||y|| + ||y||^2 = (||x|| + ||y||)^2$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the triangle inequality:

Let $x, y \in \mathbf{R}^N$ be arbitrary vectors, then:

$$||x + y||^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$$

$$= ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$\leq ||x||^2 + 2 \cdot ||x|| \cdot ||y|| + ||y||^2 = (||x|| + ||y||)^2$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the triangle inequality:

Let $x, y \in \mathbf{R}^N$ be arbitrary vectors, then:

$$||x + y||^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$$

$$= ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$\leq ||x||^2 + 2 \cdot ||x|| \cdot ||y|| + ||y||^2 = (||x|| + ||y||)^2$$



Properties of the inner product

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Proof of the triangle inequality:

Let $x, y \in \mathbf{R}^N$ be arbitrary vectors, then:

$$||x + y||^2 = \langle x + y, x + y \rangle = \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$$

$$= ||x||^2 + 2\langle x, y \rangle + ||y||^2$$

$$\leq ||x||^2 + 2 \cdot ||x|| \cdot ||y|| + ||y||^2 = (||x|| + ||y||)^2$$



Orthogonality

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $x, y \in \mathbf{R}^N$ be N -dimensional vectors.

Orthogonality

The N -dimensional vectors $x, y \in \mathbf{R}^N$ are said to be **orthogonal**, if $\langle x, y \rangle = 0$.

Pythagoras's theorem

Let $x^{(1)}, x^{(2)}, \dots, x^{(m)} \in \mathbf{R}^N$ be pairwise orthogonal vectors, then:

$$\left\| \sum_{j=1}^m x^{(j)} \right\|^2 = \sum_{j=1}^m \|x^{(j)}\|^2.$$

Orthogonality

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $x, y \in \mathbf{R}^N$ be N -dimensional vectors.

Orthogonality

The N -dimensional vectors $x, y \in \mathbf{R}^N$ are said to be **orthogonal**, if $\langle x, y \rangle = 0$.

Pythagoras's theorem

Let $x^{(1)}, x^{(2)}, \dots, x^{(m)} \in \mathbf{R}^N$ be pairwise orthogonal vectors, then:

$$\left\| \sum_{j=1}^m x^{(j)} \right\|^2 = \sum_{j=1}^m \|x^{(j)}\|^2.$$

The Gram-Schmidt orthogonalization

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $a^{(1)}, a^{(2)}, \dots, a^{(N)} \in \mathbf{R}^N$ be linearly independent vectors. Define $\tilde{e}^{(1)} := a^{(1)}$, $e^{(1)} := \frac{\tilde{e}^{(1)}}{\|\tilde{e}^{(1)}\|}$, and for $k = 2, 3, \dots, N$:

$$\tilde{e}^{(k)} := a^{(k)} - \sum_{j=1}^{k-1} \langle a^{(k)}, e^{(j)} \rangle \cdot e^{(j)}, \quad e^{(k)} := \frac{\tilde{e}^{(k)}}{\|\tilde{e}^{(k)}\|}$$

Gram-Schmidt orthogonalization

The vectors $e^{(1)}, e^{(2)}, \dots, e^{(N)}$ form an **orthonormal vector system** (i.e. $\langle e^{(k)}, e^{(j)} \rangle = 0$, if $k \neq j$, and $\|e^{(k)}\| = 1$), and the vector systems $a^{(1)}, a^{(2)}, \dots, a^{(k)}$ and $e^{(1)}, e^{(2)}, \dots, e^{(k)}$ generate the same subspace of \mathbf{R}^N for all $k = 1, 2, \dots, N$.



The Gram-Schmidt orthogonalization

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $a^{(1)}, a^{(2)}, \dots, a^{(N)} \in \mathbf{R}^N$ be linearly independent vectors.
Define $\tilde{e}^{(1)} := a^{(1)}$, $e^{(1)} := \frac{\tilde{e}^{(1)}}{\|\tilde{e}^{(1)}\|}$, and for $k = 2, 3, \dots, N$:

$$\tilde{e}^{(k)} := a^{(k)} - \sum_{j=1}^{k-1} \langle a^{(k)}, e^{(j)} \rangle \cdot e^{(j)}, \quad e^{(k)} := \frac{\tilde{e}^{(k)}}{\|\tilde{e}^{(k)}\|}$$

Gram-Schmidt orthogonalization

The vectors $e^{(1)}, e^{(2)}, \dots, e^{(N)}$ form an **orthonormal vector system** (i.e. $\langle e^{(k)}, e^{(j)} \rangle = 0$, if $k \neq j$, and $\|e^{(k)}\| = 1$), and the vector systems $a^{(1)}, a^{(2)}, \dots, a^{(k)}$ and $e^{(1)}, e^{(2)}, \dots, e^{(k)}$ generate the same subspace of \mathbf{R}^N for all $k = 1, 2, \dots, N$.

Orthogonal projection

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $X_0 \subset \mathbf{R}^N$ be a subspace spanned by the orthonormal system $e^{(1)}, e^{(2)}, \dots, e^{(M)}$. For arbitrary vector $x \in \mathbf{R}^N$:

$$x^{(0)} := \sum_{j=1}^M \langle x, e^{(j)} \rangle \cdot e^{(j)}$$

is the *orthogonal projection of x to X_0* , i.e. $(x - x_0)$ is orthogonal to X_0 .

Proof. Complete the orthonormal system by the vectors $e^{(M+1)}, \dots, e^{(N)}$ to an orthonormal basis. Then $x = \sum_{j=1}^N \langle x, e^{(j)} \rangle \cdot e^{(j)}$, therefore for each

$$k = 1, 2, \dots, M: \quad \langle x - x^{(0)}, e^{(k)} \rangle = \sum_{j=M+1}^N \langle x, e^{(j)} \rangle \cdot \langle e^{(j)}, e^{(k)} \rangle = 0.$$

Orthogonal projection

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $X_0 \subset \mathbf{R}^N$ be a subspace spanned by the orthonormal system $e^{(1)}, e^{(2)}, \dots, e^{(M)}$. For arbitrary vector $x \in \mathbf{R}^N$:

$$x^{(0)} := \sum_{j=1}^M \langle x, e^{(j)} \rangle \cdot e^{(j)}$$

is the *orthogonal projection* of x to X_0 , i.e. $(x - x_0)$ is orthogonal to X_0 .

Proof. Complete the orthonormal system by the vectors $e^{(M+1)}, \dots, e^{(N)}$ to an orthonormal basis. Then $x = \sum_{j=1}^N \langle x, e^{(j)} \rangle \cdot e^{(j)}$, therefore for each

$$k = 1, 2, \dots, M: \quad \langle x - x^{(0)}, e^{(k)} \rangle = \sum_{j=M+1}^N \langle x, e^{(j)} \rangle \cdot \langle e^{(j)}, e^{(k)} \rangle = 0.$$

Orthogonal projection

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $X_0 \subset \mathbf{R}^N$ be a subspace spanned by the orthonormal system $e^{(1)}, e^{(2)}, \dots, e^{(M)}$. For arbitrary vector $x \in \mathbf{R}^N$:

$$x^{(0)} := \sum_{j=1}^M \langle x, e^{(j)} \rangle \cdot e^{(j)}$$

is the *orthogonal projection of x to X_0* , i.e. $(x - x_0)$ is orthogonal to X_0 .

Proof. Complete the orthonormal system by the vectors $e^{(M+1)}, \dots, e^{(N)}$ to an orthonormal basis. Then $x = \sum_{j=1}^N \langle x, e^{(j)} \rangle \cdot e^{(j)}$, therefore for each

$$k = 1, 2, \dots, M: \quad \langle x - x^{(0)}, e^{(k)} \rangle = \sum_{j=M+1}^N \langle x, e^{(j)} \rangle \cdot \langle e^{(j)}, e^{(k)} \rangle = 0.$$

Orthogonal matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbf{M}_{N \times N}$ is called **orthogonal**, if its column vectors form an orthonormal system in \mathbf{R}^N .

Example: The 2-by-2 matrix $A := \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$ is an orthogonal matrix.

The inverse of orthogonal matrices can be computed in an extremely simple way:

If the matrix $A \in \mathbf{M}_{N \times N}$ is orthogonal, then $A^{-1} = A^*$



Orthogonal matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbf{M}_{N \times N}$ is called **orthogonal**, if its column vectors form an orthonormal system in \mathbf{R}^N .

Example: The 2-by-2 matrix $A := \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$ is an orthogonal matrix.

The inverse of orthogonal matrices can be computed in an extremely simple way:

If the matrix $A \in \mathbf{M}_{N \times N}$ is orthogonal, then $A^{-1} = A^*$



Orthogonal matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbf{M}_{N \times N}$ is called **orthogonal**, if its column vectors form an orthonormal system in \mathbf{R}^N .

Example: The 2-by-2 matrix $A := \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$ is an orthogonal matrix.

The inverse of orthogonal matrices can be computed in an extremely simple way:

If the matrix $A \in \mathbf{M}_{N \times N}$ is orthogonal, then $A^{-1} = A^*$



Normal matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbb{M}_{N \times N}$ is called **normal**, if it is interchangeable with its adjoint, i.e. $AA^* = A^*A$.

Examples:

- 1) All self-adjoint matrices are normal.
- 2) All orthogonal matrices are normal.

Theorem

If the matrix $A \in \mathbb{M}_{N \times N}$ is normal, then A has a system of eigenvectors which form an orthonormal system. In this basis, A has a diagonal form.



Normal matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbb{M}_{N \times N}$ is called **normal**, if it is interchangeable with its adjoint, i.e. $AA^* = A^*A$.

Examples:

- 1) All self-adjoint matrices are normal.
- 2) All orthogonal matrices are normal.

Theorem

If the matrix $A \in \mathbb{M}_{N \times N}$ is normal, then A has a system of eigenvectors which form an orthonormal system. In this basis, A has a diagonal form.



Normal matrices

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbf{M}_{N \times N}$ is called **normal**, if it is interchangeable with its adjoint, i.e. $AA^* = A^*A$.

Examples:

- 1) All self-adjoint matrices are normal.
- 2) All orthogonal matrices are normal.

Theorem

If the matrix $A \in \mathbf{M}_{N \times N}$ is normal, then A has a system of eigenvectors which form an orthonormal system. In this basis, A has a diagonal form.



Matrix norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to measure that a matrix is 'great' or 'small'?

Let $A \in \mathbb{M}_{N \times M}$ a matrix of type $N \times M$. Then the 'greatness' of A can be characterized in several ways:



Matrix norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to measure that a matrix is 'great' or 'small'?

Let $A \in \mathbf{M}_{N \times M}$ a matrix of type $N \times M$. Then the 'greatness' of A can be characterized in several ways:

Maximum norm

$$\|A\| := \max\{|A_{kj}| : 1 \leq k \leq N, 1 \leq j \leq M\}$$



Matrix norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to measure that a matrix is 'great' or 'small'?

Let $A \in \mathbf{M}_{N \times M}$ a matrix of type $N \times M$. Then the 'greatness' of A can be characterized in several ways:

Sum norm

$$\|A\| := \sum_{k=1}^N \sum_{j=1}^M |A_{kj}|$$



Matrix norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to measure that a matrix is 'great' or 'small'?

Let $A \in \mathbf{M}_{N \times M}$ a matrix of type $N \times M$. Then the 'greatness' of A can be characterized in several ways:

Frobenius norm

$$\|A\| := \sqrt{\sum_{k=1}^N \sum_{j=1}^M |A_{kj}|^2}$$



Matrix norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

How to measure that a matrix is 'great' or 'small'?

Let $A \in \mathbf{M}_{N \times M}$ a matrix of type $N \times M$. Then the 'greatness' of A can be characterized in several ways:

These norms similar to the vector norms are rarely used. The use of the **matrix norm induced by a vector norm** is much more important.

Matrix norms induced by vector norms

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Denote by $||\cdot||$ a vector norm in \mathbf{R}^N , and $||\cdot||$ a vector norm in \mathbf{R}^M (possibly of another type). The matrix norm of $A \in \mathbf{M}_{N \times M}$ **induced by the above vector norms** is defined as follows:

$$||A|| := \max\{||Ax|| : x \in \mathbf{R}^M, ||x|| \leq 1\}$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

For every $x \in \mathbf{R}^M$: $\|Ax\| \leq \|A\| \cdot \|x\|$.

For arbitrary vector $0 \neq x \in \mathbf{R}^M$, the norm of the vector $\frac{x}{\|x\|}$ equals to 1, therefore:

$$\|Ax\| = \|A \frac{x}{\|x\|}\| \cdot \|x\| \leq \|A\| \cdot \|x\|.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

For every $x \in \mathbf{R}^M$: $\|Ax\| \leq \|A\| \cdot \|x\|$.

For arbitrary vector $0 \neq x \in \mathbf{R}^M$, the norm of the vector $\frac{x}{\|x\|}$ equals to 1, therefore:

$$\|Ax\| = \|A \frac{x}{\|x\|}\| \cdot \|x\| \leq \|A\| \cdot \|x\|.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

For every $x \in \mathbf{R}^M$: $\|Ax\| \leq \|A\| \cdot \|x\|$.

For arbitrary vector $0 \neq x \in \mathbf{R}^M$, the norm of the vector $\frac{x}{\|x\|}$ equals to 1, therefore:

$$\|Ax\| = \|A \frac{x}{\|x\|}\| \cdot \|x\| \leq \|A\| \cdot \|x\|.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

For every $x \in \mathbf{R}^M$: $\|Ax\| \leq \|A\| \cdot \|x\|$.

For arbitrary vector $0 \neq x \in \mathbf{R}^M$, the norm of the vector $\frac{x}{\|x\|}$ equals to 1, therefore:

$$\|Ax\| = \|A \frac{x}{\|x\|}\| \cdot \|x\| \leq \|A\| \cdot \|x\|.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, then $\|A\| \leq C$, i.e. $\|A\|$ is the least number with this property.

In particular, for any vector x , for which $\|x\| \leq 1$:

$$\|Ax\| \leq C \cdot \|x\| \leq C,$$

therefore the maximum of the numbers $\|Ax\|$ is at most C , implying:

$$\|A\| \leq C.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, then $\|A\| \leq C$, i.e. $\|A\|$ is the least number with this property.

In particular, for any vector x , for which $\|x\| \leq 1$:

$$\|Ax\| \leq C \cdot \|x\| \leq C,$$

therefore the maximum of the numbers $\|Ax\|$ is at most C , implying:

$$\|A\| \leq C.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, then $\|A\| \leq C$, i.e. $\|A\|$ is the least number with this property.

In particular, for any vector x , for which $\|x\| \leq 1$:

$$\|Ax\| \leq C \cdot \|x\| \leq C,$$

therefore the maximum of the numbers $\|Ax\|$ is at most C , implying:

$$\|A\| \leq C.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, then $\|A\| \leq C$, i.e. $\|A\|$ is the least number with this property.

In particular, for any vector x , for which $\|x\| \leq 1$:

$$\|Ax\| \leq C \cdot \|x\| \leq C,$$

therefore the maximum of the numbers $\|Ax\|$ is at most C , implying:

$$\|A\| \leq C.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, then $\|A\| \leq C$, i.e. $\|A\|$ is the least number with this property.

In particular, for any vector x , for which $\|x\| \leq 1$:

$$\|Ax\| \leq C \cdot \|x\| \leq C,$$

therefore the maximum of the numbers $\|Ax\|$ is at most C , implying:

$$\|A\| \leq C.$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, and there exists a vector $x \neq 0$, for which the equality is valid: $\|Ax\| = C \cdot \|x\|$, then $\|A\| = C$, (i.e. not only $\|A\| \leq C$).

We know already that $\|A\| \leq C$. Moreover, from the above vector x :

$$C \cdot \|x\| = \|Ax\| \leq \|A\| \cdot \|x\|.$$

whence $C \leq \|A\|$ is also valid.



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, and there exists a vector $x \neq 0$, for which the equality is valid: $\|Ax\| = C \cdot \|x\|$, then $\|A\| = C$, (i.e. not only $\|A\| \leq C$).

We know already that $\|A\| \leq C$. Moreover, from the above vector x :

$$C \cdot \|x\| = \|Ax\| \leq \|A\| \cdot \|x\|.$$

whence $C \leq \|A\|$ is also valid.



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, and there exists a vector $x \neq 0$, for which the equality is valid: $\|Ax\| = C \cdot \|x\|$, then $\|A\| = C$, (i.e. not only $\|A\| \leq C$).

We know already that $\|A\| \leq C$. Moreover, from the above vector x :

$$C \cdot \|x\| = \|Ax\| \leq \|A\| \cdot \|x\|.$$

whence $C \leq \|A\|$ is also valid.



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

If the number $C \geq 0$ is such that for any vector $x \in \mathbf{R}^M$, the following estimation holds: $\|Ax\| \leq C \cdot \|x\|$, and there exists a vector $x \neq \mathbf{0}$, for which the equality is valid: $\|Ax\| = C \cdot \|x\|$, then $\|A\| = C$, (i.e. not only $\|A\| \leq C$).

We know already that $\|A\| \leq C$. Moreover, from the above vector x :

$$C \cdot \|x\| = \|Ax\| \leq \|A\| \cdot \|x\|.$$

whence $C \leq \|A\|$ is also valid.

Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The norm of the zero matrix is always 0, while the norm of the unit matrix is always 1 (with respect to arbitrary vector norm).

Consequently, if $N > 1$, then in $\mathbf{M}_{N \times N}$, neither the maximum norm nor the sum norm nor the Frobenius norm can be induced by any vector norm.

$$\|A\| \geq 0, \quad \text{and} \quad \|A\| = 0 \quad \text{if and only if} \quad A = 0$$

$$\|\alpha \cdot A\| = |\alpha| \cdot \|A\| \quad \text{with arbitrary number } \alpha \in \mathbf{R}$$

$$\|A + B\| \leq \|A\| + \|B\| \quad (\text{triangle inequality})$$

$$\|A \cdot B\| \leq \|A\| \cdot \|B\|$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The norm of the zero matrix is always 0, while the norm of the unit matrix is always 1 (with respect to arbitrary vector norm).

Consequently, if $N > 1$, then in $\mathbf{M}_{N \times N}$, neither the maximum norm nor the sum norm nor the Frobenius norm can be induced by any vector norm.

$$\|A\| \geq 0, \quad \text{and} \quad \|A\| = 0 \quad \text{if and only if} \quad A = 0$$

$$\|\alpha \cdot A\| = |\alpha| \cdot \|A\| \quad \text{with arbitrary number } \alpha \in \mathbb{R}$$

$$\|A + B\| \leq \|A\| + \|B\| \quad (\text{triangle inequality})$$

$$\|A \cdot B\| \leq \|A\| \cdot \|B\|$$



Induced matrix norms, basic features

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The norm of the zero matrix is always 0, while the norm of the unit matrix is always 1 (with respect to arbitrary vector norm).

Consequently, if $N > 1$, then in $\mathbf{M}_{N \times N}$, neither the maximum norm nor the sum norm nor the Frobenius norm can be induced by any vector norm.

$$\|A\| \geq 0, \quad \text{and} \quad \|A\| = 0 \quad \text{if and only if} \quad A = 0$$

$$\|\alpha \cdot A\| = |\alpha| \cdot \|A\| \quad \text{with arbitrary number } \alpha \in \mathbf{R}$$

$$\|A + B\| \leq \|A\| + \|B\| \quad (\text{triangle inequality})$$

$$\|A \cdot B\| \leq \|A\| \cdot \|B\|$$

Induced matrix norms, examples

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Row norm

Suppose that both in \mathbf{R}^N and in \mathbf{R}^M , the maximum norm is defined. Then

$$\|A\| = \max_{1 \leq k \leq N} \sum_{j=1}^M |A_{kj}|$$

Column norm

Suppose that both in \mathbf{R}^N and in \mathbf{R}^M , the sum norm is given. Then

$$\|A\| = \max_{1 \leq j \leq M} \sum_{k=1}^N |A_{kj}|$$

Induced matrix norms, examples

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Row norm

Suppose that both in \mathbf{R}^N and in \mathbf{R}^M , the maximum norm is defined. Then

$$\|A\| = \max_{1 \leq k \leq N} \sum_{j=1}^M |A_{kj}|$$

Column norm

Suppose that both in \mathbf{R}^N and in \mathbf{R}^M , the sum norm is given. Then

$$\|A\| = \max_{1 \leq j \leq M} \sum_{k=1}^N |A_{kj}|$$



Induced matrix norms, examples

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Suppose that in \mathbf{R}^N , the Euclidean norm is given. Then for every **self-adjoint** matrix $A \in \mathbf{M}_{N \times N}$:

$$\|A\| = \max_{1 \leq k \leq N} |\lambda_k|,$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of A .

Suppose that in \mathbf{R}^N , the Euclidean norm is given. Then for every matrix $A \in \mathbf{M}_{N \times N}$:

$$\|A\| = \max_{1 \leq k \leq N} \sqrt{\lambda_k},$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of the (self-adjoint, positive semidefinite) matrix A^*A .



Induced matrix norms, examples

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Suppose that in \mathbf{R}^N , the Euclidean norm is given. Then for every **self-adjoint** matrix $A \in \mathbf{M}_{N \times N}$:

$$\|A\| = \max_{1 \leq k \leq N} |\lambda_k|,$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of A .

Suppose that in \mathbf{R}^N , the Euclidean norm is given. Then for every matrix $A \in \mathbf{M}_{N \times N}$:

$$\|A\| = \max_{1 \leq k \leq N} \sqrt{\lambda_k},$$

where $\lambda_1, \dots, \lambda_N$ are the eigenvalues of the (self-adjoint, positive semidefinite) matrix A^*A .



The spectral radius

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The spectral radius

Let the eigenvalues of the matrix $A \in \mathbb{M}_{N \times N}$ be the following (gerenally complex) numbers: $\lambda_1, \dots, \lambda_N$. The **spectral radius** of the matrix A is defined by:

$$\rho(A) := \max_{1 \leq k \leq N} |\lambda_k|$$

The spectral radius is not greater than any matrix norm induced by some vector norm:

$$\rho(A) \leq \|A\|$$

Moreover, if A is **self-adjoint**, then $\rho(A) = \|A\|$, where $\|A\|$ denotes the matrix norm of A induced by the Euclidean norm.



The spectral radius

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The spectral radius

Let the eigenvalues of the matrix $A \in \mathbf{M}_{N \times N}$ be the following (gerenally complex) numbers: $\lambda_1, \dots, \lambda_N$. The **spectral radius** of the matrix A is defined by:

$$\rho(A) := \max_{1 \leq k \leq N} |\lambda_k|$$

The spectral radius is not greater than any matrix norm induced by some vector norm:

$$\rho(A) \leq \|A\|$$

Moreover, if A is **self-adjoint**, then $\rho(A) = \|A\|$, where $\|A\|$ denotes the matrix norm of A induced by the Euclidean norm.



The spectral radius

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The spectral radius

Let the eigenvalues of the matrix $A \in \mathbf{M}_{N \times N}$ be the following (gerenally complex) numbers: $\lambda_1, \dots, \lambda_N$. The **spectral radius** of the matrix A is defined by:

$$\rho(A) := \max_{1 \leq k \leq N} |\lambda_k|$$

The spectral radius is not greater than any matrix norm induced by some vector norm:

$$\rho(A) \leq \|A\|$$

Moreover, if A is **self-adjoint**, then $\rho(A) = \|A\|$, where $\|A\|$ denotes the matrix norm of A induced by the Euclidean norm.



Condition number

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Condition number

The **condition number** of a regular matrix $A \in \mathbb{M}_{N \times N}$ is defined by:

$$\text{cond}(A) := \|A\| \cdot \|A^{-1}\|$$

Obviously: $\text{cond}(A) \geq 1$, and $\text{cond}(I) = 1$.

If $A \in \mathbb{M}_{N \times N}$ is self-adjoint, then the condition number can be computed from the extremal eigenvalues:

$$\text{cond}(A) = \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Condition number

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Condition number

The **condition number** of a regular matrix $A \in \mathbb{M}_{N \times N}$ is defined by:

$$\text{cond}(A) := \|A\| \cdot \|A^{-1}\|$$

Obviously: $\text{cond}(A) \geq 1$, and $\text{cond}(I) = 1$.

If $A \in \mathbb{M}_{N \times N}$ is self-adjoint, then the condition number can be computed from the extremal eigenvalues:

$$\text{cond}(A) = \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$

Condition number

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Condition number

The **condition number** of a regular matrix $A \in \mathbf{M}_{N \times N}$ is defined by:

$$\text{cond}(A) := \|A\| \cdot \|A^{-1}\|$$

Obviously: $\text{cond}(A) \geq 1$, and $\text{cond}(I) = 1$.

If $A \in \mathbf{M}_{N \times N}$ is self-adjoint, then the condition number can be computed from the extremal eigenvalues:

$$\text{cond}(A) = \frac{|\lambda|_{\max}}{|\lambda|_{\min}}$$



Linear systems of equations

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a given matrix, and let $b \in \mathbf{R}^N$ be a given vector. Consider the equation

$$Ax = b$$

This is equivalent to the following system of linear equations with N unknowns:

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1N}x_N = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2N}x_N = b_2$$

.....

$$a_{N1}x_1 + a_{N2}x_2 + \dots + a_{NN}x_N = b_N$$

The system is **homogeneous**, if $b = 0$. In this case, $x = 0$ is always a solution (**trivial solution**). The solution x is said to be a **nontrivial solution**, if at least one component of x differs from zero.



Linear systems of equations

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a given matrix, and let $b \in \mathbf{R}^N$ be a given vector. Consider the equation

$$Ax = b$$

This is equivalent to the following system of linear equations with N unknowns:

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1N}x_N = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2N}x_N = b_2$$

.....

$$a_{N1}x_1 + a_{N2}x_2 + \dots + a_{NN}x_N = b_N$$

The system is **homogeneous**, if $b = 0$. In this case, $x = 0$ is always a solution (**trivial solution**). The solution x is said to be a **nontrivial solution**, if at least one component of x differs from zero.



Linear systems of equations

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbf{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a given matrix, and let $b \in \mathbf{R}^N$ be a given vector. Consider the equation

$$Ax = b$$

This is equivalent to the following system of linear equations with N unknowns:

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1N}x_N = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2N}x_N = b_2$$

.....

$$a_{N1}x_1 + a_{N2}x_2 + \dots + a_{NN}x_N = b_N$$

The system is **homogeneous**, if $b = \mathbf{0}$. In this case, $x = \mathbf{0}$ is always a solution (**trivial solution**). The solution x is said to be a **nontrivial solution**, if at least one component of x differs from zero.



Solvability of linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbb{M}_{N \times N}$ is regular if and only if the equation $Ax = b$ has a solution for every right-hand side. In this case, the solution is unique, namely: $x = A^{-1}b$.

The matrix $A \in \mathbb{M}_{N \times N}$ is regular if and only if the corresponding homogeneous equation $Ax = 0$ has the trivial solution only, i.e. the matrix A is singular if and only if the corresponding homogeneous equation has a nontrivial solution (in this case, infinitely many nontrivial solutions exist).



Solvability of linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbf{M}_{N \times N}$ is regular if and only if the equation $Ax = b$ has a solution for every right-hand side. In this case, the solution is unique, namely: $x = A^{-1}b$.

The matrix $A \in \mathbf{M}_{N \times N}$ is regular if and only if the corresponding homogeneous equation $Ax = \mathbf{0}$ has the trivial solution only, i.e. the matrix A is singular if and only if the corresponding homogeneous equation has a nontrivial solution (in this case, infinitely many nontrivial solutions exist).



Solvability of linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

The matrix $A \in \mathbf{M}_{N \times N}$ is regular if and only if the equation $Ax = b$ has a solution for every right-hand side. In this case, the solution is unique, namely: $x = A^{-1}b$.

The matrix $A \in \mathbf{M}_{N \times N}$ is regular if and only if the corresponding homogeneous equation $Ax = \mathbf{0}$ has the trivial solution only, i.e. the matrix A is singular if and only if the corresponding homogeneous equation has a nontrivial solution (in this case, infinitely many nontrivial solutions exist).

Perturbed linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of linear equations $Ax = b$ and the perturbed system as well: $A(x + \Delta x) = b + \Delta b$.

Hence $A\Delta x = \Delta b$, therefore $\|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$.

On the other hand: $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$, therefore

$$\frac{1}{\|x\|} \cdot \frac{1}{\|A\|} \leq \frac{1}{\|b\|}.$$

This implies:

Perturbation lemma

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \cdot \frac{\|\Delta b\|}{\|b\|}$$

The condition number characterizes how sensitive the solution of the system is to the changes in the right-hand side.

Perturbed linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of linear equations $Ax = b$ and the perturbed system as well: $A(x + \Delta x) = b + \Delta b$.

Hence $A\Delta x = \Delta b$, therefore $\|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$.

On the other hand: $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$, therefore

$$\frac{1}{\|x\|} \cdot \frac{1}{\|A\|} \leq \frac{1}{\|b\|}.$$

This implies:

Perturbation lemma

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \cdot \frac{\|\Delta b\|}{\|b\|}$$

The condition number characterizes how sensitive the solution of the system is to the changes in the right-hand side.

Perturbed linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of linear equations $Ax = b$ and the perturbed system as well: $A(x + \Delta x) = b + \Delta b$.

Hence $A\Delta x = \Delta b$, therefore $\|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$.

On the other hand: $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$, therefore

$$\frac{1}{\|x\|} \cdot \frac{1}{\|A\|} \leq \frac{1}{\|b\|}.$$

This implies:

Perturbation lemma

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \cdot \frac{\|\Delta b\|}{\|b\|}$$

The condition number characterizes how sensitive the solution of the system is to the changes in the right-hand side.

Perturbed linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of linear equations $Ax = b$ and the perturbed system as well: $A(x + \Delta x) = b + \Delta b$.

Hence $A\Delta x = \Delta b$, therefore $\|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$.

On the other hand: $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$, therefore

$$\frac{1}{\|x\|} \cdot \frac{1}{\|A\|} \leq \frac{1}{\|b\|}.$$

This implies:

Perturbation lemma

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \cdot \frac{\|\Delta b\|}{\|b\|}$$

The condition number characterizes how sensitive the solution of the system is to the changes in the right-hand side.

Perturbed linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of linear equations $Ax = b$ and the perturbed system as well: $A(x + \Delta x) = b + \Delta b$.

Hence $A\Delta x = \Delta b$, therefore $\|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$.

On the other hand: $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$, therefore

$$\frac{1}{\|x\|} \cdot \frac{1}{\|A\|} \leq \frac{1}{\|b\|}.$$

This implies:

Perturbation lemma

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \cdot \frac{\|\Delta b\|}{\|b\|}$$

The condition number characterizes how sensitive the solution of the system is to the changes in the right-hand side.



Perturbed linear systems

Numerical
Analysis

by Csaba
Gáspár

Introduction

Motivations

Vectors and
matrices

Norm and
distance in \mathbb{R}^N

Inner product

Orthogonality

Matrix norms

Linear systems

Consider the system of linear equations $Ax = b$ and the perturbed system as well: $A(x + \Delta x) = b + \Delta b$.

Hence $A\Delta x = \Delta b$, therefore $\|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$.

On the other hand: $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$, therefore

$$\frac{1}{\|x\|} \cdot \frac{1}{\|A\|} \leq \frac{1}{\|b\|}.$$

This implies:

Perturbation lemma

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \cdot \frac{\|\Delta b\|}{\|b\|}$$

The condition number characterizes how sensitive the solution of the system is to the changes in the right-hand side.