



Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Numerical Analysis Matrix decompositions

by Csaba Gáspár

Széchenyi István University

2020, autumn semester



LU decomposition by Gaussian elimination

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a regular matrix, for which the Gaussian elimination can be performed *without swapping rows*.

LU decomposition

The matrix A can be uniquely decomposed in the following form:

$$A = LU$$

where L is a **normed lower triangular matrix** (i.e. $L_{kk} = 1$, and $L_{kj} = 0$, if $j > k$), and U is an **upper triangular matrix** (i.e. $U_{kj} = 0$, whenever $j < k$).

If a linear system $Ax = b$ should be solved with a lot of different right-hand sides b , then it is much cheaper to solve the pairs of equations instead: $Ly = b$, $Ux = y$, since only forward and backward substitutions have to be performed.



LU decomposition by Gaussian elimination

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A = [a_{kj}] \in \mathbf{M}_{N \times N}$ be a regular matrix, for which the Gaussian elimination can be performed *without swapping rows*.

LU decomposition

The matrix A can be uniquely decomposed in the following form:

$$A = LU$$

where L is a **normed lower triangular matrix** (i.e. $L_{kk} = 1$, and $L_{kj} = 0$, if $j > k$), and U is an **upper triangular matrix** (i.e. $U_{kj} = 0$, whenever $j < k$).

If a linear system $Ax = b$ should be solved with a lot of different right-hand sides b , then it is much cheaper to solve the pairs of equations instead: $Ly = b$, $Ux = y$, since only forward and backward substitutions have to be performed.



Some useful statements

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

- The product of (normed) lower triangular matrix is (normed) lower triangular
- The product of (normed) upper triangular matrix is (normed) upper triangular
- The inverse of a (normed) lower triangular matrix is (normed) lower triangular (if exists)
- The inverse of a (normed) upper triangular matrix is (normed) upper triangular (if exists)



Some useful statements

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

- The product of (normed) lower triangular matrix is (normed) lower triangular
- The product of (normed) upper triangular matrix is (normed) upper triangular
- The inverse of a (normed) lower triangular matrix is (normed) lower triangular (if exists)
- The inverse of a (normed) upper triangular matrix is (normed) upper triangular (if exists)



Some useful statements

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

- The product of (normed) lower triangular matrix is (normed) lower triangular
- The product of (normed) upper triangular matrix is (normed) upper triangular
- The inverse of a (normed) lower triangular matrix is (normed) lower triangular (if exists)
- The inverse of a (normed) upper triangular matrix is (normed) upper triangular (if exists)



Some useful statements

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

- The product of (normed) lower triangular matrix is (normed) lower triangular
- The product of (normed) upper triangular matrix is (normed) upper triangular
- The inverse of a (normed) lower triangular matrix is (normed) lower triangular (if exists)
- The inverse of a (normed) upper triangular matrix is (normed) upper triangular (if exists)

LU decomposition by Gaussian elimination

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Performing the decomposition by Gaussian elimination:
At the elimination by the k th row, subtract the k th row multiplied by $l_{m,k} := a_{m,k}/a_{k,k}$ from the m th row ($m = k + 1, \dots, N$).

From these numbers $l_{m,k}$, the matrix L can be assembled:

$$L = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ l_{2,1} & 1 & 0 & \dots & 0 \\ l_{3,1} & l_{3,2} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ l_{N,1} & l_{N,2} & l_{N,3} & \dots & 1 \end{pmatrix}$$

After the elimination steps, from the original matrix A , we arrive at the matrix U .



LU decomposition by Gaussian elimination

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Performing the decomposition by Gaussian elimination:
At the elimination by the k th row, subtract the k th row multiplied by $l_{m,k} := a_{m,k}/a_{k,k}$ from the m th row ($m = k + 1, \dots, N$).

From these numbers $l_{m,k}$, the matrix L can be assembled:

$$L = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ l_{2,1} & 1 & 0 & \dots & 0 \\ l_{3,1} & l_{3,2} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ l_{N,1} & l_{N,2} & l_{N,3} & \dots & 1 \end{pmatrix}$$

After the elimination steps, from the original matrix A , we arrive at the matrix U .



LU decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Determine the LU decomposition of the matrix

$$A = \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix}$$



LU decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$\begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ . & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 7 & -14 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & 35 \end{pmatrix} = U$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} = L$$



LU decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$\begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ . & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 7 & -14 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & 35 \end{pmatrix} = U$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} = L$$



LU decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$\begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ . & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 7 & -14 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & 35 \end{pmatrix} = U$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} = L$$



LU decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$\begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ . & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 3 & -2 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ . & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 7 & -14 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & . & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & 35 \end{pmatrix} = U$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} = L$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

In the previous example, first, write down the entries of L and U which are a priori known:

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & \ell_{32} & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd row of L by the 2nd column of U , u_{22} can be computed: $u_{22} = 1$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & \ell_{32} & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd row of L by the 2nd column of U , u_{22} can be computed: $u_{22} = 1$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & u_{23} \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & \ell_{32} & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 3rd row of L by the 2nd column of U , ℓ_{32} can be computed: $\ell_{32} = 7$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & u_{23} \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & \ell_{32} & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 3rd row of L by the 2nd column of U , ℓ_{32} can be computed: $\ell_{32} = 7$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & u_{23} \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd row of L by the 3rd column of U , u_{23} can be computed: $u_{23} = -7$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & u_{23} \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd row of L by the 3rd column of U , u_{23} can be computed: $u_{23} = -7$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Finally, multiplying the 3rd row of L by the 3rd column of U ,
 u_{33} can be computed: $u_{33} = 35$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & u_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix} \begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Finally, multiplying the 3rd row of L by the 3rd column of U , u_{33} can be computed: $u_{33} = 35$.

$$U = \begin{pmatrix} 2 & -6 & 10 \\ 0 & 1 & -7 \\ 0 & 0 & 35 \end{pmatrix}$$

$$L = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ \frac{3}{2} & 7 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 2 & -6 & 10 \\ 2 & -5 & 3 \\ 3 & -2 & 1 \end{pmatrix} = A$$



LU decomposition, numerical features

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Computational cost

The necessary number of arithmetic operations of the LU decomposition of an $N \times N$ matrix is $\mathcal{O}(N^3)$.

Calculation of the determinant

Once the LU decomposition $A = LU$ has been performed, the determinant of A can be calculated as:

$$\det(A) = \det(L) \cdot \det(U) = u_{11} \cdot u_{22} \cdot \dots \cdot u_{NN}$$



LU decomposition, numerical features

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Computational cost

The necessary number of arithmetic operations of the LU decomposition of an $N \times N$ matrix is $\mathcal{O}(N^3)$.

Calculation of the determinant

Once the LU decomposition $A = LU$ has been performed, the determinant of A can be calculated as:

$$\det(A) = \det(L) \cdot \det(U) = u_{11} \cdot u_{22} \cdot \dots \cdot u_{NN}$$



The LDL^* factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a **self-adjoint, positive definite** matrix (which assures that the Gaussian elimination can be performed without swapping rows). Consider the LU decomposition of A :

$$A = LU$$

Denote by D the diagonal part of U , then $U = DU'$, where U' is a normed upper triangular matrix. Since A is self-adjoint, the uniqueness of the LU decomposition implies that

$$U' = L^*$$

therefore

$$A = LDL^*$$



The LDL^* factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a **self-adjoint, positive definite** matrix (which assures that the Gaussian elimination can be performed without swapping rows). Consider the LU decomposition of A :

$$A = LU$$

Denote by D the diagonal part of U , then $U = DU'$, where U' is a normed upper triangular matrix. Since A is self-adjoint, the uniqueness of the LU decomposition implies that

$$U' = L^*$$

therefore

$$A = LDL^*$$



The LDL^* factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a **self-adjoint, positive definite** matrix (which assures that the Gaussian elimination can be performed without swapping rows). Consider the LU decomposition of A :

$$A = LU$$

Denote by D the diagonal part of U , then $U = DU'$, where U' is a normed upper triangular matrix. Since A is self-adjoint, the uniqueness of the LU decomposition implies that

$$U' = L^*$$

therefore

$$A = LDL^*$$



The LDL^* factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

LDL^* decomposition

Every self-adjoint, positive definite matrix A can be uniquely decomposed in the following form:

$$A = LDL^*$$

where L is a normed lower triangular matrix and D is a diagonal matrix, the diagonal entries of which are positive.



The Cholesky factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint, positive definite matrix with the LDL^* decomposition:

$$A = L_0 D L_0^*$$

Then

$$A = L_0 \sqrt{D} \sqrt{D} L_0^* = (L_0 \sqrt{D}) \cdot (L_0 \sqrt{D})^* =: LL^*$$

Cholesky decomposition

Every self-adjoint, positive definite matrix A can be uniquely decomposed in the following form:

$$A = LL^*$$

where L is a not necessarily normed lower triangular matrix, with positive diagonal elements.



The Cholesky factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint, positive definite matrix with the LDL^* decomposition:

$$A = L_0 D L_0^*$$

Then

$$A = L_0 \sqrt{D} \sqrt{D} L_0^* = (L_0 \sqrt{D}) \cdot (L_0 \sqrt{D})^* =: LL^*$$

Cholesky decomposition

Every self-adjoint, positive definite matrix A can be uniquely decomposed in the following form:

$$A = LL^*$$

where L is a not necessarily normed lower triangular matrix, with positive diagonal elements.



The Cholesky factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint, positive definite matrix with the LDL^* decomposition:

$$A = L_0 D L_0^*$$

Then

$$A = L_0 \sqrt{D} \sqrt{D} L_0^* = (L_0 \sqrt{D}) \cdot (L_0 \sqrt{D})^* =: LL^*$$

Cholesky decomposition

Every self-adjoint, positive definite matrix A can be uniquely decomposed in the following form:

$$A = LL^*$$

where L is a not necessarily normed lower triangular matrix, with positive diagonal elements.



The Cholesky factorization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $A \in \mathbf{M}_{N \times N}$ be a self-adjoint, positive definite matrix with the LDL^* decomposition:

$$A = L_0 D L_0^*$$

Then

$$A = L_0 \sqrt{D} \sqrt{D} L_0^* = (L_0 \sqrt{D}) \cdot (L_0 \sqrt{D})^* =: LL^*$$

Cholesky decomposition

Every self-adjoint, positive definite matrix A can be uniquely decomposed in the following form:

$$A = LL^*$$

where L is a not necessarily normed lower triangular matrix, with positive diagonal elements.

The Cholesky decomposition, numerical features

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Computational cost

The number of necessary arithmetic operations of the Cholesky decomposition of an $N \times N$ self-adjoint, positive definite matrix is $\mathcal{O}(N^3)$.

The Cholesky decomposition is less sensitive to the roundoff errors than the LU decomposition.

If a linear system $Ax = b$ should be solved with a lot of different right-hand sides b , then it is cheaper to solve the pairs of equations instead: $Ly = b$, $L^*x = y$, since only forward and backward substitutions have to be performed.



The Cholesky decomposition, numerical features

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Computational cost

The number of necessary arithmetic operations of the Cholesky decomposition of an $N \times N$ self-adjoint, positive definite matrix is $\mathcal{O}(N^3)$.

The Cholesky decomposition is less sensitive to the roundoff errors than the LU decomposition.

If a linear system $Ax = b$ should be solved with a lot of different right-hand sides b , then it is cheaper to solve the pairs of equations instead: $Ly = b$, $L^*x = y$, since only forward and backward substitutions have to be performed.



The Cholesky decomposition, numerical features

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Computational cost

The number of necessary arithmetic operations of the Cholesky decomposition of an $N \times N$ self-adjoint, positive definite matrix is $\mathcal{O}(N^3)$.

The Cholesky decomposition is less sensitive to the roundoff errors than the LU decomposition.

If a linear system $Ax = b$ should be solved with a lot of different right-hand sides b , then it is cheaper to solve the pairs of equations instead: $Ly = b$, $L^*x = y$, since only forward and backward substitutions have to be performed.



The Cholesky decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$A := \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$$

The LU decomposition (please check):

$$A = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} 2 & 1 & 0 \\ 0 & \frac{3}{2} & 1 \\ 0 & 0 & \frac{4}{3} \end{pmatrix}$$

Thus, the Cholesky decomposition: $A = L \cdot L^*$, where

$$L = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{\frac{3}{2}} & 0 \\ 0 & 0 & \sqrt{\frac{4}{3}} \end{pmatrix} = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \frac{\sqrt{2}}{2} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \frac{2}{\sqrt{3}} \end{pmatrix}$$



The Cholesky decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$A := \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$$

The LU decomposition (please check):

$$A = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} 2 & 1 & 0 \\ 0 & \frac{3}{2} & 1 \\ 0 & 0 & \frac{4}{3} \end{pmatrix}$$

Thus, the Cholesky decomposition: $A = L \cdot L^*$, where

$$L = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{\frac{3}{2}} & 0 \\ 0 & 0 & \sqrt{\frac{4}{3}} \end{pmatrix} = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \frac{\sqrt{2}}{2} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \frac{2}{\sqrt{3}} \end{pmatrix}$$



The Cholesky decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$A := \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$$

The LU decomposition (please check):

$$A = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} 2 & 1 & 0 \\ 0 & \frac{3}{2} & 1 \\ 0 & 0 & \frac{4}{3} \end{pmatrix}$$

Thus, the Cholesky decomposition: $A = L \cdot L^*$, where

$$L = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{\frac{3}{2}} & 0 \\ 0 & 0 & \sqrt{\frac{4}{3}} \end{pmatrix} = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \frac{\sqrt{2}}{2} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \frac{2}{\sqrt{3}} \end{pmatrix}$$



The Cholesky decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

$$A := \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix}$$

The LU decomposition (please check):

$$A = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} 2 & 1 & 0 \\ 0 & \frac{3}{2} & 1 \\ 0 & 0 & \frac{4}{3} \end{pmatrix}$$

Thus, the Cholesky decomposition: $A = L \cdot L^*$, where

$$L = \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ 0 & \frac{2}{3} & 1 \end{pmatrix} \cdot \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{\frac{3}{2}} & 0 \\ 0 & 0 & \sqrt{\frac{4}{3}} \end{pmatrix} = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \frac{\sqrt{2}}{2} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \frac{2}{\sqrt{3}} \end{pmatrix}$$



Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

In the previous example, first, write down the entries of L which are a priori known:

$$L^* = \begin{pmatrix} \ell_{11} & \ell_{21} & \ell_{31} \\ 0 & \ell_{22} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} \ell_{11} & 0 & 0 \\ \ell_{21} & \ell_{22} & 0 \\ \ell_{31} & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$



Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the first row of L by the first column of L^* , ℓ_{11} can be computed: $\ell_{11} = \sqrt{2}$.

$$L^* = \begin{pmatrix} \ell_{11} & \ell_{21} & \ell_{31} \\ 0 & \ell_{22} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} \ell_{11} & 0 & 0 \\ \ell_{21} & \ell_{22} & 0 \\ \ell_{31} & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$

Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the first row of L by the first column of L^* , ℓ_{11} can be computed: $\ell_{11} = \sqrt{2}$.

$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \ell_{21} & \ell_{22} & 0 \\ \ell_{31} & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} \sqrt{2} & \ell_{21} & \ell_{31} \\ 0 & \ell_{22} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix} = A$$



Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd and 3rd rows of L by the first column of L^* , ℓ_{21} , ℓ_{31} can be computed: $\ell_{21} = \sqrt{\frac{1}{2}}$, $\ell_{31} = 0$.

$$L^* = \begin{pmatrix} \sqrt{2} & \ell_{21} & \ell_{31} \\ 0 & \ell_{22} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$

$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \ell_{21} & \ell_{22} & 0 \\ \ell_{31} & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$

Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd and 3rd rows of L by the first column of L^* , ℓ_{21} , ℓ_{31} can be computed: $\ell_{21} = \sqrt{\frac{1}{2}}$, $\ell_{31} = 0$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \ell_{22} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \ell_{22} & 0 \\ 0 & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$



Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd row of L by the 2nd column of L^* , ℓ_{22} can be computed: $\ell_{22} = \sqrt{\frac{3}{2}}$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \ell_{22} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \ell_{22} & 0 \\ 0 & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$



Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 2nd row of L by the 2nd column of L^* , ℓ_{22} can be computed: $\ell_{22} = \sqrt{\frac{3}{2}}$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \sqrt{\frac{3}{2}} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$

$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$



Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 3rd row of L by the 2nd column of L^* , ℓ_{32} can be computed: $\ell_{32} = \sqrt{\frac{2}{3}}$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \sqrt{\frac{3}{2}} & \ell_{32} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$

$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \ell_{32} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$

Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Multiplying the 3rd row of L by the 2nd column of L^* , ℓ_{32} can be computed: $\ell_{32} = \sqrt{\frac{2}{3}}$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \sqrt{\frac{3}{2}} & \sqrt{\frac{2}{3}} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$

$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$

Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Finally, multiplying the 3rd row of L by the 3rd column of L^* , ℓ_{33} can be computed: $\ell_{33} = \sqrt{\frac{4}{3}}$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \sqrt{\frac{3}{2}} & \sqrt{\frac{2}{3}} \\ 0 & 0 & \ell_{33} \end{pmatrix}$$
$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \ell_{33} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$

Cholesky decomposition by matrix multiplication

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Finally, multiplying the 3rd row of L by the 3rd column of L^* , ℓ_{33} can be computed: $\ell_{33} = \sqrt{\frac{4}{3}}$.

$$L^* = \begin{pmatrix} \sqrt{2} & \sqrt{\frac{1}{2}} & 0 \\ 0 & \sqrt{\frac{3}{2}} & \sqrt{\frac{2}{3}} \\ 0 & 0 & \sqrt{\frac{4}{3}} \end{pmatrix}$$

$$L = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ \sqrt{\frac{1}{2}} & \sqrt{\frac{3}{2}} & 0 \\ 0 & \sqrt{\frac{2}{3}} & \sqrt{\frac{4}{3}} \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{pmatrix} = A$$



The Gram-Schmidt orthogonalization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition
The Cholesky
decomposition

The QR
decomposition

Let $a_1, a_2, \dots, a_N \in \mathbf{R}^N$ be linearly independent vectors. Define $\tilde{e}_1 := a_1$, $e_1 := \frac{\tilde{e}_1}{\|\tilde{e}_1\|}$, and for $k = 2, 3, \dots, N$:

$$\tilde{e}_k := a_k - \sum_{j=1}^{k-1} \langle a_k, e_j \rangle e_j, \quad e_k := \frac{\tilde{e}_k}{\|\tilde{e}_k\|}$$

Gram-Schmidt orthogonalization

The vectors e_1, e_2, \dots, e_N form an **orthonormal vector system** (i.e. $\langle e_k, e_j \rangle = 0$, if $k \neq j$, and $\|e_k\| = 1$), and the vector systems a_1, a_2, \dots, a_k and e_1, e_2, \dots, e_k generate the same subspace of \mathbf{R}^N for all $k = 1, 2, \dots, N$.



The Gram-Schmidt orthogonalization

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Let $a_1, a_2, \dots, a_N \in \mathbf{R}^N$ be linearly independent vectors. Define $\tilde{e}_1 := a_1$, $e_1 := \frac{\tilde{e}_1}{\|\tilde{e}_1\|}$, and for $k = 2, 3, \dots, N$:

$$\tilde{e}_k := a_k - \sum_{j=1}^{k-1} \langle a_k, e_j \rangle e_j, \quad e_k := \frac{\tilde{e}_k}{\|\tilde{e}_k\|}$$

Gram-Schmidt orthogonalization

The vectors e_1, e_2, \dots, e_N form an **orthonormal vector system** (i.e. $\langle e_k, e_j \rangle = 0$, if $k \neq j$, and $\|e_k\| = 1$), and the vector systems a_1, a_2, \dots, a_k and e_1, e_2, \dots, e_k generate the same subspace of \mathbf{R}^N for all $k = 1, 2, \dots, N$.



Orthogonal matrices

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition
The Cholesky
decomposition

The QR
decomposition

The matrix $A \in \mathbf{M}_{N \times N}$ is called **orthogonal**, if its column vectors form an orthonormal system in \mathbf{R}^N .

Example: The 2-by-2 matrix $A := \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$ is an orthogonal matrix.

The inverse of orthogonal matrices can be computed in an extremely simple way:

If the matrix $A \in \mathbf{M}_{N \times N}$ is orthogonal, then $A^{-1} = A^*$



Orthogonal matrices

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition
The Cholesky
decomposition

The QR
decomposition

The matrix $A \in \mathbf{M}_{N \times N}$ is called **orthogonal**, if its column vectors form an orthonormal system in \mathbf{R}^N .

Example: The 2-by-2 matrix $A := \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$ is an orthogonal matrix.

The inverse of orthogonal matrices can be computed in an extremely simple way:

If the matrix $A \in \mathbf{M}_{N \times N}$ is orthogonal, then $A^{-1} = A^*$



Orthogonal matrices

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition
The Cholesky
decomposition

The QR
decomposition

The matrix $A \in \mathbf{M}_{N \times N}$ is called **orthogonal**, if its column vectors form an orthonormal system in \mathbf{R}^N .

Example: The 2-by-2 matrix $A := \begin{pmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{pmatrix}$ is an orthogonal matrix.

The inverse of orthogonal matrices can be computed in an extremely simple way:

If the matrix $A \in \mathbf{M}_{N \times N}$ is orthogonal, then $A^{-1} = A^*$



The QR decomposition

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition
The Cholesky
decomposition

The QR
decomposition

Every regular matrix $A \in \mathbb{M}_{N \times N}$ can be uniquely decomposed in the form $A = QR$, where Q is an orthogonal matrix, R is an upper triangular matrix with positive diagonal entries.

Denote by a_1, a_2, \dots, a_N its column vectors. Consider the orthonormal basis e_1, e_2, \dots, e_N obtained from a_1, a_2, \dots, a_N by a Gram-Schmidt orthogonalization. Then

$$Q = \left(\begin{array}{c|c|c|c} e_1 & e_2 & \dots & e_N \end{array} \right), \quad R = \begin{pmatrix} r_{11} & r_{12} & r_{13} & \dots \\ 0 & r_{22} & r_{23} & \dots \\ 0 & 0 & r_{33} & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

where $r_{kj} := \langle e_k, a_j \rangle$

After performing the QR decomposition, the solution of the linear system $Ax = b$ is extremely simple with low computational cost:

$$Rx = Q^{-1}b = Q^*b$$

The QR decomposition

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Every regular matrix $A \in \mathbf{M}_{N \times N}$ can be uniquely decomposed in the form $A = QR$, where Q is an orthogonal matrix, R is an upper triangular matrix with positive diagonal entries.

Denote by a_1, a_2, \dots, a_N its column vectors. Consider the orthonormal basis e_1, e_2, \dots, e_N obtained from a_1, a_2, \dots, a_N by a Gram-Schmidt orthogonalization. Then

$$Q = \left(\begin{array}{c|c|c|c} e_1 & e_2 & \dots & e_N \end{array} \right), \quad R = \begin{pmatrix} r_{11} & r_{12} & r_{13} & \dots \\ 0 & r_{22} & r_{23} & \dots \\ 0 & 0 & r_{33} & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

where $r_{kj} := \langle e_k, a_j \rangle$

After performing the QR decomposition, the solution of the linear system $Ax = b$ is extremely simple with low computational cost:

$$Rx = Q^{-1}b = Q^*b$$



The QR decomposition

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Every regular matrix $A \in \mathbf{M}_{N \times N}$ can be uniquely decomposed in the form $A = QR$, where Q is an orthogonal matrix, R is an upper triangular matrix with positive diagonal entries.

Denote by a_1, a_2, \dots, a_N its column vectors. Consider the orthonormal basis e_1, e_2, \dots, e_N obtained from a_1, a_2, \dots, a_N by a Gram-Schmidt orthogonalization. Then

$$Q = \left(\begin{array}{c|c|c|c} e_1 & e_2 & \dots & e_N \end{array} \right), \quad R = \begin{pmatrix} r_{11} & r_{12} & r_{13} & \dots \\ 0 & r_{22} & r_{23} & \dots \\ 0 & 0 & r_{33} & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix}$$

where $r_{kj} := \langle e_k, a_j \rangle$

After performing the QR decomposition, the solution of the linear system $Ax = b$ is extremely simple with low computational cost:

$$Rx = Q^{-1}b = Q^*b$$



The QR decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition
The Cholesky
decomposition

The QR
decomposition

Compute the QR decomposition of the matrix $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 3 \end{pmatrix}$.

$$\text{First, } a_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad a_2 = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \quad a_3 = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}.$$

Next, perform a Gram-Schmidt-orthogonalization for these vectors:

$$\tilde{e}_1 = a_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad e_1 = \begin{pmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix}$$

$$\tilde{e}_2 = a_2 - \langle a_2, e_1 \rangle \cdot e_1 = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\tilde{e}_3 = a_3 - \langle a_3, e_1 \rangle \cdot e_1 - \langle a_3, e_2 \rangle \cdot e_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \quad e_3 = \begin{pmatrix} -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix}$$

The QR decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Compute the QR decomposition of the matrix $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 3 \end{pmatrix}$.

$$\text{First, } a_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad a_2 = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \quad a_3 = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}.$$

Next, perform a Gram-Schmidt-orthogonalization for these vectors:

$$\tilde{e}_1 = a_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad e_1 = \begin{pmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix}$$

$$\tilde{e}_2 = a_2 - \langle a_2, e_1 \rangle \cdot e_1 = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\tilde{e}_3 = a_3 - \langle a_3, e_1 \rangle \cdot e_1 - \langle a_3, e_2 \rangle \cdot e_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \quad e_3 = \begin{pmatrix} -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix}$$

The QR decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Compute the QR decomposition of the matrix $A = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 3 \end{pmatrix}$.

$$\text{First, } a_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad a_2 = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \quad a_3 = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix}.$$

Next, perform a Gram-Schmidt-orthogonalization for these vectors:

$$\tilde{e}_1 = a_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad e_1 = \begin{pmatrix} 1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix}$$

$$\tilde{e}_2 = a_2 - \langle a_2, e_1 \rangle \cdot e_1 = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\tilde{e}_3 = a_3 - \langle a_3, e_1 \rangle \cdot e_1 - \langle a_3, e_2 \rangle \cdot e_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \quad e_3 = \begin{pmatrix} -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{pmatrix}$$

The QR decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Thus we have obtained:

$$Q = \begin{pmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix}$$

Moreover:

$$\begin{aligned} r_{11} &= \|\tilde{e}_1\| = \sqrt{2}, & r_{12} &= \langle e_1, a_2 \rangle = 0, & r_{13} &= \langle e_1, a_3 \rangle = 2\sqrt{2} \\ r_{22} &= \|\tilde{e}_2\| = 2, & r_{23} &= \langle e_2, a_3 \rangle = 0 \\ r_{33} &= \|\tilde{e}_3\| = \sqrt{2} \end{aligned}$$

and therefore:

$$R = \begin{pmatrix} \sqrt{2} & 0 & 2\sqrt{2} \\ 0 & 2 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix}$$

The QR decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Thus we have obtained:

$$Q = \begin{pmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix}$$

Moreover:

$$\begin{aligned} r_{11} &= \|\tilde{e}_1\| = \sqrt{2}, & r_{12} &= \langle e_1, a_2 \rangle = 0, & r_{13} &= \langle e_1, a_3 \rangle = 2\sqrt{2} \\ r_{22} &= \|\tilde{e}_2\| = 2, & r_{23} &= \langle e_2, a_3 \rangle = 0 \\ r_{33} &= \|\tilde{e}_3\| = \sqrt{2} \end{aligned}$$

and therefore:

$$R = \begin{pmatrix} \sqrt{2} & 0 & 2\sqrt{2} \\ 0 & 2 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix}$$

The QR decomposition, example

Numerical
Analysis

by Csaba
Gáspár

Matrix decom-
positions

The LU
decomposition

The Cholesky
decomposition

The QR
decomposition

Thus we have obtained:

$$Q = \begin{pmatrix} 1/\sqrt{2} & 0 & -1/\sqrt{2} \\ 0 & 1 & 0 \\ 1/\sqrt{2} & 0 & 1/\sqrt{2} \end{pmatrix}$$

Moreover:

$$\begin{aligned} r_{11} &= \|\tilde{e}_1\| = \sqrt{2}, & r_{12} &= \langle e_1, a_2 \rangle = 0, & r_{13} &= \langle e_1, a_3 \rangle = 2\sqrt{2} \\ r_{22} &= \|\tilde{e}_2\| = 2, & r_{23} &= \langle e_2, a_3 \rangle = 0 \\ r_{33} &= \|\tilde{e}_3\| = \sqrt{2} \end{aligned}$$

and therefore:

$$R = \begin{pmatrix} \sqrt{2} & 0 & 2\sqrt{2} \\ 0 & 2 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix}$$