



Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Numerical Analysis

The discrete and fast Fourier Transform

by Csaba Gáspár

Széchenyi István University

2020, autumn semester



Trigonometric Fourier series in $L_2(0, 2\pi)$

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

An arbitrary real function $f \in L_2(0, 2\pi)$ can be expressed as a trigonometric Fourier series which is convergent with respect to the $L_2(0, 2\pi)$ -norm:

$$f(x) = a_0 + \sum_{k=1}^{\infty} a_k \cos kx + \sum_{k=1}^{\infty} b_k \sin kx$$

where the coefficients can be calculated as:

$$a_0 = \frac{1}{2\pi} \int_0^{2\pi} f(x) dx,$$

$$a_k = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos kx dx, \quad b_k = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin kx dx$$



Complex exponential function

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

For any $z \in \mathbb{C}$, define the **exponential series** as:

$$e^z := \sum_{k=0}^{\infty} \frac{z^k}{k!} = 1 + \frac{z}{1!} + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots$$

Euler's formula

For every $t \in \mathbb{R}$: $e^{it} = \cos t + i \cdot \sin t$

Utilizing the well-known Taylor series of the sine and cosine functions:

$$\cos t = 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \dots, \quad \sin t = \frac{t}{1!} - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots$$

which implies that:

$$e^{it} = 1 + \frac{it}{1!} + \frac{i^2 t^2}{2!} + \frac{i^3 t^3}{3!} + \frac{i^4 t^4}{4!} + \dots = 1 + \frac{it}{1!} - \frac{t^2}{2!} - \frac{it^3}{3!} + \frac{t^4}{4!} + \dots$$

Separating the real and imaginary parts, we have the theorem.





Complex exponential function

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

For any $z \in \mathbb{C}$, define the **exponential series** as:

$$e^z := \sum_{k=0}^{\infty} \frac{z^k}{k!} = 1 + \frac{z}{1!} + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots$$

Euler's formula

For every $t \in \mathbb{R}$: $e^{it} = \cos t + i \cdot \sin t$

Utilizing the well-known Taylor series of the sine and cosine functions:

$$\cos t = 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \dots, \quad \sin t = \frac{t}{1!} - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots$$

which implies that:

$$e^{it} = 1 + \frac{it}{1!} + \frac{i^2 t^2}{2!} + \frac{i^3 t^3}{3!} + \frac{i^4 t^4}{4!} + \dots = 1 + \frac{it}{1!} - \frac{t^2}{2!} - \frac{it^3}{3!} + \frac{t^4}{4!} + \dots$$

Separating the real and imaginary parts, we have the theorem.



Complex exponential function

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

For any $z \in \mathbb{C}$, define the **exponential series** as:

$$e^z := \sum_{k=0}^{\infty} \frac{z^k}{k!} = 1 + \frac{z}{1!} + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots$$

Euler's formula

For every $t \in \mathbb{R}$: $e^{it} = \cos t + i \cdot \sin t$

Utilizing the well-known Taylor series of the sine and cosine functions:

$$\cos t = 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \dots, \quad \sin t = \frac{t}{1!} - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots$$

which implies that:

$$e^{it} = 1 + \frac{it}{1!} + \frac{i^2 t^2}{2!} + \frac{i^3 t^3}{3!} + \frac{i^4 t^4}{4!} + \dots = 1 + \frac{it}{1!} - \frac{t^2}{2!} - \frac{it^3}{3!} + \frac{t^4}{4!} + \dots$$

Separating the real and imaginary parts, we have the theorem.

The Discrete Fourier Transform (DFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

If $f_0, f_1, \dots, f_{N-1} \in \mathbf{C}$ is a finite sequence, then define its **discrete Fourier transform** as:

$$\hat{f}_k := \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Relationship with the Fourier series: Let f be a continuous function defined on the interval $[0, 2\pi)$, and denote by $f_j := f(\frac{2\pi j}{N})$. Then the sum $\frac{2\pi}{N} \cdot \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}}$ is a Riemann sum of the integral $\int_0^{2\pi} f(x) e^{ikx} dx$. Utilizing Euler's formula, we have:

$$\begin{aligned} \frac{1}{N} \hat{f}_k &= \frac{1}{N} \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \approx \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{ikx} dx = \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(x) \cos kx dx + i \cdot \frac{1}{2\pi} \int_0^{2\pi} f(x) \sin kx dx = a_k + ib_k, \end{aligned}$$

where a_k, b_k are the trigonometric Fourier coefficients.

The Discrete Fourier Transform (DFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

If $f_0, f_1, \dots, f_{N-1} \in \mathbf{C}$ is a finite sequence, then define its **discrete Fourier transform** as:

$$\hat{f}_k := \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Relationship with the Fourier series: Let f be a continuous function defined on the interval $[0, 2\pi)$, and denote by $f_j := f(\frac{2\pi j}{N})$. Then the sum $\frac{2\pi}{N} \cdot \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}}$ is a Riemann sum of the integral $\int_0^{2\pi} f(x) e^{ikx} dx$.

Utilizing Euler's formula, we have:

$$\begin{aligned} \frac{1}{N} \hat{f}_k &= \frac{1}{N} \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \approx \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{ikx} dx = \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(x) \cos kx dx + i \cdot \frac{1}{2\pi} \int_0^{2\pi} f(x) \sin kx dx = a_k + ib_k, \end{aligned}$$

where a_k, b_k are the trigonometric Fourier coefficients.

The Discrete Fourier Transform (DFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

If $f_0, f_1, \dots, f_{N-1} \in \mathbf{C}$ is a finite sequence, then define its **discrete Fourier transform** as:

$$\hat{f}_k := \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Relationship with the Fourier series: Let f be a continuous function defined on the interval $[0, 2\pi)$, and denote by $f_j := f(\frac{2\pi j}{N})$. Then the sum $\frac{2\pi}{N} \cdot \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}}$ is a Riemann sum of the integral $\int_0^{2\pi} f(x) e^{ikx} dx$. Utilizing Euler's formula, we have:

$$\begin{aligned} \frac{1}{N} \hat{f}_k &= \frac{1}{N} \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \approx \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{ikx} dx = \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(x) \cos kx dx + i \cdot \frac{1}{2\pi} \int_0^{2\pi} f(x) \sin kx dx = a_k + ib_k, \end{aligned}$$

where a_k, b_k are the trigonometric Fourier coefficients.

The Discrete Fourier Transform (DFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

If $f_0, f_1, \dots, f_{N-1} \in \mathbf{C}$ is a finite sequence, then define its **discrete Fourier transform** as:

$$\hat{f}_k := \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Relationship with the Fourier series: Let f be a continuous function defined on the interval $[0, 2\pi)$, and denote by $f_j := f(\frac{2\pi j}{N})$. Then the sum $\frac{2\pi}{N} \cdot \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}}$ is a Riemann sum of the integral $\int_0^{2\pi} f(x) e^{ikx} dx$. Utilizing Euler's formula, we have:

$$\begin{aligned} \frac{1}{N} \hat{f}_k &= \frac{1}{N} \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \approx \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{ikx} dx = \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(x) \cos kx dx + i \cdot \frac{1}{2\pi} \int_0^{2\pi} f(x) \sin kx dx = a_k + ib_k, \end{aligned}$$

where a_k, b_k are the trigonometric Fourier coefficients.

The inverse Discrete Fourier Transform (iDFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Every finite sequence can uniquely be reconstructed from its DFT, namely:

$$f_k := \frac{1}{N} \cdot \sum_{j=0}^{N-1} \hat{f}_j \cdot e^{-\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

$$\begin{aligned} \frac{1}{N} \sum_{j=0}^{N-1} \hat{f}_j e^{-\frac{2\pi i k j}{N}} &= \frac{1}{N} \sum_{j=0}^{N-1} \sum_{r=0}^{N-1} f_r e^{\frac{2\pi i r j}{N}} e^{-\frac{2\pi i k j}{N}} = \sum_{r=0}^{N-1} f_r \cdot \frac{1}{N} \sum_{j=0}^{N-1} e^{\frac{2\pi i (r-k) j}{N}} \\ &= \sum_{r=0}^{N-1} f_r \cdot \frac{1}{N} \sum_{j=0}^{N-1} z^j \quad (z := e^{\frac{2\pi i (r-k)}{N}}) \end{aligned}$$

If $r = k$, then $z = 1$, therefore the inner sum equals to 1. If $r \neq k$, then the inner sum is the sum of a finite geometric sequence, which equals to 0. This completes the proof.



The inverse Discrete Fourier Transform (iDFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Every finite sequence can uniquely be reconstructed from its DFT, namely:

$$f_k := \frac{1}{N} \cdot \sum_{j=0}^{N-1} \hat{f}_j \cdot e^{-\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

$$\begin{aligned} \frac{1}{N} \sum_{j=0}^{N-1} \hat{f}_j e^{-\frac{2\pi i k j}{N}} &= \frac{1}{N} \sum_{j=0}^{N-1} \sum_{r=0}^{N-1} f_r e^{\frac{2\pi i r j}{N}} e^{-\frac{2\pi i k j}{N}} = \sum_{r=0}^{N-1} f_r \cdot \frac{1}{N} \sum_{j=0}^{N-1} e^{\frac{2\pi i (r-k) j}{N}} \\ &= \sum_{r=0}^{N-1} f_r \cdot \frac{1}{N} \sum_{j=0}^{N-1} z^j \quad (z := e^{\frac{2\pi i (r-k)}{N}}) \end{aligned}$$

If $r = k$, then $z = 1$, therefore the inner sum equals to 1. If $r \neq k$, then the inner sum is the sum of a finite geometric sequence, which equals to 0. This completes the proof.



The inverse Discrete Fourier Transform (iDFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Every finite sequence can uniquely be reconstructed from its DFT, namely:

$$f_k := \frac{1}{N} \cdot \sum_{j=0}^{N-1} \hat{f}_j \cdot e^{-\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

$$\begin{aligned} \frac{1}{N} \sum_{j=0}^{N-1} \hat{f}_j e^{-\frac{2\pi i k j}{N}} &= \frac{1}{N} \sum_{j=0}^{N-1} \sum_{r=0}^{N-1} f_r e^{\frac{2\pi i r j}{N}} e^{-\frac{2\pi i k j}{N}} = \sum_{r=0}^{N-1} f_r \cdot \frac{1}{N} \sum_{j=0}^{N-1} e^{\frac{2\pi i (r-k) j}{N}} \\ &= \sum_{r=0}^{N-1} f_r \cdot \frac{1}{N} \sum_{j=0}^{N-1} z^j \quad (z := e^{\frac{2\pi i (r-k)}{N}}) \end{aligned}$$

If $r = k$, then $z = 1$, therefore the inner sum equals to 1. If $r \neq k$, then the inner sum is the sum of a finite geometric sequence, which equals to 0. This completes the proof.



The Discrete Fourier Transform, numerical features

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Total computational cost: $\mathcal{O}(N^2)$, which is too high!



The Fast Fourier Transform (FFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

**The Fast Fourier
Transform**

Image
compression by
DFT

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. First, let k be a 'small' index: $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_k &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$



The Fast Fourier Transform (FFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. First, let k be a 'small' index: $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_k &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$

The Fast Fourier Transform (FFT)

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. First, let k be a 'small' index: $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_k &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$

The Fast Fourier Transform (FFT)

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. First, let k be a 'small' index: $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_k &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} + e^{\frac{2\pi i k}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$



The Fast Fourier Transform (FFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

**The Fast Fourier
Transform**

Image
compression by
DFT

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. Now consider the 'great' indices having the form $N_1 + k$, where $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_{N_1+k} &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + e^{\frac{2\pi i (N_1+k)}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} - e^{\frac{2\pi i k}{N_1}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$



The Fast Fourier Transform (FFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. Now consider the 'great' indices having the form $N_1 + k$, where $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_{N_1+k} &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + e^{\frac{2\pi i (N_1+k)}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} - e^{\frac{2\pi i k}{N_1}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$

The Fast Fourier Transform (FFT)

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. Now consider the 'great' indices having the form $N_1 + k$, where $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_{N_1+k} &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + e^{\frac{2\pi i (N_1+k)}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} - e^{\frac{2\pi i k}{N_1}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$

The Fast Fourier Transform (FFT)

Denote by $F_N : \mathbf{C}^N \rightarrow \mathbf{C}^N$ the linear operator of the DFT:

$$(F_N f)_k = \sum_{j=0}^{N-1} f_j \cdot e^{\frac{2\pi i k j}{N}} \quad (k = 0, 1, \dots, N-1)$$

Assume that N is even: $N = 2N_1$. Let us separate the terms with even and odd indices in the expression of $(F_N f)_k$. Now consider the 'great' indices having the form $N_1 + k$, where $k = 0, 1, \dots, N_1 - 1$:

$$\begin{aligned} (F_N f)_{N_1+k} &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot (2\ell+1)}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} + e^{\frac{2\pi i (N_1+k)}{N}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i (N_1+k) \cdot 2\ell}{N}} = \\ &= \sum_{\ell=0}^{N_1-1} f_{2\ell} \cdot e^{\frac{2\pi i k \ell}{N_1}} - e^{\frac{2\pi i k}{N_1}} \cdot \sum_{\ell=0}^{N_1-1} f_{2\ell+1} \cdot e^{\frac{2\pi i k \ell}{N_1}} \end{aligned}$$



The Fast Fourier Transform (FFT)

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

**The Fast Fourier
Transform**

Image
compression by
DFT

For both the 'small' and 'great' indices, both sums on the right-hand sides are discrete Fourier transforms with smaller vectors. The procedure can recursively be continued, if N is a power of two.



The Fast Fourier Transform, recursive algorithm

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

With recursive invocations:

- $N_1 := N/2$

$$f^{even} := (f_0, f_2, \dots, f_{2N_1-2}), \quad f^{odd} := (f_1, f_3, \dots, f_{2N_1-1})$$

- $\hat{f}^{even} := F_{N_1} f^{even}, \quad \hat{f}^{odd} := F_{N_1} f^{odd}$

-

$$(F_N f)_k := \hat{f}_k^{even} + e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

$$(F_N f)_{N_1+k} := \hat{f}_k^{even} - e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

where for $N = 1$, $F_1 f := f$ (f has one component only)



The Fast Fourier Transform, recursive algorithm

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

With recursive invocations:

- $N_1 := N/2$

$$f^{even} := (f_0, f_2, \dots, f_{2N_1-2}), \quad f^{odd} := (f_1, f_3, \dots, f_{2N_1-1})$$

- $\hat{f}^{even} := F_{N_1} f^{even}, \quad \hat{f}^{odd} := F_{N_1} f^{odd}$



$$(F_N f)_k := \hat{f}_k^{even} + e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

$$(F_N f)_{N_1+k} := \hat{f}_k^{even} - e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

where for $N = 1$, $F_1 f := f$ (f has one component only)



The Fast Fourier Transform, recursive algorithm

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

With recursive invocations:

- $N_1 := N/2$

$$f^{even} := (f_0, f_2, \dots, f_{2N_1-2}), \quad f^{odd} := (f_1, f_3, \dots, f_{2N_1-1})$$

- $\hat{f}^{even} := F_{N_1} f^{even}, \quad \hat{f}^{odd} := F_{N_1} f^{odd}$

-

$$(F_N f)_k := \hat{f}_k^{even} + e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

$$(F_N f)_{N_1+k} := \hat{f}_k^{even} - e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

where for $N = 1$, $F_1 f := f$ (f has one component only)



The Fast Fourier Transform, recursive algorithm

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$

Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

With recursive invocations:

- $N_1 := N/2$

$$f^{even} := (f_0, f_2, \dots, f_{2N_1-2}), \quad f^{odd} := (f_1, f_3, \dots, f_{2N_1-1})$$

- $\hat{f}^{even} := F_{N_1} f^{even}, \quad \hat{f}^{odd} := F_{N_1} f^{odd}$

-

$$(F_N f)_k := \hat{f}_k^{even} + e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

$$(F_N f)_{N_1+k} := \hat{f}_k^{even} - e^{\frac{2\pi i k}{N}} \cdot \hat{f}_k^{odd} \quad (k = 0, 1, \dots, N_1 - 1)$$

where for $N = 1$, $F_1 f := f$ (f has one component only)



The Fast Fourier Transform, numerical features

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Total computational cost: $\mathcal{O}(N \cdot \log N)$, which is much smaller than the computational cost of the original DFT.

Example:

f_0 f_1 f_2 f_3 f_4 f_5 f_6 f_7

f_0 f_2 f_4 f_6 | f_1 f_3 f_5 f_7

f_0 f_4 | f_2 f_6 | f_1 f_5 | f_3 f_7

f_0 | f_4 | f_2 | f_6 | f_1 | f_5 | f_3 | f_7



The Fast Fourier Transform, numerical features

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Total computational cost: $\mathcal{O}(N \cdot \log N)$, which is much smaller than the computational cost of the original DFT.

Example:

f_0 f_1 f_2 f_3 f_4 f_5 f_6 f_7

f_0 f_2 f_4 f_6 | f_1 f_3 f_5 f_7

f_0 f_4 | f_2 f_6 | f_1 f_5 | f_3 f_7

f_0 | f_4 | f_2 | f_6 | f_1 | f_5 | f_3 | f_7



The Fast Fourier Transform, numerical features

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

Total computational cost: $\mathcal{O}(N \cdot \log N)$, which is much smaller than the computational cost of the original DFT.

Example:

f_0 f_1 f_2 f_3 f_4 f_5 f_6 f_7

f_0 f_2 f_4 f_6 | f_1 f_3 f_5 f_7

f_0 f_4 | f_2 f_6 | f_1 f_5 | f_3 f_7

f_0 | f_4 | f_2 | f_6 | f_1 | f_5 | f_3 | f_7

The 2D Discrete Fourier Transform

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

The DFT of a matrix $f \in \mathbf{M}_{N \times N}$ is the matrix $\hat{f} \in \mathbf{M}_{N \times N}$ with the following entries:

$$\hat{f}_{k,j} := \sum_{r=0}^{N-1} \sum_{s=0}^{N-1} f_{r,s} \cdot e^{\frac{2\pi i k r}{N}} e^{\frac{2\pi i j s}{N}}$$

The algorithm of the computation of the DFT of a matrix:

- For every row of the matrix f , substitute the 1D DFT of the corresponding row.
- For every column of this matrix, substitute the 1D DFT of the corresponding column.

This results in the 2D DFT of the original matrix f .



The 2D Discrete Fourier Transform

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

The DFT of a matrix $f \in \mathbf{M}_{N \times N}$ is the matrix $\hat{f} \in \mathbf{M}_{N \times N}$ with the following entries:

$$\hat{f}_{k,j} := \sum_{r=0}^{N-1} \sum_{s=0}^{N-1} f_{r,s} \cdot e^{\frac{2\pi i k r}{N}} e^{\frac{2\pi i j s}{N}}$$

The algorithm of the computation of the DFT of a matrix:

- For every row of the matrix f , substitute the 1D DFT of the corresponding row.
- For every column of this matrix, substitute the 1D DFT of the corresponding column.

This results in the 2D DFT of the original matrix f .



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform



Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

An N -by- N grayscale image can be considered an N -by- N matrix. In general, *there are a lot of Fourier coefficients that almost equal to 0.*



An 512-by-512 image and its Discrete Fourier Transform



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár

The Fast
Fourier
Transform



Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT

The idea of the image compression: *to keep the Fourier coefficients which are closer to the corners than r , and to set the others to zero.* From the truncated DFT, the image can be approximately reconstructed.



The original and truncated DFT of the image ($r = 100$)



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár


The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT



Original image



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár


The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT



Reconstructed image, $r = 20$



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár


The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT



Reconstructed image, $r = 40$



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár


The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT



Reconstructed image, $r = 60$



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár


The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT



Reconstructed image, $r = 80$



Image compression by DFT

Numerical
Analysis

by Csaba
Gáspár


The Fast
Fourier
Transform

Trigonometric
Fourier series in
 $L_2(0, 2\pi)$
Complex
exponential
function

The Discrete
Fourier
Transform

The Fast Fourier
Transform

Image
compression by
DFT



Reconstructed image, $r = 100$