

# Transcritical Bifurcation

Introducer: Yichen Lu

May 11, 2023

# Contents

## 1 Concept

## 2 Exercises

# Bifurcation

As parameters are varied,

- Fixed points can be created or destroyed,

# Bifurcation

As parameters are varied,

- Fixed points can be created or destroyed,
- or their stability can change (stable  $\leftrightarrow$  unstable).

# Transcritical Bifurcation




- A fixed point must exist for all values of a parameter and can never be destroyed

# Transcritical Bifurcation

- A fixed point must exist for all values of a parameter and can never be destroyed
- Such a fixed point may change its stability as the parameter is varied

# Transcritical Bifurcation

- A fixed point must exist for all values of a parameter and can never be destroyed
- Such a fixed point may change its stability as the parameter is varied

(a)  $r < 0$ (b)  $r = 0$ (c)  $r > 0$

# Contents

1 Concept

2 Exercises



## Exercises 3.2.1


Sketch all the qualitatively different vector fields that occur as  $r$  is varied. Show that a transcritical bifurcation occurs at a critical value of  $r$ , to be determined. Finally, sketch the bifurcation diagram of fixed points  $x^*$  vs.  $r$ .

Consider the following system of ODEs:

$$\dot{x} = rx + x^2$$


Note that  $x = 0$  is a fixed point for all value of  $r$

All the qualitatively different vector fields that occur as  $r$  is varied:




Transcritical bifurcation occurs at  $r = 0$ .

Bifurcation diagram of fixed points  $x^*$  vs.  $r$  :



## Exercises 3.2.5



According to the law of mass action of chemical kinetics, the rate of an elementary reaction is proportional to the product of the concentrations of the reactants. Then the equation for the kinetics of  $x$  is

$$\dot{x} = k_1 a x - k_{-1} x^2 - k_2 b x$$

$$\dot{x} = (k_1 a - k_2 b) x - k_{-1} x^2$$

$$\dot{x} = c_1 x - c_2 x^2$$

, where  $c_1 = k_1 a - k_2 b$  and  $c_2 = k_{-1}$ .

For  $c_1 < 0$ ,  $x^* = 0$  become stable, so  $k_1 a < k_2 b$

Thank you for listening!