

Radii of Emergent Patterns in Swarmalator Systems

Publisher: IEEE

Cite This

easyScholar文献收藏

PDF

Udo Schilcher; Christoph W. Rauter; Christian Bettstetter [All Authors](#)

27

Full

Text Views



Abstract

Document Sections

1. Introduction
2. System Model
3. Derivation of Radii
4. Numerical Evaluation
5. Conclusions and Outlook

Abstract:

For a system of swarmalators converging to different types of circular patterns, we provide expressions for the outer and inner radii of these patterns and examine their dependence on the model parameters. Derivations are made for three static patterns with an infinite number of entities and a generalized swarmalator model with parameterized attraction and repulsion kernels. Simulations of finite systems show good agreement with the asymptotic expressions.

Published in: 2023 IEEE International Conference on Autonomic Computing and Self-Organizing Systems (ACSOS)

Date of Conference: 25-29 September 2023

DOI: 10.1109/ACSOS58161.2023.00034

Model equations:

$$\begin{aligned}\dot{x}_i &= \frac{1}{N} \sum_{j \neq i}^N e_{ij} d_{ij}^{\alpha} (1 + J \cos \theta_{ij}) - e_{ij} d_{ij}^{\beta} \\ \dot{\theta}_i &= \frac{K}{N} \sum_{j \neq i}^N d_{ij}^{\gamma} \sin \theta_{ij}\end{aligned}$$


where $d_{ij} = \|x_j - x_i\|$, $\theta_{ij} = \theta_i - \theta_j$, $e_{ij} = \frac{1}{d_{ij}}(x_j - x_i)$.

Choosing $\alpha = 0$ and $\beta = \gamma = -1$ gives the original model.

In the converged state of the static patterns, the entities no longer move, so we have $\dot{x}_i = 0$, which yields:

$$r_{\text{out}}^{\alpha} \underbrace{\sum_{j \neq i}^N e_{ij} g_{ij}^{\alpha} (1 + J \cos \theta_{ij})}_{\text{Attraction } A} = r_{\text{out}}^{\beta} \underbrace{\sum_{j \neq i}^N e_{ij} g_{ij}^{\alpha}}_{\text{Repulsion } R}$$

where $r_{\text{out}} = \max_{1 \leq i \leq N} \|x_i - \bar{x}\|$, $g_{ij} = \frac{d_{ij}}{r_{\text{out}}}$.



- Pattern center \bar{x} is positioned at $(1, 0)$ and a swarmalator is present at the origin.
- using polar coordinates (r, ϕ) , hence $e_{ij} = (\cos \phi, \sin \phi)$ and $g_{ij} = r$.
- $l_{O1}(\phi) = 2 \cos \phi = \frac{2 \sin \phi \cos \phi}{\sin \phi} = \frac{\sin 2\phi}{\sin \phi}$

(a) Calculating the length $l_{O1}(\phi)$.

Due to symmetry, $\sum_{j \neq i}^N e_{ij,y} g_{ij}^\alpha = 0$, $\sum_{j \neq i}^N e_{ij,y} g_{ij}^\beta = 0$, which yields

$$r_{\text{out}}^\alpha \sum_{j \neq i}^N e_{ij,x} g_{ij}^\alpha (1 + J \cos \theta_{ij}) = r_{\text{out}}^\beta \sum_{j \neq i}^N e_{ij,x} g_{ij}^\alpha$$

Next, the authors consider the limiting case $N \rightarrow \infty$ and rewrite the sums as integrals ($dx dy = r dr d\phi$)

$$r_{\text{out}}^{\alpha-\beta} = \frac{\text{Repulsion}}{\text{Attraction}} = \frac{\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos \phi \int_0^{l_{O1}(\phi)} r^{1+\beta} dr d\phi}{\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos \phi \int_0^{l_{O1}(\phi)} (1 + J \cos \theta_{ij}) r^{1+\alpha} dr d\phi}$$

The solution in the first integration:

$$\int_0^{l_{O1}(\phi)} r^{1+m} dr = \frac{1}{2+m} l_{O1}^{2+m}(\phi) = \frac{(2 \cos \phi)^{2+m}}{2+m}$$

Supposed θ_{ij} is a constant, we have

$$\begin{aligned} r_{\text{out}}^{\alpha-\beta} &= \frac{\frac{2^{2+\beta}}{2+\beta} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^{3+\beta} \phi d\phi}{\frac{(1+J \cos \theta_{ij}) 2^{2+\alpha}}{2+\alpha} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^{3+\alpha} \phi d\phi} \\ &= \frac{2^{\beta-\alpha} (2+\alpha) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^{3+\beta} \phi d\phi}{(1+J \cos \theta_{ij}) (2+\beta) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^{3+\alpha} \phi d\phi} \end{aligned}$$

- **Static Sync.** In Static Sync State, $J \cos \theta_{ij} = J \forall (i, j)$, For $\alpha = 0$ and $\beta = -1$, we have

$$r_{\text{out}} = \frac{\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos \phi \frac{\sin(2\phi)}{\sin \phi} d\phi}{(1 + J) \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos \phi \frac{\sin^2(2\phi)}{\sin^2 \phi} d\phi} = \frac{3\pi}{8(1 + J)}$$

- **Static Async.** In Static Async State, $J \cos \theta_{ij} = 0 \forall (i, j)$, For $\alpha = 0$ and $\beta = -1$, we have

$$r_{\text{out}} = \frac{\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos \phi \frac{\sin(2\phi)}{\sin \phi} d\phi}{\frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos \phi \frac{\sin^2(2\phi)}{\sin^2 \phi} d\phi} = \frac{3\pi}{8} \approx 1.178$$