

# Real Functions in Several Variables: Volume IX

Formation of Integrals and Improper Integrals

Leif Mejlbro



Leif Mejlbro

# **Real Functions in Several Variables**

## Volume IX Transformation of Integrals and Improper Integrals



Real Functions in Several Variables: Volume IX Transformation of Integrals and  
Improper Integrals

2<sup>nd</sup> edition

© 2015 Leif Mejlbro & [bookboon.com](http://bookboon.com)

ISBN 978-87-403-0916-4

## Contents

<b>Volume I, Point Sets in <math>\mathbb{R}^n</math></b>	<b>1</b>
Preface	15
<b>Introduction to volume I, Point sets in <math>\mathbb{R}^n</math>. The maximal domain of a function</b>	<b>19</b>
<b>1 Basic concepts</b>	<b>21</b>
1.1 Introduction .....	21
1.2 The real linear space $\mathbb{R}^n$ .....	22
1.3 The vector product .....	26
1.4 The most commonly used coordinate systems .....	29
1.5 Point sets in space .....	37
1.5.1 Interior, exterior and boundary of a set .....	37
1.5.2 Starshaped and convex sets .....	40
1.5.3 Catalogue of frequently used point sets in the plane and the space .....	41
1.6 Quadratic equations in two or three variables. Conic sections .....	47
1.6.1 Quadratic equations in two variables. Conic sections .....	47
1.6.2 Quadratic equations in three variables. Conic sectional surfaces .....	54
1.6.3 Summary of the canonical cases in three variables .....	66
<b>2 Some useful procedures</b>	<b>67</b>
2.1 Introduction .....	67
2.2 Integration of trigonometric polynomials .....	67
2.3 Complex decomposition of a fraction of two polynomials .....	69
2.4 Integration of a fraction of two polynomials .....	72
<b>3 Examples of point sets</b>	<b>75</b>
3.1 Point sets .....	75
3.2 Conics and conical sections .....	104
<b>4 Formulae</b>	<b>115</b>
4.1 Squares etc. .....	115
4.2 Powers etc. .....	115
4.3 Differentiation .....	116
4.4 Special derivatives .....	116
4.5 Integration .....	118
4.6 Special antiderivatives .....	119
4.7 Trigonometric formulae .....	121
4.8 Hyperbolic formulae .....	123
4.9 Complex transformation formulae .....	124
4.10 Taylor expansions .....	124
4.11 Magnitudes of functions .....	125
<b>Index</b>	<b>127</b>

<b>Volume II, Continuous Functions in Several Variables</b>	<b>133</b>
Preface	147
Introduction to volume II, Continuous Functions in Several Variables	151
<b>5 Continuous functions in several variables</b>	<b>153</b>
5.1 Maps in general .....	153
5.2 Functions in several variables .....	154
5.3 Vector functions .....	157
5.4 Visualization of functions .....	158
5.5 Implicit given function .....	161
5.6 Limits and continuity .....	162
5.7 Continuous functions .....	168
5.8 Continuous curves .....	170
5.8.1 Parametric description .....	170
5.8.2 Change of parameter of a curve .....	174
5.9 Connectedness .....	175
5.10 Continuous surfaces in $\mathbb{R}^3$ .....	177
5.10.1 Parametric description and continuity .....	177
5.10.2 Cylindric surfaces .....	180
5.10.3 Surfaces of revolution .....	181
5.10.4 Boundary curves, closed surface and orientation of surfaces .....	182
5.11 Main theorems for continuous functions .....	185
<b>6 A useful procedure</b>	<b>189</b>
6.1 The domain of a function .....	189
<b>7 Examples of continuous functions in several variables</b>	<b>191</b>
7.1 Maximal domain of a function .....	191
7.2 Level curves and level surfaces .....	198
7.3 Continuous functions .....	212
7.4 Description of curves .....	227
7.5 Connected sets .....	241
7.6 Description of surfaces .....	245
<b>8 Formulae</b>	<b>257</b>
8.1 Squares etc. .....	257
8.2 Powers etc. .....	257
8.3 Differentiation .....	258
8.4 Special derivatives .....	258
8.5 Integration .....	260
8.6 Special antiderivatives .....	261
8.7 Trigonometric formulæ .....	263
8.8 Hyperbolic formulæ .....	265
8.9 Complex transformation formulæ .....	266
8.10 Taylor expansions .....	266
8.11 Magnitudes of functions .....	267
<b>Index</b>	<b>269</b>

<b>Volume III, Differentiable Functions in Several Variables</b>	<b>275</b>
Preface	289
Introduction to volume III, Differentiable Functions in Several Variables	293
<b>9 Differentiable functions in several variables</b>	<b>295</b>
9.1 Differentiability .....	295
9.1.1 The gradient and the differential .....	295
9.1.2 Partial derivatives .....	298
9.1.3 Differentiable vector functions .....	303
9.1.4 The approximating polynomial of degree 1 .....	304
9.2 The chain rule .....	305
9.2.1 The elementary chain rule .....	305
9.2.2 The first special case .....	308
9.2.3 The second special case .....	309
9.2.4 The third special case .....	310
9.2.5 The general chain rule .....	314
9.3 Directional derivative .....	317
9.4 $C^n$ -functions .....	318
9.5 Taylor's formula .....	321
9.5.1 Taylor's formula in one dimension .....	321
9.5.2 Taylor expansion of order 1 .....	322
9.5.3 Taylor expansion of order 2 in the plane .....	323
9.5.4 The approximating polynomial .....	326
<b>10 Some useful procedures</b>	<b>333</b>
10.1 Introduction .....	333
10.2 The chain rule .....	333
10.3 Calculation of the directional derivative .....	334
10.4 Approximating polynomials .....	336
<b>11 Examples of differentiable functions</b>	<b>339</b>
11.1 Gradient .....	339
11.2 The chain rule .....	352
11.3 Directional derivative .....	375
11.4 Partial derivatives of higher order .....	382
11.5 Taylor's formula for functions of several variables .....	404
<b>12 Formulæ</b>	<b>445</b>
12.1 Squares etc. ....	445
12.2 Powers etc. ....	445
12.3 Differentiation .....	446
12.4 Special derivatives .....	446
12.5 Integration .....	448
12.6 Special antiderivatives .....	449
12.7 Trigonometric formulæ .....	451
12.8 Hyperbolic formulæ .....	453
12.9 Complex transformation formulæ .....	454
12.10 Taylor expansions .....	454
12.11 Magnitudes of functions .....	455
<b>Index</b>	<b>457</b>

<b>Volume IV, Differentiable Functions in Several Variables</b>	<b>463</b>
Preface	477
Introduction to volume IV, Curves and Surfaces	481
<b>13 Differentiable curves and surfaces, and line integrals in several variables</b>	<b>483</b>
13.1 Introduction .....	483
13.2 Differentiable curves .....	483
13.3 Level curves .....	492
13.4 Differentiable surfaces .....	495
13.5 Special $C^1$ -surfaces .....	499
13.6 Level surfaces .....	503
<b>14 Examples of tangents (curves) and tangent planes (surfaces)</b>	<b>505</b>
14.1 Examples of tangents to curves .....	505
14.2 Examples of tangent planes to a surface .....	520
<b>15 Formulæ</b>	<b>541</b>
15.1 Squares etc. .....	541
15.2 Powers etc. .....	541
15.3 Differentiation .....	542
15.4 Special derivatives .....	542
15.5 Integration .....	544
15.6 Special antiderivatives .....	545
15.7 Trigonometric formulæ .....	547
15.8 Hyperbolic formulæ .....	549
15.9 Complex transformation formulæ .....	550
15.10 Taylor expansions .....	550
15.11 Magnitudes of functions .....	551
<b>Index</b>	<b>553</b>
<b>Volume V, Differentiable Functions in Several Variables</b>	<b>559</b>
Preface	573
Introduction to volume V, The range of a function, Extrema of a Function in Several Variables	577
<b>16 The range of a function</b>	<b>579</b>
16.1 Introduction .....	579
16.2 Global extrema of a continuous function .....	581
16.2.1 A necessary condition .....	581
16.2.2 The case of a closed and bounded domain of $f$ .....	583
16.2.3 The case of a bounded but not closed domain of $f$ .....	599
16.2.4 The case of an unbounded domain of $f$ .....	608
16.3 Local extrema of a continuous function .....	611
16.3.1 Local extrema in general .....	611
16.3.2 Application of Taylor's formula .....	616
16.4 Extremum for continuous functions in three or more variables .....	625
<b>17 Examples of global and local extrema</b>	<b>631</b>
17.1 MAPLE .....	631
17.2 Examples of extremum for two variables .....	632
17.3 Examples of extremum for three variables .....	668

17.4 Examples of maxima and minima .....	677
17.5 Examples of ranges of functions .....	769
<b>18 Formulæ</b>	<b>811</b>
18.1 Squares etc. ....	811
18.2 Powers etc. ....	811
18.3 Differentiation ....	812
18.4 Special derivatives ....	812
18.5 Integration ....	814
18.6 Special antiderivatives ....	815
18.7 Trigonometric formulæ ....	817
18.8 Hyperbolic formulæ ....	819
18.9 Complex transformation formulæ ....	820
18.10 Taylor expansions ....	820
18.11 Magnitudes of functions ....	821
<b>Index</b>	<b>823</b>
<b>Volume VI, Antiderivatives and Plane Integrals</b>	<b>829</b>
<b>Preface</b>	<b>841</b>
<b>Introduction to volume VI, Integration of a function in several variables</b>	<b>845</b>
<b>19 Antiderivatives of functions in several variables</b>	<b>847</b>
19.1 The theory of antiderivatives of functions in several variables .....	847
19.2 Templates for gradient fields and antiderivatives of functions in three variables .....	858
19.3 Examples of gradient fields and antiderivatives .....	863
<b>20 Integration in the plane</b>	<b>881</b>
20.1 An overview of integration in the plane and in the space .....	881
20.2 Introduction .....	882
20.3 The plane integral in rectangular coordinates .....	887
20.3.1 Reduction in rectangular coordinates .....	887
20.3.2 The colour code, and a procedure of calculating a plane integral .....	890
20.4 Examples of the plane integral in rectangular coordinates .....	894
20.5 The plane integral in polar coordinates .....	936
20.6 Procedure of reduction of the plane integral; polar version .....	944
20.7 Examples of the plane integral in polar coordinates .....	948
20.8 Examples of area in polar coordinates .....	972
<b>21 Formulæ</b>	<b>977</b>
21.1 Squares etc. ....	977
21.2 Powers etc. ....	977
21.3 Differentiation ....	978
21.4 Special derivatives ....	978
21.5 Integration ....	980
21.6 Special antiderivatives ....	981
21.7 Trigonometric formulæ ....	983
21.8 Hyperbolic formulæ ....	985
21.9 Complex transformation formulæ ....	986
21.10 Taylor expansions ....	986
21.11 Magnitudes of functions ....	987
<b>Index</b>	<b>989</b>

<b>Volume VII, Space Integrals</b>	<b>995</b>
Preface	1009
Introduction to volume VII, The space integral	1013
<b>22 The space integral in rectangular coordinates</b>	<b>1015</b>
22.1 Introduction .....	1015
22.2 Overview of setting up of a line, a plane, a surface or a space integral .....	1015
22.3 Reduction theorems in rectangular coordinates .....	1021
22.4 Procedure for reduction of space integral in rectangular coordinates .....	1024
22.5 Examples of space integrals in rectangular coordinates .....	1026
<b>23 The space integral in semi-polar coordinates</b>	<b>1055</b>
23.1 Reduction theorem in semi-polar coordinates .....	1055
23.2 Procedures for reduction of space integral in semi-polar coordinates .....	1056
23.3 Examples of space integrals in semi-polar coordinates .....	1058
<b>24 The space integral in spherical coordinates</b>	<b>1081</b>
24.1 Reduction theorem in spherical coordinates .....	1081
24.2 Procedures for reduction of space integral in spherical coordinates .....	1082
24.3 Examples of space integrals in spherical coordinates .....	1084
24.4 Examples of volumes .....	1107
24.5 Examples of moments of inertia and centres of gravity .....	1116
<b>25 Formulæ</b>	<b>1125</b>
25.1 Squares etc. ....	1125
25.2 Powers etc. ....	1125
25.3 Differentiation .....	1126
25.4 Special derivatives .....	1126
25.5 Integration .....	1128
25.6 Special antiderivatives .....	1129
25.7 Trigonometric formulæ .....	1131
25.8 Hyperbolic formulæ .....	1133
25.9 Complex transformation formulæ .....	1134
25.10 Taylor expansions .....	1134
25.11 Magnitudes of functions .....	1135
<b>Index</b>	<b>1137</b>
<b>Volume VIII, Line Integrals and Surface Integrals</b>	<b>1143</b>
Preface	1157
Introduction to volume VIII, The line integral and the surface integral	1161
<b>26 The line integral</b>	<b>1163</b>
26.1 Introduction .....	1163
26.2 Reduction theorem of the line integral .....	1163
26.2.1 Natural parametric description .....	1166
26.3 Procedures for reduction of a line integral .....	1167
26.4 Examples of the line integral in rectangular coordinates .....	1168
26.5 Examples of the line integral in polar coordinates .....	1190
26.6 Examples of arc lengths and parametric descriptions by the arc length .....	1201

<b>27 The surface integral</b>	<b>1227</b>
27.1 The reduction theorem for a surface integral .....	1227
27.1.1 The integral over the graph of a function in two variables .....	1229
27.1.2 The integral over a cylindric surface .....	1230
27.1.3 The integral over a surface of revolution .....	1232
27.2 Procedures for reduction of a surface integral .....	1233
27.3 Examples of surface integrals .....	1235
27.4 Examples of surface area .....	1296
<b>28 Formulæ</b>	<b>1315</b>
28.1 Squares etc. ....	1315
28.2 Powers etc. ....	1315
28.3 Differentiation ....	1316
28.4 Special derivatives ....	1316
28.5 Integration ....	1318
28.6 Special antiderivatives ....	1319
28.7 Trigonometric formulæ ....	1321
28.8 Hyperbolic formulæ ....	1323
28.9 Complex transformation formulæ ....	1324
28.10 Taylor expansions ....	1324
28.11 Magnitudes of functions ....	1325
<b>Index</b>	<b>1327</b>

[www.sylvania.com](http://www.sylvania.com)

We do not reinvent the wheel we reinvent light.

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM

**OSRAM**  
**SYLVANIA**

Click on the ad to read more

<b>Volume IX, Transformation formulæ and improper integrals</b>	<b>1333</b>
<b>Preface</b>	<b>1347</b>
<b>Introduction to volume IX, Transformation formulæ and improper integrals</b>	<b>1351</b>
<b>29 Transformation of plane and space integrals</b>	<b>1353</b>
29.1 Transformation of a plane integral .....	1353
29.2 Transformation of a space integral .....	1355
29.3 Procedures for the transformation of plane or space integrals .....	1358
29.4 Examples of transformation of plane and space integrals .....	1359
<b>30 Improper integrals</b>	<b>1411</b>
30.1 Introduction .....	1411
30.2 Theorems for improper integrals .....	1413
30.3 Procedure for improper integrals; bounded domain .....	1415
30.4 Procedure for improper integrals; unbounded domain .....	1417
30.5 Examples of improper integrals .....	1418
<b>31 Formulae</b>	<b>1447</b>
31.1 Squares etc. .....	1447
31.2 Powers etc. .....	1447
31.3 Differentiation .....	1448
31.4 Special derivatives .....	1448
31.5 Integration .....	1450
31.6 Special antiderivatives .....	1451
31.7 Trigonometric formulæ .....	1453
31.8 Hyperbolic formulæ .....	1455
31.9 Complex transformation formulæ .....	1456
31.10 Taylor expansions .....	1456
31.11 Magnitudes of functions .....	1457
<b>Index</b>	<b>1459</b>
<b>Volume X, Vector Fields I; Gauß's Theorem</b>	<b>1465</b>
<b>Preface</b>	<b>1479</b>
<b>Introduction to volume X, Vector fields; Gauß's Theorem</b>	<b>1483</b>
<b>32 Tangential line integrals</b>	<b>1485</b>
32.1 Introduction .....	1485
32.2 The tangential line integral. Gradient fields. ....	1485
32.3 Tangential line integrals in Physics .....	1498
32.4 Overview of the theorems and methods concerning tangential line integrals and gradient fields .....	1499
32.5 Examples of tangential line integrals .....	1502
<b>33 Flux and divergence of a vector field. Gauß's theorem</b>	<b>1535</b>
33.1 Flux .....	1535
33.2 Divergence and Gauß's theorem .....	1540
33.3 Applications in Physics .....	1544
33.3.1 Magnetic flux .....	1544
33.3.2 Coulomb vector field .....	1545
33.3.3 Continuity equation .....	1548
33.4 Procedures for flux and divergence of a vector field; Gauß's theorem .....	1549
33.4.1 Procedure for calculation of a flux .....	1549
33.4.2 Application of Gauß's theorem .....	1549
33.5 Examples of flux and divergence of a vector field; Gauß's theorem .....	1551
33.5.1 Examples of calculation of the flux .....	1551
33.5.2 Examples of application of Gauß's theorem .....	1580

<b>34 Formulæ</b>	<b>1619</b>
34.1 Squares etc.	1619
34.2 Powers etc.	1619
34.3 Differentiation	1620
34.4 Special derivatives	1620
34.5 Integration	1622
34.6 Special antiderivatives	1623
34.7 Trigonometric formulæ	1625
34.8 Hyperbolic formulæ	1627
34.9 Complex transformation formulæ	1628
34.10 Taylor expansions	1628
34.11 Magnitudes of functions	1629
<b>Index</b>	<b>1631</b>
<b>Volume XI, Vector Fields II; Stokes's Theorem</b>	<b>1637</b>
<b>Preface</b>	<b>1651</b>
<b>Introduction to volume XI, Vector fields II; Stokes's Theorem; nabla calculus</b>	<b>1655</b>
<b>35 Rotation of a vector field; Stokes's theorem</b>	<b>1657</b>
35.1 Rotation of a vector field in $\mathbb{R}^3$	1657
35.2 Stokes's theorem	1661
35.3 Maxwell's equations	1669
35.3.1 The electrostatic field	1669
35.3.2 The magnostatic field	1671
35.3.3 Summary of Maxwell's equations	1679
35.4 Procedure for the calculation of the rotation of a vector field and applications of Stokes's theorem	1682



**Deloitte.**

Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

© Deloitte & Touche LLP and affiliated entities.



Click on the ad to read more

35.5 Examples of the calculation of the rotation of a vector field and applications of Stokes's theorem .....	1684
35.5.1 Examples of divergence and rotation of a vector field .....	1684
35.5.2 General examples .....	1691
35.5.3 Examples of applications of Stokes's theorem .....	1700
<b>36 Nabla calculus</b>	<b>1739</b>
36.1 The vectorial differential operator $\nabla$ .....	1739
36.2 Differentiation of products .....	1741
36.3 Differentiation of second order .....	1743
36.4 Nabla applied on $\mathbf{x}$ .....	1745
36.5 The integral theorems .....	1746
36.6 Partial integration .....	1749
36.7 Overview of Nabla calculus .....	1750
36.8 Overview of partial integration in higher dimensions .....	1752
36.9 Examples in nabla calculus .....	1754
<b>37 Formulæ</b>	<b>1769</b>
37.1 Squares etc. ....	1769
37.2 Powers etc. ....	1769
37.3 Differentiation .....	1770
37.4 Special derivatives .....	1770
37.5 Integration .....	1772
37.6 Special antiderivatives .....	1773
37.7 Trigonometric formulæ .....	1775
37.8 Hyperbolic formulæ .....	1777
37.9 Complex transformation formulæ .....	1778
37.10 Taylor expansions .....	1778
37.11 Magnitudes of functions .....	1779
<b>Index</b>	<b>1781</b>
<b>Volume XII, Vector Fields III; Potentials, Harmonic Functions and Green's Identities</b>	<b>1787</b>
<b>Preface</b>	<b>1801</b>
<b>Introduction to volume XII, Vector fields III; Potentials, Harmonic Functions and Green's Identities</b>	<b>1805</b>
<b>38 Potentials</b>	<b>1807</b>
38.1 Definitions of scalar and vectorial potentials .....	1807
38.2 A vector field given by its rotation and divergence .....	1813
38.3 Some applications in Physics .....	1816
38.4 Examples from Electromagnetism .....	1819
38.5 Scalar and vector potentials .....	1838
<b>39 Harmonic functions and Green's identities</b>	<b>1889</b>
39.1 Harmonic functions .....	1889
39.2 Green's first identity .....	1890
39.3 Green's second identity .....	1891
39.4 Green's third identity .....	1896
39.5 Green's identities in the plane .....	1898
39.6 Gradient, divergence and rotation in semi-polar and spherical coordinates .....	1899
39.7 Examples of applications of Green's identities .....	1901
39.8 Overview of Green's theorems in the plane .....	1909
39.9 Miscellaneous examples .....	1910

<b>40 Formulæ</b>	<b>1923</b>
40.1 Squares etc.	1923
40.2 Powers etc.	1923
40.3 Differentiation	1924
40.4 Special derivatives	1924
40.5 Integration	1926
40.6 Special antiderivatives	1927
40.7 Trigonometric formulæ	1929
40.8 Hyperbolic formulæ	1931
40.9 Complex transformation formulæ	1932
40.10 Taylor expansions	1932
40.11 Magnitudes of functions	1933
<b>Index</b>	<b>1935</b>

**WHY  
WAIT FOR  
PROGRESS?**

**DARE TO DISCOVER**

---

Discovery means many different things at Schlumberger. But it's the spirit that unites every single one of us. It doesn't matter whether they join our business, engineering or technology teams, our trainees push boundaries, break new ground and deliver the exceptional. If that excites you, then we want to hear from you.

**Dare to discover.**  
[careers.slb.com/recentgraduates](http://careers.slb.com/recentgraduates)



**Schlumberger**

## Preface

The topic of this series of books on “*Real Functions in Several Variables*” is very important in the description in e.g. *Mechanics* of the real 3-dimensional world that we live in. Therefore, we start from the very beginning, modelling this world by using the coordinates of  $\mathbb{R}^3$  to describe e.g. a motion in space. There is, however, absolutely no reason to restrict ourselves to  $\mathbb{R}^3$  alone. Some motions may be rectilinear, so only  $\mathbb{R}$  is needed to describe their movements on a line segment. This opens up for also dealing with  $\mathbb{R}^2$ , when we consider plane motions. In more elaborate problems we need higher dimensional spaces. This may be the case in *Probability Theory* and *Statistics*. Therefore, we shall in general use  $\mathbb{R}^n$  as our abstract model, and then restrict ourselves in examples mainly to  $\mathbb{R}^2$  and  $\mathbb{R}^3$ .

For rectilinear motions the familiar *rectangular coordinate system* is the most convenient one to apply. However, as known from e.g. *Mechanics*, circular motions are also very important in the applications in engineering. It becomes natural alternatively to apply in  $\mathbb{R}^2$  the so-called *polar coordinates* in the plane. They are convenient to describe a circle, where the rectangular coordinates usually give some nasty square roots, which are difficult to handle in practice.

Rectangular coordinates and polar coordinates are designed to model each their problems. They supplement each other, so difficult computations in one of these coordinate systems may be easy, and even trivial, in the other one. It is therefore important always in advance carefully to analyze the geometry of e.g. a domain, so we ask the question: Is this domain best described in rectangular or in polar coordinates?

Sometimes one may split a problem into two subproblems, where we apply rectangular coordinates in one of them and polar coordinates in the other one.

It should be mentioned that in *real life* (though not in these books) one cannot always split a problem into two subproblems as above. Then one is really in trouble, and more advanced mathematical methods should be applied instead. This is, however, outside the scope of the present series of books.

The idea of polar coordinates can be extended in two ways to  $\mathbb{R}^3$ . Either to *semi-polar* or *cylindric coordinates*, which are designed to describe a cylinder, or to *spherical coordinates*, which are excellent for describing spheres, where rectangular coordinates usually are doomed to fail. We use them already in daily life, when we specify a place on Earth by its longitude and latitude! It would be very awkward in this case to use rectangular coordinates instead, even if it is possible.

Concerning the contents, we begin this investigation by modelling point sets in an  $n$ -dimensional Euclidean space  $E^n$  by  $\mathbb{R}^n$ . There is a subtle difference between  $E^n$  and  $\mathbb{R}^n$ , although we often identify these two spaces. In  $E^n$  we use *geometrical methods* without a coordinate system, so the objects are independent of such a choice. In the coordinate space  $\mathbb{R}^n$  we can use ordinary calculus, which in principle is not possible in  $E^n$ . In order to stress this point, we call  $E^n$  the “abstract space” (in the sense of calculus; not in the sense of geometry) as a warning to the reader. Also, whenever necessary, we use the colour black in the “abstract space”, in order to stress that this expression is theoretical, while variables given in a chosen coordinate system and their related concepts are given the colours blue, red and green.

We also include the most basic of what mathematicians call *Topology*, which will be necessary in the following. We describe what we need by a function.

Then we proceed with limits and continuity of functions and define continuous curves and surfaces, with parameters from subsets of  $\mathbb{R}$  and  $\mathbb{R}^2$ , resp..

Continue with (partial) differentiable functions, curves and surfaces, the chain rule and Taylor's formula for functions in several variables.

We deal with maxima and minima and extrema of functions in several variables over a domain in  $\mathbb{R}^n$ . This is a very important subject, so there are given many worked examples to illustrate the theory.

Then we turn to the problems of integration, where we specify four different types with increasing complexity, *plane integral, space integral, curve (or line) integral and surface integral*.

Finally, we consider *vector analysis*, where we deal with vector fields, Gauß's theorem and Stokes's theorem. All these subjects are very important in theoretical Physics.

The structure of this series of books is that each subject is *usually* (but not always) described by three successive chapters. In the first chapter a brief theoretical theory is given. The next chapter gives some practical guidelines of how to solve problems connected with the subject under consideration. Finally, some worked out examples are given, in many cases in several variants, because the standard solution method is seldom the only way, and it may even be clumsy compared with other possibilities.

I have as far as possible structured the examples according to the following scheme:

**A Awareness**, i.e. a short description of what is the problem.

**D Decision**, i.e. a reflection over what should be done with the problem.

**I Implementation**, i.e. where all the calculations are made.

**C Control**, i.e. a test of the result.

This is an ideal form of a general procedure of solution. It can be used in any situation and it is not linked to Mathematics alone. I learned it many years ago in the Theory of Telecommunication in a situation which did not contain Mathematics at all. The student is recommended to use it also in other disciplines.

From high school one is used to immediately to proceed to **I. Implementation**. However, examples and problems at university level, let alone situations in real life, are often so complicated that it in general will be a good investment also to spend some time on the first two points above in order to be absolutely certain of what to do in a particular case. Note that the first three points, **ADI**, can always be executed.

This is unfortunately not the case with **C Control**, because it from now on may be difficult, if possible, to check one's solution. It is only an extra securing whenever it is possible, but we cannot include it always in our solution form above.

I shall on purpose not use the logical signs. These should in general be avoided in Calculus as a shorthand, because they are often (too often, I would say) misused. Instead of  $\wedge$  I shall either write "and", or a comma, and instead of  $\vee$  I shall write "or". The arrows  $\Rightarrow$  and  $\Leftrightarrow$  are in particular misunderstood by the students, so they should be totally avoided. They are not telegram short hands, and from a logical point of view they usually do not make sense at all! Instead, write in a plain language what you mean or want to do. This is difficult in the beginning, but after some practice it becomes routine, and it will give more precise information.

When we deal with multiple integrals, one of the possible pedagogical ways of solving problems has been to colour variables, integrals and upper and lower bounds in blue, red and green, so the reader by the colour code can see in each integral what is the variable, and what are the parameters, which

do not enter the integration under consideration. We shall of course build up a hierarchy of these colours, so the order of integration will always be defined. As already mentioned above we reserve the colour black for the theoretical expressions, where we cannot use ordinary calculus, because the symbols are only shorthand for a concept.

The author has been very grateful to his old friend and colleague, the late Per Wennerberg Karlsson, for many discussions of how to present these difficult topics on real functions in several variables, and for his permission to use his textbook as a template of this present series. Nevertheless, the author has felt it necessary to make quite a few changes compared with the old textbook, because we did not always agree, and some of the topics could also be explained in another way, and then of course the results of our discussions have here been put in writing for the first time.

The author also adds some calculations in MAPLE, which interact nicely with the theoretic text. Note, however, that when one applies MAPLE, one is forced first to make a geometrical analysis of the domain of integration, i.e. apply some of the techniques developed in the present books.

The theory and methods of these volumes on "Real Functions in Several Variables" are applied constantly in higher Mathematics, Mechanics and Engineering Sciences. It is of paramount importance for the calculations in *Probability Theory*, where one constantly integrate over some point set in space.

It is my hope that this text, these guidelines and these examples, of which many are treated in more ways to show that the solutions procedures are not unique, may be of some inspiration for the students who have just started their studies at the universities.

Finally, even if I have tried to write as careful as possible, I doubt that all errors have been removed. I hope that the reader will forgive me the unavoidable errors.

Leif Mejlbro  
March 21, 2015

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



Click on the ad to read more



## Introduction to volume IX, Transformation formulæ and improper integrals

This is the ninth volume in the series of books on *Real Functions in Several Variables*.

In Chapter 29 we investigate how to change variables in the integrals in the plane or space. It is shown that the previous chapters are special cases of this general theory. In particular, we obtain a new introduction of the various weight functions. Formally, Chapter 29 would suffice for the theory, but we have chosen for pedagogical reasons to describe separately the plane integral in rectangular or polar coordinates, the space integral in rectangular, semi-polar or spherical coordinates, the line integral and the surface integral, because then we can identify the weight function, which should be used in each case.

In Chapter 30 we look at the cases, where  $f : A \rightarrow \mathbb{R}$  is continuous, but  $A$  is either unbounded or not closed. In this case the integrand may tend to  $\pm\infty$ , when  $\mathbf{x}$  approaches a boundary point. Such integrals called improper integrals.

One should for improper integrals always split the integrand  $f$  into its positive and negative parts, i.e.

$$f(\mathbf{x}) = f_+(\mathbf{x}) - f_-(\mathbf{x}),$$

where

$$f_+(\mathbf{x}) := \begin{cases} f(\mathbf{x}) & \text{if } f(\mathbf{x}) > 0, \\ 0 & \text{if } f(\mathbf{x}) \leq 0, \end{cases} \quad f_-(\mathbf{x}) := \begin{cases} 0 & \text{if } f(\mathbf{x}) > 0, \\ -f(\mathbf{x}) & \text{if } f(\mathbf{x}) \leq 0, \end{cases}$$

where for technical reasons both  $f_+$  and  $f_-$  are nonnegative. Then we discuss the possible definition of

$$\int_A f(\mathbf{x}) d\mu, \quad \text{provided that } f(\mathbf{x}) \geq 0 \text{ for all } \mathbf{x} \in A,$$

i.e. for  $f$  nonnegative. Or, alternatively, split the domain  $A = A_+ \cup A_-$ , where  $f > 0$  on  $A_+$  and  $f \leq 0$  on  $A_-$ .

If both  $f_+$  and  $f_-$  have finite values of their (proper or improper) integrals, then we say that

$$\int_A f(\mathbf{x}) d\mu := \int_A f_+(\mathbf{x}) d\mu - \int_A f_-(\mathbf{x}) d\mu = \int_{A_+} f(\mathbf{x}) d\mu + \int_{A_-} f(\mathbf{x}) d\mu \quad (\in \mathbb{R}),$$

is convergent.

All the integrals above, plane, space and surface integrals, can be improper. To decide whether they are convergent or not, we split them as above and then use a truncation technique and finally let the truncations shrink towards the “singular points” on the (intrinsic) boundary of  $A$  to see if we have convergence or divergence.



## 29 Transformation of plane and space integrals

### 29.1 Transformation of a plane integral

We shall in this section see how we can integrate a plane integral by using a change of variables.

Consider two bounded and closed plane sets  $B, D \subset \mathbb{R}^2$ , and let

$$\mathbf{r} = (r_1, r_2) : D \rightarrow \mathbb{R}^2$$

be a  $C^1$  vector function, which satisfies

- 1) The vector function  $\mathbf{r}$  maps  $D$  onto  $B$ , i.e.  $\mathbf{r}(D) = B$ , so it is surjective.
- 2) The vector function  $\mathbf{r}$  is injective almost everywhere.

We use the coordinates  $(x, y) \in B$  and  $(u, v) \in D$ , so we have

$$x = r_1(u, v) \quad \text{and} \quad y = r_2(u, v).$$

If we consider  $B$  as a *surface*, and not just a plane set, then  $\int_B f(x, y) dS$  can be viewed as a *surface integral*, so we get from Chapter 27 that the reduction formula is

$$\int_B f(x, y) dx dy = \int_D f(r_1(u, v), r_2(u, v)) \cdot \|\mathbf{N}(u, v)\| du dv.$$

When the plane domain (surface)  $B$  is imbedded in  $\mathbb{R}^3$ , we can use the following rectangular description,

$$B = \{(x, y, 0) \mid x = r_1(u, v) \text{ and } y = r_2(u, v) \text{ for } (u, v) \in D\}.$$

Then the *normal vector*  $\mathbf{N}$  is parallel with the  $z$ -axis, and we get

$$\mathbf{N}(u, v) = \det \begin{pmatrix} \frac{\partial r_1}{\partial u} & \frac{\partial r_1}{\partial v} & 0 \\ \frac{\partial r_2}{\partial u} & \frac{\partial r_2}{\partial v} & 0 \\ \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \end{pmatrix} = \det \begin{pmatrix} \frac{\partial r_1}{\partial u} & \frac{\partial r_1}{\partial v} \\ \frac{\partial r_2}{\partial u} & \frac{\partial r_2}{\partial v} \end{pmatrix} \mathbf{e}_3,$$

so we have calculated the weight function, which is the absolute value of the so-called *Jacobian*, defined by

$$\frac{\partial(x, y)}{\partial(u, v)} := \det \begin{pmatrix} \frac{\partial r_1}{\partial u} & \frac{\partial r_1}{\partial v} \\ \frac{\partial r_2}{\partial u} & \frac{\partial r_2}{\partial v} \end{pmatrix}, \quad \|\mathbf{N}(u, v)\| = \left| \frac{\partial(x, y)}{\partial(u, v)} \right|.$$

The requirement from Chapter 27 is that the weight function is  $\neq 0$  almost everywhere. We can therefore formulate (and again without a correct proof) the following theorem,

**Theorem 29.1** The transformation theorem for a plane integral. Let  $(x, y) \in B$  and  $(u, v) \in D$ , where  $B$  and  $D$  are bounded and closed sets in the  $(x, y)$ -plane, the  $(u, v)$ -plane resp.. Assume that

$$\mathbf{r} = (r_1, r_2) : D \rightarrow \mathbb{R}^2$$

is a  $C^1$  vector function, such that


- 1) The vector function maps  $D$  onto  $B$ , i.e.  $\mathbf{r}(D) = B$ , and  $x = r_1(u, v)$  and  $y = r_2(u, v)$ .
- 2) The vector function  $\mathbf{r}$  is injective almost everywhere in  $D$ .
- 3) The Jacobian is  $\neq 0$  almost everywhere, i.e.

$$\frac{\partial(x, y)}{\partial(u, v)} := \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \neq 0 \quad \text{almost everywhere.}$$

If  $f: B \rightarrow \mathbb{R}$  is a continuous function, then we have the reduction formula

$$\int_B f(x, y) dx dy = \int_D f(r_1(u, v), r_2(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv.$$

I joined MITAS because  
I wanted **real responsibility**



Real work  
International opportunities  
Three work placements

**Month 16**  
I was a construction supervisor in the North Sea advising and helping foremen solve problems

The Graduate Programme for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)





It is for mnemotechnical reasons that we use the old-fashioned notation of the *Jacobian*. The reason is that we then remember that intuitively  $du\,dv$  in the “numerator” is “cancelled” by the symbol “ $\partial(u, v)$ ” in the “denominator”, leaving “ $\partial(x, y)$ ” in the “numerator”, which is more or less the same as  $dx\,dy$  on the left hand side of the transformation formula. This incorrect notation reminds us that the *Jacobian* is a function of  $(u, v)$ , which is more difficult to derive, when we use a more correct notation.

As a simple check, let us consider the change from rectangular coordinate in the plane to polar coordinates, so  $\mathbf{r}$  is given by

$$x = \varrho \cos \varphi \quad \text{and} \quad y = \varrho \sin \varphi.$$

Then

$$\frac{\partial(x, y)}{\partial(\varrho, \varphi)} = \det \begin{pmatrix} \frac{\partial x}{\partial \varrho} & \frac{\partial x}{\partial \varphi} \\ \frac{\partial y}{\partial \varrho} & \frac{\partial y}{\partial \varphi} \end{pmatrix} = \det \begin{pmatrix} \cos \varphi & -\varrho \sin \varphi \\ \sin \varphi & \varrho \cos \varphi \end{pmatrix} = \varrho,$$

which is precisely the weight function we found previously in Chapter 20, when we used polar coordinates, so we have again derived the well-known formula

$$\int_B f(x, y) dx dy = \int_D f(\varrho \cos \varphi, \varrho \sin \varphi) \varrho d\varrho d\varphi.$$

## 29.2 Transformation of a space integral

Since in the previous section

$$\|\mathbf{N}(u, v)\| = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \left| \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix} \right|$$

is the area of the parallelogram formed by the tangents of the two parameter curves, this gives us a clue of what we should expect in three

The analogue in three dimensions is a map  $\mathbf{r} = (r_1, r_2, r_3) : D \rightarrow B$ , where we use the notation

$$(x, y, z) = (r_1(u, v, w), r_2(u, v, w), r_3(u, v, w)),$$

and where we assume that  $\mathbf{r}$  is surjective, and injective almost everywhere.

The *Jacobian* is here

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} := \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{pmatrix}.$$

It is well-known from *Linear Algebra* that the absolute value of the *Jacobian* is the volume of the parallelepipedum spanned by the three tangents of the three parameter curves at a given point. In other words, our parallelepipeda are the building stones, which we use when we build up the 3-dimensional integral, and their volumes, the absolute value of the Jacobian, form the weight function.

The above makes the following theorem plausible. We quote it – as usual without a correct proof.

The advertisement features a man in a green jacket walking down a city street at night, looking towards the right. The background shows blurred city lights and buildings. On the left, the IE business school logo is visible. On the right, a graphic indicates "#1 European Business School FINANCIAL TIMES 2013". A speech bubble contains the hashtag #gobeyond. Below the image, a banner reads "MASTER IN MANAGEMENT". The text "Because achieving your dreams is your greatest challenge." is followed by a description of the program's goals: "IE Business School's Master in Management taught in English, Spanish or bilingually, trains young high performance professionals at the beginning of their career through an innovative and stimulating program that will help them reach their full potential." A bulleted list highlights features: "Choose your area of specialization.", "Customize your master through the different options offered.", and "Global Immersion Weeks in locations such as London, Silicon Valley or Shanghai." The tagline "Because you change, we change with you." is at the bottom. Contact information includes the website [www.ie.edu/master-management](http://www.ie.edu/master-management), email [mim.admissions@ie.edu](mailto:mim.admissions@ie.edu), and social media links for Facebook, Twitter, LinkedIn, YouTube, and Instagram. A green button at the bottom right encourages interaction with the ad.

**Theorem 29.2** Transformation theorem for a space integral Let  $(x, y, z) \in B$  and  $(u, v, w) \in D$ , where  $B \subset \mathbb{R}^3$  and  $D \subset \mathbb{R}^3$  are bounded and closed sets. Assume that

$$\mathbf{r} = (r_1, r_2, r_3) : D \rightarrow \mathbb{R}^3$$

is a  $C^1$  vector function, such that

1) The vector function  $\mathbf{r}$  maps  $D$  onto  $B$ , i.e.  $\mathbf{r}(D) = B$ , and

$$x = r_1(u, v, w), \quad y = r_2(u, v, w), \quad z = r_3(u, v, w).$$

2) The function  $\mathbf{r}$  is injective almost everywhere.

3) The Jacobian is  $\neq 0$  almost everywhere, i.e.

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} := \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{pmatrix} \neq 0 \quad \text{almost everywhere.}$$

If  $f : B \rightarrow \mathbb{R}$  is a continuous function, then the reduction formula is

$$\int_B f(x, y, z) dx dy dz = \int_D f(r_1(u, v, w), r_2(u, v, w), r_3(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw.$$

Let us check this formula by transforming the integral in rectangular coordinates into spherical coordinates, i.e.  $\mathbf{r}$  is here specified by

$$x = r \sin \theta \cos \varphi, \quad y = r \sin \theta \sin \varphi, \quad z = r \cos \theta.$$

Then the Jacobian is

$$\begin{aligned} \frac{\partial(x, y, z)}{\partial(r, \theta, \varphi)} &= \det \begin{pmatrix} \sin \theta \cos \varphi & r \cos \theta \cos \varphi & -r \sin \theta \sin \varphi \\ \sin \theta \cos \varphi & r \cos \theta \sin \varphi & r \cos \theta \cos \varphi \\ \cos \theta & -r \sin \theta & 0 \end{pmatrix} \\ &= \cos \theta \det \begin{pmatrix} r \cos \theta \cos \varphi & -r \sin \theta \sin \varphi \\ r \cos \theta \sin \varphi & r \sin \theta \cos \varphi \end{pmatrix} + r \sin \theta \det \begin{pmatrix} \sin \theta \cos \varphi & -r \sin \theta \sin \varphi \\ \sin \theta \sin \varphi & r \sin \theta \cos \varphi \end{pmatrix} \\ &= r^2 \cos^2 \theta \sin \theta \det \begin{pmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{pmatrix} + r^2 \sin^3 \theta \det \begin{pmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{pmatrix} \\ &= r^2 \sin \theta \{ \cos^2 \theta + \sin^2 \theta \} = r^2 \sin \theta, \end{aligned}$$

so we have indeed obtained the same weight function as we found in Chapter 24.

### 29.3 Procedures for the transformation of plane or space integrals

All the reduction formulæ in the previous chapters are special cases of more general formulæ. The presentations were using the classical coordinate systems: the *rectangular, polar, semi-polar, and the spherical* coordinate systems. When the coordinate system under consideration is *not* one of these we must use the general formulæ from the present section instead.

#### A. Dimension 2.

- 1) Find a suitable parameter representation  $(x, y) = \mathbf{r}(u, v)$ ,  $(u, v) \in D$ .

Sketch the parametric domain  $D$  and argue *briefly* that  $\mathbf{r}(u, v)$  is injective, with the exception of e.g. a *finite* number of points. (More precisely one can neglect a so-called null set; which usually is not defined in elementary courses in Calculus).

Show also that the *range* is  $\mathbf{r}(D) = B$ .

- 2) Calculate the *Jacobian*

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}.$$

Note here that one is *forced* to find  $x$  and  $y$  as *functions of*  $(u, v)$  in 1) in order to calculate the Jacobian.

The *area element* is

$$dS = dx dy = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv,$$

where the area element  $dx dy$  is lying in  $B$ , while the area element  $du dv$  is lying in the parametric domain  $D$ .

- 3) Insert the result and calculate the right hand side by previous known methods in the expression

$$\int_B f(x, y) dx dy = \int_D f(\mathbf{r}(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv.$$

**Remark 29.1** Usually  $u$  and  $v$  are given as functions in  $x$  and  $y$  instead of the form we shall use:

$$(29.1) \quad u = U(x, y) \quad \text{and} \quad v = V(x, y).$$

Then one has to solve these equations with respect to  $x$  and  $y$ . If we in this way obtain a unique solution, then we have at the same time implicitly proved that the map is injective. Apply furthermore (29.1) to find the images of the boundary curves of  $B$ , thereby finding the boundary of  $D$ . Finally, the parametric domain  $D$  is identified.  $\diamond$

### B. Dimension 3.

Formally the procedure is the same as in section A with obvious modifications due to the higher dimension.

- 1) Find a suitable parametric representation  $(x, y, z) = \mathbf{r}(u, v, w)$ ,  $(u, v, w) \in D$ . (This will usually be given, possibly in the form  $u = U(x, y, z)$ ,  $v = V(x, y, z)$ ,  $w = W(x, y, z)$ . If so, solve these equations with respect to  $x, y, z$ ). Sketch the parametric domain  $D$  and argue (briefly) that the mapping  $\mathbf{r}(u, v, w)$  is injective almost everywhere.  
 Show that the range is  $\mathbf{r}(D) = B$ . Cf. also the remarks to section A.
- 2) Calculate the **Jacobian**

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}.$$

Then the *volume element* is

$$d\Omega = dx dy dz = \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw,$$

where one must be careful not to forget the numerical signs of the Jacobian.

- 3) Insert and calculate the right hand side by means of one of the previous methods in the formula

$$\int_B f(x, y, z) dx dy dz = \int_D f(\mathbf{r}(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw.$$

### 29.4 Examples of transformation of plane and space integrals

#### Example 29.1

##### A. Calculate the plane integral

$$I = \int_B \cos\left(\frac{y-x}{y+x}\right) dx dy$$

over the trapeze shown on the figure.

- D. A direct calculation applying one of the usual reduction theorems is not possible, because none of the forms

$$\int \cos\left(\frac{y-x}{y+x}\right) dx = \int \cos\left(\frac{2y}{y+x} - 1\right) dx = \int \cos\left(1 - \frac{2x}{y+x}\right) dy$$

can be integrated within the realm of our known functions. The situation is even worse in polar coordinates. Therefore, the only possibility left is to find a convenient transform, such that the integrand becomes more easy to handle.




Figure 29.1: The trapeze  $B$ .

The unpleasant thing is of course the fraction  $\frac{y-x}{y+x}$ . One idea would be to introduce the numerator as a new variable, and the denominator as another new variable. If we do this, then we must show that we obtain a *unique* correspondence between the domain  $B$  and a *parametric domain*  $D$ , which also should be found. Finally we shall find the *Jacobian*. When we have found all the terms in the transformation formula, then calculate the integral.

"I studied English for 16 years but...  
...I finally learned to speak it in just six lessons"  
Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download



Click on the ad to read more

**Remark 29.2** This time we see that it is here quite helpful to start the discussion in  $\mathbf{D}$ , which is not common knowledge from high school. First we discuss the problem. Based on this discussion we make a decision on the further procedure.  $\diamond$

- I. According to  $\mathbf{D}$ . we choose the numerator and the denominator as our new variables. Most people would here choose the *numerator* as  $u$  and the *denominator* as  $v$ , so we shall do the same, although it can be shown that we shall get simpler calculations if we interchange the definition of  $u$  and  $v$ .

We therefore put as the most natural choice

$$(29.2) \text{ numerator : } u = y - x \quad \text{and} \quad \text{denominator : } v = y + x.$$

Then we shall prove that this gives a *one-to-one* correspondence. This means that we for any given  $u$  and  $v$  obtain unique solutions  $x$  and  $y$ :

$$x = \frac{v - y}{2} \quad \text{and} \quad y = \frac{u + v}{2}.$$

Obviously the transform is *continuous* both ways. Since  $B$  is *closed and bounded*, the range  $D$  by this transform is again *closed and bounded*, cf. the important *second main theorem for continuous functions*.

Since the transform is one-to-one *everywhere*, the boundary  $\partial B$  is mapped one-to-one onto the boundary  $\partial D$ . This is expressed in the following way:

- 1) The line  $x + y = 1$  corresponds by (29.2) to  $v = 1$ .
- 2) The line  $y = x$ , i.e.  $y - x = 0$ , corresponds by (29.2) to  $u = 0$ .
- 3) The line  $y + x = 4$  corresponds by (29.2) to  $v = 4$ .
- 4) The line  $y = 3x$  corresponds to  $\frac{u+v}{2} = 3 \frac{v-u}{2}$ , i.e. to  $v = 2u$ .




Figure 29.2: The parametric domain  $D$ . The oblique line has the equation  $v = 2u$  or  $u = \frac{1}{2}v$  as its representation.

The only *closed and bounded* domain in the  $(u, v)$ -plane, which has the new boundary curves as its boundary is  $D$  as indicated on the figure. In practice one draws the lines  $v = 1$ ,  $u = 0$ ,  $v = 4$  and

$v = 2u$  and use the figure to find out where the bounded set  $D$  is situated, such that the boundary consists of parts of all four lines.

Then we calculate the *weight function*  $\left| \frac{\partial(x, y)}{\partial(u, v)} \right|$ .

The “old fashioned” notation above indicates that we shall use the transform, where  $x$  and  $y$  (in the “numerator”) are functions of  $u$  and  $v$  (in the “denominator”), i.e.

$$x = \frac{1}{2}v - \frac{1}{2}u \quad \text{and} \quad y = \frac{1}{2}u + \frac{1}{2}v.$$

This gives us the *Jacobian*

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{vmatrix} = \frac{1}{2} \cdot \frac{1}{2} \begin{vmatrix} -1 & 1 \\ 1 & 1 \end{vmatrix} = \frac{1}{4} \cdot (-2) = -\frac{1}{2}.$$

It follows that the Jacobian is *negative*, hence the *weight function* becomes

$$\left| \frac{\partial(x, y)}{\partial(u, v)} \right| = \frac{1}{2}.$$

**Remark 29.3** If we here interchange  $u$  and  $v$  in (29.2), then we obtain that the Jacobian becomes *positive*. ◇

We have now come to the *reduction formula*,

$$I = \int_B \cos\left(\frac{y-x}{y+x}\right) dx dy = \int_D \cos\left(\frac{u}{v}\right) \cdot \frac{1}{2} du dv.$$

Note that both sides here are *abstract plane integrals*.

We see on the right hand side that  $\int \cos\left(\frac{u}{v}\right) dv$  cannot be integrated within the realm of our known arsenal of functions. But  $\int \cos\left(\frac{u}{v}\right) du$  can! Therefore, when we reduce the plane integral on the right hand side we put the  $u$ -integral as the inner integral. Then

$$I = \frac{1}{2} \int_D \cos\left(\frac{u}{v}\right) du dv = \frac{1}{2} \int_1^4 \left\{ \int_0^{\frac{v}{2}} \cos\left(\frac{u}{v}\right) du \right\} dv.$$

When  $v \neq 0$  is kept constant, we get from the inner integral

$$\int_0^{\frac{v}{2}} \cos\left(\frac{u}{v}\right) du = \left[ v \sin\left(\frac{u}{v}\right) \right]_0^{\frac{v}{2}} = v \sin\left(\frac{v}{2} \cdot \frac{1}{v}\right) = \sin\left(\frac{1}{2}\right) \cdot v,$$

where  $\sin\left(\frac{1}{2}\right)$  is a constant, which shall *not* be found explicitly! (Note that at  $\frac{1}{2}$  radian is *not* equal to  $\frac{\pi}{2}$ ).

Finally we get by insertion

$$I = \frac{1}{2} \int_1^4 \sin\left(\frac{1}{2}\right) \cdot v \, dv = \frac{1}{2} \sin\left(\frac{1}{2}\right) \cdot \left[\frac{1}{2} v^2\right]_1^4 = \frac{15}{4} \sin\left(\frac{1}{2}\right). \quad \diamond$$

Excellent Economics and Business programmes at:

 university of  
groningen





**“The perfect start  
of a successful,  
international career.”**

[www.rug.nl/feb/education](http://www.rug.nl/feb/education)

**CLICK HERE**  
to discover why both socially  
and academically the University  
of Groningen is one of the best  
places for a student to be



Click on the ad to read more

**Example 29.2**

A. Let

$$A = \{(x, y, z) \mid 0 \leq x \leq 1, 0 \leq y \leq \sqrt{x}, -y \leq z \leq y\}.$$

calculate

$$I = \int_A \frac{\exp((2-y-z)^3)}{4+y+z} d\Omega.$$




Figure 29.3: The domain  $A$ . Note the different scales on the axes.

D. Let us start by pulling out the teeth of this big and horrible example! Its purpose is *only* to demonstrate that even apparent incalculable integrals in some cases nevertheless can be calculated by using a “convenient transform”. This example is from a textbook, where earlier students got the wrong impression that “every application of the transformation theorem looks like this example”, which is not true. Without this extra comment this example will send a *wrong* message to the reader.

Let us first discuss, how we can find a reasonable transform. I shall follow more or less the way of thinking which the author of this example must have used, the first time it was created.

At the end of this example I shall describe the very modest requirements which may be demanded of the students. In other words, this example should only be used as an *inspiration* for other similar problems which may occur in practice.

I. Let us start by looking at the *geometry* of  $A$ . The projection  $B$  of  $A$  onto the  $(x, y)$ -plane is

$$B = \{(x, y) \mid 0 \leq x \leq 1, 0 \leq y \leq \sqrt{x}\}.$$

Since  $A$  for every  $x \in ]0, 1]$  is cut into an isosceles rectangular triangle

$$\Delta_x = \{(y, z) \mid 0 \leq y \leq \sqrt{x}, -y \leq z \leq y\},$$

it is easy to sketch  $A$ , cf. a previous figure.




Figure 29.4: The projection  $B$  of the domain  $A$  onto the  $(x, y)$ -plane.

Then the integrand

$$\frac{\exp((2-y-z)^3)}{4+y+z}$$

should be “straightened out”. A reasonable guess would be to introduce

$$u = y + z.$$

**Remark 29.4** Once we have gone through all the calculations it can be seen that

$$\tilde{u} = \frac{y+z}{2},$$

would be a better choice, because then we would get rid of a lot of irritating constants. Of pedagogical reasons we shall *not* here use the most optimal transform, but instead the transform which one would expect the student to choose.  $\diamond$

Since we do not get further information from the integrand, we shall turn to the domain  $A$ . The boundary of  $A$  is (almost) determined by putting equality sign into the definition of  $A$  instead of  $\leq$ . First everything is written in a “binary” way in the definition of  $A$ ,

$$\begin{aligned} A &= \{(x, y, z) \mid 0 \leq x \leq 1 \wedge 0 \leq y \leq \sqrt{x} \wedge -y \leq z \leq y\} \\ &= \{(x, y, z) \mid 0 \leq x \wedge x \leq 1 \wedge 0 \leq y \wedge 0 \leq \sqrt{x} - y \wedge 0 \leq y + z \wedge 0 \leq y - z\}, \end{aligned}$$

i.e. every condition which is defining  $A$  contains only one inequality sign and one of the sides of the inequality is a constant.

We see that there are composed expressions in the latter three conditions,

$$\sqrt{x} - y \geq 0, \quad y + z \geq 0, \quad y - z \geq 0.$$

Since we already have chosen  $u = y + z \geq 0$ , we get the inspiration of choosing the new variables

$$(29.3) \quad u = y + z \geq 0, \quad v = y - z \geq 0, \quad w = \sqrt{x} - y \geq 0,$$

where we have taken the most ugly term,  $\sqrt{x} - y$  and put it equal to  $w$ , i.e.

$$w = \sqrt{x} - y.$$

We note that we by these choices have obtained that  $u, v, w \geq 0$ , and that equality signs must correspond to boundary points in the  $(u, v, w)$ -space for the parametric domain  $D$ .

Next we show that the transform (29.3) is one-to-one. i.e. we shall express  $x, y, z$  uniquely by  $u, v, w$ . We get immediately from the first two equations that

$$y = \frac{u+v}{2} \quad \text{and} \quad z = \frac{u-v}{2}.$$

From the third equation we get

$$\sqrt{x} = w + y = w + \frac{u+v}{2} = \frac{1}{2}(u+v+2w),$$

which obviously is  $\geq 0$ , because  $u, v, w \geq 0$ . Therefore, by squaring,

$$x = \frac{1}{4}(u+v+2w)^2.$$

Thus,  $x, y, z$  are uniquely determined by  $u, v, w$ , so the transform is *one-to-one*.

Since the transform and its inverse are both *continuous* and the domain  $A$  is *closed and bounded*, it follows from the *second main theorem for continuous functions* that  $D$  is also *closed and bounded*. It follows from the binary representation of  $A$  that  $\partial A$  is a subset of the union of the surfaces  $x = 0, x = 1, y = 0, \sqrt{x} - y = 0, y + z \leq 0$  and  $y - z = 0$ . These are now investigated one by one.




Figure 29.5: The projection of the parametric domain  $D$  in the  $(u, v)$ -plane.

- 1) The plane  $x = 0$  corresponds to  $\frac{1}{4}(u+v+2w)^2 = 0$ . Since  $u, v, w \geq 0$ , we only get  $(u, v, w) = (0, 0, 0)$ , which is in agreement with the figure of  $A$ , because the plane  $x = 0$  just cuts  $A$  in  $\mathbf{0}$ .

2) The plane  $x = 2$  corresponds to  $\frac{1}{4}(u + v + w)^2 = 1$ , i.e.  $\frac{1}{2}(u + v + 2w) = +1$ , from which

$$w = 1 - \frac{u + v}{2} \geq 0.$$

Here we have again used that  $u, v, w \geq 0$ . Note that we also get that

$$u + v \leq 2.$$

3) The plane  $y = 0$  corresponds to  $\frac{1}{2}(u + v) = 0$ , i.e.  $u + v = 0$ .

4) The remaining conditions have been found previously for  $u = 0$ ,  $v = 0$  and  $w = 0$ .

## American online LIGS University

is currently enrolling in the  
Interactive Online BBA, MBA, MSc,  
DBA and PhD programs:

- ▶ enroll by September 30th, 2014 and
- ▶ save up to 16% on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive Online education
- ▶ visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to find out more!

Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education.  
[More info here.](#)



Click on the ad to read more




Figure 29.6: The parametric domain  $D$ .

Summing up we find that the *parametric domain* is given by

$$\begin{aligned} D &= \left\{ (u, v, w) \mid 0 \leq u, 0 \leq u + v \leq 2, 0 \leq w \leq 1 - \frac{u+v}{2} \right\} \\ &= \left\{ (u, v, w) \mid 0 \leq u \leq 2, 0 \leq v \leq 2 - u, 0 \leq w \leq 1 - \frac{u+v}{2} \right\} \\ &= \left\{ (u, v, w) \mid (u, v) \in B, 0 \leq w \leq 1 - \frac{u+v}{2} \right\}, \end{aligned}$$

where the projection  $B$  in the  $(u, v)$ -plane is given by

$$B = \{(u, v) \mid 0 \leq u \leq 2, 0 \leq v \leq 2 - u\},$$

so  $B$  and  $D$  are now easily sketched.

By the chosen transform the *integrand* is carried over into

$$\frac{\exp((2-y-z)^3)}{4+y+z} = \frac{\exp((2-u)^2)}{4+u}.$$

Then we calculate the *weight function*  $\left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right|$ . First note that

$$\begin{aligned} \frac{\partial(x, y, z)}{\partial(u, v, w)} &= \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix} = \begin{vmatrix} \frac{1}{2}(u+v+2w) & \frac{1}{2}(u+v+2w) & u+v+2w \\ \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{vmatrix} \\ &= (u+v+2w) \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}(u+v+2w). \end{aligned}$$

Since  $u, v, w \geq 0$  in  $D$ , we see that the *weight function* is

$$\left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| = +\frac{1}{2}(u + v + 2w).$$

This is only 0 for  $(u, v, w) = (0, 0, 0)$  in  $D$ , i.e. in just one point, which is a null-set (without a positive volume). Therefore we may continue with the transformation theorem in its abstract form:

$$\begin{aligned} I &= \int_A \frac{\exp((2-y-z)^3)}{4+y+z} dx dy dz \\ &= \int_D \frac{\exp((2-u)^3)}{4+u} \frac{1}{2}(u+v+2w) du dv dw. \end{aligned}$$

Here it is obvious that we shall *not* start by integrating after  $u$ . If we choose  $u$  as the last (i.e. the outer) variable of integration, then we get by one of the reduction theorems that

$$(29.4) \quad I = \frac{1}{2} \int_0^2 \frac{\exp((2-u)^3)}{4+u} \left\{ \int_{B(u)} (u+v+2w) dv dw \right\} du,$$

where  $B(u)$  is the intersection of  $D$  for  $u$  constant, i.e.

$$B(u) = \left\{ (v, w) \mid 0 \leq v \leq 2-u, 0 \leq w \leq 1 - \frac{u+v}{2} \right\}.$$

We calculate for fixed  $u \in [0, 2]$  the inner integral in (29.4) by first integrating vertically with respect to  $w$ :

$$\int_{B(u)} (u+v+2w) dv dw = \int_0^{2-u} \left\{ \int_0^{1-\frac{u+v}{2}} (u+v+2w) dw \right\} dv.$$

We calculate the inner integral

$$\begin{aligned} \int_0^{1-\frac{u+v}{2}} (u+v+2w) dw &= [(u+v)w + w^2]_0^{1-\frac{u+v}{2}} = [w(u+v+w)]_0^{1-\frac{u+v}{2}} \\ &= \left\{ 1 - \frac{u+v}{2} \right\} \left\{ 1 + \frac{u+v}{2} \right\} = 1 - \frac{1}{4}(u+v)^2. \end{aligned}$$

By insertion we next get for fixed  $u$  that

$$\begin{aligned} \int_{B(u)} (u+v+2w) dw dv &= \int_0^{2-u} \left\{ 1 - \frac{1}{4}(u+v)^2 \right\} dv = \left[ v - \frac{1}{12}(u+v)^3 \right]_{v=0}^{2-u} \\ &= -(u-2) + \frac{1}{12}(u^3 - 2^3) \\ &= \frac{1}{12}(u-2)\{-12 + u^2 + 2u + 4\} \\ &= \frac{1}{12}(u-2)\{u^2 + 2u - 8\} = \frac{1}{12}(u-2)^2(u+4). \end{aligned}$$

Note that we have found all factors. When this result is put into (29.4), we get the reduction

$$\begin{aligned} I &= \frac{1}{2} \int_0^2 \frac{\exp((2-u)^3)}{4+u} \left\{ \int_{B(u)} (u+v+2w) dv dw \right\} du \\ &= \frac{1}{2} \int_0^2 \frac{\exp((2-u)^3)}{4+u} \cdot \frac{1}{12} (u-2)^2 (u+4) du \\ &= \frac{1}{24} \int_0^2 \exp((2-u)^3) (u-2)^2 du. \end{aligned}$$

Now, choose the substitution  $t = (2-u)^3$ . Then  $dt = -3(2-u)^2 du$ , and hence

$$(u-2)^2 du = -\frac{1}{3} dt.$$

Finally we get

$$\begin{aligned} I &= \frac{1}{24} \int_0^2 \exp((2-u)^3) (u-2)^2 du = \frac{1}{24} \int_{(2-0)^3}^{(2-2)^3} \exp(t) \cdot \left(-\frac{1}{3}\right) dt \\ &= \frac{1}{72} \int_0^8 e^t dt = \frac{e^8 - 1}{72}. \end{aligned}$$

**Remark 29.5** It is obvious from this example, that the application of transformation theorems is not an easy job. Therefore, one will usually be *given* the transform which should be applied,

$$u = f(x, y, z), \quad v = g(x, y, z), \quad w = h(x, y, z).$$

Then the task for the reader can be described in the following points:

1) *Solve* the equations after  $x, y, z$ , (from this follows automatically that the transform is *one-to-one*),

$$x = F(u, v, w), \quad y = G(u, v, w), \quad z = H(u, v, w).$$

2) *Identify* the parametric domain; use here the second main theorem and that a boundary in most cases by a continuous transform again is mapped into a part of the boundary.

3) *Calculate* the weight function  $\left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right|$  from the expressions found in 1).

4) *Reduce* the integrand

$$\Phi(x, y, z) = \Phi(F, G, H) = \Psi(u, v, w);$$

5) *Set up* the abstract reduction formula,

$$\int_A \Phi(x, y, z) dx dy dz = \int_B \Psi(u, v, w) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw.$$

6) *Reduce* the right hand side of 5) by known methods, usually in rectangular coordinates, though semi-polar coordinates may occur, and calculate the value of the resulting integral.

It should be of no surprise that in general even this very simple type of example may be fairly large.◊

**Example 29.3** Let  $B$  be the trapeze which is bounded by the coordinate axes and the lines given by the equations  $x + y = 1$  and  $x + y = \frac{1}{2}$ . Compute the plane integral

$$\int_B \exp\left(\frac{y}{x+y}\right) dx dy$$

by introducing the new variable  $(u, v) = (x + y, x - y)$ .

**A** Transformation of a plane integral.

**D** Compute the Jacobian and find the new domain  $D$ .




Figure 29.7: The domain  $B$  in the  $XY$ -plane.




Figure 29.8: The domain  $D$  after the transformation to the  $UV$ -plane.

**I** From

$$(x, y) = \Phi(u, v) = \left( \frac{u+v}{2}, \frac{u-v}{2} \right),$$

follows that

$$J_{\Phi} = \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{1}{2} & d^{\frac{1}{2}} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2},$$

and

$$D = \left\{ (u, v) \mid \frac{1}{2} \leq u \leq 1, -u \leq v \leq u \right\}.$$

Then by the formula of transformation,

$$\begin{aligned} \int_B \exp\left(\frac{y}{x+y}\right) dx dy &= \int_D \exp\left(\frac{u-v}{2u}\right) \cdot \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv \\ &= \frac{1}{2} \int_{\frac{1}{2}}^1 \sqrt{e} \left\{ \int_{-u}^u \exp\left(-\frac{v}{2u}\right) dv \right\} du = \frac{\sqrt{e}}{2} \int_{\frac{1}{2}}^1 (-2u) \left[ \exp\left(-\frac{v}{2u}\right) \right]_{v=-u}^u du \\ &= -\sqrt{e} \int_{\frac{1}{2}}^1 u \cdot \left( \frac{1}{\sqrt{e}} - \sqrt{e} \right) du = (e-1) \int_{\frac{1}{2}}^1 u du = \frac{3}{8} (e-1). \end{aligned}$$

..... Alcatel-Lucent 

[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?

One generation's transformation is the next's status quo.  
 In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".



Click on the ad to read more

**Example 29.4** Let  $B$  denote set in the first quadrant, which is bounded by the curves  $xy = 1$  and  $xy = 2$  and by the lines  $y = x$  and  $y = 4x$ . Sketch  $B$  and compute the plane integral

$$\int_B x^2 y^2 dx dy$$

by introducing the new variables  $(u, v) = \left( xy, \frac{y}{x} \right)$ .

**A** Transformation of a plane integral.

**D** Sketch  $B$ . Find den inverse function

$$(x, y) = (x(u, v), y(u, v)) = \Phi(u, v),$$

and find the corresponding domain  $D$  in the  $UV$ -plane. Calculate the Jacobian and finally transform the plane integral.




Figure 29.9: The domain  $B$  in the  $XY$ -plane.

**I** If  $u = xy$  and  $v = \frac{y}{x}$  and  $x, y > 0$ , then  $u, v > 0$ , and

$$x(u, v) = \sqrt{\frac{u}{v}}, \quad y(u, v) = \sqrt{uv}.$$

The domain  $D$  is given by

$$1 \leq xy = u \leq 2 \quad \text{and} \quad 1 \leq \frac{y}{x} = v \leq 4,$$

hence

$$D = \{(u, v) \mid 1 \leq u \leq 2, 1 \leq v \leq 4\} = [1, 2] \times [1, 4],$$

i.e. a rectangle in the  $UV$ -plane, which it is no need to sketch.

Finally, the Jacobian is

$$\begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{2\sqrt{uv}} & -\frac{1}{2} \cdot \frac{1}{v} \sqrt{\frac{u}{v}} \\ \frac{1}{2} \sqrt{\frac{v}{u}} & \frac{1}{2} \cdot \frac{1}{u} \sqrt{\frac{u}{v}} \end{vmatrix} = \frac{1}{4} \left\{ \frac{1}{\sqrt{uv}} \sqrt{\frac{u}{v}} + \sqrt{\frac{v}{u}} \cdot \frac{1}{v} \sqrt{\frac{u}{v}} \right\} = \frac{1}{4} \left\{ \frac{1}{v} + \frac{1}{v} \right\} = \frac{1}{2v}.$$

We get by the transformation formula of the plane integral

$$\begin{aligned}\int_B x^2 y^2 dx dy &= \int_D u^2 \cdot \frac{1}{2v} du dv = \frac{1}{2} \int_1^2 u^2 du \cdot \int_1^4 \frac{1}{v} dv \\ &= \frac{1}{2} \left[ \frac{1}{3} u^3 \right]_1^2 \cdot [\ln v]_1^4 = \frac{1}{6} (8 - 1) \ln 4 = \frac{7}{3} \ln 2.\end{aligned}$$

**Example 29.5** Find the area of the set in the first quadrant, which is bounded by the curves

$$xy = 4, \quad xy = 8, \quad xy^3 = 5, \quad xy^3 = 15,$$

by introducing the new variables  $u = xy$  and  $v = xy^3$ .

**A** Area of a set computed by a transformation of a plane integral.

**D** Find the transformed domain  $D$  in the  $UV$ -plane and the inverse functions  $x(u, v)$  and  $y(u, v)$  by this transformation. Calculate the Jacobian and apply the transformation formula to find the area.




Figure 29.10: The domain  $D$  in the  $XY$ -plane. (Different scales on the axes).

**I** Let  $B$  be the given set in the first quadrant. Then  $x, y > 0$  for  $(x, y) \in B$ . It follows immediately that we by the transformation get the domain

$$D = [4, 8] \times [5, 15].$$

From  $u = xy, v = xy^3, u > 0$  and  $v > 0$  follows  $y^2 = \frac{v}{u}$  and  $x^2 = \frac{u^3}{v}$ , i.e.

$$y = +\sqrt{\frac{v}{u}}, \quad \text{and} \quad x = +\sqrt{\frac{u^3}{v}}.$$

Then we get the Jacobian,

$$J(u, v) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{3}{2} \sqrt{\frac{u}{v}} & -\frac{1}{2} \sqrt{\frac{u^3}{v^3}} \\ -\frac{1}{2} \sqrt{\frac{v}{u^3}} & \frac{1}{2} \sqrt{\frac{1}{uv}} \end{vmatrix} = \frac{3}{4} \frac{1}{v} - \frac{1}{4} \frac{1}{v} = \frac{1}{2v} > 0.$$

Hence the area is

$$\text{area}(B) = \int_B dx dy = \int_D J(u, v) du dv = \int_4^8 du \cdot \int_4^1 5 \frac{1}{2v} dv = \frac{4}{2} [\ln v]_5^{15} = 2 \ln 3.$$

**Example 29.6** Find the area of the set in the first quadrant, which is bounded by the curves

$$y = x^3, \quad y = 4x^3, \quad x = y^3, \quad x = 4y^3,$$

by introducing the new variables

$$u = \frac{y}{x^3}, \quad v = \frac{x}{y^3}.$$

**A** Area of a set by a transformation of a plane integral.

**D** Sketch the domain  $B$ . Then find  $D$  and  $x(u, v)$  and  $y(u, v)$  by the transformation. Calculate the Jacobian and apply the transformation formula to find the area.




Figure 29.11: The domain  $B$  in the  $XY$ -plane.

**I** The curves  $y = x^3$  and  $x = y^3$  intersect at  $(x, y) = (1, 1)$ . The curves  $y = 4x^3$  and  $x = 4y^3$  intersect at  $(x, y) = \left(\frac{1}{2}, \frac{1}{2}\right)$ . It follows that if the transformation exists and is bijective, then

$$D = [1, 4] \times [1, 4].$$

Clearly,  $x > 0$  and  $y > 0$ , and hence  $u > 0$  and  $v > 0$ . We shall now try to solve the equations

$$u = \frac{y}{x^3} \quad \text{and} \quad v = \frac{x}{y^3} \quad \text{for } u, v \in [1, 4].$$

From

$$u^3 v = \frac{y^3}{x^9} \cdot \frac{x}{y^3} = \frac{1}{x^8}$$

follows that

$$x = u^{-\frac{3}{8}} v^{-\frac{1}{8}}, \quad \text{and similarly } y = u^{-\frac{1}{8}} v^{-\frac{3}{8}}.$$

The Jacobian is

$$\begin{aligned} J(u, v) &= \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} \\ \frac{\partial x}{\partial v} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} -\frac{3}{8} u^{-\frac{11}{8}} v^{-\frac{1}{8}} & -\frac{1}{8} u^{-\frac{9}{8}} v^{-\frac{3}{8}} \\ -\frac{1}{8} u^{-\frac{3}{8}} v^{-\frac{9}{8}} & -\frac{3}{8} u^{-\frac{1}{8}} v^{-\frac{11}{8}} \end{vmatrix} \\ &= \frac{9}{64} u^{-\frac{3}{2}} v^{-\frac{3}{2}} - \frac{1}{64} u^{-\frac{3}{2}} v^{-\frac{3}{2}} = \frac{1}{8} u^{-\frac{3}{2}} v^{-\frac{3}{2}}. \end{aligned}$$

We get the area by applying the transformation formula

$$\begin{aligned} \text{area}(B) &= \int_B dS = \frac{1}{8} \int_1^4 u^{-\frac{3}{2}} du \cdot \int_1^4 v^{-\frac{3}{2}} dv = \frac{1}{8} \left\{ \int_1^4 t^{-\frac{3}{2}} dt \right\}^2 \\ &= \frac{1}{8} \left\{ \left[ -\frac{2}{\sqrt{t}} \right]_1^4 \right\}^2 = \frac{1}{8} (2 - 1)^2 = \frac{1}{8}. \end{aligned}$$



**Maastricht University** *Leading in Learning!*

**Join the best at  
 the Maastricht University  
 School of Business and  
 Economics!**



**Visit us and find out why we are the best!  
 Master's Open Day: 22 February 2014**

**Top master's programmes**

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

Maastricht  
 University is  
 the best specialist  
 university in the  
 Netherlands  
 (Elsevier)

[www.mastersopenday.nl](http://www.mastersopenday.nl)



Click on the ad to read more

**Example 29.7** Let  $B \subset \mathbb{R}^2$  be given by

$$0 \leq x, \quad 0 \leq y, \quad \sqrt{x} + \sqrt{y} \leq 1.$$

find the area of  $B$  and the plane integral

$$I = \int_B \exp \left[ (\sqrt{x} + \sqrt{y})^4 \right] dx dy$$

by introducing the new variables

$$u = \sqrt{x} + \sqrt{y}, \quad v = \sqrt{x} - \sqrt{y}.$$

**A** Transformation of a plane integral.

**D** Sketch  $B$ ; find  $x$  and  $y$  as functions of  $u$  and  $v$ ; calculate the Jacobian; find the domain of the parameters  $(u, v) \in A$ ; finally, apply the transformation theorem.




Figure 29.12: The domain  $A$  in the  $(X, Y)$ -plane.

**I** If we put  $u = \sqrt{x} + \sqrt{y}$  and  $v = \sqrt{x} - \sqrt{y}$ , then

$$2\sqrt{x} = u + v \quad \text{and} \quad 2\sqrt{y} = u - v,$$

hence

$$x = \frac{1}{4}(u+v)^2 \quad \text{and} \quad y = \frac{1}{4}(u-v)^2.$$

Then we get the Jacobian

$$\begin{aligned} \frac{\partial(x, y)}{\partial(u, v)} &= \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{2}(u+v) & \frac{1}{2}(u+v) \\ \frac{1}{2}(u-v) & -\frac{1}{2}(u-v) \end{vmatrix} \\ &= \frac{1}{2}(u+v) \cdot \frac{1}{2}(u-v) \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} = -\frac{1}{2}(u^2 - v^2). \end{aligned}$$

We shall then find the domain of the new parameters  $A$ :

- 1) The boundary part  $x = 0$  corresponds to  $u + v = 0$ .
- 2) The boundary part  $y = 0$  corresponds to  $u - v = 0$ .
- 3) The boundary part  $\sqrt{x} + \sqrt{y} = 1$  corresponds to  $u = 1$ .

Since a closed and bounded set by the second main theorem of continuous functions is mapped into a closed and bounded set by this continuous change of variables, the new domain is the triangle  $A$  on the figure.




Figure 29.13: The domain  $A$  in the  $(U, V)$ -plane.

Note that the Jacobian is *negative* on  $A$ , so this time we shall need the absolute values in the formula.

By the transformation theorem,

$$\begin{aligned} \text{area}(B) &= \int_B dx dy = \int_A \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv = \frac{1}{2} \int_A (u^2 - v^2) du dv \\ &= \frac{1}{2} \int_0^1 \left\{ \int_{-u}^u (u^2 - v^2) dv \right\} du = \frac{1}{2} \int_0^1 \left[ u^2 v - \frac{1}{3} v^3 \right]_{-u}^u du \\ &= \frac{1}{2} \int_0^1 \left( 2u^3 - \frac{2}{3} u^3 \right) du = \frac{2}{3} \int_0^1 u^3 du = \frac{1}{6}, \end{aligned}$$

and

$$\begin{aligned} I &= \int_B \exp((\sqrt{x} + \sqrt{y})^4) dx dy = \frac{1}{2} \int_A \exp(u^4) \cdot (u^2 - v^2) du dv \\ &= \frac{1}{2} \int_0^1 \left\{ \int_{-u}^u \exp(u^4) \cdot (u^2 - v^2) dv \right\} du = \frac{2}{3} \int_0^1 \exp(u^4) \cdot u^3 du \\ &= \frac{1}{6} \int_0^1 e^t dt = \frac{e-1}{6}. \end{aligned}$$

**Example 29.8** Define a vector field  $\mathbf{r} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  in the following way,

$$\mathbf{r}(u, v) = (e^u \cos v, e^u \sin v).$$

Prove that the Jacobian  $J_{\mathbf{r}}$  is different from zero almost everywhere, and that  $\mathbf{r}$  is not injective.

**A** Jacobian and a non-injective transformation.

**D** Calculate and exploit the periodicity.

**I** The Jacobian is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} e^u \cos v & -e^u \sin v \\ e^u \sin v & e^u \cos v \end{vmatrix} = e^{2u} \neq 0.$$

Then note that  $(u, v) = (u_0, v_0 + 2p\pi)$ ,  $p \in \mathbb{Z}$ , are all mapped into the same point

$$(x, y) = (e^{u_0} \cos v_0, e^{u_0} \sin v_0),$$

so the transformation is not injective.

REMARK. We may add that  $\mathbb{R}^2$  by  $\mathbf{r}$  is mapped (infinitely often) onto  $\mathbb{R}^2 \setminus \{(0, 0)\}$ .

**Example 29.9** Define a vector field  $\mathbf{r} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  as follows:

$$\mathbf{r}(u, v) = (u^2 - v^2, 2uv).$$

Prove that the Jacobian  $J_{\mathbf{r}}$  is different from zero almost everywhere and that  $\mathbf{r}$  is not injective.

**A** Jacobian and a non-injective transformation.

**D** Calculate the Jacobian and find two different  $(u, v)$ -points which are mapped into the same  $(x, y)$ .

**I** The Jacobian is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 2u & -2v \\ 2v & 2u \end{vmatrix} = 4(u^2 + v^2) \neq 0 \quad \text{for } (u, v) \neq (0, 0).$$

Clearly,  $(u, v)$  and  $(-u, -v)$  are mapped into the same point,

$$(x, y) = (u^2 - v^2, 2uv),$$

so the map is not injective for  $(u, v) \neq (0, 0)$ .

**Example 29.10** Let  $B$  be the parallelogram of vertices  $(0, 0)$ ,  $(1, -1)$ ,  $(2, 1)$  and  $(3, 0)$ . Compute the plane integral

$$I = \int_B \frac{\cos(\frac{1}{2}\pi(x+y))}{1+x-2y} dx dy$$

by introducing the new variables

$$u = x + y, \quad v = x - 2y.$$

**A** Plane integral by a change of variables and the transformation formula.

**D** Sketch  $B$  and find the domain  $D$ . Calculate the Jacobian and insert into the formula.



The advertisement features a woman teacher smiling and interacting with two young students (a boy and a girl) who are looking at a laptop screen. The background is yellow with orange swirling lines. The e-Learning for Kids logo is in the top left. In the bottom right, there's a green oval containing text about their impact: "The number 1 MOOC for Primary Education", "Free Digital Learning for Children 5-12", and "15 Million Children Reached". Below this, a box contains the text: "About e-Learning for Kids" followed by a detailed description of the organization's mission and success.

**About e-Learning for Kids** Established in 2004, e-Learning for Kids is a global nonprofit foundation dedicated to fun and free learning on the Internet for children ages 5 - 12 with courses in math, science, language arts, computers, health and environmental skills. Since 2005, more than 15 million children in over 190 countries have benefitted from eLessons provided by EFK! An all-volunteer staff consists of education and e-learning experts and business professionals from around the world committed to making difference. eLearning for Kids is actively seeking funding, volunteers, sponsors and courseware developers; get involved! For more information, please visit [www.e-learningforkids.org](http://www.e-learningforkids.org).



Click on the ad to read more




Figure 29.14: The parallelogram  $B$ .

**I** It follows from the figure that

$$u = x + y \in [0, 3] \quad \text{and} \quad v = x - 2y \in [0, 3],$$

and the new domain is the square  $D = [0, 3] \times [0, 3]$ .

From

$$x = \frac{2}{3}u + \frac{1}{3}v \quad \text{and} \quad y = \frac{1}{3}u - \frac{1}{3}v,$$

follows that the Jacobian is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{1}{3} \end{vmatrix} = -\frac{1}{9}.$$

When we finally put everything into the transformation formula, then

$$\begin{aligned} I &= \int_B \frac{\cos(\frac{1}{2}\pi(x+y))}{1+x-2y} dx dy = \int_D \frac{\cos(\frac{1}{2}\pi \cdot u)}{1+v} \left| -\frac{1}{9} \right| du dv \\ &= \frac{1}{9} \int_0^3 \cos\left(\frac{\pi}{2}u\right) du \cdot \int_0^3 \frac{dv}{1+v} = \frac{1}{9} \cdot \frac{2}{\pi} \left[ \sin\left(\frac{\pi}{2}u\right) \right]_0^3 \cdot [\ln(1+v)]_0^3 \\ &= \frac{2}{9\pi} \left\{ \sin\left(\frac{3\pi}{2}\right) - 0 \right\} \cdot \{\ln 4 - \ln 1\} = -\frac{4}{9\pi} \ln 2. \end{aligned}$$

**Example 29.11** Let  $B$  be the plane set which is bounded by the  $X$ -axis and the line of equation  $y = x$  and an arc of the parabola given by

$$5x = 4 + y^2, \quad y \in [0, 1].$$

Calculate the plane integral

$$I = \int_B \cos \left[ \left( \sqrt{\frac{5}{4}x + y} + \sqrt{\frac{5}{4}x - y} \right)^4 \right] dx dy$$

by introducing the new variables  $(u, v)$  given by

$$5x = u^2 + v^2, \quad 2y = uv, \quad -u \leq v \leq u.$$

**A** Plane integral by a change of variables and the transformation formula.

**D** Sketch  $B$  and find the new domain  $D$ . Calculate the Jacobian and put everything into the formula.




Figure 29.15: The point set  $B$ .

**I** It follows from  $5x = u^2 + v^2$  and  $2y = uv$  that

$$5x + 4y = u^2 + v^2 + 2uv = (u + v)^2, \quad 5x - 4y = u^2 + v^2 - 2uv = (u - v)^2.$$

Since  $|v| \leq u$ , we get from this

$$u + v = +\sqrt{5x + 4y} \quad \text{and} \quad u - v = +\sqrt{5x - 4y},$$

hence

$$u = \frac{\sqrt{5x + 4y} + \sqrt{5x - 4y}}{2} \quad \text{and} \quad v = \frac{\sqrt{5x + 4y} - \sqrt{5x - 4y}}{2}.$$

Then we determine the boundary curves of the new domain.

1) If  $y = x$ ,  $x \in [0, 1]$ , then

$$u = \frac{\sqrt{9x} + \sqrt{x}}{2} = 2\sqrt{x} \quad \text{and} \quad v = \frac{\sqrt{9x} - \sqrt{x}}{2} = \sqrt{x},$$

so this boundary curve is transformed into  $v = \frac{1}{2}u$ . Then by a small consideration,  $u \in [0, 2]$ .

2) If  $y = 0$ ,  $x \in \left[0, \frac{4}{5}\right]$  on the  $X$ -axis, then  $v = 0$  and  $u = \sqrt{5x} \in [0, 2]$ .

3) If finally  $5x = 4 + y^2$ ,  $y \in [0, 1]$ , then

$$4 + y^2 = u^2 + v^2 \quad \text{and} \quad 4y = 2uv,$$

i.e.

$$(u + v)^2 = (y + 2)^2 \quad \text{and} \quad (u - v)^2 = (2 - y)^2,$$

thus

$$u + v = y + 2 \geq 0 \quad \text{and} \quad u - v = 2 - y \geq 0,$$

or  $u = 2$  and  $v = y \in [0, 1]$ . Then we can sketch the new domain (a triangle).




Figure 29.16: The new domain  $D$ .

Since

$$x = \frac{1}{5}(u^2 + v^2), \quad y = \frac{1}{2}uv,$$

we get the Jacobian

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{2}{5}u & \frac{2}{5}v \\ \frac{1}{2}v & \frac{1}{2}u \end{vmatrix} = \frac{1}{5}(u^2 - v^2) \geq 0.$$

Finally, since

$$\frac{5}{4}x + y = \frac{1}{4}(5x + 4y) = \frac{1}{4}(u^2 + v^2 + 2uv) = \left(\frac{u+v}{2}\right)^2,$$

and similarly,

$$\frac{5}{4}x - y = \left(\frac{u-v}{2}\right)^2,$$

we get the plane integral

$$\begin{aligned} I &= \int_B \cos \left[ \left( \sqrt{\frac{5}{4}x+y} + \sqrt{\frac{5}{4}x-y} \right)^4 \right] dx dy \\ &= \int_D \cos \left[ \left( \frac{u+v}{2} + \frac{u-v}{2} \right)^4 \right] \cdot \frac{1}{5} (u^2 - v^2) du dv = \int_D \cos(u^4) \cdot \frac{1}{5} (u^2 - v^2) du dv \\ &= \int_0^2 \cos(u^4) \left\{ \int_0^{\frac{u}{2}} \frac{1}{5} (u^2 - v^2) dv \right\} du = \int_0^2 \cos(u^4) \cdot \frac{1}{5} \left\{ u^2 \cdot \frac{u}{2} - \frac{1}{3} \cdot \left( \frac{u^3}{8} \right) \right\} du \\ &= \frac{1}{5} \cdot \frac{11}{24} \int_0^2 \cos(u^4) u^3 du = \frac{11}{480} [\sin(u^4)]_0^2 = \frac{11}{480} \sin 16. \end{aligned}$$



## Empowering People. Improving Business.

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

[www.bi.edu/master](http://www.bi.edu/master)



**Example 29.12** Let  $B$  be the plane point set which is bounded by the  $X$ -axis and the line of equation  $y = \frac{1}{2}x$ , and the branches of the hyperbola,

$$x^2 - y^2 = 1, \quad x > 0, \quad \text{and} \quad x^2 - y^2 = 4, \quad x > 0.$$

Compute the plane integral

$$I = \int_B \frac{x+y}{x-y} \exp(x^2 - y^2) \, dx \, dy$$

by introducing the new variables  $(u, v)$  given by

$$x = u \cosh v, \quad y = u \sinh v.$$

**A** Transformation of a plane integral.

**D** Sketch the domain  $B$ , and find the domain  $D$  of the new variables, and compute the Jacobian. Finally, insert everything into the transformation formula.




Figure 29.17: The domain  $B$ .

**I** If  $y = 0$ ,  $x > 0$ , then  $v = 0$  and  $x = u$ , hence the segment on the  $X$ -axis is transformed onto a segment on the  $U$ -axis.

If  $y = \frac{1}{2}x$ , then  $u \sinh v = \frac{1}{2}u \cosh v$ , i.e.

$$\tanh v = \frac{1}{2} = \frac{e^v - e^{-v}}{e^v + e^{-v}} = \frac{e^{2v} - 1}{e^{2v} + 1},$$

or  $e^{2v} + 1 = 2e^{2v} - 2$ , thus  $e^{2v} = 3$ , and hence  $v = \frac{1}{2} \ln 3$ , and  $u$  is a “free” variable.

If  $x^2 - y^2 = 1$   $x > 0$ , then  $u^2 = 1$ , and since  $u > 0$ , we must have  $u = 1$ .

If  $x^2 - y^2 = 4$ ,  $x > 0$ , then  $u^2 = 4$ , and since  $u > 0$ , we must have  $u = 2$ .

Summarizing, the new domain is the rectangle

$$D = [1, 2] \times \left[0, \frac{1}{2} \ln 3\right].$$

Then the Jacobian is computed,

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \cosh v & u \sinh v \\ \sinh v & u \cosh v \end{vmatrix} = u > 0.$$

By the transformation formula,

$$\begin{aligned} I &= \int_B \frac{x+y}{x-y} \exp(x^2 - y^2) dx dy = \int_D \frac{u(\cosh v + \sinh v)}{u \cosh v - \sinh v} \exp(u^2) \cdot u du dv \\ &= \int_D e^{2v} \exp(u^2) u du dv = \int_1^2 \exp(u^2) u du \cdot \int_0^{\frac{1}{2} \ln 3} e^{2v} dv \\ &= \frac{1}{2} [\exp(u^2)]_1^2 \cdot \frac{1}{2} [e^{2v}]_0^{\frac{1}{2} \ln 3} = \frac{1}{4} (e^4 - e) \cdot (3 - 1) = \frac{e}{2} (e^3 - 1). \end{aligned}$$

**Example 29.13** A triangle  $B$  in the  $(X, Y)$ -plane is given by the inequalities

$$x + y \geq 1, \quad 2y - x \leq 2, \quad y - 2x \geq -2.$$

By introducing

$$(29.5) \quad u = x + y, \quad v = x - y,$$

we get a map from the  $(X, Y)$ -plane onto the  $(U, V)$ -plane.

1) Prove that the image  $D$  in the  $(U, V)$ -plane of  $B$  by this map is given by

$$1 \leq u \leq 4, \quad u - 4 \leq 3v \leq 4 - u,$$

and sketch  $D$ .

2) Calculate the plane integral

$$\int_B \frac{3}{x+y} dx dy$$

by introducing the new variables given by (29.5).

**A** Transformation of a plane integral.

**D** Find the domain of the new variables  $D$  and compute the Jacobian, and then finally insert into the formula.

**I** 1) It follows from (29.13) that

$$x = \frac{u+v}{2} \quad \text{and} \quad y = \frac{u-v}{2}.$$

Hence




Figure 29.18: The new domain  $D$ .

- a)  $x + y \geq 1$  is transformed into  $u \geq 1$ ,
- b)  $2y - x \leq 2$  is transformed into  $u - 3v \leq 4$ ,
- c)  $y - 2x \geq -2$  is transformed into  $u + 3v \leq 4$ .

We get by a rearrangement,  $u - 4 \leq 3v \leq 4 - u$ , hence  $u \leq 4$ , and

$$D = \{(u, v) \mid 1 \leq u \leq 4, u - 4 \leq 3v \leq 4 - u\}.$$

We can here exploit that it is given that  $B$  is a triangle and thus bounded. The transformation (29.13) is continuous, so  $D$  is connected and bounded, and then we can sketch the three boundary lines and identify the image as the bounded part.

2) The Jacobian is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}.$$

Then by the transformation formula,

$$\begin{aligned} \int_B \frac{3}{x+y} dx dy &= \int_D \frac{3}{u} \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv = \frac{3}{2} \int_1^4 \left\{ \int_{-\frac{4-u}{3}}^{\frac{4-u}{3}} dv \right\} du \\ &= \frac{3}{2} \int_1^4 \frac{1}{u} \cdot 2 \cdot \frac{4-u}{3} du = \int_1^4 \left( \frac{4}{u} - 1 \right) du = 4 \ln 4 - 3 = 8 \ln 2 - 3. \end{aligned}$$

**Example 29.14** Let  $B$  be the bounded domain which is given by the inequalities

$$e^{-x} \leq y \leq 2e^{-x}, \quad e^x \leq y \leq 2e^x.$$

1. Sketch  $B$ .

If we put

$$(29.6) \quad u = y e^x, \quad v = y e^{-x},$$

we get a map of the  $(X, Y)$ -plane into the  $(U, V)$ -plane.

2. Prove that the image of  $B$  by this map is the square  $[1, 2] \times [1, 2]$ .

3. Calculate the plane integral

$$I = \int_B 4y^2 \exp(y^2 + x) \, dx \, dy$$

by introducing the new variables given by (29.6).

A Transformation of a plane integral.

D Follow the guidelines supplied by a calculation of the Jacobian before everything is put into the transformation formula.

ALTERNATIVELY, one can actually in this case compute the plane integral directly.

## Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!

Get Help Now



Go to [www.helpmyassignment.co.uk](http://www.helpmyassignment.co.uk) for more info

**Helpmyassignment**




Figure 29.19: The domain  $B$ .

**I 1)** Let us first find the intersection point of the boundary curves of  $B$ .

a) If  $y = e^{-x} = 2e^x$ , then  $x = -\frac{1}{2} \ln 2$  and hence  $y = \sqrt{2}$ .

b) If  $y = e^x = 2e^{-x}$ , then  $x = \frac{1}{2} \ln 2$  and hence  $y = \sqrt{2}$ .

c) The remaining two intersection points are immediately seen to be  $(0, 1)$  and  $(0, 2)$ .

Then it is easy to sketch the domain  $B$ , even if one does not have MAPLE at hand.

2) By the change of variables  $u = ye^x$  and  $v = ye^{-x}$ ,

a)  $y = e^x$  and  $x \in \left[0, \frac{1}{2} \ln 2\right]$  is transformed into  $v = 1$  and  $u = e^{2x} \in [1, 2]$ ,

b)  $y = 2e^{-x}$  and  $x \in \left[0, \frac{1}{2} \ln 2\right]$  is transformed into  $u = 2$  and  $v = 2e^{2x} \in [1, 2]$ ,

c)  $y = 2e^x$  and  $x \in \left[-\frac{1}{2} \ln 2, 0\right]$  is transformed into  $v = 2$  and  $u = 2e^{2x} \in [1, 2]$ ,

d)  $y = e^{-x}$  and  $x \in \left[-\frac{1}{2} \ln 2, 0\right]$  is transformed into  $u = 1$  and  $v = e^{-2x} \in [1, 2]$ .

Thus we get the new domain  $D = [1, 2] \times [1, 2]$  in the  $(U, V)$ -plane.

3) Then we find  $x$  and  $y$  as functions of  $u$  and  $v$ :

From  $y \geq 1$  and  $u, v \geq 1$ , follows that

$$\frac{u}{v} = e^{2x}, \quad \text{dvs.} \quad x = \frac{1}{2} \ln \left( \frac{u}{v} \right) = \frac{1}{2} \ln u - \frac{1}{2} \ln v.$$

From  $v = ye^{-x}$ , follows that

$$y = ve^x = v \sqrt{\frac{u}{v}} = \sqrt{uv}.$$

This gives the Jacobian

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{1}{2u} & -\frac{1}{2v} \\ \frac{1}{2}\sqrt{\frac{u}{v}} & \frac{1}{2}\sqrt{\frac{u}{v}} \end{vmatrix} = \frac{1}{4} \left( \frac{1}{\sqrt{uv}} + \frac{1}{\sqrt{uv}} \right) = \frac{1}{2} \frac{1}{\sqrt{uv}} > 0.$$

When we insert into the transformation formula, we get

$$\begin{aligned}
 I &= \int_B 4y^2 \exp(y^2 + x) dx dy = \int_D 4uv \exp\left(uv + \frac{1}{2} \ln\left(\frac{u}{v}\right)\right) \frac{1}{2} \frac{1}{\sqrt{uv}} du dv \\
 &= \int_D 4uv \exp(uv) \cdot \sqrt{\frac{u}{v}} \cdot \frac{1}{2} \frac{1}{\sqrt{uv}} du dv = \int_D 2u \exp(uv) du dv \\
 &= 2 \int_1^2 \left\{ \int_1^2 u \exp(uv) dv \right\} du = 2 \int_1^2 [\exp(uv)]_{v=1}^2 du \\
 &= 2 \int_1^2 (e^{2u} - e^u) du = [e^{2u} - 2e^u]_1^2 = e^4 - 2e^2 - e^2 + 2e = e^4 - 3e^2 + 2e.
 \end{aligned}$$

ALTERNATIVELY, it is actually possible to calculate the plane integral directly without using the transformation theorem. First write  $B = B_1 \cup B_2$ , as an (almost) disjoint union where

$$B_1 = \left\{ (x, y) \mid \sqrt{2} \leq y \leq 2, \ln\left(\frac{y}{2}\right) \leq x \leq \ln\left(\frac{2}{y}\right) \right\}$$

and

$$B_2 = \left\{ (x, y) \mid 1 \leq y \leq \sqrt{2}, \ln\left(\frac{1}{y}\right) \leq x \leq \ln y \right\}.$$

We have the following natural splitting,

$$\int_B 4y^2 \exp(y^2 + x) dx dy = I_1 + I_2,$$

where

$$\begin{aligned}
 I_1 &= \int_{B_1} 4y^2 \exp(y^2 + x) dx dy = \int_{\sqrt{2}}^2 \left\{ \int_{\ln(\frac{y}{2})}^{\ln(\frac{2}{y})} 4y^2 \exp(y^2) e^x dx \right\} dy \\
 &= \int_{\sqrt{2}}^2 4y^2 \exp(y^2) \cdot \left( \frac{2}{y} - \frac{y}{2} \right) dy = \int_{\sqrt{2}}^2 (8y - 2y^3) \exp(y^2) dy \\
 &= \int_{\sqrt{2}}^2 2(4 - y^2) \exp(y^2) 2y dy = \int_{t=2}^4 (4-t)e^t dt = [(5-t)e^t]_2^4 = e^4 - 3e^2,
 \end{aligned}$$

and

$$\begin{aligned}
 I_2 &= \int_{B_2} 4y^2 \exp(y^2 + x) dx dy = \int_1^{\sqrt{2}} \left\{ \int_{\ln(\frac{1}{y})}^{\ln y} 4y^2 \exp(y^2) e^x dx \right\} dy \\
 &= \int_1^{\sqrt{2}} 4y^2 \exp(y^2) \cdot \left( y - \frac{1}{y} \right) dy = \int_1^{\sqrt{2}} (4y^3 - 4y) \exp(y^2) dy \\
 &= \int_1^{\sqrt{2}} (2y^2 - 2) \exp(y^2) \cdot 2y dy = 2 \int_1^2 (t-1)e^t dt = 2[(t-2)e^t]_1^2 = 2e.
 \end{aligned}$$

Summarizing we get

$$\int_B 4y^2 \exp(y^2 + x) dx dy = I_1 + I_2 = e^4 - 3e^2 + 2e.$$

**Example 29.15** Let  $A$  denote the tetrahedron, which is bounded by the four planes of the equations

$$x + y = 1, \quad y + z = 1, \quad z + x = 1, \quad x + y + z = 1.$$

Calculate the space integral

$$I = \int_A (x+y)(y+z) dx dy dz$$

by introducing the new variables

$$u = 1 - x - y, \quad v = 1 - y - z, \quad w = 1 - z - x.$$

**A** Transformation of a space integral.

Find the Jacobian and the limits of  $u$ ,  $v$  and  $w$ .

**I** We derive from

$$dx dy dz = \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| du dv dw = \left| \frac{\partial(u, v, w)}{\partial(x, y, z)} \right|^{-1} du dv dw$$

and

$$\frac{\partial(u, v, w)}{\partial(x, y, z)} = \begin{vmatrix} -1 & -1 & 0 \\ 0 & -1 & -1 \\ -1 & 0 & -1 \end{vmatrix} = -1 - 1 + 0 - 0 - 0 - 0 = -2$$

that the weight function is

$$\left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| = \left| \frac{\partial(u, v, w)}{\partial(x, y, z)} \right|^{-1} = \frac{1}{2}.$$

The integrand is

$$(x+y)(y+z) = (1-u)(1-v).$$

Considering the limits of  $u$ ,  $v$  and  $w$  we see that

$$\begin{aligned} x + y = 1 &\text{ corresponds to } u = 1 - x - y = 0, \\ y + z = 1 &\text{ corresponds to } v = 1 - y - z = 0, \\ z + x = 1 &\text{ corresponds to } w = 1 - z - x = 0. \end{aligned}$$

From

$$u + v + w = 3 - 2(x + y + z),$$

follows that

$$x + y + z = 1 \text{ corresponds to } u + v + w = 1.$$

Finally, the tetrahedron lies in the first octant of the  $XYZ$ -space, where  $x + y \leq 1$ ,  $y + z \leq 1$  and  $z + x \leq 1$ . Hence the domain in the  $UVW$ -space is

$$\begin{aligned} B &= \{(u, v, w) \mid u \geq 0, v \geq 0, w \geq 0, u + v + w \leq 1\} \\ &= \{(u, v, w) \mid 0 \leq u \leq 1, 0 \leq v \leq 1 - u, 0 \leq w \leq 1 - u - v\}. \end{aligned}$$

By this transformation followed by a reduction in rectangular coordinates we get

$$\begin{aligned}
 I &= \int_A (x+y)(y+z) dx dy dz = \int_B (1-u)(1-v) \cdot \frac{1}{2} du dv dw \\
 &= \frac{1}{2} \int_0^1 (1-u) \left\{ \int_0^{1-u} (1-v) \left\{ \int_0^{1-u-v} dw \right\} dv \right\} du \\
 &= \frac{1}{2} \int_0^1 (1-u) \left\{ \int_0^{1-u} (1-v)(1-u-v) dv \right\} du \\
 &= \frac{1}{2} \int_0^1 (1-u) \left\{ \int_0^{1-u} \{(v-1)^2 + u(v-1)\} dv \right\} du \\
 &= \frac{1}{2} \int_0^1 (1-u) \left[ \frac{1}{3}(v-1)^3 + \frac{u}{2}(v-1)^2 \right]_0^{1-u} du \\
 &= \frac{1}{2} \int_0^1 (1-u) \left\{ \frac{1}{3} - \frac{1}{3}u^2 + \frac{u}{2} \cdot u^2 - \frac{u}{2} \right\} du \\
 &= \frac{1}{2} \int_0^1 (1-u) \left\{ \frac{1}{3} + \frac{1}{6}u^3 - \frac{u}{2} \right\} du = \frac{1}{12} \int_0^1 (1-u)(2+u^3-3u) du \\
 &= \frac{1}{12} \int_0^1 \{2+u^3-3u-2u-u^4+3u^2\} du = \frac{1}{12} \left\{ 2 + \frac{1}{4} - \frac{3}{2} - \frac{2}{2} - \frac{1}{5} + \frac{3}{3} \right\} \\
 &= \frac{1}{12} \cdot \frac{1}{30} (60 + 15 - 45 - 6) = \frac{1}{12} \cdot \frac{1}{30} \cdot 24 = \frac{1}{15}.
 \end{aligned}$$

## Brain power

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations.

Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering.  
 Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**



Click on the ad to read more

**Example 29.16** Let  $A$  be the closed point set in  $\mathbb{R}^3$ , which is bounded by the four elliptic paraboloids of the equations

$$(1) \quad z = \frac{3}{2} - \frac{1}{6}x^2 - \frac{2}{3}y^2$$

$$(2) \quad z = \frac{1}{2} - \frac{1}{2}x^2 - 2y^2,$$

$$(3) \quad z = -1 + \frac{1}{4}x^2 + y^2,$$

$$(4) \quad z = -2 + \frac{1}{8}x^2 + \frac{1}{2}y^2.$$

The point set  $A$  intersects the  $ZX$ -plane in a point set  $B_1$ , and the  $YZ$ -plane in a point set  $B_2$ .

- 1) Sketch  $B_1$  and  $B_2$ .
- 2) Calculate the volume  $\text{Vol}(A)$  and the space integral

$$I = \int_A \frac{1}{\sqrt{x^2 + 4y^2 + z^2}} dx dy dz$$

by introducing the new variables  $(u, v, w)$ , such that

$$x = \sqrt{uv} \cos w, \quad y = \frac{1}{2} \sqrt{uv} \sin w, \quad z = \frac{1}{2}(u - v),$$

where

$$u, v \in [0, +\infty[, \quad w \in [0, 2\pi].$$

**A** Transformation of a space integral.

**D** First sketch  $B_1$  (put  $y = 0$ ) and  $B_2$  (put  $x = 0$ ). Then apply the transformation formula, i.e. calculate the weight function and change variables.




Figure 29.20: The set  $B_1$  is the union of the two “oblique” quadrilateral sets.




Figure 29.21: The set  $B_2$  is the union of the two “oblique” quadrilateral sets.

I 1) By putting  $y = 0$ , we get in the  $XZ$ -plane the four parabolas

$$z = \frac{3}{2} - \frac{1}{6}x^2, \quad z = \frac{1}{2} - \frac{1}{2}x^2, \quad z = -1 + \frac{1}{4}x^2, \quad z = -2 + \frac{1}{8}x^2,$$

and it is easy to sketch  $B_1$ .

By putting  $x = 0$ , we get in the  $YZ$ -plane

$$z = \frac{3}{2} - \frac{2}{3}y^2, \quad z = \frac{1}{2} - 2y^2, \quad z = -1 + y^2, \quad z = -2 + \frac{1}{2}y^2,$$

and it is easy to sketch  $B_2$ .

2) Let

$$x = \sqrt{uv} \cos w, \quad y = \frac{1}{2}\sqrt{uv} \sin w, \quad z = \frac{1}{2}(u - v),$$

where  $u, v \geq 0$  and  $w \in [0, 2\pi]$ . We shall first find the image of  $A$  by this transformation.

a) By insertion into the boundary surface

$$z = \frac{3}{2} - \frac{1}{6}x^2 - \frac{2}{3}y^2$$

we get

$$\frac{1}{2}(u - v) = \frac{3}{2} - \frac{1}{6}uv \cos^2 w - \frac{1}{6}uv \sin^2 w = \frac{3}{2} - \frac{1}{6}uv,$$

hence  $3(u - v) = 9 - uv$ , which is reformulated as

$$uv + 3u = u(v + 3) = 9 + 3v = 3(v + 3).$$

It follows from  $v \geq 0$  that  $u = 3$ , hence this boundary surface is mapped into a part of the plane  $u = 3$ .

b) By insertion into the boundary surface

$$z = \frac{1}{2} - \frac{1}{2}x^2 - 2y^2$$

we get

$$\frac{1}{2}(u-v) = \frac{1}{2} - \frac{1}{2}uv\cos^2 w - \frac{1}{2}uv\sin^2 w = \frac{1}{2} - \frac{1}{2}uv,$$

i.e.  $u-v = 1-uv$ , and thus

$$uv + u = u(v+1) = v+1.$$

From  $v \geq 0$  follows that  $u = 1$ , hence the boundary surface is mapped into a part of the plane  $u = 1$ .

c) We get by insertion into the boundary surface

$$z = -1 + \frac{1}{4}x^2 + y^2$$

that

$$\frac{1}{2}(u-v) = -1 + \frac{1}{4}uv\cos^2 w + \frac{1}{4}uv\sin^2 w = -1 + \frac{1}{4}uv,$$

thus  $2(u-2) = uv - 4$ , and hence

$$uv + 2v = v(u+2) = 2u + 4 = 2(u+2).$$

It follows from  $u \geq 0$  that  $v = 2$ , so the boundary surface is mapped into a part of the plane  $v = 2$ .

d) We get by insertion into the boundary surface

$$z = -2 + \frac{1}{8}x^2 + \frac{1}{2}y^2$$

that

$$\frac{1}{2}(u-v) = -2 + \frac{1}{8}uv\cos^2 w + \frac{1}{8}uv\sin^2 w = -2 + \frac{1}{8}uv,$$

thus  $4(u-v) = -16 + uv$ , and hence

$$uv + 4v = v(u+4) = 4u + 16 = 4(u+4).$$

It follows from  $u \geq 0$  that  $v = 4$ , so the boundary surface is mapped into a part of the plane  $v = 4$ .

The only condition on  $w$  is that  $(\cos w, \sin w)$  shall encircle the unit circle only once, so  $w \in [0, 2\pi]$ .

By the transformation  $A$  is mapped onto the set

$$B = [1, 3] \times [2, 4] \times [0, 2\pi].$$

Then we calculate the Jacobian (for  $u, v > 0$ )

$$\begin{aligned}\frac{\partial(x, y, z)}{\partial(u, v, w)} &= \begin{vmatrix} \frac{1}{2}\sqrt{\frac{v}{u}}\cos w & \frac{1}{2}\sqrt{\frac{u}{v}}\cos w & -\sqrt{uv}\sin w \\ \frac{1}{4}\sqrt{\frac{v}{u}}\sin w & \frac{1}{4}\sqrt{\frac{u}{v}}\sin w & \frac{1}{2}\sqrt{uv}\cos w \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{vmatrix} \\ &= \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{1}{2} \sqrt{uv} \begin{vmatrix} \sqrt{\frac{v}{u}}\cos w & \sqrt{\frac{u}{v}}\cos w & -2\sin w \\ \sqrt{\frac{v}{u}}\sin w & \sqrt{\frac{u}{v}}\sin w & 2\cos w \\ 1 & -1 & 0 \end{vmatrix} \\ &= \frac{1}{16}\sqrt{uv} \cdot 2 \begin{vmatrix} \sqrt{\frac{v}{u}}\cos w & \left(\sqrt{\frac{u}{v}} + \sqrt{\frac{v}{u}}\right)\cos w & -\sin w \\ \sqrt{\frac{v}{u}}\sin w & \left(\sqrt{\frac{u}{v}} + \sqrt{\frac{v}{u}}\right)\sin w & \cos w \\ 1 & 0 & 0 \end{vmatrix} \\ &= \frac{1}{8}\sqrt{uv} \left( \sqrt{\frac{u}{v}} + \sqrt{\frac{v}{u}} \right) \begin{vmatrix} \cos w & -\sin w \\ \sin w & \cos w \end{vmatrix} = \frac{1}{8}(u+v) > 0.\end{aligned}$$



## What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvologistics.com](http://www.volvologistics.com). We look forward to getting to know you!

**VOLVO**  
AB Volvo (publ)  
[www.volvologistics.com](http://www.volvologistics.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA



Click on the ad to read more

We get by the transformation formula,

$$\begin{aligned}
 \text{Vol}(A) &= \int_A dx dy dz = \int_B \frac{\partial(x, y, z)}{\partial(u, v, w)} du dv dw = \int_0^{2\pi} \left\{ \int_2^4 \left\{ \int_1^3 \frac{1}{8} (u+v) du \right\} dv \right\} dw \\
 &= \frac{1}{8} \cdot 2\pi \int_2^4 \left[ \frac{u^2}{2} + uv \right]_{u=1}^3 dv = \frac{\pi}{8} \int_2^4 \{9 + 6v - 1 - 2v\} dv \\
 &= \frac{\pi}{8} \int_2^4 \{4v + 8\} dv = \frac{\pi}{8} [2v^2 + 8v]_2^4 = \frac{\pi}{4} [v^2 + 4v]_2^4 \\
 &= \frac{\pi}{4} \{16 + 16 - 4 - 8\} = \pi \{4 + 4 - 1 - 2\} = 5\pi.
 \end{aligned}$$

Let us turn to the space integral. Since

$$x^2 + 4y^2 + z^2 = uv \cos^2 w + uv \sin^2 w + \frac{1}{4} (u-v)^2 = \frac{1}{4} \{(u-v)^2 + 4uv\} = \frac{1}{4} (u+v)^2,$$

and  $u+v > 0$ , the integrand is transformed into

$$\frac{1}{\sqrt{x^2 + 4y^2 + z^2}} = \frac{2}{u+v}.$$

Finally, by the transformation formula,

$$\begin{aligned}
 \int_A \frac{1}{\sqrt{x^2 + 4y^2 + z^2}} dx dy dz &= \int_B \frac{1}{u+v} \cdot \frac{\partial(x, y, z)}{\partial(u, v, w)} du dv dw \\
 &= \int_B \frac{2}{u+v} \cdot \frac{u+v}{8} du dv dw = \frac{1}{4} \int_B du dv dw = \frac{1}{4} \text{Vol}(B) \\
 &= \frac{1}{4} \cdot 2 \cdot 2 \cdot 2\pi = 2\pi.
 \end{aligned}$$

**Example 29.17** We can write the formula of transformation of a space integral in the following way,

$$\int_{\tilde{\Omega}} f(\tilde{\mathbf{x}}) d\tilde{\Omega} = \int_{\Omega} f(\tilde{\mathbf{x}}(\mathbf{x})) |J(\mathbf{x})| d\Omega,$$

where  $J$  is the determinant of that matrix, the elements of which are

$$\frac{\partial \tilde{\mathbf{x}}_i}{\partial x_j}, \quad i, j \in \{1, 2, 3\}.$$

We shall in particular interpret  $\tilde{\Omega}$  as created from  $\Omega$  by a translation and a deformation. This means that to the point  $\mathbf{x}$  we let correspond a point  $\tilde{\mathbf{x}}$  given by

$$\tilde{\mathbf{x}} = \mathbf{x} + \mathbf{u}(\mathbf{x}),$$

where  $\mathbf{u}$  is the displacement vector field. When  $\mathbf{u}$  is a constant vector, we get a translation. However, in general  $\mathbf{u}$  varies in space (as indicated by the notation), such that we have a combination of a translation and a deformation.

- 1.** Compute  $J(x, y, z)$  by introducing  $\mathbf{u} = (u_x, u_y, u_z)$ .

In the Theory of Elasticity the deformations are often small in the sense that the derivative of  $\mathbf{u}$  is small, so we can reject all there products.

- 2.** Prove that by this assumption,  $J = 1 + \operatorname{div} \mathbf{u}$ .

- 3.** Finally, prove that the divergence is the relative increase of the volume corresponding to the deformation.

**A** Transformation of space integrals.

**D** Calculate the first approximation of the Jacobian.

**I** 1) The transformation is given by

$$\tilde{x}_1 = x + u_x(\mathbf{x}),$$

$$\tilde{x}_2 = y + u_y(\mathbf{x}), \quad \mathbf{x} = (x, y, z),$$

$$\tilde{x}_3 = z + u_z(\mathbf{x}),$$

hence

$$\begin{aligned} J(x, y, z) &= \begin{vmatrix} 1 + \frac{\partial u_x}{\partial x} & \frac{\partial u_y}{\partial x} & \frac{\partial u_z}{\partial x} \\ \frac{\partial u_x}{\partial y} & 1 + \frac{\partial u_y}{\partial y} & \frac{\partial u_z}{\partial y} \\ \frac{\partial u_x}{\partial z} & \frac{\partial u_y}{\partial z} & 1 + \frac{\partial u_z}{\partial z} \end{vmatrix} \\ &= 1 + \frac{\partial u_x}{\partial x} + \frac{\partial u_y}{\partial y} + \frac{\partial u_z}{\partial z} + \text{products of higher order.} \end{aligned}$$

2) If we remove all products of higher order, then we get

$$J = 1 + \operatorname{div} \mathbf{u}.$$

3) The geometrical interpretation of  $J$  is given by

$$d\tilde{\Omega} = |J(\mathbf{x})| d\Omega,$$

where  $d\tilde{\Omega}$  and  $d\Omega$  are considered as infinitesimal volumes corresponding to each other. This may possibly be clarified by

$$\Delta\tilde{\Omega} \approx |J(\mathbf{x})| \Delta\Omega.$$

Assume that  $\operatorname{div} \mathbf{u}$  is small, so higher order terms can be rejected. Then

$$J = 1 + \operatorname{div} \mathbf{u} > 0,$$

and thus

$$d\tilde{\Omega} = \{1 + \operatorname{div} \mathbf{u}\} d\Omega.$$

The factor  $1 + \operatorname{div} \mathbf{u}$  indicates the quotient between the two infinitesimal volumes, so  $\operatorname{div} \mathbf{u}$  can be interpreted as the relative signed increase of the volume.

**Example 29.18** Let  $A \subset \mathbb{R}^3$  be given by

$$0 \leq x, \quad 0 \leq y, \quad 0 \leq z, \quad \sqrt{x} + \sqrt{y} + \sqrt{z} \leq 1.$$

Compute the volume of  $A$  and the space integral

$$I = \int_A \exp \left[ (\sqrt{x} + \sqrt{y} + \sqrt{z})^6 \right] dx dy dz$$

by introducing the new variables

$$u = \sqrt{x} + \sqrt{y}, \quad v = \sqrt{x} - \sqrt{y}, \quad w = \sqrt{x} + \sqrt{y} + \sqrt{z}.$$

**A** Transformation of space integrals.

**D** Find the inverse transformation and compute the Jacobian before the transformation formula is applied.

**I** We derive from

$$2\sqrt{x} = u + v, \quad 2\sqrt{y} = u - v, \quad \sqrt{z} = w - u,$$

that

$$x = \frac{1}{4}(u+v)^2, \quad y = \frac{1}{4}(u-v)^2, \quad z = (w-u)^2.$$

Then find the parametric domain  $B$  in the  $(u, v, w)$ -space.

1) The boundary surface  $x = 0$  is mapped into the plane  $v = -u$ .




Figure 29.22: The domain  $A$ .

- 2) The boundary surface  $y = 0$  is mapped into the plane  $v = u$ .
- 3) The boundary surface  $z = 0$  is mapped into the plane  $w = u$ .
- 4) The boundary surface  $\sqrt{x} + \sqrt{y} + \sqrt{z} = 1$  is mapped into the plane  $w = 1$ .

The set  $A$  is closed and bounded, and the transformation is continuous. It therefore follows from the second main theorem for continuous functions that  $A$  is transformed into the closed and bounded parametric domain

$$B = \{(u, v, w) \mid 0 \leq w \leq 1, 0 \leq u \leq w, -u \leq v \leq u\}.$$

Then the Jacobian is given by

$$\begin{aligned} \frac{\partial(x, y, z)}{\partial(u, v, w)} &= \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix} = \begin{vmatrix} \frac{1}{2}(u+v) & \frac{1}{2}(u+v) & 0 \\ \frac{1}{2}(u-v) & -\frac{1}{2}(u-v) & 0 \\ -2(w-u) & 0 & 2(w-u) \end{vmatrix} \\ &= \frac{1}{2}(u+v) \cdot \frac{1}{2}(u-v) \cdot 2(w-u) \begin{vmatrix} 1 & 1 & 0 \\ 1 & -1 & 0 \\ -1 & 0 & 1 \end{vmatrix} \\ &= \frac{1}{2} (u^2 - v^2) (w-u) \cdot (-2) = -(u^2 - v^2) (w-u). \end{aligned}$$

Since  $v^2 \leq u^2$  and  $u \leq w$  in  $B$ , it follows from the transformation formula that

$$\begin{aligned}\text{vol}(A) &= \int_A d\Omega = \int_V |(-u^2 - v^2)(w - u)| du dv dw \\ &= \int_0^1 \left\{ \int_0^w \left[ \int_{-u}^u (u^2 - v^2)(w - u) dv \right] du \right\} dw \\ &= \int_0^1 \left\{ \int_0^{w(w-u)} \frac{4}{3} u^3 du \right\} dw = \int_0^1 \left\{ \frac{4}{3} \int_0^w (wu^3 - u^4) du \right\} dw \\ &= \int_0^1 \frac{4}{3} \left[ \frac{1}{4} wu^4 - \frac{1}{5} u^5 \right]_{u=0}^w dw = \int_0^1 \frac{1}{15} w^5 dw = \frac{1}{90},\end{aligned}$$

and

$$\begin{aligned}I &= \int_A \exp \left[ (\sqrt{x} + \sqrt{y} + \sqrt{z})^6 \right] d\Omega \\ &= \int_0^1 \exp(w^6) \left\{ \int_0^w \left[ \int_{-u}^u (u^2 - v^2)(w - u) dv \right] du \right\} dw \\ &= \int_0^1 \frac{1}{15} \exp(w^6) \cdot w^5 dw = \frac{1}{90} \int_0^1 e^t dt = \frac{e-1}{90},\end{aligned}$$

where we also found that

$$\int_0^w \left\{ \int_{-u}^u (u^2 - v^2)(w - u) dv \right\} du = \frac{1}{15} w^5.$$

**Example 29.19** Let  $B$  be the triangle given by  $x \geq 0$ ,  $y \geq 0$ ,  $x + y \leq 1$ . Compute the improper plane integral

$$I = \int_B \exp\left(\frac{x-y}{x+y}\right) dS$$

by introducing the new variables  $(u, v) = (x + y, x - y)$ .

**A** Transformation of an improper plane integral.

**D** The integrand is not defined at  $(x, y) = (0, 0) \in B$ . Otherwise, the integrand is positive, so in the worst case we shall only get that the value becomes  $+\infty$ .

Find  $x$  and  $y$  expressed by  $u$  and  $v$ . Find the parametric domain in the  $(u, v)$ -plane. Compute the Jacobian. Finally, insert into the transformation formula, check if the singularity has any effect and compute.




Figure 29.23: The domain of integration  $B$ .

**I** The transformation is continuous with a continuous inverse:

$$\begin{aligned} u &= x + y && \text{and} & v &= x - y, \\ x &= \frac{1}{2}(u + v) && \text{and} & y &= \frac{1}{2}(u - v). \end{aligned}$$

Furthermore,  $B$  is closed and bounded, so by using the second main theorem for continuous functions we conclude that the image, i.e. the new parametric domain  $A$  in the  $(u, v)$ -plane is also closed and bounded. It therefore suffices to find the images of the boundary curves.

- 1)  $x = 0$  is mapped into  $u + v = 0$ , i.e. into  $v = -u$ .
- 2)  $y = 0$  is mapped into  $u - v = 0$ , i.e. into  $v = u$ .
- 3)  $x + y = 1$  is mapped into  $u = 1$ .

It follows that  $A$  in the  $(u, v)$ -plane is the triangle which is defined by these three lines, hence

$$A = \{(u, v) \mid 0 \leq u \leq 1, -u \leq v \leq u\}.$$

Then compute the Jacobian,




Figure 29.24: The parametric domain  $A$  in the  $(u, v)$ -plane.

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}.$$

Finally, by putting into the transformation formula where we also have in mind that the integral is an improper one of a positive integrand:

$$\begin{aligned} I &= \int_B \exp\left(\frac{x-y}{x+y}\right) dS = \int_A \exp\left(\frac{v}{u}\right) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv \\ &= \frac{1}{2} \int_0^1 \left\{ \int_{-u}^u \exp\left(\frac{v}{u}\right) dv \right\} du = \frac{1}{2} \lim_{\varepsilon \rightarrow 0+} \int_\varepsilon^1 \left\{ \int_{-u}^u \exp\left(\frac{v}{u}\right) dv \right\} du \\ &= \frac{1}{2} \lim_{\varepsilon \rightarrow 0+} \int_\varepsilon^1 u \left[ \exp\left(\frac{v}{u}\right) \right]_{v=-u}^u du = \frac{1}{2} \lim_{\varepsilon \rightarrow 0+} \int_\varepsilon^1 u (e - e^{-1}) du \\ &= \sinh 1 \int_0^1 u du = \frac{\sinh 1}{2}. \end{aligned}$$

**Example 29.20** Let  $A$  be the tetrahedron which is bounded by the four planes of the equations

$$x + y + z = 0, \quad x + y - z = 0, \quad x - y - z = 0, \quad 2x - z = 1.$$

Calculate the space integral

$$I = \int_A (x + y + z)(x + y - z)(x - y - z) dx dy dz$$

by introducing the new variables

$$u = x + y + z, \quad v = x + y - z, \quad w = x - y - z.$$

**A** Transformation of a space integral.

**D** Find  $x, y, z$  expressed by  $u, v, w$ . Then find the parametric domain  $B$  in the  $(u, v, w)$ -space which is uniquely mapped onto  $A$ . Compute the Jacobian, and finally, apply the transformation formula.

**I** It follows from

$$u = x + y + z, \quad v = x + y - z, \quad w = x - y - z,$$

that

$$u + w = 2x, \quad \text{i.e.} \quad x = \frac{1}{2}(u + w).$$

Then

$$u - v = 2z, \quad \text{i.e.} \quad z = \frac{1}{2}(u - v)$$

and

$$v - w = 2y, \quad \text{i.e.} \quad y = \frac{1}{2}(v - w).$$

Summarizing,

$$\begin{aligned} u &= x + y + z, & v &= x + y - z, & w &= x - y - z, \\ x &= \frac{1}{2}(u + w), & y &= \frac{1}{2}(v - w), & z &= \frac{1}{2}(u - v), \end{aligned}$$

and the Jacobian is

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix} = \begin{vmatrix} \frac{1}{2} & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{vmatrix} = \frac{1}{8} \begin{vmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 1 & -1 & 0 \end{vmatrix} = \frac{1}{8}(-1 - 1) = -\frac{1}{4}.$$

We shall find the images of the boundary surfaces of the tetrahedron:

- 1)  $x + y + z = 0$  is mapped into  $u = 0$ .
- 2)  $x + y - z = 0$  is mapped into  $v = 0$ .
- 3)  $x - y - z = 0$  is mapped into  $w = 0$ .
- 4)  $2x - z = 1$ , i.e.  $2 = 4x - 2z$ , is mapped into

$$2 = 2u + 2w - u + v = u + v + 2w, \quad \text{i.e.} \quad u + v + 2w = 2.$$




Figure 29.25: The transformed parametric domain  $B$ .

The inverse transformation is continuous, and  $A$  is closed and bounded. Hence,  $A$  is transformed into a new tetrahedron  $B$  as indicated on the figure. Note that  $B$  is cut at the height  $w \in [0, 1[$  in the triangle

$$B(w) = \{(u, v) \mid u \geq 0, v \geq 0, u + v \leq 2(1 - w)\}.$$

This can be exploited in the calculation of the transformed integral by the method of slicing.

This e-book  
*is made with*  
**SetaPDF**


**SETASIGN**





PDF components for PHP developers

[www.setasign.com](http://www.setasign.com)



According to the transformation theorem,

$$\begin{aligned} I &= \int_A (x+y+z)(x+y-z)(x-y-z) dx dy dz \\ &= \int_B uvw \left| \frac{\partial(x,y,z)}{\partial(u,v,w)} \right| du dv dw = \frac{1}{4} \int_0^1 w \left\{ \int_{B(w)} uv dS \right\} dw. \end{aligned}$$

Then calculate the integral over the slice at height  $w$ ,

$$\begin{aligned} \int_{B(w)} uv dS &= \int_0^{2(1-w)} u \left\{ \int_0^{2(1-w)-u} v dv \right\} du = \int_0^{2(1-w)} u \cdot \frac{1}{2} \{2(1-w) - u\}^2 du \\ &= \frac{1}{2} \int_0^{2(1-w)} \{4(1-w)^2 u - 4(1-w)u^2 + u^3\} du \\ &= \frac{1}{2} \left[ 2(1-w)^2 u^2 - \frac{4}{3}(1-w)u^3 + \frac{1}{4}u^4 \right]_0^{2(1-w)} \\ &= \frac{1}{2} \left\{ 2(1-w)^2 \cdot 4(1-w)^2 - \frac{4}{3}(1-w) \cdot 8(1-w)^3 + \frac{1}{4} \cdot 16(1-w)^4 \right\} \\ &= \frac{1}{2} (1-w)^4 \left\{ 8 - \frac{32}{3} + 4 \right\} = \frac{1}{2} \cdot \frac{4}{3} (1-w)^4 = \frac{2}{3} (w-1)^4. \end{aligned}$$

Finally, by insertion,

$$\begin{aligned} I &= \frac{1}{4} \int_0^1 w \left\{ \int_{B(w)} uv dS \right\} dw = \frac{1}{4} \cdot \frac{2}{3} \int_0^1 w(w-1)^4 dw \\ &= \int 16 \int_0^1 \{(w-1)^5 + (w-1)^4\} dw = \frac{1}{6} \left[ \frac{1}{6}(w-1)^6 + \frac{1}{5}(w-1)^5 \right]_0^1 \\ &= \frac{1}{6} \left\{ -\frac{1}{6} - \frac{1}{5}(-1) \right\} = \frac{1}{6} \left( \frac{1}{5} - \frac{1}{6} \right) = \frac{1}{180}. \end{aligned}$$

**Example 29.21** Let  $K$  denote the closed ball of centrum  $(1, 1, 1)$  and radius  $\sqrt{3}$ . We construct a subset  $A \subset K$  by only keeping those points from  $K$  in  $A$ , which furthermore satisfy  $r \geq 1$  and lie in the first octant.

Calculate the space integral

$$I = \int_A \frac{1}{r^6} d\Omega$$

by introducing the new variables

$$u = \frac{x}{r^2}, \quad v = \frac{y}{r^2}, \quad w = \frac{z}{r^2}.$$

- A** Transformation of a space integral. This is the ‘simplest’ non-trivial example in the three dimensional space. We shall see that even in this case the computations grow very big.
- D** First find  $A$ , and then the parametric domain  $D$  of the variables  $(u, v, w)$ . Compute the Jacobian, and finally also the transformed integral.




Figure 29.26: The boundary surface of  $A$  in each of the three planes  $x = 0$ ,  $y = 0$ , or  $z = 0$ .

**I** The set  $A$  is described by

$$A = \{(x, y, z) \mid (x-1)^2 + (y-1)^2 + (z-1)^2 \leq 3, x^2 + y^2 + z^2 \geq 1, x \geq 0, y \geq 0, z \geq 0\}.$$

The boundary surface in each of the planes  $x = 0$ ,  $y = 0$  and  $z = 0$  is indicated on the figure.

Then check the image in the  $(u, v, w)$ -space of each of the boundary surfaces in the  $(x, y, z)$ -space.

Clearly, the boundary surface  $x^2 + y^2 + z^2 = 1$  is mapped into  $u^2 + v^2 + w^2 = 1$ , and they both lie in the first octant.

Then check the transformation of the boundary surface

$$(x-1)^2 + (y-1)^2 + (z-1)^2 = 3.$$

If we put

$$R^2 = u^2 + v^2 + w^2 = \frac{x^2}{r^4} + \frac{y^2}{r^4} + \frac{z^2}{r^4} = \frac{1}{r^2},$$

then

$$x = u \cdot r^2 = \frac{u}{R^2}, \quad y = \frac{v}{R^2}, \quad z = \frac{w}{R^2},$$

hence by insertion

$$(u - R^2)^2 + (v - R^2)^2 + (w - R^2)^2 = 3R^4,$$

and thus by a computation

$$\begin{aligned} 3R^4 &= u^2 - 2uR^2 + R^4 + v^2 - 2vR^2 + R^4 + w^2 - 2wR^2 + R^4 \\ &= (u^2 + v^2 + w^2) - 2(u + v + w)R^2 + 3R^4 \\ &= R^2 - 2(u + v + w)R^2 + 3R^4 = 3R^4 + R^2\{1 - 2(u + v + w)\}. \end{aligned}$$

Now,  $R^2 = u^2 + v^2 + w^2 = \frac{1}{r^2} > 0$ , to this is reduced to the equation of a plane surface in the first octant,

$$u + v + w = \frac{1}{2}.$$




Figure 29.27: The domain  $D$  lies in the first octant between the two surfaces.

We conclude that  $D$  is that part of the closed first octant, which also lies between the plane  $u + v + w = \frac{1}{2}$  and the sphere  $u^2 + v^2 + w^2 = 1$ .

Since we have

$$u + v + w = R\{\sin\theta(\cos\varphi + \sin\varphi) + \cos\theta\}$$

in spherical coordinates

$$u = R \sin\theta \cos\varphi, \quad v = R \sin\theta \sin\varphi, \quad w = R \cos\theta,$$

we get the following description of  $D$  in spherical coordinates

$$D = \left\{ (R, \varphi, \theta) \mid [2\{\sin \theta(\cos \varphi + \sin \varphi) + \cos \theta\}]^{-1} \leq R \leq 1, 0 \leq \varphi, \theta \leq \frac{\pi}{2} \right\}.$$

Then calculate the Jacobian  $\frac{\partial(x, y, z)}{\partial(u, v, w)}$ , where we use that

$$(x, y, z) = \left( \frac{u}{R^2}, \frac{v}{R^2}, \frac{w}{R^2} \right).$$

First note that e.g.

$$\frac{\partial}{\partial u} \left( \frac{1}{R^2} \right) = -\frac{1}{R^4} \cdot \frac{\partial R^2}{\partial u} = -\frac{2u}{R^4},$$

and similarly of symmetric reasons,

$$\frac{\partial}{\partial v} \left( \frac{1}{R^2} \right) = -\frac{2v}{R^4} \quad \text{and} \quad \frac{\partial}{\partial w} \left( \frac{1}{R^2} \right) = -\frac{2w}{R^4},$$



[www.sylvania.com](http://www.sylvania.com)

We do not reinvent the wheel we reinvent light.

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM

**OSRAM**  
**SYLVANIA**



Click on the ad to read more

Then

$$\begin{aligned}
 \frac{\partial(x, y, z)}{\partial(u, v, w)} &= \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix} = \begin{vmatrix} \frac{1}{R^2} - \frac{2u^2}{R^4} & -\frac{2uv}{R^4} & -\frac{2uw}{R^4} \\ -\frac{2uv}{R^4} & \frac{1}{r^2} - \frac{2v^2}{R^4} & -\frac{2vw}{R^4} \\ -\frac{2uw}{R^4} & -\frac{2vw}{R^4} & \frac{1}{R^2} - \frac{2w^2}{R^4} \end{vmatrix} \\
 &= \frac{1}{R^{12}} \begin{vmatrix} -u^2 + v^2 + w^2 & -2uv & -2uw \\ -2uv & u^2 - v^2 + w^2 & -2vw \\ -2uw & -2vw & u^2 + v^2 - w^2 \end{vmatrix} \\
 &= \frac{1}{R^{12}} \left\{ (R^2 - 2u^2)(R^2 - 2v^2)(R^2 - 2w^2) - 8u^2v^2w^2 - 8u^2v^2w^2 \right. \\
 &\quad \left. - 4u^2w^2(R^2 - 2v^2) - 4v^2w^2(R^2 - 2u^2) - 4u^2v^2(R^2 - 2w^2) \right\} \\
 &= \frac{1}{R^{12}} \left\{ R^6 - 2(u^2 + v^2 + w^2)R^4 + 4R^2(u^2v^2 + u^2w^2 + v^2w^2) - 24u^2v^2w^2 \right. \\
 &\quad \left. - 4R^2(u^2w^2 + v^2w^2 + u^2v^2) + 8u^2v^2w^2 + 8u^2v^2w^2 + 8u^2v^2w^2 \right\} \\
 &= \frac{1}{R^{12}} \{-R^6\} = -\frac{1}{R^6}.
 \end{aligned}$$

Finally, we get by the transformation theorem and a consideration of a volume that

$$\begin{aligned}
 I &= \int_A \frac{1}{r^6} d\Omega = \int_D R^6 \cdot \frac{1}{R^6} d\omega = \int_D d\omega = \text{vol}(D) \\
 &= \frac{1}{8} \cdot \frac{4\pi}{3} - \frac{1}{3} \cdot \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{\pi}{6} - \frac{1}{48},
 \end{aligned}$$

in which the slicing method is latently applied.

## 30 Improper integrals

### 30.1 Introduction

In some cases it is possible to extend the various forms of integrals considered in the previous volumes to situations which are not covered by the theorems already given. We considered earlier (abstract) integrals symbolized by

$$\int_A f(\mathbf{x}) d\mu,$$

where  $A$  is closed and bounded, and the function  $f : A \rightarrow \mathbb{R}$  is continuous. We shall in this chapter investigate what can be done if these conditions above are not all fulfilled.

Let us start with a useful technical trick, which will be very important in the discussion below. We shall split the function  $f$  under consideration into its positive part  $f_+$  and its negative part  $f_-$ , and then continue by only discussing the case where  $f$  is nonnegative.

Given any function  $f : A \rightarrow \mathbb{R}$ , we define  $f_+$  and  $f_-$  in the following way,

$$f_+(\mathbf{x}) := \begin{cases} f(\mathbf{x}), & \text{if } f(\mathbf{x}) > 0, \\ 0, & \text{if } f(\mathbf{x}) \leq 0, \end{cases} \quad f_-(\mathbf{x}) := \begin{cases} -f(\mathbf{x}), & \text{if } f(\mathbf{x}) < 0, \\ 0, & \text{if } f(\mathbf{x}) \geq 0. \end{cases}$$

Then clearly both  $f_+$  and  $f_-$  are *nonnegative*, and

$$f(\mathbf{x}) = f_+(\mathbf{x}) - f_-(\mathbf{x}) \quad \text{and} \quad f_+(\mathbf{x}) \cdot f_-(\mathbf{x}) = 0 \quad \text{for all } \mathbf{x} \in A.$$

Furthermore, the splitting of  $f$ , satisfying the two conditions above, is unique.

Now, if we can prescribe (finite) values of both  $\int_A f_+(\mathbf{x}) d\mu$  and  $\int_A f_-(\mathbf{x}) d\mu$ , then we simply define the value of the improper integral  $\int_A f(\mathbf{x}) d\mu$  by putting

$$\int_A f(\mathbf{x}) d\mu = \int_A f_+(\mathbf{x}) d\mu - \int_A f_-(\mathbf{x}) d\mu.$$

This means that we in the following only shall consider *nonnegative integrands*,  $f \geq 0$ , when we extend the plane, space, line or surface integrals to the case of improper integrals, where either  $f$  is not continuous, or  $A$  is either not bounded or not closed.

Once we have restricted the investigation to nonnegative functions  $f \geq 0$ , the idea of defining the improper integral  $\int_A f(\mathbf{x}) d\mu$  goes as follows. We choose a sequence of *nested closed and bounded sets*, i.e.

$$A_1 \subset A_2 \subset \cdots \subset A_n \subset \cdots \subset A,$$

such that the given nonnegative function  $f$  is continuous on every  $A_n$ , and such that  $A_n \rightarrow A$  in some sense for  $n \rightarrow +\infty$ . Since  $f$  is continuous on every  $A_n$ , and  $A_n$  is closed and bounded, the integral of  $f$  over each  $A_n$  does indeed exist, and since  $f \geq 0$  is nonnegative, and the sequence  $A_n$  is nested, i.e.  $A_n \subset A_{n+1}$  for every  $n \in \mathbb{N}$ , we conclude that the sequence

$$\int_{A_1} f(\mathbf{x}) d\mu \leq \int_{A_2} f(\mathbf{x}) d\mu \leq \cdots \leq \int_{A_n} f(\mathbf{x}) d\mu \leq \cdots,$$

of real numbers is (weakly) increasing, hence it is either convergent, or tending towards  $+\infty$ . This is the reason for only considering nonnegative functions.

It is not hard to see that any nested increasing sequence of closed and bounded subsets of  $A$ , on which  $f \geq 0$  is continuous, will do. We still have to explain, what is meant by  $A_n \rightarrow A$  for  $n \rightarrow +\infty$ .

If

$$\bigcup_{n=1}^{+\infty} A_n = A,$$

then the immediate definition of the improper integral of  $f$  over  $A$  becomes

$$\int_A f(\mathbf{x}) d\mu = \lim_{n \rightarrow +\infty} \int_{A_n} f(\mathbf{x}) d\mu.$$

If the limit is convergent towards the value  $I \in \mathbb{R}$ , we say that the improper integral is convergent and

$$\int_A f(\mathbf{x}) d\mu = I.$$

If the limit is  $+\infty$ , we say that the improper integral is divergent.



**Deloitte.**

Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

© Deloitte & Touche LLP and affiliated entities.



Click on the ad to read more

However, measures are only determined modulo null sets, so we may weaken the definition to the requirement that only

$$\bigcup_{n=1}^{+\infty} A_n = A \setminus M, \quad \text{where } \mu(M) = 0, \quad \text{i.e. } M \text{ is a null set.}$$

Then we can still use the definitions above of the improper integral, because the “missing values” from  $f(\mathbf{x})$  over the null set  $M$  have by definition only the weight 0, so we only exclude 0.

### 30.2 Theorems for proper integrals

A further analysis of the improper integrals shows that we can divide those considered here into two types, which may not be disjoint,

- 1) Unbounded continuous integrand.
- 2) Unbounded domain of integration.

Only these cases will be relevant for us, although the classification above is not complete. There exist improper integrals, which cannot be broken down to a finite number of cases of the types above, but they will be outside the realm of this chapter.

The proofs of the following theorems have already been sketched in Section 30.1. We shall be satisfied with these sketches and not bother with more precise proofs.

**Theorem 30.1 . Improper integral with unbounded integrand.** Let  $A \subset \mathbb{R}^k$  be a bounded set, and let  $M \subset A$  be a null set,  $\mu(M) = 0$ . Assume that

$$f : A \setminus M \rightarrow [0, +\infty[$$

is a nonnegative continuous and unbounded function. Choose an increasing (nested) sequence of sets

$$A_1 \subset A_2 \subset \cdots \subset A_n \subset \cdots \subset A,$$

such that

$$M \subset A \setminus A_n \quad \text{for all } n \in \mathbb{N}, \quad \text{and} \quad \mu(A \setminus A_n) \rightarrow 0 \quad \text{for } n \rightarrow +\infty.$$

Then either

$$\lim_{n \rightarrow +\infty} \int_{A_n} f(\mathbf{x}) d\mu = +\infty,$$

and we call the improper integral divergent, or,

$$\lim_{n \rightarrow +\infty} \int_{A_n} f(\mathbf{x}) d\mu = I < +\infty,$$

is convergent, in which case we say that the improper integral is convergent of the value

$$\int_A f(\mathbf{x}) d\mu := \lim_{n \rightarrow +\infty} \int_{A_n} f(\mathbf{x}) d\mu.$$

**Theorem 30.2 Improper integral with unbounded domain of integration.** Assume that  $A \subseteq \mathbb{R}^k$  is unbounded, and that  $f : A \rightarrow \mathbb{R}$  is a nonnegative and continuous function. Choose an increasing (nested) sequence of closed and bounded sets,

$$A_1 \subset A_2 \subset \cdots \subset A_n \subset \cdots \subset A, \quad \text{such that } \bigcup_{n=1}^{+\infty} A_n = A.$$

then either

$$\lim_{n \rightarrow +\infty} \int_{A_n} f(\mathbf{x}) d\mu = +\infty,$$

and the improper integral of  $f$  over  $A$  is said to be divergent, or it is convergent of the value

$$\int_A f(\mathbf{x}) d\mu = \lim_{n \rightarrow +\infty} \int_{A_n} f(\mathbf{x}) d\mu < +\infty.$$

In the applications we usually have to combine these two theorems.

### 30.3 Procedure for improper integrals; bounded domain

**Problem 30.1** Calculate the integral

$$\int_A f(\mathbf{x}) dS,$$

where the domain  $A$  is bounded, and where either  $A$  is not closed, or the integrand  $f(\mathbf{x})$  is not defined for all  $\mathbf{x} \in A$ .

We choose in this description for convenience  $A \subset \mathbb{R}^2$ . The procedure is analogous for any  $A \subset \mathbb{R}^k$ .

**Idea:** Approximate  $\int_A f(\mathbf{x}) dS$  by  $\int_{A_n} f(\mathbf{x}) dS$ , where

- 1) The function  $f(\mathbf{x})$  is continuous for all  $x \in A_n$ .
- 2) All sets  $A_n \subset A$  are closed and bounded,  $n \in \mathbb{N}$ .
- 3)  $\text{area}(A \setminus A_n) \rightarrow 0$  for  $n \rightarrow +\infty$ .

**Remark 30.1** Even if we are approaching the solution by this idea, it is not enough. We shall below supply it with some sufficient conditions in the description of the procedure.  $\diamond$

#### Procedure.

- 1) Examine whether  $f(\mathbf{x})$  has a continuous extension  $F(\mathbf{x})$  to  $\overline{A}$ . If this is the case, then

$$\int_A f(\mathbf{x}) dS = \int_{\overline{A}} F(\mathbf{x}) dS,$$

where the right hand side is calculated by well-known methods, and the problem is solved.

- 2) If  $f(\mathbf{x})$  does not have a continuous extension to  $\overline{A}$ , we continue by an *analysis of the sign of  $f(\mathbf{x})$* . This step is extremely *important!* Divide  $A$  into two subsets

$$A^+ = \{(x, y) \in A \mid f(x, y) \geq 0\} \quad \text{and} \quad A^- = \{(x, y) \in A \mid f(x, y) \leq 0\}.$$

If  $f(x, y)$  is continuous where it is defined, then this division is most easily performed by finding the *null curves* i.e. the curves where  $f(x, y) = 0$ . These curves divide the domain into open sub-domains. Due to the continuity the sign of  $f$  is constant in each of these sub-domains, so in order to find the sign one just has to apply  $f$  to one point in each sub-domain. The curves where  $f(x, y) = 0$  can afterwards be included in any of the two sets  $A^+$  or  $A^-$ , and even in both of them, because since the integrand is 0 on these curves they will not contribute to the final result.

- 3) Let us first consider  $A^+$ . If  $A^+ = \emptyset$ , then go to 4) below. If on the other hand  $A^+ \neq \emptyset$ , we choose a convenient increasing sequence of *closed and bounded* subsets  $A_\varepsilon^+ \subset A^+$ , such that

$$\text{area}(A^+ \setminus A_\varepsilon^+) \rightarrow 0 \quad \text{for } \varepsilon \rightarrow 0+.$$

Calculate

$$I_\varepsilon^+ = \int_{A_\varepsilon^+} f(x, y) dS \quad (\geq 0).$$

If  $I_\varepsilon^+ \rightarrow +\infty$  for  $\varepsilon \rightarrow 0+$ , then the improper integral is *divergent*.

If on the other hand  $\lim_{\varepsilon \rightarrow 0} I_\varepsilon^+ < +\infty$  (there are only these two possibilities, because  $I_\varepsilon^+$  increases, when  $\varepsilon$  decreases towards 0), then

$$\int_{A^+} f(x, y) dS = \lim_{\varepsilon \rightarrow 0^+} \int_{A_\varepsilon^+} f(x, y) dS \geq 0.$$

- 4) Then consider the other subset  $A^-$ . If  $A^- = \emptyset$ , go to 5).

If on the other hand,  $A^- \neq \emptyset$ , choose a convenient increasing sequence of *closed and bounded* subsets  $A_\varepsilon^- \subset A^-$ , such that

$$\text{area}(A^- \setminus A_\varepsilon^-) \rightarrow 0 \quad \text{for } \varepsilon \rightarrow 0^+.$$

Calculate

$$I_\varepsilon^- = \int_{A_\varepsilon^-} f(x, y) dS \quad (\leq 0).$$

If  $I_\varepsilon^- \rightarrow -\infty$  for  $\varepsilon \rightarrow 0$ , then the improper integral is *divergent*.

If instead  $\lim_{\varepsilon \rightarrow 0} I_\varepsilon^- > -\infty$  (there are only these two possibilities), then

$$\int_{A^-} f(x, y) dS = \lim_{\varepsilon \rightarrow 0^+} \int_{A_\varepsilon^-} f(x, y) dS \quad (\leq 0).$$

- 5) It is only when we have obtained *convergence* in both 3) and 4) that we can conclude that the improper integral is *convergent* with the value

$$\int_A f(x, y) dS = \int_{A^+} f(x, y) dS + \int_{A^-} f(x, y) dS.$$

**WHY WAIT FOR PROGRESS?**

**DARE TO DISCOVER**

Discovery means many different things at Schlumberger. But it's the spirit that unites every single one of us. It doesn't matter whether they join our business, engineering or technology teams, our trainees push boundaries, break new ground and deliver the exceptional. If that excites you, then we want to hear from you.

**Dare to discover.**  
[careers.slb.com/recentgraduates](http://careers.slb.com/recentgraduates)

**Schlumberger**



Click on the ad to read more

### 30.4 Procedure for improper integrals; unbounded domain

**Problem 30.2** Calculate the integral

$$\int_A f(x, y) dS,$$

where the domain of integration  $A \subseteq \mathbb{R}^2$  is unbounded.

**Procedure:**

- 1) Examine whether  $f$  has a continuous extension to  $\overline{A} \setminus \{\infty\}$ . If this is not the case, remove suitable bounded neighbourhoods of the “sick” points, and treat each of these neighbourhoods as previously.

The residual set which we here denote by  $A$  is then treated in the following way.

- 2) The important *analysis of the sign of the integrand*. Divide  $A$  into the two sets

$$A^+ = \{(x, y) \in A \mid f(x, y) \geq 0\} \quad \text{and} \quad A^- = \{(x, y) \in A \mid f(x, y) \leq 0\}.$$

The sets  $A^+$  and  $A^-$  are found by analyzing the null curves, i.e. the set defined by the equation  $f(x, y) = 0$ .

- 3) Let us first consider  $A^+$ . If  $A^+ = \emptyset$ , then go to 4) in the following.

If  $A^+ \neq \emptyset$ , we choose one of the possible methods of trimming the set

$$\text{a) Rectangular: } A_n^+ = [-n, n]^2 \cap A^+, \quad \text{b) Polar: } A_R^+ = \overline{K}(\mathbf{0}; R) \cap A^+,$$

and we calculate

$$\text{a) Rectangular: } I_n^+ = \int_{A_n^+} f(x, y) dS, \quad \text{b) Polar: } I_R^+ = \int_{A_R^+} f(x, y) dS.$$

If  $I_n^+ \rightarrow +\infty$  for  $n \rightarrow +\infty$ , resp.  $I_R^+ \rightarrow +\infty$  for  $R \rightarrow +\infty$ , then the improper integral is *divergent*.

Otherwise,

$$\text{a) Rectangular: } \int_{A^+} f(x, y) dS = \lim_{n \rightarrow +\infty} \int_{A_n^+} f(x, y) dS.$$

$$\text{b) Polar: } \int_{A^+} f(x, y) dS = \lim_{R \rightarrow +\infty} \int_{A_R^+} f(x, y) dS.$$

Then go to 4).

- 4) Next turn to  $A^-$ . If  $A^- = \emptyset$ , then go to 5).

If  $A^- \neq \emptyset$ , choose one of the trimming possibilities

$$\text{a) Rectangular: } A_n^- = [-n, n]^2 \cap A^-, \quad \text{b) Polar: } A_R^- = \overline{K}(\mathbf{0}; R) \cap A^-,$$

and calculate

$$\text{a) Rectangular: } I_n^- = \int_{A_n^-} f(x, y) dS, \quad \text{b) Polar: } I_R^- = \int_{A_R^-} f(x, y) dS.$$

If  $I_n^- \rightarrow -\infty$  for  $n \rightarrow +\infty$ , resp.  $I_R^- \rightarrow -\infty$  for  $R \rightarrow +\infty$ , then the improper integral is *divergent*.

Otherwise,

a) Rectangular:  $\int_{A^-} f(x, y) dS = \lim_{n \rightarrow +\infty} \int_{A_n^-} f(x, y) dS.$

b) Polar:  $\int_{A^-} f(x, y) dS = \lim_{R \rightarrow +\infty} \int_{A_R^-} f(x, y) dS.$

5) If we obtain *convergence* in both 3) and 4), then the improper integral is *convergent* with the value

$$\int_A f(x, y) dS = \int_{A^+} f(x, y) dS + \int_{A^-} f(x, y) dS.$$

**Remark 30.2** If we put

$$A^0 = \{(x, y) \in A \mid f(x, y) = 0\},$$

it is seen that

$$\int_{A^0} f(x, y) dS = \int_{A^0} 0 dS = 0,$$

so it is of no importance whether we include  $A^0$  in our calculations, or not. The contribution from  $A^0$  is always 0.  $\diamond$

### 30.5 Examples of improper integrals

#### Example 30.1

A. Let  $B = [0, 1]^2$  be the unit square. Examine whether the improper plane integral

$$I = \int_B \frac{1}{y - x - 1} dS$$

is convergent. If this is the case, find its value.

- D. The domain  $B$  is closed and bounded; but the integrand is not defined in all points of  $B$ . Cut away open neighbourhoods of the points where the integrand is not defined; note that the integrand does not change sign; calculate the integral over any of the truncated domains and finally go to the limit.
- I. The denominator must never be zero, so we have to avoid the line  $y = x + 1$ . This line cuts  $B$  at the point  $(0, 1)$ , which must be removed from the domain of integration.

Then we realize that  $\frac{1}{y - x - 1} < 0$  everywhere in  $B \setminus \{(0, 1)\}$ , hence the integrand does not change sign in the part of  $B$ , where it is defined. Therefore we shall not further divide the domain according to the positive and the negative part of the integrand.

Our next step is to truncate  $B$ , such that the singular point  $(0, 1)$  does not lie in any of the closed and bounded domains  $B(\varepsilon)$ , and such that

$$\text{areal}(B \setminus B(\varepsilon)) \rightarrow 0 \quad \text{for } \varepsilon \rightarrow 0.$$




Figure 30.1: The domain  $B$  with the oblique singular line  $y = x + 1$  and the singular point  $(0, 1)$ , and a convenient truncation parallel to the  $Y$ -axis.

The most reasonable truncations among many possibilities are given by

$$B(\varepsilon) = [\varepsilon, 1] \times [0, 1] \quad \text{and} \quad \tilde{B}(\varepsilon) = [0, 1] \times [\varepsilon, 1],$$

for  $0 < \varepsilon < 1$ . We shall here choose the first one. Then by a reduction,

$$I(\varepsilon) = \int_{B(\varepsilon)} \frac{1}{y-x-1} dS = \int_{\varepsilon}^1 \left\{ \int_0^1 \frac{1}{y-x-1} dy \right\} dx.$$

We calculate the inner integral,

$$\int_0^1 \frac{1}{y-x-1} dy = [\ln |y-x-1|]_0^1 = \ln |-x| - \ln |-x-1| = \ln x - \ln(x+1).$$

Then insert the result followed by a partial integration,

$$\begin{aligned} I(\varepsilon) &= \int_{\varepsilon}^1 \{1 \cdot \ln x - 1 \cdot \ln(x+1)\} dx \\ &= [x \ln x]_{\varepsilon}^1 - \int_{\varepsilon}^1 x \cdot \frac{1}{x} dx - [(x+1) \ln(x+1)]_{\varepsilon}^1 + \int_{\varepsilon}^1 (x+1) \cdot \frac{1}{x+1} dx. \end{aligned}$$

The clever trick is here to choose  $\int dx = x$  in the first partial integration, and  $\int dx = x+1$  in the second one. In fact, the antiderivatives are only determined modulo a constant. By this trick we get

$$\begin{aligned} I(\varepsilon) &= 1 \cdot \ln 1 - \varepsilon \ln \varepsilon - 2 \ln 2 + (1+\varepsilon) \ln(1+\varepsilon) \\ &= -2 \ln 2 - \varepsilon \cdot \ln \varepsilon = (1+\varepsilon) \ln(1+\varepsilon). \end{aligned}$$

Since area  $B \setminus B(\varepsilon) = \varepsilon \rightarrow 0$  for  $\varepsilon \rightarrow 0+$ , this is the right limit. It follows from the magnitude of functions that

$$\varepsilon \ln \varepsilon \rightarrow 0 \quad \text{for } \varepsilon \rightarrow 0+,$$

and since

$$(1 + \varepsilon) \cdot \ln(1 + \varepsilon) \rightarrow 1 \cdot \ln 1 = 0 \quad \text{for } \varepsilon \rightarrow 0+,$$

we conclude that

$$\begin{aligned} I &= \lim_{\varepsilon \rightarrow 0+} I(\varepsilon) \\ &= -2 \ln 2 - \lim_{\varepsilon \rightarrow 0+} \varepsilon \cdot \ln \varepsilon + \lim_{\varepsilon \rightarrow 0+} (1 + \varepsilon) \cdot \ln(1 + \varepsilon) \\ &= -2 \ln 2, \end{aligned}$$

i.e. the improper integral is *convergent* and its value is

$$I = \int_B \frac{1}{y - x - 1} dS = -2 \ln 2 < 0.$$

- C. A *very weak control*. The integrand is negative, where it is defined. Hence the result should also be negative, which is seen to be true. (Note that this is only catching errors, where we end up with a *positive* result. Negative wrong results cannot be traced in this way). ◇

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
 We will appreciate and reward both your enthusiasm and talent.  
 Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



Click on the ad to read more

**Example 30.2**

A. Let  $\mathcal{F}$  be the half-sphere  $\mathcal{F}$  in the upper half plane of radius  $a > 0$ , i.e. in spherical coordinates,

$$r = a, \quad 0 \leq \theta \leq \frac{\pi}{2}, \quad 0 \leq \varphi \leq 2\pi.$$

Examine whether the improper surface integral

$$I = \int_{\mathcal{F}} \frac{1}{z} dS$$

is convergent or divergent.




Figure 30.2: The surface  $\mathcal{F}$  for  $a = 1$ .

D. Surface integrals can also be improper!

In the formulation of the task there is a hint of using *spherical* coordinates; though there exist also quite reasonable variants of solutions in *semi-polar* and *rectangular* coordinates.

The points where the integrand is not defined lie on the circle

$$\{(x, y, 0) \mid x^2 + y^2 = a^2\}$$

in the  $(x, y)$ -plane. Hence by the truncations we shall stay away from the  $(x, y)$ -plane. On the residual part of the surface we see that the integrand  $\frac{1}{z} > 0$ , so no further division of the domain is needed concerning the sign of the integrand.

I 1. *Spherical coordinates.* In this case we truncate  $\mathcal{F}$  to  $\mathcal{F}(\varepsilon)$  by

$$\mathcal{F}(\varepsilon) : \quad r = a, \quad 0 \leq \theta \leq \frac{\pi}{2} - \varepsilon, \quad 0 \leq \varphi \leq 2\pi, \quad \text{where } 0 < \varepsilon < \frac{\pi}{2}.$$

A geometrical consideration gives that

$$\text{area}(\mathcal{F} \setminus \mathcal{F}(\varepsilon)) \rightarrow 0 \quad \text{for } \varepsilon \rightarrow 0+,$$

and as mentioned in D. the integrand is positive, so we shall not divide the domain further.

By a reduction of the ordinary *surface integral* over  $\mathcal{F}(\varepsilon)$  we get

$$\begin{aligned} I(\varepsilon) &= \int_{\mathcal{F}(\varepsilon)} \frac{1}{z} dS = \int_0^{2\pi} \left\{ \int_0^{\frac{\pi}{2}-\varepsilon} \frac{1}{a \cos \theta} \cdot a^2 \sin \theta d\theta \right\} d\varphi \\ &= a \cdot 2\pi \int_0^{\frac{\pi}{2}-\varepsilon} \frac{1}{\cos \theta} \cdot \sin \theta d\theta = 2\pi a [-\ln \cos \theta]_0^{\frac{\pi}{2}-\varepsilon} \\ &= 2\pi a \left\{ \ln \cos 0 - \ln \cos \left( \frac{\pi}{2} - \varepsilon \right) \right\} \\ &= -2\pi a \ln \left\{ \cos \frac{\pi}{2} \cdot \cos \varepsilon + \sin \frac{\pi}{2} \cdot \sin \varepsilon \right\} \\ &= -2\pi a \ln \{0 + \sin \varepsilon\} = -2\pi a \ln \sin \varepsilon. \end{aligned}$$

Since  $\sin \varepsilon \rightarrow 0+$  for  $\varepsilon \rightarrow 0+$ , we have  $\ln \sin \varepsilon \rightarrow -\infty$  for  $\varepsilon \rightarrow 0+$ , i.e.  $I(\varepsilon) \rightarrow -(-\infty) = +\infty$  for  $\varepsilon \rightarrow 0+$ , and the improper surface integral is *divergent*.

**I 2.** *Semi-polar coordinates.* Here the surface is described by

$$\mathcal{F}: \quad 0 \leq z \leq a \quad \text{and} \quad \varrho = \sqrt{a^2 - z^2},$$

i.e.  $\varrho^2 + z^2 = a^2$ ,  $\varrho \geq 0$ . The surface is truncated in the following way,

$$\mathcal{F}(\varepsilon): \quad \varepsilon \leq z \leq a, \quad \varrho = \sqrt{a^2 - z^2}, \quad \text{where } 0 < \varepsilon < a,$$

and it follows geometrically that area  $(\mathcal{F} \setminus \mathcal{F}(\varepsilon)) \rightarrow 0$  for  $\varepsilon \rightarrow 0+$ .

The surface integral over  $\mathcal{F}(\varepsilon)$  is reduced in the following

$$I(\varepsilon) = \int_{\mathcal{F}(\varepsilon)} \frac{1}{z} dS = \int_{\varepsilon}^a \frac{1}{z} \left\{ \int_{B(z)} ds \right\} dz = \int_{\varepsilon}^a \frac{1}{z} \text{length}\{B(z)\} dz,$$

because  $\mathcal{F}(\varepsilon)$  is cut at height  $z \in [\varepsilon, a]$  in a circle  $B(z)$  of radius  $\varrho = \sqrt{a^2 - z^2}$ , such that the inner integral is the length of  $B(z)$ , i.e.

$$2\pi \varrho = 2\pi \sqrt{a^2 - z^2}.$$

Then we get by insertion

$$I(\varepsilon) = 2\pi \int_{\varepsilon}^a \frac{1}{z} \sqrt{a^2 - z^2} dz.$$

When  $z$  is small, then  $\sqrt{a^2 - z^2} \approx a$ , hence the integrand is  $\approx \frac{a}{z}$ . This function cannot be integrated from 0, so this gives a hint that we may have divergence. Let us prove this.

The integrand is positive everywhere. If  $0 < z \leq \frac{\sqrt{3}}{2} a$ , then

$$\sqrt{a^2 - z^2} \geq \sqrt{a^2 - \frac{3}{4} a^2} = \frac{a}{2}.$$

We then have for  $0 < \varepsilon < \frac{\sqrt{3}}{2}a$  the following estimates

$$\begin{aligned} I(\varepsilon) &= 2\pi \int_{\varepsilon}^a \frac{1}{z} \sqrt{a^2 - z^2} dz \geq 2\pi \int_{\varepsilon}^{\frac{\sqrt{3}}{2}a} \frac{1}{z} \sqrt{a^2 - z^2} dz \\ &\geq 2\pi \int_{\varepsilon}^{\frac{\sqrt{3}}{2}a} \frac{a}{2} \cdot \frac{1}{z} dz = \pi a \int_{\varepsilon}^{\frac{\sqrt{3}}{2}a} \frac{dz}{z} \\ &= \pi a [\ln z]_{\varepsilon}^{\frac{\sqrt{3}}{2}a} = \pi a \left\{ \ln \left( \frac{\sqrt{3}}{2}a \right) - \ln \varepsilon \right\} \rightarrow +\infty \quad \text{for } \varepsilon \rightarrow \infty, \end{aligned}$$

and we conclude that the improper surface integral is *divergent*.

**I 3.** *Rectangular coordinates.* In this case we consider  $\mathcal{F}$  as the graph of the function

$$z = f(x, y) = \sqrt{a^2 - x^2 - y^2}, \quad (x, y) \in E,$$

where the parametric domain is

$$E = \{(x, y) \mid x^2 + y^2 \leq a^2\}.$$

The natural truncation of the domain is here

$$E_\varepsilon = \{(x, y) \mid x^2 + y^2 \leq (a - \varepsilon)^2\}, \quad 0 < \varepsilon < a,$$

and it is obvious that  $\text{area}(E \setminus E_\varepsilon) \rightarrow 0$  for  $\varepsilon \rightarrow 0+$ .

I joined MITAS because  
I wanted **real responsibility**



The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)

**Month 16**  
I was a construction supervisor in the North Sea advising and helping foremen solve problems

Real work  
International opportunities  
Three work placements





 Click on the ad to read more

The *weight function* is in the case of a graph given by

$$\begin{aligned}\sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} &= \sqrt{1 + \left(\frac{x}{\sqrt{a^2 - x^2 - y^2}}\right)^2 + \left(\frac{y}{\sqrt{a^2 - x^2 - y^2}}\right)^2} \\ &= \frac{a}{\sqrt{a^2 - x^2 - y^2}}.\end{aligned}$$

Then we get by the reduction formula,

$$\begin{aligned}I(\varepsilon) &= \int_{\mathcal{F}(\varepsilon)} \frac{1}{z} dS = \int_{E_\varepsilon} \frac{1}{\sqrt{a^2 - x^2 - y^2}} \cdot \frac{a}{\sqrt{a^2 - x^2 - y^2}} dx dy \\ &= a \int_{E_\varepsilon} \frac{1}{a^2 - x^2 - y^2} dx dy = a \int_0^{2\pi} \left\{ \int_0^{a-\varepsilon} \frac{1}{a^2 - \varrho^2} \cdot \varrho d\varrho \right\} d\varphi \\ &= a \cdot 2\pi \int_0^{(a-\varepsilon)^2} \frac{1}{a^2 - t} \cdot \frac{1}{2} dt = \pi a \left[ -\ln |(a^2 - t)| \right]_0^{(a-\varepsilon)^2} \\ &= \pi a \{ \ln a^2 - \ln \{a^2 - (a - \varepsilon)^2\} \} \\ &= \pi a \{ 2 \ln a - \ln \{(2a - \varepsilon)\varepsilon\} \} \\ &= 2\pi a \ln a - \pi a \ln(2a - \varepsilon) - \pi a \ln \varepsilon \\ &\rightarrow 2\pi a \ln a - \pi a \ln 2a - (-\infty) = +\infty \quad \text{for } \varepsilon \rightarrow 0+, \end{aligned}$$

and we conclude again that the improper surface integral is *divergent*.

**Remark 30.3** In this case we could use all the three classical coordinate systems. Note that the three proofs are totally different in their arguments. ◇

### Example 30.3

A. Let  $B$  be the disc given by  $x^2 + y^2 \leq a^2$ . Find all values of  $\alpha \in \mathbb{R}$ , for which the (proper or improper) plane integral

$$J(\alpha) = \int_B (a^2 - y^2 - x^2)^\alpha dS$$

is convergent. In case of convergence, find its value.

This example is of the same type as

$$\int_0^a t^\alpha dt = \begin{cases} \frac{1}{\alpha+1} a^{\alpha+1} & \text{for } \alpha > -1, \\ \text{divergent} & \text{for } \alpha \leq -1, \end{cases}$$

and

$$\int_a^{+\infty} t^\alpha dt = \begin{cases} \text{divergent} & \text{for } \alpha \geq -1, \\ \frac{1}{|\alpha+1|} \cdot \frac{1}{a^{|\alpha+1|}} & \text{for } \alpha < -1, \end{cases}$$

known from the *Theory of Functions in One Variable*.




Figure 30.3: The disc  $B$  and the truncation  $B(p)$  for  $a = 1$ .

**D.** *Dimensional considerations* are here extremely useful. In fact,  $x, y \sim a$  and  $\int_B \cdots dS \sim a^2$ , hence

$$J(\alpha) \sim a^{2\alpha} \cdot a^2 = a^{2(\alpha+1)}.$$

In order to get convergence we must have

$$a^{2(\alpha+1)} \rightarrow 0 \quad \text{for } a \rightarrow 0+,$$

i.e.  $\alpha + 1 > 1$ , or  $\alpha > -1$ . Since

$$a^{2(\alpha+1)} \rightarrow +\infty \quad \text{for } a \rightarrow 0+, \text{ when } \alpha < -1,$$

we may expect divergence in this case. (A rough argument is the following: The integral of a positive term tends to  $+\infty$ , when the domain is shrunk. This is only possible, when we *start* with the value  $+\infty$ , i.e. with divergence).

The integral is proper, when  $\alpha \geq 0$ , and it is improper when  $\alpha < 0$ . When the integrand is defined, it is positive, so we shall not bother with an extra division of the domain according to the positive and negative part of the integrand.

If  $\alpha < 0$ , then the integrand is not defined on the boundary (the circle)  $x^2 + y^2 = a^2$ .

The considerations above indicate that the value  $\alpha = -1$  divides convergence and divergence. For that reason we start by first examining this case.

**I.** When we truncate, it is of no importance whether the integral is proper or improper. Since the integrand is positive (of fixed sign) we shall not make any further division of the domain. We truncate by the definition

$$B(p) = \{(x, y) \mid x^2 + y^2 \leq (pa)^2\}, \quad 0 < p < 1,$$

and we note that  $\text{area}\{B \setminus B(p)\} \rightarrow 0$  for  $p \rightarrow 1-$ .

1) If  $\alpha = -1$ , we get by using *polar* coordinates in  $B(p)$  that

$$\begin{aligned} J(-1; p) &= \int_{B(p)} \frac{1}{a^2 - x^2 - y^2} dS = \int_0^{2\pi} \left\{ \int_0^{pa} \frac{1}{a^2 - \varrho^2} \cdot \varrho d\varrho \right\} d\varphi \\ &= 2\pi \int_0^{(pa)^2} \frac{1}{a^2 - t^2} \cdot \frac{1}{2} dt = \pi \left[ -\ln |a^2 - t^2| \right]_0^{(pa)^2} \\ &= \pi \{ \ln a^2 - \ln (a^2 - p^2 a^2) \} = \pi \ln \left\{ \frac{a^2}{a^2(1 - p^2)} \right\} \\ &= \pi \ln \left( \frac{1}{1 - p^2} \right) \rightarrow +\infty \quad \text{for } p \rightarrow -1. \end{aligned}$$

We therefore conclude that

$$J(-1) = \int_B \frac{dS}{a^2 - x^2 - y^2} = \lim_{p \rightarrow 1^-} J(-1; p) = +\infty,$$

is divergent.

2) If  $\alpha < -1$ , i.e.  $\alpha + 1 < 0$ , then we can use 1) in the following rearrangements and estimates,

$$\begin{aligned} (a^2 - x^2 - y^2)^\alpha &= (a^2 - x^2 - y^2)^{\alpha+1} \cdot \frac{1}{a^2 - x^2 - y^2} \\ &= a^{2(\alpha+1)} \cdot \left\{ 1 - \frac{x^2 + y^2}{a^2} \right\}^{\alpha+1} \cdot \frac{1}{a^2 - x^2 - y^2} \\ &= a^{2(\alpha+1)} \cdot \frac{1}{\left\{ 1 - \frac{x^2 + y^2}{a^2} \right\}^{|\alpha+1|}} \cdot \frac{1}{a^2 - x^2 - y^2} \\ &\geq a^{2(\alpha+1)} \cdot \frac{1}{a^2 - x^2 - y^2}, \end{aligned}$$

because

$$\left\{ 1 - \frac{x^2 + y^2}{a^2} \right\}^{|\alpha+1|} \leq 1.$$

Then it follows from 1) that

$$\begin{aligned} J(\alpha; p) &= \int_{B(p)} (a^2 - x^2 - y^2)^\alpha dS \geq a^{2(\alpha+1)} \int_{B(p)} \frac{1}{a^2 - x^2 - y^2} dS \\ &= a^{2(\alpha+1)} J(-1; p) \rightarrow +\infty \quad \text{for } p \rightarrow 1^-, \end{aligned}$$

i.e.  $J(\alpha; p) \rightarrow +\infty$  for  $p \rightarrow 1^-$ , and we have got *divergence* for  $\alpha < -1$ , and hence also for  $\alpha \leq -1$ .

3) If  $\alpha > -1$ , i.e.  $\alpha + 1 > 0$ , we get by using *polar* coordinates in  $B(p)$  that

$$\begin{aligned} J(\alpha; p) &= \int_{B(p)} (a^2 - x^2 - y^2)^\alpha dS = \int_0^{2\pi} \left\{ \int_{black0}^{pa} (a^2 - \varrho^2)^\alpha \cdot \varrho d\varrho \right\} d\varphi \\ &= 2\pi \int_{a^2}^{a^2(1-p^2)} t^\alpha \cdot \left( -\frac{1}{2} \right) dt = \pi \left[ -\frac{1}{\alpha+1} t^{\alpha+1} \right]_{a^2}^{a^2(1-p^2)} \\ &= \frac{\pi}{\alpha} \left\{ a^{2(\alpha+1)} - a^{2(\alpha+1)} \cdot (1 - p^2)^{\alpha+1} \right\} \\ &= \frac{\pi}{\alpha+1} a^{2(\alpha+1)} \left\{ 1 - (1 - p^2)^{\alpha+1} \right\}. \end{aligned}$$

Since  $\alpha + 1 > 0$  and  $1 - p^2 \rightarrow 0+$  for  $p \rightarrow 1-$ , it follows that

$$(1 - p^2)^{\alpha+1} \rightarrow 0 \quad \text{for } p \rightarrow 1-.$$

Hence the integral is *convergent* for  $\alpha > -1$  with the value

$$J(\alpha) = \int_B (a^2 - x^2 - y^2)^\alpha dS = \frac{\pi}{\alpha + 1} a^{2(\alpha+1)}.$$

Summing up we have proved that

$$J(\alpha) = \int_B (a^2 - x^2 - y^2)^\alpha dS = \begin{cases} \frac{\pi}{\alpha + 1} a^{2(\alpha+1)} & \text{for } \alpha > -1; \text{ convergence;} \\ \infty & \text{for } \alpha \leq -1; \text{ divergence.} \end{cases}$$

When the integrand has fixed sign, we allow ourselves to put the value equal to  $+\infty$  (positive integrand) or  $-\infty$  (negative integrand).

Note that we shall *not* allow this notation, if both the positive part and the negative part are infinite, because  $\infty - \infty$  does not make sense. We shall return to this in Example 30.5.  $\diamond$

The advertisement features a man walking down a city street at night, looking towards the right. In the background, there are tall buildings and blurred lights. On the left, the IE business school logo is visible. On the right, a speech bubble contains the hashtag #gobeyond. Above the man, a graphic shows the ranking "#1 European Business School" from Financial Times 2013. Below the man, a banner reads "MASTER IN MANAGEMENT". A descriptive text block states: "Because achieving your dreams is your greatest challenge. IE Business School's Master in Management taught in English, Spanish or bilingually, trains young high performance professionals at the beginning of their career through an innovative and stimulating program that will help them reach their full potential." A bulleted list highlights: "Choose your area of specialization.", "Customize your master through the different options offered.", and "Global Immersion Weeks in locations such as London, Silicon Valley or Shanghai.". At the bottom, a tagline says "Because you change, we change with you." and provides contact information: www.ie.edu/master-management, mim.admissions@ie.edu, and social media links for Facebook, Twitter, LinkedIn, YouTube, and Instagram.



Click on the ad to read more

**Example 30.4**

A. In advanced technical literature one often sees the improper 1-dimensional integral

$$\int_0^\infty e^{-x^2} dx.$$

Obvious applications can be found in Probability and Statistics (the normal distribution); but one can also find it in many other places (the heat equation, diffusion). We shall find the value of this important integral.

D. The integral cannot be calculated by methods from the elementary calculus. It is fairly easy to prove that it is convergent. In fact, if we introduce the function

$$f(t) = (1+t)e^{-t} \quad \text{with } f'(t) = -te^{-t} < 0 \quad \text{for } t > 0,$$

then it follows, because  $f(t)$  is decreasing for  $t > 0$ , that we have

$$(1+t)e^{-t} \leq f(0) = 1, \quad \text{i.e. } e^{-t} \leq \frac{1}{1+t} \text{ for } t \geq 0.$$

If we put  $t = x^2$ , we get

$$e^{-x^2} \leq \frac{1}{1+x^2},$$

hence

$$0 < \int_0^n e^{-x^2} dx \leq \int_0^n \frac{dx}{1+x^2} = \text{Arctan } n \rightarrow \frac{\pi}{2} \quad \text{for } n \rightarrow +\infty.$$

This proves the convergence, and also the estimate

$$(0 \leq) \quad \int_0^\infty e^{-x^2} dx \leq \frac{\pi}{2}.$$

However, we still have not found the exact value.

We shall show that it is possible to find the value by using methods from the *Theory of Functions in Several Variables*. The trick is instead to consider the improper plane integral

$$I = \iint_B \exp(-x^2 - y^2) dS, \quad \text{where } B = [0, +\infty[^2 \text{ is the first quadrant.}$$

The integrand is defined and positive everywhere, so we shall not make any further division of the domain according to the sign of the integrand.

The domain of integration is unbounded, so we must truncate it in a bounded way. We have two obvious possibilities of doing this, depending on whether we consider *polar* or *rectangular* coordinates. The idea is to use both of them, because we by using the *polar* coordinates obtain the *value* of the integral  $I$ , and by using the *rectangular* coordinates we obtain the connection to the integral under consideration.




Figure 30.4: The domain  $Q(R)$  for  $R = 1$ .

- 1) *Polar coordinates.* We truncate by taking the intersection of  $B$  and discs of radius  $R$ ,

$$Q(R) = \{(x, y) \mid x \geq 0, y \geq 0, x^2 + y^2 \leq R^2\}.$$

We reach every point in the first quadrant  $B$  by taking the limit  $R \rightarrow +\infty$ .

When we apply the reduction theorem in polar coordinates over  $Q(R)$  we get

$$\begin{aligned} I(R) &= \int_{Q(R)} \exp(-x^2 - y^2) \, dS = \int_0^{\frac{\pi}{2}} \left\{ \int_0^R \exp(-\varrho^2) \varrho \, d\varrho \right\} d\varphi \\ &= \frac{\pi}{2} \int_0^{R^2} e^{-t} \cdot \frac{1}{2} dt = \frac{\pi}{4} [e^{-t}]_0^{R^2} = \frac{\pi}{4} \{1 - \exp(-R^2)\}. \end{aligned}$$

Since  $\exp(-R^2) \rightarrow 0$  for  $R \rightarrow +\infty$ , we conclude that the improper integral is *convergent* with the value

$$I = \int_B \exp(-x^2 - y^2) \, dS = \lim_{R \rightarrow \infty} I(R) = \frac{\pi}{4}.$$

- 2) *Rectangular coordinates.* The truncation is here

$$R(n) = \{(x, y) \mid 0 \leq x \leq n, 0 \leq y \leq n\} = [0, n]^2.$$

We get every point in the first quadrant  $B$  by taking the limit  $n \rightarrow +\infty$ .




Figure 30.5: The domain  $R(n)$  for  $n = 1$ .

When we apply the rectangular reduction theorem over  $R(n)$  we get

$$\begin{aligned} J(n) &= \int_{R(n)} \exp(-x^2 - y^2) \, dS = \int_0^n \left\{ \int_0^n e^{-x^2} \cdot e^{-y^2} \, dx \right\} dy \\ &= \int_0^n e^{-x^2} \, dx \cdot \int_0^n e^{-y^2} \, dy = \left\{ \int_0^n e^{-t^2} \, dt \right\}^2. \end{aligned}$$

- 3) *Summary.* According to 1) the improper integral is convergent, hence the limit can be taken in 2). Since the limit is the same, no matter which truncation we are using, we must have

$$I = \frac{\pi}{4} = \int_B \exp(-x^2 - y^2) \, dS = \lim_{n \rightarrow \infty} J(n) = \left\{ \int_0^\infty e^{-t^2} \, dt \right\}^2.$$

Since  $\int_0^\infty e^{-t^2} \, dt > 0$ , we finally get the value of the integral

$$\int_0^\infty e^{-t^2} \, dt = \frac{\sqrt{\pi}}{2}.$$

**Remark 30.4** It follows that

$$I = \frac{\sqrt{\pi}}{2} \leq \frac{\pi}{2},$$

cf. a remark in section D.  $\diamond$

**Example 30.5**

- A. Examine whether the improper plane integral

$$\int_{\mathbb{R}^2} \frac{xy}{(1+x^2)(1+y^2)} dS$$

is convergent or divergent.

- D. An analysis of the sign shows that the integrand is positive in the first and the third quadrant, while it is negative in the second and the fourth quadrant, so we shall also divide the domain according to the sign. Here we shall also demonstrate the *wrong* argument where one forgets to divide according to the sign of the integrand. This shows that one has to be careful here.
- I. We see that the integrand changes its sign if we reflect it in either the  $x$ -axis or in the  $y$ -axis. This shows that the integral over *any* bounded set  $B$ , which is symmetric with respect to at least one of the axes, is 0,

$$\int_B \frac{xy}{(1+x^2)(1+y^2)} dS = 0.$$

The usual truncations in discs or centred squares satisfy this symmetry with respect to both axes, so if one is not too careful one will *erroneously* conclude in the limit that the plane integral is “convergent with the value  $\lim_{n \rightarrow \infty} 0 = 0$ ”.

**"I studied English for 16 years but...  
...I finally learned to speak it in just six lessons"**

Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download




Figure 30.6: The truncation  $B(n)$  for  $n = 1$ .

We shall prove that this conclusion is *not* correct. We choose this time the truncation

$$B(n) = [0, n] \times [0, n],$$

i.e. we only consider a subset of the set, where the integrand is  $\geq 0$ . If something goes wrong here in the first quadrant, then it also is wrong in any bigger subset of  $[0, n]^2 \cup [-n, 0]^2$ .

By reduction over  $B(n)$  we get

$$\begin{aligned} \int_{B(n)} \frac{xy}{(1+x^2)(1+y^2)} dS &= \int_0^n \left\{ \int_0^n \frac{xy}{(1+x^2)(1+y^2)} dx \right\} dy \\ &= \int_0^n \frac{x}{1+x^2} dx \cdot \int_0^n \frac{y}{1+y^2} dy = \left\{ \int_0^n \frac{t}{1+t^2} dt \right\}^2 \\ &= \left\{ \left[ \frac{1}{2} \ln(1+t^2) \right]_0^n \right\}^2 = \frac{1}{4} \{ \ln(1+n^2) \}^2 \\ &\rightarrow +\infty \quad \text{for } n \rightarrow \infty. \end{aligned}$$

This shows that the improper integral is *divergent*, thus our first argument must be wrong! (Of the type  $\infty - \infty$ ).

**Remark 30.5** We shall demonstrate how wrong this illegal method is. If we choose the “skew” truncation

$$Q(a, n) = [-an, n] \times [-an, n], \quad a > 0 \text{ constant},$$

we still get  $\mathbb{R}^2$  by taking the limit  $n \rightarrow +\infty$ . We get by a rectangular reduction,

$$\begin{aligned} \int_{Q(a,n)} \frac{xy}{(1+x^2)(1+y^2)} dS &= \left\{ \int_{-an}^n \frac{t}{1+t^2} dt \right\}^2 = \left\{ \left[ \frac{1}{2} \ln(1+t^2) \right]_{-an}^n \right\}^2 \\ &= \left\{ \frac{1}{2} \ln \left( \frac{1+n^2}{1+a^2 n^2} \right) \right\}^2 = \left\{ \frac{1}{2} \ln \left( \frac{1+\frac{1}{n^2}}{a^2 + \frac{1}{n^2}} \right) \right\}^2 \\ &\rightarrow \{\ln a\}^2 \quad \text{for } n \rightarrow \infty. \end{aligned}$$

Every calculation is correct. The only thing which is *wrong* is that the assumptions of taking the limit (with respect to the conclusion of convergence) are *not* satisfied. We note that  $\{\ln a\}^2$  go through the whole interval  $[0, +\infty[$ , when  $a$  go through  $\mathbb{R}_+$ , which means that we can obtain any  $q \geq 0$  as a candidate for a limit of the improper plane integral, which is nonsense.

If we instead use the truncations

$$R(a, n) = [-n, an] \times [-an, n], \quad a > 0 \text{ constant},$$

we obtain analogously all negative numbers as possible limits. But *if* the limit exists, then it is unique! Hence the improper plane integral is *divergent*.  $\diamond$

Excellent Economics and Business programmes at:

 university of  
groningen





**“The perfect start  
of a successful,  
international career.”**

[www.rug.nl/feb/education](http://www.rug.nl/feb/education)

**CLICK HERE**  
to discover why both socially  
and academically the University  
of Groningen is one of the best  
places for a student to be



Click on the ad to read more

**Example 30.6** Given the meridian curve  $\mathcal{M}$  of the parametric description

$$\varrho = a \cos t, \quad z = a\{\ln(1 + \sin t) - \ln \cos t - \sin t\}, \quad t \in \left[0, \frac{\pi}{2}\right[$$

When this is rotated we obtain a surface of revolution  $\mathcal{O}$  (half of the pseudo-sphere), which stretches into infinity along the positive part of the  $Z$ -axis.

Find the integral which gives the area of that part of  $\mathcal{O}$ , which corresponds to  $[0, T]$ , where  $T < \frac{\pi}{2}$ .

Then find the area of the pseudo-sphere by letting  $T \rightarrow \frac{\pi}{2}$ .

**A** Surface area of an infinite surface of revolution; improper surface integral.

**D** First find the curve element  $ds$  on  $\mathcal{M}$ . Then compute the surface area of  $\mathcal{O}_T$ , i.e. the surface corresponding to  $t \in [0, T]$ , where  $T < \frac{\pi}{2}$ . This means that we shall calculate

$$2\pi \int_{\mathcal{O}_T} \varrho(t) ds.$$

Finally, take the limit  $T \rightarrow \frac{\pi}{2}$ .

**I** First calculate

$$\begin{aligned} \mathbf{r}'(t) &= a \left( -\sin t, \frac{\cos t}{1 + \sin t} + \frac{\sin t}{\cos t} - \cos t \right) = a \left( -\sin t, \frac{\cos t \cdot (1 - \sin t)}{1 - \sin^2 t} + \frac{\sin t}{\cos t} - \cos t \right) \\ &= a \left( -\sin t, \frac{1 - \sin t + \sin t - \cos^2 t}{\cos t} \right) = a \sin t \cdot (-1, \tan t). \end{aligned}$$

Hence

$$ds = \|\mathbf{r}'(t)\| dt = a |\sin t| \sqrt{1 + \tan^2 t} dt = a \left| \frac{\sin t}{\cos t} \right| dt = a \tan t dt,$$




Figure 30.7: The meridian curve of the pseudo-sphere.

and accordingly,

$$\begin{aligned}\text{area}(\mathcal{O}_T) &= 2\pi \int_{\mathcal{O}_T} \varrho(t) ds = 2\pi \int_0^T a \cos t \cdot a \tan t dt \\ &= 2\pi a^2 \int_0^T \sin t dt = 2\pi a^2 (1 - \cos T).\end{aligned}$$

Finally, by taking the limit we find the improper surface area

$$\text{area}(\mathcal{O}) = \lim_{T \rightarrow \frac{\pi}{2}^-} \text{area}(\mathcal{O}_T) = 2\pi a^2.$$

REMARK. Note that the “half” pseudo-sphere” has the same surface area as the usual upper half sphere of radius  $a$ . ◇

**Example 30.7** Check in each of the following cases if the given surface integral is convergent or divergent; in case of convergency, find the value.

- 1) The surface integral  $\int_{\mathcal{F}} \frac{1}{(a+4z)^2} dS$  over the surface  $\mathcal{F}$  given by  $az = x^2 + y^2$ ,  $(x, y) \in \mathbb{R}^2$ .
- 2) The surface integral  $\int_{\mathcal{F}} \frac{x^2}{z^2 + a^2} dS$  over the surface  $\mathcal{F}$  given by  $x^2 + y^2 = a^2$ ,  $z \in \mathbb{R}$ .
- 3) The surface integral  $\int_{\mathcal{F}} y^2 \exp\left(-\frac{|z|}{a}\right) dS$  over the surface  $\mathcal{F}$  given by  $x^2 + y^2 = a^2$ ,  $z \in \mathbb{R}$ .
- 4) The surface integral  $\int_{\mathcal{F}} \frac{1}{z(x+y)} dS$  over the surface  $\mathcal{F}$  given by  $z = \sqrt{2xy}$ ,  $(x, y) \in [a, +\infty[^2$ .

**A** Improper surface integral.

**D** First analyze why the integral is improper. Then truncate the surface and split it into the positive and the negative part of the integrand. Finally take the limit.

**I** 1) The surface is a paraboloid of revolution.

$$z = \frac{1}{a} (x^2 + y^2) = \frac{1}{a} \varrho \geq 0.$$

The integrand is  $\geq \frac{1}{a^2} > 0$  everywhere on the surface.

The surface is described as the graph of the equation  $z = \frac{1}{a} (x^2 + y^2)$ , so the weight function becomes

$$\sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} = \sqrt{1 + \left(\frac{2x}{a}\right)^2 + \left(\frac{2y}{a}\right)^2} = \sqrt{1 + \frac{4}{a^2} \varrho^2}.$$

We choose the truncated domain in polar coordinates as  $0 \leq \varrho \leq R$ . It follows from the above that the area element is

$$dS = \sqrt{1 + \frac{4}{a^2} \varrho^2} \varrho d\varrho d\varphi,$$

hence the surface integral over the truncated surface  $\mathcal{F}_R$  is

$$\begin{aligned}\int_{\mathcal{F}_R} \frac{1}{(a+4z)^2} dS &= \int_0^{2\pi} \left\{ \int_0^R \frac{1}{\left(a + \frac{4\varrho^2}{a}\right)^2} \sqrt{1 + \frac{4}{a^2}} \varrho d\varrho \right\} d\varphi \\ &= \frac{2\pi}{a^2} \int_0^R \left\{ 1 + \frac{4}{a^2} \varrho^2 \right\}^{-\frac{3}{2}} \cdot \frac{1}{2} \cdot \frac{a^2}{4} \cdot \frac{4}{a^2} \cdot 2\varrho d\varrho \\ &= \frac{2\pi}{a^2} \cdot \frac{a^2}{8} \left[ -\frac{2}{\sqrt{1 + \frac{4}{a^2} \varrho^2}} \right]_0^R = \frac{\pi}{2} \left( 1 - \frac{1}{\sqrt{1 + \frac{4}{a^2} R^2}} \right).\end{aligned}$$

This expression clearly converges for  $R \rightarrow +\infty$ , hence the improper surface integral is convergent of the value

$$\int_{\mathcal{F}} \frac{1}{(a+4z)^2} dS = \lim_{R \rightarrow +\infty} \frac{\pi}{2} \left( 1 - \frac{1}{\sqrt{1 + \frac{4}{a^2} R^2}} \right) = \frac{\pi}{2}.$$

- 2) The surface is an infinite cylinder surface with the circle in the  $XY$ -plane of centrum  $(0, 0)$  and radius  $a$  as its leading curve. When we use semi-polar coordinates we get

$$x = a \cos \varphi, \quad y = a \sin \varphi, \quad z = z, \quad \varphi \in [0, 2\pi], \quad z \in \mathbb{R},$$

and

$$dS = a d\varphi dz.$$

The integrand is positive, so we choose the truncation  $|z| \leq A$ . Then

$$\begin{aligned}\int_{\mathcal{F}_A} \frac{x^2}{z^2 + a^2} dS &= \int_0^{2\pi} \left\{ \int_{black-A}^A \frac{a^2 \cos^2 \varphi}{z^2 + a^2} \cdot a dz \right\} d\varphi \\ &= \int_0^{2\pi} \cos^2 \varphi d\varphi \cdot a^2 \int_{-A}^A \frac{1}{1 + \left(\frac{z}{a}\right)^2} \cdot \frac{1}{a} dz = a^2 \pi \cdot 2 \operatorname{Arctan} \left( \frac{A}{a} \right).\end{aligned}$$

This expression converges for  $A \rightarrow +\infty$ , and we conclude that the improper surface integral is convergent with the value

$$\int_{\mathcal{F}} \frac{x^2}{z^2 + a^2} dS = \lim_{A \rightarrow +\infty} a^2 \pi \cdot 2 \operatorname{Arctan} \left( \frac{A}{a} \right) = a^2 \pi^2.$$

- 3) By using semi-polar coordinates it is seen that

$$x = a \cos \varphi, \quad y = a \sin \varphi, \quad z = z, \quad \varphi \in [0, 2\pi], \quad z \in \mathbb{R}.$$

The surface element is

$$dS = a d\varphi dz.$$

The integrand is positive everywhere, so we choose the truncation  $|z| \leq A$ . Then

$$\begin{aligned}\int_{\mathcal{F}_A} y^2 \exp\left(-\frac{|z|}{a}\right) dS &= \int_0^{2\pi} \left\{ \int_{-A}^A a^2 \sin^2 \varphi \cdot \exp\left(-\frac{|z|}{a}\right) a dz \right\} d\varphi \\ &= a^3 \cdot \pi \cdot 2 \int_0^A \exp\left(-\frac{z}{a}\right) dz = 2a^4 \pi \cdot \left[ -\exp\left(-\frac{z}{a}\right) \right]_0^A \\ &= 2\pi a^4 \left\{ 1 - \exp\left(-\frac{A}{a}\right) \right\}.\end{aligned}$$

This expression is clearly convergent for  $A \rightarrow +\infty$ , hence the improper surface integral is convergent with the value

$$\int_{\mathcal{F}} y^2 \exp\left(-\frac{|z|}{a}\right) dS = \lim_{A \rightarrow +\infty} \int_{\mathcal{F}_A} y^2 \exp\left(-\frac{|z|}{a}\right) dS = 2\pi a^4.$$

- 4) When the surface is the graph of  $z = \sqrt{2xy}$  for  $x, y \geq a$ , then the surface element is

$$\begin{aligned}dS &= \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy = \sqrt{1 + \frac{y}{2x} + \frac{x}{2y}} dx dy \\ &= \sqrt{\frac{2xy + y^2 + x^2}{2xy}} dx dy = \frac{x+y}{\sqrt{2xy}} dx dy.\end{aligned}$$

## American online LIGS University

is currently enrolling in the  
Interactive Online BBA, MBA, MSc,  
DBA and PhD programs:

- enroll by **September 30th, 2014** and
- save up to **16%** on the tuition!
- pay in 10 installments / 2 years
- Interactive Online education
- visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to find out more!

Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education.  
[More info here.](#)



Click on the ad to read more

The integrand is given on the surface  $\mathcal{F}$  by

$$\frac{1}{z(x+y)} = \frac{1}{\sqrt{2xy}(x+y)}$$

which is clearly positive, because  $x, y \geq a$ .

For every  $A > a$  we define the truncation by  $a \leq x, y \leq A$ . Then the surface integral over the corresponding truncated surface  $\mathcal{F}_A$  is

$$\begin{aligned} \int_{\mathcal{F}_A} \frac{1}{z(x+y)} dS &= \int_a^A \left\{ \int_a^A \frac{1}{\sqrt{2xy}(x+y)} \cdot \frac{x+y}{\sqrt{2xy}} dx \right\} dy \\ &= \frac{1}{2} \left\{ \int_a^A \frac{dx}{x} \right\} \cdot \left\{ \int_a^A \frac{dy}{y} \right\} = \frac{1}{2} \left\{ [\ln t]_a^A \right\}^2 = \frac{1}{2} \left\{ \ln \left( \frac{A}{a} \right) \right\}^2 \rightarrow +\infty \quad \text{for } A \rightarrow +\infty. \end{aligned}$$

We conclude that the improper surface integral is divergent.

- 5) The surface is the same as in **Example 30.7.4**, so the surface element is

$$dS = \frac{x+y}{\sqrt{2xy}} dx dy.$$

The integrand is on the surface  $\mathcal{F}$  given by

$$\frac{1}{z^2 xy} = \frac{1}{2xy \cdot xy} = \frac{1}{2} \cdot \frac{1}{x^2} \cdot \frac{1}{y^2}.$$

This is positive, so we shall again use the truncation  $a \leq x, y \leq A$ . Then

$$\begin{aligned} \int_{\mathcal{F}_A} \frac{1}{z^2 xy} dS &= \int_a^A \left\{ \int_a^A \frac{1}{2} \cdot \frac{1}{x^2} \cdot \frac{1}{y^2} \cdot \frac{x+y}{\sqrt{2xy}} dx dy \right\} dy \\ &= \frac{1}{2\sqrt{2}} \int_a^A x^{-\frac{3}{2}} dx \cdot \int_a^A y^{-\frac{5}{2}} dy + \frac{1}{2\sqrt{2}} \int_a^A x^{-\frac{5}{2}} dx \cdot \int_a^A y^{-\frac{3}{2}} dy \\ &= \frac{1}{\sqrt{2}} \left[ -\frac{2}{\sqrt{x}} \right]_a^A \cdot \left[ -\frac{2}{3} \cdot \frac{1}{y\sqrt{y}} \right]_a^A = \frac{2\sqrt{2}}{3} \left( \frac{1}{\sqrt{a}} - \frac{1}{\sqrt{A}} \right) \cdot \left( \frac{1}{a\sqrt{a}} - \frac{1}{A\sqrt{A}} \right) \\ &\rightarrow \frac{2\sqrt{2}}{3} \cdot \frac{1}{\sqrt{a}} \cdot \frac{1}{a\sqrt{a}} = \frac{2\sqrt{2}}{3a^2} \quad \text{for } A \rightarrow +\infty. \end{aligned}$$

The improper surface integral converges towards the value

$$\int_{\mathcal{F}} \frac{1}{z^2 xy} dS = \frac{2\sqrt{2}}{3a^2}.$$

**Example 30.8** Check in each of the following cases if the given surface integral is convergent or divergent; in case of convergency, find its value.

Let  $\mathcal{S}$  denote the sphere of centrum  $(0, 0, 0)$  and radius  $a$ , while  $\mathcal{F}$  is given by  $az = x^2 + y^2$ ,  $x^2 + y^2 \leq a^2$ .

$$1) \int_{\mathcal{S}} \frac{1}{a-z} dS,$$

$$2) \int_{\mathcal{S}} \sqrt{\frac{a}{|z|}} dS,$$

$$3) \int_{\mathcal{F}} \frac{1}{a-z} dS,$$

$$4) \int_{\mathcal{F}} \sqrt{\frac{a}{z}} dS.$$

**A** Improper surface integrals.

**D** Analyze why the integral is improper. Since the integrands are  $\geq 0$  in all cases, we shall only find some nice truncations of the surface.

**I** 1) Since  $|z| \leq a$  on  $\mathcal{S}$ , the integrand is  $\frac{1}{a-z} > 0$  on  $\mathcal{S} \setminus \{(0, 0, a)\}$ . The integrand tends towards  $+\infty$ , when  $(x, y, z) \rightarrow (0, 0, a)$  on  $\mathcal{S}$ .

When we use spherical coordinates on  $\mathcal{S}$ ,

$$x = a \cos \varphi \cdot \sin \theta, \quad y = a \sin \varphi \cdot \sin \theta, \quad z = a \cos \theta,$$

for

$$\varphi \in [0, 2\pi], \quad \theta \in [0, \pi],$$

it is well-known that

$$dS = a^2 \sin \theta d\varphi d\theta.$$

The singular point  $(0, 0, a)$  corresponds to  $\theta = 0$ , hence we choose the truncation  $\theta \in [\varepsilon, \pi]$ , where  $\varepsilon > 0$  corresponds to the subsurface  $\mathcal{S}_\varepsilon$ . When we integrate over  $\mathcal{S}_\varepsilon$  we get

$$\begin{aligned} \int_{\mathcal{S}_\varepsilon} \frac{1}{a-z} dS &= \int_0^{2\pi} \left\{ \int_\varepsilon^\pi \frac{1}{a-a \cos \theta} a^2 \sin \theta d\theta \right\} d\varphi \\ &= 2\pi a \int_\varepsilon^\pi \frac{\sin \theta}{1-\cos \theta} d\theta = 2\pi a [\ln(1-\cos \theta)]_\varepsilon^\pi \\ &= 2\pi a \{ \ln 2 - \ln(1-\cos \varepsilon) \} = 2\pi a \ln \frac{2}{2 \sin^2 \frac{\varepsilon}{2}} \\ &= 4a\pi \ln \frac{1}{\sin \frac{\varepsilon}{2}} \rightarrow +\infty \quad \text{for } \varepsilon \rightarrow 0+, \end{aligned}$$

and the improper surface integral is divergent.

2) In this case the integrand is  $> 0$  on  $\mathcal{S}_0$ , where  $\mathcal{S}_0$  is the set of points on  $\mathcal{S}$ , which is not contained in the  $XY$ -plane, where the integrand is not defined. We use again spherical coordinates. Due to the symmetry it suffices to consider the domain

$$\mathcal{S}_\varepsilon : \quad \varphi \in [0, 2\pi] \quad \text{and} \quad \theta \in \left[0, \frac{\pi}{2} - \varepsilon\right].$$

It follows by insertion of  $z = a \cos \theta$  and  $dS = a^2 \sin \theta d\theta d\varphi$ , that

$$\begin{aligned} \int_{S_\varepsilon} \sqrt{\frac{a}{|z|}} dS &= \int_0^{2\pi} \left\{ \int_0^{\frac{\pi}{2}-\varepsilon} \sqrt{\frac{a}{a \cos \theta}} \cdot a^2 \sin \theta d\theta \right\} d\varphi \\ &= 2\pi a^2 \int_0^{\frac{\pi}{2}-\varepsilon} \frac{\sin \theta}{\sqrt{\cos \theta}} d\theta = 2\pi a^2 \left[ -2\sqrt{\cos \theta} \right]_0^{\frac{\pi}{2}-\varepsilon} \\ &= 4\pi a^2 \{1 - \sqrt{\sin \varepsilon}\} \rightarrow 4\pi a^2 \quad \text{for } \varepsilon \rightarrow 0+. \end{aligned}$$

We conclude that the improper surface integral is convergent.

Of symmetric reasons the value is

$$\int_S \sqrt{\frac{a}{|z|}} dS = \lim_{\varepsilon \rightarrow 0+} 2 \int_{S_\varepsilon} \sqrt{\frac{a}{|z|}} dS = 2 \cdot 4\pi a^2 = 8\pi a^2.$$

- 3) The surface is the graph of  $z = \frac{1}{a}(x^2 + y^2) = \frac{\varrho^2}{a}$ , so the area element is

$$\begin{aligned} dS &= \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy = \sqrt{1 + \left(\frac{2x}{a}\right)^2 + \left(\frac{2y}{a}\right)^2} dx dy \\ &= \frac{1}{a} \sqrt{a^2 + 4(x^2 + y^2)} dx dy = \frac{1}{a} \sqrt{a^2 + 4\varrho^2} \cdot \varrho d\varrho d\varphi. \end{aligned}$$

The integrand is the same as in **Example 30.8.1**, and since  $z \leq a$  on  $\mathcal{F}$ , it is positive for  $z < a$ . We choose the truncation in polar coordinates by

$$\mathcal{F}_\varepsilon : \quad 0 \leq \varrho \leq a - \varepsilon, \quad \varphi \in [0, 2\pi].$$

Then by insertion,

$$\begin{aligned} \int_{\mathcal{F}_\varepsilon} \frac{1}{a-z} dS &= \int_0^{2\pi} \left\{ \int_0^{a-\varepsilon} \frac{1}{a - \frac{\varrho^2}{a}} \cdot \frac{1}{a} \sqrt{a^2 + 4\varrho^2} \varrho d\varrho \right\} d\varphi \\ &= 2\pi \int_0^{a-\varepsilon} \frac{\varrho}{a^2 - \varrho^2} \sqrt{a^2 + 4\varrho^2} d\varrho \geq a\pi \int_0^{a-\varepsilon} \frac{1}{a^2 - \varrho^2} \cdot 2\varrho d\varrho \\ &= a\pi [-\ln(a^2 - \varrho^2)]_0^{a-\varepsilon} \\ &= a\pi \{\ln a^2 - \ln(a^2 - (a - \varepsilon)^2)\} \rightarrow +\infty \end{aligned}$$

for  $\varepsilon \rightarrow 0+$ , and the improper surface integral is divergent.

- 4) The singular point is  $(0, 0, 0)$ . We choose the truncation

$$\mathcal{F}_\varepsilon : \quad \varphi \in [0, 2\pi], \quad \varrho \in [\varepsilon, a],$$

and

$$z = \frac{\varrho^2}{a} > 0, \quad dS = \frac{1}{a} \sqrt{a^2 + 4\varrho^2} \varrho d\varrho d\varphi.$$

Then by insertion

$$\begin{aligned}
 \int_{\mathcal{F}_\varepsilon} \sqrt{\frac{a}{z}} dS &= \int_0^{2\pi} \left\{ \int_\varepsilon^a \sqrt{\frac{a^2}{\varrho^2} \cdot \frac{1}{a} \sqrt{a^2 + 4\varrho^2} \varrho d\varrho} \right\} d\varphi \\
 &= 2\pi a \int_\varepsilon^a \sqrt{1 + \left(\frac{2\varrho}{a}\right)^2} d\varrho \quad \left[ \frac{2\varrho}{a} = \sinh t \right] \\
 &= 2\pi a \int_{\varrho=\varepsilon}^a \sqrt{1 + \sinh^2 t} \cdot \frac{a}{2} \cosh t dt = \pi a^2 \int_{\text{Arsinh } \frac{2\varepsilon}{a}}^{\text{Arsinh } 2} \cosh^2 t dt \\
 &= \frac{\pi a^2}{2} \int_{\text{Arsinh } \frac{2\varepsilon}{a}}^{\text{Arsinh } 2} (1 + \cosh 2t) dt = \frac{\pi a^2}{2} \left[ t + \frac{1}{2} \sinh 2t \right]_{\text{Arsinh } \frac{2\varepsilon}{a}}^{\text{Arsinh } 2} \\
 &= \frac{\pi a^2}{2} \left\{ \text{Arsinh } 2 - \text{Arsinh } \frac{2\varepsilon}{a} \right\} + \frac{\pi a^2}{2} \left[ \sinh t \sqrt{1 + \sinh^2 t} \right]_{\text{Arsinh } \frac{2\varepsilon}{a}}^{\text{Arsinh } 2} \\
 &\rightarrow \frac{\pi a^2}{2} \text{Arsinh } 2 + \frac{\pi a^2}{2} \{ \ln(2 + \sqrt{5}) + 2\sqrt{5} \}
 \end{aligned}$$

for  $\varepsilon \rightarrow 0+$ , and the improper surface integral converges towards the value

$$\int_{\mathcal{F}} \sqrt{\frac{a}{z}} dS = \frac{\pi a^2}{2} \{ \ln(2 + \sqrt{5}) + 2\sqrt{5} \}.$$

..... Alcatel-Lucent 

[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?



One generation's transformation is the next's status quo.  
In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".



Click on the ad to read more

**Example 30.9** Check if the surfaces of the bodies of revolution of the leading curves of the equations

$$y^2(a-x) = x^3 \quad (\text{rectangular}) \quad \text{and} \quad \varrho = \frac{a^2}{a^2 + z^2}, \quad z \in \mathbb{R}, \quad (\text{Semi-polar coordinates}),$$

can be given a finite area.

(The values shall not be computed).

**A** Improper surface integrals.

**D** Since we are only dealing with areas, the integrand is automatically positive. Truncate suitably before the computation of the surface integral, and then take the limit.

**I** 1) The curve  $\mathcal{K}$  of the equation

$$y^2(a-x) = x^3$$

is rotated around the asymptote  $x = a$ .

For symmetric reasons it suffices to consider  $y \geq 0$ , thus

$$y = x \sqrt{\frac{x}{a-x}} = x^{\frac{3}{2}}(a-x)^{-\frac{1}{2}}.$$

One easily sees that

$$\frac{dy}{dx} = \frac{1}{2} \sqrt{\frac{x}{(a-x)^3}} \cdot (3a-2x).$$

The length of the circle  $C_x$  (around the line  $x = a$ ) at the height  $y(x)$  is  $2\pi(a-x)$ , [In fact,  $0 \leq x < a$ ].

If we truncate at the height  $y(x_0)$  corresponding to some  $x_0 \in [0, a[$ , and remember the symmetry around  $y = 0$  we get the corresponding surface area,

$$\begin{aligned} & 2 \int_0^{x_0} \text{length}(C_x) \cdot \frac{dy}{dx} dx \\ &= 2 \int_0^{x_0} 2\pi(a-x) \cdot \frac{1}{2} \sqrt{\frac{x}{(a-x)^3}} \cdot (3a-2x) dx = 2\pi \int_0^{x_0} \sqrt{\frac{x}{a-x}} \{a+2(a-x)\} dx \\ &= 2\pi \int_0^{x_0} \left\{ a \sqrt{\frac{x}{a-x}} + 2\sqrt{x(a-x)} \right\} dx, \quad 0 < x_0 < a. \end{aligned}$$

We conclude that the surface has a finite area. The only problem is the term  $\sqrt{\frac{x}{a-x}}$  in the integrand, and

$$0 \leq \sqrt{\frac{x}{a-x}} \leq \sqrt{a} \cdot \frac{1}{\sqrt{a-x}} \quad \text{for } 0 < x < a,$$

and

$$\int_0^{x_0} \frac{dx}{\sqrt{a-x}} = [-2\sqrt{a-x}]_0^{x_0} = 2\{\sqrt{a} - \sqrt{a-x_0}\},$$

which converges towards  $2\sqrt{a}$  for  $x_0 \rightarrow a$ . Since the area of the surface is smaller than this value, we conclude that the improper surface integral exists.

REMARK. It is in fact possible to find the exact value. When we put  $t = \frac{x}{a-x}$  we get

$$x = \frac{at}{t+1} = a - \frac{a}{t+1},$$

hence

$$dx = \frac{a}{(t+1)^2} dt,$$

and

$$\begin{aligned} 2\pi \int a \sqrt{\frac{x}{a-x}} dx &= 2\pi a \int \sqrt{t} \cdot \frac{a}{(t+1)^2} dt = 2\pi a^2 \int u \cdot \frac{1}{(u^2+1)^2} \cdot 2u du \\ &= 2\pi a^2 \left\{ -\frac{u}{u^2+1} + \int \frac{1}{u^2+1} du \right\} = 2\pi a^2 \left\{ \arctan u - \frac{u}{u^2+1} \right\} \\ &= 2\pi a^2 \left\{ \arctan \sqrt{\frac{x}{a-x}} - \frac{\sqrt{\frac{x}{a-x}}}{\frac{x}{a-x} + 1} \right\} = 2\pi a^2 \left\{ \arctan \sqrt{\frac{x}{a-x}} - \frac{1}{a} \sqrt{x(a-x)} \right\}, \end{aligned}$$

hence by taking the limit

$$2\pi \int_0^a a \sqrt{\frac{x}{a-x}} dx = 2\pi a^2 \left\{ \frac{\pi}{2} - 0 \right\} = \pi^2 a^2.$$

The latter integral is calculated by noting that  $y = \sqrt{x(a-x)}$  for  $0 \leq x \leq a$  describes a half circle of centrum  $\frac{a}{2}$  and radius  $\frac{a}{2}$ , hence

$$4\pi \int_0^a \sqrt{x(a-x)} dx = 4\pi \cdot \frac{1}{2} \cdot \pi \left(\frac{a}{2}\right)^2 = \frac{\pi^2 a^2}{2}.$$

Summarizing, the improper surface area is convergent, and its value is

$$\pi^2 a^2 + \frac{\pi^2 a^2}{2} = \frac{3}{2} \pi^2 a^2. \quad \diamond$$

- 2) When the curve  $\varrho = \frac{a^2}{a^2 + z^2}$ ,  $z \in \mathbb{R}$ , is rotated around the  $Z$ -axis, we get an infinite surface which at the height  $z$  is cut into a circle  $C(x)$  of radius  $\varrho(z)$ , hence

$$\text{length}(C(x)) = 2\pi\varrho = \frac{2\pi a^3}{a^2 + z^2}.$$

When we put

$$\mathcal{F}_k = \{(x, y, z) \in \mathcal{F} \mid |z| \leq ka\}, \quad k > 0,$$

then we get

$$\begin{aligned} \text{area}(\mathcal{F}_k) &= 2 \int_0^{ka} \text{length}(C(x)) dz = 4\pi a^3 \int_0^{ka} \frac{1}{a^2 + z^2} dz \\ &= 4\pi a^2 \int_0^k \frac{1}{1+t^2} dt = 4\pi a^2 \arctan k \\ &\rightarrow 4\pi a^2 \cdot \frac{\pi}{2} = 2\pi^2 a^2 \quad \text{for } k \rightarrow +\infty. \end{aligned}$$

The improper surface area exists and its value is

$$\text{area}(\mathcal{F}) = 2\pi^2 a^2.$$

**Example 30.10** A surface  $\mathcal{F}$  is given by the equation

$$z = 1 + x^2 - y^2, \quad (x, y) \in \mathbb{R}^2.$$

1. Indicate the type of the surface and its vertices.
2. Find an equation of the tangent plane of  $\mathcal{F}$  through the point  $(2, 1, 4)$ .

Let  $q$  be a positive number. Let  $\mathcal{F}(q)$  denote the subset of  $\mathcal{F}$ , which is given by

$$z = 1 + x^2 - y^2, \quad x^2 + y^2 \leq q^2.$$

3. Compute the surface integral

$$I(q) = \int_{\mathcal{F}(q)} \frac{1}{(z + 3x^2 + 5y^2)^{3/2}} dS.$$

4. Explain shortly why

$$I = \int_{\mathcal{F}} \frac{1}{(z + 3x^2 + 5y^2)^{3/2}} dS$$

is an improper surface integral and prove that  $I$  is divergent.

- A** Surface; tangent plane; surface integral; improper surface integral.
- D** Identify the type of the surface; e.g. set up a parametric description (or use a formula) and find the field of the normal vectors. Calculate the surface integral by a reduction theorem. Note that the integrand is positive, and finally take the limit.




Figure 30.8: The surface  $\mathcal{F}(q)$  for  $q = 3$  with the projection  $D(q)$  onto the  $(x, y)$ -plane.

I 1) It follows from the rearrangement

$$z - 1 = x^2 - y^2$$

that the surface is an equilateral hyperbolic paraboloid with its vertex at  $(0, 0, 1)$ .

2) It follows from the parametric description

$$\mathbf{r}(x, y) = (x, y, 1 + x^2 - y^2), \quad (x, y) \in \mathbb{R}^2,$$

that

$$\frac{\partial \mathbf{r}}{\partial x} \times \frac{\partial \mathbf{r}}{\partial y} = \begin{vmatrix} \mathbf{e}_x & \mathbf{e}_y & \mathbf{e}_z \\ 1 & 0 & 2x \\ 0 & 1 & -2y \end{vmatrix} = (-2x, 2y, 1).$$

Then we check if the point  $(2, 1, 4)$  lies on  $\mathcal{F}$ :

$$1 + x^2 - y^2 = 1 + 4 - 1 = 4 = z,$$

thus  $(2, 1, 4) \in \mathcal{F}$ .



**Maastricht University** *Leading in Learning!*

**Join the best at  
the Maastricht University  
School of Business and  
Economics!**

**Top master's programmes**

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

**Visit us and find out why we are the best!  
Master's Open Day: 22 February 2014**

Maastricht  
University is  
the best specialist  
university in the  
Netherlands  
(Elsevier)

[www.mastersopenday.nl](http://www.mastersopenday.nl)



Click on the ad to read more

The normal vector is in this point

$$(-2x, 2y, 1) = (-4, 2, 1) = \mathbf{N},$$

and an equation of the tangent plane is

$$\begin{aligned} 0 &= \mathbf{N} \cdot (x - 2, y - 1, z - 4) = (-4, 2, 1) \cdot (x - 2, y - 1, z - 4) \\ &= -4x + 2y + z + 8 - 2 - 4 = -4x + 2y + z + 2, \end{aligned}$$

hence by a rearrangement,

$$z = 4x - 2y - 2.$$

- 3) The parametric domain for  $\mathcal{F}(q)$  is the disc in the  $(x, y)$ -plane

$$D(q) = \{(x, y) \mid x^2 + y^2 \leq q^2\}.$$

Since  $z = 1 + x^2 - y^2$  on  $\mathcal{F}(q)$ , it follows by the theorem of reduction that

$$\begin{aligned} I(q) &= \int_{\mathcal{F}(q)} \frac{1}{(z+3x^2+5y^2)^{3/2}} dS = \int_{D(q)} \frac{\|\mathbf{N}(x, y)\|}{(1+4x^2+4y^2)^{3/2}} dx dy \\ &= \int_{D(q)} \frac{(1+4x^2+4y^2)^{1/2}}{(1+4x^2+4y^2)^{3/2}} dx dy = \int_{D(q)} \frac{1}{1+4(x^2+y^2)} dx dy \\ &= \int_0^{2\pi} \left\{ \int_0^q \frac{1}{1+4\rho^2} \rho d\rho \right\} d\varphi = 2\pi \cdot \frac{1}{8} \ln(1+4q^2) = \frac{\pi}{4} \ln(1+4q^2). \end{aligned}$$

- 4) Now  $\mathcal{F}$  is unbounded, so  $I$  is an improper surface integral. The integrand is positive on  $\mathcal{F}$ , hence it suffices to take the limit  $q \rightarrow +\infty$  for  $I(q)$ . Then

$$I = \lim_{q \rightarrow +\infty} I(q) = \frac{\pi}{4} \lim_{q \rightarrow +\infty} \ln(1+4q^2) = +\infty,$$

which proves that the improper surface integral is divergent.

## 31 Formulæ

Some of the following formulæ can be assumed to be known from high school. It is highly recommended that one *learns most of these formulæ in this appendix by heart*.

### 31.1 Squares etc.

The following simple formulæ occur very frequently in the most different situations.

$$\begin{aligned} (a+b)^2 &= a^2 + b^2 + 2ab, & a^2 + b^2 + 2ab &= (a+b)^2, \\ (a-b)^2 &= a^2 + b^2 - 2ab, & a^2 + b^2 - 2ab &= (a-b)^2, \\ (a+b)(a-b) &= a^2 - b^2, & a^2 - b^2 &= (a+b)(a-b), \\ (a+b)^2 &= (a-b)^2 + 4ab, & (a-b)^2 &= (a+b)^2 - 4ab. \end{aligned}$$

### 31.2 Powers etc.

**Logarithm:**

$$\begin{aligned} \ln |xy| &= \ln |x| + \ln |y|, & x, y \neq 0, \\ \ln \left| \frac{x}{y} \right| &= \ln |x| - \ln |y|, & x, y \neq 0, \\ \ln |x^r| &= r \ln |x|, & x \neq 0. \end{aligned}$$

**Power function, fixed exponent:**

$$(xy)^r = x^r \cdot y^r, x, y > 0 \quad (\text{extensions for some } r),$$

$$\left( \frac{x}{y} \right)^r = \frac{x^r}{y^r}, x, y > 0 \quad (\text{extensions for some } r).$$

**Exponential, fixed base:**

$$\begin{aligned} a^x \cdot a^y &= a^{x+y}, & a > 0 & \quad (\text{extensions for some } x, y), \\ (a^x)^y &= a^{xy}, & a > 0 & \quad (\text{extensions for some } x, y), \end{aligned}$$

$$a^{-x} = \frac{1}{a^x}, a > 0, \quad (\text{extensions for some } x),$$

$$\sqrt[n]{a} = a^{1/n}, a \geq 0, \quad n \in \mathbb{N}.$$

**Square root:**

$$\sqrt{x^2} = |x|, \quad x \in \mathbb{R}.$$

**Remark 31.1** It happens quite frequently that students make errors when they try to apply these rules. They must be mastered! In particular, as one of my friends once put it: “If you can master the square root, you can master everything in mathematics!” Notice that this innocent looking square root is one of the most difficult operations in Calculus. Do not forget the *absolute value!* ◇

### 31.3 Differentiation

Here are given the well-known rules of differentiation together with some rearrangements which sometimes may be easier to use:

$$\{f(x) \pm g(x)\}' = f'(x) \pm g'(x),$$

$$\{f(x)g(x)\}' = f'(x)g(x) + f(x)g'(x) = f(x)g(x) \left\{ \frac{f'(x)}{f(x)} + \frac{g'(x)}{g(x)} \right\},$$

where the latter rearrangement presupposes that  $f(x) \neq 0$  and  $g(x) \neq 0$ .

If  $g(x) \neq 0$ , we get the usual formula known from high school

$$\left\{ \frac{f(x)}{g(x)} \right\}' = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}.$$

It is often more convenient to compute this expression in the following way:

$$\left\{ \frac{f(x)}{g(x)} \right\} = \frac{d}{dx} \left\{ f(x) \cdot \frac{1}{g(x)} \right\} = \frac{f'(x)}{g(x)} - \frac{f(x)g'(x)}{g(x)^2} = \frac{f(x)}{g(x)} \left\{ \frac{f'(x)}{f(x)} - \frac{g'(x)}{g(x)} \right\},$$

where the former expression often is *much easier* to use in practice than the usual formula from high school, and where the latter expression again presupposes that  $f(x) \neq 0$  and  $g(x) \neq 0$ . Under these assumptions we see that the formulæ above can be written

$$\frac{\{f(x)g(x)\}'}{f(x)g(x)} = \frac{f'(x)}{f(x)} + \frac{g'(x)}{g(x)},$$

$$\frac{\{f(x)/g(x)\}'}{f(x)/g(x)} = \frac{f'(x)}{f(x)} - \frac{g'(x)}{g(x)}.$$

Since

$$\frac{d}{dx} \ln |f(x)| = \frac{f'(x)}{f(x)}, \quad f(x) \neq 0,$$

we also name these the *logarithmic derivatives*.

Finally, we mention the rule of **differentiation of a composite function**

$$\{f(\varphi(x))\}' = f'(\varphi(x)) \cdot \varphi'(x).$$

We first differentiate the function itself; then the insides. This rule is a 1-dimensional version of the so-called *Chain rule*.

### 31.4 Special derivatives.

**Power like:**

$$\frac{d}{dx} (x^\alpha) = \alpha \cdot x^{\alpha-1}, \quad \text{for } x > 0, \quad (\text{extensions for some } \alpha).$$

$$\frac{d}{dx} \ln |x| = \frac{1}{x}, \quad \text{for } x \neq 0.$$

**Exponential like:**

$$\begin{aligned}\frac{d}{dx} \exp x &= \exp x, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} (a^x) &= \ln a \cdot a^x, && \text{for } x \in \mathbb{R} \text{ and } a > 0.\end{aligned}$$

**Trigonometric:**

$$\begin{aligned}\frac{d}{dx} \sin x &= \cos x, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \cos x &= -\sin x, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \tan x &= 1 + \tan^2 x = \frac{1}{\cos^2 x}, && \text{for } x \neq \frac{\pi}{2} + p\pi, p \in \mathbb{Z}, \\ \frac{d}{dx} \cot x &= -(1 + \cot^2 x) = -\frac{1}{\sin^2 x}, && \text{for } x \neq p\pi, p \in \mathbb{Z}.\end{aligned}$$

**Hyperbolic:**

$$\begin{aligned}\frac{d}{dx} \sinh x &= \cosh x, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \cosh x &= \sinh x, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \tanh x &= 1 - \tanh^2 x = \frac{1}{\cosh^2 x}, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \coth x &= 1 - \coth^2 x = -\frac{1}{\sinh^2 x}, && \text{for } x \neq 0.\end{aligned}$$

**Inverse trigonometric:**

$$\begin{aligned}\frac{d}{dx} \text{Arcsin } x &= \frac{1}{\sqrt{1-x^2}}, && \text{for } x \in ]-1, 1[, \\ \frac{d}{dx} \text{Arccos } x &= -\frac{1}{\sqrt{1-x^2}}, && \text{for } x \in ]-1, 1[, \\ \frac{d}{dx} \text{Arctan } x &= \frac{1}{1+x^2}, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \text{Arccot } x &= \frac{1}{1+x^2}, && \text{for } x \in \mathbb{R}.\end{aligned}$$

**Inverse hyperbolic:**

$$\begin{aligned}\frac{d}{dx} \text{Arsinh } x &= \frac{1}{\sqrt{x^2+1}}, && \text{for } x \in \mathbb{R}, \\ \frac{d}{dx} \text{Arcosh } x &= \frac{1}{\sqrt{x^2-1}}, && \text{for } x \in ]1, +\infty[, \\ \frac{d}{dx} \text{Artanh } x &= \frac{1}{1-x^2}, && \text{for } |x| < 1, \\ \frac{d}{dx} \text{Arcoth } x &= \frac{1}{1-x^2}, && \text{for } |x| > 1.\end{aligned}$$

**Remark 31.2** The derivative of the trigonometric and the hyperbolic functions are to some extent exponential like. The derivatives of the inverse trigonometric and inverse hyperbolic functions are power like, because we include the logarithm in this class. ◇

### 31.5 Integration

The most obvious rules are dealing with linearity

$$\int \{f(x) + \lambda g(x)\} dx = \int f(x) dx + \lambda \int g(x) dx, \quad \text{where } \lambda \in \mathbb{R} \text{ is a constant,}$$

and with the fact that differentiation and integration are “inverses to each other”, i.e. modulo some arbitrary constant  $c \in \mathbb{R}$ , which often tacitly is missing,

$$\int f'(x) dx = f(x).$$

If we in the latter formula replace  $f(x)$  by the product  $f(x)g(x)$ , we get by reading from the right to the left and then differentiating the product,

$$f(x)g(x) = \int \{f(x)g(x)\}' dx = \int f'(x)g(x) dx + \int f(x)g'(x) dx.$$

Hence, by a rearrangement

**The rule of partial integration:**

$$\int f'(x)g(x) dx = f(x)g(x) - \int f(x)g'(x) dx.$$

The differentiation is moved from one factor of the integrand to the other one by changing the sign and adding the term  $f(x)g(x)$ .

**Remark 31.3** This technique was earlier used a lot, but is almost forgotten these days. It must be revived, because MAPLE and pocket calculators apparently do not know it. It is possible to construct examples where these devices cannot give the exact solution, unless you first perform a partial integration yourself. ◇

**Remark 31.4** This method can also be used when we estimate integrals which cannot be directly calculated, because the antiderivative is not contained in e.g. the catalogue of MAPLE. The idea is by a succession of partial integrations to make the new integrand smaller. ◇

**Integration by substitution:**

If the integrand has the special structure  $f(\varphi(x)) \cdot \varphi'(x)$ , then one can change the variable to  $y = \varphi(x)$ :

$$\int f(\varphi(x)) \cdot \varphi'(x) dx = " \int f(\varphi(x)) d\varphi(x) " = \int_{y=\varphi(x)} f(y) dy.$$

### Integration by a monotonous substitution:

If  $\varphi(y)$  is a *monotonous* function, which maps the  $y$ -interval *one-to-one* onto the  $x$ -interval, then

$$\int f(x) dx = \int_{y=\varphi^{-1}(x)} f(\varphi(y))\varphi'(y) dy.$$

**Remark 31.5** This rule is usually used when we have some “ugly” term in the integrand  $f(x)$ . The idea is to put this ugly term equal to  $y = \varphi^{-1}(x)$ . When e.g.  $x$  occurs in  $f(x)$  in the form  $\sqrt{x}$ , we put  $y = \varphi^{-1}(x) = \sqrt{x}$ , hence  $x = \varphi(y) = y^2$  and  $\varphi'(y) = 2y$ .  $\diamond$

## 31.6 Special antiderivatives

Power like:

$$\int \frac{1}{x} dx = \ln|x|, \quad \text{for } x \neq 0. \quad (\text{Do not forget the numerical value!})$$

$$\int x^\alpha dx = \frac{1}{\alpha+1} x^{\alpha+1}, \quad \text{for } \alpha \neq -1,$$

$$\int \frac{1}{1+x^2} dx = \arctan x, \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right|, \quad \text{for } x \neq \pm 1,$$

$$\int \frac{1}{1-x^2} dx = \operatorname{Artanh} x, \quad \text{for } |x| < 1,$$

$$\int \frac{1}{1-x^2} dx = \operatorname{Arcoth} x, \quad \text{for } |x| > 1,$$

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x, \quad \text{for } |x| < 1,$$

$$\int \frac{1}{\sqrt{1-x^2}} dx = -\arccos x, \quad \text{for } |x| < 1,$$

$$\int \frac{1}{\sqrt{x^2+1}} dx = \operatorname{Arsinh} x, \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{1}{\sqrt{x^2+1}} dx = \ln(x + \sqrt{x^2+1}), \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{x}{\sqrt{x^2-1}} dx = \sqrt{x^2-1}, \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{1}{\sqrt{x^2-1}} dx = \operatorname{Arcosh} x, \quad \text{for } x > 1,$$

$$\int \frac{1}{\sqrt{x^2-1}} dx = \ln|x + \sqrt{x^2-1}|, \quad \text{for } x > 1 \text{ eller } x < -1.$$

There is an error in the programs of the pocket calculators TI-92 and TI-89. The *numerical signs* are missing. It is obvious that  $\sqrt{x^2-1} < |x|$  so if  $x < -1$ , then  $x + \sqrt{x^2-1} < 0$ . Since you cannot take the logarithm of a negative number, these pocket calculators will give an error message.

**Exponential like:**

$$\int \exp x \, dx = \exp x, \quad \text{for } x \in \mathbb{R},$$

$$\int a^x \, dx = \frac{1}{\ln a} \cdot a^x, \quad \text{for } x \in \mathbb{R}, \text{ and } a > 0, a \neq 1.$$

**Trigonometric:**

$$\int \sin x \, dx = -\cos x, \quad \text{for } x \in \mathbb{R},$$

$$\int \cos x \, dx = \sin x, \quad \text{for } x \in \mathbb{R},$$

$$\int \tan x \, dx = -\ln |\cos x|, \quad \text{for } x \neq \frac{\pi}{2} + p\pi, \quad p \in \mathbb{Z},$$

$$\int \cot x \, dx = \ln |\sin x|, \quad \text{for } x \neq p\pi, \quad p \in \mathbb{Z},$$

$$\int \frac{1}{\cos x} \, dx = \frac{1}{2} \ln \left( \frac{1 + \sin x}{1 - \sin x} \right), \quad \text{for } x \neq \frac{\pi}{2} + p\pi, \quad p \in \mathbb{Z},$$

$$\int \frac{1}{\sin x} \, dx = \frac{1}{2} \ln \left( \frac{1 - \cos x}{1 + \cos x} \right), \quad \text{for } x \neq p\pi, \quad p \in \mathbb{Z},$$

$$\int \frac{1}{\cos^2 x} \, dx = \tan x, \quad \text{for } x \neq \frac{\pi}{2} + p\pi, \quad p \in \mathbb{Z},$$

$$\int \frac{1}{\sin^2 x} \, dx = -\cot x, \quad \text{for } x \neq p\pi, \quad p \in \mathbb{Z}.$$

**Hyperbolic:**

$$\int \sinh x \, dx = \cosh x, \quad \text{for } x \in \mathbb{R},$$

$$\int \cosh x \, dx = \sinh x, \quad \text{for } x \in \mathbb{R},$$

$$\int \tanh x \, dx = \ln \cosh x, \quad \text{for } x \in \mathbb{R},$$

$$\int \coth x \, dx = \ln |\sinh x|, \quad \text{for } x \neq 0,$$

$$\int \frac{1}{\cosh x} \, dx = \operatorname{Arctan}(\sinh x), \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{1}{\cosh x} \, dx = 2 \operatorname{Arctan}(e^x), \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{1}{\sinh x} \, dx = \frac{1}{2} \ln \left( \frac{\cosh x - 1}{\cosh x + 1} \right), \quad \text{for } x \neq 0,$$

$$\int \frac{1}{\sinh x} dx = \ln \left| \frac{e^x - 1}{e^x + 1} \right|, \quad \text{for } x \neq 0,$$

$$\int \frac{1}{\cosh^2 x} dx = \tanh x, \quad \text{for } x \in \mathbb{R},$$

$$\int \frac{1}{\sinh^2 x} dx = -\coth x, \quad \text{for } x \neq 0.$$

### 31.7 Trigonometric formulæ

The trigonometric formulæ are closely connected with circular movements. Thus  $(\cos u, \sin u)$  are the coordinates of a point  $P$  on the unit circle corresponding to the angle  $u$ , cf. figure A.1. This geometrical interpretation is used from time to time.




Figure 31.1: The unit circle and the trigonometric functions.

#### The fundamental trigonometric relation:

$$\cos^2 u + \sin^2 u = 1, \quad \text{for } u \in \mathbb{R}.$$

Using the previous geometric interpretation this means according to *Pythagoras's theorem*, that the point  $P$  with the coordinates  $(\cos u, \sin u)$  always has distance 1 from the origo  $(0, 0)$ , i.e. it is lying on the boundary of the circle of centre  $(0, 0)$  and radius  $\sqrt{1} = 1$ .

#### Connection to the complex exponential function:

The *complex exponential* is for imaginary arguments defined by

$$\exp(iu) := \cos u + i \sin u.$$

It can be checked that the usual functional equation for  $\exp$  is still valid for complex arguments. In other word: The definition above is extremely conveniently chosen.

By using the definition for  $\exp(iu)$  and  $\exp(-iu)$  it is easily seen that

$$\cos u = \frac{1}{2}(\exp(iu) + \exp(-iu)),$$

$$\sin u = \frac{1}{2i}(\exp(iu) - \exp(-iu)),$$

**Moivre's formula:** We get by expressing  $\exp(inu)$  in two different ways:

$$\exp(inu) = \cos nu + i \sin nu = (\cos u + i \sin u)^n.$$

**Example 31.1** If we e.g. put  $n = 3$  into Moivre's formula, we obtain the following typical application,

$$\begin{aligned} \cos(3u) + i \sin(3u) &= (\cos u + i \sin u)^3 \\ &= \cos^3 u + 3i \cos^2 u \cdot \sin u + 3i^2 \cos u \cdot \sin^2 u + i^3 \sin^3 u \\ &= \{\cos^3 u - 3 \cos u \cdot \sin^2 u\} + i\{3 \cos^2 u \cdot \sin u - \sin^3 u\} \\ &= \{4 \cos^3 u - 3 \cos u\} + i\{3 \sin u - 4 \sin^3 u\} \end{aligned}$$

When this is split into the real- and imaginary parts we obtain

$$\cos 3u = 4 \cos^3 u - 3 \cos u, \quad \sin 3u = 3 \sin u - 4 \sin^3 u. \quad \diamond$$

**Addition formulæ:**

$$\begin{aligned} \sin(u+v) &= \sin u \cos v + \cos u \sin v, \\ \sin(u-v) &= \sin u \cos v - \cos u \sin v, \\ \cos(u+v) &= \cos u \cos v - \sin u \sin v, \\ \cos(u-v) &= \cos u \cos v + \sin u \sin v. \end{aligned}$$

**Products of trigonometric functions to a sum:**

$$\begin{aligned} \sin u \cos v &= \frac{1}{2} \sin(u+v) + \frac{1}{2} \sin(u-v), \\ \cos u \sin v &= \frac{1}{2} \sin(u+v) - \frac{1}{2} \sin(u-v), \\ \sin u \sin v &= \frac{1}{2} \cos(u-v) - \frac{1}{2} \cos(u+v), \\ \cos u \cos v &= \frac{1}{2} \cos(u-v) + \frac{1}{2} \cos(u+v). \end{aligned}$$

**Sums of trigonometric functions to a product:**

$$\begin{aligned} \sin u + \sin v &= 2 \sin\left(\frac{u+v}{2}\right) \cos\left(\frac{u-v}{2}\right), \\ \sin u - \sin v &= 2 \cos\left(\frac{u+v}{2}\right) \sin\left(\frac{u-v}{2}\right), \\ \cos u + \cos v &= 2 \cos\left(\frac{u+v}{2}\right) \cos\left(\frac{u-v}{2}\right), \\ \cos u - \cos v &= -2 \sin\left(\frac{u+v}{2}\right) \sin\left(\frac{u-v}{2}\right). \end{aligned}$$

**Formulæ of halving and doubling the angle:**

$$\sin 2u = 2 \sin u \cos u,$$

$$\cos 2u = \cos^2 u - \sin^2 u = 2 \cos^2 u - 1 = 1 - 2 \sin^2 u,$$

$$\sin \frac{u}{2} = \pm \sqrt{\frac{1 - \cos u}{2}} \quad \text{followed by a discussion of the sign,}$$

$$\cos \frac{u}{2} = \pm \sqrt{\frac{1 + \cos u}{2}} \quad \text{followed by a discussion of the sign,}$$

### 31.8 Hyperbolic formulæ

These are very much like the trigonometric formulæ, and if one knows a little of Complex Function Theory it is realized that they are actually identical. The structure of this section is therefore the same as for the trigonometric formulæ. The reader should compare the two sections concerning similarities and differences.

**The fundamental relation:**

$$\cosh^2 x - \sinh^2 x = 1.$$

**Definitions:**

$$\cosh x = \frac{1}{2} (\exp(x) + \exp(-x)), \quad \sinh x = \frac{1}{2} (\exp(x) - \exp(-x)).$$

**“Moivre’s formula”:**

$$\exp(x) = \cosh x + \sinh x.$$

This is trivial and only rarely used. It has been included to show the analogy.

**Addition formulæ:**

$$\begin{aligned}\sinh(x+y) &= \sinh(x)\cosh(y) + \cosh(x)\sinh(y), \\ \sinh(x-y) &= \sinh(x)\cosh(y) - \cosh(x)\sinh(y), \\ \cosh(x+y) &= \cosh(x)\cosh(y) + \sinh(x)\sinh(y), \\ \cosh(x-y) &= \cosh(x)\cosh(y) - \sinh(x)\sinh(y).\end{aligned}$$

**Formulæ of halving and doubling the argument:**

$$\begin{aligned}\sinh(2x) &= 2\sinh(x)\cosh(x), \\ \cosh(2x) &= \cosh^2(x) + \sinh^2(x) = 2\cosh^2(x) - 1 = 2\sinh^2(x) + 1, \\ \sinh\left(\frac{x}{2}\right) &= \pm\sqrt{\frac{\cosh(x)-1}{2}} \quad \text{followed by a discussion of the sign,} \\ \cosh\left(\frac{x}{2}\right) &= \sqrt{\frac{\cosh(x)+1}{2}}.\end{aligned}$$

**Inverse hyperbolic functions:**

$$\begin{aligned}\text{Arsinh}(x) &= \ln\left(x + \sqrt{x^2 + 1}\right), \quad x \in \mathbb{R}, \\ \text{Arcosh}(x) &= \ln\left(x + \sqrt{x^2 - 1}\right), \quad x \geq 1, \\ \text{Artanh}(x) &= \frac{1}{2}\ln\left(\frac{1+x}{1-x}\right), \quad |x| < 1, \\ \text{Arcoth}(x) &= \frac{1}{2}\ln\left(\frac{x+1}{x-1}\right), \quad |x| > 1.\end{aligned}$$

### 31.9 Complex transformation formulæ

$$\begin{aligned}\cos(ix) &= \cosh(x), & \cosh(ix) &= \cos(x), \\ \sin(ix) &= i \sinh(x), & \sinh(ix) &= i \sin x.\end{aligned}$$

### 31.10 Taylor expansions

The generalized binomial coefficients are defined by

$$\binom{\alpha}{n} := \frac{\alpha(\alpha - 1) \cdots (\alpha - n + 1)}{1 \cdot 2 \cdots n},$$

with  $n$  factors in the numerator and the denominator, supplied with

$$\binom{\alpha}{0} := 1.$$

The Taylor expansions for *standard functions* are divided into *power like* (the radius of convergency is finite, i.e. = 1 for the standard series) and *exponential like* (the radius of convergency is infinite).

**Power like:**

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n, \quad |x| < 1,$$

$$\frac{1}{1+x} = \sum_{n=0}^{\infty} (-1)^n x^n, \quad |x| < 1,$$

$$(1+x)^n = \sum_{j=0}^n \binom{n}{j} x^j, \quad n \in \mathbb{N}, x \in \mathbb{R},$$

$$(1+x)^{\alpha} = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n, \quad \alpha \in \mathbb{R} \setminus \mathbb{N}, |x| < 1,$$

$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}, \quad |x| < 1,$$

$$\text{Arctan}(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}, \quad |x| < 1.$$

**Exponential like:**

$$\exp(x) = \sum_{n=0}^{\infty} \frac{1}{n!} x^n, \quad x \in \mathbb{R}$$

$$\exp(-x) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{n!} x^n, \quad x \in \mathbb{R}$$

$$\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n+1)!} x^{2n+1}, \quad x \in \mathbb{R},$$

$$\sinh(x) = \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} x^{2n+1}, \quad x \in \mathbb{R},$$

$$\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{1}{(2n)!} x^{2n}, \quad x \in \mathbb{R},$$

$$\cosh(x) = \sum_{n=0}^{\infty} \frac{1}{(2n)!} x^{2n}, \quad x \in \mathbb{R}.$$

### 31.11 Magnitudes of functions

We often have to compare functions for  $x \rightarrow 0+$ , or for  $x \rightarrow \infty$ . The simplest type of functions are therefore arranged in an hierarchy:

- 1) logarithms,
- 2) power functions,
- 3) exponential functions,
- 4) faculty functions.

When  $x \rightarrow \infty$ , a function from a higher class will always dominate a function from a lower class. More precisely:

**A)** A *power function* dominates a *logarithm* for  $x \rightarrow \infty$ :

$$\frac{(\ln x)^\beta}{x^\alpha} \rightarrow 0 \quad \text{for } x \rightarrow \infty, \quad \alpha, \beta > 0.$$

**B)** An *exponential* dominates a *power function* for  $x \rightarrow \infty$ :

$$\frac{x^\alpha}{a^x} \rightarrow 0 \quad \text{for } x \rightarrow \infty, \quad \alpha, a > 1.$$

**C)** The *faculty function* dominates an *exponential* for  $n \rightarrow \infty$ :

$$\frac{a^n}{n!} \rightarrow 0, \quad n \rightarrow \infty, \quad n \in \mathbb{N}, \quad a > 0.$$

**D)** When  $x \rightarrow 0+$  we also have that a *power function* dominates the *logarithm*:

$$x^\alpha \ln x \rightarrow 0-, \quad \text{for } x \rightarrow 0+, \quad \alpha > 0.$$



## Index

- absolute value 162
- acceleration 490
- addition 22
- affinity factor 173
- Ampère-Laplace law 1671
- Ampère-Maxwell's law 1678
- Ampère's law 1491, 1498, 1677, 1678, 1833
- Ampère's law for the magnetic field 1674
- angle 19
- angular momentum 886
- angular set 84
- annulus 176, 243
- anticommutative product 26
- antiderivative 301, 847
- approximating polynomial 304, 322, 326, 336, 404, 488, 632, 662
- approximation in energy 734
- Archimedes's spiral 976, 1196
- Archimedes's theorem 1818
- area 887, 1227, 1229, 1543
- area element 1227
- area of a graph 1230
- asteroid 1215
- asymptote 51
- axial moment 1910
- axis of revolution 181
- axis of rotation 34, 886
- axis of symmetry 49, 50, 53
- barycentre 885, 1910
- basis 22
- bend 486
- bijective map 153
- body of revolution 43, 1582, 1601
- boundary 37–39
- boundary curve 182
- boundary curve of a surface 182
- boundary point 920
- boundary set 21
- bounded map 153
- bounded set 41
- branch 184
- branch of a curve 492
- Brownian motion 884
- cardioid 972, 973, 1199, 1705
- Cauchy-Schwarz's inequality 23, 24, 26
- centre of gravity 1108
- centre of mass 885
- centrum 66
- chain rule 305, 333, 352, 491, 503, 581, 1215, 1489, 1493, 1808
- change of parameter 174
- circle 49
- circular motion 19
- circulation 1487
- circulation theorem 1489, 1491
- circumference 86
- closed ball 38
- closed differential form 1492
- closed disc 86
- closed domain 176
- closed set 21
- closed surface 182, 184
- closure 39
- clothoid 1219
- colour code 890
- compact set 186, 580, 1813
- compact support 1813
- complex decomposition 69
- composite function 305
- conductivity of heat 1818
- cone 19, 35, 59, 251
- conic section 19, 47, 54, 239, 536
- conic sectional conic surface 59, 66
- connected set 175, 241
- conservation of electric charge 1548, 1817
- conservation of energy 1548, 1817
- conservation of mass 1548, 1816
- conservative force 1498, 1507
- conservative vector field 1489
- continuity equation 1548, 1569, 1767, 1817
- continuity 162, 186
- continuous curve 170, 483
- continuous extension 213
- continuous function 168
- continuous surfaces 177
- contraction 167
- convective term 492
- convex set 21, 22, 41, 89, 91, 175, 244
- coordinate function 157, 169
- coordinate space 19, 21

- Cornu's spiral 1219  
Coulomb field 1538, 1545, 1559, 1566, 1577  
Coulomb vector field 1585, 1670  
cross product 19, 163, 169, 1750  
cube 42, 82  
current density 1678, 1681  
current 1487, 1499  
curvature 1219  
curve 227  
curve length 1165  
curved space integral 1021  
cusp 486, 487, 489  
cycloid 233, 1215  
cylinder 34, 42, 43, 252  
cylinder of revolution 500  
cylindric coordinates 15, 21, 34, 147, 181, 182, 289, 477, 573, 841, 1009, 1157, 1347, 1479, 1651, 1801  
cylindric surface 180, 245, 247, 248, 499, 1230  
degree of trigonometric polynomial 67  
density 885  
density of charge 1548  
density of current 1548  
derivative 296  
derivative of inverse function 494  
Descartes'a leaf 974  
dielectric constant 1669, 1670  
difference quotient 295  
differentiability 295  
differentiable function 295  
differentiable vector function 303  
differential 295, 296, 325, 382, 1740, 1741  
differential curves 171  
differential equation 369, 370, 398  
differential form 848  
differential of order  $p$  325  
differential of vector function 303  
diffusion equation 1818  
dimension 1016  
direction 334  
direction vector 172  
directional derivative 317, 334, 375  
directrix 53  
Dirichlet/Neumann problem 1901  
displacement field 1670  
distribution of current 886  
divergence 1535, 1540, 1542, 1739, 1741, 1742  
divergence free vector field 1543  
dodecahedron 83  
domain 153, 176  
domain of a function 189  
dot product 19, 350, 1750  
double cone 252  
double point 171  
double vector product 27  
eccentricity 51  
eccentricity of ellipse 49  
eigenvalue 1906  
elasticity 885, 1398  
electric field 1486, 1498, 1679  
electrical dipole moment 885  
electromagnetic field 1679  
electromagnetic potentials 1819  
electromotive force 1498  
electrostatic field 1669  
element of area 887  
elementary chain rule 305  
elementary fraction 69  
ellipse 48–50, 92, 113, 173, 199, 227  
ellipsoid 56, 66, 110, 197, 254, 430, 436, 501, 538, 1107  
ellipsoid of revolution 111  
ellipsoidal disc 79, 199  
ellipsoidal surface 180  
elliptic cylindric surface 60, 63, 66, 106  
elliptic paraboloid 60, 62, 66, 112, 247  
elliptic paraboloid of revolution 624  
energy 1498  
energy density 1548, 1818  
energy theorem 1921  
entropy 301  
Euclidean norm 162  
Euclidean space 19, 21, 22  
Euler's spiral 1219  
exact differential form 848  
exceptional point 594, 677, 920  
expansion point 327  
explicit given function 161  
extension map 153  
exterior 37–39  
exterior point 38  
extremum 580, 632  
Faraday-Henry law of electromagnetic induction 1676  
Fick's first law of diffusion 297

- Fick's law 1818  
field line 160  
final point 170  
fluid mechanics 491  
flux 1535, 1540, 1549  
focus 49, 51, 53  
force 1485  
Fourier's law 297, 1817  
function in several variables 154  
functional matrix 303  
fundamental theorem of vector analysis 1815  
  
Gaussian integral 938  
Gauß's law 1670  
Gauß's law for magnetism 1671  
Gauß's theorem 1499, 1535, 1540, 1549, 1580, 1718,  
    1724, 1737, 1746, 1747, 1749, 1751, 1817,  
    1818, 1889, 1890, 1913  
Gauß's theorem in  $\mathbb{R}^2$  1543  
Gauß's theorem in  $\mathbb{R}^3$  1543  
general chain rule 314  
general coordinates 1016  
general space integral 1020  
general Taylor's formula 325  
generalized spherical coordinates 21  
generating curve 499  
generator 66, 180  
geometrical analysis 1015  
global minimum 613  
gradient 295, 296, 298, 339, 847, 1739, 1741  
gradient field 631, 847, 1485, 1487, 1489, 1491,  
    1916  
gradient integral theorem 1489, 1499  
graph 158, 179, 499, 1229  
Green's first identity 1890  
Green's second identity 1891, 1895  
Green's theorem in the plane 1661, 1669, 1909  
Green's third identity 1896  
Green's third identity in the plane 1898  
  
half-plane 41, 42  
half-strip 41, 42  
half disc 85  
harmonic function 426, 427, 1889  
heat conductivity 297  
heat equation 1818  
heat flow 297  
height 42  
helix 1169, 1235  
  
Helmholtz's theorem 1815  
homogeneous function 1908  
homogeneous polynomial 339, 372  
Hopf's maximum principle 1905  
hyperbola 48, 50, 51, 88, 195, 217, 241, 255, 1290  
hyperbolic cylindric surface 60, 63, 66, 105, 110  
hyperbolic paraboloid 60, 62, 66, 246, 534, 614,  
    1445  
hyperboloid 232, 1291  
hyperboloid of revolution 104  
hyperboloid of revolution with two sheets 111  
hyperboloid with one sheet 56, 66, 104, 110, 247,  
    255  
hyperboloid with two sheets 59, 66, 104, 110, 111,  
    255, 527  
hysteresis 1669  
  
identity map 303  
implicit given function 21, 161  
implicit function theorem 492, 503  
improper integral 1411  
improper surface integral 1421  
increment 611  
induced electric field 1675  
induction field 1671  
infinitesimal vector 1740  
infinity, signed 162  
infinity, unspecified 162  
initial point 170  
injective map 153  
inner product 23, 29, 33, 163, 168, 1750  
inspection 861  
integral 847  
integral over cylindric surface 1230  
integral over surface of revolution 1232  
interior 37–40  
interior point 38  
intrinsic boundary 1227  
isolated point 39  
Jacobian 1353, 1355  
  
Kronecker symbol 23  
  
Laplace equation 1889  
Laplace force 1819  
Laplace operator 1743  
latitude 35  
length 23  
level curve 159, 166, 198, 492, 585, 600, 603

- level surface 198, 503  
limit 162, 219  
line integral 1018, 1163  
line segment 41  
Linear Algebra 627  
linear space 22  
local extremum 611  
logarithm 189  
longitude 35  
Lorentz condition 1824
- Maclaurin's trisectrix 973, 975  
magnetic circulation 1674  
magnetic dipole moment 886, 1821  
magnetic field 1491, 1498, 1679  
magnetic flux 1544, 1671, 1819  
magnetic force 1674  
magnetic induction 1671  
magnetic permeability of vacuum 1673  
magnostatic field 1671  
main theorems 185  
major semi-axis 49  
map 153  
MAPLE 55, 68, 74, 156, 171, 173, 341, 345, 350, 352–354, 356, 357, 360, 361, 363, 364, 366, 368, 374, 384–387, 391–393, 395–397, 401, 631, 899, 905–912, 914, 915, 917, 919, 922–924, 926, 934, 935, 949, 951, 954, 957–966, 968, 971–973, 975, 1032–1034, 1036, 1037, 1039, 1040, 1042, 1053, 1059, 1061, 1064, 1066–1068, 1070–1072, 1074, 1087, 1089, 1091, 1092, 1094, 1095, 1102, 1199, 1200  
matrix product 303  
maximal domain 154, 157  
maximum 382, 579, 612, 1916  
maximum value 922  
maximum-minimum principle for harmonic functions 1895  
Maxwell relation 302  
Maxwell's equations 1544, 1669, 1670, 1679, 1819  
mean value theorem 321, 884, 1276, 1490  
mean value theorem for harmonic functions 1892  
measure theory 1015  
Mechanics 15, 147, 289, 477, 573, 841, 1009, 1157, 1347, 1479, 1651, 1801, 1921  
meridian curve 181, 251, 499, 1232  
meridian half-plane 34, 35, 43, 181, 1055, 1057, 1081
- method of indefinite integration 859  
method of inspection 861  
method of radial integration 862  
minimum 186, 178, 579, 612, 1916  
minimum value 922  
minor semi-axis 49  
mmf 1674  
Möbius strip 185, 497  
Moivre's formula 122, 264, 452, 548, 818, 984, 1132, 1322, 1454, 1626, 1776, 1930  
monopole 1671  
multiple point 171
- nabla 296, 1739  
nabla calculus 1750  
nabla notation 1680  
natural equation 1215  
natural parametric description 1166, 1170  
negative definite matrix 627  
negative half-tangent 485  
neighbourhood 39  
neutral element 22  
Newton field 1538  
Newton-Raphson iteration formula 583  
Newton's second law 1921  
non-oriented surface 185  
norm 19, 23  
normal 1227  
normal derivative 1890  
normal plane 487  
normal vector 496, 1229
- octant 83  
Ohm's law 297  
open ball 38  
open domain 176  
open set 21, 39  
order of expansion 322  
order relation 579  
ordinary integral 1017  
orientation of a surface 182  
orientation 170, 172, 184, 185, 497  
oriented half line 172  
oriented line 172  
oriented line segment 172  
orthonormal system 23
- parabola 52, 53, 89–92, 195, 201, 229, 240, 241  
parabolic cylinder 613

- parabolic cylindric surface 64, 66  
paraboloid of revolution 207, 613, 1435  
parallelepipedum 27, 42  
parameter curve 178, 496, 1227  
parameter domain 1227  
parameter of a parabola 53  
parametric description 170, 171, 178  
parfrac 71  
partial derivative 298  
partial derivative of second order 318  
partial derivatives of higher order 382  
partial differential equation 398, 402  
partial fraction 71  
Peano 483  
permeability 1671  
piecewise  $C^k$ -curve 484  
piecewise  $C^n$ -surface 495  
plane 179  
plane integral 21, 887  
point of contact 487  
point of expansion 304, 322  
point set 37  
Poisson's equation 1814, 1889, 1891, 1901  
polar coordinates 15, 19, 21, 30, 85, 88, 147, 163,  
172, 213, 219, 221, 289, 347, 388, 390,  
477, 573, 611, 646, 720, 740, 841, 936,  
1009, 1016, 1157, 1165, 1347, 1479, 1651,  
1801  
polar plane integral 1018  
polynomial 297  
positive definite matrix 627  
positive half-tangent 485  
positive orientation 173  
potential energy 1498  
pressure 1818  
primitive 1491  
primitive of gradient field 1493  
prism 42  
Probability Theory 15, 147, 289, 477, 573, 841,  
1009, 1157, 1347, 1479, 1651, 1801  
product set 41  
projection 23, 157  
proper maximum 612, 618, 627  
proper minimum 612, 613, 618, 627  
pseudo-sphere 1434  
Pythagoras's theorem 23, 25, 30, 121, 451, 547,  
817, 983, 1131, 1321, 1453, 1625, 1775,  
1929  
quadrant 41, 42, 84  
quadratic equation 47  
range 153  
rectangle 41, 87  
rectangular coordinate system 29  
rectangular coordinates 15, 21, 22, 147, 289, 477,  
573, 841, 1009, 1016, 1079, 1157, 1165,  
1347, 1479, 1651, 1801  
rectangular plane integral 1018  
rectangular space integral 1019  
rectilinear motion 19  
reduction of a surface integral 1229  
reduction of an integral over cylindric surface 1231  
reduction of surface integral over graph 1230  
reduction theorem of line integral 1164  
reduction theorem of plane integral 937  
reduction theorem of space integral 1021, 1056  
restriction map 153  
Riccati equation 369  
Riesz transformation 1275  
Rolle's theorem 321  
rotation 1739, 1741, 1742  
rotational body 1055  
rotational domain 1057  
rotational free vector field 1662  
rules of computation 296  
saddle point 612  
scalar field 1485  
scalar multiplication 22, 1750  
scalar potential 1807  
scalar product 169  
scalar quotient 169  
second differential 325  
semi-axis 49, 50  
semi-definite matrix 627  
semi-polar coordinates 15, 19, 21, 33, 147, 181,  
182, 289, 477, 573, 841, 1009, 1016, 1055,  
1086, 1157, 1231, 1347, 1479, 1651, 1801  
semi-polar space integral 1019  
separation of the variables 853  
signed curve length 1166  
signed infinity 162  
simply connected domain 849, 1492  
simply connected set 176, 243  
singular point 487, 489  
space filling curve 171  
space integral 21, 1015

- specific capacity of heat 1818  
sphere 35, 179  
spherical coordinates 15, 19, 21, 34, 147, 179, 181, 289, 372, 477, 573, 782, 841, 1009, 1016, 1078, 1080, 1081, 1157, 1232, 1347, 1479, 1581, 1651, 1801  
spherical space integral 1020  
square 41  
star-shaped domain 1493, 1807  
star shaped set 21, 41, 89, 90, 175  
static electric field 1498  
stationary magnetic field 1821  
stationary motion 492  
stationary point 583, 920  
Statistics 15, 147, 289, 477, 573, 841, 1009, 1157, 1347, 1479, 1651, 1801  
step line 172  
Stokes's theorem 1499, 1661, 1676, 1679, 1746, 1747, 1750, 1751, 1811, 1819, 1820, 1913  
straight line (segment) 172  
strip 41, 42  
substantial derivative 491  
surface 159, 245  
surface area 1296  
surface integral 1018, 1227  
surface of revolution 110, 111, 181, 251, 499  
surjective map 153
- tangent 486  
tangent plane 495, 496  
tangent vector 178  
tangent vector field 1485  
tangential line integral 861, 1485, 1598, 1600, 1603  
Taylor expansion 336  
Taylor expansion of order 2, 323  
Taylor's formula 321, 325, 404, 616, 626, 732  
Taylor's formula in one dimension 322  
temperature 297  
temperature field 1817  
tetrahedron 93, 99, 197, 1052  
Thermodynamics 301, 504  
top point 49, 50, 53, 66  
topology 15, 19, 37, 147, 289, 477, 573, 841, 1009, 1157, 1347, 1479, 1651, 1801  
torus 43, 182–184  
transformation formulæ 1353  
transformation of space integral 1355, 1357  
transformation theorem 1354  
trapeze 99
- triangle inequality 23, 24  
triple integral 1022, 1053
- uniform continuity 186  
unit circle 32  
unit disc 192  
unit normal vector 497  
unit tangent vector 486  
unit vector 23  
unspecified infinity 162
- vector 22  
vector field 158, 296, 1485  
vector function 21, 157, 189  
vector product 19, 26, 30, 163, 169, 1227, 1750  
vector space 21, 22  
vectorial area 1748  
vectorial element of area 1535  
vectorial potential 1809, 1810  
velocity 490  
volume 1015, 1543  
volumen element 1015
- weight function 1081, 1229, 1906  
work 1498
- zero point 22  
zero vector 22
- ( $r, s, t$ )-method 616, 619, 633, 634, 638, 645–647, 652, 655  
 $C^k$ -curve 483  
 $C^n$ -functions 318  
1-1 map 153