



HUST

ĐẠI HỌC BÁCH KHOA HÀ NỘI
HANOI UNIVERSITY OF SCIENCE AND TECHNOLOGY

ONE LOVE. ONE FUTURE.



ĐẠI HỌC
BÁCH KHOA HÀ NỘI
HANOI UNIVERSITY
OF SCIENCE AND TECHNOLOGY

FUNDAMENTALS OF OPTIMIZATION

Integer Linear Programming

ONE LOVE. ONE FUTURE.

CONTENT

- Relaxation and Bound
- Branch and Bound
- Cutting plane
- Gomory Cut
- Branch and Cut

RELAXATION AND BOUND

- Given an Integer Program (IP)
- Find decreasing sequence of upper bounds

$$\bar{z}_1 > \bar{z}_1 > \dots > \bar{z}_s \geq z$$

- Find increasing sequence of lower bounds

$$\underline{z}_1 < \underline{z}_1 < \dots < \underline{z}_t \leq z$$

- Algorithm stop when $\bar{z}_s - \underline{z}_t \leq \varepsilon$

RELAXATION AND BOUND

- Primal bounds
 - Every feasible solution $x^* \in X$ provides a lower bound of the maximization problem: $\underline{z} = cx^* \leq z$
 - Example: in TSP, every closed tour is a upper bound of the objective function (as TSP is a minimization problem)

RELAXATION AND BOUND

- Dual bounds
 - Finding upper bounds for a maximization problem (or lower bounds for a minimization problem) gives dual bounds of the objective
- **Definition** A problem $(RP) z^R = \max\{f(x): x \in T \subseteq R^n\}$ is a relaxation of $(IP) z = \max\{c^T x: x \in X \subseteq Z^n\}$ if:
 - $X \subseteq T$
 - $f(x) \geq c^T x, \forall x \in X$


RELAXATION AND BOUND

- Linear Relaxation

- $Z^{LP} = \max\{c^T x : x \in P\}$ with $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ is a linear relaxation program of the (IP) $\max\{c^T x : x \in P \cap \mathbb{Z}^n\}$


BRANCH AND BOUND

- Feasible region of P_0 is divided into feasible regions of P_1 and P_2 : $X(P_0) = X(P_1) \cup X(P_2)$




BRANCH AND BOUND

- Feasible region of P_0 is divided into feasible regions of P_1 and P_2 : $X(P_0) = X(P_1) \cup X(P_2)$



BRANCH AND BOUND

- Feasible region of P_0 is divided into feasible regions of P_1 and P_2 : $X(P_0) = X(P_1) \cup X(P_2)$



BRANCH AND BOUND

- Branch and Bound schema


```
Initial problem  $S$  with formulation  $P$  on a list  $L$ ;  
Incumbent  $x^*$  is initialized with primal bound  $\underline{z} = -INF$ ;  
while  $L$  not empty do {  
    Select a problem  $S^i$  with formulation  $P^i$  from  $L$ ;  
    Solve LP relaxation over  $P^i$  got dual bound  $\bar{z}^i$  and solution  $x^i(LP)$ ;  
    if  $\bar{z}^i \leq \underline{z}$  then continue; // prune by dual bound;  
    if  $x^i(LP)$  integer then{  
         $\underline{z} = \bar{z}^i$  ;  
         $x^* = x^i(LP)$ ;  
    }else{  
        select a component  $x_i$  of  $x^i(LP)$  whose value  $\lambda_i$  is fractional;  
         $P_1^i = P^i \cup (x_i \leq \lfloor \lambda_i \rfloor)$ ,  $P_2^i = P^i \cup (x_i \geq \lceil \lambda_i \rceil)$ ;  
        add  $P_1^i$  and  $P_2^i$  to  $L$ ;  
    }  
}  
return  $x^*$ ;
```

CUTTING PLANE

- Given a (MIP) $\max\{c^T z: z \in X\}$
- Inequality $\pi z \leq \pi_0$ is called a valid inequality if $\pi z \leq \pi_0$ is true for all $z \in X$
- Finding valid inequalities allows us to narrow the search space, transform the (MIP) to a corresponding (LP) in which an optimal solution to (LP) is an optimal solution to the original (MIP)

CUTTING PLANE

- Example, consider a MIP with $X = \{(x_1, x_2): x_1 \leq 10x_2, 0 \leq x_1 \leq 14, x_2 \in \mathbb{Z}_+^1\}$
- Red lines represent X
- $x_1 \leq 6 + 4x_2$ is a valid inequality (dashed line)



CUTTING PLANE

- Example: **Integer Rounding**

- Consider feasible region $X = P \cap \mathbb{Z}^3$ where $P = \{x \in \mathbb{R}_+^3 : 5x_1 + 9x_2 + 13x_3 \geq 19\}$

- From $5x_1 + 9x_2 + 13x_3 \geq 19$ we have $x_1 + \frac{9}{5}x_2 + \frac{13}{5}x_3 \geq \frac{19}{5}$

$$\rightarrow x_1 + 2x_2 + 3x_3 \geq \frac{19}{5}$$

- As x_1, x_2, x_3 are integers, so we have

$$x_1 + 2x_2 + 3x_3 \geq \left\lceil \frac{19}{5} \right\rceil = 4 \text{ (this is a valid inequality for } X)$$

GOMORY CUT

- (IP) $\max \{cx: Ax = b, x \geq 0 \text{ and integer}\}$
- Solve corresponding linear programming relaxation (LP) $\max \{cx: Ax = b, x \geq 0\}$
- Suppose with an optimal basis, the (LP) is rewritten in the form

$$\overline{a}_{00} + \sum_{j \in J_N} \overline{a}_{0j} x_j \rightarrow \max$$

$$x_{B_u} + \sum_{j \in J_N} \overline{a}_{uj} x_j = \overline{a}_{u0}, u = 1, 2, \dots, m$$

$$x \geq 0 \text{ and integer}$$

with $\overline{a}_{0j} \leq 0$ (as these coefficients corresponds to a maximizer), and $\overline{a}_{u0} \geq 0$

GOMORY CUT

- (IP) $\max \{cx: Ax = b, x \geq 0 \text{ and integer}\}$
- Solve corresponding linear programming relaxation (LP) $\max \{cx: Ax = b, x \geq 0\}$
- Suppose with an optimal basis, the (LP) is rewritten in the form

$$\overline{a}_{00} + \sum_{j \in J_N} \overline{a}_{0j} x_j \rightarrow \max$$

$$x_{B_u} + \sum_{j \in J_N} \overline{a}_{uj} x_j = \overline{a}_{u0}, u = 1, 2, \dots, m$$

$$x \geq 0 \text{ and integer}$$

with $\overline{a}_{0j} \leq 0$ (as these coefficients corresponds to a maximizer), and $\overline{a}_{u0} \geq 0$

- If the basic optimal solution x^* is not integer, then there exists some row u with $\overline{a}_{u0}$ is not integer
 $\rightarrow$ Create a Gomory cut $x_{B_u} + \sum_{j \in J_N} \lfloor \overline{a}_{uj} \rfloor x_j \leq \lfloor \overline{a}_{u0} \rfloor$ (1)

- **Example.** Consider the Integer Linear Program (ILP)

$$f(x_1, x_2, x_3, x_4, x_5) = x_1 + x_2 \rightarrow \max$$

$$2x_1 + x_2 + x_3 = 8$$

$$3x_1 + 4x_2 + x_4 = 24$$

$$x_1 - x_2 + x_5 = 2$$

$$x_1, x_2, x_3, x_4, x_5 \geq 0 \text{ and integer}$$

- Solve the corresponding (LP)

$$f(x_1, x_2, x_3, x_4, x_5) = x_1 + x_2 \rightarrow \max$$

$$2x_1 + x_2 + x_3 = 8$$

$$3x_1 + 4x_2 + x_4 = 24$$

$$x_1 - x_2 + x_5 = 2$$

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

→ Optimal solution (1.6, 4.8, 0, 0, 5.2) with $J_B = (1, 2, 5)$ and $J_N = (3, 4)$

Rewrite the original (ILP)

$$f(x_1, x_2, x_3, x_4, x_5) = 6.4 - 0.2x_3 - 0.2x_4 \rightarrow \max$$

$$x_1 + 0.8x_3 - 0.2x_4 = 1.6$$

$$x_2 - 0.6x_3 + 0.4x_4 = 4.8$$

$$x_5 - 1.4x_3 + 0.6x_4 = 5.2$$

$$x_1, x_2, x_3, x_4, x_5 \geq 0 \text{ and integer}$$

GOMORY CUT

- Add gomory cut

- $x_1 + 0.8x_3 - 0.2x_4 = 1.6 \rightarrow \text{add gomory cut: } x_1 + 0x_3 - x_4 \leq 1$
- $x_2 - 0.6x_3 + 0.4x_4 = 4.8 \rightarrow \text{add gomory cut: } x_2 - x_3 + 0x_4 \leq 4$
- $x_5 - 1.4x_3 + 0.6x_4 = 5.2 \rightarrow \text{add gomory cut: } x_5 - 2x_3 + 0x_4 \leq 5$

A graphic on the left side of the slide. It features a dark blue background with a large, stylized circular pattern of red dots. The dots are arranged in concentric, slightly irregular rings, creating a textured, halftone-like effect. In the center of this pattern, the word "HUST" is written in a bold, white, sans-serif font.

HUST

THANK YOU !